Next-to-Leading Order QCD corrections to the Lifetime Difference of B_s Mesons

Alexander Lenz *

Technische Universität München Physik Department D-85747 Garching Germany E-mail: alenz@feynman.t30.physik.tu-muenchen.de

ABSTRACT: In this talk we present a calculation of the dacay rate difference in the neutral $B_s - \overline{B}_s$ system, $\Delta\Gamma_{B_s}$, in next-to-leading order (NLO) QCD. We find a sizeable decrease compared to leading-order (LO) estimates: $(\Delta\Gamma/\Gamma)_{B_s} = (f_{B_s}/210MeV)^2[0.006B(m_b)+0.150B_S(m_b)-0.0063]$ in terms of the bag parameters B and B_S in the NDR scheme. We put special emphasize on the theoretical and physical implications of this quantity.

1. Non-expert-introduction

As there were many students in the audience we will start with an elementary introduction. Neutral mesons are well known from lectures at the university and were mentioned here several times e.g. in [1, 2, 3, 4]. As in the K-system we have in the B_s -system flavour eigenstates which are defined by their quark content.

$$|B_s\rangle = (\bar{b}s) \; ; \; |\overline{B}_s\rangle = (b\bar{s}) \, .$$
 (1.1)

The *mass eigenstates* are linear combinations of the flavour eigenstates

$$|B_H\rangle = p|B_s\rangle - q|\overline{B}_s\rangle \tag{1.2}$$

$$|B_L\rangle = p|B_s\rangle + q|\overline{B}_s\rangle \tag{1.3}$$

with the normalization condition $|p|^2 + |q|^2 = 1$. B_H and B_L are the physical states. They have definite masses and lifetimes, but no definite CPquantum numbers. The mass eigenstates are in general mixtures of CP-odd and CP-even eigenstates.

The time evolution of the physical states is described by a simple Schrödinger equation

$$i\partial_t \vec{B} = \hat{H}\vec{B} \tag{1.4}$$

with

$$\vec{B} = \begin{pmatrix} |B_s\rangle \\ |\overline{B}_s\rangle \end{pmatrix}; \ \hat{H} = \begin{pmatrix} M - \frac{i}{2}\Gamma & M_{12} - \frac{i}{2}\Gamma_{12} \\ M_{12}^* - \frac{i}{2}\Gamma_{12}^* & M - \frac{i}{2}\Gamma \end{pmatrix}$$
(1.5)

To find the mass eigenstates and the eigenvalues of the mass operator and the dacay rate operator we have to diagonalize the hamiltonian. We get

$$\Delta M_B = M_H - M_L = 2Re(Q) \qquad (1.6)$$

$$\Delta \Gamma_B = \Gamma_L - \Gamma_H = 4Im(Q) \qquad (1.7)$$

with

$$Q = \sqrt{(M_{12} - \frac{i}{2}\Gamma_{12})(M_{12}^* - \frac{i}{2}\Gamma_{12}^*)} .$$
 (1.8)

If we neglect CP violation and expand in m_b^2/m_t^2 we can write with a very good precision

$$\Delta M_B = 2|M_{12}| \tag{1.9}$$

$$\Delta \Gamma_B = -2\Gamma_{12} . \tag{1.10}$$

^{*}in collaboration with Martin Beneke, Theory Division, CERN, CH-1211 Geneva 23, Switzerland; Gerhard Buchalla, Theory Division, CERN, CH-1211 Geneva 23, Switzerland; Christoph Greub, Institut für Theoretische Physik, Universität Bern, Sidlerstrasse 5, CH-3012 Berne, Switzerland; Ulrich Nierste, Fermilab, Theory Division MS106, IL-60510-500, USA.

The different neutral meson systems gave rise to important contributions to the field of high energy physics. In 1964 Christenson, Cronin, Fitch and Turlay [5] discovered indirect CP-violation¹ in the $K^0 - \overline{K}^0$ -system. The mass difference in the $B_d - \overline{B}_d$ -system was the first experimen-

Each of these terms can be expanded in a power series in the strong coupling constant

$$\Gamma_i = \Gamma_i^{(0)} + \frac{\alpha_s}{\pi} \Gamma_i^{(1)} + \cdots$$
 (1.12)

So $\Delta\Gamma_B$ has the following form

$$\Delta\Gamma_B = \frac{\Lambda^3}{m_b^3} \left(\Gamma_3^{(0)} + \frac{\alpha_s}{\pi} \Gamma_3^{(1)} + \ldots \right) + \frac{\Lambda^4}{m_b^4} \left(\Gamma_4^{(0)} + \ldots \right)$$
(1.13)

After this short introduction for non-experts we motivate the special interest in the quantity $\Delta\Gamma_{B_s}$.

2. Motivation

From a physical point of view one wants to know the exact value of the decay rate difference, because

- $(\Delta\Gamma/\Gamma)_{B_s}$ is expected to be large. LO [6] estimates give values up to 20%. This is on the border of the experimental visibility [7];
- a big value of ΔΓ_{B_s} would enable us to do novel studies of CP-violation without the need of tagging [8]. Tagging is a major expermintal difficulty in B-physics;
- in the ratio $\Delta\Gamma_{B_s}/\Delta M_{B_s}$ some of the nonperturbative parameters cancel [9, 10]. So we can get theoretically clean information on ΔM_{B_s} from a measurement of $\Delta\Gamma_{B_s}$;
- the decay rate difference can be used to search for non SM-physics. In [11] it was shown that $\Delta\Gamma_{new \ physics} \leq \Delta\Gamma_{SM}$.

In order to fullfill this physics program we need a relieable prediction in the standard model. Therefore we need in addition to the LO estimate $\Gamma_3^{(0)}$, which was calculated in [6]

the 1/m_b-corrections Γ₄⁽⁰⁾. They have been calculated by [9];

The Heavy Quark Expansion (HQE) is the theoretical framework to handle inclusive B-decays. It allows us to expand the dacay rate in the following way

important bound on the parameters of the CKM

matrix.

tal hint for a very large top quark mass, before

the indirect determination at LEP and before the

discovery at Tevatron. As m_t is by now quite well known, we can extract the CKM parameter $|V_{td}V_{tb}|$ from ΔM_{B_d} . The determination of the CKM parameters is crucial for a test of our un-

derstanding of the standard model and for the search for new physics. The mass difference in the $B_s - \overline{B}_s$ -system is not measured yet, but we have a lower limit from which we already get an

$$\Gamma = \Gamma_0 + \left(\frac{\Lambda}{m_b}\right)^2 \Gamma_2 + \left(\frac{\Lambda}{m_b}\right)^3 \Gamma_3 + \cdots \quad (1.11)$$

Here we have an systematic expansion in the small parameter Λ/m_b . The different terms have the following physical interpretations:

- Γ_0 : The leading term is described by the decay of a free quark (parton model), we have no non-perturbative corrections.
- Γ_1 : In the derivation of eq. (1.11) we make an operator product expansion. From dimensional reasons we do not get an operator which would contribute to this order in the HQE.²
- Γ₂: First non-perturbative corrections arise at the second order in the expansion due to the kinetic and the chromomagnetic operator. They can be regarded as the first terms in a non-relativistic expansion.

¹Indirect or equivalently mixing induced CP violation means that the physical states $K_{S/L}$ are not pure CPeigenstates. There is a big contribution of one CP-parity and a tiny of the opposite CP-parity. If the small contribution dacays, one speaks of indirect CP violation.

 $^{^{2}}$ Strictly spoken we get one operator of the appropriate dimension, but with the equations of motion we can incorporate it in the leading term.

- the non-perturbative matrix elements for the $\Delta B = 2$ operators, which arise in the calculation. Here a relieable prediction is still missing;
- the NLO QCD corrections to the leading term in the $1/m_b$ expansion, $\Gamma_3^{(1)}$. This was the aim of our work [10]. Besides the better accuracy and a reduction of the μ dependence there is a very important point: NLO-QCD correction are needed for the proper matching of the perturbative calculation to lattice calculations.

From a technical point of view this calculation was very interesting because

- our result provides the first calculation of perturbative QCD corrections beyond leading logarithmic order to spectator effects in the HQE. Soft gluon emmision from the spectator *s* quark leads to power-like infrared singularities in individual contributions. As a conceptual test of the HQE the final result has to be infrared finite [12].
- a crucial point in the derivation of the HQE is the validity of the operator product expansion. This assumption is known under the name quark hadron duality and can be tested via a comparison of theory and experiment. A recent discussion of that subject can be found in [13].

In the next chapter we will describe the calculation.

3. Calculation

The width difference in the $B^0-\overline{B}^0$ -system is defined as

$$\Delta \Gamma = \Gamma_L - \Gamma_H = -2\Gamma_{21} . \qquad (3.1)$$

The off-diagonal element of decay-width matrix can be related to the so-called transition operator \mathcal{T} via

$$\Gamma_{21} = \frac{1}{2M_{B_S}} \langle \bar{B}_S | \mathcal{T} | B_S \rangle \tag{3.2}$$

with

$$\mathcal{T} = \operatorname{Im} i \int d^4x \ T \ \mathcal{H}_{eff}(x) \mathcal{H}_{eff}(0) \ . \tag{3.3}$$

In \mathcal{T} we have a double insertion of the effective hamiltonian with the standard form [14]

$$\mathcal{H}_{eff} = \frac{G_F}{\sqrt{2}} V_{cb}^* V_{cs} \left(\sum_{r=1}^6 C_r Q_r + C_8 Q_8 \right) \ . \ (3.4)$$

 G_F denotes the Fermi constant, V_{pq} are the CKM matrix elements and Q_i are local $\Delta B = 1$ operators. The Wilson coefficients C_i describe the short distance physics and are known to NLO QCD.

Formally we proceed now with an operator product expansion of that product of two hamiltonians. In real life one has to calculate diagrams of the following form:



Figure 1: The imaginary part of massive two loop diagram of that form has to be calculated.

One can do the calculation in two different ways (we did it in both ways, to have a check):

• calculate the imaginary part of the two loop integrals

 or

• use Cutkosky rules and calculate virtual and real one loop corrections, followed by a phase space integration.

The result in LO QCD has the following form

$$\mathcal{T} = -\frac{G_F^2 m_b^2}{12\pi} \left(V_{cb}^* V_{cs} \right)^2 \left[G(z)Q + G_S(z)Q_S \right]$$
(3.5)

with $z = m_c^2/m_b^2$ and the $\Delta B = 2$ operators

$$Q = (\bar{b}_i s_i)_{V-A} (\bar{b}_j s_j)_{V-A}$$

$$Q_S = (\bar{b}_i s_i)_{S-P} (\bar{b}_j s_j)_{S-P} . \qquad (3.6)$$

In principle we have more operators, but we can reduce them to the two operators above with the use of Fierz identities 3 .

Equation (3.5) is an example of an operator product expansion of equation (3.3). We have reduced the double insertion of $\Delta B = 1$ operators, which appear in \mathcal{H}_{eff} , to a single insertion of an $\Delta B = 2$ operator. In principle we have integrated out the internal charm quarks in figure 1. For the NLO calculation we have to match the $\Delta B = 1$ double insertion with gluon exchange to a $\Delta B = 2$ insertion with gluon exchange. This means, we have to calculate the following diagrams:



Figure 2: All diagrams that have to be calculated for the NLO QCD determination of $\Delta \Gamma_{B_s}$.

These diagrams can be classified in the following way:

- $E_1 E_4$: Virtual one loop corrections to a $\Delta B = 2$ operator insertion.
- $D_1 D_{10}$: Imaginary part of virtual two loop corrections to a double insertion of $\Delta B = 1$ operators.
- D_{11}, D_{12} : Penguin contributions to the $\Delta B = 1$ double insertion.

The calculation of all these diagrams gives us the NLO QCD result.

4. Results

The result in NLO is:

$$\mathcal{T} = -\frac{G_F^2 m_b^2}{12\pi} \left(V_{cb}^* V_{cs} \right)^2 \left[G(z)Q - G_S(z)Q_S \right]$$
(4.1)

with the following numerical values for the Wilson coefficients

μ	$m_b/2$	m_b	$2m_b$
$G^{(0)}$	0.013	0.047	0.097
G	0.023	0.030	0.036
$G_{S}^{(0)}$	1.622	1.440	1.292
\overline{G}_S	0.743	0.937	1.018

with

$$G = G^{(0)} + \frac{\alpha}{4\pi} G^{(1)} . \qquad (4.2)$$

Here one can see two important points. First, the value for G_S is numerical dominant and second, the NLO values are considerably smaller than the LO values.

For the final result we parametrise the matrix elements of the $\Delta B = 2$ operators in the following way:

$$\langle \bar{B}_s | Q | B_s \rangle = \frac{8}{3} f_{B_s}^2 M_{B_s}^2 B$$

 $\langle \bar{B}_s | Q_S | B_s \rangle = -\frac{5}{3} f_{B_s}^2 M_{B_s}^2 \frac{M_{B_s}^2}{(\bar{m}_b + \bar{m}_s)^2} B_S .$

B and B_S are so-called bag parameters, f_{B_s} is the decay constant. The values of these parameters have to be determined by non-perturbative methods like lattice simulations. \bar{m}_q denotes the running quark mass in the \overline{MS} -scheme.

With the following input parameters

$$m_b = 4.8 \text{GeV} \quad \left(\frac{m_c}{m_b}\right)^2 = 0.085 \quad \bar{m}_s = 0.2 \text{GeV}$$
$$M_{B_s} = 5.37 \text{GeV} \quad B(B_s \mapsto X e \nu) = 0.104$$
$$(4.3)$$

we obtain for the relative dacay rate difference

$$\left(\frac{\Delta\Gamma}{\Gamma}\right)_{B_s} = \left(\frac{f_{B_s}}{210 \text{MeV}}\right)^2$$
$$[0.006B(m_b) + 0.150B_S(m_b) - 0.063]$$
$$(4.4)$$

A definitive determination of the two bag parameters is still missing. From the literature [15] we were able to extract **preliminary** values for the bag parameters

$$B(m_b) = 0.9$$
 $B_S(m_b) = 0.75$. (4.5)

 $^{^{3}\}mathrm{This}$ reduction is relativelyy tricky. For details see [10].

With that numbers at hand we obtain as a final result

$$\left[\left(\frac{\Delta\Gamma}{\Gamma}\right)_{B_s} = \left(\frac{f_{B_s}}{210\text{MeV}}\right)^2 \left(0.054^{+0.016}_{-0.032} \pm ???\right)\right].$$
(4.6)

The question marks remind us that we do not know the uncertainties in the numerical values for the bag parameters.

5. Disscussion and outlook

The LO estimate for the relative decay rate difference $\Delta\Gamma_{B_s}/\Gamma_{B_s} = \mathcal{O}(20\%)$ is considerably reduced due to several effects:

- the 1/m_b corrections are sizeable and give an absolute reduction of about - 6.3 % [9].
- the pure NLO QCD corrections are sizeable, too and give an absolute reduction of about - 4.8 % [10].
- with the NLO QCD corrections at hand we can perform a proper matching to the (preliminary) lattice calculations for the bag parameters. This tells us that we have to use a low value for the bag parameters, i.e. $B_S(m_b) = 0.75$ [10, 15]. Compared to the naive estimate $B_S = 1$, this is another absolute reduction of about - 3.8 %.

Unfortunately the value of $\Delta\Gamma_{B_s}/\Gamma_{B_s}$ has been pinned down to a value of about 5%. The LO prediction was just a the border of experimental visibility [7]. Now we will have to wait for the forthcoming experiments like HERA-B, Tevatron (run II) and LHC.

Another application of our calculation are inclusive indirect CP-asymmetries in the $b \rightarrow u\bar{u}d$ channel. For the complete NLO prediction of this quantity, Γ_{12} in the B_d system was missing. We get this value from our calculation with a trivial exchange of the CKM parameters and the limit $m_c \mapsto 0$. This allows a determination of the CKM-angle α [10, 16].

Acknowledgements. I want to thank the organizers of the Corfu Summer Institute on Elementary Particle Physics for their successful work, M. Beneke, G. Buchalla, C. Greub and U. Nierste for the pleasant collaboration and A.J. Buras for proofreading the manuscript.

References

- [1] A.Ali, these proceedings.
- [2] Branco, these proceedings.
- [3] Bertolini, these proceedings.
- [4] R. Rückl, these proceedings.
- [5] J.H. Christenson, J.W. Cronin, V. L. Fitch and R. Turlay, Phys. Rev. Lett. **13**, (1964), 138.
- [6] J.S. Hagelin, Nucl. Phys. B193, (1981), 123;
 E. Franco, M. Lusignoli and A. Pugliese, Nucl. Phys. B194, (1982), 403; L.L. Chau, Phys. Rep. 95, (1983), 1; A.J. Buras, W. Słominski and H. Steger, Nucl. Phys. B245, (1984), 369; M.B. Voloshin, N.G. Uraltsev, V.A. Khoze and M.A. Shifman, Sov. J. Nucl. Phys. 46, (1987), 112; A. Datta, E.A. Paschos and U. Türke, Phys. Lett. B196, (1987), 382; A. Datta, E.A. Paschos and Y.L. Wu, Nucl. Phys. B311, (1988), 35.
- [7] K. Hartkorn and H.-G. Moser, MPI-PhE/98-21, appears in European Physical Journal C.
- [8] I. Dunietz, Phys. Rev. D52, (1995), 3048. R.
 Fleischer and I. Dunietz, Phys. Lett. B387, (1996), 361; Phys. Rev. D55, (1997), 259.
- [9] M. Beneke, G. Buchalla and I. Dunietz, Phys. Rev. D54, (1996), 4419.
- [10] M. Beneke, G. Buchalla, C. Greub, A. Lenz and U. Nierste, hep-ph/9808385; to appear in Phys. Lett. B.
- [11] Y. Grossman, Phys. Lett. **B380**, 99 (1996).
- [12] I. Bigi and N. Uraltsev, Phys. Lett. B280, (1992), 271.
- [13] I. Bigi, M. Shifman, N. Uraltsev, A. Vainshtein Phys.Rev. **D59**, (1999), 054011.
- [14] G. Buchalla, A.J. Buras and M.E. Lautenbacher, Rev. Mod. Phys. 68, (1996), 1125.
- [15] R. Gupta, T. Bhattacharya and S.R. Sharpe, Phys. Rev. D55, (1997), 4036.
 J.M. Flynn and C.T. Sachrajda, [heplat/9710057], in *Heavy Flavours (2nd ed.)*, eds. A.J. Buras and M. Lindner, World Scientific, Singapore.
- [16] M. Beneke, G. Buchalla and I. Dunietz, Phys. Lett. B393, (1997), 132.