

Quantum Unique Ergodicity for Cayley graphs of quasirandom groups

Michael Magee, Joe Thomas, and Yufei Zhao

Abstract

A finite group G is called C -quasirandom (by Gowers) if all non-trivial irreducible complex representations of G have dimension at least C . For any unit ℓ^2 function on a finite group we associate the *quantum probability measure* on the group given by the absolute value squared of the function.

We show that if a group is highly quasirandom, in the above sense, then any Cayley graph of this group has an orthonormal eigenbasis of the adjacency operator such that the quantum probability measures of the eigenfunctions put close to the correct proportion of their mass on suitably selected subsets of the group that are not too small.

Contents

1	Introduction	1
1.1	QUE on manifolds	6
1.2	Outline of the paper	7
2	Background	7
2.1	Representation theory of finite groups	7
2.2	Cayley graphs	8
3	Random basis construction	8
3.1	Non self-dual representations	8
3.2	Self-dual representations	9
3.2.1	Real representations.	9
3.2.2	Quaternionic representations.	9
4	Deterministic error term for mean zero functions	10
5	Probabilistic ingredients	12
5.1	Large deviations estimates	12
5.2	Random matrix estimates	14
6	Proof of main results	17

1 Introduction

The main question of quantum chaos is to what extent ‘chaotic’ features of the geodesic flow on a manifold (for example, ergodicity, exponential mixing, etc.) manifest themselves in the corresponding quantized system; that is, the L^2 Laplace-Beltrami operator and its eigenvalues and eigenfunctions. One of the main questions here is whether the quantum probability measures associated to eigenfunctions of the Laplacian have unique weak-

limits (semiclassical measures) as the corresponding eigenvalue tends to infinity. If there is a unique limit, the manifold is called quantum uniquely ergodic.

In this paper, we work with *graphs* instead of manifolds and prove results in the spirit of quantum unique ergodicity for certain families $\{\mathcal{G}_i\}_{i \in \mathcal{I}}$, $\mathcal{I} \subset \mathbf{N}$ of d -regular graphs, with $d \geq 3$ fixed. We will always write \mathcal{G}_i to refer to such a family of graphs. We write V_i for the vertex set of \mathcal{G}_i , let $N \stackrel{\text{def}}{=} |V_i|$ and assume $N \rightarrow \infty$ as $i \rightarrow \infty$. Each \mathcal{G}_i has an adjacency matrix that has rows and columns indexed by V_i , a 1 in entry (x, y) if there is an edge between x and y , and 0 otherwise; we view this as an operator on $\ell^2(V_i)$. *In this paper, ℓ^2 norms will be defined with respect to the counting measure.*

Given an element $\varphi \in \ell^2(V_i)$ with $\|\varphi\|_{\ell^2} = 1$, which will usually be an eigenfunction of the adjacency operator of \mathcal{G}_i , we associate to φ the *quantum probability measure*¹ μ_φ on V_i defined by

$$\mu_\varphi \stackrel{\text{def}}{=} \sum_{v \in V_i} |\varphi(v)|^2 \delta_v,$$

where δ_v is the unit mass atom at v . Note that $\|\varphi\|_{\ell^2} = 1$ implies μ_φ is a probability measure.

We will say *quantum unique ergodicity* (QUE) holds for a sequence of adjacency operator eigenfunctions $\varphi_i \in \ell^2(V_i)$ with $\|\varphi_i\|_{\ell^2} = 1$ and a sequence of subsets $A_i \subset V_i$ if

$$\mu_{\varphi_i}[A_i] \rightarrow \frac{|A_i|}{|V_i|} = \frac{|A_i|}{N}$$

as $i \rightarrow \infty$. It is very hard in general to establish this bound for all A_i , so we will restrict to A_i that are not too small.

Suppose that G is a finite group and S is a symmetric subset of G , then we will denote the Cayley graph associated to the pair (G, S) by $\text{Cay}(G, S)$. We write \hat{G} for the equivalence classes of irreducible representations of G , and define

$$\mathfrak{D}(G) \stackrel{\text{def}}{=} \min_{(\rho, V) \in \hat{G} - \text{triv}} \dim V;$$

i.e. the smallest dimension of a non-trivial representation of G . Then in the language of Gowers from [Gow08], G is $\mathfrak{D}(G)$ -*quasirandom*². The first main theorem of the paper is the following.

Theorem 1.1. *Let G_i be finite groups with $|G_i| \xrightarrow{i \rightarrow \infty} \infty$, $S_i \subseteq G_i$ be symmetric subsets ($S_i = S_i^{-1}$), $\mathcal{G}_i = \text{Cay}(G_i, S_i)$ and $t_i > 0$. Moreover, let $M_i \in \mathbf{N}$ be such that*

$$2M_i \sum_{(\pi, V) \in \hat{G}_i - \text{triv}} (\dim V)^2 \left(6e^{-\frac{t_i \sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right) < 1, \quad (1.1)$$

and let $f_i^j : V_i \rightarrow \mathbb{R}$ be any collection of functions for $j = 1, \dots, M_i$ and $i \in \mathbf{N}$. Then, there exist orthonormal bases \mathcal{B}_i of $\ell^2(G_i)$ of real-valued eigenfunctions of \mathcal{G}_i such that for every

¹From the point of view of quantum mechanics, this is the probability density function.

²Before the formal naming of this property by Gowers, the property of a group G being $|G|^\delta$ -quasirandom was used to prove eigenvalue bounds in works of Sarnak and Xue [SX91] and Bourgain and Gamburd [BG08].

$\varphi \in \mathcal{B}_i$ and $j = 1, \dots, M_i$,

$$\left| \mu_\varphi[f_i^j] - \frac{\sum_{g \in G_i} f_i^j(g)}{|G_i|} \right| \leq t_i \frac{\|f_i^j\|_{\ell^2}}{\sqrt{|G_i|}}. \quad (1.2)$$

If $f_i^j = \mathbf{1}_{A_i^j}$ for some subsets $A_i^j \subseteq V_i$ then

$$\left| \mu_\varphi[A_i^j] - \frac{|A_i^j|}{|G_i|} \right| \leq t_i \frac{\sqrt{|A_i^j|}}{\sqrt{|G_i|}}; \quad (1.3)$$

which in particular implies that $\mu_\varphi[A_i^j]$ is asymptotic to $\frac{|A_i^j|}{|G_i|}$ as $i \rightarrow \infty$ whenever $\frac{t_i^2 |G_i|}{|A_i^j|} = o_{i \rightarrow \infty}(1)$.

Remark 1.2. The proof of Theorem 1.1 is slightly easier if one only wants complex orthonormal eigenbases; see Remark 6.4 at the end of the paper. In this case, one can also take the functions f_i^j to be complex-valued.

The condition (1.1) involving M_i and t_i displays a dependence between the desired strength of the QUE bound in (1.2), and the number of functions that one simultaneously wishes the bound to hold for. With knowledge on the size and number of irreducible representations of the group, one can be more precise with values for t_i and M_i .

The most simple case of this is as follows. For groups with $\mathfrak{D}(G) \geq \log^2(|G|)$ one can obtain at least logarithmic improvement in (1.2) while taking the number of functions to be polynomial in the size of the group.

Corollary 1.3. *Let $\varepsilon > 0$, and suppose that G is a finite group satisfying $\mathfrak{D}(G) \geq \log^2(|G|)$. Moreover, let $S \subseteq G$ be a symmetric subset and $\mathcal{G} = \text{Cay}(G, S)$. Then given $M \in \mathbf{N}$ satisfying $M \leq \min\left(\frac{1}{24}|G|^\varepsilon, \frac{1}{8}|G|^{-1}e^{\frac{\mathfrak{D}(G)}{12}}\right)$, and functions $f_i : V \rightarrow \mathbb{R}$ for $i = 1, \dots, M$, there exists an orthonormal basis \mathcal{B} of $\ell^2(G)$ of real-valued eigenfunctions of \mathcal{G} such that for every $\varphi \in \mathcal{B}$ and $i = 1, \dots, M$,*

$$\left| \mu_\varphi[f_i] - \frac{\sum_{g \in G} f_i(g)}{|G|} \right| \leq 64 \frac{(\varepsilon + 1) \log(|G|)}{\sqrt{\mathfrak{D}(G)}} \frac{\|f_i\|_{\ell^2}}{\sqrt{|G|}}. \quad (1.4)$$

Remark 1.4. The proof of Theorem 1.1 shows that if e.g. $\mathfrak{D}(G_i) \geq |G_i|^\alpha$ with $\alpha > 0$ as it is in cases of interest (see below), if we only want to obtain

$$\left| \mu_\varphi[f_i^j] - \frac{\sum_{g \in G_i} f_i^j(g)}{|G_i|} \right| = o\left(\frac{\|f_i^j\|_{\ell^2}}{\sqrt{|G_i|}}\right)$$

above then we can actually take $M_i \geq e^{c|G_i|^\beta}$ for $c, \beta > 0$ depending on α , i.e. take the number of functions f_i^j to be super-polynomial in $|G_i|$.

Example 1.5. If \mathcal{I} are the prime numbers, $G_p = \text{PSL}_2(\mathbb{F}_p)$, and \mathcal{G}_p are any Cayley graphs of $\text{PSL}_2(\mathbb{F}_p)$ with respect to symmetric generators, then a result of Frobenius gives

$$\mathfrak{D}(\text{PSL}_2(\mathbb{F}_p)) \geq \frac{p-1}{2},$$

and $|G_p| \approx p^3$. So in this setting, Theorem 1.1 gives that for any finite collection $A_p^1, \dots, A_p^m \subset V_p$ with $|A_p^j| \gg p^{2+\epsilon}$, there are real orthonormal eigenbases of $\ell^2(\text{PSL}_2(\mathbb{F}_p))$

such that for any elements φ_p of these bases,

$$\mu_{\varphi_p}[A_p^j] = \frac{|A_p^j|}{|\mathrm{PSL}_2(\mathbb{F}_p)|} (1 + O(p^{-\epsilon}))$$

as $p \rightarrow \infty$.

When $\mathfrak{D}(G)$ is polynomial in $|G|$, we can also obtain a quantum unique ergodicity result for partitions of the group into sets whose sizes are on scales of the order $|G|^{1-\eta}$ for some $\eta > 0$ dependent upon on the size of $\mathfrak{D}(G)$.

Corollary 1.6. *Let G be a finite group, $S \subseteq G$ be a symmetric subset and $\mathcal{G} = \mathrm{Cay}(G, S)$. Suppose that there exists an absolute constant $s > 0$ such that $\mathfrak{D}(G) \geq |G|^s$ and let $\eta = s - \varepsilon$ for any $0 < \varepsilon < s$. Let $A_i \subseteq G$ be a collection of subsets partitioning G with sizes satisfying $c|G|^{1-\eta} \leq |A_i| \leq C|G|^{1-\eta}$ for some absolute constants $c, C > 0$. Then, for $|G|$ sufficiently large (dependent only upon c and η) there is an orthonormal eigenbasis \mathcal{B} of the adjacency operator of G such that for every i and every $\varphi \in \mathcal{B}$,*

$$\left| \mu_\varphi[A_i] - \frac{|A_i|}{|G|} \right| \leq \frac{K \log |G|}{|G|^{\frac{1}{2}\varepsilon}} \frac{|A_i|}{|G|},$$

where $K > 0$ is a constant dependent only upon c .

So far we have dealt with groups that are at least $\log^2(|G|)$ -quasirandom. One key feature of the condition (1.1) is that it enables us to go beyond $\mathfrak{D}(G) \geq \log^2(|G|)$. This pertains to the important class of examples where G_n is either the alternating group $\mathrm{Alt}(n)$ or the symmetric group $\mathrm{Sym}(n)$.

Proposition 1.7. *Let $G_n = \mathrm{Alt}(n)$ or $\mathrm{Sym}(n)$, $S_n \subseteq G_n$ be symmetric subsets and $\mathcal{G}_n = \mathrm{Cay}(G_n, S_n)$. Then given $M_n \in \mathbf{N}$ satisfying $M_n = o_{n \rightarrow \infty}(n)$ and functions $f_i^n : V_n \rightarrow \mathbb{R}$ for $i = 1, \dots, M_n$, there exists an orthonormal basis \mathcal{B}_n of $\ell^2(G_n)$ of real-valued eigenfunctions of \mathcal{G}_n such that for every $\varphi \in \mathcal{B}_n$, $i = 1, \dots, M_n$, and n sufficiently large*

$$\left| \mu_\varphi[f_i] - \frac{\sum_{g \in G_n} f_i^n(g)}{|G_n|} \right| \leq 192 \frac{\log(n)}{\sqrt{n}} \frac{\|f_i^n\|_{\ell^2(G_n)}}{\sqrt{|G_n|}}. \quad (1.5)$$

The proof of Theorem 1.1 revolves around the fact that all eigenspaces of Cayley graphs arise from some irreducible representation of the group and hence have multiplicities at least the dimension of this corresponding representation. This leads to a dichotomy: either the eigenspace is trivial (which we can deal with directly) or has large dimension if the group is suitably quasirandom. In the latter case, this allows one to choose a random basis for the eigenspace using a random matrix of large dimension which is reflected in the condition (1.1).

We describe in §3 a random model for real eigenbases of Cayley graphs that arise from products of the classical compact groups with their Haar measures. This model was used by Sah, Sawhney, and Zhao in [SSZ20] to show the existence of eigenbases of Cayley graphs with close to optimal ℓ^∞ bounds. What we prove here is the following.

Theorem 1.8. *Let G be a finite group, $S \subseteq G$ be a symmetric subset and $\mathcal{G} = \mathrm{Cay}(G, S)$. Let $M \in \mathbf{N}$ and let $f_1, \dots, f_M \in \ell^2(G)$ be a collection of real-valued functions. Then, for*

any $t > 0$, with probability at least

$$1 - 2M \sum_{(\pi, V) \in \hat{G} - \text{triv}} (\dim V)^2 \left(6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right),$$

if \mathcal{B} is a random real orthonormal eigenbasis of \mathcal{G} as in §3, then for any $\varphi \in \mathcal{B}$ and any $i = 1, \dots, M$, we have

$$\left| \mu_\varphi[f_i] - \frac{\sum_{g \in G} f_i(g)}{|G|} \right| \leq t \frac{\|f_i\|_{\ell^2}}{\sqrt{|G|}}. \quad (1.6)$$

As indicated by Corollaries 1.3 and 1.6, it is good to know that there are an abundance of $|G|^\delta$ -quasirandom groups for $0 < \delta < 1$. Indeed, for *finite simple groups of Lie type* with rank r over finite fields, it is shown in the proof of [BGGT15, Prop. 3.2] (see also Remark 1.3.6 of [Tao15]) using earlier work of [LS74, SZ93] that such groups are $|G|^\delta$ -quasirandom with δ depending only on the rank r . We refer to [BGGT15, §5.2] for the precise definition of these groups. As such, the values of t in Theorems 1.1 and 1.8 can be taken to have decay that is *polynomial* in $|G|$ for this wide class of groups (see Corollary 1.3).

Let us now discuss the strength of the upper bound obtained in Theorems 1.1 and 1.8. Since the sum of squares of the dimensions of the irreducible representations of a group equal the size of the group,

$$\mathfrak{D}(G) \leq |G|^{\frac{1}{2}}$$

which means that the best possible value we could possibly obtain for the right hand side of (1.3) or (1.6) is

$$\frac{C \log(|G|)}{|G|^{\frac{3}{4}}} \|f\|_{\ell^2}.$$

This is still a factor of $|G|^{\frac{1}{4}}$ off from what is known about random regular graphs: recently Bauerschmidt, Huang, and Yau [BHY19] obtained a very strong version of QUE for random regular graphs with respect to the uniform model of fixed degree and number of vertices³.

Theorem 1.9 (Bauerschmidt-Huang-Yau [BHY19, Cor. 13]). *Let $d \gg 1$ and let \mathcal{G}_n be a uniformly random d -regular graph on n vertices. Suppose $f_n : V_n \rightarrow \mathbb{R}$, then with probability tending to one as $n \rightarrow \infty$, for any eigenfunction $\varphi \in \ell^2(V_n)$ of the adjacency operators of \mathcal{G}_n with eigenvalues λ_n satisfying $|\lambda_n \pm 2\sqrt{d-1}| > (\log n)^{-\frac{3}{2}}$,*

$$\left| \sum_{v \in V_n} f_n(v) |\varphi(v)|^2 - \frac{\sum_{v \in V_n} f_n(v)}{n} \right| \leq \frac{(\log n)^{250}}{n} \sqrt{\sum_{v \in V_n} |f_n(v)|^2}.$$

The first result about equidistribution of quantum probability measures of eigenfunctions⁴ on graphs was obtained by Anantharaman and Le Masson in [ALM15, Thm. 1].

Theorem 1.10 ([ALM15, Thm. 1]). *Let \mathcal{G}_i be d -regular, $d > 3$, and $N \stackrel{\text{def}}{=} |V(\mathcal{G}_i)| \rightarrow \infty$ as $i \rightarrow \infty$. Suppose that the sequence \mathcal{G}_i form a family of uniform expanders and converge to the infinite d -regular tree in the sense of Benjamini and Schramm [BS01]. Let $\{\varphi_j^{(i)}\}_{j=1}^N$ be an orthonormal basis of eigenfunctions of the adjacency operator of \mathcal{G}_i . Let $f_i : V_i \rightarrow \mathbb{C}$*

³See also [BKY17] for the case of growing degree.

⁴Strictly speaking, Theorem 1.10 is a result about Quantum *Ergodicity* rather than QUE.

be a sequence of functions with $\|f_i\|_\infty \leq 1$, then for any $\delta > 0$

$$\frac{1}{N} \left| \left\{ j \in [1, N] : \left| \sum_{v \in V_i} f_i(v) |\varphi_j^{(i)}(v)|^2 - \frac{1}{N} \sum_{v \in V_i} f_i(v) \right| > \delta \right\} \right| \rightarrow 0 \quad (1.7)$$

as $i \rightarrow \infty$.

For related results of quantum ergodicity on quantum graphs, see for example [BW16, AISW21]. See also the recent work of Naor, Sah, Sawhney and Zhao [NSSZ22] in the Cayley graph setting, where they prove an incomparable quantum ergodicity result, rather than quantum unique ergodicity.

1.1 QUE on manifolds

Because the type of results of the current paper draw their inspiration from analogous questions about manifolds, we include a brief discussion of the state of the art results in that setting.

Let M be a closed and connected Riemannian manifold and let $\{\varphi_j\}_{j \geq 1}$ be an orthonormal basis of $L^2(M)$ consisting of Laplacian eigenfunctions with corresponding eigenvalues $0 = \lambda_1 < \lambda_2 \leq \dots \rightarrow \infty$. A central question is the *quantum unique ergodicity conjecture* of Rudnick and Sarnak [RS94]. This says that if M is negatively curved, then the quantum probability measures of the eigenfunctions weak-* converge as $i \rightarrow \infty$ to the normalized Riemannian volume form. A more general statement of this conjecture involving microlocal lifts can be found in the survey article of Sarnak [Sar11]. For manifolds without negative curvature, there are counterexamples to this conjecture as illustrated for example by Hassel [Has10] for certain ergodic billiards, building upon earlier numerical work by O'Connor and Heller [OH88].

Despite counterexamples demonstrating that ergodicity alone is insufficient for quantum unique ergodicity, there is numerical evidence to support the conjecture in the presence of negative curvature [AS93, HR92]. In addition, there are striking results of Anantharaman and Nonnenmacher [AN07, Ana08] and Dyatlov and Jin [DJ18] regarding the entropy and support of possible limits of quantum probability measures. Moreover, Lindenstrauss [Lin06] (with an extension by Soundararajan [Sou10] for the non-compact case), proved that the quantum unique ergodicity conjecture holds for Hecke-Laplace eigenfunctions on arithmetic surfaces.

For closed Riemannian manifolds in general, ergodicity of the geodesic flow alone is sufficient to prove a weaker result known as quantum ergodicity. This result exhibits the existence of a density one subsequence of the quantum probability measures that weak-* converges to the normalized volume measure [Šni74, Zel87, Col85]. Theorem 1.10 above can be seen as a natural graph analogue of this weaker property. In the manifold setting, quantum ergodicity has also been investigated for random bases. For example, in [Zel92] it is shown that random (Haar unitary) eigenbases of the Laplacian for $L^2(S^2)$ are quantum ergodic with probability one, despite the standard basis of spherical harmonics failing to have this property. This is upgraded to quantum unique ergodicity in [Van97]. Similarly, quantum ergodicity and quantum mixing properties have been studied for random bases (not necessarily eigenbases) for general compact Riemannian manifolds [Zel96, Zel14] as well as quantum unique ergodicity [Map13].

1.2 Outline of the paper

The remainder of the paper proceeds as follows. In §2 we give an overview of the relevant representation theoretic background and outline the construction of Cayley graphs and how the adjacency operator acts through representation theory. In §3 we describe the random bases we use throughout the paper. In §4 we give a deterministic bound on the quantities

$$\left| \mu_\varphi[f] - \frac{\sum_{g \in G} f(g)}{|G|} \right|$$

featuring in the main results. In §5 we give first some basic large deviations estimates for sums of independent random variables, and then apply these to obtain concentration results for tensor products of random matrices from the classical compact groups. Finally, in §6 we prove Theorem 1.8 by combining the deterministic error estimate and our random matrix results.

2 Background

2.1 Representation theory of finite groups

We begin by outlining basic concepts in representation theory. A more complete background can be found in [FH91].

Let G denote a finite group. We consider unitary representations of G . These are pairs (π, V) where V is a finite-dimensional complex Hilbert space and $\pi : G \rightarrow GL(V)$ is a homomorphism such that $\pi(g)$ is unitary for each $g \in G$. When clear, we will just refer to π or V as a representation. We will denote the trivial representation of G by $(\text{triv}, \mathbb{C})$, where \mathbb{C} has the standard inner product and $\text{triv}(g)$ is the identity for all $g \in G$.

The group algebra $\mathbb{C}[G]$ is the ring of formal complex linear combinations of elements of G . We identify $\mathbb{C}[G]$ with $\ell^2(G)$ throughout the paper. Any representation (π, V) of G linearly extends to $\pi : \mathbb{C}[G] \rightarrow \text{End}(V)$ making V a $\mathbb{C}[G]$ module.

Recall that a representation (π, V) is irreducible if there are no proper subspaces of V that are invariant under $\pi(g)$ for all $g \in G$. Two representations (π_1, V_1) and (π_2, V_2) are equivalent if there is a unitary isomorphism $T : V_1 \rightarrow V_2$ that intertwines the representations: $T \circ \pi_1(g) = \pi_2(g) \circ T$ for all $g \in G$. We will denote the unitary dual of G by \hat{G} , it is the collection of equivalence classes of irreducible representations of G . We will not make any distinction between an equivalence class in \hat{G} and an element of the equivalence class; hence we will freely write (π, V) , π , $V \in \hat{G}$.

Given a representation (π, V) of G , the dual representation will be denoted by $(\check{\pi}, \check{V})$. Here, \check{V} is the dual space of V equipped with the inner product arising from that of V on the corresponding Riesz representation vectors, and $\check{\pi}$ is defined by $[\check{\pi}(g)\alpha](v) = \alpha(\pi(g^{-1})v)$ for all $g \in G$ and $v \in V$. If (π, V) is irreducible, then so is $(\check{\pi}, \check{V})$.

Given $(\pi, V) \in \hat{G}$, and $v_1, v_2 \in V$, the matrix coefficient

$$\Phi_{v_1, v_2}^V \stackrel{\text{def}}{=} \langle \pi(g)v_2, v_1 \rangle$$

is in $\ell^2(G)$. This extends bilinearly to a map $\Phi^V : \check{V} \otimes V \rightarrow \ell^2(G)$. The inner product on $\ell^2(G)$ is given by

$$\langle f_1, f_2 \rangle \stackrel{\text{def}}{=} \sum_{g \in G} f_1(g) \overline{f_2(g)}.$$

The space $\ell^2(G)$ is a bimodule for $G \times G$ (under left and right multiplication) and the induced map

$$\Phi \stackrel{\text{def}}{=} \bigoplus_{(\pi, V) \in \hat{G}} \frac{\sqrt{\dim V}}{\sqrt{|G|}} \Phi^V : \bigoplus_{(\pi, V) \in \hat{G}} \check{V} \otimes V \rightarrow \ell^2(G) \quad (2.1)$$

is a unitary bimodule isomorphism by the Peter-Weyl theorem. We also have the Plancherel formula

$$\|f\|_2^2 = \frac{1}{|G|} \sum_{(\pi, V) \in \hat{G}} \dim V \|\pi(f)\|_{\text{HS}}^2, \quad (2.2)$$

where $\|\pi(f)\|_{\text{HS}}^2 \stackrel{\text{def}}{=} \text{tr}_V(\pi(f)\pi(f)^*)$.

2.2 Cayley graphs

Let G be a finite group and let $S = \{s_1, s_1^{-1}, \dots, s_d, s_d^{-1}\}$ be a symmetric subset in G such that $|S| = 2d$. The Cayley graph $\text{Cay}_0(G, S)$ is the directed graph with an edge between g and h if $gs = h$ for some $s \in S$. The directed edges of $\text{Cay}_0(G, S)$ have a pairing arising from matching edges arising from $gs = h$ with the edge arising from $g = hs^{-1}$; the quotient by this equivalence relation is the undirected Cayley graph $\text{Cay}(G, S)$, which is a $2d$ -regular graph. The adjacency operator on $\ell^2(G)$ can be written as

$$\mathcal{A}[f](g) = \sum_{i=1}^d \left(f(gs_i) + f(gs_i^{-1}) \right) = \rho(A)[f](g),$$

where ρ is the right regular representation and

$$A \stackrel{\text{def}}{=} \sum_{i=1}^d (s_i + s_i^{-1}) \in \mathbb{C}[G].$$

3 Random basis construction

In this section we will outline the construction of the bases of eigenfunctions for the adjacency operator. The idea is to exploit the decomposition of $\ell^2(G)$ as the direct sum $\bigoplus_{(\pi, V) \in \hat{G}} \check{V} \otimes V$. To obtain a basis of real-valued functions, one must select the basis inside each irreducible representation dependent upon whether the representation is non-self dual, real or quaternionic as we explain below.

3.1 Non self-dual representations

We start with the case that (π, V) is an irreducible representation that is not equivalent to its dual representation $(\check{\pi}, \check{V})$. Due to their non-equivalence, both $\check{V} \otimes V$ and $V \otimes \check{V}$ appear as distinct summands in the decomposition of $\ell^2(G)$ as the direct sum $\bigoplus_{(\theta, W) \in \hat{G}} \check{W} \otimes W$. We will thus seek an orthonormal basis of $(\check{V} \otimes V) \oplus (V \otimes \check{V})$. As before, let $\{v_k^V\}$ be an orthonormal basis of V consisting of eigenvectors of $\pi(A)$. Moreover, let $\{w_j^V\}$ be any orthonormal basis of V . Then the collection

$$\left\{ \frac{1}{\sqrt{2}}(\check{w}_j^V \otimes v_k^V + w_j^V \otimes \check{v}_k^V), \frac{1}{i\sqrt{2}}(\check{w}_j^V \otimes v_k^V - w_j^V \otimes \check{v}_k^V) : j, k = 1, \dots, \dim V \right\}$$

forms an orthonormal basis of $(\check{V} \otimes V) \oplus (V \otimes \check{V})$. Moreover, they correspond to functions in $\ell^2(G)$

$$\begin{aligned} x_{k,j}^V(g) &\stackrel{\text{def}}{=} \frac{\sqrt{\dim V}}{\sqrt{2}\sqrt{|G|}} (\langle \pi(g)v_k^V, w_j^V \rangle + \langle \check{\pi}(g)\check{v}_k^V, \check{w}_j^V \rangle) = \frac{\sqrt{2\dim V}}{\sqrt{|G|}} \operatorname{Re}(\langle \pi(g)v_k^V, w_j^V \rangle), \\ y_{k,j}^V(g) &\stackrel{\text{def}}{=} \frac{\sqrt{\dim V}}{i\sqrt{2}\sqrt{|G|}} (\langle \pi(g)v_k^V, w_j^V \rangle - \langle \check{\pi}(g)\check{v}_k^V, \check{w}_j^V \rangle) = \frac{\sqrt{2\dim V}}{\sqrt{|G|}} \operatorname{Im}(\langle \pi(g)v_k^V, w_j^V \rangle), \end{aligned}$$

which are real-valued functions with unit L^2 -norm that are mutually orthogonal.

To randomize this basis, we randomize the choice of the basis $\{w_j^V\}_j$. We fix an orthonormal basis $\{e_j^V\}_j$ of V and then given a Haar random unitary operator $u \in U(V)$, we set $w_j^V = ue_j^V$ for each $j = 1, \dots, \dim V$.

3.2 Self-dual representations

A complex irreducible representation that is equivalent to its dual has a conjugate-linear intertwining map $J : V \rightarrow V$ such that $J^2 = \pm \operatorname{Id}$. In the case $J^2 = \operatorname{Id}$ the representation is called *real* and in case $J^2 = -\operatorname{Id}$ the representation is called *quaternionic* [FH91]. It is not hard to check using uniqueness (up to scalars) of the π -invariant inner product on V that for all $v, w \in V$

$$\langle v, w \rangle = \langle J(w), J(v) \rangle. \quad (3.1)$$

3.2.1 Real representations.

In this case, J defines a real structure for V . That is, $V = V_J \oplus iV_J$ where $V_J = \{v \in V : J(v) = v\}$ is a real vector space. It follows from (3.1) that $\langle \bullet, \bullet \rangle$ restricts to a real valued symmetric inner product on V_J , and the inner product on V is obtained from this one by extension of scalars from \mathbb{R} to \mathbb{C} .

Since J intertwines with π , for each $g \in G$ we have $\pi(g) : V_J \rightarrow V_J$, and so $\pi(A)$ is a symmetric operator on $(V_J, \langle \bullet, \bullet \rangle)$. Let $\{v_k^V\}$ denote an orthonormal basis of $\pi(A)$ eigenvectors in V_J with respect to the real inner product. By extension of scalars, these also form an orthonormal eigenbasis of $\pi(A)$ acting on V .

Fix an orthonormal basis $\{e_j^V\}$ of V_J . Choosing a Haar random orthogonal matrix $o \in O(V)$ we let $w_j^V \stackrel{\text{def}}{=} oe_j^V$ for each $1 \leq j \leq \dim V$. The corresponding real random basis of $\rho(A)$ eigenvectors in $\ell^2(G)$ is given by

$$\varphi_{kj}^V(g) \stackrel{\text{def}}{=} \frac{\sqrt{\dim V}}{\sqrt{|G|}} \langle \pi(g)v_k^V, w_j^V \rangle.$$

These are the image under the inclusion $\check{V} \otimes V \rightarrow \ell^2(G)$ of the vectors $\check{w}_j^V \otimes v_k^V$ (this makes it clear that they are $\rho(A)$ eigenvectors).

3.2.2 Quaternionic representations.

Next, suppose that (π, V) is a quaternionic representation of G . In this case, (3.1) implies

$$\langle v, J(v) \rangle = \langle J^2(v), J(v) \rangle = -\langle v, J(v) \rangle$$

hence $\langle v, J(v) \rangle = 0$ for any $v \in V$. This implies $\dim V$ is even and since $\pi(A)$ is Hermitian and commutes with J we can find an orthonormal basis of V of eigenvectors of $\pi(A)$ of the form $\{v_k^V, J(v_k^V)\}_{k=1}^{\frac{1}{2}\dim V}$.

Fix an orthonormal basis $\{e_j^V\}$ of V . Choosing a Haar random unitary matrix $u \in u(V)$ we let $w_j^V \stackrel{\text{def}}{=} ue_j^V$ for each $1 \leq j \leq \dim V$. The corresponding real random basis of $\rho(A)$ eigenvectors in $\ell^2(G)$ is given by

$$\begin{aligned} x_{kj}^V(g) &\stackrel{\text{def}}{=} \frac{\sqrt{2\dim V}}{\sqrt{|G|}} \operatorname{Re}(\langle \pi(g)v_k^V, w_j^V \rangle), \\ y_{kj}^V(g) &\stackrel{\text{def}}{=} \frac{\sqrt{2\dim V}}{\sqrt{|G|}} \operatorname{Im}(\langle \pi(g)v_k^V, w_j^V \rangle). \end{aligned}$$

These are the image under the inclusion $\check{V} \otimes V \rightarrow \ell^2(G)$ of the vectors

$$\begin{aligned} x_{kj}^V &\stackrel{\text{def}}{=} \frac{1}{\sqrt{2}}(\check{w}_j^V \otimes v_k^V + \overline{J(w_j^V)} \otimes J(v_k^V)), \\ y_{kj}^V &\stackrel{\text{def}}{=} \frac{1}{i\sqrt{2}}(\check{w}_j^V \otimes v_k^V - \overline{J(w_j^V)} \otimes J(v_k^V)), \end{aligned}$$

and thus clearly they are $\rho(A)$ eigenvectors.

Putting together all of the different cases for the type of the representation π , the random model for the real-valued eigenbasis of $\ell^2(G)$ has underlying topological space

$$X = \prod_{\substack{\{(\pi, V), (\check{\pi}, \check{V})\} \subseteq \hat{G} \\ \pi \text{ non-self-dual pair}}} U(V) \prod_{\substack{(\pi, V) \in \hat{G} \\ \pi \text{ self-dual and quaternionic}}} U(V) \prod_{\substack{(\pi, V) \in \hat{G} \\ \pi \text{ self-dual and real}}} O(V),$$

equipped with the product probability measure

$$\mathbb{P} = \prod_{\substack{\{(\pi, V), (\check{\pi}, \check{V})\} \subseteq \hat{G} \\ \pi \text{ non-self-dual pair}}} \mathbb{P}_{U(V)} \prod_{\substack{(\pi, V) \in \hat{G} \\ \pi \text{ self-dual and quaternionic}}} \mathbb{P}_{U(V)} \prod_{\substack{(\pi, V) \in \hat{G} \\ \pi \text{ self-dual and real}}} \mathbb{P}_{O(V)}, \quad (3.2)$$

where $\mathbb{P}_{U(V)}$ is the Haar probability measure on the unitary operators $U(V)$ of V , and $\mathbb{P}_{O(V)}$ is the Haar probability measure on the orthogonal operators $O(V)$ of V .

4 Deterministic error term for mean zero functions

In this section we will derive an upper bound for

$$\left| \sum_{g \in G} f(g) |\varphi(g)|^2 - \frac{1}{|G|} \sum_{g \in G} f(g) \right|, \quad (4.1)$$

where φ is one of the eigenbasis elements of $\ell^2(G)$ described in the previous section, and f is a real-valued function on the group G . In fact, we will further make the assumption that

$$\sum_{g \in G} f(g) = 0,$$

so that we can instead just bound

$$\left| \sum_{g \in G} f(g) |\varphi(g)|^2 \right|.$$

This can be done without any loss of generality since given a non-zero mean function, we can consider $f - \frac{1}{|G|} \sum_{g \in G} f(g)$ which has zero mean, and then a bound on the above quantity for this zero mean function provides a bound on the desired difference since

$$\begin{aligned} \left| \sum_{g \in G} \left(f(g) - \frac{1}{|G|} \sum_{h \in G} f(h) \right) |\varphi(g)|^2 \right| &= \left| \sum_{g \in G} f(g) |\varphi(g)|^2 - \frac{1}{|G|} \sum_{h \in G} f(h) \sum_{g \in G} |\varphi(g)|^2 \right| \\ &= \left| \sum_{g \in G} f(g) |\varphi(g)|^2 - \frac{1}{|G|} \sum_{g \in G} f(g) \right|, \end{aligned}$$

as the eigenfunction φ is normalized with respect to the counting measure. The bounds we will obtain later will involve $\|f\|_{\ell^2}$, but since the mean of f is just the Fourier component of f corresponding to the constant eigenfunction, we have $\|f - \frac{1}{|G|} \sum_{g \in G} f(g)\|_{\ell^2} \leq \|f\|_{\ell^2}$ and so any bounds depending on the ℓ^2 -norm of the zero mean function can just be bounded by the ℓ^2 -norm of the function itself.

Now, recall that there were three types of functions in the eigenbasis dependent upon the type of irreducible representation that they come from. In the case of complex irreducible representations that are not real we have the following two types given by real and imaginary parts of matrix coefficients

Type 1 - Real Part

$$\varphi(g) = \frac{\sqrt{\dim V}}{\sqrt{2}\sqrt{|G|}} (\langle \pi(g)v, w \rangle + \langle \check{\pi}(g)\check{v}, \check{w} \rangle),$$

Type 2 - Imaginary Part

$$\varphi(g) = \frac{\sqrt{\dim V}}{i\sqrt{2}\sqrt{|G|}} (\langle \pi(g)v, w \rangle - \langle \check{\pi}(g)\check{v}, \check{w} \rangle).$$

In the case of a complex irreducible representation that is real we have the following type of basis element

Type 3 - Real Matrix Coefficient

$$\varphi(g) = \frac{\sqrt{\dim V}}{\sqrt{|G|}} \langle \pi(g)v, w \rangle.$$

In each of the above types, (π, V) is an irreducible unitary representation and $v, w \in V$ are unit vectors.

We will show the following.

Proposition 4.1. *Let $\varphi : G \rightarrow \mathbb{R}$ be one of types 1, 2 or 3, and let $f : G \rightarrow \mathbb{R}$ have zero*

mean. If φ is of type 1 or type 2, then

$$\left| \sum_{g \in G} f(g) |\varphi(g)|^2 \right| \leq \left| \operatorname{Re} \left(\left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \check{\pi})(g) (v \otimes \check{v}), w \otimes \check{w} \right\rangle \right) \right| \\ + \left| \operatorname{Re} \left(\left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \pi)(g) (v \otimes v), w \otimes w \right\rangle \right) \right|,$$

and if φ is of type 3, then

$$\left| \sum_{g \in G} f(g) |\varphi(g)|^2 \right| \leq \left| \operatorname{Re} \left(\left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \check{\pi})(g) (v \otimes \check{v}), w \otimes \check{w} \right\rangle \right) \right|.$$

Proof. Suppose that φ is type 1. Then,

$$\begin{aligned} |\varphi(g)|^2 &= \frac{1}{2} \frac{\dim V}{|G|} (\langle (\pi \otimes \pi)(g) (v \otimes v), w \otimes w \rangle + \langle (\check{\pi} \otimes \check{\pi})(g) (\check{v} \otimes \check{v}), \check{w} \otimes \check{w} \rangle \\ &\quad + \langle (\check{\pi} \otimes \pi)(g) (\check{v} \otimes v), \check{w} \otimes w \rangle + \langle (\pi \otimes \check{\pi})(g) (v \otimes \check{v}), w \otimes \check{w} \rangle) \\ &= \frac{\dim V}{|G|} (\operatorname{Re} (\langle (\pi \otimes \pi)(g) (v \otimes v), w \otimes w \rangle) + \operatorname{Re} (\langle (\pi \otimes \check{\pi})(g) (v \otimes \check{v}), w \otimes \check{w} \rangle)). \end{aligned} \tag{4.2}$$

The result is then an immediate application of the triangle inequality using the fact that f is real-valued. The proof for type 2 functions is essentially the same, and the proof for type 3 is even simpler (one only needs to deal with $\pi \otimes \check{\pi}$ terms). \square

5 Probabilistic ingredients

In this section, we outline some results that we will use in §6 when bounding the probability that our random bases have the properties of Theorems 1.1 and 1.8.

5.1 Large deviations estimates

We begin by recalling that the χ -squared distribution with k -degrees of freedom, denoted by χ_k^2 , has probability density function

$$f_k(x) = \frac{x^{\frac{k}{2}-1} e^{-\frac{x}{2}}}{2^{\frac{k}{2}} \Gamma\left(\frac{k}{2}\right)} \mathbf{1}_{\{x>0\}}. \tag{5.1}$$

If Z_1, \dots, Z_k are independent standard normal random variables, then

$$\sum_{i=1}^k Z_i^2 \sim \chi_k^2.$$

In this article we will use the following results regarding independent χ_1^2 and χ_2^2 random variables.

Lemma 5.1. *If X_1, \dots, X_N are independent χ_1^2 -distributed random variables, then*

$$\mathbb{P}\left(\sum_{i=1}^N X_i \leq \frac{N}{2}\right) \leq e^{-\frac{N}{12}}.$$

Proof. By exponential Chebyshev, for any $A > 0$

$$\begin{aligned} \mathbb{P}\left(\sum_{i=1}^N X_i \leq t\right) &\leq e^{At} \mathbb{E}\left[e^{-A \sum_{i=1}^N X_i}\right] = e^{At} \prod_{i=1}^N \mathbb{E}\left[e^{-AX_i}\right] \\ &= e^{At} \prod_{i=1}^N \frac{1}{\sqrt{2\pi}} \int_0^\infty x^{-\frac{1}{2}} e^{-x(\frac{1}{2}+A)} dx = e^{At} \left(\frac{1}{\sqrt{1+2A}}\right)^N. \end{aligned}$$

Taking $t = \frac{N}{2}$ and $A = \frac{1}{2}$ (so that $A - \log(1+2A) \leq -\frac{1}{6}$) we obtain the stated result. \square

Lemma 5.2. *Suppose that $(a_1, \dots, a_N) \in \mathbb{R}^N$ and there exist constants $A, C > 0$ such that*

1. $\sum_{i=1}^N a_i = 0$,
2. $\sum_{i=1}^N a_i^2 \leq C$, and
3. $|a_i| \leq A$ for each $1 \leq i \leq N$.

Then,

(i) *If X_1, \dots, X_N are independent χ_1^2 -distributed random variables then for all $t > 0$,*

$$\mathbb{P}\left(\left|\sum_{i=1}^N a_i X_i\right| \geq t\right) \leq 2 \left(\frac{At}{C} + 1\right)^{\frac{C}{2A^2}} e^{-\frac{t}{2A}}.$$

(ii) *If X_1, \dots, X_N are independent χ_2^2 -distributed random variables then for all $t > 0$,*

$$\mathbb{P}\left(\left|\sum_{i=1}^N a_i X_i\right| \geq t\right) \leq 2 \left(\frac{At}{2C} + 1\right)^{\frac{C}{A^2}} e^{-\frac{t}{2A}}.$$

Remark 5.3. Note that condition (3) in Lemma 5.2 immediately follows from condition (2) since we must have $|a_i| \leq \sqrt{C}$ for all $1 \leq i \leq N$. Likewise, condition (2) follows from condition (3) with $C = A^2 N$.

Proof. We start with (i). Notice that

$$\mathbb{P}\left(\left|\sum_{i=1}^N a_i X_i\right| \geq t\right) = \mathbb{P}\left(\sum_{i=1}^N a_i X_i \geq t\right) + \mathbb{P}\left(-\sum_{i=1}^N a_i X_i \geq t\right).$$

Now for any $\varepsilon \in [0, \frac{1}{2A})$, exponential Chebyshev inequality along with independence of the

X_i and the formula (5.1) implies that

$$\begin{aligned}
& \mathbb{P}\left(\sum_{i=1}^N a_i X_i \geq t\right) \\
& \leq e^{-t\varepsilon} \mathbb{E}\left(\exp\left(\varepsilon \sum_{i=1}^N a_i X_i\right)\right) = e^{-t\varepsilon} \prod_{i=1}^N \mathbb{E} \exp(\varepsilon a_i X_i) \\
& = e^{-t\varepsilon} \prod_{i=1}^N \frac{1}{\sqrt{2\pi}} \int_0^\infty x^{-\frac{1}{2}} e^{-\frac{x}{2}(1-2\varepsilon a_i)} dx = e^{-t\varepsilon} \prod_{i=1}^N \frac{1}{\sqrt{1-2\varepsilon a_i}} \\
& = e^{-t\varepsilon} \exp\left(-\frac{1}{2} \sum_{i=1}^N \log(1-2\varepsilon a_i)\right) = e^{-t\varepsilon} \exp\left(\varepsilon \sum_{i=1}^N a_i + \frac{1}{2A^2} \sum_{i=1}^N a_i^2 \sum_{n=2}^\infty \frac{(2A\varepsilon)^n \left(\frac{a_i}{A}\right)^{n-2}}{n}\right).
\end{aligned}$$

The final equality follows from $|2\varepsilon a_i| < 1$. Now by assumption (3), we have $|\frac{a_i}{A}|^{n-2} \leq 1$ and so using assumptions (1) and (2) we have

$$\begin{aligned}
\mathbb{P}\left(\sum_{i=1}^N a_i X_i \geq t\right) & \leq e^{-t\varepsilon} \exp\left(\frac{C}{2A^2} \sum_{n=2}^\infty \frac{(2A\varepsilon)^n}{n}\right) \\
& = e^{-t\varepsilon} \exp\left(\log\left((1-2A\varepsilon)^{-\frac{C}{2A^2}}\right) - \frac{C\varepsilon}{A}\right) \\
& = \frac{e^{-\varepsilon(t+\frac{C\varepsilon}{A})}}{(1-2A\varepsilon)^{\frac{C}{2A^2}}},
\end{aligned}$$

the second equality following from the fact that $|2A\varepsilon| < 1$. We now choose $\varepsilon \in [0, \frac{1}{2A})$ that minimizes this upper bound. This can readily be seen to be given by

$$\varepsilon = \frac{1}{2A} - \frac{\frac{C}{2A^2}}{t + \frac{C}{A}} \in \left[0, \frac{1}{2A}\right).$$

We hence obtain the upper bound

$$\mathbb{P}\left(\sum_{i=1}^N a_i X_i \geq t\right) \leq \left(\frac{At}{C} + 1\right)^{\frac{C}{2A^2}} e^{-\frac{t}{2A}}.$$

The same bound applies to $\mathbb{P}\left(-\sum_{i=1}^N a_i X_i \geq t\right)$ since we may set $b_i = -a_i$ and then $(b_1, \dots, b_N) \in \mathbb{R}^N$ satisfies assumptions (1), (2) and (3) so that the above computations still hold.

The proof of (ii) follows identically but using the probability density function $f_2(x)$ rather than $f_1(x)$. \square

5.2 Random matrix estimates

Lemma 5.4. *Suppose that V is an n -dimensional complex Hermitian inner product space with orthonormal basis $\{e_i\}_{i=1}^n$, and u is a Haar random unitary matrix in $U(V)$. Then,*

1. *For any fixed vector $\beta = \sum_{1 \leq i, j \leq n} \beta_{ij} e_i \otimes \check{e}_j \in V \otimes \check{V}$ with $\beta_{ij} \in \mathbb{C}$, $\sum_{1 \leq i, j \leq n} |\beta_{ij}|^2 \leq C$ and $\sum_{i=1}^n \beta_{ii} = 0$, for any $1 \leq k \leq n$, and any $T > 0$ we have*

$$\mathbb{P}_{u \in U(V)}(|\langle \beta, u e_k \otimes \check{u} \check{e}_k \rangle| \geq T) \leq 6e^{-\frac{nT}{32\sqrt{C}}} + 2e^{-\frac{n}{6}}.$$

2. For any fixed vector $\alpha = \sum_{1 \leq i, j \leq n} \alpha_{ij} e_i \otimes e_j \in V \otimes V$ with $\alpha_{ij} \in \mathbb{C}$ and $\sum_{1 \leq i, j \leq n} |\alpha_{ij}|^2 \leq C$, for any $1 \leq k \leq n$, and any $T > 0$ we have

$$\mathbb{P}_{u \in U(V)} (|\operatorname{Re} \langle \alpha, u e_k \otimes u e_k \rangle| \geq T) \leq 6e^{-\frac{nT}{32\sqrt{C}}} + 2e^{-\frac{n}{6}}.$$

Proof. Proof of Part 1. We have $|\langle \beta, u e_k \otimes \widetilde{u e_k} \rangle| = |\langle u^{-1} M u e_k, e_k \rangle|$ where $M \in \operatorname{End}(V)$ is the operator defined by $M(e_j) = \sum_i \beta_{ij} e_i$. The conditions on β imply that M has zero trace and Hilbert-Schmidt norm bounded by \sqrt{C} .

Write $M = H_1 + iH_2$ where $H_1 \stackrel{\text{def}}{=} \frac{1}{2}(M + M^*)$ and $H_2 \stackrel{\text{def}}{=} \frac{1}{2i}(M - M^*)$ are Hermitian operators. We have $\|H_1\|_{\text{HS}}^2 + \|H_2\|_{\text{HS}}^2 = \|M\|_{\text{HS}}^2 \leq C$ and hence

$$\operatorname{tr}(H_1) = \operatorname{tr}(H_2) = 0, \quad \|H_1\|_{\text{HS}}^2, \|H_2\|_{\text{HS}}^2 \leq C.$$

Also,

$$\mathbb{P}(|\langle \beta, u e_k \otimes \widetilde{u e_k} \rangle| \geq T) \leq \sum_{i=1,2} \mathbb{P}\left(|\langle u^{-1} H_i u e_k, e_k \rangle| \geq \frac{T}{2}\right). \quad (5.2)$$

Since each H_i is Hermitian, it is conjugate to a real diagonal matrix D_i with the same Hilbert-Schmidt norm and trace zero by a unitary operator, and by bi-invariance of Haar measure, we obtain

$$\mathbb{P}\left(|\langle u^{-1} H_i u e_k, e_k \rangle| \geq \frac{T}{2}\right) \leq \mathbb{P}\left(|\langle u^{-1} D_i u e_1, e_1 \rangle| \geq \frac{T}{2}\right).$$

We treat only D_1 as the bound for D_2 is the same. Thus we can assume that $H_1 = D_1 = \operatorname{diag}(\lambda_1, \dots, \lambda_{\dim V})$ with

$$\sum_i \lambda_i = 0, \quad \sum_i |\lambda_i|^2 \leq C,$$

and we have

$$|\langle u^{-1} D_i u e_1, e_1 \rangle| = \sum_i \lambda_i |u_{i1}|^2.$$

As is well-known⁵ the entries $u_{i1} = \frac{1}{\sqrt{N}} \eta_i$ where η_i are independent standard complex normal random variables and

$$N \stackrel{\text{def}}{=} \sum_{i=1}^n |\eta_i|^2 = \frac{1}{2} \sum_{i=1}^{2n} Y_i$$

where Y_i are independent χ_1^2 random variables. Hence by Lemma 5.1

$$\mathbb{P}\left(N \leq \frac{n}{2}\right) \leq e^{-\frac{n}{6}}. \quad (5.3)$$

Thus with probability at least $1 - e^{-\frac{n}{6}}$, we have $N \geq \frac{n}{2}$. We have

$$|\langle u^{-1} D_i u e_1, e_1 \rangle| = \frac{1}{2N} \left| \sum_{i=1}^n \lambda_i X_i \right|$$

⁵A Haar random unitary matrix can be obtained by considering a random matrix whose entries are i.i.d. standard complex normal random variables, and then making the columns orthonormal by a Gram-Schmidt procedure on the columns (see for example §§1.2 of [Mec19]). Carrying out this algorithm starting with the 1st column just normalizes the column.

where X_i are independent χ_2^2 distributed random variables, and so by Lemma 5.2 Part (ii) with $C = C$ and $A = \sqrt{C}$

$$\mathbb{P}\left(\left|\sum_{i=1}^n \lambda_i X_i\right| \geq \frac{nT}{4}\right) \leq \left(2 + \frac{nT}{4\sqrt{C}}\right) e^{-\frac{nT}{8\sqrt{C}}}. \quad (5.4)$$

Combining (5.3) and (5.4) then gives

$$\mathbb{P}\left(|\langle u^{-1} D_i u e_1, e_1 \rangle| \geq \frac{T}{2}\right) \leq \left(2 + \frac{nT}{4\sqrt{C}}\right) e^{-\frac{nT}{8\sqrt{C}}} + e^{-\frac{n}{6}} \leq 3e^{-\frac{nT}{32\sqrt{C}}} + e^{-\frac{n}{6}}.$$

Part 1 then follows from (5.2).

Proof of Part 2. This is similar except here we let $M \in \text{End}(V)$ be the operator defined by $M(e_j) = \sum_i \alpha_{ij} e_i$ and write $M = S + R$ with $S \stackrel{\text{def}}{=} \frac{1}{2}(M + M^T)$ and $R \stackrel{\text{def}}{=} \frac{1}{2}(M - M^T)$ where transpose is defined with respect to the real inner product $\text{Re}\langle \bullet, \bullet \rangle$. We have $u^T R u = 0$ so R makes no contribution to $\langle \alpha, u e_k \otimes u e_k \rangle$.

The rest of the proof follows analogous lines to the proof of part 1, diagonalizing the real and imaginary parts of S by orthogonal (unitary) matrices. This leads to bounding $\mathbb{P}\left(|\text{Re}(\sum_{i=1}^n \lambda_i \eta_i^2)| \geq \frac{nT}{4}\right)$ and $\mathbb{P}\left(|\text{Im}(\sum_{i=1}^n \lambda'_i \eta_i^2)| \geq \frac{nT}{4}\right)$ where $\lambda_i, \lambda'_i \in \mathbb{R}$, $\sum \lambda_i^2, \sum (\lambda'_i)^2 \leq C$ and η_i , $1 \leq i \leq n$ are independent standard complex normals. For the first we have

$$\begin{aligned} \mathbb{P}\left(\left|\text{Re}\left(\sum_{i=1}^n \lambda_i \eta_i^2\right)\right| \geq \frac{nT}{4}\right) &= \mathbb{P}\left(\left|\sum_{i=1}^n \lambda_i (x_i^2 - y_i^2)\right| \geq \frac{nT}{4}\right) \\ &= \mathbb{P}\left(\left|\sum_{i=1}^n \lambda_i (X_i - Y_i)\right| \geq \frac{nT}{2}\right) \end{aligned}$$

where this time, X_i and Y_i are independent χ_1^2 distributed random variables. One can apply Lemma 5.2 Part (i) with

$$a_i \stackrel{\text{def}}{=} \begin{cases} \lambda_i & \text{if } i = 1, \dots, n, \\ -\lambda_{i-n} & \text{if } i = n+1, \dots, 2n \end{cases}$$

to obtain

$$\mathbb{P}\left(\left|\text{Re}\left(\sum_{i=1}^n \lambda_i \eta_i^2\right)\right| \geq \frac{nT}{4}\right) \leq \left(2 + \frac{nT}{2\sqrt{C}}\right) e^{-\frac{nT}{4\sqrt{C}}}.$$

Dealing with $\mathbb{P}\left(|\text{Im}(\sum_{i=1}^n \lambda'_i \eta_i^2)| \geq \frac{nT}{4}\right)$ is similar. These lead to the stated result. \square

Lemma 5.5. *Suppose that V is a real inner product space with $n \stackrel{\text{def}}{=} \dim V$. Then for any fixed vector $\beta = \sum_{1 \leq i, j \leq n} \beta_{ij} e_i \otimes \check{e}_j \in V \otimes \check{V}$, $\sum_{1 \leq i, j \leq n} |\beta_{ij}|^2 \leq C$ and $\sum_{i=1}^n \beta_{ii} = 0$, for any $1 \leq k \leq n$, and any $T > 0$ we have*

$$\mathbb{P}_{o \in O(V)}(|\langle \beta, o e_k \otimes \widetilde{o e_k} \rangle| \geq T) \leq 6e^{-\frac{nT}{32\sqrt{C}}} + 2e^{-\frac{n}{12}}.$$

Proof. This is just the real version of Lemma 5.4 Part 1. The proof is along exactly the same lines, using that the first column of an orthogonal random matrix is obtained by choosing independent standard real normal random variables as the entries, and then normalizing. Accordingly, one ends up using Lemma 5.2 Part (ii). \square

6 Proof of main results

Let $(\pi, V) \in \hat{G}$ be an irreducible representation of G , $1 \leq j, k \leq n \stackrel{\text{def}}{=} \dim V$, and $f : G \rightarrow \mathbb{R}$ have zero mean. The randomness of the basis enters into the error term given in Proposition 4.1 via the quantities

$$\left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g)(\pi \otimes \check{\pi})(g)(v_k^V \otimes \check{v}_k^V), w_j^V \otimes \check{w}_j^V \right\rangle, \\ \left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g)(\pi \otimes \pi)(g)(v_k^V \otimes v_k^V), w_j^V \otimes w_j^V \right\rangle.$$

Accordingly, let $v \stackrel{\text{def}}{=} v_k^V$, $e_i \stackrel{\text{def}}{=} e_i^V$ and

$$x \stackrel{\text{def}}{=} \frac{\dim V}{|G|} \sum_{g \in G} f(g)(\pi \otimes \check{\pi})(g)(v \otimes \check{v}), \quad y \stackrel{\text{def}}{=} \frac{\dim V}{|G|} \sum_{g \in G} f(g)(\pi \otimes \pi)(g)(v \otimes v). \quad (6.1)$$

We write

$$x \stackrel{\text{def}}{=} \sum_{i,j=1}^n x_{ij} e_i \otimes \check{e}_j, \quad y \stackrel{\text{def}}{=} \sum_{i,j=1}^n y_{ij} e_i \otimes e_j \quad (6.2)$$

for some $x_{ij}, y_{ij} \in \mathbb{C}$. The vectors x and y satisfy the following properties.

Lemma 6.1. *Let x be defined as in (6.1) and (6.2). Then,*

$$(i) \sum_{i,j=1}^n |x_{ij}|^2, \sum_{i,j=1}^n |y_{ij}|^2 \leq \frac{\|f\|_{\ell^2}^2 \dim V}{|G|}, \text{ and}$$

$$(ii) \sum_{i=1}^n x_{ii} = 0.$$

Proof. Using (6.2), we see that $\sum_{i,j} |x_{ij}|^2 = \langle x, x \rangle$, and so computing this inner product with the expression (6.1), we obtain

$$\begin{aligned} \sum_{i,j} |x_{ij}|^2 &= \frac{(\dim V)^2}{|G|^2} \sum_g \sum_h f(g)f(h) \langle (\pi \otimes \check{\pi})(g)(v \otimes \check{v}), (\pi \otimes \check{\pi})(h)(v \otimes \check{v}) \rangle \\ &= \frac{(\dim V)^2}{|G|^2} \sum_g \sum_h f(g)f(h) |\langle \pi(g)v, \pi(h)v \rangle|^2 \\ &\leq \frac{(\dim V)^2}{|G|^2} \sum_g \sum_h \frac{|f(g)|^2 + |f(h)|^2}{2} |\langle \pi(g)v, \pi(h)v \rangle|^2 \\ &= \frac{(\dim V)^2}{|G|} \sum_g |f(g)|^2 \frac{1}{|G|} \sum_h |\langle \pi(g)v, \pi(h)v \rangle|^2 \\ &= \frac{\dim V}{|G|} \|f\|_{\ell^2}^2, \end{aligned}$$

with the last equality following from Schur orthogonality. The same bound holds for y since

$$\sum_{i,j} |y_{ij}|^2 = \frac{(\dim V)^2}{|G|^2} \sum_g \sum_h f(g)f(h) (\langle \pi(g)v, \pi(h)v \rangle)^2.$$

To prove (ii), we see from (6.2) that $\sum_i x_{ii} = \langle x, \sum_i e_i \otimes \check{e}_i \rangle$. Computing this inner product

with (6.1) we obtain

$$\begin{aligned}
\sum_i x_{ii} &= \frac{\dim V}{|G|} \sum_g f(g) \sum_i \langle \pi(g)v, e_i \rangle \langle \check{\pi}(g)\check{v}, \check{e}_i \rangle \\
&= \frac{\dim V}{|G|} \sum_g f(g) \left\langle \pi(g)v, \sum_i \langle \pi(g)v, e_i \rangle e_i \right\rangle \\
&= \frac{\dim V}{|G|} \sum_g f(g) \langle \pi(g)v, \pi(g)v \rangle \\
&= \frac{\dim V}{|G|} \sum_g f(g) = 0,
\end{aligned}$$

since f has mean zero. \square

The following bound applies to the error terms that arise from random basis elements coming from complex non-self-dual or quaternionic representations (type 1 or type 2 in the previous language) in Proposition 4.1.

Proposition 6.2. *Let $(\pi, V) \in \hat{G}$ be an irreducible representation of G that is either complex non-self-dual or quaternionic. Then, for any $t > 0$ and indices $1 \leq i, j \leq n \stackrel{\text{def}}{=} \dim V$,*

$$\begin{aligned}
\mathbb{P}_{u \in U(V)} \left(\left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \check{\pi})(g) (v_i^V \otimes \check{v}_i^V), u e_j^V \otimes \widetilde{u e_j^V} \right\rangle \right| \geq t \frac{\|f\|_{\ell^2}}{2\sqrt{|G|}} \right) \\
\leq 6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{6}},
\end{aligned}$$

and

$$\begin{aligned}
\mathbb{P}_{u \in U(V)} \left(\left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \pi)(g) (v_i^V \otimes v_i^V), u e_j^V \otimes u e_j^V \right\rangle \right| \geq t \frac{\|f\|_{\ell^2}}{2\sqrt{|G|}} \right) \\
\leq 6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{6}}.
\end{aligned}$$

Proof. This follows by combining the respective parts of Lemma 5.4 and Lemma 6.1, with $C = \frac{\|f\|_{\ell^2}^2 \dim V}{|G|}$ and $T = t \frac{\|f\|_{\ell^2}}{2\sqrt{|G|}}$. \square

The next bound applies to the other error terms coming from real representations.

Proposition 6.3. *Let $(\pi, V) \in \hat{G}$ be a self-dual real irreducible representation of G and $1 \leq i, j \leq n \stackrel{\text{def}}{=} \dim V$, then for any $t > 0$,*

$$\begin{aligned}
\mathbb{P}_{o \in O(V_J)} \left(\left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \check{\pi})(g) (v_i^V \otimes \check{v}_i^V), o e_j^V \otimes \widetilde{o e_j^V} \right\rangle \right| \geq t \frac{\|f\|_{\ell^2}}{2\sqrt{|G|}} \right) \\
\leq 6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}}.
\end{aligned}$$

Proof. Let

$$x \stackrel{\text{def}}{=} \frac{\dim V}{|G|} \sum_{g \in G} f(g) (\pi \otimes \check{\pi})(g) (v_i^V \otimes \check{v}_i^V).$$

Expanding x over the basis $\{e_i^V \otimes \check{e}_j^V\}_{i,j}$ of $V \otimes \check{V}$ we obtain

$$x = \sum_{i,j=1}^n x_{ij} e_i^V \otimes \check{e}_j^V,$$

for some $x_{ij} \in \mathbb{C}$. Because in this case, the inner product is extended from a real inner product on the real subspace V_J (cf. §§3.2.1), and all $oe_j \in V_J$, we have

$$\operatorname{Re} \left\langle x, oe_j^V \otimes \widetilde{oe_j^V} \right\rangle = \langle \beta, oe_j^V \otimes \widetilde{oe_j^V} \rangle$$

where $\beta = \sum_{i,j=1}^n \beta_{ij} e_i^V \otimes \check{e}_j^V$, $\beta_{ij} \stackrel{\text{def}}{=} \operatorname{Re}(x_{ij})$. We thus have $\sum_{ij} |\beta_{ij}|^2 \leq \sum_{ij} |x_{ij}|^2 \leq \frac{\|f\|_2^2 \dim V}{|G|}$ and $\sum_i \beta_{ii} = 0$ using Lemma 6.1. We can apply Lemma 5.5 to get the result. \square

We are now ready to combine the probabilistic estimates of Propositions 6.2 and 6.3 with the deterministic error estimate of Proposition 4.1 to prove Theorem 1.8.

Proof of Theorem 1.8. Recall the probability space (X, \mathbb{P}) and the notation used for the elements of the random eigenbasis constructed in §3. For each $k = 1, \dots, M$, we define $\tilde{f}_k = f_k - \frac{1}{|G|} \sum_{g \in G} f_k(g)$ and set

$$\begin{aligned} F_1(\pi, i, j, f_k) &\stackrel{\text{def}}{=} \left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} \tilde{f}_k(g) (\pi \otimes \pi)(g) (v_i^V \otimes v_j^V), ue_j^V \otimes ue_j^V \right\rangle \right|, \\ F_2(\pi, i, j, f_k) &\stackrel{\text{def}}{=} \left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} \tilde{f}_k(g) (\pi \otimes \check{\pi})(g) (v_i^V \otimes \check{v}_j^V), ue_j^V \otimes \widetilde{ue_j^V} \right\rangle \right|, \\ F_3(\pi, i, j, f_k) &\stackrel{\text{def}}{=} \left| \operatorname{Re} \left\langle \frac{\dim V}{|G|} \sum_{g \in G} \tilde{f}_k(g) (\pi \otimes \check{\pi})(g) (v_i^V \otimes \check{v}_j^V), oe_j^V \otimes \widetilde{oe_j^V} \right\rangle \right|, \end{aligned}$$

where in F_1 and F_2 we assume V is not real and in F_3 we assume that V is a real representation. In all cases, we may assume that π is non-trivial since this is a one dimensional representation with corresponding eigenspace spanned by the constant function for which the desired estimates trivially hold.

Let \mathcal{E}_t denote the event that some $F_1(\pi, i, j, f_k)$ or $F_2(\pi, i, j, f_k)$ with (π, V) complex or quaternionic, or some $F_3(\pi, i, j, f_k)$ with (π, V) real satisfies

$$F_\ell(\pi, i, j, f_k) > t \frac{\|\tilde{f}_k\|_{\ell^2}}{2\sqrt{|G|}}.$$

By carrying out a union bound over all $\pi \in \hat{G} - \text{triv}$, all functions f_1, \dots, f_M in the collection, and $1 \leq i, j \leq \dim V$ with the estimates from Propositions 6.2 and 6.3, we obtain

$$\mathbb{P}(\mathcal{E}_t) \leq 2M \sum_{(\pi, V) \in \hat{G} - \text{triv}} (\dim V)^2 \left(6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right) \quad (6.3)$$

Now assume we have a basis $\mathcal{B} \subset \ell^2(V(G))$ that is not in \mathcal{E}_t , and let $\varphi \in \mathcal{B}$. Then, for any

of the functions f_k , since $\|\tilde{f}_k\|_2 \leq \|f\|_2$ we obtain from Proposition 4.1 that

$$\left| \sum_{g \in G} f_k(g) |\varphi(g)|^2 - \frac{1}{|G|} \sum_{g \in G} f_k(g) \right| \leq t \frac{\|f_k\|_{\ell^2}}{\sqrt{|G|}}.$$

This completes the proof. \square

The proof of Theorem 1.1 is then immediate.

Remark 6.4. The proofs of Theorems 1.1 and 1.8 simplify if one only wishes to consider complex-valued eigenbases. The construction of these bases is similar to §3. Indeed, given an irreducible representation (π, V) of G , let $\{v_i^V\}$ be an orthonormal basis of V consisting of $\pi(A)$ eigenvectors and let $\{w_j^V\}$ be any other orthonormal basis of V . The collection

$$\{\check{w}_j^V \otimes v_i^V : i, j = 1, \dots, \dim V\}$$

forms an orthonormal basis of $\check{V} \otimes V$ corresponding to the following orthonormal adjacency operator eigenfunctions

$$\varphi_{i,j}^V(g) \stackrel{\text{def}}{=} \frac{\sqrt{\dim V}}{\sqrt{|G|}} \langle \pi(g) v_i^V, w_j^V \rangle$$

in $\ell^2(G)$. To randomize this basis, we fix an orthonormal basis $\{e_j^V\}_j$ of V and then given a Haar random unitary operator $u \in U(V)$, we set $w_j^V = u e_j^V$ for each $j = 1, \dots, \dim V$.

The upper bound obtained in Proposition 4.1 for type 3 basis elements then holds for the collection $\{\varphi_{i,j}^V\}$ but with the real part in the upper bound replaced by the absolute value; the proof of this is analogous. Expanding the vectors in the inner product for this upper bound as in §6, we recover Lemma 6.1 identically. Thus, we can combine Lemma 6.1 and part 1 of Lemma 5.4 to prove the same probabilistic bound in the first part of Proposition 6.2 (without the real part) for the $\varphi_{i,j}^V$. Theorems 1.1 and 1.8 then follow via a union bound over the irreducible representations and basis vectors as in the proof of Theorem 1.8. In fact, in the complex-valued basis case, one may take the functions to be complex-valued.

Proof of Corollary 1.3. We use Theorem 1.1 with $t = 64(\varepsilon + 1) \frac{\log(|G|)}{\sqrt{\mathfrak{D}(G)}}$. Then,

$$2M \sum_{(\pi, V) \in \hat{G} - \text{triv}} (\dim V)^2 \left(6e^{-\frac{t\sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right) < 12M|G|e^{-(\varepsilon+1)\log(|G|)} + 4M|G|e^{-\frac{\mathfrak{D}(G)}{12}},$$

and so requiring that both terms in this summation are less than $\frac{1}{2}$ gives the required bound on M for a basis satisfying (1.4) to exist. \square

Proof of Corollary 1.6. Since the collection of subsets A_i satisfy the bound $c|G|^{1-\eta} \leq |A_i| \leq C|G|^{1-\eta}$ on their size, there are at most $\frac{1}{c}|G|^\eta$ of them. We take $t = \frac{128 \log |G|}{\sqrt{\mathfrak{D}(G)}}$ so that when $|G|$ is sufficiently large (dependent only upon c and η), we have $12\frac{1}{c}|G|^{\eta-1} + 4\frac{1}{c}|G|^{\eta+1}e^{-\frac{1}{12}|G|^{\eta+\varepsilon}} < 1$. Thus by Theorem 1.1 if one takes the functions to be the at most $\frac{1}{c}|G|^\eta$ indicator functions on the sets A_i , there exists an orthonormal eigenbasis \mathcal{B} such that

$$\left| \mu_\varphi[A_i] - \frac{|A_i|}{|G|} \right| \leq \frac{128 \log |G|}{|G|^{\frac{1}{2}\eta + \frac{1}{2}\varepsilon}} \frac{\sqrt{|A_i|}}{\sqrt{|G|}} \leq \frac{128 \log |G|}{\sqrt{c}|G|^{\frac{1}{2}\varepsilon}} \frac{|A_i|}{|G|},$$

for every $\varphi \in \mathcal{B}$ and each set A_i , with the last inequality following from $\sqrt{|A_i|} \leq \frac{|A_i|}{\sqrt{|A_i|}} \leq \frac{|A_i|}{\sqrt{c|G|^{\frac{1}{2}-\frac{1}{2}\eta}}}$. \square

Proof of Proposition 1.7. For $\text{Sym}(n)$, we firstly note that the sign representation is one-dimensional and the corresponding eigenfunctions are spanned by the function assigning 1 to even permutations and -1 to odd permutations and thus these eigenfunctions already satisfy the QUE bound exactly after normalization. When doing the randomization in the proof of Theorem 1.8, we thus only require the union bound to run over the non-sign and non-trivial permutations. In other words, for $\text{Sym}(n)$, Theorem 1.1 holds when

$$2M_n \sum_{(\pi, V) \in \widehat{\text{Sym}(n)} - \{\text{triv}, \text{sign}\}} (\dim V)^2 \left(6e^{-\frac{t_n \sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right) < 1,$$

instead. Now, consider $t_n = 192 \frac{\log(n-1)}{\sqrt{n-1}}$. Since $\dim V \geq n-1$ for all non-trivial and non-sign irreducible representations (π, V) we have that $e^{-\frac{t_n \sqrt{\dim V}}{64}} \leq e^{-3 \log(\dim V)}$. Moreover, for $n \geq 24$ and (π, V) non-sign and non-trivial we have $(\dim V)^2 e^{-\frac{\dim V}{12}} \leq (\dim V)^{-1} n^3 e^{-\frac{n}{12}}$. It follows that

$$\begin{aligned} & 2M_n \sum_{(\pi, V) \in \widehat{\text{Sym}(n)} - \{\text{triv}, \text{sign}\}} (\dim V)^2 \left(6e^{-\frac{t_n \sqrt{\dim V}}{64}} + 2e^{-\frac{\dim V}{12}} \right) \\ & < 2M_n \left(\left(\sum_{(\pi, V) \in \widehat{\text{Sym}(n)}} (\dim V)^{-1} \right) - 2 \right) \left(6 + 2n^3 e^{-\frac{n}{12}} \right). \end{aligned} \quad (6.4)$$

The quantity $\sum_{(\pi, V) \in \widehat{\text{Sym}(n)}} (\dim V)^{-1}$ is precisely the Witten Zeta function of the symmetric group at 1. By [Lul96, MP02, LS04, Gam06] it is known that

$$\sum_{(\pi, V) \in \widehat{\text{Sym}(n)}} (\dim V)^{-1} = 2 + O(n^{-1})$$

and so (6.4) is $O(M_n(n^{-1} + n^2 e^{-\frac{n}{12}}))$. Thus, taking $M_n = o_{n \rightarrow \infty}(n)$ is sufficient for the existence of a basis satisfying (1.5).

In the case of $\text{Alt}(n)$, we note that any irreducible representation corresponds to two irreducible representations of $\text{Sym}(n)$ and so $\sum_{(\pi, V) \in \widehat{\text{Alt}(n)}} (\dim V)^{-1} = 1 + O(n^{-1})$. In addition, $\mathfrak{D}(\text{Alt}(n)) \geq n-1$ and so an identical argument to the case for $\text{Sym}(n)$ (this time there is no sign representation) gives the same result for $\text{Alt}(n)$. \square

Acknowledgements

MM and JT were supported by funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 949143). YZ was supported by funding from NSF CAREER award DMS-2044606, a Sloan Research Fellowship, and the MIT Solomon Buchsbaum Fund. We thank Sean Eberhard for useful discussions about this project. We thank Assaf Naor, Ashwin Sah, and Mehtaab Sawhney for their comments on a previous version of this paper that helped us improve it.

References

- [AISW21] Nalini Anantharaman, Maxime Ingremeau, Mostafa Sabri, and Brian Winn, *Quantum Ergodicity for Expanding Quantum Graphs in the Regime of Spectral Delocalization*, Journal de Mathématiques Pures et Appliquées **151** (2021), 28–98. [1](#)
- [ALM15] Nalini Anantharaman and Etienne Le Masson, *Quantum Ergodicity on Large Regular Graphs*, Duke Mathematical Journal **164** (2015), no. 4, 723–765. MR 3322309 [1](#), [1.10](#)
- [AN07] Nalini Anantharaman and Stéphane Nonnenmacher, *Half-delocalization of eigenfunctions for the Laplacian on an Anosov manifold*, Annales de l’institut Fourier **57** (2007), no. 7, 2465–2523 (eng). [1.1](#)
- [Ana08] Nalini Anantharaman, *Entropy and the Localization of Eigenfunctions*, Annals of Mathematics **168** (2008), no. 2, 435–475. MR 2434883 [1.1](#)
- [AS93] Ralf Aurich and Frank Steiner, *Statistical Properties of Highly Excited Quantum Eigenstates of a Strongly Chaotic System*, Physica D: Nonlinear Phenomena **64** (1993), no. 1, 185–214. [1.1](#)
- [BG08] Jean Bourgain and Alex Gamburd, *Uniform expansion bounds for Cayley graphs of $SL_2(\mathbb{F}_p)$* , Ann. of Math. (2) **167** (2008), no. 2, 625–642. MR 2415383 [2](#)
- [BGGT15] Emmanuel Breuillard, Ben J Green, Robert M Guralnick, and Terence Tao, *Expansion in finite simple groups of Lie type*, Journal of the European Mathematical Society **17** (2015), no. 6, 1367–1434. [1](#)
- [BHY19] Roland Bauerschmidt, Jiaoyang Huang, and Horng-Tzer Yau, *Local Kesten-McKay Law for Random Regular Graphs*, Communications in Mathematical Physics **369** (2019), no. 2, 523–636. MR 3962004 [1](#), [1.9](#)
- [BKY17] Roland Bauerschmidt, Antti Knowles, and Horng-Tzer Yau, *Local Semicircle Law for Random Regular Graphs*, Communications on Pure and Applied Mathematics **70** (2017), no. 10, 1898–1960. [3](#)
- [BS01] Itai Benjamini and Oded Schramm, *Recurrence of Distributional Limits of Finite Planar Graphs*, Electronic Journal of Probability **6** (2001), 1 – 13. [1.10](#)
- [BW16] Matthew Brammall and B. Winn, *Quantum Ergodicity for Quantum Graphs without Back-Scattering*, Annales Henri Poincaré **17** (2016), no. 6, 1353–1382. [1](#)
- [Col85] Yves Colin de Verdière, *Ergodicité et Fonctions Propres Du Laplacien*, Communications in Mathematical Physics **102** (1985), no. 3, 497–502. MR 818831 [1.1](#)
- [DJ18] Semyon Dyatlov and Long Jin, *Semiclassical Measures on Hyperbolic Surfaces Have Full Support*, Acta Mathematica **220** (2018), no. 2, 297–339. [1.1](#)
- [FH91] William Fulton and Joe Harris, *Representation Theory: A First Course*, Springer Science & Business Media, 1991. [2.1](#), [3.2](#)
- [Gam06] Alex Gamburd, *Poisson–Dirichlet distribution for random Belyi surfaces*, The Annals of Probability **34** (2006), no. 5, 1827 – 1848. [6](#)
- [Gow08] William Gowers, *Quasirandom Groups*, Combinatorics, Probability and Computing **17** (2008), no. 3, 363–387. [1](#)
- [Has10] Andrew Hassell, *Ergodic Billiards That Are Not Quantum Unique Ergodic*, Annals of Mathematics **171** (2010), no. 1, 605–619. MR 2630052 [1.1](#)
- [HR92] Dennis A. Hejhal and Barry N. Rackner, *On the Topography of Maass Waveforms for $PSL(2, \mathbb{Z})$* , Experimental Mathematics **1** (1992), no. 4, 275–305. [1.1](#)
- [Lin06] Elon Lindenstrauss, *Invariant measures and arithmetic quantum unique ergodicity*, Annals of Mathematics (2006), 165–219. [1.1](#)
- [LS74] Vicente Landazuri and Gary M Seitz, *On the minimal degrees of projective representations of the finite Chevalley groups*, Journal of Algebra **32** (1974), no. 2, 418–443. [1](#)
- [LS04] Martin W. Liebeck and Aner Shalev, *Fuchsian groups, coverings of riemann surfaces, subgroup growth, random quotients and random walks*, Journal of Algebra **276** (2004), no. 2, 552–601. [6](#)
- [Lul96] Nathan Anton Mikerin Lulov, *Random walks on the symmetric group generated by conjugacy classes*, Ph.D. thesis, USA, 1996, AAI9710450. [6](#)

- [Map13] Kenneth Maples, *Quantum unique ergodicity for random bases of spectral projections*, arXiv:1306.3329 (2013). [1.1](#)
- [Mec19] Elizabeth S Meckes, *The random matrix theory of the classical compact groups*, vol. 218, Cambridge University Press, 2019. [5](#)
- [MP02] Thomas W. Müller and Jan-Christoph Puchta, *Character theory of symmetric groups and subgroup growth of surface groups*, Journal of the London Mathematical Society **66** (2002), no. 3, 623–640. [6](#)
- [NSSZ22] Assaf Naor, Ashwin Sah, Mehtaab Sawhney, and Yufei Zhao, *Cayley graphs that have a quantum ergodic eigenbasis*, arXiv:2207.05527 (2022). [1](#)
- [OH88] Patrick W. O’Connor and Eric J. Heller, *Quantum Localization for a Strongly Classically Chaotic System*, Physical Review Letters **61** (1988), no. 20, 2288–2291. [1.1](#)
- [RS94] Zeév Rudnick and Peter Sarnak, *The Behaviour of Eigenstates of Arithmetic Hyperbolic Manifolds*, Communications in Mathematical Physics **161** (1994), no. 1, 195–213. MR 1266075 [1.1](#)
- [Sar11] Peter Sarnak, *Recent Progress on the Quantum Unique Ergodicity Conjecture*, American Mathematical Society. Bulletin. New Series **48** (2011), no. 2, 211–228. MR 2774090 [1.1](#)
- [Šni74] Alexander I. Šnirel’man, *Ergodic Properties of Eigenfunctions*, Uspehi Mat. Nauk **29** (1974), no. 6(180), 181–182. MR 0402834 [1.1](#)
- [Sou10] Kannan Soundararajan, *Quantum unique ergodicity for $SL_2(\mathbb{Z}) \backslash \mathbb{H}$* , Annals of Mathematics **172** (2010), no. 2, 1529–1538. [1.1](#)
- [SSZ20] Ashwin Sah, Mehtaab Sawhney, and Yufei Zhao, *Cayley Graphs Without a Bounded Eigenbasis*, International Mathematics Research Notices (2020), rnaa298. [1](#)
- [SX91] Peter Sarnak and Xiao Xi Xue, *Bounds for multiplicities of automorphic representations*, Duke Math. J. **64** (1991), no. 1, 207–227. MR 1131400 [2](#)
- [SZ93] Gary M. Seitz and Alexander E. Zaleskii, *On the Minimal Degrees of Projective Representations of the Finite Chevalley Groups, II*, Journal of Algebra **158** (1993), no. 1, 233–243. [1](#)
- [Tao15] Terence Tao, *Expansion in finite simple groups of Lie type*, vol. 164, American Mathematical Soc., 2015. [1](#)
- [Van97] Jeffrey M. VanderKam, *L^∞ norms and quantum ergodicity on the sphere*, International Mathematics Research Notices **1997** (1997), no. 7, 329–347. [1.1](#)
- [Zel87] Steven Zelditch, *Uniform Distribution of Eigenfunctions on Compact Hyperbolic Surfaces*, Duke Mathematical Journal **55** (1987), no. 4, 919–941. MR 916129 [1.1](#)
- [Zel92] ———, *Quantum ergodicity on the sphere*, Communications in Mathematical Physics **146** (1992), no. 1, 61–71. [1.1](#)
- [Zel96] Steve Zelditch, *A random matrix model for quantum mixing*, International Mathematics Research Notices **1996** (1996), no. 3, 115–137. [1.1](#)
- [Zel14] ———, *Quantum ergodicity of random orthonormal bases of spaces of high dimension*, Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences **372** (2014), no. 2007, 20120511. [1.1](#)

Michael Magee,
 Department of Mathematical Sciences,
 Durham University,
 Lower Mountjoy, DH1 3LE Durham,
 United Kingdom
michael.r.magee@durham.ac.uk

Joe Thomas,
 Department of Mathematical Sciences,
 Durham University,

Lower Mountjoy, DH1 3LE Durham,
United Kingdom
`joe.thomas@durham.ac.uk`

Yufei Zhao,
Department of Mathematics,
Massachusetts Institute of Technology,
Cambridge, MA 02139,
USA
`yufeiz@mit.edu`