

Anomalies of non-Abelian finite groups via cobordism

Joe Davighi,^a Ben Gripaios^b and Nakarin Lohitsiri^c

^a*Physics Institute, University of Zurich,
Zurich, Switzerland*

^b*Cavendish Laboratory, University of Cambridge,
Cambridge, U.K.*

^c*Department of Mathematical Sciences, Durham University,
Durham, U.K.*

E-mail: joedavighi@gmail.com, gripaios@hep.phy.cam.ac.uk,
[nakarin.lohitsiri@durham.ac.uk](mailto:nakaran.lohitsiri@durham.ac.uk)

ABSTRACT: We use cobordism theory to analyse anomalies of finite non-abelian symmetries in 4 spacetime dimensions. By applying the method of ‘anomaly interplay’, which uses functoriality of cobordism and naturality of the η -invariant to relate anomalies in a group of interest to anomalies in other (finite or compact Lie) groups, we derive the anomaly for every representation in many examples motivated by flavour physics, including S_3 , A_4 , Q_8 , and $SL(2, \mathbb{F}_3)$.

In the case of finite abelian groups, it is well known that anomalies can be ‘truncated’ in a way that has no effect on low-energy physics, by means of a group extension. We extend this idea to non-abelian symmetries. We show, for example, that a system with A_4 symmetry can be rendered anomaly-free, with only one-third as many fermions as naïvely required, by passing to a larger symmetry. As another example, we find that a well-known model of quark and lepton masses utilising the $SL(2, \mathbb{F}_3)$ symmetry is anomalous, but that the anomaly can be cancelled by enlarging the symmetry to a $\mathbb{Z}/3$ extension of $SL(2, \mathbb{F}_3)$.

KEYWORDS: Anomalies in Field and String Theories, Discrete Symmetries, Theories of Flavour, Topological Field Theories

ARXIV EPRINT: [2207.10700](https://arxiv.org/abs/2207.10700)

Contents

1	Introduction	2
2	The cobordism classification of anomalies for finite groups	4
2.1	A tool: the Frobenius-Schur indicator	6
2.2	A strategy: anomaly interplay	6
3	Bordism group results	8
4	Computing finite group anomalies by anomaly interplay	9
4.1	$G = A_4$	9
4.2	$G = Q_8$	11
4.3	$G = \text{SL}(2, \mathbb{F}_3)$	13
4.3.1	Anomaly interplay with $\mathbb{Z}/3$ abelianisation	13
4.3.2	Anomaly interplay with Q_8 derived subgroup	15
4.3.3	Anomaly interplay with $\text{U}(2)$: proof ‘by Standard Model’	15
4.3.4	Example: an anomalous flavour physics model	17
4.4	Dihedral group symmetry	17
5	Not all anomalies are equal: anomaly-truncation by group extensions	18
5.1	General formalism	18
5.2	Known example: abelian finite anomalies	19
5.3	Extending A_4 by $\mathbb{Z}/3$	20
5.4	Extending $\text{SL}(2, \mathbb{F}_3)$ by $\mathbb{Z}/3$	23
5.4.1	Example (continued): anomaly truncation in a flavour physics model	25
6	Summary	25
A	Spin bordism groups $\Omega_d^{\text{Spin}}(BG)$ for various non-abelian finite group G	26
A.1	$G = \mathbb{Z}/2 \times \mathbb{Z}/2$	27
A.2	$G = A_4$	28
A.3	The quaternion group $G = Q_8$	28
A.4	$G = \text{SL}(2, \mathbb{F}_3)$	29
A.5	The dihedral group $G = D_{2n}$	31
B	TQFTs for Q_8 structure	35
C	Bordism generators of $\Omega_5^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3))$ and $\Omega_5^{\text{Spin}}(BQ_8)$	39
D	Decomposition of Standard Model representation	40

1 Introduction

Chiral symmetries of massless fermions typically exhibit anomalies, which play an important role in quantum field theory (QFT). If the chiral symmetry is gauged, then anomalies must cancel, modulo a coupling to a topological quantum field theory (TQFT). If the chiral symmetry is global, then there can be mixed ‘ABJ anomalies’ [1, 2] that render the classical symmetry broken, and there can also be ‘t Hooft anomalies [3] that prevents us from coupling background gauge fields to global symmetries. Due to their topological nature, ‘t Hooft anomalies are not renormalised and so provide powerful tools for analysing QFTs, especially in régimes of strong coupling.

It is now known that all such chiral fermion anomalies are captured by cobordism groups [4–9], where ‘cobordism’ is a particular cohomology theory.¹ Given a symmetry type,² an appropriate cobordism group classifies both perturbative (‘local’) anomalies, i.e. those computed via one-loop Feynman diagrams, and also non-perturbative (‘global’) anomalies [10] in that symmetry. While local anomalies are easily analysed by more pedestrian means, in the case of global anomalies the rigorous cobordism classification supersedes previous arguments based on homotopy [11], and allows anomalies to be analysed systematically using the tools of algebraic topology.

This cobordism classification of anomalies has been used in many contexts in recent years, with applications in both high energy physics (including string theory) and condensed matter physics. On the high energy side, applications include the elucidation of an order 16 anomaly in $\frac{\text{Spin} \times \mathbb{Z}/4}{\mathbb{Z}/2}$ symmetry [12] and related anomalies [13], a new SU(2) anomaly for non-spin spacetimes [14], anomalies (and their absence) in the Standard Model (SM) and Beyond the SM [15–20], examples in 6d [11, 21] and 8d [22, 23], anomalies in duality groups [24, 25], anomaly cancellation in heterotic string theory [26, 27], and a newly discovered anomaly in Type IIB string theory [28].

In this paper, we apply the rigorous cobordism classification of anomalies to the case of finite non-abelian symmetries in 4d. Through anomaly inflow, it has been shown that fermion anomalies are given by the exponentiated η -invariant evaluated on a manifold in one dimension higher [9]. For a finite group G , in 4d, there are no perturbative anomalies and so the exponentiated η -invariant becomes a 5d bordism invariant. Thus, if fermions are defined using an ordinary spin structure, then the possible anomalies are elements of the (abelian) group of homomorphisms from the bordism group $\Omega_5^{\text{Spin}}(BG)$ to $U(1)$, given by the exponentiated η -invariant. The first step in characterising the anomalies is thus to find that bordism group. In the abelian case, the bordism groups were explored by Hsieh

¹For us, the word ‘bordism’ will always refer to a covariant functor (that was once upon a time introduced as ‘cobordism’), while we reserve the word ‘cobordism’ to denote its contravariant dual. Roughly speaking, the elements in a bordism group are equivalence classes of manifolds, equipped with certain structures, where two manifolds are considered equivalent if there exists a manifold in one dimension higher joining them. By cobordism groups, we mean the (shifted) Anderson dual to bordism, or in a special case the Pontryagin dual.

²The ‘symmetry type’, defined in the technical sense of Definition 2.4 in Freed and Hopkins’ ref. [8], includes not only the internal symmetry but also the spacetime symmetry, allowing for possible quotients between the two. As is usual in the study of anomalies, we assume Euclidean signature throughout.

in ref. [29] (see also [16]), and a smattering of non-abelian finite group anomalies have been studied in the literature.³

We here compute $\Omega_d^{\text{Spin}}(BG)$, in degrees $d = 1, \dots, 6$, for a variety of non-abelian finite groups (see table 1). The examples that we choose to showcase our methods are not random, but chosen because of their applications to flavour physics, where the groups S_3 , A_4 , Q_8 , and $\text{SL}(2, \mathbb{F}_3)$ have been widely used in explanations of the quark flavour hierarchies and/or neutrino mass and mixing data [31–36].

We then proceed to compute the anomalies, by determining the exponentiated η -invariant for each irreducible representation (henceforth *irrep*) of G , for the chosen examples listed in table 1 in 4d. We use a variety of tools to perform this task completely. We find that S_3 and Q_8 are completely anomaly free, while A_4 and $\text{SL}(2, \mathbb{F}_3)$ suffer from anomalies of order 9.⁴ However, the moral of the story is that the anomalies should be computable for any group of interest, if one knows enough tricks of algebraic topology.

The main tool we use for deriving these anomalies is the fact that the exponentiated η -invariant is a natural transformation between two appropriate functors (see section 2.2), through which the (unknown) anomalies in symmetry G are related to (known) anomalies in a different group H , where G and H are related by a homomorphism. This mapping between anomalies in different symmetry types, dubbed ‘anomaly interplay’ [37–40], is particularly useful when there are non-trivial maps going in both directions between G and H ; for example, this occurs when G is a semi-direct product $K \rtimes H$, as is the case in several of our examples e.g. $\text{SL}(2, \mathbb{F}_3) \cong Q_8 \rtimes \mathbb{Z}/3$.

In the context of particle physics, there is a subtlety pertaining to finite group symmetries that is not seen for connected Lie groups. The subtlety, which was first appreciated in the context of \mathbb{Z}/k symmetries by Banks and Dine [41] (see [29] for a discussion from the bordism perspective), is that anomalies in a finite symmetry G are not ‘invariant’ under extensions of G , where such extensions cannot necessarily be distinguished from G by the low-energy physicist. In particular, a representation \mathbf{r} of G whose exponentiated η -invariant is non-trivial can pull back to a representation of a group extension $H \twoheadrightarrow G$ with a trivial exponentiated η -invariant. For example, take $G = \mathbb{Z}/3$, and let the (reducible) representation \mathbf{r} denote three copies of the fundamental representation of $\mathbb{Z}/3$. While the representation \mathbf{r} has an anomaly $\exp(2\pi i/3)$, meaning that naïvely three such fermions would be required to be anomaly-free, the representation \mathbf{r} nevertheless pulls back to an anomaly-free representation of the extension $\mathbb{Z}/9 \twoheadrightarrow \mathbb{Z}/3$.

The final goal of this paper is to extend this idea of ‘truncating’ finite anomalies by a group extension to the case of finite non-abelian groups. In both cases where we find non-trivial anomalies of order 9, namely for $G = A_4$ and $G = \text{SL}(2, \mathbb{F}_3)$, we show that the anomaly is truncated to an order 3 anomaly by a non-trivial extension of G by $\mathbb{Z}/3$. The anomalies that are left are consistent with the recent ref. [42] (see also [43, 44]), which analysed the ‘linear part’ of finite gauge anomalies using character theory.

³Examples include quaternionic groups in 6d [16] and dihedral groups that mix with spacetime symmetry [30].

⁴By the *order* of an anomaly, we mean the order of the image of the exponentiated η -invariant in the appropriate cobordism group (which, for us, is always a finite group). This notion will be made precise in section 2.

There are some new mathematical results in this paper, including:

- Calculation of the odd torsion for $\Omega_d^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3))$ for $d \leq 6$ (appendix A.4);
- Proof that $\Omega_5^{\text{Spin}}(BD_{2n}) = 0$ for any $n \in \mathbb{Z}$ (appendix A.5);
- Calculation of $\Omega_d^{\text{Spin}}(BD_{2n})$ for $d \leq 5$ when n is a power of 2 (appendix A.5);
- Computations of the exponentiated η -invariant for all complex (and pseudo-real) irreps of A_4 and $\text{SL}(2, \mathbb{F}_3)$ in sections 4.1 and 4.3.

The structure of the paper is as follows. In section 2 we review the precise connection between chiral fermion anomalies and η -invariants and the consequent classification of anomalies using cobordism, specialising to the case of finite groups. We review the tools of anomaly interplay on which we rely. In section 3 we record all our spin-bordism computations, for a selection of finite non-abelian groups in degrees 1 through 6 — the calculations themselves are collected in appendix A. In section 4 we use anomaly interplay to compute η -invariants for these groups, and thus derive complete information about anomalies. In section 5 we formally introduce the notion of ‘anomaly truncation’ by finite group extensions, and show how this works out in our two non-trivial examples, before summarising in section 6.

2 The cobordism classification of anomalies for finite groups

By now it is well-established that fermionic anomalies are captured by (co)bordism groups. We begin by briefly reviewing why this is the case. Consider a system of massless chiral fermions $\{\psi\}$ on a d -dimensional closed manifold Σ_d , with a metric g , spin structure s , and principal G -bundle with connection A . For now, G may be either a compact Lie group or a finite group.

Recently, Witten and Yonekura proved [9] a long-suspected formula [4, 5, 7], which is that the phase of the fermionic partition function obtained by integrating over $\{\psi\}$ is equal to the exponentiated η -invariant of Atiyah, Patodi and Singer [45–47], evaluated on a manifold X_{d+1} whose boundary is Σ_d and to which all the structure extends.⁵ We write $Z_\psi[\Sigma, (A, g, s)] = |Z_\psi| \exp(2\pi i \eta(X_{d+1}))$.

For this partition function to make sense as a description of local physics in d -dimensions, it should not depend on the choice of auxiliary manifold X_{d+1} , nor on the particular choice of extension of structure thereto. Thus, if we had chosen another such extension X'_{d+1} with structure, then we must have $\exp(2\pi i \eta(X'_{d+1})) = \exp(2\pi i \eta(X_{d+1}))$. Now, since X and X' share a boundary on which the structures agree, one can glue X_{d+1} and $-X'_{d+1}$, where the minus sign denotes orientation reversal, to form a closed $(d+1)$ -manifold

⁵If the relevant bordism group in degree d is non-vanishing, then there are manifolds for which no extension exists. A 4d example is the K3 manifold, for which there is no extension of the spin structure to a 5-manifold bounded by K3. For such non-nullbordant spacetimes, the partition function suffers from an ambiguity that corresponds to assigning a generalised theta angle to each generator of the bordism group Ω_d [48].

\overline{X}_{d+1} . The exponentiated η -invariant obeys a gluing law [5], which implies our previous condition is equivalent to requiring $\exp(2\pi i \eta(\overline{X}_{d+1})) = 1$. To be fully consistent with locality, one thus requires that

$$\exp(2\pi i \eta(\overline{X}_{d+1})) = 1 \quad (2.1)$$

for *all* closed $(d+1)$ -manifolds \overline{X} with structure.

The condition (2.1) implies that the theory is anomaly-free in the more traditional sense. This is because, given any gauge transformation $A \rightarrow A^g$ of the background connection (possibly combined with a diffeomorphism φ of Σ_d), one can construct a mapping torus $T_{d+1}^g \cong \Sigma_d \times_g S_1$ by taking a cylinder and gluing the ends using the gauge transformation (plus diffeomorphism). By cutting and gluing, one sees that the variation of the partition function is given by $Z[\Sigma, (A, g, s)]/Z[\varphi(\Sigma), \varphi(A^g, g, s)] = \exp(2\pi i \eta(T_{d+1}^g))$, where we have used the fact that the modulus of the partition is necessarily anomaly-free. The mapping torus T_{d+1}^g is a closed $(d+1)$ -manifold with structure (A, g, s) , on which the exponentiated η -invariant is trivial by (2.1). Thus, the locality condition implies complete anomaly-freedom, from both perturbative and non-perturbative anomalies in the traditional sense, but is much stronger.⁶

In this paper we are interested in the special case of theories with symmetry group G being a *finite group*, for which the analysis of anomalies simplifies. Let us further assume for now that $d = 4$.⁷ In this scenario, the fermion anomaly given by the exponentiated η -invariant⁸ becomes a bordism invariant. This can be easily seen by evaluating the exponentiated η -invariant on a closed manifold \overline{X}_5 that bounds Y_6 . The Atiyah-Patodi-Singer (APS) index theorem tells us that

$$\exp(2\pi i \eta(X_5)) = \exp\left(2\pi i \int_{Y_6} \Phi_6\right), \quad (2.2)$$

where Φ_6 , the anomaly polynomial, is a closed 6-form with integer periods built from the curvature of Y_6 and the field strength of the connection A . For finite G , A is flat and consequently Φ_6 vanishes.⁹ Thus, the exponentiated η -invariant evaluated on a pair of bordant manifolds X_5 and X'_5 must be equal, since

$$\frac{\exp(2\pi i \eta(X_5))}{\exp(2\pi i \eta(X'_5))} = \exp(2\pi i \eta(X_5 \sqcup (-X'_5))) = 1, \quad (2.3)$$

⁶Such a condition is also thought to be needed ultimately for consistently quantizing gravity; see e.g. [49, 50] for related arguments.

⁷In other dimensions, e.g. in 2d or 6d, the situation is complicated somewhat due to the presence of pure gravitational anomalies. In practice, however, these pose no significant issue for analysing anomalies, because one can trivially cancel such pure gravity anomalies by adding enough neutral fermions in the opposite chirality.

⁸Like any anomaly theory, the exponentiated η -invariant for a more general symmetry structure is an example of a reflection positive invertible quantum field theory. Much is known mathematically about the space of such theories — for example, in the case of anomaly theories that are moreover topological, they are rigorously classified in terms of stable homotopy [8].

⁹Because we are in $d = 4$, there is no ‘pure gravitational anomaly’ terms in the anomaly polynomial that do not depend on the field strength of A .

where we used the fact that $X_5 \sqcup (-X'_5)$ is a boundary to get the last equality. Therefore, the exponentiated η -invariant defines an element of the group of homomorphisms from the bordism group $\Omega_5^{\text{Spin}}(BG)$ to $U(1)$, which we denote

$$\mathcal{U}_{\text{Spin}}^6(BG) := \text{Hom}\left(\Omega_5^{\text{Spin}}(BG), U(1)\right), \quad (2.4)$$

adopting the notation of e.g. [51]. This abelian group is pure torsion,¹⁰ implying that the anomaly is of finite order, as is always the case for a global anomaly. In the rest of the paper, we call $\mathcal{U}_{\text{Spin}}^6(BG)$ a *cobordism group*.¹¹

Clearly, we can specify the exponentiated η -invariant by giving its values of a set of generators of $\Omega_5^{\text{Spin}}(BG)$ for a single representation from each equivalence class of irreps of G . From there, the complete set of conditions for anomaly cancellation in a 4d finite G gauge theory can be obtained. While this is in general a daunting task, for the abelian case, in which $G = \prod_i \mathbb{Z}/k$ is a product of cyclic groups, this task has been completed in ref. [29]. One goal of this paper is to extend these computations to some non-abelian finite groups, such as S_3 , A_4 , Q_8 , and $\text{SL}(2, \mathbb{F}_3)$.

2.1 A tool: the Frobenius-Schur indicator

One elementary fact that we can easily leverage is that, in 4d, chiral fermions transforming in real representations are anomaly-free. For finite groups, it is especially straightforward to check whether a representation \mathbf{r} is real simply from its character $\chi(g)$.¹² To wit, one computes the Frobenius-Schur indicator,

$$\iota_{\mathbf{r}} := \frac{1}{|G|} \sum_{g \in G} \chi(g^2), \quad (2.6)$$

which equals 1, 0, or -1 when \mathbf{r} is a real, complex, or pseudo-real irrep, respectively. To compute the anomaly for a complex or pseudo-real representation, in 4d, requires less trivial methods.

2.2 A strategy: anomaly interplay

The main strategy that we use to calculate anomalies is to exploit ‘anomaly interplay’. The idea is to pullback anomalies from one symmetry type to another, by viewing the exponentiated η -invariant as a natural transformation between contravariant functors. (Here we

¹⁰This can be seen from looking at the E^2 page of the Atiyah-Hirzebruch spectral sequence for the spin bordism group of BG associated to the fibration $\text{pt} \rightarrow BG \rightarrow BG$:

$$E_{p,q}^2 = H_p(BG; \Omega_q^{\text{Spin}}(\text{pt})) \Rightarrow \Omega_{p+q}^{\text{Spin}}(BG). \quad (2.5)$$

On the diagonal $p+q=5$, the only possibly non-trivial entries are of the form $H_i(BG; A)$ where $i > 0$ and $A = \mathbb{Z}$ or $\mathbb{Z}/2$. These homology groups are finite when G is finite. Since $|\Omega_5^{\text{Spin}}(BG)| \leq \left| \bigoplus_{p+q=5} E_{p,q}^2 \right|$, we obtain the desired result.

¹¹This agrees with a more general (and more widely used) usage of the term cobordism to mean the (shifted) Anderson dual of the bordism group [8, 38, 52]. When G is finite and $d=4$, as we consider here, the two coincide.

¹²For compact Lie groups things are only slightly more involved, with the Frobenius-Schur indicator then being given by the integral $\iota_{\mathbf{r}} = \int_{g \in G} \chi(g^2) d\mu$, where $d\mu$ is the Haar measure normalised such that $\int d\mu = 1$.

use the term ‘anomaly’ and the exponentiated η -invariant interchangeably.) Physics-wise, this harks back to work of Elitzur and Nair [53] who, following Witten [54], showed that the 4d SU(2) anomaly can be derived by ‘pulling back’ a local anomaly from e.g. SU(3). Anomaly interplay is rather trivially defined with a little category theory, which we now introduce before continuing.

Let FinGp be the usual category of finite groups and let Ab be the usual category of abelian groups. Then $\mathcal{U}_{\text{Spin}}^6(B\cdot)$ defines a contravariant functor $\text{FinGp} \rightarrow \text{Ab}$. A second contravariant functor $\text{FinGp} \rightarrow \text{Ab}$ is given by the representation ring $RU(\cdot)$, where we forget the ring multiplication, regarding $RU(G)$ just as an abelian group. Concretely, $RU(G)$ is a free abelian group with basis given by (isomorphism classes of) irreps of G and sum given by direct sum of representations. Writing $\mathbb{Z}\langle \mathbf{r}_i \rangle$ for the free abelian group generated by an irrep \mathbf{r}_i , we can write $RU(G)$ explicitly as

$$RU(G) = \bigoplus_i \mathbb{Z}\langle \mathbf{r}_i \rangle, \quad (2.7)$$

where i runs through all irreps of G . We emphasise that, for our purposes, (2.7) should be thought of as a free abelian group, and we forget its ring structure. This permits us to consider subgroups thereof.

The exponentiated η -invariant is a natural transformation from the functor $RU(\cdot)$ to the functor $\mathcal{U}_{\text{Spin}}^6(B\cdot)$ [39, 40]. This means that, given a homomorphism $f : G \rightarrow K$ between finite groups, there are pullbacks induced on both these functors,

$$RU(f) : RU(K) \rightarrow RU(G), \quad (2.8)$$

$$\mathcal{U}(f) : \mathcal{U}_{\text{Spin}}^6(BK) \rightarrow \mathcal{U}_{\text{Spin}}^6(BG), \quad (2.9)$$

such that the diagram

$$\begin{array}{ccc} RU(K) & \xrightarrow{RU(f)} & RU(G) \\ \exp(2\pi i \eta)_K \downarrow & & \downarrow \exp(2\pi i \eta)_G \\ \mathcal{U}_{\text{Spin}}^6(BK) & \xrightarrow{\mathcal{U}(f)} & \mathcal{U}_{\text{Spin}}^6(BG) \end{array} \quad (2.10)$$

commutes (where we use the usual subscript notation to denote a component of the natural transformation $\exp(2\pi i \eta)$). By ‘anomaly interplay’ we simply refer to the constraints placed on anomalies of G that follow from this commutative diagram. This version of anomaly interplay, in which both groups involved are finite, has been applied to the study of anomalies in two-dimensions in refs. [39, 40].¹³

One particular trick we will frequently employ is to pullback whole short exact sequences that relate a non-abelian group G to *abelian* finite groups. For example, we are

¹³Another version of anomaly interplay between anomalies of finite groups and anomalies of compact Lie groups has been used in refs. [37, 38]. In fact, at one point in this paper (section 4.3.3) we will need to go outside the framework we have set up, and consider a compact Lie group, namely U(2), instead of a finite group. But since we do so for a representation of U(2) which has no local anomalies, very similar arguments apply.

p	1	2	3	4	5	6
$\tilde{\Omega}_p^{\text{Spin}}(BS_3)$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/8 \times \mathbb{Z}/3$	0	0	0
$\tilde{\Omega}_p^{\text{Spin}}(BA_4)$	$\mathbb{Z}/3$	$\mathbb{Z}/2$	$\mathbb{Z}/12$	0	$\mathbb{Z}/9$	$\mathbb{Z}/2$
$\tilde{\Omega}_p^{\text{Spin}}(BQ_8)$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/8 \times (\mathbb{Z}/4)^2$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$
$\tilde{\Omega}_p^{\text{Spin}}(BSL(2, \mathbb{F}_3))$	$\mathbb{Z}/3$	0	$\mathbb{Z}/8 \times \mathbb{Z}/3$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \times \mathbb{Z}/9$	$\mathbb{Z}/2$
$\tilde{\Omega}_p^{\text{Spin}}(BD_{2n}), \ n \text{ odd}$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/8 \times \mathbb{Z}/n$	0	0	0
$\tilde{\Omega}_p^{\text{Spin}}(BD_{2n}), \ n = 2^{k+1}$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^3$	$(\mathbb{Z}/8)^2 \times \mathbb{Z}/(2n)$	$(\mathbb{Z}/2)^2$	0	$\mathbb{Z}/2$

Table 1. Reduced spin-bordism groups in degrees 1 through 6 of BG , for various finite non-abelian groups G . The full (i.e. unreduced) spin bordism groups are obtained as $\Omega_p^{\text{Spin}}(BG) \cong \tilde{\Omega}_p^{\text{Spin}}(BG) \oplus \Omega_p^{\text{Spin}}(\text{pt})$.

guaranteed a short exact sequence defining the ‘abelianisation’ of G :

$$0 \rightarrow G' \rightarrow G \rightarrow G/G' \rightarrow 0. \quad (2.11)$$

Here, G' is the derived subgroup of G , i.e. the normal subgroup of G generated by commutators $[g, h] := ghg^{-1}h^{-1}$. The quotient group G/G' is necessarily abelian, and is called the abelianisation of G . By pulling back such sequences, particularly in cases where they split, often allows us to completely derive the anomalies in G from known building blocks, i.e. from anomalies in the abelian case.

Before we begin in earnest, we introduce a few more notions that will be useful in what follows. Firstly, the *order of the anomaly of a representation \mathbf{r}* is the order of the image of \mathbf{r} under the exponentiated η -invariant. We can also define a slightly more general notion, which is the *order of the anomaly of a finite group G* , to be simply the order of the image of the exponentiated η -invariant in $\mathcal{U}_{\text{Spin}}^6(BG)$. Lastly, we define the *subgroup of anomaly-free representations* $RU_0(G)$ to be

$$RU_0(G) := \ker(\exp(2\pi i \eta)) \subset RU(G) \quad (2.12)$$

for any finite group G . This notion will be especially useful when we come to discuss the ‘truncation’ of finite group anomalies in section 5.

3 Bordism group results

To compute the relevant bordism groups, we rely on standard spectral sequence techniques, using a combination of the Atiyah-Hirzebruch spectral sequence (AHSS) and the Adams spectral sequence (ASS). We tabulate our results for the (reduced) spin-bordism groups of BG for various non-abelian finite groups G in table 1.

The computations themselves, some of which can be found in existing literature, are recorded in appendix A. These bordism computations provide the platform from which we systematically, and comprehensively, study anomalies in 4d for various finite G symmetries.

	1	4	4	3
A_4	1	σ	σ^2	τ
1	1	1	1	1
1'	1	ω	ω^2	1
$\overline{\mathbf{1'}}$	1	ω^2	ω	1
3	3	0	0	-1

Table 2. Character table of A_4 , the alternating group of order 4. Conjugacy classes are labelled by a representative element with the numbers of elements in each class atop. ω is a cube root of unity.

4 Computing finite group anomalies by anomaly interplay

4.1 $G = A_4$

For our first example we consider $G = A_4$, the group of even permutations of four elements, equivalently the orientation-preserving symmetries of a regular tetrahedron. This group has often been used in flavour model building, particularly for controlling neutrino masses and mixing angles (e.g. [33, 34]).

There are three one-dimensional irreps and one three-dimensional irrep of A_4 . We denote these irreps by their dimensions, as **1**, **1'**, $\overline{\mathbf{1'}}$, and **3**, respectively. The character table (which can be obtained from e.g. the computer program **GAP** [55] or from ref. [56]) is given in table 2, where we use a presentation of A_4 with two generators σ and τ of order 2 and 3 satisfying $(\tau\sigma)^3 = 1$. The representation ring $RU(A_4)$ for the group A_4 can then be written explicitly as

$$RU(A_4) = \mathbb{Z} \langle \mathbf{1} \rangle \oplus \mathbb{Z} \langle \mathbf{1'} \rangle \oplus \mathbb{Z} \langle \overline{\mathbf{1'}} \rangle \oplus \mathbb{Z} \langle \mathbf{3} \rangle. \quad (4.1)$$

In appendix A.2 we record the spin bordism groups of BA_4 , taken from [57]. Of relevance to 4d anomalies, we have $\Omega_5^{\text{Spin}}(BA_4) \cong \mathbb{Z}/9$. Since there are no pure gravitational anomalies in 4d, the cobordism group that classifies anomalies is simply

$$\mathcal{U}_{\text{Spin}}^6(BA_4) \cong \Omega_5^{\text{Spin}}(BA_4) \cong \mathbb{Z}/9 \quad (4.2)$$

also. In this section we calculate the anomalies for $G = A_4$ in 4d, using anomaly interplay. We will compute the anomaly associated with every irrep of A_4 , and from there write down the general condition for A_4 anomaly cancellation.

To begin, recall that the derived subgroup of A_4 is the Klein four-group, $A'_4 = \mathbb{Z}/2 \times \mathbb{Z}/2$. The abelianisation short exact sequence is

$$0 \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2 \rightarrow A_4 \rightarrow \mathbb{Z}/3 \rightarrow 0. \quad (4.3)$$

This sequence moreover splits, meaning that A_4 can be realised as a semi-direct product

$$A_4 = (\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/3, \quad (4.4)$$

so there is a map $j : \mathbb{Z}/3 \rightarrow A_4$ that embeds $\mathbb{Z}/3$ as a (non-normal) subgroup of A_4 . We therefore obtain the following commutative diagram, where the horizontal row is exact,

$$\begin{array}{ccccccc}
 0 & \longrightarrow & \mathbb{Z}/2 \times \mathbb{Z}/2 & \xrightarrow{i} & A_4 & \xrightarrow{\pi} & \mathbb{Z}/3 \longrightarrow 0 \\
 & & & & \uparrow j & \nearrow & \\
 & & & & \mathbb{Z}/3 & &
 \end{array} \quad (4.5)$$

For the other groups involved in this diagram, the relevant cobordism groups are $\mathcal{U}_{\text{Spin}}^6(B(\mathbb{Z}/2 \times \mathbb{Z}/2)) \cong \Omega_5^{\text{Spin}}(B(\mathbb{Z}/2 \times \mathbb{Z}/2)) = 0$ [58] (see appendix A.1), and $\mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3) \cong \Omega_5^{\text{Spin}}(B\mathbb{Z}/3) \cong \mathbb{Z}/9$.

Applying the representation ring functor and the cobordism functor to the right-hand triangle in (4.5), and relating the two via the $\exp(2\pi i \eta)$ natural transformation, we get the following commutative diagram:

$$\begin{array}{ccccc}
 RU(A_4) & \xleftarrow{\pi^*} & & RU(\mathbb{Z}/3) & \\
 \downarrow \exp(2\pi i \eta) & \searrow j^* & & \nearrow & \downarrow \\
 & RU(\mathbb{Z}/3) & & & \\
 \mathcal{U}_{\text{Spin}}^6(BA_4) \cong \mathbb{Z}/9 & \xleftarrow{\pi^*} & & \mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3) \cong \mathbb{Z}/9 & \\
 & \searrow j^* & & \nearrow & \\
 & \mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3) \cong \mathbb{Z}/9 & & &
 \end{array} \quad (4.6)$$

Notice that all groups in the lower triangle are isomorphic to $\mathbb{Z}/9$, and that since $j^* \circ \pi^* = \text{id}$, both these maps must be group isomorphisms. Thus, if we know the anomalies for representations of $\mathbb{Z}/3$, and we know how to map representations of A_4 to and from representations of $\mathbb{Z}/3$, then commutativity of (4.6) allows us to deduce the anomaly for every irrep of A_4 .¹⁴

We first recap the anomalies for $\mathbb{Z}/3$. In general, all irreps of the cyclic group \mathbb{Z}/n are one-dimensional. There is an irrep taking $1 \bmod n \in \mathbb{Z}/n$ to $\exp(2\pi i q/n)$ for each $q = 0, 1, \dots, n-1$, denoted here by $[q]_n$. In particular, for $\mathbb{Z}/3$, we have three irreps, namely the trivial representation $[0]_3$, and the fundamental representation $[1]_3$ together with its complex conjugate $[2]_3$. The exponentiated η -invariant maps, along the middle vertical arrow in (4.6), the fundamental representation $[1]_3$ to $1 \bmod 9$ in $\mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3)$ [29].

We now relate the representations of A_4 and those of $\mathbb{Z}/3$ under j^* and π^* . Recall that j^* maps an A_4 representation to a $\mathbb{Z}/3$ representation. From the character table, it is

¹⁴We emphasise that, in order to unambiguously relate A_4 anomalies to $\mathbb{Z}/3$ anomalies in this way, it is crucial that there are maps j^* and π^* going in both directions, which follows from the right-splitting of (4.3). (And this property is not even enough in general, if the bordism groups differ — see section 4.3 for the relevant example of $\text{SL}(2, \mathbb{F}_3) = Q_8 \rtimes \mathbb{Z}/3$.)

clear that

$$\begin{aligned}
 j^* : RU(A_4) &\rightarrow RU(\mathbb{Z}/3) : \\
 \mathbf{1} &\mapsto [0]_3, \\
 \mathbf{1}' &\mapsto [1]_3, \\
 \overline{\mathbf{1}'} &\mapsto [2]_3.
 \end{aligned} \tag{4.7}$$

For the three-dimensional representation $\mathbf{3}$, there is a basis where the order-3 generator σ of a $\mathbb{Z}/3$ subgroup acts by the 3×3 traceless matrix $\sigma = \text{diag}(1, \omega, \omega^2)$, where $\omega = \exp(2\pi i/3)$. Therefore,

$$j^* : \mathbf{3} \rightarrow [0]_3 \oplus [1]_3 \oplus [2]_3 \tag{4.8}$$

for the triplet representation.

By the anomaly interplay diagram (4.6), we can then conclude that the trivial representation $\mathbf{1}$ and the three-dimensional irrep $\mathbf{3}$ of A_4 are anomaly free, while the irreps $\mathbf{1}'$ and $\overline{\mathbf{1}'}$ have non-trivial anomalies of order 9 with the opposite sign. Our result is consistent with the fact that chiral fermions in real representations cannot contribute to any anomaly in 4d [7]. (The triplet of A_4 is a real representation, which can be verified directly from the character table by applying the Frobenius-Schur indicator (2.6).) The condition for A_4 anomaly cancellation in 4d is thus

$$n(\mathbf{1}') - n(\overline{\mathbf{1}'}) = 0 \bmod 9 \quad (A_4 \text{ anomaly cancellation in 4d}), \tag{4.9}$$

where $n(\mathbf{r})$ denotes the number of left-handed fermions minus the number of right-handed fermions in the representation \mathbf{r} . There is no constraint on the number of (anomaly-free) triplet representations. This information can be restated in a more succinct manner in terms of the subgroup of non-anomalous representations $RU_0(A_4)$ of $RU(A_4)$. Our result shows that this group is given by

$$RU_0(A_4) = \mathbb{Z} \langle \mathbf{1} \rangle \oplus \mathbb{Z} \langle \mathbf{1}' \oplus \overline{\mathbf{1}'} \rangle \oplus \mathbb{Z} \langle \mathbf{1}^{\oplus 9} \rangle \oplus \mathbb{Z} \langle \mathbf{3} \rangle. \tag{4.10}$$

The full anomaly discussion above also enables us to relate to the notion of anomaly-free subgroups used in refs. [42, 44]. Notice that there is an order 9 anomaly for A_4 if and only if the fermions are in either the $\mathbf{1}'$ representation or its complex conjugate, which do not transform under the normal subgroup $\mathbb{Z}/2 \times \mathbb{Z}/2$. On the other hand, the irrep $\mathbf{3}$ that transforms non-trivially under $\mathbb{Z}/2 \times \mathbb{Z}/2$ has no anomaly. Thus, the anomaly-free subgroup of the alternating group A_4 is its normal subgroup $\mathbb{Z}/2 \times \mathbb{Z}/2$, in agreement with the result of [42].

In section 5.3 we will tackle the issue of group extensions. We will there learn that the order 9 anomaly (4.9) can in fact be truncated down to an order 3 anomaly by a group extension of A_4 by $\mathbb{Z}/3$.

4.2 $G = Q_8$

Next, we turn to the quaternion group $G = Q_8$, which has been used for example in quark mass model building [31]. It is a multiplicative group of order 8 whose elements consist of

	1	1	2	2	2
Q_8	$\{1\}$	$\{-1\}$	$\{i, -i\}$	$\{j, -j\}$	$\{k, -k\}$
1	1	1	1	1	1
1_i	1	1	1	-1	-1
1_j	1	1	-1	1	-1
1_k	1	1	-1	-1	1
2	2	-2	0	0	0

Table 3. Character table of Q_8 , the quaternion group. The column are labelled by the conjugacy classes with the numbers of elements in each class atop.

1, and the unit quaternions i, j, k ($i^2 = j^2 = k^2 = -1$, $ij = k$), together with their additive inverses. There are three non-trivial 1-dimensional irreps, $\mathbf{1}_i$, $\mathbf{1}_j$, $\mathbf{1}_k$, which are the sign representations with kernels $\{\pm 1, \pm i\}$, $\{\pm 1, \pm j\}$, $\{\pm 1, \pm k\}$, respectively. The only other non-trivial irrep is 2-dimensional, denoted **2**. The character table for Q_8 is given in table 3.

From the bordism perspective, this example appears to be rather interesting; the derived subgroup is $Q'_8 \cong \mathbb{Z}/2$ and the abelianisation is $Q_8/Q'_8 \cong \mathbb{Z}/2 \times \mathbb{Z}/2$, both of which are anomaly-free. But $G = Q_8$ is not anomaly-free, with

$$\tilde{\Omega}_5^{\text{Spin}}(BQ_8) = \mathbb{Z}/2 \times \mathbb{Z}/2. \quad (4.11)$$

Thus, the anomalies for Q_8 , and more generally for any finite group G , cannot be ‘pieced together’ from the anomalies in its abelianisation and in its derived subgroup (with which one can ‘build’ G via the SES $G' \rightarrow G \rightarrow G/G'$), considered separately.

By applying the Frobenius-Schur indicator (2.6) to each of the irreps, we learn that all 1-dimensional irreps of Q_8 are real, while the 2-dimensional irrep is pseudo-real. Thus, the singlets are anomaly-free, but there could be an order 2 anomaly from fermions in the doublet representation. It turns out, however, that the η -invariant evaluated on generators of $\Omega_5^{\text{Spin}}(BQ_8)$ vanishes [59]. Thus,

$$\text{All } Q_8 \text{ representations are anomaly-free in 4d.} \quad (4.12)$$

In this case, $RU_0(Q_8)$ is the whole of $RU(Q_8)$.

We remark that this is consistent with the anomaly interplay with $\mathbb{Z}/4$, using a subgroup embedding $\pi : \mathbb{Z}/4 \rightarrow Q_8$ (which is the only possibly-non-trivial anomaly interplay diagram involving a subgroup or quotient of Q_8). Under the pullback to the representation ring, one can check that every irrep of Q_8 pulls back to an anomaly-free representation of $\mathbb{Z}/4$.

There nonetheless remains a curiosity with this Q_8 example, which is that the bordism group classifying anomalies is not zero, but $\mathbb{Z}/2 \times \mathbb{Z}/2$. Even though there are no anomalies, the Q_8 structure allows non-trivial TQFT couplings that detect these bordism classes. We offer a partial account of these TQFTs in appendix B, where we show that one of the bordism invariants is

$$\text{Arf} \cup xy^2, \quad (4.13)$$

$\text{SL}(2, \mathbb{F}_3)$	1	1	4	4	6	4	4
	$\left\{\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}\right\}$	$\left\{\begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}\right\}$
1	1	1	1	1	1	1	1
1'	1	1	ω	ω^2	1	ω	ω^2
$\overline{\mathbf{1'}}$	1	1	ω^2	ω	1	ω^2	ω
2	2	-2	-1	-1	0	1	1
2'	2	-2	$-\omega$	$-\omega^2$	0	ω	ω^2
$\overline{\mathbf{2'}}$	2	-2	$-\omega^2$	$-\omega$	0	ω^2	ω
3	3	3	0	0	-1	0	0

Table 4. Character table for the group $\text{SL}(2, \mathbb{F}_3)$. The columns are labelled by representatives of conjugacy classes, with the numbers of elements in each class written above.

where x, y are degree 1 generators of $H^\bullet(BQ_8; \mathbb{Z}/2)$ and Arf is the Arf invariant of the spin structure. This phenomenon that free fermions do not account for every possible anomaly given by the cobordism group has also been observed in 2d when the symmetry group is the dihedral group D_8 [40].

4.3 $G = \text{SL}(2, \mathbb{F}_3)$

We now consider the group $G = \text{SL}(2, \mathbb{F}_3)$ of special linear matrices with elements in the finite field \mathbb{F}_3 , that was used in flavour physics model building in refs. [35, 36]. This group has order-24, with 7 conjugacy classes of element and thus 7 irreps, with the character table shown in table 4.

By computing the Frobenius-Schur indicator (2.6), we immediately learn that the irreps **1**, and **3** are real and hence anomaly-free, while the irrep **2** is pseudo-real and suffers at most an order 2 anomaly. The other four irreps (two singlets, and two doublets) are, however, complex. In this section we analyse their anomalies using cobordism and anomaly interplay. This will be the most elaborate example we consider, in which a variety of techniques are needed to eventually arrive at the complete result (4.21).

4.3.1 Anomaly interplay with $\mathbb{Z}/3$ abelianisation

Using the bordism calculation in appendix A.4, we have that anomalies are here classified by the cobordism group

$$\mathcal{U}_{\text{Spin}}^6(B\text{SL}(2, \mathbb{F}_3)) \cong \Omega_5^{\text{Spin}}(B\text{SL}(2, \mathbb{F}_3)) \cong \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/9. \quad (4.14)$$

The derived subgroup of $\text{SL}(2, \mathbb{F}_3)$ is the quaternion group Q_8 , with abelianisation $\text{SL}(2, \mathbb{F}_3)/Q_8 \cong \mathbb{Z}/3$. Indeed, the short exact sequence¹⁵

$$0 \rightarrow Q_8 \xrightarrow{i} \text{SL}(2, \mathbb{F}_3) \xrightarrow{\pi} \mathbb{Z}/3 \rightarrow 0 \quad (4.15)$$

¹⁵We remark that there is also a short exact sequence $\mathbb{Z}/2 \hookrightarrow \text{SL}(2, \mathbb{F}_3) \twoheadrightarrow A_4$, which follows from the fact that $\text{SL}(2, \mathbb{F}_3)$ is a double cover of A_4 . This sequence is less useful for our purposes because, unlike the abelianisation sequence (4.15), it is not right-split.

is right-split, meaning that $\mathrm{SL}(2, \mathbb{F}_3)$ has a semi-direct product structure

$$\mathrm{SL}(2, \mathbb{F}_3) = Q_8 \rtimes \mathbb{Z}/3, \quad (4.16)$$

and that the abelianisation $\mathbb{Z}/3$ is also a subgroup, with subgroup embedding $j : \mathbb{Z}/3 \rightarrow \mathrm{SL}(2, \mathbb{F}_3)$. Recall from our previous examples that 4d anomalies for $G = \mathbb{Z}/3$ are classified by the cobordism group $\mathcal{U}_{\mathrm{Spin}}^6(B\mathbb{Z}/3) \cong \mathbb{Z}/9$, and that $\mathcal{U}_{\mathrm{Spin}}^6(BQ_8) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$.

We will use these maps to study anomalies in the $\mathrm{SL}(2, \mathbb{F}_3)$ symmetry by anomaly interplay. As usual, by applying the representation ring and cobordism functors to these maps, and relating the two via the exponentiated η -invariant, we obtain the following anomaly interplay commutative diagram¹⁶

$$\begin{array}{ccccc}
 RU(Q_8) & \xleftarrow{i^*} & RU(\mathrm{SL}(2, \mathbb{F}_3)) & \xleftarrow{\pi^*} & RU(\mathbb{Z}/3) \\
 \downarrow \scriptstyle \exp(2\pi i \eta) & & \downarrow & \searrow j^* & \downarrow \\
 & & & RU(\mathbb{Z}/3) & \\
 \mathcal{U}_{\mathrm{Spin}}^6(BQ_8) & \xleftarrow{i^*} & \mathcal{U}_{\mathrm{Spin}}^6(B\mathrm{SL}(2, \mathbb{F}_3)) & \xleftarrow{\pi^*} & \mathcal{U}_{\mathrm{Spin}}^6(B\mathbb{Z}/3) \\
 & & \searrow j^* & \downarrow & \\
 & & & \mathcal{U}_{\mathrm{Spin}}^6(B\mathbb{Z}/3) &
 \end{array} \quad (4.17)$$

As groups, the entries in the ‘lower plane’ are

$$\begin{array}{ccccc}
 (\mathbb{Z}/2)^2 & \xleftarrow{i^*} & (\mathbb{Z}/2)^2 \times \mathbb{Z}/9 & \xleftarrow{\pi^*} & \mathbb{Z}/9 \\
 & & \searrow j^* & \downarrow & \\
 & & & \mathbb{Z}/9 &
 \end{array} \quad (4.18)$$

This diagram should be contrasted with the equivalent diagram (4.6) for the case $G = A_4$. In that case, the presence of maps π^* and j^* going in both directions, and the fact that the cobordism groups in the triangle were all isomorphic, guaranteed that π^* and j^* were isomorphisms. This meant that the anomalies for each irrep of A_4 could be deduced directly by decomposing to irreps of $\mathbb{Z}/3$, and computing the anomalies thereof. Here, the extra factor of $(\mathbb{Z}/2)^2$ in the cobordism group for $\mathrm{SL}(2, \mathbb{F}_3)$ prevents such a direct method, and we will need extra ingredients to complete our analysis.

Proceeding, we first need to know the maps between representation rings in the upper plane. We can restrict our attention to the four complex irreps, and the one pseudo-real irrep, since only these can have anomalies. The pullback of the complex representations of $\mathrm{SL}(2, \mathbb{F}_3)$ is

$$j^* : \mathbf{1}' \mapsto [1]_3, \quad \overline{\mathbf{1}'} \mapsto [2]_3, \quad \mathbf{2}' \mapsto [0]_3 \oplus [2]_3, \quad \overline{\mathbf{2}'} \mapsto [0]_3 \oplus [1]_3 \quad (4.19)$$

¹⁶The homological version of this anomaly interplay diagram was a crucial ingredient in our calculation of the bordism group $\Omega_5^{\mathrm{Spin}}(B\mathrm{SL}(2, \mathbb{F}_3))$, as recorded in appendix A.4.

under the subgroup embedding j . On the other hand, the pullback of complex representations of $\mathbb{Z}/3$ under the quotient map π is

$$\pi^* : [1]_3 \mapsto \mathbf{1}', \quad [2]_3 \mapsto \overline{\mathbf{1}}'. \quad (4.20)$$

Recall that the $[0]_3$ of $\mathbb{Z}/3$ is anomaly-free, while the $[1]_3$ and $[2]_3$ contribute 1 mod 9 and -1 mod 9 anomalies respectively.

While we observed above that the maps j^* and π^* are not isomorphisms, as they were in the $G = A_4$ example of section 4.1, the fact that $j^* \circ \pi^*$ is the identity map is still extremely useful; it uniquely fixes the order 9 part of the anomaly for every irrep of $\mathrm{SL}(2, \mathbb{F}_3)$. Thence, using the maps between representations above, we can swiftly extract the following necessary condition for anomaly cancellation:

$$n(\mathbf{1}') - n(\overline{\mathbf{1}}') - n(\mathbf{2}') + n(\overline{\mathbf{2}}') = 0 \bmod 9. \quad (4.21)$$

Moreover, we can deduce that the anomalies of the irreps $\mathbf{1}'$ and $\overline{\mathbf{1}}'$, which (by (4.20)) are the only non-trivial irreps in the image of π^* , contain no $\mathbb{Z}/2 \times \mathbb{Z}/2$ element because π^* is a homomorphism. Thus, so far we have computed the anomaly for all the 1-dimensional and 3-dimensional irreps of $\mathrm{SL}(2, \mathbb{F}_3)$.

Unlike the $G = A_4$ example, however, this analysis does not yet fix the anomaly completely; we must determine the $\mathbb{Z}/2 \times \mathbb{Z}/2$ part of the anomaly for all three of the 2-dimensional irreps. We need a new strategy to finish the task. It will turn out that the $\mathbb{Z}/2 \times \mathbb{Z}/2$ factor of $\Omega_5^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$ is not probed by the η -invariant, just like the case for the quaternion group Q_8 considered in section 4.2.

4.3.2 Anomaly interplay with Q_8 derived subgroup

To see why this is the case, we will first argue that $\exp(2\pi i \eta)_{\mathrm{SL}(2, \mathbb{F}_3)}(\mathbf{2})$ is trivial on any generator of the $\mathbb{Z}/2 \times \mathbb{Z}/2$ subgroup of $\Omega_5^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$, and so, by the fact that the irrep $\mathbf{2}$ is pseudo-real and can only have order 2 anomalies, it is anomaly-free. To proceed, let (X, f) denote a spin 5-manifold X equipped with a certain $\mathrm{SL}(2, \mathbb{F}_3)$ -bundle $f : X \rightarrow \mathrm{BSL}(2, \mathbb{F}_3)$, that is a generator of $\mathbb{Z}/2 \times \mathbb{Z}/2 \subset \Omega_5^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$. It turns out that (X, f') , where $f' : X \rightarrow BQ_8$ is naturally induced from f , is also a generator of $\Omega_5^{\mathrm{Spin}}(BQ_8) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$, as we show in appendix C. By the pullback $i^* : \mathcal{U}_{\mathrm{Spin}}^6(\mathrm{BSL}(2, \mathbb{F}_3)) \rightarrow \mathcal{U}_{\mathrm{Spin}}^6(BQ_8)$, we thus infer that

$$\exp(2\pi i \eta)_{\mathrm{SL}(2, \mathbb{F}_3)}(\mathbf{2}, X) = \exp(2\pi i \eta)_{Q_8}(\mathbf{2}, X), \quad (4.22)$$

where we used the character table (table 4) to deduce that the pullback of the irrep $\mathbf{2}$ of $\mathrm{SL}(2, \mathbb{F}_3)$ under the embedding i is the irrep $\mathbf{2}$ of Q_8 . The r.h.s. of this equation is trivial due to a result by Botvinnik and Gilkey [59], as used in section 4.2.

4.3.3 Anomaly interplay with $U(2)$: proof ‘by Standard Model’

All that remains is to finish computing the anomaly for the complex 2-dimensional irreps. To show that these do not contribute any order 2 anomaly, we now study an anomaly

interplay between $\mathrm{SL}(2, \mathbb{F}_3)$ and $\mathrm{U}(2)$. Recall that the group $\mathrm{SL}(2, \mathbb{F}_3)$ is a semi-direct product between the quaternion group Q_8 and $\mathbb{Z}/3$ where $\mathbb{Z}/3$ acts on Q_8 by cyclically permuting the quaternionic units i, j, k . More precisely, we can write

$$\mathrm{SL}(2, \mathbb{F}_3) = \langle i, j, \sigma | i^4 = \sigma^3 = 1, j^2 = i^2, j i j^{-1} = i^1, \sigma i \sigma^{-1} = j, \sigma j \sigma^{-1} = i j \rangle. \quad (4.23)$$

Using this, we now find an explicit embedding of $\mathrm{SL}(2, \mathbb{F}_3)$ as a subgroup of $\mathrm{U}(2)$, defining a group homomorphism $s : \mathrm{SL}(2, \mathbb{F}_3) \rightarrow \mathrm{U}(2)$ such that

$$s : i \mapsto \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad j \mapsto \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \sigma \mapsto e^{\pi i/12} \begin{pmatrix} 1 & -1 \\ -i & -i \end{pmatrix}. \quad (4.24)$$

It is straightforward to verify that the relations determining the structure of $\mathrm{SL}(2, \mathbb{F}_3)$ are all satisfied with this choice of embedding.

We do not need the full machinery of anomaly interplay between local and global anomalies used in ref. [38] because we only consider $\mathrm{U}(2)$ representations that are free of local anomalies. The only thing we will make use of here is that anomaly-free representations remain anomaly-free under the pullback. Let us denote an irrep of $\mathrm{U}(2) = (\mathrm{SU}(2) \times \mathrm{U}(1)) / (\mathbb{Z}/2)$ by (\mathbf{n}, q) , where \mathbf{n} is the irrep under the $\mathrm{SU}(2)$ subgroup and q is the $\mathrm{U}(1)$ charge satisfying $q - (n - 1) \in 2\mathbb{Z}$.

Consider Weyl fermions in a well-known anomaly-free representation of $\mathrm{U}(2)$, namely one generation of the Standard Model fermion content whose representation under the electroweak gauge group $\mathrm{U}(2)$ reads¹⁷

$$\mathbf{r}_{\mathrm{SM}} = (\mathbf{2}, 1)^{\oplus 3} \oplus (\mathbf{2}, -3) \oplus (\mathbf{1}, -4)^{\oplus 3} \oplus (\mathbf{1}, 2)^{\oplus 3} \oplus (\mathbf{1}, 6). \quad (4.25)$$

Restricting to the subgroup $\mathrm{SL}(2, \mathbb{F}_3)$ under the embedding s given in (4.24), the representation \mathbf{r}_{SM} decomposes in terms of the irreps of $\mathrm{SL}(2, \mathbb{F}_3)$ as (see appendix D)

$$s^* \mathbf{r}_{\mathrm{SM}} = \mathbf{2}'^{\oplus 3} \oplus \mathbf{2} \oplus \overline{\mathbf{1}}'^{\oplus 6} \oplus \mathbf{1}. \quad (4.26)$$

Note that in this representation, $n(\mathbf{1}') - n(\overline{\mathbf{1}}') - n(\mathbf{2}') + n(\overline{\mathbf{2}}') = -9$, consistent with the anomaly cancellation condition (4.21).

Now, let's evaluate the η -invariant in this representation on a generator X of $\mathbb{Z}/2 \times \mathbb{Z}/2 \subset \Omega_5^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$ (which is also a manifold equipped with a $\mathrm{U}(2)$ -bundle). Since we know that this representation is anomaly-free, we get

$$1 = \exp(2\pi i \eta)_{\mathrm{SL}(2, \mathbb{F}_3)}(s^* \mathbf{r}_{\mathrm{SM}}, X) = \exp(2\pi i \eta)_{\mathrm{SL}(2, \mathbb{F}_3)}(\mathbf{2}'^{\oplus 3}, X), \quad (4.27)$$

where we use the fact that all other irreps in $s^* \mathbf{r}_{\mathrm{SM}}$ has no order 2 anomaly in the last equality. This immediately tells us that the irrep $\mathbf{2}'$ and its conjugate are free of any anomaly of order 2.

¹⁷Here, local physics allows us to choose the global form of the gauge group to be $\mathrm{SU}(3) \times \mathrm{U}(2)$. For other possibilities of the gauge group for the algebra $\mathfrak{g} = \mathfrak{su}(3) \oplus \mathfrak{su}(2) \oplus \mathfrak{u}(1)$ and ways to distinguish between them, see e.g. ref. [60].

Putting everything together, we have proven that all representations of $\text{SL}(2, \mathbb{F}_3)$ have no order 2 anomaly, and moreover computed the anomalies associated with every irrep. Thus, the condition

$$n(\mathbf{1}') - n(\overline{\mathbf{1}'}') - n(\mathbf{2}') + n(\overline{\mathbf{2}'}') = 0 \bmod 9 \quad (\text{SL}(2, \mathbb{F}_3) \text{ anomaly cancellation in 4d}) \quad (4.21)$$

is both necessary and sufficient for $\text{SL}(2, \mathbb{F}_3)$ anomaly cancellation. The anomaly cancellation condition above is equivalent to saying that the non-anomalous representations are generated by the set

$$\mathbf{1}, \quad \mathbf{2}, \quad \mathbf{3}, \quad \mathbf{1}' \oplus \overline{\mathbf{1}'}', \quad \mathbf{2}' \oplus \overline{\mathbf{2}'}', \quad \mathbf{1}' \oplus \overline{\mathbf{2}'}', \quad \mathbf{1}'^{\oplus 9}, \quad (4.28)$$

which can be taken as a basis for $RU_0(\text{SL}(2, \mathbb{F}_3))$.

4.3.4 Example: an anomalous flavour physics model

One can apply our result to compute the anomaly in various flavour physics models that use an $\text{SL}(2, \mathbb{F}_3)$ symmetry. For an explicit example, we consider the model of Feruglio et al. [36] for quark and lepton masses. We denote the fermion representation there used, recorded in table 2 of ref. [36], by \mathbf{r}_{BSM} . The leptons are charged in triplet and singlet representations of $\text{SL}(2, \mathbb{F}_3)$, to mimic the construction of A_4 -based models for lepton masses and mixings. The doublet representations are also utilised, to unify the first two generations of quarks, while the third family quarks transform in singlets.

Within the leptons alone, one can check that anomalies cancel. But summing over the quarks, there are 12 left-handed Weyl fermions transforming in one of the complex doublet irreps, which we can take to be the irrep $\overline{\mathbf{2}'}'$ without loss of generality. From the results of this section, the anomaly is therefore

$$\exp(2\pi i \eta)_{\text{SL}(2, \mathbb{F}_3)}(\mathbf{r}_{\text{BSM}}) = \exp(2\pi i/3). \quad (4.29)$$

Interestingly, the anomaly is non-zero, corresponding to the element $3 \bmod 9 \in \mathcal{U}_6^{\text{Spin}}(B\text{SL}(2, \mathbb{F}_3))$. The $\text{SL}(2, \mathbb{F}_3)$ symmetry is therefore anomalous. Perhaps more interestingly, this is not the end of the story, as we shall see in section 5.4.1; it turns out this anomaly can be ‘truncated’ (in a sense soon to be defined) by passing to a larger symmetry group that surjects onto $\text{SL}(2, \mathbb{F}_3)$.

4.4 Dihedral group symmetry

Our last example is the symmetric group S_3 , whose anomaly is also analysed in ref. [42] by a different method. From the bordism perspective this example is especially simple, because the fact that

$$\Omega_5^{\text{Spin}}(BS_3) = 0 \quad (4.30)$$

means there can be no anomalies in 4d. To see this, recall that S_3 is isomorphic to the dihedral group of order 6, D_6 . Further, as we explain in appendix A.5, it can be shown that

$$\Omega_5^{\text{Spin}}(BD_{2n}) = 0 \quad (4.31)$$

for any integer n , so any such dihedral group symmetry (of which S_3 is a special case) is anomaly-free in 4d.

5 Not all anomalies are equal: anomaly-truncation by group extensions

In the previous section we computed the anomalies associated with all representations for certain non-abelian finite groups. But there is a subtle issue that we must confront when considering anomalies for finite groups. This is the issue of whether a finite group anomaly can be ‘truncated’ (by which we mean, very roughly, that the order of the (finite) anomaly for a given representation is reduced) by an extension of the symmetry. This idea was first appreciated, for the case of abelian finite groups, by Banks and Dine [41] (and discussed from the bordism perspective in [29]).

In section 5.1 we put this notion of ‘anomaly truncation’ on a rigorous footing. We then review the known example of Banks and Dine in section 5.2, recasting the analysis in our language, before considering examples with non-abelian finite symmetry groups in sections 5.3 and 5.4.

5.1 General formalism

Consider a group extension $f : H \twoheadrightarrow G$ of G . Since this is a surjection, for any element of G there is a corresponding element in H , and so we retain the original symmetry of the theory. One could then ask about the anomaly of the system of fermions considered as transforming in a representation of H instead of the original G .

As discussed in section 2, the homomorphism f induces the commutative diagram

$$\begin{array}{ccc} RU(G) & \xrightarrow{RU(f)} & RU(H) \\ \exp(2\pi i\eta)_G \downarrow & & \downarrow \exp(2\pi i\eta)_H \\ \mathcal{U}_{\text{Spin}}^6(BG) & \xrightarrow{\mathcal{U}(f)} & \mathcal{U}_{\text{Spin}}^6(BH) \end{array} \quad (5.1)$$

Clearly, an anomaly-free representation \mathbf{r} of G remains anomaly-free after getting pulled back to a representation of H since the commutativity implies that

$$\exp(2\pi i\eta)_H (RU(f)(\mathbf{r})) = \mathcal{U}(f) \circ \exp(2\pi i\eta)_G (\mathbf{r}) = 0. \quad (5.2)$$

It is also possible that this equation is satisfied for an anomalous representation \mathbf{r} of G . In other words, an anomalous representation may become anomaly-free after the group extension. In that case, we say that the anomaly is *truncated*, because the order of the image of $\exp(2\pi i\eta)_H \circ RU(f)$ is generally smaller than the order of the image of $\exp(2\pi i\eta)_G$.

This anomaly truncation can be quantified as follows. Recall that

$$RU_0(G) = \ker(\exp(2\pi i\eta)_G) \subset RU(G) \quad (5.3)$$

is the subgroup of anomaly-free representations inside $RU(G)$. We can similarly define the subgroup of representations of G that become anomaly-free after being pulled back along f to be

$$RU_0(G; H, f) := \ker(\mathcal{U}(f) \circ \exp(2\pi i\eta)_G) \subset RU(G). \quad (5.4)$$

Clearly, $RU_0(G)$ is a subgroup of $RU_0(G; H, f)$ since all anomaly-free representations are pulled back to anomaly-free representations. Thus, the anomaly truncation occurs when the quotient

$$Q(G; H, f) = RU_0(G; H, f) / RU_0(G) \quad (5.5)$$

is non-trivial. We call $Q(G; H, f)$ the *anomaly truncation quotient*.

5.2 Known example: abelian finite anomalies

To give this rather formal description some teeth, we start by revisiting the known example of anomaly truncation for a finite abelian group [29, 41].

For concreteness, let us consider a 4d theory with symmetry $G = \mathbb{Z}/3$, that we would like to interpret as a low-energy effective field theory. As seen in section 4.1, the anomalies are captured by the cobordism group $\mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3) \cong \Omega_5^{\text{Spin}}(B\mathbb{Z}/3) \cong \mathbb{Z}/9$, which we take to be generated by the exponentiated η -invariant for the fundamental $[1]_3$ representation. For a generic set of LH Weyl fermions the condition for $\mathbb{Z}/3$ anomaly cancellation is [29].

$$n([1]_3) - n([2]_3) = 0 \bmod 9. \quad (5.6)$$

This can be derived, for example, by embedding $\mathbb{Z}/3 \hookrightarrow \text{U}(1)$. From that perspective, the mod 9 condition (5.6) comes from the cubic $\text{U}(1)$ anomaly cancellation condition (which implies the ‘coarser’ linear condition, $n([1]_3) - n([2]_3) = 0 \bmod 3$).

Now consider extending $\mathbb{Z}/3$ via

$$\mathbb{Z}/3 \xhookrightarrow{\alpha} \mathbb{Z}/9 \xrightarrow{p} \mathbb{Z}/3. \quad (5.7)$$

The pullback of representations is simply multiplication of charges by three, *viz.*

$$p^* : RU(\mathbb{Z}/3) \rightarrow RU(\mathbb{Z}/9) : [q]_3 \mapsto [3q]_9. \quad (5.8)$$

Physically, the key point is that this extension of the symmetry group from $\mathbb{Z}/3$ to $\mathbb{Z}/9$ is undetectable from the low-energy perspective. Indeed, there may lurk UV states with discrete charges that are finer multiples of the charges of the low-energy degrees of freedom (in this case e.g. $q_{UV} = [1]_9$), such that it is $\mathbb{Z}/9$ (or indeed any extension of $\mathbb{Z}/3$) that acts faithfully in the UV.

For a general $\mathbb{Z}/9$ gauge theory, with a set of LH Weyls in the representation $\bigoplus_i [Q_i]_9$ of $\mathbb{Z}/9$, let us define the anomaly coefficients $\mathcal{A}_{\text{cub}} := \sum_i Q_i^3$ and $\mathcal{A}_{\text{grav}} := \sum_i Q_i$ (so named because they pullback from the cubic and mixed-gravitational anomalies under embedding $\mathbb{Z}/9$ in $\text{U}(1)$). The anomaly associated to this fermion spectrum is [29]

$$\left(\mathcal{A}_{\text{cub}} \bmod 27, \frac{1}{3} (\mathcal{A}_{\text{cub}} - \mathcal{A}_{\text{grav}}) \bmod 3 \right) \in \mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/9) \cong \mathbb{Z}/27 \times \mathbb{Z}/3. \quad (5.9)$$

A set of LH fermions in the representation $\bigoplus_i [q_i]_3$ of $\mathbb{Z}/3$ pulls back to $\bigoplus_i [3q_i]_9$ of $\mathbb{Z}/9$. Clearly, the $\mathbb{Z}/27$ -valued element of the anomaly always vanishes after this pullback, since $(3q_i)^3 = 27q_i^3 \implies \mathcal{A}_{\text{cub}} = 0 \bmod 27$. This leaves only the order 3 anomaly, which vanishes if and only if $\sum_i q_i = 0 \bmod 3$, which is equivalent to the condition

$$n([1]_3) - n([2]_3) = 0 \bmod 3 \quad (5.10)$$

for fermions in the original $\mathbb{Z}/3$ theory. By comparison with the original condition (5.6), we see that the order of the anomaly is truncated from 9 to 3 by the symmetry extension.

We can state the anomaly truncation more clearly as the enlargement of the subgroup of anomaly-free representations. Originally, $RU_0(\mathbb{Z}/3)$ is generated by $[0]_3$, $[1]_3 \oplus [2]_3$, and

$[1]_3^{\oplus 9}$. After the group extension to $\mathbb{Z}/9$, the subgroup of anomaly-free representations becomes $RU_0(\mathbb{Z}/3)'$, now generated by $[0]_3$, $[1]_3 \oplus [2]_3$, and $[1]_3^{\oplus 3}$. Thus, the anomaly truncation quotient is

$$Q(\mathbb{Z}/3; \mathbb{Z}/9, p) = \frac{\mathbb{Z} \langle [1]_3^{\oplus 3} \rangle}{\mathbb{Z} \langle [1]_3^{\oplus 9} \rangle} \cong \mathbb{Z}/3. \quad (5.11)$$

Note that the remaining condition (5.10) is itself the result of pulling back the linear mixed $U(1)$ -gravitational anomaly along $\mathbb{Z}/9 \hookrightarrow U(1)$. Since that condition is *linear* in the charges, it is invariant under further symmetry extensions of $\mathbb{Z}/9$ by any \mathbb{Z}/r , which are tantamount simply to rescalings of all charges by r .

At least in this example, where G is abelian, the anomaly truncation is equivalent to cancelling the anomaly by coupling to a topological quantum field theory (TQFT), as discussed e.g. in ref. [16] (section 4.6.2). The TQFT discussed therein has the effect of restricting the allowed $\mathbb{Z}/3$ bundles to those that lift to $\mathbb{Z}/9$ bundles. It will be interesting to see if this version of events, in which anomaly truncation via symmetry extension is explicitly realised by coupling to a TQFT, carries over to the case where G is non-abelian, as in the examples we consider next. We will explore this more in future work.

5.3 Extending A_4 by $\mathbb{Z}/3$

Let us return to the $G = A_4$ anomalies in 4d, for which we have derived in section 4.1 that the non-trivial singlet irreps $\mathbf{1}'$ and $\overline{\mathbf{1}}'$ contribute anomalies of $\pm 1 \bmod 9$. The anomaly-free subgroup of A_4 was identified to be the derived subgroup $A'_4 \cong \mathbb{Z}/2 \times \mathbb{Z}/2$, with the anomaly ‘coming from’ its abelianisation $A_4/A'_4 \cong \mathbb{Z}/3$. Now, as we reviewed in section 5.2 for the case of $\mathbb{Z}/3$ anomalies in 4d, the order 9 anomaly is truncated to an order 3 anomaly by extending the symmetry from $\mathbb{Z}/3$ to $p : \mathbb{Z}/9 \rightarrow \mathbb{Z}/3$. As discussed, such a symmetry extension is undetectable from the low-energy perspective, and so it is only the reduced mod 3 condition that *must* be satisfied by the $\mathbb{Z}/3$ -charged IR degrees of freedom.

Since the order 9 gauge anomaly associated with the $\mathbf{1}'$ irrep of $G = A_4$ comes from its $\mathbb{Z}/3$ subgroup, in a precise sense captured by the anomaly interplay maps (4.6), it is plausible that this order 9 anomaly could also be truncated to, say, an order 3 anomaly by some finite group extension of A_4 . By analogy with the $G = \mathbb{Z}/3$ case above, we consider the extension

$$0 \rightarrow \mathbb{Z}/3 \rightarrow K \rightarrow A_4 \rightarrow 0, \quad (5.12)$$

of A_4 by $\mathbb{Z}/3$, where K is the semi-direct product

$$K = (\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/9. \quad (5.13)$$

The injective map from $\mathbb{Z}/3$ takes the element $1 \bmod 3 \in \mathbb{Z}/3$ to the element $((0 \bmod 2, 0 \bmod 2), 3 \bmod 9)$ of $(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/9$. The resulting group K can also be described as a different group extension,

$$0 \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2 \rightarrow K \rightarrow \mathbb{Z}/9 \rightarrow 0, \quad (5.14)$$

which splits because K is a semi-direct product. Armed with both these extensions, we know that K sits at the centre of the following commutative diagram of short exact

sequences:

$$\begin{array}{ccccc}
 0 & \longrightarrow & \mathbb{Z}/2 \times \mathbb{Z}/2 & \xlongequal{\quad} & \mathbb{Z}/2 \times \mathbb{Z}/2 \\
 \downarrow & & \downarrow i' & & \downarrow i \\
 \mathbb{Z}/3 & \xhookrightarrow{\alpha'} & K & \xrightarrow{p'} & A_4 \\
 \parallel & & \downarrow \pi' & & \downarrow \pi \\
 \mathbb{Z}/3 & \xhookrightarrow{\alpha} & \mathbb{Z}/9 & \xrightarrow{p} & \mathbb{Z}/3
 \end{array} \quad . \quad (5.15)$$

Importantly for the calculation that follows, the fact that both the middle and right columns are in fact split extensions means there are two maps ‘going the other way’, labelled $j : \mathbb{Z}/3 \rightarrow A_4$ (a map we have already used extensively above) and $j' : \mathbb{Z}/9 \rightarrow K$, satisfying $j \circ \pi = \text{id}$ and $j' \circ \pi' = \text{id}$.¹⁸

For every one of the maps in the commutative diagram (5.15), functorality implies an induced pullback on both the representation ring $RU(\cdot)$ and cobordism $\mathcal{U}_{\text{Spin}}^6(B\cdot)$, with commuting squares involving the exponentiated η -invariant that maps $RU(\cdot) \rightarrow \mathcal{U}_{\text{Spin}}^6(B\cdot)$, as drawn in (2.10). For example, there is a commuting square:

$$\begin{array}{ccc}
 RU(A_4) & \xrightarrow{p'^*} & RU(K) \\
 \exp(2\pi i \eta)_{A_4} \downarrow & & \downarrow \exp(2\pi i \eta)_K \\
 \mathcal{U}_{\text{Spin}}^6(BA_4) & \xrightarrow{p'^*} & \mathcal{U}_{\text{Spin}}^6(BK)
 \end{array} \quad (5.16)$$

If we focus on the maps in the bottom right square of eq. (5.15), this gives rise to the following commutative ‘cube’,

$$\begin{array}{ccccc}
 & & RU(K) & & \\
 & \nearrow p'^* & & \nwarrow j'^* & \\
 RU(A_4) & & & & RU(\mathbb{Z}/9) \\
 & \searrow j^* & & \nearrow p^* & \\
 & & RU(\mathbb{Z}/3) & & \\
 & \nearrow p'^* & & \nwarrow j'^* & \\
 \mathcal{U}_{\text{Spin}}^6(BA_4) & & \mathcal{U}_{\text{Spin}}^6(BK) & & \mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/9) \\
 & \searrow j^* & & \nearrow p^* & \\
 & & \mathcal{U}_{\text{Spin}}^6(B\mathbb{Z}/3) & &
 \end{array} \quad (5.17)$$

where each ‘vertical face’ is a commutative square of the kind (2.10).

¹⁸Note that the middle row, which describes K as an extension of A_4 , cannot split; if it did, that would imply a right-to-left map from $\mathbb{Z}/3$ to $\mathbb{Z}/9$ in the bottom row whose composition with p is the identity, meaning that the bottom row splits also — which it does not.

Now, we wish to follow the fate of the representation $\mathbf{1}' \in RU(A_4)$, to see if its anomaly is truncated by passing to the extension K . Our computation of section 4.1 means that

$$\exp(2\pi i \eta)_{A_4} : RU(A_4) \rightarrow \mathcal{U}_{\text{Spin}}^6(BA_4) : \mathbf{1}' \mapsto 1 \bmod 9. \quad (5.18)$$

We wish to compute $\exp(2\pi i \eta)_K(p'^*(\mathbf{1}')) \in \mathcal{U}_{\text{Spin}}^6(BK)$, i.e. the anomaly associated with the pullback of the $\mathbf{1}' \in RU(A_4)$ to a K -representation. This equals $p'^*(\exp(2\pi i \eta)_{A_4}(\mathbf{1}'))$ by commutativity (of the ‘back left’ face of (5.17)). To find this anomaly pullback, we use our knowledge of the anomaly interplay maps j^* and p^* in the lower face of (5.17). Recalling $j^*(\mathbf{1}') = [1]_3 \in RU(\mathbb{Z}/3)$, we have

$$j^* : [\exp(2\pi i \eta)_{A_4}(\mathbf{1}') = \exp(2\pi i/9)] \mapsto [\exp(2\pi i \eta)_{\mathbb{Z}/3}([1]_3) = \exp(2\pi i/9)], \quad (5.19)$$

using the results of section 4.1. Now, using $p^*([1]_3) = [3]_9 \in RU(\mathbb{Z}/9)$, we have

$$p^* : [\exp(2\pi i \eta)_{\mathbb{Z}/3}([1]_3) = \exp(2\pi i/9)] \mapsto [\exp(2\pi i \eta)_{\mathbb{Z}/9}([3]_9) = \exp(2\pi i/3)], \quad (5.20)$$

by the calculation in section 5.2.

Thus, if we restrict to the image of the anomalous representation $\mathbf{1}' \in RU(A_4)$ in the lower square, we have

$$\begin{array}{ccc}
 & ? & \\
 p'^* \nearrow & & \nwarrow j'^* \\
 1 \bmod 9 & & 1 \bmod 3 \\
 j^* \searrow & & \nearrow p^* \\
 & 1 \bmod 9 &
 \end{array}
 \quad (5.21)$$

where the question mark denotes $\exp(2\pi i \eta)_K(p'^*(\mathbf{1}'))$, the anomaly that we wish to pin down. Thanks to the existence of both maps j'^* and π'^* satisfying $j'^* \circ \pi'^* = \text{id}$ (i.e. thanks to the splitting of $\mathbb{Z}/2 \times \mathbb{Z}/2 \hookrightarrow K \twoheadrightarrow \mathbb{Z}/9$), we infer that the unknown anomaly phase is

$$\exp(2\pi i \eta)_K(p'^*(\mathbf{1}')) = \exp(2\pi i/3). \quad (5.22)$$

Thus, by passing to the extension K , we see that the order 9 anomaly associated with the non-trivial singlets of A_4 can be truncated to an order 3 anomaly.

Alternatively, we could phrase this result in terms of the anomaly-free representations. After the group extension, the subgroup $RU_0(A_4)$ of anomaly-free representations becomes $RU_0(A_4; K, p')$, now generated by

$$\mathbf{1}, \quad \mathbf{3}, \quad \mathbf{1}' \oplus \overline{\mathbf{1}'}, \quad \mathbf{1}'^{\oplus 3}. \quad (5.23)$$

By taking the quotient with $RU_0(A_4)$ given in (4.10), it can be easily seen that the anomaly truncation quotient $Q(A_4; K, p')$ is isomorphic to $\mathbb{Z}/3$, showing that the order of the anomaly is ‘down’ by a third.

We emphasise that, to calculate this anomaly, we did not even need to know the pullback representation $p'^*(\mathbf{1}') \in RU(K)$, nor did we need to know the cobordism group $\mathcal{U}_{\text{Spin}}^6(BK)$. The ‘gridlock’ properties of so many commutative squares, thanks to the two different short exact sequences for K , are powerful enough to completely fix the value of the anomaly that we wanted.

5.4 Extending $\mathrm{SL}(2, \mathbb{F}_3)$ by $\mathbb{Z}/3$

Let's now consider the symmetry group $\mathrm{SL}(2, \mathbb{F}_3)$ and explore a possibility of anomaly truncation by group extension. First recall that, as shown in section 4.3, the only anomalies from Weyl fermions in representations of this symmetry group are order 9 anomalies, which directly comes from the order 9 anomalies associated with its $\mathbb{Z}/3$ subgroup. It is thus probable that these order 9 anomalies can be truncated in a similar manner to that of the symmetry group A_4 discussed in the previous subsection.

We extend $\mathrm{SL}(2, \mathbb{F}_3)$ non-trivially by $\mathbb{Z}/3$ to the group

$$L = Q_8 \rtimes \mathbb{Z}/9 \quad (5.24)$$

that fits inside the commutative diagram

$$\begin{array}{ccccc} 0 & \longrightarrow & Q_8 & \xlongequal{\quad} & Q_8 \\ \downarrow & & \downarrow i' & & \downarrow i \\ \mathbb{Z}/3 & \xleftarrow{\alpha'} & L & \xrightarrow{p'} & \mathrm{SL}(2, \mathbb{F}_3) \\ \parallel & & \downarrow \pi' & & \downarrow \pi \\ \mathbb{Z}/3 & \xleftarrow{\alpha} & \mathbb{Z}/9 & \xrightarrow{p} & \mathbb{Z}/3 \end{array} \quad (5.25)$$

where the injective map α' maps $1 \bmod 3$ in $\mathbb{Z}/3$ to $(0, 3 \bmod 9)$ in L , and α is multiplication by 3. Again, applying the functors $RU(\cdot)$ and $\mathcal{U}_{\mathrm{Spin}}^6(\cdot)$ to the lower-right square of the grid (5.25), we obtain two commutative squares linked together by the natural transformation $\exp(2\pi i \eta)$ to form a commutative cube

$$\begin{array}{ccccc} & & RU(L) & & \\ & \nearrow p'^* & & \nwarrow j'^* & \\ RU(\mathrm{SL}(2, \mathbb{F}_3)) & & & & RU(\mathbb{Z}/9) \\ & \searrow j^* & & \nearrow p^* & \\ & & RU(\mathbb{Z}/3) & & \\ & & \downarrow & & \\ & & \mathcal{U}_{\mathrm{Spin}}^6(BL) & & \\ & \nearrow p'^* & & \nwarrow j'^* & \\ \mathcal{U}_{\mathrm{Spin}}^6(B\mathrm{SL}(2, \mathbb{F}_3)) & & & & \mathcal{U}_{\mathrm{Spin}}^6(B\mathbb{Z}/9) \\ & \searrow j^* & & \nearrow p^* & \\ & & \mathcal{U}_{\mathrm{Spin}}^6(B\mathbb{Z}/3) & & \end{array} \quad (5.26)$$

As the irrep $\mathbf{1}' \in RU(\mathrm{SL}(2, \mathbb{F}_3))$ is pulled back by j^* to the irrep $[1]_3 \in RU(\mathbb{Z}/3)$, exactly as in section 5.3, the argument for finding the anomaly $\exp(2\pi i \eta)_L(p'^*(\mathbf{1}')) \in$

$\mathcal{U}_{\text{Spin}}^6(BL)$ goes exactly the same way. Thus, we obtain

$$\exp(2\pi i \eta)_L(p'^*(\mathbf{1}')) = \exp(2\pi i/3). \quad (5.27)$$

We now repeat the argument to find the anomaly of the irrep $\mathbf{2}' \in RU(\text{SL}(2, \mathbb{F}_3))$ pulled back to the representation $p'^*(\mathbf{2}')$ of L . Under various maps on the top square of the commutative cube (5.26), we get

$$\begin{array}{ccccc}
 & & p'^*(\mathbf{2}') & & \\
 & \nearrow p'^* & & \nwarrow j'^* & \\
 \mathbf{2}' & & & & [0]_9 \oplus [6]_9 \\
 & \searrow j^* & & \nearrow \pi'^* & \\
 & & [0]_3 \oplus [2]_3 & & \\
 & \nwarrow \pi^* & & \nearrow p^* &
 \end{array} \quad (5.28)$$

where we find $p^*([2]_3) = [6]_9$ by requiring that the action of $1 \bmod 3 \in \mathbb{Z}/3$ in the representation $[2]_3$, given by $\exp(4\pi i/3)$, is the same as the action of $1 \bmod 9 \in \mathbb{Z}/9$ in the $p^*([1]_3) = [q]_9$ representation given by $\exp(2\pi i q/9)$. Under the natural transformation, this commutative square is pushed downward to the bottom square of the cube (5.26) that reads

$$\begin{array}{ccccc}
 & & ? & & \\
 & \nearrow p'^* & & \nwarrow j'^* & \\
 -1 \bmod 9 & & & & -1 \bmod 3 \\
 & \searrow j^* & & \nearrow \pi'^* & \\
 & & -1 \bmod 9 & & \\
 & \nwarrow \pi^* & & \nearrow p^* &
 \end{array} \quad (5.29)$$

where the question mark denotes $\exp(2\pi i \eta)_L(p'^*(\mathbf{2}'))$, the anomaly that we wish to compute. Since $j'^* \circ \pi'^* = \text{id}$ due to the semi-direct product structure of L , we can infer from this commutative square that

$$\exp(2\pi i \eta)_L(p'^*(\mathbf{2}')) = \exp(-2\pi i/3), \quad (5.30)$$

and the anomaly cancellation condition (4.21) is truncated to

$$n(\mathbf{1}') - n(\overline{\mathbf{1}}') - n(\mathbf{2}') + n(\overline{\mathbf{2}}') = 0 \bmod 3. \quad (5.31)$$

Equivalently, after the group extension, the subgroup of anomaly-free representations becomes $RU_0(\text{SL}(2, \mathbb{F}_3); L, p')$ generated by

$$\mathbf{1}, \quad \mathbf{2}, \quad \mathbf{3}, \quad \mathbf{1}' \oplus \overline{\mathbf{1}}', \quad \mathbf{1}' \oplus \overline{\mathbf{1}}', \quad \mathbf{1}' \oplus \overline{\mathbf{2}}', \quad \mathbf{1}'^{\oplus 3}. \quad (5.32)$$

Comparing this with the original $RU_0(\text{SL}(2, \mathbb{F}_3))$ whose generators are given in (4.28), the anomaly truncation quotient is again isomorphic to $\mathbb{Z}/3$, indicating that the order of the anomaly is truncated by a third.

5.4.1 Example (continued): anomaly truncation in a flavour physics model

Recall that in section 4.3.4 we computed the anomaly in an $\mathrm{SL}(2, \mathbb{F}_3)$ symmetry for a fermion representation $\mathbf{r}_{\mathrm{BSM}}$ utilised in a particular flavour physics model [36]. The anomaly was found to be the phase $\exp(2\pi i/3)$, corresponding to an order 3 anomaly in the group $\mathcal{U}_6^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$.

We have just shown that all order 3 anomalies in $\mathcal{U}_6^{\mathrm{Spin}}(\mathrm{BSL}(2, \mathbb{F}_3))$ map to zero under the pullback induced by the $\mathbb{Z}/3$ group extension $p' : L \twoheadrightarrow \mathrm{SL}(2, \mathbb{F}_3)$. In our notation, we can write

$$\mathbf{r}_{\mathrm{BSM}} \notin \mathrm{RU}_0(\mathrm{SL}(2, \mathbb{F}_3)), \quad \text{but} \quad (5.33)$$

$$\mathbf{r}_{\mathrm{BSM}} \in \mathrm{RU}_0(\mathrm{SL}(2, \mathbb{F}_3); L, p'). \quad (5.34)$$

Physically, this means that, while the theory as set out in [36] is anomalous *per se*, it becomes anomaly-free (for the same fermion content) by considering an enlargement of the symmetry to the $\mathbb{Z}/3$ extension $p' : L \twoheadrightarrow \mathrm{SL}(2, \mathbb{F}_3)$, which is otherwise physically indistinguishable at low energies from $\mathrm{SL}(2, \mathbb{F}_3)$.

6 Summary

In this work, we investigated global anomalies of 4d chiral fermions charged under some finite non-abelian symmetry groups G , using the rigorous tool of the cobordism classification. Motivated in part by their relevance to flavour physics, we considered as examples the alternating group A_4 , the quaternion group Q_8 , the special linear group $\mathrm{SL}(2, \mathbb{F}_3)$, and the symmetric group S_3 .

For each of these groups, we determined the associated bordism group $\Omega_5^{\mathrm{Spin}}(BG)$ that classifies all possible anomalies, and then evaluated the η -invariant to precisely obtain the anomalies for chiral fermions charged in each representation of these groups. Our methods can be adapted to many other finite groups of interest.

When there is a non-trivial anomaly, the order of the anomaly can be further ‘truncated’ to an effective anomaly of lower finite order by passing to an extension of the original symmetry. After putting this notion of ‘anomaly truncation’ on a mathematically sound footing, we worked out the possible anomaly truncations for all the cases that we considered. We conclude this paper by summarising our specific results.

The quaternion group Q_8 and the dihedral groups D_{2n} (with a special case $S_3 = D_6$) are anomaly-free. There are order 9 anomalies for the groups A_4 and $\mathrm{SL}(2, \mathbb{F}_3)$ due to chiral fermions in complex representations of these groups; fermions in the real or pseudo-real representations do not contribute to any anomaly. The anomalies can be concisely expressed as anomaly cancellation conditions. Let $n(\mathbf{r})$ denote the number of left-handed fermions in the representation \mathbf{r} minus the number of right-handed fermions in the same representation. Then the anomaly cancellation condition reads

$$n(\mathbf{1}') - n(\overline{\mathbf{1}'}) = 0 \bmod 9 \quad (6.1)$$

for the group A_4 , and

$$n(\mathbf{1}') - n(\overline{\mathbf{1}'}) - n(\mathbf{2}') + n(\overline{\mathbf{2}'}) = 0 \bmod 9 \quad (6.2)$$

for the group $\mathrm{SL}(2, \mathbb{F}_3)$. The labels for the irreps are those listed in the character tables 2 and 4. As an example, we show that the flavour physics model in ref. [36] has an order 3 anomaly.

Both A_4 and $\mathrm{SL}(2, \mathbb{F}_3)$ can be extended by $\mathbb{Z}/3$ to $(\mathbb{Z}/2 \times \mathbb{Z}/2) \rtimes \mathbb{Z}/9$ and $Q_8 \rtimes \mathbb{Z}/9$, respectively, with the effect of truncating these order 9 anomalies down to order 3 anomalies. The anomaly cancellation conditions above remain the same apart from being taken modulo 3 instead of modulo 9. Under such a symmetry extension, the flavour physics model of ref. [36] thus becomes anomaly-free.

Acknowledgments

We are grateful to Arun Debray for divulging some helpful tricks for computing spectral sequences. JD has received funding from the European Research Council (ERC) under the European Union’s Horizon 2020 research and innovation programme under grant agreement 833280 (FLAY), and by the Swiss National Science Foundation (SNF) under contract 200020-204428. BG is supported by STFC consolidated grant ST/T000694/1. NL is supported by an STFC consolidated grant in ‘Particles, Strings and Cosmology’, and by the Royal Society.

A Spin bordism groups $\Omega_d^{\mathrm{Spin}}(BG)$ for various non-abelian finite group G

In this appendix, we collect facts about the non-abelian discrete groups considered in the main part of the paper.

There is an isomorphism between the spin bordism group and the connective real K-theory,

$$\Omega_d^{\mathrm{Spin}}(X) \cong ko_d(X), \quad (\text{A.1})$$

when $d \leq 7$, by virtue of the Anderson-Brown-Peterson theorem [61] which proves that the Atiyah-Bott-Shapiro map [62] $MT\mathrm{Spin} \rightarrow ko$ is 7-connected [30]. Here, ko is the connective real K-theory spectrum and $MT\mathrm{Spin}$ is the stable version of the Madsen-Tillmann spectrum of the spin group which represent the ko -homology and the spin bordism, respectively, according to Brown’s representability theorem [63, 64]. There are already many computations and results on the connective real K-theory of a space X when X is the classifying space of a discrete group, which are nicely summarised in ref. [57]. The bordism groups quoted in this appendix will be mainly from this reference.

When there is an ambiguity in the existing ko -computations, or when there is no result at all, we make use of both the Atiyah-Hirzebruch spectral sequence (AHSS) and the Adams spectral sequence (ASS). For an introduction to the use of these spectral sequences for calculating bordism groups in physics context, see e.g. refs. [16, 17] for the AHSS, and ref. [65] for the ASS.

d	0	1	2	3	4	5	6
$\tilde{\Omega}_d^{\text{Spin}}(B\mathbb{Z}/2 \times B\mathbb{Z}/2)$	0	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^3$	$\mathbb{Z}/4 \times (\mathbb{Z}/8)^2$	$(\mathbb{Z}/2)^2$	0	$\mathbb{Z}/2$

Table 5. The reduced spin bordism groups of $B\mathbb{Z}/2 \times B\mathbb{Z}/2$ up to degree 6.

A.1 $G = \mathbb{Z}/2 \times \mathbb{Z}/2$

The ko -homology groups for BG when $G = \mathbb{Z}/2 \times \mathbb{Z}/2$ were first computed in ref. [58] (which computed the ko -homology groups more generally for BG when $G = (\mathbb{Z}/2)^k$). The calculations were also used in physics literature in recent years (see e.g. [66]).

We first decompose $B\mathbb{Z}/2 \times B\mathbb{Z}/2$ into a wedge sum of different factors as

$$B\mathbb{Z}/2 \times B\mathbb{Z}/2 = (B\mathbb{Z}/2) \vee (B\mathbb{Z}/2) \vee (B\mathbb{Z}/2 \wedge B\mathbb{Z}/2)$$

and get

$$\widetilde{ko}_d(B\mathbb{Z}/2 \times B\mathbb{Z}/2) \cong \widetilde{ko}_d(B\mathbb{Z}/2) \oplus \widetilde{ko}_d(B\mathbb{Z}/2) \oplus \widetilde{ko}_d((B\mathbb{Z}/2)^{\wedge 2}), \quad (\text{A.2})$$

where we use the notation

$$X^{\wedge k} = \underbrace{X \wedge \dots \wedge X}_{k \text{ times}} \quad (\text{A.3})$$

to denote the k -fold wedge sum of X . $\widetilde{ko}_d((B\mathbb{Z}/2)^{\wedge k})$ were calculated in ref. [58] to be

$$\widetilde{ko}_d((B\mathbb{Z}/2)^{\wedge k}) \cong \begin{cases} \mathbb{Z}/2 \times (\mathbb{Z}/2)^{C(k,d)}, & d = 8l + 1, \quad k < 4l + 2, \\ & d = 8l + 2, \quad k < 4l + 3, \\ \mathbb{Z}/2^{4l+4-k} \times (\mathbb{Z}/2)^{C(k,d)}, & d = 8l + 3, \quad k < 4l + 4, \\ \mathbb{Z}/2^{4l+5-k} \times (\mathbb{Z}/2)^{C(k,d)}, & d = 8l + 7, \quad k < 4l + 5, \\ (\mathbb{Z}/2)^{C(k,d)}, & \text{otherwise.} \end{cases} \quad (\text{A.4})$$

Here, $C(k, d)$ denotes the coefficient of the monomial t^d in the infinite series

$$P_k(t) = \frac{t^k (1 - (1 - t)^{k-1}) Q_k(t)}{(1 - t^4)(1 + t^3)(1 - t)^{k-1}},$$

where

$$Q_k(t) = \begin{cases} t^{k-1}, & k = 0, 1 \bmod 4, \\ t^{k-2} (1 + t - t^2 + t^3 - t^5), & k = 2 \bmod 4, \\ t^{k-3} (1 + t^3 - t^5), & k = 3 \bmod 4. \end{cases}$$

Combining all the results, specialising to $k = 2$ and using the isomorphism between Ω^{Spin} and ko , the reduced spin bordism groups of $B\mathbb{Z}/2 \times B\mathbb{Z}/2$ are given in table 5 (see also figure 12.3.3 of ref. [57]).

A.2 $G = A_4$

Consider A_4 , the alternating group of order 4. Its derived subgroup A'_4 is $\mathbb{Z}/2 \times \mathbb{Z}/2$, giving rise to the short exact sequence

$$0 \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2 \rightarrow A_4 \rightarrow \mathbb{Z}/3 \rightarrow 0. \quad (\text{A.5})$$

and take X to be its classifying space BG . As outlined above, we obtain the spin bordism groups X through the isomorphism with the ko -homology groups of X , whose calculations can be found in more quantities in mathematics literature. $ko_d(BA_4)$ have been computed in ref. [57], with results nicely summarised in section 7.7.E. therein. Using the isomorphism to write their results as spin bordism and leaving out the factors corresponding to the spin bordism groups of a point, $\tilde{\Omega}_d^{\text{Spin}}(\text{pt})$, the reduced spin bordism groups of BA_4 are given by

$$\begin{aligned} \tilde{\Omega}_1^{\text{Spin}}(BA_4) &\cong \mathbb{Z}/3, \\ \tilde{\Omega}_2^{\text{Spin}}(BA_4) &\cong \mathbb{Z}/2, \\ \tilde{\Omega}_3^{\text{Spin}}(BA_4) &\cong \mathbb{Z}/12, \\ \tilde{\Omega}_4^{\text{Spin}}(BA_4) &= 0, \\ \tilde{\Omega}_5^{\text{Spin}}(BA_4) &\cong \mathbb{Z}/9, \\ \tilde{\Omega}_6^{\text{Spin}}(BA_4) &\cong \mathbb{Z}/2. \end{aligned} \quad (\text{A.6})$$

A.3 The quaternion group $G = Q_8$

Let's now consider the quaternion group Q_8 , a multiplicative group of order 8 whose elements are the basic quaternions $1, i, j, k$ together with their additive inverses $-1, -i, -j, -k$. Its derived subgroup Q'_8 is isomorphic to $\mathbb{Z}/2$, and Q_8/Q'_8 is isomorphic to the Klein four-group, putting Q_8 inside the short exact sequence

$$0 \rightarrow \mathbb{Z}/2 \rightarrow Q_8 \rightarrow \mathbb{Z}/2 \times \mathbb{Z}/2 \rightarrow 0. \quad (\text{A.7})$$

The calculations for $\widetilde{ko}_\bullet(BQ_8)$ were carried out in ref. [67]. The authors of [67] first decomposed BQ_8 stably as [68]

$$BQ_8 \cong BSL(2, \mathbb{F}_3) \vee \Sigma^{-1} (BS^3/BN) \vee \Sigma^{-1} (BS^3/BN), \quad (\text{A.8})$$

again localised at the prime 2. Here $BSL(2, \mathbb{F}_3)$ is the classifying space of $SL(2, \mathbb{F}_3)$, the group of 2×2 unit determinant matrices with entries in the finite field \mathbb{F}_3 , while N is the normaliser of the maximal torus in S^3 (whose group structure is inherited from the quaternion multiplication, and can be thought of as the group $SU(2)$). They then calculated the ko theory (localised at 2) for each factor, as given in eqs. (A.9) and (A.10) with $n \geq 0$:

$$\widetilde{ko}_{\epsilon+8n}BSL_2(\mathbb{F}_3) \cong \begin{cases} \mathbb{Z}/2^{4n+3} \oplus \mathbb{Z}/2^{2n}, & \epsilon = 3, \\ \mathbb{Z}/2, & \epsilon = 4, 6, 8, 10, \\ \mathbb{Z}/2 \oplus \mathbb{Z}/2, & \epsilon = 5, 9, \\ \mathbb{Z}/2, & \epsilon = 6, \\ \mathbb{Z}/2^{4n+6} \oplus \mathbb{Z}/2^{2n}, & \epsilon = 7. \end{cases} \quad (\text{A.9})$$

n	1	2	3	4	5	6
$\tilde{\Omega}_n^{\text{Spin}}(BQ_8)$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/8 \times (\mathbb{Z}/4)^2$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$

Table 6. The reduced spin bordism groups for BQ_8 in degrees lower than 7.

$$\widetilde{ko}_{\epsilon+8n}(\Sigma^{-1}BS^3/BN) \cong \begin{cases} \mathbb{Z}/2, & \epsilon = 1, 2, \\ \mathbb{Z}/2^{2n+2}, & \epsilon = 3, 7, \\ 0, & \epsilon = 4, 5, 6, 8. \end{cases} \quad (\text{A.10})$$

We combine these results and summarise the spin bordism groups of BQ_8 in table 6 for convenience (see also section 7.4.C of ref. [57]). From this identification, we can deduce that

$$\Omega_5^{\text{Spin}}(BQ_8) \cong \mathbb{Z}/2 \times \mathbb{Z}/2. \quad (\text{A.11})$$

A.4 $G = \text{SL}(2, \mathbb{F}_3)$

The group

$$\text{SL}(2, \mathbb{F}_3) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \mid a, b, c, d \in \mathbb{F}_3, ad - bc = 1 \right\} \quad (\text{A.12})$$

is a group extension of $\mathbb{Z}/3$ by the quaternion group Q_8 ,

$$0 \rightarrow Q_8 \rightarrow \text{SL}(2, \mathbb{F}_3) \rightarrow \mathbb{Z}/3 \rightarrow 0. \quad (\text{A.13})$$

We know from [67] that at 2-localisation,

$$\widetilde{ko}_5(B\text{SL}(2, \mathbb{F}_3))_2^\wedge \cong \mathbb{Z}/2 \times \mathbb{Z}/2, \quad (\text{A.14})$$

but since there is a 3-torsion present in $\text{SL}(2, \mathbb{F}_3)$, there is a possibility that there is a non-trivial odd-torsion piece in the spin bordism of $B\text{SL}(2, \mathbb{F}_3)$. To find the odd torsion, we consider the Atiyah-Hirzebruch spectral sequence for the fibration $\text{pt} \rightarrow B\text{SL}(2, \mathbb{F}_3) \rightarrow B\text{SL}(2, \mathbb{F}_3)$,

$$E_{p,q}^2 = H_p(B\text{SL}(2, \mathbb{F}_3), \Omega_q^{\text{Spin}}(\text{pt})) \Rightarrow \Omega_{p+q}^{\text{Spin}}(B\text{SL}(2, \mathbb{F}_3)). \quad (\text{A.15})$$

To construct the entries on the E^2 page, we need to know both the integral and mod 2 homology groups of $B\text{SL}(2, \mathbb{F}_3)$. Using the isomorphism between the singular homology of BG and the group homology of G , $H_\bullet(BG; \mathbb{Z}) \cong \mathcal{H}_\bullet(G, \mathbb{Z})$, with help from GAP [55], we obtain the homology groups of $B\text{SL}(2, \mathbb{F}_3)$ as shown in table The E^2 page of the AHSS (A.15) can be constructed as shown in figure 1. The entries on the diagonal $p + q = 5$ already stabilise on this page. To determine the 5th bordism group, however, we need to solve the non-trivial extension problem. Using the result from the 2-localisation above, we narrow down the extensions to two options:

$$\Omega_5^{\text{Spin}}(B\text{SL}(2, \mathbb{F}_3)) \cong \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/9 \quad \text{or} \quad \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/3 \times \mathbb{Z}/3. \quad (\text{A.16})$$




Figure 2. The $\mathcal{A}(1)$ -module structure and Adams chart for calculating spin bordism of BD_{2n} , for odd n .

to $\mathbb{Z}/2$ or 0 respectively. Thus we learn from the AHSS that $\Omega_6^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3)) \cong \mathbb{Z}/2$ or 0. Computation of $\text{ko}(\text{BSL}(2, \mathbb{F}_3))$ at 2-completion in ref. [67], which we have quoted in eq. (A.9), tells us that this differential must be trivial and $\Omega_6^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3))$ must be isomorphic to $\mathbb{Z}/2$.

A.5 The dihedral group $G = D_{2n}$

In this appendix, we will show that $\Omega_5^{\text{Spin}}(BD_{2n})$ vanishes for any n .

Odd n . Consider first the case with odd n . The integral cohomology ring and the mod 2 cohomology ring of BD_{2n} are given by [69]

$$H^\bullet(BD_{2n}; \mathbb{Z}) \cong \frac{\mathbb{Z}[a_2, d_4]}{(2a_2, nd_4)}, \quad H^\bullet(BD_{2n}; \mathbb{Z}/2) \cong \mathbb{Z}/2[u_1], \quad (\text{A.20})$$

where the subscript on each indeterminate denotes its degree. We will now apply the Adams spectral sequence

$$E_2^{s,t} = \text{Ext}_{\mathcal{A}(1)}^{s,t}(\tilde{H}^\bullet(BD_{2n}; \mathbb{Z}/2), \mathbb{Z}/2) \Rightarrow \tilde{\Omega}_{t-s}^{\text{Spin}}(BD_{2n})_2^\wedge, \quad t-s \leq 7, \quad (\text{A.21})$$

to find 2-torsion factors of the spin bordism groups of BD_{2n} . As an $\mathcal{A}(1)$ -module, $H^\bullet(BD_{2n}; \mathbb{Z}/2)$ has the structure as depicted by a cell diagram in figure 2a. Discarding the summand $\mathbb{Z}/2$ which is not part the reduced mod 2 cohomology, we obtain the Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(\tilde{H}^\bullet(BD_{2n}; \mathbb{Z}/2), \mathbb{Z}/2)$ as shown in figure 2b, whence the 2-torsion factors in the spin bordism groups of BD_{2n} can be read off:

$$\begin{aligned} \tilde{\Omega}_1^{\text{Spin}}(BD_{2n})_2^\wedge &= \tilde{\Omega}_2^{\text{Spin}}(BD_{2n})_2^\wedge \cong \mathbb{Z}/2, & \tilde{\Omega}_3^{\text{Spin}}(BD_{2n})_2^\wedge &\cong \mathbb{Z}/8, \\ \tilde{\Omega}_4^{\text{Spin}}(BD_{2n})_2^\wedge &= \tilde{\Omega}_5^{\text{Spin}}(BD_{2n})_2^\wedge = \tilde{\Omega}_6^{\text{Spin}}(BD_{2n})_2^\wedge = 0. \end{aligned} \quad (\text{A.22})$$

We complete our computation by working out the possible odd-torsion through the Atiyah-Hirzebruch spectral sequence (AHSS). Applying the universal coefficient theorem

i	0	1	2	3	4	5	6	7
$H_i(BD_{2n}; \mathbb{Z})$	\mathbb{Z}	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \times \mathbb{Z}/n$	0	$\mathbb{Z}/2$	0	\mathbb{Z}/n
$H_i(BD_{2n}; \mathbb{Z}/2)$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$

Table 8. Integral and mod 2 homology groups of BD_{2n} with odd n .

E^2 page								E^3 page							
5	0	0	0	0	0	0	*	5	0	0	0	0	0	0	*
4	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \times \mathbb{Z}/n$	0	$\mathbb{Z}/2$	*	4	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \times \mathbb{Z}/n$	0	$\mathbb{Z}/2$	*
3	0	0	0	0	0	0	*	3	0	0	0	0	0	0	*
2	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	*	2	0	$\mathbb{Z}/2$	0	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	*
1	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	$\mathbb{Z}/2$	*	1	0	$\mathbb{Z}/2$	$\mathbb{Z}/2$	0	0	0	*
0	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \times \mathbb{Z}/n$	0	$\mathbb{Z}/2$	0	0	0	$\mathbb{Z}/2$	0	$\mathbb{Z}/2 \times \mathbb{Z}/n$	0	0	0
	0	1	2	3	4	5	6		0	1	2	3	4	5	6

Figure 3. Pages E^2 and E^3 of the AHSS for $\tilde{\Omega}_d^{\text{Spin}}(BD_{2n})$ with odd n .

to the cohomology groups given in eq. (A.20), we obtain the homology groups, as shown in table 8. We can then use this data to construct the corresponding AHSS for the fibration $\text{pt} \rightarrow BD_{2n} \rightarrow BD_{2n}$ converging to the reduced bordism groups,

$$E_{p,q}^2 = \tilde{H}_p(BD_{2n}; \Omega_q^{\text{Spin}}(\text{pt})) \Rightarrow \tilde{\Omega}_{p+q}^{\text{Spin}}(BD_{2n}), \quad (\text{A.23})$$

as shown in figure 3. The non-trivial differentials on the E^2 page are denoted by the red arrows. These differentials can be deduced from the fact that the differentials d_2 acting on the 0th and 1st rows in the E^2 page of the AHSS for the fibration $\text{pt} \rightarrow BG \rightarrow BG$ converging to a spin bordism group is dual to the action of the Steenrod square Sq^2 on the mod 2 cohomology [70, 71]. The mod 2 cohomology ring of BD_{2n} with n odd is generated by a single generator of degree one, so the action of the Steenrod squares on any element of the cohomology ring can be deduced from the axioms.

The entries with $p + q \leq 6$ all stabilise by the E^4 page, where we have used the consistency with the Adams spectral sequence calculation above to deduce that $d_3 : E_{4,2}^3 \rightarrow E_{1,4}^3$ is the identity map. The computation shows us that for $d \leq 6$, there is an odd-torsion factor in $\Omega_d^{\text{Spin}}(BD_{2n})$ only when $d = 3$:

$$\Omega_3^{\text{Spin}}(BD_{2n}) \cong \mathbb{Z}/n \times \mathbb{Z}/8, \quad n \text{ odd}. \quad (\text{A.24})$$

The resulting spin bordism groups of BD_{2n} , with odd n , are summarised in table 1.

Even n . The calculation of the spin bordism groups of BD_{2n} when n is even is more involved. For n even, the cohomology ring is given by [69]

$$H^\bullet(BD_{2n}; \mathbb{Z}) \cong \mathbb{Z}[a_2, b_2, c_3, d_4]/I, \quad (\text{A.25})$$




Figure 4. The $\mathcal{A}(1)$ -module structure and Adams chart for calculating spin bordism of BD_{2n} , for even n .

where the subscripts on the generators denote the degree, and I is the ideal generated by $2a_2$, $2b_2$, $2c_3$, nd_4 , $b_2^2 + a_2b_2 + (n^2/4)d_4$, and $c_3^2 + a_2d_4$. Writing this out explicitly, we have (for $p \neq 0$)

$$H^p(BD_{2n}; \mathbb{Z}) \cong \begin{cases} (\mathbb{Z}/2)^{\frac{p-1}{2}}, & p = 1, 3 \bmod 4 \\ (\mathbb{Z}/2)^{\frac{p+2}{2}}, & p = 2 \bmod 4 \\ \mathbb{Z}/n \times (\mathbb{Z}/2)^{p/2}, & p = 0 \bmod 4 \end{cases} \quad (\text{A.26})$$

The mod 2 cohomology ring is given by [69]

$$H^*(BD_{2n}; \mathbb{Z}/2) \cong \frac{\mathbb{Z}/2[u_1, v_1, w_2]}{(u_1^2 + u_1v_1 + (n/2)w_2)}, \quad n \text{ even}, \quad (\text{A.27})$$

with $\text{Sq}^1(w_2) = v_1w_2$ (the action of the Steenrod squares on other generators follows automatically from the axioms). Using the same strategy as the odd n case, we first construct the Adams spectral sequence (A.21) to find the 2-torsion factors. As a $\mathcal{A}(1)$ -module, the structure of $H^*(BD_{2n}; \mathbb{Z}/2)$ is given in figure 4a. Ignoring the summand $\mathbb{Z}/2$ which gives us the spin bordism groups of a point, the corresponding the Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(\tilde{H}^*(BD_{2k+2}; \mathbb{Z}/2), \mathbb{Z}/2)$ is shown in figure 4b. We can see right away that

$$\tilde{\Omega}^{\text{Spin}}(BD_{2n})_2^\wedge \cong \mathbb{Z}/2 \times \mathbb{Z}/2, \quad \tilde{\Omega}_5^{\text{Spin}}(BD_{2n})_2^\wedge = 0. \quad (\text{A.28})$$

However, we cannot determine the differentials from the column $t-s=4$ to the $t-s=3$ in general, nor can we solve the extension problem in the column $t-s=2$. Finally, the AHSS for the fibration $\text{pt} \rightarrow BD_{2n} \rightarrow BD_{2n}$,

$$E_{p,q}^2 = H_p(BD_{2n}; \Omega_q^{\text{Spin}}(\text{pt})) \Rightarrow \Omega_{p+q}^{\text{Spin}}(BD_{2n}), \quad (\text{A.29})$$

whose E^2 page is shown in figure 5, tells us that there is no odd-torsion in the spin bordism group of BD_{2n} of degree $d=5$. Hence, we can conclude that

$$\tilde{\Omega}_5^{\text{Spin}}(BD_{2n}) = 0, \quad (\text{A.30})$$

for any integer n . Similarly, we also know that $\tilde{\Omega}_1^{\text{Spin}}(BD_{2n}) = (\mathbb{Z}/2)^2$.

E^2 page

5	0	0	0	0	0	0	0
4	\mathbb{Z}	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \times \mathbb{Z}/n$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^3$
3	0	0	0	0	0	0	0
2	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^3$	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^5$	$(\mathbb{Z}/2)^6$	$(\mathbb{Z}/2)^7$
1	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^3$	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^5$	$(\mathbb{Z}/2)^6$	$(\mathbb{Z}/2)^7$
0	\mathbb{Z}	$(\mathbb{Z}/2)^2$	$\mathbb{Z}/2$	$(\mathbb{Z}/2)^2 \times \mathbb{Z}/n$	$(\mathbb{Z}/2)^2$	$(\mathbb{Z}/2)^4$	$(\mathbb{Z}/2)^3$
	0	1	2	3	4	5	6

Figure 5. The E^2 page of the Atiyah-Hirzebruch spectral sequence for calculating $\Omega_d^{\text{Spin}}(BD_{2n})$ when n is even.

d	2	3	4	5	6
$\widetilde{ko}_d(BD_{2^{k+2}})$	$(\mathbb{Z}/2)^c$	$ \widetilde{ko}_3(BD_{2^{k+2}}) = 2^{k+8}$	$(\mathbb{Z}/2)^2$	0	$\mathbb{Z}/2$

Table 9. The reduced ko -homology of $BD_{2^{k+2}}$ from degree 2 through 6.

One can do better for $n = 2^{k+1}$ and find also the bordism groups in lower degrees. We proceed by first making use of the isomorphism between $\tilde{\Omega}_d^{\text{Spin}}(BD_{2n})$ and $\widetilde{ko}_d(BD_{2n})$ and note that the K-theory computation has been carried out using the Bockstein spectral sequence in ref. [57], which we reproduce in table 9, where the power c are determined up to a range: $3 \leq c \leq 2^k + 5$.

The first thing we notice from the table above is that

$$\tilde{\Omega}_5^{\text{Spin}}(BD_{2^{k+2}}) \cong \widetilde{ko}_5(BD_{2^{k+2}}) = 0, \quad (\text{A.31})$$

agreeing with our calculation above. Moreover, the K-theory results also resolves the extension problem in the column $t - s = 2$ in the ASS shown in figure 4b, giving us

$$\tilde{\Omega}_2^{\text{Spin}}(BD_{2^{k+2}}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2 \times \mathbb{Z}/2. \quad (\text{A.32})$$

The only remaining spin bordism groups of $BD_{2^{k+2}}$ in degrees lower than 5 to be worked out are $\tilde{\Omega}_3^{\text{Spin}}(BD_{2^{k+2}})$ and $\tilde{\Omega}_4^{\text{Spin}}(BD_{2^{k+2}})$. The connective KO-theory computation in ref. [57], done via the Bockstein spectral sequence, only gives the order of the group: $|\Omega_3^{\text{Spin}}(BD_{2^{k+2}})| = 2^{k+8}$. Fortunately, we can use this information to deduce the differentials from the column $t - s = 4$ in the ASS of figure 4b and precisely determine the bordism groups, as follows.

From the computations of the connective real K-theory for $BD_{2^{k+1}}$ in ref. [57], we know that

$$\tilde{\Omega}_4^{\text{Spin}}(BD_{2^{k+2}}) \cong \mathbb{Z}/2 \times \mathbb{Z}/2. \quad (\text{A.33})$$

Therefore, there is bound to be a series of differentials d_r on the r^{th} page out of the h_0 -tower in the $t - s = 4$ column, where r is still unknown. The differentials will hit part of




Figure 6. Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(\tilde{H}^\bullet(BD_8; \mathbb{Z}/2), \mathbb{Z}/2)$.

the h_0 -tower in the $t - s = 3$ column, truncating the extension down to $\mathbb{Z}/2^{r+1}$. The third spin bordism group then has the order $2^{3+3+r+1} = 2^{r+7}$. Again, we know from ref. [57] that the third spin bordism group of BD_{2k+2} is 2^{k+8} . Thus, we deduce that

$$r = k + 1.$$

For example, the Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(\tilde{H}^\bullet(BD_8; \mathbb{Z}/2))$ is shown in figure 6. We can then conclude that

$$\tilde{\Omega}_3^{\text{Spin}}(BD_{2k+2}) \cong (\mathbb{Z}/8)^2 \times \mathbb{Z}/2^{k+2}. \quad (\text{A.34})$$

These results are summarised in table 1.

B TQFTs for Q_8 structure

In section 4.2 it was shown that there is no global anomaly from fermions in any representation of Q_8 , because the η -invariant vanishes identically when evaluated on a generator of $\Omega_5^{\text{Spin}}(BQ_8)$ [59]. We might then be tempted to conclude that the factors $\mathbb{Z}/2 \times \mathbb{Z}/2$ in $\Omega_5^{\text{Spin}}(BQ_8)$ must be associated with ‘pure bosonic anomalies’ that descend from mod 2 cohomology. Indeed, the mod 2 cohomology ring of BQ_8 is [67]

$$H^\bullet(BQ_8, \mathbb{Z}/2) \cong \frac{\mathbb{Z}/2[x, y, w]}{(x^2 + xy + y^2, x^2y + xy^2)}, \quad (\text{B.1})$$

where $\deg(x) = \deg(y) = 1$ while $\deg(w) = 4$, and so there are exactly two elements in the degree 5 cohomology group, xw and yw . However, it does not follow that these cohomology classes are the cobordism generators that we are looking for. It turns out that the cobordism generators of the $\mathbb{Z}/2 \times \mathbb{Z}/2$ depend on the spin structure.

One way to see this is to look at the Atiyah-Hirzebruch spectral sequence

$$E_{p,q}^2 = H_p(BQ_8; \Omega_q^{\text{Spin}}(\text{pt})) \Rightarrow \Omega_{p+q}^{\text{Spin}}(BQ_8) \quad (\text{B.2})$$

for the fibration $\text{pt} \rightarrow BQ_8 \rightarrow BQ_8$, as shown in figure 7. Since Sq^2 acts trivially on every generator of $H^\bullet(BQ_8; \mathbb{Z}/2)$, it follows that all differentials on the E^2 page that act on the

E^2 page

5	0	0	0	0	0	0	0	0
4	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	\mathbb{Z}_2^2	0	\mathbb{Z}_8
3	0	0	0	0	0	0	0	0
2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2
1	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2
0	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	\mathbb{Z}_2^2	0	\mathbb{Z}_8
	0	1	2	3	4	5	6	7

Figure 7. The E^2 page of the AHSS for calculating $\Omega_{\bullet}^{\text{Spin}}(BQ_8)$.

E^3 page

5	0	0	0	0	0	0	0	0
4	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	*	0	*
3	0	0	0	0	0	0	0	0
2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	*	*
1	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	*
0	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	\mathbb{Z}_2^2	0	\mathbb{Z}_8
	0	1	2	3	4	5	6	7

E^4 page

5	0	0	0	0	0	0	0	0
4	\mathbb{Z}	\mathbb{Z}_2	0	\mathbb{Z}_8	0	*	0	*
3	0	0	0	0	0	0	0	0
2	\mathbb{Z}_2	\mathbb{Z}_2^2	0	\mathbb{Z}_2	0	\mathbb{Z}_2^2	*	*
1	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	*
0	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	0	0	\mathbb{Z}_8
	0	1	2	3	4	5	6	7

Figure 8. The E^3 and E^4 pages of the AHSS for $\Omega_{\bullet}^{\text{Spin}}(BQ_8)$.

0th and 1st rows must also be trivial. To work out some differentials on the E^3 page, we compare the AHSS with known results. Now, we know from the K-theory computation that $\tilde{\Omega}_4^{\text{Spin}}(BQ_8) \cong \mathbb{Z}/2$. On the $p + q = 4$ diagonal, there are entries $E_{2,2}^3 \cong (\mathbb{Z}/2)^2$ and $E_{3,1}^3 \cong \mathbb{Z}/2$. Out of these two, the entry $E_{3,1}^3$ already stabilises since there are no more non-trivial differentials into or emanating from it. Thus, it is this entry that gives the resulting $\mathbb{Z}/2$ factor in $\tilde{\Omega}_4^{\text{Spin}}(BQ_8)$, and the entry $E_{2,2}^3$ must be killed by the differential

$$d_3 : E_{5,0}^3 \cong (\mathbb{Z}/2)^2 \rightarrow E_{2,2}^3 \cong (\mathbb{Z}/2)^2 \quad (\text{B.3})$$

as shown in figure 8.

The $\mathbb{Z}/2$ factors from the entries $E_{4,1}^3$ and $E_{3,2}^3$ also stabilise at this page. Since $\Omega_5^{\text{Spin}}(BQ_8) \cong (\mathbb{Z}/2)^2$, we can deduce that the entry $E_{1,4}^3$ must be completely killed by differentials, either successively from $E_{4,2}^3$ and $E_{5,1}^4$ (which is the situation shown in figure 8), or in one go by d_4 from $E_{5,1}^4$, as shown in figure 9. We can then clearly see that the entries contributing to the $\mathbb{Z}/2$ factors in $\Omega_5^{\text{Spin}}(BQ_8)$ depends on the $\mathbb{Z}/2$ factors in $\Omega_1^{\text{Spin}}(\text{pt})$ and $\Omega_2^{\text{Spin}}(\text{pt})$, which need the existence of the spin structure as claimed.

E^3 page									E^4 page								
5	0	0	0	0	0	0	0	0	5	0	0	0	0	0	0	0	0
4	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	*	0	*	4	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	*	0	*
3	0	0	0	0	0	0	0	0	3	0	0	0	0	0	0	0	0
2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	*	*	2	\mathbb{Z}_2	\mathbb{Z}_2^2	0	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	*	*
1	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	*	1	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	\mathbb{Z}_2	\mathbb{Z}_2	\mathbb{Z}_2^2	\mathbb{Z}_2^2	*
0	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	\mathbb{Z}	\mathbb{Z}_2^2	0	\mathbb{Z}_8	0	0	0	\mathbb{Z}_8
	0	1	2	3	4	5	6	7		0	1	2	3	4	5	6	7

Figure 9. Another possibility for the elimination of the entry $E_{1,4}$.

The $\mathbb{Z}/2$ in the entry $E_{3,2}^\infty$ from AHSS computation above seems to suggest that the bordism generator for this $\mathbb{Z}/2$ factor is a product of the torus with periodic spin structure around both 1-cycles and a 3-manifold that is a generator of the $\mathbb{Z}/8$ factor in $\Omega_3^{\text{Spin}}(BQ_8)$. Indeed, this observation is correct, but we need to examine the Adams spectral sequence to confirm it.

The Adams spectral sequence (ASS) for our purpose reads

$$E_2^{s,t} = \text{Ext}_{\mathcal{A}(1)}^{s,t}(H^\bullet(BQ_8; \mathbb{Z}/2), \mathbb{Z}/2) \Rightarrow \Omega_{t-s}^{\text{Spin}}(BQ_8)_2^\wedge, \quad (\text{B.4})$$

with the input being the mod 2 cohomology ring of BQ_8 , given by eq. (B.1). Using GAP [55], the action of $\mathcal{A}(1)$ on the generators of $H^\bullet(BQ_8; \mathbb{Z}/2)$ that cannot be obtained from the Axioms reads

$$\text{Sq}^1 w = 0, \quad \text{Sq}^2 w = 0. \quad (\text{B.5})$$

This information is enough to determine the structure of $H^\bullet(BQ_8; \mathbb{Z}/2)$ as an $\mathcal{A}(1)$ -module, which we present in figure 10 below. Algebraically, we can write


$$H^\bullet(BQ_8; \mathbb{Z}/2) \cong \mathbb{Z}/2 \oplus \Sigma S \oplus \Sigma S \oplus \Sigma^3 \mathbb{Z}/2 \oplus \Sigma^4 \mathbb{Z}/2 \oplus \Sigma^5 S \oplus \Sigma^5 S \oplus \dots, \quad (\text{B.6})$$

where S is the “stick” $\mathcal{A}(1)$ -module. Since we already know the Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(\mathbb{Z}/2, \mathbb{Z}/2)$, we only need calculate the Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(S, \mathbb{Z}/2)$ to obtain the E_2 page of the ASS for $H^\bullet(BQ_8; \mathbb{Z}/2)$.

Again, we put the module S in the middle of a short exact sequence, which reads

$$0 \rightarrow \Sigma \mathbb{Z}/2 \rightarrow S \rightarrow \mathbb{Z}/2 \rightarrow 0, \quad (\text{B.7})$$

which pictorially looks like



This short exact sequence induces a long exact sequence in the Ext-functor [65], whose connecting morphism

$$\delta : \text{Ext}_{\mathcal{A}(1)}^{s,t}(\Sigma \mathbb{Z}/2, \mathbb{Z}/2) \rightarrow \text{Ext}_{\mathcal{A}(1)}^{s,t}(\mathbb{Z}/2, \mathbb{Z}/2), \quad (\text{B.8})$$




Figure 10. The $\mathcal{A}(1)$ -module structure of $H^\bullet(BQ_8; \mathbb{Z}/2)$ up to degree 5.




Figure 11. The Adams chart for $\text{Ext}_{\mathcal{A}(1)}^{s,t}(S, \mathbb{Z}/2)$.

which translates into a differentials of slope $(-1, 1)$ in the Adams chart where the charts for $\Sigma\mathbb{Z}/2$ and $\mathbb{Z}/2$ are drawn together as shown in figure 11. We thus obtain the Adams chart for the stick module S as shown on the right of the figure.

We now combine the Adams chart of S with the Adams chart of $\mathbb{Z}/2$, and ignoring the $\mathbb{Z}/2$ cell at the bottom that only gives us the factors corresponding to $\Omega_{\bullet}^{\text{Spin}}(\text{pt})$. Then, the Adams charts for the E_2 , E_3 , E_4 pages of the Adams spectral sequence that computes the reduced spin bordism groups of BQ_8 is shown in figure 12. The differentials are deduced in such a way that the resulting bordism groups are compatible with the results from K-theory computation in [67].

In the Adams chart, the vertical line indicates the non-trivial extension by $\mathbb{Z}/2$ and the slanted line of slope 1 indicates a Cartesian product by a circle with periodic spin structure.¹⁹ In the final page of figure 12, we see that one $\mathbb{Z}/2$ factor in $\Omega_5^{\text{Spin}}(BQ_8)$ is at the end of a sequence of two slanted lines starting at the generator of the $\mathbb{Z}/8$ factor in $\Omega_3^{\text{Spin}}(BQ_8)$.²⁰ Ref. [59] showed that a generator of the $\mathbb{Z}/8$ factor can be taken to be the quotient $S^3/\tau(Q_8)$, where τ is the 2-dimensional representation of Q_8 , which can be

¹⁹We thank Arun Debray for helpful discussions about this.

²⁰We know that the element at the beginning of the sequence of two slanted lines generates $\mathbb{Z}/8$ because it is also a starting point of two successive vertical lines.




Figure 12. The Adams spectral sequence calculating the reduced spin bordism groups of BQ_8 .

thought of as a map $\tau : Q_8 \rightarrow \text{SU}(2) \cong S^3$ given explicitly by

$$\tau(\pm 1) = \pm \mathbb{I}_2, \quad \tau(\pm i) = \pm i \sigma_3, \quad \tau(\pm j) = \pm i \sigma_2, \quad \tau(\pm k) = \pm i \sigma_1, \quad (\text{B.9})$$

σ_i being the usual Pauli matrices. Thus, one of the two bordism generators of $\Omega_5^{\text{Spin}}(BQ_8)$ is

$$T_{PP}^2 \times S^3 / \tau(Q_8), \quad (\text{B.10})$$

as hinted in the AHSS, where T_{PP}^2 is a torus with periodic spin structure around both 1-cycles. The corresponding bordism invariant is

$$\text{Arf} \cup xy^2, \quad (\text{B.11})$$

where Arf is the Arf invariant of the spin structure.

Notice that the cohomology classes xw and yw , identified at the start of this appendix, are trivial on this generator of the bordism group: we can factorise e.g. $xw(T_{PP}^2 \times S^3 / \tau(Q_8)) = x(S_P^1)w(S_P^1 \times S^3 / \tau(Q_8))$ which vanishes because the S^1 does not come with a non-trivial Q_8 bundle, and so the cohomology class $x \in H^1(BQ_8; \mathbb{Z}/2)$ is trivial when evaluated on it.²¹ The same argument works for yw . While not giving an explicit calculation, we expect a similar story to play out for the second independent bordism invariant.

C Bordism generators of $\Omega_5^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3))$ and $\Omega_5^{\text{Spin}}(BQ_8)$

In this appendix, we will argue that the manifolds generating the factor $\mathbb{Z}/2 \times \mathbb{Z}/2$ of $\Omega_5^{\text{Spin}}(\text{BSL}(2, \mathbb{F}_3))$ are also generators of $\Omega_5^{\text{Spin}}(BQ_8) \cong \mathbb{Z}/2 \times \mathbb{Z}/2$.

Since we are only concerned with a mod 2 effect, we can work at 2-completion without loss of generality. A generator of the $\mathbb{Z}/2 \times \mathbb{Z}/2$ is given by a spin 5-manifold X together with a map

$$f : X \rightarrow \text{BSL}(2, \mathbb{F}_3). \quad (\text{C.1})$$

²¹To see this, note that $x(S_P^1)$ is shorthand for the pairing $\langle f^*(x), S_P^1 \rangle$, where $f^* : H^\bullet(BQ_8; \mathbb{Z}/2) \rightarrow H^\bullet(S_P^1; \mathbb{Z}/2)$ is the pullback of the classifying map $f : S_P^1 \rightarrow BQ_8$. Since the Q_8 -bundle on S_P^1 is trivial, f is trivial, and so does $f^*(x)$.

	1	1	4	4	6	4	4
(θ_1, θ_2)	$(0, 0)$	$(\pi, -\pi)$	$(0, -\frac{2\pi}{3})$	$(0, \frac{2\pi}{3})$	$(\frac{\pi}{2}, -\frac{\pi}{2})$	$(\pi, \frac{\pi}{3})$	$(-\pi, -\frac{\pi}{3})$
(\mathbf{n}, q)	n	$n(-1)^{n-1}$	$e^{-\frac{\pi i q}{3}} \frac{\sin \frac{n\pi}{3}}{\sin \frac{\pi}{3}}$	$e^{\frac{\pi i q}{3}} \frac{\sin \frac{n\pi}{3}}{\sin \frac{\pi}{3}}$	$\sin \frac{n\pi}{2}$	$e^{\frac{2\pi i q}{3}} \frac{\sin \frac{n\pi}{3}}{\sin \frac{\pi}{3}}$	$e^{-\frac{2\pi i q}{3}} \frac{\sin \frac{n\pi}{3}}{\sin \frac{\pi}{3}}$

Table 10. Characters for the irreps of $U(2)$ restricted to the subgroup $SL(2, \mathbb{F}_3)$ with the subgroup embedding (4.24).

Note that the subgroup embedding $Q_8 \hookrightarrow SL(2, \mathbb{F}_3)$ induces a map $i : BQ_8 \rightarrow BSL(2, \mathbb{F}_3)$. Moreover, at 2-completion, there is a stable decomposition [68]

$$BQ_8 \cong BSL(2, \mathbb{F}_3) \vee \Sigma^{-1} (BS^3 / BN) \vee \Sigma^{-1} (BS^3 / BN), \quad (C.2)$$

where N is the normaliser of a maximal torus in S^3 , and \vee is the wedge sum. Thus, the map f naturally induces a map $f' : X \rightarrow BQ_8$ such that $i \circ f' = f$ and we can also think of X as equipped with a Q_8 -bundle. Now, suppose that (X, f') is nullbordant, that is, there exists a spin 6-manifold Y with a map $g : Y \rightarrow BQ_8$ such that $\partial Y = X$ and $g|_{\partial} = f'$. Using the embedding $Q \hookrightarrow SL(2, \mathbb{F}_3)$, we now have a spin 6-manifold Y whose boundary is X with a map $i \circ g : Y \rightarrow BSL(2, \mathbb{F}_3)$ such that $i \circ g|_{\partial} = i \circ f' = f$. Hence, (X, f) is nullbordant, which is a contradiction. Therefore, manifolds that generate the $\mathbb{Z}/2 \times \mathbb{Z}/2$ factor in $\Omega_5^{\text{Spin}}(BSL(2, \mathbb{F}_3))$ also generate $\Omega_5^{\text{Spin}}(BQ_8)$.

D Decomposition of Standard Model representation

In this short appendix we derive the decomposition (4.26) used in our proof of the anomaly structure of the group $SL(2, \mathbb{F}_3)$. It can be deduced via character theory as follows. First, note that any element of the group $U(2)$ is conjugate to a diagonal matrix of the form $\text{diag}(e^{i\theta_1}, e^{i\theta_2})$, so we can label a conjugacy class by a pair of angles (θ_1, θ_2) . The character of the conjugacy class (θ_1, θ_2) in the irrep (\mathbf{n}, q) is given by the *Weyl character formula*

$$\chi_{n,k}(\theta_1, \theta_2) = e^{i(q-n+1)\frac{\theta_1+\theta_2}{2}} \left[\frac{e^{in\theta_1} - e^{in\theta_2}}{e^{i\theta_1} - e^{i\theta_2}} \right]. \quad (D.1)$$

Each conjugacy class of the subgroup $SL(2, \mathbb{F}_3)$ under the subgroup embedding (4.24) can now be alternatively labelled by (θ_1, θ_2) , given in table 10, with the conjugacy classes arranged in the same order as the character table of $SL(2, \mathbb{F}_3)$ (table 4). Applying the Weyl character formula above to each of these conjugacy classes, we arrive at the character table of the $U(2)$ irrep (\mathbf{n}, q) restricted to the subgroup $SL(2, \mathbb{F}_3)$. The result is again tabulated in table 10. We can now use the orthogonality of the characters to obtain the decomposition for each irrep (\mathbf{n}, q) . Applying the outlined procedure to the irreps in the Standard Model representation (4.25), we easily obtain the decomposition (4.26) as claimed.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited. SCOAP³ supports the goals of the International Year of Basic Sciences for Sustainable Development.

References

- [1] J.S. Bell and R. Jackiw, *A PCAC puzzle: $\pi^0 \rightarrow \gamma\gamma$ in the σ model*, *Nuovo Cim. A* **60** (1969) 47 [INSPIRE].
- [2] S.L. Adler, *Axial vector vertex in spinor electrodynamics*, *Phys. Rev.* **177** (1969) 2426 [INSPIRE].
- [3] G. 't Hooft, *Naturalness, chiral symmetry, and spontaneous chiral symmetry breaking*, *NATO Sci. Ser. B* **59** (1980) 135 [INSPIRE].
- [4] E. Witten, *Global gravitational anomalies*, *Commun. Math. Phys.* **100** (1985) 197 [INSPIRE].
- [5] X.-z. Dai and D.S. Freed, *eta invariants and determinant lines*, *J. Math. Phys.* **35** (1994) 5155 [Erratum *ibid.* **42** (2001) 2343] [hep-th/9405012] [INSPIRE].
- [6] A. Kapustin, R. Thorngren, A. Turzillo and Z. Wang, *Fermionic Symmetry Protected Topological Phases and Cobordisms*, *JHEP* **12** (2015) 052 [arXiv:1406.7329] [INSPIRE].
- [7] E. Witten, *Fermion Path Integrals And Topological Phases*, *Rev. Mod. Phys.* **88** (2016) 035001 [arXiv:1508.04715] [INSPIRE].
- [8] D.S. Freed and M.J. Hopkins, *Reflection positivity and invertible topological phases*, *Geom. Topol.* **25** (2021) 1165 [arXiv:1604.06527] [INSPIRE].
- [9] E. Witten and K. Yonekura, *Anomaly Inflow and the η -Invariant*, in *The Shoucheng Zhang Memorial Workshop*, 9, 2019 [arXiv:1909.08775] [INSPIRE].
- [10] E. Witten, *An $SU(2)$ Anomaly*, *Phys. Lett. B* **117** (1982) 324 [INSPIRE].
- [11] J. Davighi and N. Lohitsiri, *Omega vs. pi, and 6d anomaly cancellation*, *JHEP* **05** (2021) 267 [arXiv:2012.11693] [INSPIRE].
- [12] Y. Tachikawa and K. Yonekura, *Why are fractional charges of orientifolds compatible with Dirac quantization?*, *SciPost Phys.* **7** (2019) 058 [arXiv:1805.02772] [INSPIRE].
- [13] Z. Wan and J. Wang, *Higher anomalies, higher symmetries, and cobordisms I: classification of higher-symmetry-protected topological states and their boundary fermionic/bosonic anomalies via a generalized cobordism theory*, *Ann. Math. Sci. Appl.* **4** (2019) 107 [arXiv:1812.11967] [INSPIRE].
- [14] J. Wang, X.-G. Wen and E. Witten, *A New $SU(2)$ Anomaly*, *J. Math. Phys.* **60** (2019) 052301 [arXiv:1810.00844] [INSPIRE].
- [15] D.S. Freed, *Pions and Generalized Cohomology*, *J. Diff. Geom.* **80** (2008) 45 [hep-th/0607134] [INSPIRE].
- [16] I. García-Etxebarria and M. Montero, *Dai-Freed anomalies in particle physics*, *JHEP* **08** (2019) 003 [arXiv:1808.00009] [INSPIRE].
- [17] J. Davighi, B. Gripaios and N. Lohitsiri, *Global anomalies in the Standard Model(s) and Beyond*, *JHEP* **07** (2020) 232 [arXiv:1910.11277] [INSPIRE].
- [18] Z. Wan and J. Wang, *Beyond Standard Models and Grand Unifications: Anomalies, Topological Terms, and Dynamical Constraints via Cobordisms*, *JHEP* **07** (2020) 062 [arXiv:1910.14668] [INSPIRE].
- [19] J. Davighi and J. Tooby-Smith, *Electroweak flavour unification*, arXiv:2201.07245 [INSPIRE].

- [20] J. Wang, Z. Wan and Y.-Z. You, *Proton stability: From the standard model to beyond grand unification*, *Phys. Rev. D* **106** (2022) 025016 [[arXiv:2204.08393](#)] [[INSPIRE](#)].
- [21] Y. Lee and Y. Tachikawa, *Some comments on 6D global gauge anomalies*, *PTEP* **2021** (2021) 08B103 [[arXiv:2012.11622](#)] [[INSPIRE](#)].
- [22] I. García-Etxebarria, H. Hayashi, K. Ohmori, Y. Tachikawa and K. Yonekura, *8d gauge anomalies and the topological Green-Schwarz mechanism*, *JHEP* **11** (2017) 177 [[arXiv:1710.04218](#)] [[INSPIRE](#)].
- [23] Y. Lee and K. Yonekura, *Global anomalies in 8d supergravity*, *JHEP* **07** (2022) 125 [[arXiv:2203.12631](#)] [[INSPIRE](#)].
- [24] N. Seiberg, Y. Tachikawa and K. Yonekura, *Anomalies of Duality Groups and Extended Conformal Manifolds*, *PTEP* **2018** (2018) 073B04 [[arXiv:1803.07366](#)] [[INSPIRE](#)].
- [25] C.-T. Hsieh, Y. Tachikawa and K. Yonekura, *Anomaly of the Electromagnetic Duality of Maxwell Theory*, *Phys. Rev. Lett.* **123** (2019) 161601 [[arXiv:1905.08943](#)] [[INSPIRE](#)].
- [26] Y. Tachikawa, *Topological modular forms and the absence of a heterotic global anomaly*, *PTEP* **2022** (2022) 04A107 [[arXiv:2103.12211](#)] [[INSPIRE](#)].
- [27] Y. Tachikawa and M. Yamashita, *Topological modular forms and the absence of all heterotic global anomalies*, [arXiv:2108.13542](#) [[INSPIRE](#)].
- [28] A. Debray, M. Dierigl, J.J. Heckman and M. Montero, *The anomaly that was not meant IIB*, [arXiv:2107.14227](#) [[INSPIRE](#)].
- [29] C.-T. Hsieh, *Discrete gauge anomalies revisited*, [arXiv:1808.02881](#) [[INSPIRE](#)].
- [30] A. Debray, *Invertible phases for mixed spatial symmetries and the fermionic crystalline equivalence principle*, [arXiv:2102.02941](#) [[INSPIRE](#)].
- [31] P.H. Frampton and O.C.W. Kong, *Quark Mass Textures within a Finite Nonabelian Dicyclic Group*, *Phys. Rev. Lett.* **75** (1995) 781 [[hep-ph/9502395](#)] [[INSPIRE](#)].
- [32] A. Aranda, C.D. Carone and R.F. Lebed, *U(2) flavor physics without U(2) symmetry*, *Phys. Lett. B* **474** (2000) 170 [[hep-ph/9910392](#)] [[INSPIRE](#)].
- [33] E. Ma and G. Rajasekaran, *Softly broken A_4 symmetry for nearly degenerate neutrino masses*, *Phys. Rev. D* **64** (2001) 113012 [[hep-ph/0106291](#)] [[INSPIRE](#)].
- [34] G. Altarelli and F. Feruglio, *Tri-bimaximal neutrino mixing from discrete symmetry in extra dimensions*, *Nucl. Phys. B* **720** (2005) 64 [[hep-ph/0504165](#)] [[INSPIRE](#)].
- [35] P.D. Carr and P.H. Frampton, *Group Theoretic Bases for Tribimaximal Mixing*, [hep-ph/0701034](#) [[INSPIRE](#)].
- [36] F. Feruglio, C. Hagedorn, Y. Lin and L. Merlo, *Tri-bimaximal Neutrino Mixing and Quark Masses from a Discrete Flavour Symmetry*, *Nucl. Phys. B* **775** (2007) 120 [Erratum *ibid.* **836** (2010) 127] [[hep-ph/0702194](#)] [[INSPIRE](#)].
- [37] J. Davighi and N. Lohitsiri, *Anomaly interplay in U(2) gauge theories*, *JHEP* **05** (2020) 098 [[arXiv:2001.07731](#)] [[INSPIRE](#)].
- [38] J. Davighi and N. Lohitsiri, *The algebra of anomaly interplay*, *SciPost Phys.* **10** (2021) 074 [[arXiv:2011.10102](#)] [[INSPIRE](#)].
- [39] A. Grigoletto and P. Putrov, *Spin-cobordisms, surgeries and fermionic modular bootstrap*, [arXiv:2106.16247](#) [[INSPIRE](#)].

- [40] A. Grigoletto, *Anomalies of fermionic CFTs via cobordism and bootstrap*, [arXiv:2112.01485](#) [[INSPIRE](#)].
- [41] T. Banks and M. Dine, *Note on discrete gauge anomalies*, *Phys. Rev. D* **45** (1992) 1424 [[hep-th/9109045](#)] [[INSPIRE](#)].
- [42] B. Gripaios, *Gauge anomalies of finite groups*, *Phys. Rev. D* **105** (2022) 105008 [[arXiv:2201.11801](#)] [[INSPIRE](#)].
- [43] J. Talbert, *Pocket Formulae for Non-Abelian Discrete Anomaly Freedom*, *Phys. Lett. B* **786** (2018) 426 [[arXiv:1804.04237](#)] [[INSPIRE](#)].
- [44] T. Kobayashi and H. Uchida, *Anomaly of non-Abelian discrete symmetries*, *Phys. Rev. D* **105** (2022) 036018 [[arXiv:2111.10811](#)] [[INSPIRE](#)].
- [45] M.F. Atiyah, V.K. Patodi and I.M. Singer, *Spectral asymmetry and Riemannian geometry. 1*, *Math. Proc. Camb. Philos. Soc.* **77** (1975) 43.
- [46] M.F. Atiyah, V.K. Patodi and I.M. Singer, *Spectral asymmetry and Riemannian geometry. 2*, *Math. Proc. Camb. Philos. Soc.* **78** (1976) 405.
- [47] M.F. Atiyah, V.K. Patodi and I.M. Singer, *Spectral asymmetry and Riemannian geometry. 3*, *Math. Proc. Camb. Philos. Soc.* **79** (1976) 71.
- [48] D.S. Freed and G.W. Moore, *Setting the quantum integrand of M-theory*, *Commun. Math. Phys.* **263** (2006) 89 [[hep-th/0409135](#)] [[INSPIRE](#)].
- [49] J. McNamara and C. Vafa, *Cobordism Classes and the Swampland*, [arXiv:1909.10355](#) [[INSPIRE](#)].
- [50] H. Ooguri and T. Takayanagi, *Cobordism Conjecture in AdS*, [arXiv:2006.13953](#) [[INSPIRE](#)].
- [51] J. Kaidi, J. Parra-Martinez, Y. Tachikawa, with a mathematical appendix by A. Debray, *Topological Superconductors on Superstring Worldsheets*, *SciPost Phys.* **9** (2020) 10 [[arXiv:1911.11780](#)] [[INSPIRE](#)].
- [52] D.S. Freed, *Short-range entanglement and invertible field theories*, [arXiv:1406.7278](#) [[INSPIRE](#)].
- [53] S. Elitzur and V.P. Nair, *Nonperturbative Anomalies in Higher Dimensions*, *Nucl. Phys. B* **243** (1984) 205 [[INSPIRE](#)].
- [54] E. Witten, *Global Aspects of Current Algebra*, *Nucl. Phys. B* **223** (1983) 422 [[INSPIRE](#)].
- [55] The GAP Group, *GAP — Groups, Algorithms, and Programming, Version 4.11.0*, (2020).
- [56] W. Fulton and J. Harris, *Representation Theory*, Springer, New York, U.S.A. (2004).
- [57] R. Bruner, J. Greenlees and A. M. Society, *Connective Real K-Theory of Finite Groups*, *Mathematical Surveys and Monographs*, American Mathematical Society (2010).
- [58] C.-Y. Yu, *The connective real K-theory of elementary abelian 2-groups*, Ph.D. Thesis, University of Notre Dame, Notre Dame, U.S.A. (1995).
- [59] B. Botvinnik and P. Gilkey, *An analytic computation of $ko_{4\nu-1}(BQ_8)$* , *Topol. Methods Nonlinear Anal.* **6** (1995) 127.
- [60] D. Tong, *Line Operators in the Standard Model*, *JHEP* **07** (2017) 104 [[arXiv:1705.01853](#)] [[INSPIRE](#)].

- [61] D.W. Anderson, E.H. Brown, Jr. and F.P. Peterson, *The structure of the Spin cobordism ring*, *Ann. Math.* **86** (1967) 271.
- [62] M. Atiyah, R. Bott and A. Shapiro, *Clifford modules*, *Topology* **3** (1964) 3.
- [63] E.H. Brown, *Cohomology theories*, *Ann. Math.* **75** (1962) 467
- [64] J. Adams, *A variant of E.H. Brown's representability theorem*, *Topology* **10** (1971) 185.
- [65] A. Beaudry and J. A. Campbell, *A guide for computing stable homotopy groups*, [arXiv:1801.07530](#).
- [66] M. Guo, K. Ohmori, P. Putrov, Z. Wan and J. Wang, *Fermionic Finite-Group Gauge Theories and Interacting Symmetric/Crystalline Orders via Cobordisms*, *Commun. Math. Phys.* **376** (2020) 1073 [[arXiv:1812.11959](#)] [[INSPIRE](#)].
- [67] D. Bayen and R.R. Bruner, *Real connective k-theory and the quaternion group*, *Trans. Am. Math. Soc.* **348** (1996) 2201.
- [68] J. Martino and S. Priddy, *Classification of bg for groups with dihedral or quaternion sylow 2-subgroups*, *J. Pure Appl. Algebra* **73** (1991) 13.
- [69] D. Handel, *On products in the cohomology of the dihedral groups*, *Tohoku Math. J.* **45** (1993) 13.
- [70] P. Teichner, *Topological four-manifolds with finite fundamental group*, Ph.D. Thesis, Johannes-Gutenberg Universität, Mainz, Germany (1992).
- [71] P. Teichner, *On the signature of four-manifolds with universal covering spin*, *Math. Ann.* **295** (1993) 745 [[INSPIRE](#)].