

TOPICAL REVIEW • OPEN ACCESS

Remote sensing for monitoring tropical dryland forests: a review of current research, knowledge gaps and future directions for Southern Africa

To cite this article: Ruusa M David *et al* 2022 *Environ. Res. Commun.* **4** 042001

View the [article online](#) for updates and enhancements.

You may also like

- [Irrigation offsets wheat yield reductions from warming temperatures](#)
Jesse Tack, Andrew Barkley and Nathan Hendricks
- [Distinct vegetation response to drying and wetting trends across an aridity threshold](#)
Wei Zhao, Xiubo Yu, Yu Liu et al.
- [The trend shift caused by ecological restoration accelerates the vegetation greening of China's drylands since the 1980s](#)
Zidong Li, Shuai Wang, Changjia Li et al.

Environmental Research Communications



TOPICAL REVIEW

OPEN ACCESS

RECEIVED
17 November 2021REVISED
19 January 2022ACCEPTED FOR PUBLICATION
8 March 2022PUBLISHED
20 April 2022

Original content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.



Remote sensing for monitoring tropical dryland forests: a review of current research, knowledge gaps and future directions for Southern Africa

Ruusa M David , Nick J Rosser and Daniel N M Donoghue*

Department of Geography, Durham University, Science Laboratories, DH1 3LE Durham, United Kingdom

* Author to whom any correspondence should be addressed.

E-mail: rdavid5@wisc.edu, n.j.rosser@durham.ac.uk and danny.donoghue@durham.ac.uk**Keywords:** Remote sensing, Dryland forests, Forest monitoring, Synthetic Aperture Radar (SAR), Optical, Systematic review

Abstract

Climate change, manifest via rising temperatures, extreme drought, and associated anthropogenic activities, has a negative impact on the health and development of tropical dryland forests. Southern Africa encompasses significant areas of dryland forests that are important to local communities but are facing rapid deforestation and are highly vulnerable to biome degradation from land uses and extreme climate events. Appropriate integration of remote sensing technologies helps to assess and monitor forest ecosystems and provide spatially explicit, operational, and long-term data to assist the sustainable use of tropical environment landscapes. The period from 2010 onwards has seen the rapid development of remote sensing research on tropical forests, which has led to a significant increase in the number of scientific publications. This review aims to analyse and synthesise the evidence published in peer review studies with a focus on optical and radar remote sensing of dryland forests in Southern Africa from 1997–2020. For this study, 137 citation indexed research publications have been analysed with respect to publication timing, study location, spatial and temporal scale of applied remote sensing data, satellite sensors or platforms employed, research topics considered, and overall outcomes of the studies. This enabled us to provide a comprehensive overview of past achievements, current efforts, major research topics studies, EO product gaps/challenges, and to propose ways in which challenges may be overcome. It is hoped that this review will motivate discussion and encourage uptake of new remote sensing tools (e.g., Google Earth Engine (GEE)), data (e.g., the Sentinel satellites), improved vegetation parameters (e.g., red-edge related indices, vegetation optical depth (VOD)) and methodologies (e.g., data fusion or deep learning, etc.), where these have potential applications in monitoring dryland forests.

1. Introduction

1.1. Tropical dryland forest

Approximately 40% of the Earth's tropical and subtropical land surface is covered by open or closed forests. Of this, tropical dryland forests (TDFs) account for the largest share at 42%; the remaining 33% is moist forest, and only 25% is rain forest (Murphy and Lugo 1986, Janzen 1988). The largest proportion of dryland forests ecosystems are found in Africa, accounting for 60%–80% of the total biome area (three times the area covered by African rain forest) (figure 1) (Bullock *et al* 1995, Bodart *et al* 2013). Dryland forests hold a significant amount of terrestrial organic carbon that may contribute more to climate mitigation and adaptation than previously appreciated (Valentini *et al* 2014). Dryland forests also provide diverse ecosystem services, including water regulation and erosion control, the provision of food, fuel, and tourism opportunities (Djoudi *et al* 2015, Schröder *et al* 2021). On the other hand, dryland forests are subject to prolonged dry seasons and their rate of conversion to secondary forests has historically been higher than other tropical forest types




Figure 1. The graphic illustration shows the relative distribution of tropical dry forests. Reproduced with permission from FAO (1999).

(Pennington *et al* 2018). According to the Intergovernmental Panel on Climate Change (IPCC), these changes have impacts on carbon emissions to the atmosphere and forest biodiversity loss that reduce adaptive capacity and resilience to the impact of high temperatures and varying precipitation (IPCC 2014).

The definition of ‘dryland forest’ remains debatable and controversial, which contributes to the difficulty in accurately assessing and measuring its distribution patterns and status (Blackie *et al* 2014). The lack of a clear and comprehensive understanding of general terms including ‘drylands’ and ‘forests’ makes it a challenge to explicitly define dryland forests (Charles *et al* 2015). Given the fact that dryland forests progressively grade into other vegetation types such as moist tropical forests, woodlands, and savannas, also makes clear definitions complex (Putz and Redford 2010). Walter and Burnett (1971) noted that the accuracy of estimates of all tropical forest areas is constrained by uncertainty in the distribution of open woodlands in dryland areas, which are extensive in Africa, Australia, and Latin America.

In the scientific literature, many different names have been applied to tropical dryland forests, including savanna forests, Sudanian woodland and miombo woodland in Africa, monsoon forest in Asia, neotropical dry forests in South America (Linares-Palomino *et al* 2011, Suresh *et al* 2011, Chidumayo 2013). The neotropical dry forests in South America have a plethora of names from ‘caatinga’ in northeast Brazil, to ‘bosque tropical caducifolio’ in Mexico, and ‘cuabal’ in Cuba, which in part hinders comparisons (Sánchez-Azofeifa *et al* 2005, Mayes *et al* 2017). For example, Dexter *et al* (2015) identified dry deciduous forest in India (Suresh *et al* 2011), miombo woodland in southern Africa (Chidumayo 2013), and deciduous dipterocarp forest in continental Asia (Bunyavejchewin *et al* 2011) as a form of savanna, and not TDFs, despite the formal classification as TDFs by these studies, and the FAO (FAO 2001). The Caatinga and Chaco vegetation in Latin America is also considered by some authors as part of the dry forests (Pennington and Ratter 2006, Gasparri and Grau 2009), although Olson *et al* (2001) classifies these regions as a shrubland ecosystem.

There are several definitions currently available for TDFs, but there is still a lack of consensus in developing a common understanding. Mooney *et al* (1995) defined TDFs as forests occurring in the tropical regions characterized by pronounced seasonality in rainfall, where there are several months of severe, or even absolute drought. Sánchez-Azofeifa *et al* (2005) broadly defined TDFs as a vegetation type typically dominated by deciduous trees (at least 50% of trees present are drought deciduous), where the mean annual temperature is ≥ 25 °C, total annual precipitation ranges between 700 and 2000 mm, and there are three or more dry months every year (precipitation < 100 mm per month). A widely accepted definition is that of the FAO, which has identified TDFs as a Global Ecological Zone (GEZ), experiencing a tropical climate, with a dry period of 5 to 8 months and annual rainfall ranges from 500 to 1500 mm; GEZ includes the drier type mbo and Sudanian woodlands, savannah (Africa), caatinga and chaco (South America), and dry deciduous dipterocarp forest and woodlands (Asia) (FAO 2001). For the scope of this present review, we followed the FAO (2001) definition of

TDFs because it recognises forests occurring in the dry tropical climate globally including areas with relatively open canopies such as woodlands, and woody stands, then those based entirely on climate definitions. The growing body of evidence suggests that the current climate does not define the biogeography of TDFs or determine biome distributions (Staver *et al* 2011, Sunderland *et al* 2015), particularly in the context of future unprecedented climate change (IPCC 2007). If climates become sufficiently warmer and drier in the tropics, dry forests may expand into areas that are currently dominated by moist tropical forests (Putz and Redford 2010).

1.2. Recent research trends on tropical dry forests

1.2.1. Geographical research trends on tropical dry forests

Studies have pointed out that dryland forests generally receive a lower number of scientific publications and are under-represented in research in comparison with tropical moist forests (Miles *et al* 2006, Quesada *et al* 2009). Global reviews on dryland forests addressed the imbalance in the geographical coverage of dryland forest publications using remote sensing with certain tropical countries such as Latin America receiving the highest publications on dryland forests in comparison to most places in Africa (Blackie *et al* 2014, Schröder *et al* 2021). To investigate the geographical distribution of tropical dry forest studies, we initially searched for publications in ISI web of knowledge and Scopus on tropical dryland forests from Asia, Africa, America, and Australia. This search was conducted by using the keywords 'Dry Forest', 'Dryland Forest', 'Savan* Woodland', 'Savan* Tree', 'Dryland Vegetation', 'Dry Vegetation', 'Satellite', 'Remote Sensing', 'Optical', 'Radar', 'Image', 'SAR', 'Earth Observation', 'country/continent e.g., Africa'. In the search period from 1997 to 2020, we identified 1662 papers for Africa, 1639 for Australia, 1338 for America, and 1134 for Asia. In Africa, when we narrowed the search to individual countries, the results showed that about 743 publications are from the Republic of South Africa (RSA) while 355 publications were from the Sahel region of Nigeria. We also investigated scientific publications from other Southern African countries with dryland forest and 369 publications were identified, including from Botswana (87), Zimbabwe (69), Mozambique (60), Namibia (68), Zambia (49), Angola (24), Lesotho (6), Swaziland (5). When we combined the scientific publications from the above 8 Southern African countries, the results were 369 publications, indicating that publications on dryland forests for the Republic of South Africa were 2.01 times higher than all 8 Southern African countries combined. These results confirm that much less progress has been made in developing objective methods for assessing the rates of deforestation/conservation and threats to dryland forests ecosystems in most Southern African countries except for the Republic of South Africa.

The dryland forests in other parts of the world like Latin America are increasingly well studied at local, regional, national and continental scale, particularly with regards to carbon/biomass (Marin-Spiotta *et al* 2008, Chazdon *et al* 2016), fire (Pereira Júnior *et al* 2014, White 2019, Campos-Vargas and Vargas-Sanabria 2021), climate change (Mendivelso *et al* 2014, Castro *et al* 2018, González *et al* 2021), floristic and diversity composition (Gillespie *et al* 2000, Alvarez-Añorve *et al* 2012), ecosystem services (Castillo *et al* 2005, Paruelo *et al* 2016), Payment for Environmental Services (PES) (Alcañiz and Gutierrez 2020, Corbera *et al* 2009), novel conservation approaches (e.g., sustainable intensification for protected/conservation areas) (Méndez *et al* 2007, Reynolds *et al* 2016) and has the most comprehensive forest change/deforestation and biophysical aspects including species population changes, with extensive use of remote sensing (Trejo and Dirzo 2000, Gasparri and Grau 2009, Portillo-Quintero *et al* 2012, Stan and Sanchez-Azofeifa 2019, do Espírito-Santo *et al* 2020). In terms of reviews, many remote sensing reviews are providing valuable information on TDF's biophysical, ecological and socioeconomic at a regional level of Latin America (Castro *et al* 2003, Sánchez-Azofeifa *et al* 2003, Sánchez-Azofeifa *et al* 2005, Metternicht *et al* 2010, Portillo 2010, Sanchez-Azofeifa *et al* 2013, Quijas *et al* 2019, Stan and Sanchez-Azofeifa 2019), and Australia (Fensham and Fairfax 2002, Lawley *et al* 2016, Moore *et al* 2016). Also, reviews of current progress on dryland forests in individual countries can be found in many neotropics countries such as Mexico (Castillo *et al* 2005, Curry 2020), Venezuela (Fajardo *et al* 2005, Rodríguez *et al* 2008), and Costa Rica (Frankie *et al* 2004, Stoner *et al* 2004) enabling the identification of knowledge gaps and aiding in the development of a policy-relevant approach to conservation of these forests (Miles *et al* 2006).

Latin America is one of the best-represented areas for remote sensing research in dryland forests, for example, Portillo-Quintero and Sánchez-Azofeifa (2010) utilised remote sensing data at continental America, dryland forests ecoregion, and neotropics countries to show that 66% of tropical dry forest in the region has already been converted and that in some countries the conversion rate is as high as 86% and 95%, respectively. Aide *et al* (2012) using Moderate Resolution Imaging Spectroradiometer (MODIS) satellite data estimated that 200,000 km² of woody vegetation of Latin American and the Caribbean region were lost due to deforestation between 2001 and 2010. Nanni *et al* (2019) utilised MODIS satellite data at 250 m spatial resolution to assess reforestation at the regional level and reported that the reforestation hotspots cover 167,667.7 km² (7.6%) of Latin America between 2001 and 2014. While there are continental studies in Africa utilising remote sensing on biophysical parameters such as biomass/deforestation (Bodart *et al* 2013, Bouvet *et al* 2018), as compared to

Latin America, these studies may not consider the empirical observations of dryland forests extent/change per region or country level. In addition, most continental studies in Africa rather focus the attention on tropical rainforest in Central Africa (e.g., core Congolese forest) which may under-represent dryland forest (e.g., Aleman *et al* 2018). Global applications often report general land use/cover change which results in inaccurate or poor estimates of dryland forest (Aleman *et al* 2018, Smith *et al* 2019).

Several studies using optical and passive microwave instruments in the African Sahel (Olsson *et al* 2005, Horion *et al* 2014, Brandt *et al* 2016, Tian *et al* 2017) has reported that the density/size of woody vegetation stands have increased, with few areas in northern Nigeria reported to experience logging and agricultural expansion into forest reserves. Deforestation in Southern Africa is a major concern, with ca. 1.4 million ha of net forest loss annually, contributing to increased land degradation and the ensuing impacts on the balance of ecosystem function (Lesolle 2012). A global study by Tian *et al* (2017) utilising the optical Normalized Difference Vegetation (NDVI) index and passive microwave VOD across tropical drylands has reported a decreasing trend in woody vegetation in Southern African countries such as Botswana and Zimbabwe. Mitchard and Flintrop. (2013) conducted a coarse-scale analysis of changes in woody vegetation from 1982 to 2006 using NDVI time series from the Global Inventory Modeling and Mapping Studies (GIMMS) dataset and found that significant woody encroachment is occurring in most west African countries, but, in contrast, in Southern Africa, a rapid reduction in woody vegetation (deforestation) is occurring. Bodart *et al* (2013) used Landsat satellite imagery between 1990 and 2000 to estimate forest cover and forest cover changes in the African continent and found that 84% of the total deforested area occurred in the dry ecosystems of the Southern African region, with large spatially concentrated areas of forest loss found in Angola, Mozambique, Tanzania, Zambia and Zimbabwe, and isolated hotspots found in Nigeria and the border of the humid forest in Ghana. While such global and continental level studies are useful to highlight and reinforce the need to direct more attention and resources to these threatened/poorly studied ecosystems, research efforts on forest change/deforestation and climate change impacts of dryland forests at the regional level of Southern Africa are much harder to come by (Blackie *et al* 2014).

1.2.2. Remote Sensing approaches research trends in tropical dry forests

In recent decades, satellite remote sensing or Earth observation (EO) has proved a valuable tool in forest ecology, owing to its capability to perform systematic, frequent, and synoptic observation of the Earth, resulting in large data volumes and multiple datasets at varying spatial and temporal scales (Donoghue 2002, Zhu 2017). There are several sensors including multi-spectral scanners, laser scanners (LiDAR), hyper-spectral scanners as well as satellite-borne Synthetic Aperture Radar (SAR), that provide information on the colour and structure of forest environments (Donoghue 2002). EO has been applied to mapping the distribution, changes in cover, and condition including deforestation, desertification, fire damage, and climate impact (Smith *et al* 2019, Dogru *et al* 2020). Additionally, these data have been used to estimate biophysical characteristics such as total above ground biomass (AGB), leaf area index (LAI), woody area index, tree diameter, and canopy height which are key inputs into a variety of ecological models, as well as calculations of carbon balance and primary production (Donoghue 2000, Göksel *et al* 2018, Barbosa *et al* 2014). The continuous forest metrics obtained using EO data can be extracted at leaf and crown level to evaluate spectral elements of leaf or species properties and at stand-level and plot-level, or beyond to understand the variation between and among species, and through time (Muraoka and Koizumi 2009). Monitoring of dryland forest cover and forest metrics using EO data also helps to improve our understanding of the ecological drivers behind land cover change dynamics (Veldkamp and Lambin 2001, Chambers *et al* 2007).

Biomass has extensively been estimated based on the spectral reflectance values from two or more wavelengths, and the sensitivity of optical and near-infrared wavelengths to photosynthetic canopy cover has long been used for vegetation analyses (Rouse 1974, Tucker 1979). Spectral vegetation indices (VIs), including the NDVI index, are commonly used as a proxy of vegetation cover and have been shown to relate closely to LAI, biomass, and the fraction of photosynthetically active radiation absorbed by vegetation (fAPAR) (Curran 1980). Several well-known limitations of NDVI for robust estimation of biomass in drylands exist. NDVI is sensitive to green components and insensitive to woody components where the majority of carbon is stored (Tucker 1979). Also, AGB production is not always uniformly linked to either greenness or plant structure (herbaceous and woody compositions), as moisture content and vegetation species composition have been shown to impact the biomass-NDVI relationship (Asner *et al* 2009, Wessels *et al* 2006). These observations may help explain reportedly weak relationships between NDVI and tropical forest canopies, particularly for areas with complex and high vegetation amounts as in TDFs (Foody *et al* 2001, Sader *et al* 1989). For example, Madonsela *et al* (2018) investigated the interactions between seasonal NDVI and woody canopy cover in the savanna of the Kruger National Park (NP) to model tree species diversity using a factorial model and found that the interaction between NDVI and woody canopy cover was insignificant. These challenges have led to the development of alternative formulations which include correction factors or constants introduced to account for or minimize,

the varying background reflectance (Gitelson *et al* 1996, Huete *et al* 1999). The Enhanced Vegetation Index (EVI) is a modification of NDVI that provides complementary information about the spatial and temporal variations of vegetation while minimizing many of the contamination problems present in the NDVI, such as those associated with canopy background and atmospheric influences (Huete *et al* 2002). Other closely related indices include the Simple Ratio (SR), the Green Normalized Difference Vegetation Index (GNDVI), Soil-Adjusted Vegetation Index (SAVI) amongst others. Xue and Su (2017) provide a detailed review of vegetation indices.

Although vegetation monitoring has been largely based on the multispectral 'greenness' indices, which have proven invaluable for monitoring biophysical and biogeochemical parameters, it has been widely reported in the literature that they suffer from several weaknesses in dryland ecosystems (Tian *et al* 2016, Shi *et al* 2008). Other remote sensing systems such as the passive microwave-based satellite systems capture the biomass signal in the parameter termed vegetation optical depth (VOD) which has been used to monitor changes in vegetation dynamics (Andela *et al* 2013, Brandt *et al* 2018a, 2018b). Unlike the optical remote sensing-based vegetation indices that are sensitive to chlorophyll abundance and photosynthetically active biomass of the leaves, the vegetation information (e.g., VOD) deriving from passive microwave instruments is sensitive to the water content in the total aboveground vegetation, including both the canopy (e.g. woody plant foliage) and non-green woody (e.g. plant stems and branches) components due to greater penetration and sensitivity (Shi *et al* 2008, Liu *et al* 2011). The passive microwave observations VOD is relatively insensitive to signal degradation from solar illumination and atmospheric effects and provide a valuable alternative tool for rapid monitoring of carbon stocks and their changes (Jones *et al* 2011). One of the advantages of passive microwave-derived VOD is that it continues to distinguish biomass variations at a relatively high biomass density, as compared to optical-based vegetation indices which are likely to become saturated over dense canopies (Jones *et al* 2011, Liu *et al* 2015). The main disadvantage of passive microwave observations is the relatively coarse spatial resolution (>10km), as compared to satellite data in the visible and near-infrared parts of the spectrum; however, these data still have highly useful applications at regional and global scales (Liu *et al* 2015, Rahmoune *et al* 2013, Owe *et al* 2001). Some recent global and local studies from Latin America and Africa in the dryland ecosystems found VOD to be more robust against the NDVI drawbacks of saturation effect and continues to distinguish structural differences for vegetation with a near-closed canopy when used as a proxy for vegetation productivity (van Marle *et al* 2016, Cui *et al* 2015, Liu *et al* 2011, Tian *et al* 2016). Apart from the VOD and NDVI, an intercomparison between several vegetation indices including other passive microwave-based vegetation indices, such as the Microwave Polarization Difference Index (MPDI) (Becker and Choudhury 1988), and the Microwave Vegetation Indices (MVIs) (Shi *et al* 2008) would be of benefit in monitoring dryland biomes.

Due to the inherent trade-offs between spatial and temporal resolution in EO data, and geographic coverage, vegetation patterns on both spatial and temporal domains have been revealed by various technological advances resulted in the growing availability of remote sensing data and methods (Toth and Józków 2016, Zhou *et al* 2020). The application of non-parametric machine learning regression algorithms, such as decision trees, random forests (RF), support vector machines (SVMs), and k-nearest neighbour have become more predominant and demonstrate the ability to outperform widely used parametric approaches, such as polynomial and multiple linear regression variables used with remotely sensed data in a forest environment (Breiman 2001, Latifi *et al* 2010). More recently, deep learning, a branch of machine learning that stems from cognitive and information theories (e.g., convolutional neural network (CNN) founded by Schmidhuber (2015) has been highlighted as a feasible approach for handling complex data in remote sensing including large-scale image recognition, semantic segmentation, classification, and object detection tasks (Kattenborn *et al* 2021, Shafaey *et al* 2018). Non-parametric machine and deep learning models are sufficiently versatile to uncover complicated nonlinear relationships and able to extract combinations of the input data that are difficult to describe explicitly by humans, particularly, in areas with high structural variability such as dryland forests (Hastie *et al* 2009, Shao *et al* 2017). Deep learning has been used by many remote sensing studies to provide in-depth forest investigation from the perspectives of hyperspectral image analysis, interpretation of SAR/ LiDAR images, interpretation of high-resolution satellite images and classification, and multimodal data fusion (e.g., the fusion of Hyperspectral, SAR, LiDAR and optical data (Guirado *et al* 2020, Kussul *et al* 2017, Shao *et al* 2017, Liao *et al* 2018, Trier *et al* 2018, Narine *et al* 2019). Improved techniques in remote sensing such as VOD, and machine and deep learning have been utilised to estimate dryland forest attributes globally and other dryland ecosystems, however, very few of these focused on the local and regional scale of Southern Africa (e.g., Symeonakis *et al* 2020). The uncertainties reported in many dryland forests studies (Bastin *et al* 2017), could be decreased following further development, application, and comparison of these improved approaches in future works at local, regional, continental studies in Southern Africa and other dryland forest ecosystems. Critically, an increase in the spatial, spectral, and radiometric resolution of satellite sensors, increased availability of EO data and computational resources combined with the machine or deep learning techniques would enhance the potential dryland forest information to be exploited (Ali *et al* 2015). For a detailed review of machine learning and deep learning for remote sensing and Sustainable Development Goals, see Zhu *et al* (2017) and Holloway and Mengersen (2018).

1.3. Review focus justification

The majority of the residents of Southern Africa are poor and about 75% of them live in rural areas with high reliance on dryland forests (Bond *et al* 2010). Additionally, these dryland areas display a high susceptibility to bush encroachment (O'Connor *et al* 2014) and economic reliance on tourism (Ferreira 2004) and forest products (Kamwi *et al* 2020), which means that both agriculture and tourism development encroach on the dryland forests, resulting in loss of forest biodiversity and land degradation (Eva *et al* 2006, Petheram *et al* 2006). Across Southern Africa, sustainable management of dryland ecosystems is hindered by complex land tenure due to historical legacy, weak links between policy and woodland use and management, and cultural drivers (Dewees 1994, Balint and Mashinya 2006). Also, the dryland ecosystems of Southern Africa are dominated by private land ownership, a high concentration of wildlife and human populations, and agriculture where TDFs occur (Child *et al* 2012). This review focuses on Southern Africa because there is a gap in knowledge on carbon storage, biomass, and the long-term trend of forest distribution and degradation in dryland forests. Much of the research on dryland forests in Southern Africa has concentrated on livelihoods, ecosystem services, energy supply and demand, food security, livelihoods and community forest management, and conservation/development trade-offs (e.g., Chidumayo and Marunda 2010, Chidumayo and Gumbo 2010, Chidumayo 2019, Djoudi *et al* 2015, Dewees 1994, Du Preez 2014, Ryan *et al* 2016), leaving forests highly vulnerable to deforestation and degradation (Keenan 2015). The social and economic aspects are important given the large numbers of African people that rely on dry forests for their livelihoods and a range of goods and services. However, the gap in biophysical aspects, threats status, and adaptation to climate change identified for Southern African TDFs at the regional and national level (Blackie *et al* 2014, Sunderland *et al* 2015), presents an urgent need for an assessment of the effectiveness of the EO scientific foundation on current understanding of TDFs in Southern Africa; this can aid in the development of policy-relevant approaches and long-term, regional perspective for planning and conservation of the TDFs.



With the prospects of multiple free datasets from optical and SAR sensors being available; combining information from optical sensors on photosynthetic activity (e.g., through various vegetation indices) with SAR-derived information on forest structure and volume brings the benefits of higher spectral resolution, and compensating for the shortcomings of using single data products alone. Based on this hypothesis, this review focuses on examining the studies using optical and SAR sensors, both individually and the combination of the two types of EO data in monitoring tropical forests. While forest distribution, carbon storage, and reducing emissions from deforestation and forest degradation (REDD+) related research exists in African dryland forests, the geographical focus has tended to be confined to several West/Central African countries, whereas Southern Africa is relatively poorly analysed (Lewis *et al* 2013, Sunderland *et al* 2015). Although numerous reviews have been conducted discussing the application of optical and radar remote sensing, they are either concentrated on mangroves forests (Kuenzer *et al* 2011, Wang *et al* 2019), rain forests (Dupuis *et al* 2020), or ecosystem services (Barbosa *et al* 2015). To date, reviews on remote sensing and EO in Southern Africa have focused on research conducted in the Republic of South Africa (Hoffman and Todd 2000, Mutanga *et al* 2009, Mutanga *et al* 2016).

As shown in figure 2, the climate threats coupled with a growing human population and future anticipated changes in land use are predicted to lead to severe dry forest biome shifts and degradation across the whole of Southern Africa, hence the need to expand the geographical scope of this review from previous work (IPCC 2014, King 2014). This paper provides a systematic review of the scientific literatures related to the use of Earth observation data including SAR and optical sensors used to study dryland forests, with a focus on Southern Africa. To achieve this, we present examples from the literature that summarise past achievements, current efforts, and knowledge gaps. The objectives of this review are to (i) to provide a detailed overview of the current approaches and limitations for monitoring dryland forests using optical and radar remote sensing data. (ii) to provide a critical evaluation and synthesis of the literature monitoring dryland forests using remote sensing data and discuss how EO data can contribute to dryland forest monitoring and forest conservation in Southern Africa. (iii) to identify knowledge gaps and make recommendations for research that will enhance monitoring of dryland forests using remote sensing data.

2. Remote sensing applications in dryland forest

2.1. Optical data

In broad terms, the satellite platforms developed over the past 40 years (since 1972) have carried two broad types of sensor systems; passive optical and active synthetic aperture radar (SAR). Successful change detection and parameter estimation over tropical dryland forests require: (a) correct selection and application of sensor type; (b) coupling with field observation data for calibration and validation, and (c) data integration and appropriate techniques for modelling (figure 3). Optical sensors have been widely used for land cover and forest resource mapping, providing access to long-term data dating back to the launch of Landsat ERTS (Earth Resources



Technology Satellite) satellites in 1972. Landsat and several other coarse/medium spatial resolution optical sensor missions (National Oceanic and Atmospheric Administration (NOAA) - Advanced Very High-Resolution Radiometer (AVHRR); Indian Remote Sensing Satellites-1C/1D (ISRO-IRS-1C/D); the National Aeronautics and Space Administration (NASA) - Aqua/Terra- MODIS; Sentinel-2) provide well-calibrated, nadir-viewing, near-global systematic coverage which have built up a valuable archive of image data that can be used to analyse ecosystem dynamics (Donoghue 2000, Congalton 2018). In 2014, ESA launched the Multispectral Instrument (MSI) onboard Sentinel-2 as part of its Copernicus EO mission. Sentinel-2 MSI uses two identical satellite sensors to measure the Earth's reflected radiance with a revisit time of 5 d and a high spatial resolution of 10–20 m pixel size. The length of the Sentinel-2 archive is short (from 2015), compared to the Landsat mission from 1972–present, NOAA-AVHRR 1979–present, Satellite Pour l'Observation de la Terre VEGETATION (SPOT/VGT) (1998–present), IRS-1C/1D (ISRO-IRS-1C/D) (1995–2010), ENVISAT - Medium Resolution Imaging Spectrometer (MERIS) (2002–2010), the NASA - MODIS (2000–present) and the French Space Agency (CNES-Centre national d'études spatiales) high-resolution SPOT satellite constellation (6 m–20 m pixel size) - SPOT-1 in 1986–1990, SPOT-2 in 1990–2009, SPOT-3 in 1993–2009; SPOT-4 in 1990–2013; SPOT-5 in 2002–present; SPOT-6 in 2012–present; SPOT-7 in 2014–present. The VEGETATION 1 (VGT 1) (1998–2012) and VEGETATION 2 (VGT 2) (2002–2014) instrument on the SPOT 4 and SPOT 5 (SPOT/VGT) satellites provided global daily monitoring of vegetation cover, and its successor the European PROBA-V satellite (2013–present), with a pixel size of 1 km, 300 m and 100 m are supplied by the VEGETATION image Processing Centre (CTIV) of VITO (Belgium), which can be accessed through the internet site <http://free.vgt.vito.be>. Although a large number of satellite sensors have been launched that are capable of observing land dynamics, and their pixel size has increased from 80 m of the Landsat-1 to 0.41–1.65 m of the GeoEye-1 satellites (Aguilar *et al* 2013), very few sensors provide well-calibrated multispectral, nadir-viewing observations and even fewer systematically capture all global data and provide a long-term archive of data free of charge to the public. Except for AVHRR and Landsat, no other sensor or sensor line offers the chance of long-term monitoring of an area to be monitored back in time to the 1970s, covering about four decades.

There are several non-systematic commercial high-resolution satellites that allow the detection of individual trees or populations. Maxar Technologies Inc. launched 4 very high resolution satellites - WorldView-1 in 2007, WorldView-2 in 2009, WorldView-3 in 2010, and WorldView-4 in 2019 that acquire images with spatial resolution of 0.5, 0.41, and 0.31 m, respectively. From 2009 onward, Planet labs launched a swarm of micro-satellites including PlanetScope (PS), RapidEye (RE), and SkySat (SS) Earth-imaging constellations with multispectral imaging capability with the aim of acquiring daily image capture for any part of the world at a spatial resolution of 3.125 m to 6.5 m (Marta 2018). In 2011 and 2012, the Space Agency of France (CNES) launched the Pléiades—high resolution optical imaging satellite constellation (Pléiades-1A and Pléiades-1B), with a high spatial resolution of 0.7–2.8 m. Other very high resolution commercial space imaging satellites include Earlybird (1997), EROS-A (1998), IKONOS (1999), QuickBird (2001), OrbView (2001), GeoEye (2008) (Maglione 2016). In Africa, South Africa started satellite developments in the 1990s, with the successful launch of SunSat-1 with a spatial resolution of 15 m in 1999 and SumbandilaSat low orbit satellite with a high spatial resolution of 6.25 m in 2009 (Cho *et al* 2012, Mutanga *et al* 2016). While the first Nigerian satellite, a microsatellite called NigeriaSat-1, was successfully launched into low earth orbit in 2003, followed by Nigeriasat-2 with a higher spatial resolution of 2.5–5 m, built by Surrey Satellite Technology Limited (SSTL) of UK (Agbaje 2010).

Nevertheless, the use of data acquired by higher spatial resolution optical sensors, particularly at regional and global scales, can be limited by their relatively high cost, huge data volumes, and low frequency of data acquisition compounded further in tropical regions where cloud cover is prevalent (Zhu and Woodcock 2012, Lehmann *et al* 2015). The temporal resolution of sensors has also increased from, for example, 16 d for Landsat to nearly 1 d for the NOAA-AVHRR, NASA-Aqua/Terra-MODIS, SPOT/VGT, and/or ENVISAT-MERIS data, but with a coarse spatial resolution of 250 m to 1 km (Arino *et al* 2007, Herold *et al* 2008). Although lacking high spatial detail, the daily temporal resolution of such sensors enables frequent estimation of deforestation, detection of disturbances using dense time series data, and enables gaps due to cloud cover to be overcome (Mbow *et al* 2015). It is important to mention that the acquisitions of some satellites such as IRS-1C/1D, and MERIS ceased operations, however, the Sentinel, MODIS, NOAA-AVHRR, SPOT, SPOT-VGT (PROBA-V), and Landsat series continue to operate, with ongoing continuity of data collection ensured with the recent launch of Landsat-9 in September 2021.

2.2. Synthetic aperture radar (SAR)

SAR sensors for civilian applications first appeared in 1978 with NASA's SeaSat but have grown in importance as a tool for forest studies. SAR sensors can operate at different frequencies and polarisations; these system parameters provide information on the roughness and scattering properties of forest canopies and data can be

captured day and night independent of weather conditions (Durdin *et al* 1989). Since SAR can penetrate cloud, rain, smoke, and haze, and it is a valuable source of data when atmospheric conditions hamper optical data capture, particularly in the tropical dryland forest such as Southern Africa where the cloud and smoke from forest fires are prominent features (Le Canut *et al* 1996). Radar signals are sensitive to moisture, variations, surface roughness, and vegetation structure properties, whereas data from optical systems use characteristics related to reflected solar illumination or surface temperature (for thermal infrared sensors) as a basis for discrimination of the land cover (Kasischke *et al* 1997, Mitchard *et al* 2009). Cloud cover-free SAR images have great potential in the dryland tropical areas but have been used less often for forest monitoring applications compared to optical imagery, partly because of the scarcity of data (Castro *et al* 2003). Since the launch of the Sentinel-1A and B, dense SAR time-series data are now available over tropical forest areas freely and openly, with systematic acquisitions at a 10 m spatial resolution and a 6–12 d revisit time (dependent on the location) in all weather conditions.

Over the last 30 years, several satellite-borne SAR has been launched, including the United State Spaceborne Imaging Radar-Synthetic Aperture Radar (SIR-C/X-SAR), European Remote Sensing (ERS-1/-2), ESA's Envisat ASAR (Advanced Synthetic Aperture Radar), Advanced Synthetic Aperture Radar (ASAR), Japanese Earth Resources Satellite (JERS-1), Advanced Land Observation Satellite (ALOS/PALSAR-1/-2), German TerraSAR-X, Italy's Cosmo SkyMed, and the Canadian RADARSAT-1/-2 (Shimada 2018). Depending on the sensor configuration, a single channel (wavelength/frequency) or multiple channels may be recorded in either single or multiple polarizations. Generally, studies have reported that the longer the wavelength (e.g., P (30–100 cm) and L (15–30 cm)), the further is its penetration into the forest and the greater the importance of scattering beyond the upper canopy (Huang *et al* 2015). Besides the greater sensitivity of longer radar wavelengths to forest structure, different studies indicate that cross-polarized backscatter (HV-horizontally transmitted, and vertically received, VH-vertically transmitted and horizontally received) often exhibits greater sensitivity to forest biomass than like-polarized backscatter (co-polarized bands: HH-horizontally transmitted and horizontally received, VV-vertically transmitted and vertically received) (Kasischke *et al* 1997).

2.3. Limitations of optical and radar, and benefits of combining sensors

Despite the different generations and types of satellite sensors, no one sensor currently meets fully the requirements of a comprehensive forest resource assessment EO system. The selection of an appropriate source of data requires first the identification of the ecological question being asked, identification of the limitations and advantages of each sensor. The varying temporal, spatial, spectral, and radiometric resolutions unique to the individual sensor system, result in different advantages and disadvantages to the monitoring of dryland ecosystems (Lu 2006). Optical data are limited in the monitoring of this forest type. For example (1) cloud and smoke severely limit the use of optical products (Le Canut *et al* 1996); (2) Dramatic seasonal changes in the dryland forests conditions including droughts and leaf shedding make it unsuitable for systematic all-season monitoring of this forest type (Boggs 2010). One of the reasons for this is associated with the seasonality of the tropical vegetation: during the wet season, cloud-free satellite imagery is difficult to acquire, while during the dry season when the imagery is more available, the leaf-off configuration of the forest causes misclassification with savanna shrubland or grassland; (3) Optical data is sensitive at the early stages of growth but as forest canopies close, reflected radiation is no longer sensitive to biomass as the reflectance signal saturates at higher biomass values (Lu 2006); (4) Passive optical sensors only detect the surface top layer, meaning that forest canopy obscures the understory, and similarly grasses/crops obscure soil; (5) Changes in the spectral properties of the soil and atmosphere can also hinder the inference of forest cover properties (Wang *et al* 1998, Santos *et al* 2002).

Similarly, there are a number of challenges to analysing and interpreting radar images for tropical forest applications, which include: (1) Difficulty in interpreting radar backscatter, including, for example, speckle, which is unwanted random noise inherent in all SAR images, which may increase measurement uncertainty and make interpretation difficult (Klogo *et al* 2013); (2) Topography is a major limitation in mountainous regions due to geometric and radiometric effects such as radar shadowing caused by foreshortening and layover when the satellite is not able to illuminate the whole ground surface (Mitchard *et al* 2009); (3) SAR observations often lack a long-term and dense time series because they demand a relatively high energy provision on satellite platforms. Until recently, satellite-based SAR data for multi-temporal assessments over large areas were constrained by low spatial and temporal coverage at medium resolution, although this now may be overcome with acquisitions from the recently launched C-band Sentinel-1 and L-band ALOS-2 satellite missions (Reiche *et al* 2016).

Rather than using EO data from a single satellite sensor, the synergy of remotely sensed data from multiple sensors, particularly SAR systems with those acquired by optical sensors, has been shown to be beneficial for forest resource assessment (Lehmann *et al* 2015). Because optical data is capable of measuring the reflectance of the topmost layer of the forest canopy and SAR data deliver useful within-canopy biophysical parameters


without being affected by cloud cover and weather conditions, one dataset may compensate for the shortcomings of the other (Reiche *et al* 2016). Previous research indicated that integration of optical and radar can improve land and forest cover characterisation (Symeonakis *et al* 2018a). For example, the fusion of optical and radar sensor data has the potential to improve AGB estimation because it may compensate for the mixed pixels and data saturation problems in a tropical forest area. In addition to the spectral synergy afforded, the cloud penetrating capability of microwave radar sensors allows areas that have missing optical data to be included in analyses, particularly if multi-temporal methods are being employed (Reiche *et al* 2016).

3. Methodology

This review focused on scientific papers studying tropical dryland forests and made use of remote sensing data to monitor and estimate changes in dryland forests. Airborne remote sensing studies were excluded from this review process, since the review's major focus lies on satellite Earth observation of dryland forests and because the acquisition of airborne sensors have low area coverage and high cost per unit area of ground coverage (e.g., the airborne hyperspectral images), making them spatially and temporally limited in most African countries. The systematic search approach taken to querying the literature was carried out by making use of selective keyword searches in the form of structured queries using field tags and Boolean operators through the Web of Science (<http://apps.webofknowledge.com>) and Scopus (<http://www.scopus.com>) databases. At each query, terms, and keywords such as 'Dryland forests', 'Savan*', 'Woodland', 'Tree', 'Vegetation', 'Satellite', 'Remote Sensing', 'Optical', 'Radar', 'Image', 'SAR', and 'Earth Observation' were used to produce an extensive list of articles, where * is a wildcard search. The results were further refined with keywords such as 'Forest change', 'Degradation', 'Deforestation', 'Trend', 'Biodiversity', 'Phenology', 'Biomass', 'Structural parameter', and also keywords representing the countries in Southern Africa, such as 'Botswana', 'Namibia', 'Mozambique', 'South Africa', to provide a comparison in terms of the numbers of studies undertaken across the region. Within the context of this review, all research articles were categorized into eight categories, including: 'Land-use/land-cover', 'Forest cover/types', 'Biomass', 'Forest structure', 'Biodiversity/habitats', 'Phenology', 'Plant traits', and 'Disturbances'. Articles with a publication date between 1997 and 2020 were considered, capturing a period of two decades within the review, based on a broad set of inclusion criteria:

1. The paper should address dryland forests and remote sensing as either main or secondary subjects.
2. The selection terms and keywords should exist as a whole in at least one of the fields: title, keywords, and abstract.
3. The paper should be published in a peer-reviewed scientific journal.
4. The paper should be written in the English language.

During our data extraction process and literature search, we aimed to find studies meeting the criteria for peer-reviewed publications, available through the chosen indexed bibliographic databases. For this reason, our literature search did not include general non-scientific reports, books, grey literature, thesis documents or dissertations, extended abstracts, or presentations. The initial steps of the search process returned 1,478 published articles. Additional publications were added to the total set of studies by identifying relevant literature found in the reference lists of these selected papers that conform to the inclusion criteria. The review methodology was guided by the Guidelines for Systematic Review and Evidence Synthesis in Environmental Management (Collaboration for Environmental Evidence 2013). A systematic review and meta-analysis were undertaken and framed based on the PICO (population, intervention, comparison, outcomes) model (McKenzie *et al* 2019) and reported using PRISMA (Preferred Reporting Items for Systematic reviews and Meta-Analyses) flow diagram (Moher *et al* 2009). The 1,478 articles were reduced to 870 articles as we selected for inclusion in the review only the studies that had a full text available in English, papers published in peer-reviewed journals, and removing all repetitions across databases. Initially, the titles and abstracts were screened to assess eligibility by searching for predefined keywords and terms of the abstract or summary, identifying terms 'dry or dryland forests' and the country or countries where the research took place. In this way, studies not conducted in Southern Africa or dryland forests were filtered out, which reduced papers from 870 to 599 papers. The screening was followed by a full-text assessment that reduced the papers to 270 by excluding studies that, for example, mentioned the term 'dryland forest' once in the abstract but did not investigate dryland forests, as outlined in the PRISMA flow diagram in figure 4. The search was subsequently refined by assigning the papers to each of the study aims they addressed and to each category for the variables identified in the search protocol, reviewing the methodologies of each publication, excluding them from further analysis if they did not meet the inclusion criteria on review. These steps reduced the total number of entries to 137 scientific publications. The



selected literature was reviewed systematically, searching for specific information regarding the publication temporal development, study location, remote sensing sensor/platform used, spatial and temporal coverage, remote sensing product (e.g., biophysical indices) used, and application areas of the study (e.g., land cover, forest biomass). The parameters used to extract relevant information from the remaining 137 identified scientific publications are in table 1. Figure 4 is a PRISMA schematic representation of the methodology used and the derivation of the final number of articles selected.

4. Results

4.1. Temporal development of publications and author affiliations

After the literature search, we found that the cumulative number of published research papers integrating remote sensing data in dryland forests of Southern Africa grew exponentially from 2 in 1997 to 155 in 2020. The temporal development of the 137 investigated research articles is illustrated in figure 5. The graphic shows that the number of studies has increased significantly over the last 23 years, with the majority of the studies published from 2013. More than 105 (80%) of articles were published from 2009 to 2020 and only 4 (3%) of articles were published before 2000. The growth in number is also related to the increased availability of remote sensing platforms, sensors, data, for example, Landsat 8 in 2013 and Sentinel satellite in 2014, respectively.




Figure 5. Number of papers included in the review integrating remote sensing and dryland forests in Southern Africa published annually between 1997 and 2020.

Table 1. Parameters used to extract relevant information for this review.

General information

Paper Id
 1st author's institution
 Research institute city
 Publication year
 Publishing Journal
 Journal category
 No of Citation
 Study type

Site specific information

Location of the study area
 Study country
 Forest management area
 Predominant forest type


Information on remote sensing data

Sensor Type
 Instrument name
 Image resolution
 Time period observed
 Temporal resolution of EO data
 Database used

Information on research

Research topic considered:
 Forest cover/type, disturbance, phenology, biodiversity/habitats, plant traits, land cover/land use
 Parameters examined in the study
 Examined object scale
 Applied methodology
 Information on validation and accuracy of results
 Database used

In the review, we have only considered studies within Southern Africa; however, the majority of first authors, 83 (61%) of 137 investigated papers, are mainly scientists from international research institutions outside of the focus region, mainly the USA, UK, Portugal, Germany, and The Netherlands (figure 6). Conversely, the majority of first author institutions from Africa, 37 (27%) of published papers, were from RSA research institutions. The state funded research institutions in Southern Africa shown in figure 6 include South African Council for



Scientific and Industrial Research (CSIR), South African National Space Agency (SANSa), Water Resource Commission of South Africa, South Africa Agricultural Research Council, Range and Forage Institute, Botswanan Harry Oppenheimer Okavango Research Centre, Desert Research Foundation of Namibia, and Namibia Ministry of Environment and Tourism. Considering the 137 studies conducted, about 120 (90%) of the first authors are affiliated with either International and RSA institutions, but no first authors were from Zambia, Lesotho, or Angola.

4.2. Spatial coverage, spatial extent, and investigated protected areas

Looking at the spatial scale of the study areas, we distinguished between studies done at a local community level in a single country, termed local scale, and studies done at more than one local community or province termed regional scale. We also considered studies done at the national level and the whole of Southern Africa. If a study covered more than three countries, it was counted as an analysis of Southern Africa. The spatial extent of the studies in the review is shown in figure 7. The majority 88 (64%) of the investigated studies focused on a local scale, despite the need for regional scale information on dryland forest distribution. From figure 7, out of 137 investigated research papers, 20 (15%) and 13 (9%) research papers covered regional and national scales, respectively. Only 10 (7%) out of 137 research papers dealt with transboundary protected areas, while 6 (4%) of research papers were covering Southern African, considering the region as a whole, using mainly multispectral data of large spatial resolution of 1km to 8km (MODIS, SPOT, and AVHRR) to generate information on phenology, and vegetation condition (fire or drought), as shown in figure 7.

From figure 8, it is evident that considerable gaps in geographical focus of research on tropical dryland forests mapping still exist in Southern Africa. With respect to spatial coverage of the research, most studies, 50 (36%) of research papers were carried out in RSA, followed by Namibia and Botswana, with 22 (16%) and 18 (13%) of research papers, respectively. Swaziland, Angola, and Lesotho were the least frequently investigated, each with < 10 papers. Angolan dryland forests are even less well studied with 4 (6%) of research papers, despite being found extensively in that country. Figure 8 also shows the location of the most frequently studied protected areas. By far, the most studied was the Kruger National Park (NP) in RSA, involving research by local and foreign researchers from as far afield as the USA, the UK, and beyond. With this interest in the Kruger NP, there is, unfortunately, a lack of attention on other conservation areas and parks in Southern Africa. Kruger NP was the only subject of more than one-third, 23 (37%) of the 61 of all reviewed papers on protected areas. The second most frequently studied protected areas are the Etosha NP in Namibia with 6 (8%) of papers, Chobe NP with 4 (7%) of papers, and Kwando, Kavango and Zambezi transboundary NP with 8 (13%) of papers). Malipati Safari Area, South Luangwa NP, Gorongosa NP, and Central Kalahari Game Reserve were each studied 3 (5%) and 2 (3%) times.




Figure 7. Spatial extent of investigated studies.





Figure 8. Number of studies per country and National Park in Southern Africa. (Note: The data are not scaled to the proportion of dryland forest area of countries, and National Parks with fewer or no publications are not shown. Source: FAO (1999). Reproduced with permission from FAO.

To identify land surface changes and the drivers behind these, as well as short- and long-term trends, it is essential that EO temporal coverage has sufficiently frequent revisit periods and resolutions. Nonetheless, this is not an easy task since the availability of remote sensing data for long-term monitoring is constrained by sensor



characteristics (e.g., revisit time) and environmental factors (e.g., cloud cover). Looking at the temporal resolution of the EO datasets used, we distinguished between data acquired at a single point in time on a monthly basis, termed mono-temporal analyses, and on a single annual basis, termed mono-annual analyses. We also considered multi-temporal and multi-annual to separate monthly and yearly analyses studies. From figure 9 it is seen that the majority of published material has focused on a single temporal period. The majority of studies involved mapping over two or more years (multi-temporal/multi-annual) comparing images at two or more different times, with a bi-temporal approach based on discrete classification (e.g., Chiteculo *et al* 2018, Matavire *et al* 2015, Coetzer-Hanack *et al* 2016). Although the bi-temporal approach is mathematically simple and does not require large data storage, it is less useful compared to the time series approach that can provide a more comprehensive understanding of the complexity of the Earth's land surface dynamics. Very few studies feature time series analysis, which is required to perform continuous long-term monitoring of changes in a tropical forest ecosystem. The majority of articles on time series analysed multi-annual data, which masks within-year variations, as compared to the detail provided at a monthly temporal scale (e.g., Wessels *et al* 2006, Verlinden and Laamanen 2006a, Akinyemi and Kgomo 2019, Venter *et al* 2020). Only 22 (16%) out of the 137 studies analysed more than 15 years and only 11 (8%) studies covered more than 20 years using monthly time series (e.g., Bunting *et al* 2018, Schultz *et al* 2018).

4.3. Research topics

We have classified the large number of research topics into eight broad categories that cover the diversity of research into dryland forests. The eight categories, and the number of studies belonging to each of them, are shown in figure 10.

4.3.1. Land cover/land use

Land-cover change is one of the most researched areas using EO in Southern Africa, with 36 (23%) publications making it the second most common topic. We considered land-use/cover describing land surface classification, typically represented in thematic maps of different dryland vegetation. Land-use/cover changes with a specific focus on other dryland vegetation such as rangelands, grassland, coastal vegetation, or plantation forests without covering dryland forests were excluded. The majority of publications on land-use/land-cover used optical data. For example, Landsat data have been used by more than 90% of publications, except Daskin *et al* (2016) and Hüttich *et al* (2011) which used RapidEye and MODIS data. Only one publication used a combination of Radar and optical data (Symeonakis *et al* 2018a). Sentinel data have not been utilised for land cover and land use study in the reviewed papers, probably due to the relatively recent availability of these data. Looking at scale, the majority of papers on land-cover change focused on the local scale in Southern Africa, but there is still a general lack of synthesis of land-use/cover change assessment at the regional, national or subcontinental scale (figure 7).

4.3.2. Forest cover/type

The majority of publications, 46 (31%) of studies cover the topic 'Forest cover/type'. The forest cover/type comprises the generation of a forest/non-forest mask (Dlamini 2017, Heckel *et al* 2020), forest cover change




Figure 10. Research topic categories of reviewed articles between 1997 and 2020. Note that some studies cover different topics, which may result in multiple entries.




Figure 11. Number of studies based upon platform and sensor type. Note that studies investigating forest change with multiple platforms were counted multiple times.

estimation (Erkkilä and Löfman 1999, Ringrose et al 2002), forest type discrimination between dryland forests (McCarthy et al 2005), forest health assessment (Herrero et al 2020), woody cover (Boggs 2010, Ibrahim et al 2018), and tree species classification (Hüttich et al 2009, Adelabu et al 2013). The majority of forest type/cover mapping was undertaken with optical multi-spectral data including Landsat, MODIS, and AVHRR and a few studies used high-resolution data such as RapidEye, GeoEye, and WorldView. On the other hand, a few studies on forest cover/type mapping used a combination of multispectral and spaceborne SAR data (X-band, C-band, and L-band) such as Landsat and JERS-1 (Bucini et al 2009), Landsat and ALOS PALSAR (Higginbottom et al 2018, Naidoo et al 2016) and Sentinel-1 and -2 (Heckel et al 2020) (figure 11).

A few studies on forest cover/type mapping relied on field data (Bucini et al 2009, Ibrahim et al 2018, Schultz et al 2018) or forest inventory plots (Heckel et al 2020). Most studies did not include detailed field measurements (species composition, density, frequency, dominance, and basal area, percentage soil cover, total height) and had very few field samples (Gessner et al 2013). Other studies relied on high resolution EO data (Dlamini 2017,




Figure 12. Research topic by country. Note that the order of the mentioned topics has changed when compared to figure 10 as some studies were conducted in several countries.

Higginbottom *et al* 2018), and published maps (Westinga *et al* 2020) as reference data to validate their results. The majority of studies did not perform any form of accuracy assessment or validation of quantitative estimates (e.g., Campo-Bescós *et al* 2013, Harris *et al* 2014). Forest cover and species mapping is essential for many forestry-related tasks and play a key role in sustainable forest management; the importance of these topics can be seen in the fact that they are addressed across all countries in Southern Africa, with the majority of studies conducted in RSA, followed by Namibia and Botswana (figure 12).

4.3.3. Forest biomass and structures

Fifteen research papers (10%) studied forest biomass, and fourteen publications (10%) assessed 'forest structure'. Studies on biomass included the estimation of AGB (Mutanga and Rugege 2006, Dube *et al* 2018), and changes in carbon stock (Gara *et al* 2017). Some of the publications used National Forest Inventory (NFI) data (Verbesselt *et al* 2007, Halperin *et al* 2016), and field-based samples (Tsalyuk *et al* 2017, Mareya *et al* 2018) to estimate biomass in Southern Africa.

Forest structure in the review includes research on stand structure (Mathieu *et al* 2013), canopy cover (Erkkilä and Löfman 1999, Huemmrich *et al* 2005), canopy gaps (Cho *et al* 2015), and stand density (Adjorlolo and Mutanga 2013). The majority of studies on 'forest structure' in Southern Africa dealt with canopy cover (e.g., Yang and Prince 2000, Adjorlolo *et al* 2014). Very few studies considered vertical forest structure including tree height and tree crown diameter (e.g., Verlinden and Laamanen 2006b). Mareya *et al* (2018) utilised freely available high resolution Google satellite imagery in combination with object-based image analysis (OBIA) to estimate tree crown areas in miombo forests and found the overall accuracy to be low and unsuitable when high accuracy is required. Some of the 'forest structure' publications are also assigned to the research topic 'biomass', which discusses the relevance of forest structure for biomass (Meyer *et al* 2014). Forest structure is also a very important parameter when it comes to habitat suitability, species diversity, biodiversity estimation, and conservation studies and thus some publications cover both topics (e.g., Akinyemi and Kgomo 2019).

The methods applied in the biomass and forest structure publications are diverse. Most studies employed some sort of regression analysis between *in situ* field data and EO data, with the most popular methods being random forests, support vector machines, kriging, linear and generalised linear models (Mutanga and Rugege 2006, Carreiras *et al* 2013, Halperin *et al* 2016, Wingate *et al* 2018, Berger *et al* 2019). Williams *et al* (2013) utilised the simple ensemble model to analyse biomass dynamics and found that biomass distributions can diagnose disturbance processes in miombo woodlands. Most studies utilised NDVI index in dryland forest mapping to correlate with biomass (Wessels *et al* 2006, Gizachew *et al* 2016), but very few studies considered other vegetation indices such as red-edge (RE)-computed indices (e.g., Gara *et al* 2016, Dube *et al* 2018). For the most part, optical sensors were used to derive forest biomass and structures, only four papers utilised radar data, and one paper used a combination of radar and optical data to estimate biomass (Wingate *et al* 2018). More

research is needed to explore the improvement of forest AGB and forest structure estimation through multi-sensor (optical and radar) data fusion.

4.3.4. Climate change and disturbances

Here we refer to dryland forests stress monitoring (e.g., damage due to fire, climate/weather-related hazards including drought events, floods, extreme temperatures as part of climate change and disturbances). Twenty-one papers (13%) investigated disturbances to forest cover. Among the different forms of disturbance, fire damage was the most commonly studied (Mayr *et al* 2018, Pricope and Binford 2012, Roy *et al* 2019, Silva *et al* 2003). In the context of threats of climate change, other disturbances included drought (Urban *et al* 2018, Lawal *et al* 2019, Marumbwa *et al* 2021) and floods (Pricope *et al* 2015). A regional studies Lawal *et al* (2019) used gridded climate data from the Climate Research Unit and GMMS NDVI to characterise the impact of drought to vegetation in southern Africa from 1981 to 2005; They found that the responses of vegetation varied according to season and biome, and showed that droughts had extensive impacts over the central parts of South Africa and Namibia, and the southern border of Botswana and the western parts of Zambia. In this review, we only considered studies that investigated climate change in terms of temperature/drought in dryland forests where satellite data are a primary or secondary source of data. Although there are a number of studies on climate change modelling in Southern Africa, the results show that there is a striking lack of studies investigating climate change into dryland forest change and stress monitoring.

The sensors used to detect disturbances differs, with most studies using MODIS (Alleaume *et al* 2005, Chongo *et al* 2007, Archibald *et al* 2009, Giglio *et al* 2009), two publications used SPOT-VGT (Silva *et al* 2003, Verbesselt *et al* 2006), and one Landsat and Sentinel-2 (Roy *et al* 2019). Only two publications utilised SAR data. Mathieu *et al* (2019) investigated SAR Sentinel-1A C-band images for detecting surface fires in the Kruger NP, while Williams *et al* (2013) used ALOS PALSAR to analyse known disturbance agents in tropical woodlands in Mozambique. The research by Urban *et al* (2018) used Sentinel-1 SAR time series NDVI from Sentinel-2 and Landsat-8 to derive surface moisture for drought monitoring in the Kruger NP between 2015 and 2017. A combination/fusion of SAR and Optical data for detecting disturbances is not tested by any study. Only one study used field data as input data for validation (Alleaume *et al* 2005), while two studies used forest inventory data (Verbesselt *et al* 2006, Verlinden and Laamanen 2006a).

4.3.5. Biodiversity, plant traits, and phenology

Twelve (8%) of the reviewed publications dealt with research questions in the context of forest biodiversity. Almost half of the papers on forest biodiversity examined plant species diversity (Adjorlolo *et al* 2014, Mapfumo *et al* 2016, Chapungu *et al* 2020). Others looked at animal species and habitat suitability (e.g., Cáceres *et al* (2015) for birds, Ducheyne *et al* (2009) for tsetse flies, impala (Van Bommel *et al* 2006), and elephants (Marston *et al* 2020). Forest biodiversity is often related to structural canopy parameters. Most studies, nine (75%) of twelve used Landsat to derive parameters such as plant canopy height, species occurrence, richness, and diversity. Three (25%) of the studies used MODIS data (e.g., Fullman and Bunting (2014) used MODIS at 250 m pixel resolution and a Moving Standard Deviation Index (MSDI) to detect elephant-modified vegetation along the Chobe riverfront in Botswana; Akinyemi and Kgomo (2019) utilised 1 km spatial resolution of SPOT - VGT and PROBA-V annual time series of 18 years to understand species diversity and richness assessment based on the Vegetation Degradation Index in Palapye Botswana.; Adjorlolo *et al* (2014) investigated the utility of SPOT-5 multispectral data to assess tree equivalents and total leaf mass to model grazing and browsing capacity in KwaZul-Natal province in RSA.

Five papers (3%) dealt with different plant characteristics, known as plant functional traits. These include canopy chlorophyll content (Cho *et al* 2012), leaf nitrogen concentration (Cho *et al* 2013), and vegetation water content (Verbesselt *et al* 2006), and LAI (Scholes *et al* 2004). Plant functional traits including vegetation biophysical and biochemical properties (e.g., pigment levels, nitrogen content) are often related to patterns of biodiversity. Huemmrich *et al* (2005) explored monthly MODIS data at 1 km spatial resolution over two years to estimate LAI and FAPAR and found that ground-measured LAI values correspond well with MODIS LAI, and showed a discrepancy with FAPAR. Cho *et al* (2012) utilised variogram analysis and the red edge shift from SumbandilaSat and SPOT 5 to estimate canopy chlorophyll content in Dukuduku forest in Southern Africa and found that SumbandilaSat provides additional information for quantifying stress in vegetation as compared to SPOT image data. All studies on plant traits were undertaken at the local scale.

Looking at research categories per country, biodiversity/habitat publications were mainly undertaken in Botswana and RSA (figure 12). All studies in the context of forest biodiversity and plant traits covered only mono-temporal and multi-annual classifications. Only two studies utilised multi-annual time series (Verbesselt *et al* 2006, Akinyemi and Kgomo 2019), and one study used MODIS multi-temporal time series over two years (Huemmrich *et al* 2005). All of these studies focused on a coarse resolution of 1 km.

Phenology is also strongly linked to plant traits, but analysis puts more emphasis on the seasonal variations including growing season (green-up date) (Archibald and Scholes 2007, Whitecross *et al* 2017), end of the season, and length of the season (Davis *et al* 2017). To date, phenological research in Southern African dryland forests is limited, and more than half of the published papers on phenology focused only on examples from RSA. In the few studies that have analysed phenology, most studies dealt with estimating leaf flush and early-greening dates (Chidumayo 2001, Higgins *et al* 2011). For example, Archibald and Scholes (2007) developed an intricate algorithm that used MODIS NDVI products and field-based parameter estimates to predict green-up dates for grass and tree components at a site in the Kruger NP in RSA. Jolly and Running (2004) compared a water balance model to a 3-year NDVI time series and found the deviation between the onset of leaf flush predicted by the model and empirical data was between 10 and 40 d.

5. Discussion

5.1. Temporal extent

In this article, we have synthesized the current research with EO on dryland forests, with a particular focus on Southern Africa. Although the volume of scientific literature has demonstrated a sharp increase, the use of remote sensing is still limited, and up until 2013, the number of publications on this topic was relatively small. Substantial research on the dryland forests of Southern Africa is mainly based on single-date observations, and comparing classified images at two or more different times. Maps that relate successive land cover change between two dates typically lack information regarding underlying processes and do not enable insights on the nature of the transformations present, such as the rate or persistence of change (Lambin *et al* 2003). Time series analysis on dryland forests, which enables tracking changes is scarce, only 22 (16%) out of 137 studies feature time series lengths that exceed 15 years and only 11 (8%) studies that cover more than 20 years. Longer time series of remote sensing data afford the ability to assess the dynamics of forest structures, biodiversity, degradation, disturbance from climatic extremes, and change in phenology, in which a gap still exists.

5.2. Spatial scale

Another finding that stands out from our analyses is that there are very few studies at the national and regional levels. Despite new sensor and EO data availability, it is clear that a systematic and consistent regional monitoring of dryland forests is not yet fully exploited and is still in its infancy in Southern Africa. In fact, the majority of publications 88 (64%) concentrated their research efforts on local scale investigations (figure 7). Desanker and Justice (2001) and Geist (2002) also emphasised that Southern Africa is limited to local-scale studies, thereby lacking a simultaneous analysis of the impacts of these changes at a larger scale. To fully assess regional and long-term implications for tropical dryland forest change studies, analyses on large(r) scales are needed, ideally with higher spatial resolutions and longer temporal duration.

5.3. Accuracy assessment

Through evaluation of the literature, we identified that the assessment of accuracy for thematic/classified maps and statistical data to be another important issue, with only 54 (39%) of the studies appearing to have performed some form of accuracy assessment. Our results show there is limited information on sources of error and uncertainty levels of the estimates provided by most studies. We found that most forest and vegetation-related scientific outputs in Southern Africa are not yet strongly linked to field measurements and forest inventory data. Among the reviewed studies, very few studies utilized field test sites/ ground-based independent datasets for accuracy assessment, while other studies estimated uncertainties using other procedures e.g., using a sample of finer spatial resolution remote sensing data, or did not report the map uncertainty. Some studies employed root-mean-square error to assess model accuracy (RMSE) (e.g., Adjorlolo and Mutanga 2013, Higginbottom *et al* 2018), while many studies used an error matrix to assess map uncertainties, which was employed for instance (e.g., Hüttich *et al* 2011, Adelabu *et al* 2013). However, some studies used sample points below the desirable target number of validation points per class (e.g., Cabral *et al* 2011), while studies briefly mentioned that a confusion matrix was calculated but did not report how many sample points were used for validation (e.g., Chagumaira *et al* 2016). Congalton. (1988) suggests planning to collect a minimum of 50 samples for each map class for maps of less than 1 million acres in size with less than 12 classes. It has been empirically confirmed that a good balance between statistical validity and practicality for larger area maps or more complex maps can be achieved with about 75 to 100 sample sites per class (Congalton and Green 2009).

Globally, owing to TDFs low commercial importance in comparison to other tropical forests such as moist forest, they are often not assessed by field surveys, or surveyed regularly by governments (Keenan 2015). Independent validation data for dryland forest estimations are rarely available because acquiring appropriate field survey data is a time-consuming and expensive task. In Southern Africa, these areas are often remote and

dangerous to visit in the field, due to the hazard posed by wildlife and if present, unexploded landmines, almost impracticable to obtain independent validation data for large(r) area studies, especially for many protected areas. Despite challenges to obtain ground-based observation, effective integration of these data and remote sensing methods will be key to accurately mapping and monitoring dryland forest across a range of spatial scales and in reporting the accuracy of models. However, the applicability of remotely measured geospatial data is reliant on quality, and translating remote sensing data into accurate and meaningful information is often a challenge prone to errors (Congalton 2009, Donoghue 2002). In this context, it is critical to ensure the validity of these data and their suitability for each particular application, particularly where coarse spatial maps can be misleading. In addition, characterising dryland forest for large areas of Africa cannot entirely rely on global and pantropical monitoring studies for dry forest estimation because global forest monitoring generally underestimates, and in some instances overestimates, dryland biomes (Bastin *et al* 2017).

5.4. Research topics and geographical focus

The classification of studies into eight broad subject categories revealed forest cover/types 41 (26%) and land cover/land use 36 (23%) to be the most commonly researched topics. Topics receiving less attention included phenology, plant traits, biodiversity/habitats, and disturbances with regards to climate change (figure 10). With regards to disturbances, fire damage was the most commonly studied but there is a missing body of literature on the climate change impact on the composition, biodiversity, and ecological health of dry forest ecosystems in most countries of Southern Africa. We also found an interesting, non-uniform spatial distribution of dryland vegetation and forest studies using spaceborne remote sensing, particularly when considering disparities among countries and across protected areas. The distribution of research categories by country reveals that RSA is, by far the most studied nation across all categories in Southern Africa (figure 8). It should be noted that care should be taken here not to assume that the number of studies equates to research quality, which remains difficult to articulate from a review of this nature. However, the dryland forests of Mozambique, Lesotho, Swaziland, and Zambia are noticeably very poorly studied. Studies on the dryland forests of Angola are even less frequent, receiving relatively little global attention, and the few studies conducted on its forests were mostly conducted by researchers from Portuguese Universities (Leite *et al* 2018, Catarino *et al* 2020). The focus of publications tended to be biased towards conservation and national parks, particularly as a large proportion of studies were undertaken in the Kruger NP, leaving many other private and international protected areas relatively understudied. Transboundary conservation areas, such as Kavango-Zambezi (KAZA), have received relatively little attention but merit further research in terms of the vast dryland forests extent, biodiversity, species abundance and diversity, and the potential for this area to form important corridor areas for wildlife animals. There is a further concern as a result of such gaps because some of the dryland forests, and species to which they are home, notably in countries like Angola and Zambia, are listed on the IUCN red list and would almost certainly merit Alliance for Zero Extinction (AZE) ranking (Cumming 2008). Furthermore, future efforts to estimate important variables such as forest cover and biomass need not be restricted by country boundaries. Future studies, based on medium-fine resolution EO and validated with field data, will provide information to improve our understanding of African dryland vegetation and its management.

5.5. Vegetation indices, optical, SAR, and fusion of optical and SAR sensors

The most commonly used vegetation index was the NDVI, with more than half of the studies, 84 (54%) of papers utilising this index, but only 13 (8%) of papers used EVI index and SAVI index. Other vegetation indices such as the GNDVI index and Sentinel red-edge related indices and passive microwave observations such as Vegetation Optical Depth were not utilised in studies considered in this review. One major problem commonly encountered in the less studied ecosystems, such as dryland forests, is that of generalizing or transferring knowledge and methods derived from remotely sensed imagery over both space and time (Foody *et al* 2003). For example, commonly used vegetation indices and classification schemes are in general mainly been calibrated on other, better-studied ecosystems, such as temperate or rain forests, and this has led to poor accuracy results when extrapolated, to for example, tropical dryland forests. This phenomenon justifies the importance of utilizing a range of vegetation indices when studying dryland forests using EO data. Imagery from optical sensors is most commonly used, out of all sensor types, providing the data used in 90% of papers reviewed, followed by SAR data with 6%. The fusion of optical and radar data was rarely used, with only 4% of publications exploring this. The most frequently used platforms are Landsat, followed by MODIS and AVHRR. Imagery taken by the Sentinel-1/2 satellites only makes up a small portion of the remote sensing data on dryland forests. For example, Sentinel-2 was only used by 2% of investigated studies, but this may reflect the relatively short period (since 2015) when these data have been available.

5.6. Remote sensing platforms and cloud-based computing

Most of the EO data used in the publications reviewed were downloaded, and are available at no cost from a number of online portals, including the Oak Ridge National Laboratory (ORNL), the United States Geological Survey (USGS) Distributed Active Archive System (DAAC) and Earth Explorer (EE) tool. The lack of remote sensing research centers in most Southern African research institutions may contribute to limit the number of African Scientists engaged in monitoring forests resources. For example, most studies in RSA made use of remote sensing data through the University of the Witwatersrand, Satellite Application Centre (SAC), the South African National Space Agency (SANSa), and the Council of Science and Industrial Research (CSIR). The development of remote sensing capacity at local universities has inevitably contributed to RSA universities and research institutions conducting the majority of studies in Southern Africa (figure 6). To improve EO data access, and the skills to handle and interpret this across Southern Africa, there is a need to increase the number of local institutions that distribute the remote sensing data, and who have the capacity to access and use innovative web-based platforms such as the Google Earth Engine (GEE) and Amazon Web Services to overcome some of the logistical and financial constraints of this type of research.

Southern African countries face considerable technical challenges with remote sensing, particularly in respect to REDD+-related research on dryland forests monitoring. Freely available tools, for example, the cloud-based geospatial analysis platform Google Earth Engine (GEE), make it easier to access powerful computing resources for processing and analysing pre-processed large-scale datasets (Shelestov *et al* 2017). However, only nine papers (6%) out of 137 used GEE to access or analyse remote sensing data. The 'near real-time' remote sensing data offered by GEE is of particular interest for monitoring changes and automating the analysis of time-series, when detecting and tracking trends in surface reflectance properties. With increasing spatio-temporal coverage of satellite data and computational platforms that reduce the need for costly local infrastructure (e.g., GEE), there is an opportunity to overcome the limitations previously enforced by large volumes of data and the scale of analysis, whereby our knowledge of dryland forest dynamics can be improved in the upcoming years.

6. Conclusion

This review summarizes research progress towards the use and integration of remote sensing data within the context of monitoring dryland forests in Southern Africa, using a systematic review methodology that focused on 137 most relevant research articles. We have reviewed the temporal and spatial coverage of these studies, their application area, and the remote sensing platforms and sensors used. Based on the results, the following conclusions can be drawn. There are a broad range of topics covered by research on dryland forests, from which land-use/land-cover and forest cover and disturbances from the fire were the most frequently studied. However, there is still a relative lack of studies assessing dryland forest structure, phenology, biodiversity/habitats, plant traits, and disturbance from climatic extremes, suggesting additional research is required. The majority of studies relied on single-date or annual data and bi-temporal discrete classification; only a very few studies employed time series analysis.

We consider some of the limitations of the research reviewed, which indicates a need for more frequent use of field and inventory data, a greater use of validation/accuracy assessments, and testing other vegetation indices beyond NDVI and EVI such as the Vegetation Optical Depth and Sentinel-2 red-edge related indices. In addition, further improvements should focus on for extensive combination and fusion of SAR and optical data in order to have a temporally and spatially consistent data set necessary for several applications in dryland forests. Given the state of decline of woody vegetation condition in Southern Africa, long-term monitoring of monthly time series of EO data at regional and transboundary scale clearly hold potential to capture dryland forests dynamics and to understand their current status and future trends. A significant move from EO predictions that are extremely site-dependent to large(r) ecoregional level monitoring approach that integrates a range of remotely-sensed data of sufficiently high spatial and temporal resolution with field measurements and using machine/deep learning models could provide a sound basis for assessing dryland forest-related changes and dynamics. Information inferred from these kinds of models would be extremely useful for the current knowledge, management and conservation of the dryland forests as well as for understanding their responses to disturbance (natural or anthropogenic) and climatic change at regional to sub-continental level. Finally, there is significant geographical heterogeneity in study coverage; whilst there is substantial research on the forests in the Kruger NP and across RSA, the same cannot be said for other areas of Southern Africa. The EO interventions not only assess deforestation rate, but also support other forest related REDD + activities such as sustainable forest management which reduce forest degradation and enhance forest carbon stocks at a range of scales, transcending both provincial and national boundaries e.g., Kavango-Zambezi Transfrontier Conservation Area (KAZA TFCA). Nevertheless, REDD+-related research on dryland forests in most Southern African countries

and protected areas has been limited, with clear gaps across Angola, Mozambique, Zambia, and Zimbabwe. Finally, Africa has the potential to emulate other continents, such as Latin America, that have made notable progress in employing freely available remote sensing data to monitor tropical dryland forest area change and biomass on a large scale.

Acknowledgments

Many thanks to Mark Reed for his comments on an earlier draft. We are grateful to the anonymous reviewers whose insightful comments improved the manuscript substantially.

Data availability statement

The data that support the findings of this study are available upon reasonable request from the authors.

Conflicts of interest

The authors declare no conflict of interest.

Funding

This work was supported Commonwealth Scholarship Commission Ph.D grant number: NACS-2017–409 from 2017–2020, Geography doctoral program at Durham University. The authors would like to acknowledge the support provided through the Rapid Evidence Synthesis Training (REST) programme. REST was organised and delivered through a collaboration between the University of Leeds, The University of Newcastle, and the N8 AgriFood Programme and supported by Research England QR-SPF funds from The University of Leeds and University of York.

ORCID iDs

Ruusa M David  <https://orcid.org/0000-0002-7892-9878>

Nick J Rosser  <https://orcid.org/0000-0002-1435-2512>

Daniel N M Donoghue  <https://orcid.org/0000-0002-9931-9083>

References

- Adelabu S, Mutanga O, Adam E and Cho M A 2013 Exploiting machine learning algorithms for tree species classification in a semi-arid woodland using RapidEye image *J. Appl. Remote Sens.* **7** 073480
- Adjorlolo C, Botha J O, Mhangara P, Mutanga O and Odindi J 2014 Integrating remote sensing and conventional grazing/browsing models for modelling carrying capacity in southern African rangelands *Remote Sensing for Agriculture, Ecosystems, and Hydrology XVI* 9239
- Adjorlolo C and Mutanga O 2013 Integrating remote sensing and geostatistics to estimate woody vegetation in an African savanna *Journal of Spatial Science* **58** 305–22
- Agbaje G I 2010 *Nigeria in Space—an Impetus For Rapid Mapping of the Country for Sustainable Development Planning. A Report Prepared for FIG Congress 2010 Facing the Challenges Building the Capacity Sydney (Australia)* (https://fig.net/resources/proceedings/fig_proceedings/fig2010/papers/ts03h/ts03h_agbaje_4621.pdf) (accessed 20 January 2021)
- Aguilar M, Saldaña M and Aguilar F 2013 GeoEye-1 and WorldView-2 pan-sharpened imagery for object-based classification in urban environments *Int. J. Remote Sens.* **34** 2583–606
- Aide T M, Clark M L, Grau H R, López-Carr D, Levy M A, Redo D, Bonilla-Moheno M, Riner G, Andrade-Núñez M J and Muñiz M 2012 Deforestation and reforestation of Latin America and the Caribbean (2001–2010) *Biotropica* **45** 262–71
- Akinyemi F O and Kgomo M O 2019 Vegetation dynamics in African drylands: an assessment based on the vegetation degradation index in an agro-pastoral region of Botswana *Regional Environmental Change* **19** 2027–39
- Alcañiz I and Gutierrez R A 2020 Between the global commodity boom and subnational state capacities: payment for environmental services to fight deforestation in Argentina *Global Environmental Politics* **20** 38–59
- Aleman J C, Jarzyna M A and Staver A C 2018 Forest extent and deforestation in tropical Africa since 1900 *Nature Ecology & Evolution* **2** 26–33
- Ali I, Greifeneder F, Stamenkovic J, Neumann M and Notarnicola C 2015 Review of machine learning approaches for biomass and soil moisture retrievals from remote sensing data *Remote Sensing* **7** 16398–421
- Alleaume S, Hély C, Le Roux J, Korontzi S, Swap R J, Shugart H H and Justice C O 2005 Using MODIS to evaluate heterogeneity of biomass burning in southern African savannahs: a case study in Etosha *Int. J. Remote Sens.* **26** 4219–37
- Alvarez-Añorve M Y, Quesada M, Sánchez-Azofeifa G A, Avila-Cabadilla L D and Gamon J A 2012 Functional regeneration and spectral reflectance of trees during succession in a highly diverse tropical dry forest ecosystem *American Journal of Botany* **99** 816–26

- Andela N, Liu Y Y, Van Dijk A I J M, De Jeu R A M and McVicar T R 2013 Global changes in dryland vegetation dynamics (1988–2008) assessed by satellite remote sensing: comparing a new passive microwave vegetation density record with reflective greenness data *Biogeosciences* **10** 6657–76
- Archibald S, Roy D P, van Wilgen B W and Scholes R J 2009 What limits fire? An examination of drivers of burnt area in Southern Africa *Global Change Biol.* **15** 613–30
- Archibald S and Scholes R J 2007 Leaf green-up in a semi-arid African savanna - separating tree and grass responses to environmental cues *Journal of Vegetation Science* **18** 583–94
- Arino O et al 2007 Globcover—a global land cover service with MERIS *In Proc. of the ENVISAT Symp.* 23–7
- Balint P J and Mashinya J 2006 The decline of a model community-based conservation project: governance, capacity, and devolution in Mahenye, Zimbabwe *Geoforum* **37** 805–15
- Barbosa J, Broadbent E and Bitencourt M 2014 Remote sensing of aboveground biomass in tropical secondary forests: a review *International Journal of Forestry Research* **2014**
- Bastin J F et al 2017 The extent of forest in dryland biomes *Science* **356** 635–8
- Becker F and Choudhury B J 1988 Relative sensitivity of normalized difference vegetation index (NDVI) and microwave polarization difference index (MPDI) for vegetation and desertification monitoring *Remote Sens. Environ.* **24** 297–311
- Berger C, Werner S, Wigley-Coetsee C, Smit I and Schullius C 2019 Multi-temporal sentinel-1 data for wall-to-wall herbaceous biomass mapping in Kruger National Park, South Africa—first results *IGARSS 2019-2019 IEEE Int. Geoscience and Remote Sensing Symp. IEEE* **7358–60**
- Blackie R et al 2014 Tropical dry forests *The State of Global Knowledge and Recommendations for Future Research (Discussion Paper)* (Bogor Indonesia: Cifor)
- Bodart C, Brink A B, Donnay F, Lupi A, Mayaux P and Achard F 2013 Continental estimates of forest cover and forest cover changes in the dry ecosystems of Africa between 1990 and 2000 *Journal of Biogeography* **40** 1036–47
- Boggs G S 2010 Assessment of SPOT 5 and QuickBird remotely sensed imagery for mapping tree cover in savannas *Int. J. Appl. Earth Obs. Geoinf.* **12** 217–24
- Bond Ivan, Chambwerwa M, Jones B, Chundama M and Nhantumbo I 2010 *REDD+ in dryland forests: issues and prospects for pro-poor REDD in the miombo woodlands of southern Africa* (21) (London: IIED)
- Bouvet A, Mermoz S, Le Toan T, Villard L, Mathieu R, Naidoo L and Asner G P 2018 An above-ground biomass map of African savannas and woodlands at 25 m resolution derived from ALOS PALSAR *Remote Sens. Environ.* **206** 156–73
- Brandt M et al 2018a Reduction of tree cover in West African woodlands and promotion in semi-arid farmlands *Nat. Geosci.* **11** 328–33
- Brandt M et al 2018b Satellite passive microwaves reveal recent climate-induced carbon losses in African drylands *Nature ecology & evolution* **2** 827–35
- Brandt M, Hiernaux P, Rasmussen K, Mbow C, Kergoat L, Tagesson T, Ibrahim Y Z, Wélé A, Tucker C J and Fensholt R 2016 Assessing woody vegetation trends in Sahelian drylands using MODIS based seasonal metrics *Remote Sens. Environ.* **183** 215–25
- Breiman L 2001 Random forests *Mach. Learn.* **45** 5–32
- Bucini G, Saatchi S, Hanan N, Boone R B and Smit I 2009 Woody cover and heterogeneity in the savannas of the Kruger National Park, South Africa 2009 *IEEE International Geoscience and Remote Sensing Symposium. IEEE* **4** 1V–334
- Bullock S H, Mooney H A and Medina E 1995 *Seasonally Dry Tropical Forests* (Great Britain: Cambridge University Press)
- Bunting E L, Southworth J, Herrero H, Ryan S J and Waylen P 2018 Understanding long-term savanna vegetation persistence across three drainage basins in Southern Africa *Remote Sensing* **10** 1013
- Bunyavejchewin S, Baker P J and Davies S J 2011 Seasonally dry tropical forests in continental Southeast Asia: structure, composition and dynamics *The Ecology and Conservation of Seasonally Dry Forests in Asia*. (United State: Smithsonian Institution Scholarly Press) 9–35
- Cabral A I R, Vasconcelos M J, Oom D and Sardinha R 2011 Spatial dynamics and quantification of deforestation in the central-plateau woodlands of Angola (1990–2009) *Appl. Geogr.* **31** 1185–93
- Cáceres A, Melo M, Barlow J, Cardoso P, Maiato F and Mills M S L 2015 Threatened birds of the angolan central escarpment: distribution and response to habitat change at Kumbira Forest *Oryx* **49** 727–34
- Campo-Bescós M A, Muñoz-Carpena R, Southworth J, Zhu L, Waylen P R and Bunting E 2013 Combined spatial and temporal effects of environmental controls on long-term monthly NDVI in the Southern Africa Savanna *Remote Sensing* **5** 6513–38
- Campos-Vargas C and Vargas-Sanabria D 2021 Assessing the probability of wildfire occurrences in a neotropical dry forest *Écoscience* **28** 159–69
- Carreiras J, Melo J B and Vasconcelos M J 2013 Estimating the above-ground biomass in miombo savanna woodlands (Mozambique, East Africa) using L-band synthetic aperture radar data *Remote Sensing* **5** 1524–48
- Castillo A, Magaña A, Pujadas A, Martínez L and Godínez C 2005 Understanding the interaction of rural people with ecosystems: a case study in a tropical dry forest of Mexico *Ecosystems* **8** 630–43
- Castro K, Sanchez-Azofeifa G A and Rivard B 2003 Monitoring secondary tropical forests using space-borne data: implications for Central America *Int. J. Remote Sens.* **24** 1853–94
- Castro S M, Sanchez-Azofeifa G A and Sato H 2018 Effect of drought on productivity in a Costa Rican tropical dry forest *Environ. Res. Lett.* **13** 045001
- Catarino S, Romeiras M M, Figueira R, Aubard V, Silva J and Pereira J 2020 Spatial and temporal trends of burnt area in Angola: implications for natural vegetation and protected area management *Diversity* **12** 307
- Chagumaira C, Rurinda J, Nezomba H, Mtambanengwe F and Mapfumo P 2016 Use patterns of natural resources supporting livelihoods of smallholder communities and implications for climate change adaptation in Zimbabwe *Environment, development and sustainability* **18** 237–55
- Chambers J Q, Asner G P, Morton D C, Anderson L O, Saatchi S S, Espírito-Santo F D, Palace M and Souza C Jr 2007 Regional ecosystem structure and function: ecological insights from remote sensing of tropical forests *Trends Ecol. Evol.* **22** 414–23
- Chapungu L, Nhamo L, Gatti R C and Chitakira M 2020 Quantifying changes in plant species diversity in a savanna ecosystem through observed and remotely sensed data *Sustainability* **12** 2345
- Charles-Dominique T, Staver A C, Midgley G F and Bond W J 2015 Functional differentiation of biomes in an African savanna/forest mosaic *S. Afr. J. Bot.* **101** 2–90
- Chazdon R L et al 2016 Carbon sequestration potential of second-growth forest regeneration in the Latin American tropics *Sci. Adv.* **2** e1501639
- Chidumayo E 2001 Climate and phenology of savanna vegetation in southern Africa *Journal of Vegetation Science* **12** 347–54
- Chidumayo E. and Marunda C 2010 Dry forests and woodlands in sub-Saharan Africa: Context and challenges. *In The Dry Forests and Woodlands of Africa* (New York: Routledge)

- Chidumayo E N 2013 Forest degradation and recovery in a miombo woodland landscape in Zambia: 22 years of observations on permanent sample plots *Forest Ecology and Management* **291** 154–61
- Chidumayo E N and Gumbo D J 2010 *The Dry Forests and Woodlands of Africa: Managing for Products and Services* (New York: Earthscan)
- Chidumayo E N 2019 Management implications of tree growth patterns in miombo woodlands of Zambia *Forest Ecology and Management* **436** 105–16
- Child B A, Musengezi J, Parent G D and Child G F 2012 The economics and institutional economics of wildlife on private land in Africa *Pastoralism: Research, Policy and Practice* **2** 1–32
- Chiteculo V, Abdollahnejad A, Panagiotidis D, Surový P and Sharma R P 2018 Defining deforestation patterns using satellite images from 2000 and 2017: assessment of forest management in miombo forests—a case study of Huambo Province in Angola *Sustainability* **11** 98
- Cho M A, Debbab P, Mutanga O, Dudeni-Tlhoneb N, Magadla T and Khuluseb S A 2012 Potential utility of the spectral red-edge region of SumbandilaSat imagery for assessing indigenous forest structure and health *Int. J. Appl. Earth Obs. Geoinf.* **16** 85–93
- Cho M A, Malahlela O and Ramoelo A 2015 Assessing the utility WorldView-2 imagery for tree species mapping in South African subtropical humid forest and the conservation implications: Dukuduku forest patch as case study *Int. J. Appl. Earth Obs. Geoinf.* **38** 349–57
- Cho M A, Ramoelo A, Debbab P, Mutanga O, Mathieu R, van Deventer H and Ndlovu N 2013 Assessing the effects of subtropical forest fragmentation on leaf nitrogen distribution using remote sensing data *Landscape Ecology* **28** 1479–91
- Chongo D, Nagasawa R, Ould Cherif Ahmed A and Perveen M F 2007 Fire monitoring in savanna ecosystems using MODIS data: a case study of Kruger National Park, South Africa *Landscape and Ecological Engineering* **3** 79–88
- Coetzer-Hanack K L, Witkowski E T F and Erasmus B F 2016 Thresholds of change in a multi-use conservation landscape of South Africa: historical land-cover, future transformation and consequences for environmental decision-making *Environ. Conserv.* **43** 253–62
- Collaboration for Environmental Evidence 2013 Guidelines for systematic review and evidence synthesis in environmental management *Environmental Evidence* **4.2** 1–82 (<https://www.environmentalevidence.org/wp-content/uploads/2018/02/Review-guidelines-version-4.2-final-update-1.pdf>)
- Congalton R G 2009 Accuracy and error analysis of global and local maps: Lessons learned and future considerations. *Remote Sensing of Global Croplands for Food Security* ed Prasad. Thenkabail, G. John and Hugh Lyon (Florida, USA: CRC Press) pp 441–58
- Congalton R G 1988 A comparison of sampling schemes used in generating error matrices for assessing the accuracy of maps generated from remotely sensed data *Photogramm. Eng. Remote Sens.* **54** 593–600
- Congalton R G 2018 Landsat's enduring legacy: pioneering global land observations from space *Photogrammetric Engineering & Remote Sensing* **84** 9–10
- Congalton R G and Green K 2009 *Assessing the Accuracy of Remotely Sensed Data: Principles and Practices* 3rd (Florida, USA: CRC press) (<https://doi.org/10.1201/9780429052729>)
- Corbera E, Soberanis C G and Brown K 2009 Institutional dimensions of Payments for Ecosystem Services: An analysis of Mexico's carbon forestry programme *Ecological Economics* **68** 743–61
- Cui Q, Shi J, Du J, Zhao T and Xiong C 2015 An approach for monitoring global vegetation based on multiangular observations from SMOS *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **8** 604–16
- Cumming D H M 2008 Large scale conservation planning and priorities for the Kavango-Zambezi transfrontier conservation area *A report prepared for Conservation International.* (http://wcs-ahead.org/kaza/kaza_tfca_large_scale_planning_final_7nov08_logo.pdf/) (accessed 23 August 2019)
- Curran P 1980 Multispectral remote sensing of vegetation amount *Prog. Phys. Geog.* **4** 315–41
- Curry T R 2020 Mexico's tropical dry forests *Reference Module in Earth Systems and Environmental Science.* ed A L Shober and R O Maguire (Amsterdam, Netherlands: Elsevier) (<https://doi.org/10.1016/B978-0-12-821139-7.00007-6>)
- Daskin J H, Stalmans M and Pringle R M 2016 Ecological legacies of civil war: 35-year increase in savanna tree cover following wholesale large-mammal declines *Journal of Ecology* **104** 79–89
- Davis C L, Hoffman M T and Roberts W 2017 Long-term trends in vegetation phenology and productivity over Namaqualand using the GIMMS AVHRR NDVI3g data from 1982 to 2011 *S. Afr. J. Bot.* **111** 76–85
- de Araujo Barbosa C C, Atkinson P M and Dearing J A 2015 Remote sensing of ecosystem services: a systematic review *Ecol. Indic.* **52** 430–43
- Desanker P V and Justice C O 2001 Africa and global climate change: critical issues and suggestions for further research and integrated assessment modeling *Climate Research* **17** 93–103
- Deweese P A 1994 Social and economic aspects of miombo woodland management in southern Africa: options and opportunities for research *CIFOR Occasional Paper* **2** 1–28
- Dexter K G et al 2015 Floristics and biogeography of vegetation in seasonally dry tropical regions *International Forestry Review* **17** 10–32
- Djoudi H, Vergles E, Blackie R, Koame C K and Gautier D 2015 Dry forests, livelihoods and poverty alleviation: understanding current trends *International Forestry Review* **17** 54–69
- Dlamini W M 2017 Mapping forest and woodland loss in Swaziland: 1990–2015 *Remote Sensing Applications: Society and Environment* **5** 45–53
- do Espírito-Santo M M, Rocha A M, Leite M E, Silva J O, da Silva L P and Sanchez-Azofeifa G A 2020 Biophysical and socioeconomic factors associated to deforestation and forest recovery in Brazilian tropical dry forests *Frontiers in Forests and Global Change* **3** 141
- Dogru A O, Goksel C, David R M, Tolunay D, Sözen S and Orhon D 2020 Detrimental environmental impact of large scale land use through deforestation and deterioration of carbon balance in Istanbul Northern Forest Area *Environmental Earth Sciences* **79** 1–13
- Donoghue D N 2000 Remote sensing: sensors and applications *Prog. Phys. Geog.* **24** 407–14
- Donoghue D N 2002 Remote sensing: environmental change *Prog. Phys. Geog.* **26** 144–51
- Dube T, Gara T W, Mutanga O, Sibanda M, Shoko C, Murwira A, Masocha M, Ndaimani H and Hatendi C M 2018 Estimating forest standing biomass in savanna woodlands as an indicator of forest productivity using the new generation WorldView-2 sensor *Geocarto Int.* **33** 178–88
- Ducheyne E, Mweempwa C, De Pus C, Vernieuwe H, De Deken R, Hendrickx G and Van den Bossche P 2009 The impact of habitat fragmentation on tsetse abundance on the plateau of eastern Zambia *Preventive Veterinary Medicine* **91** 11–8
- Du Preez Mari-Lise 2014 *Southern Africa's Dryland Forests, Climate Change and the Water-Energy-Food Security Nexus* (<https://www.africanportal.org/publications/southern-africas-dryland-forests-climate-change-and-the-water-energy-food-security-nexus/>)
- Dupuis C, Lejeune P, Michez A and Fayolle A 2020 How can remote sensing help monitor tropical moist forest degradation?—a systematic review *Remote Sensing* **12** 1087
- Durden S L, Van Zyl J J and Zebker H A 1989 Modeling and observation of the radar polarization signature of forested areas *IEEE Trans. Geosci. Remote Sens.* **27** 290–301

- Erkkilä A and Löfman S 1999 Forest cover change in the Ohangwena region, northern Namibia: a case study based on multitemporal Landsat images and aerial photography *Southern African Forestry Journal* **184** 25–32
- Eva H D, Brink A and Simonetti D 2006 Monitoring land cover dynamics in sub-Saharan Africa. Institute for Environmental and Sustainability Office for Official Publication of the European Communities EUR 22498 EN Luxembourg
- Fajardo L, Gonzalez V, Nassar J M, Lacabana P, Portillo Q, C A, Carrasquel F and Rodriguez J P 2005 Tropical dry forests of Venezuela: characterization and current conservation status 1 *Biotropica: The Journal of Biology and Conservation* **37** 531–46
- FAO 1999 Tropical forest management techniques: a review of the sustainability of forest management practices in tropical countries from (<https://fao.org/documents/card/en/c/10c5f896-03f8-536d-b253-aebb36dd2cef/>), accessed 24 March 2021)
- FAO 2001 *FRA 2000: Global Ecological Zoning for the Global Forest Resources Assessment 2000, Final Report* (Rome: Food and Agriculture Organization of the United Nations, Forestry Department) (<http://fao.org/3/ad652e/ad652e00.htm/>) (accessed 19 March 2021)
- Fensham R J and Fairfax R J 2002 Aerial photography for assessing vegetation change: a review of applications and the relevance of findings for Australian vegetation history *Australian Journal of Botany* **50** 415–29
- Ferreira Sanette 2004 Problems associated with tourism development in Southern Africa: The case of Transfrontier Conservation Areas *GeoJournal* **60** 301–10
- Footy G M, Cutler M E, McMorrow J, Pelz D, Tangki H, Boyd D S and Douglas I 2001 Mapping the biomass of Bornean tropical rain forest from remotely sensed data *Global Ecology and Biogeography* **10** 379–87
- Footy G M, Boyd D S and Cutler M E 2003 Predictive relations of tropical forest biomass from Landsat TM data and their transferability between regions *Remote Sens. Environ.* **85** 463–74
- Frankie G W, Mata A and Vinson S B (ed) 2004 *Biodiversity Conservation In Costa Rica: Learning The Lessons In A Seasonal Dry Forest*. (Berkeley and Los Angeles, California: Univ of California Press)
- Fullman T J and Bunting E L 2014 Analyzing vegetation change in an elephant-impacted landscape using the moving standard deviation index *Land* **3** 74–104
- Gara T W, Murwira A, Dube T, Sibanda M, Rwasoka D T, Ndaimani H, Chivhenge E and Hatendi C M 2017 Estimating forest carbon stocks in tropical dry forests of Zimbabwe: exploring the performance of high and medium spatial-resolution multispectral sensors *Southern Forests: a Journal of Forest Science* **79** 31–40
- Gara T W, Murwira A and Ndaimani H 2016 Predicting forest carbon stocks from high resolution satellite data in dry forests of Zimbabwe: exploring the effect of the red-edge band in forest carbon stocks estimation *Geocarto Int.* **31** 176–92
- Gasparri N I and Grau H R 2009 Deforestation and fragmentation of Chaco dry forest in NW Argentina (1972–2007) *Forest Ecology and Management* **258** 913–21
- Geist H J 2002 Causes and pathways of land change in Southern Africa during the past 300 years. Moving from simplifications to generality and complexity *Erdkunde* **56** 144–56
- Gessner U, Machwitz M, Conrad C and Dech S 2013 Estimating the fractional cover of growth forms and bare surface in savannas. A multi-resolution approach based on regression tree ensembles *Remote Sens. Environ.* **129** 90–102
- Giglio L, Loboda T, Roy D P, Quayle B and Justice C O 2009 An active-fire based burned area mapping algorithm for the MODIS sensor *Remote Sens. Environ.* **113** 408–20
- Gillespie T W, Grijalva A and Farris C N 2000 Diversity, composition, and structure of tropical dry forests in Central America *Plant ecology* **147** 37–47
- Gitelson A A, Kaufman Y J and Merzlyak M N 1996 Use of a green channel in remote sensing of global vegetation from EOS-MODIS *Remote Sens. Environ.* **58** 289–98
- Gizachew B, Solberg S, Næsset E, Gobakken T, Bollandsås O M, Breidenbach J, Zahabu E and Mauya E W 2016 Mapping and estimating the total living biomass and carbon in low-biomass woodlands using Landsat 8 CDR data *Carbon Balance Manage.* **11** 1–14
- Göksel Ç, David R M and Dogru A O 2018 Environmental monitoring of spatio-temporal changes in northern Istanbul using remote sensing and GIS *International Journal of Environment and Geoinformatics* **5** 94–103
- González M, R et al 2021 Diverging functional strategies but high sensitivity to an extreme drought in tropical dry forests *Ecology Letters* **24** 451–63
- Guirado Emilio, Blanco-Sacristán Javier, Rodríguez-Caballero Emilio, Tabik Siham, Alcaraz-Segura Domingo, Martínez-Valderrama Jaime and Cabello Javier 2020 Mask R-CNN and OBIA Fusion Improves the Segmentation of Scattered Vegetation in Very High-Resolution Optical Sensors *Sensors* **21** 320
- Halperin J, LeMay V, Chidumayo E, Verchot L and Marshall P 2016 Model-based estimation of above-ground biomass in the miombo ecoregion of Zambia *Forest Ecosystems* **3** 1–17
- Harris A, Carr A S and Dash J 2014 Remote sensing of vegetation cover dynamics and resilience across southern Africa *Int. J. Appl. Earth Obs. Geoinf.* **28** 131–9
- Hastie T, Tibshirani R and Friedman J 2009 (2008) *The Elements of Statistical Learning; Data Mining, Inference and Prediction*. (New York: Springer)
- Heckel K, Urban M, Schratz P, Mahecha M D and Schimmlius C 2020 Predicting forest cover in distinct ecosystems: the potential of multi-source Sentinel-1 and -2 data fusion *Remote Sensing* **12** 302
- Herold M, Mayaux P, Woodcock C E, Baccini A and Schimmlius C 2008 Some challenges in global land cover mapping: an assessment of agreement and accuracy in existing 1 km datasets *Remote Sens. Environ.* **112** 2538–56
- Herrero H, Southworth J, Muir C, Khatami R, Bunting E and Child B 2020 An evaluation of vegetation health in and around Southern African National Parks during the 21st Century (2000–2016) *Applied Sciences* **10** 2366
- Higginbottom T P, Symeonakis E, Meyer H and van der Linden S 2018 Mapping fractional woody cover in semi-arid savannas using multi-seasonal composites from Landsat data *ISPRS J. Photogramm. Remote Sens.* **139** 88–102
- Higgins S I, Delgado-Cartay M D, February E C and Combrink H J 2011 Is there a temporal niche separation in the leaf phenology of savanna trees and grasses? *Journal of Biogeography* **38** 2165–75
- Hoffman M T and Todd S 2000 A national review of land degradation in South Africa: the influence of biophysical and socio-economic factors *Journal of Southern African Studies* **26** 743–58
- Holloway J and Mengersen K 2018 Statistical machine learning methods and remote sensing for sustainable development goals: a review *Remote Sensing* **10** 1365
- Horion S, Fensholt R, Tagesson T and Ehammer A 2014 Using earth observation-based dry season NDVI trends for assessment of changes in tree cover in the Sahel *Int. J. Remote Sens.* **35** 2493–515
- Huang W, Sun G, Ni W, Zhang Z and Dubayah R 2015 Sensitivity of multi-source SAR backscatter to changes in forest aboveground biomass *Remote Sensing* **7** 9587–609

- Huemmerich K F, Privette J L, Mukelabai M, Myneni R B and Knyazikhin Y 2005 Time-series validation of MODIS land biophysical products in a Kalahari woodland, Africa *Int. J. Remote Sens.* **26** 4381–98
- Huete A, Didan K, Miura T, Rodriguez E P, Gao X and Ferreira L G 2002 Overview of the radiometric and biophysical performance of the MODIS vegetation indices *Remote Sens. Environ.* **83** 195–213
- Huete A, Justice C and Van Leeuwen W 1999 MODIS vegetation index (MOD13) *Algorithm Theoretical Basis Document* **3** 295–309
- Hüttich C, Gessner U, Herold M, Strohbach B J, Schmidt M, Keil M and Dech S 2009 On the suitability of MODIS time series metrics to map vegetation types in dry savanna ecosystems: a case study in the Kalahari of NE Namibia *Remote Sensing* **1** 620–43
- Hüttich C, Herold M, Strohbach B J and Dech S 2011 Integrating *in situ*, Landsat, and MODIS data for mapping in Southern African savannas: experiences of LCCS-based land-cover mapping in the Kalahari in Namibia *Environ. Monit. Assess.* **176** 531–47
- Ibrahim S, Balzter H, Tansey K, Tsutsumida N and Mathieu R 2018 Estimating fractional cover of plant functional types in African savannah from harmonic analysis of MODIS time-series data *Int. J. Remote Sens.* **39** 2718–45
- IPCC 2007 *Climate Change 2007: An Assessment of the Intergovernmental Panel on Climate Change*. (Cambridge: Cambridge University Press)
- IPCC 2014 *Climate change 2014—Impacts, adaptation and vulnerability: regional aspects*. Niang, I., O.C. Ruppel, M.A. Abdrabo, A. Essel, C. Lennard, J. Padgham, and P. Urquhart, 2014: Africa *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* ed V R Barros et al (Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press) 1199–265
- Janzen D H 1988 Tropical dry forests *Biodiversity* **15** 130–7
- Jolly W M and Running S W 2004 Effects of precipitation and soil water potential on drought deciduous phenology in the Kalahari *Global Change Biol.* **10** 303–8
- Jones M O, Jones L A, Kimball J S and McDonald K C 2011 Satellite passive microwave remote sensing for monitoring global land surface phenology *Remote Sens. Environ.* **115** 1102–14
- Kamwi Jonathan Mutau, Endjala Junias and Siyambango Nguza 2020 Dependency of rural communities on non-timber forest products in the dry lands of southern Africa: A case of Mukwe Constituency, Kavango East Region, Namibia *Trees, Forests and People* **2** 100022
- Kasischke E S, Melack J M and Dobson M C 1997 The use of imaging radars for ecological applications—A review *Remote Sens. Environ.* **59** 141–56
- Kattenborn T, Leitloff J, Schiefer F and Hinz S 2021 Review on Convolutional Neural Networks (CNN) in vegetation remote sensing *ISPRS J. Photogramm. Remote Sens.* **173** 24–49
- Keenan Rodney J 2015 Climate change impacts and adaptation in forest management: a review *Annals of Forest Science* **72** 145–67
- King E 2014 Southern Africa's dryland forests and climate change adaptation. a policy briefing prepared for governance of Africa's resources programme (https://media.africaportal.org/documents/saia_spb__91_king_20140702.pdf/) (accessed 23 August 2020)
- Klogo G S, Gasonoo A and Ampomah I K 2013 On the performance of filters for reduction of speckle noise in SAR images off the coast of the Gulf of Guinea arXiv:1312.2383
- Kuenzer C, Bluemel A, Gebhardt S, Quoc T V and Dech S 2011 Remote sensing of mangrove ecosystems: a review *Remote Sensing* **3** 878–928
- Kussul N, Lavreniuk M, Skakun S and Shelestov A 2017 Deep learning classification of land cover and crop types using remote sensing data *IEEE Geosci. Remote Sens. Lett.* **14** 778–82
- Lambin E F, Geist H J and Lepers E 2003 Dynamics of land-use and land-cover change in tropical regions *Annual Review of Environment and Resources* **28** 205–41
- Latifi H, Nothdurft A and Koch B 2010 Non-parametric prediction and mapping of standing timber volume and biomass in a temperate forest: application of multiple optical/LiDAR-derived predictors *Forestry* **83** 395–407
- Lawal S, Lennard C, Jack C, Wolski P, Hewitson B and Abiodun B 2019 The observed and model-simulated response of southern African vegetation to drought *Agric. For. Meteorol.* **279** 107698
- Lawley V, Lewis M, Clarke K and Ostendorf B 2016 Site-based and remote sensing methods for monitoring indicators of vegetation condition: an Australian review *Ecol. Indic.* **60** 1273–83
- Le Canut P, Andreae M O, Harris G W, Wienhold F G and Zenker T 1996 Airborne studies of emissions from savanna fires in southern Africa: 1. Aerosol emissions measured with a laser optical particle counter *Journal of Geophysical Research: Atmospheres* **101** 23615–30
- Lehmann E A, Caccetta P, Lowell K, Mitchell A, Zhou Z S, Held A, Milne T and Tapley I 2015 SAR and optical remote sensing: assessment of complementarity and interoperability in the context of a large-scale operational forest monitoring system *Remote Sens. Environ.* **156** 335–48
- Leite A, Cáceres A, Melo M, Mills M S L and Monteiro A T 2018 Reducing emissions from deforestation and forest degradation in Angola: insights from the scarp forest conservation 'hotspot' *Land Degrad. Dev.* **29** 4291–300
- Lesolle D 2012 SADC policy paper on climate change: assessing the policy options for SADC member states *SADC Secretariat, Policy, Planning, Resource Mobilisation Directorate. Gaborone, Botswana* 56 (http://www.sadc.int/files/9113/6724/7724/SADC_Policy_Paper_Climate_Change_EN_1.pdf)
- Lewis S L et al 2013 Above-ground biomass and structure of 260 African tropical forests *Philosophical Transactions of the Royal Society B: Biological Sciences* **368** 20120295
- Liao W, V, Coillie F, Gao L, Li L, Zhang B and Chanussot J 2018 Deep learning for fusion of APEX hyperspectral and full-waveform LiDAR remote sensing data for tree species mapping *IEEE Access* **6** 68716–29
- Linares-Palomino R, Oliveira-Filho A T and Pennington R T 2011 Neotropical seasonally dry forests: diversity, endemism, and biogeography of woody plants *In Seasonally Dry Tropical Forests*. (Washington, DC: Island Press) 3–21
- Liu Y Y, Van Dijk A I, De Jeu R A, Canadell J G, McCabe M F, Evans J P and Wang G 2015 Recent reversal in loss of global terrestrial biomass *Nat. Clim. Change* **5** 470–4
- Liu Y Y, De Jeu R A, McCabe M F, Evans J P and Van Dijk A I 2011 Global long-term passive microwave satellite-based retrievals of vegetation optical depth *Geophys. Res. Lett.* **38** L18402
- Lu D 2006 The potential and challenge of remote sensing-based biomass estimation *Int. J. Remote Sens.* **27** 1297–328
- Madonsela S, Cho M A, Ramoelo A, Mutanga O and Naidoo L 2018 Estimating tree species diversity in the savannah using NDVI and woody canopy cover *Int. J. Appl. Earth Obs. Geoinf.* **66** 106–15
- Maglione P 2016 Very high resolution optical satellites: an overview of the most commonly used *American Journal of Applied Sciences* **13** 91
- Mapfumo R B, Murwira A, Masocha M and Andriani R 2016 The relationship between satellite-derived indices and species diversity across African savanna ecosystems *Int. J. Appl. Earth Obs. Geoinf.* **52** 306–17
- Mareya H T, Tagwireyi P, Ndaimani H, Gara T W and Gwenzi D 2018 Estimating tree crown area and aboveground biomass in miombo woodlands from high-resolution RGB-only imagery *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **11** 868–75

- Marín-Spiotta E, Cusack D F, Ostertag R and Silver W L 2008 Trends in above and belowground carbon with forest regrowth after agricultural abandonment in the neotropics *In Post-Agricultural Succession in the Neotropics*. (New York, NY: Springer) pp 22–72
- Marston C G, Wilkinson D M, Sponheimer M, Codron D, Codron J and O'Regan H J 2020 'Remote' behavioural ecology: do megaherbivores consume vegetation in proportion to its presence in the landscape? *PeerJ* **8** e8622
- Marta S 2018 *Planet Imagery Product Specifications*. (San Francisco, CA, USA: Planet Labs) 91
- Marumbwa F M, Cho M A and Chirwa P W 2021 Geospatial analysis of meteorological drought impact on Southern Africa biomes *Int. J. Remote Sens.* **42** 2155–73
- Matawire M M, Sibanda M and Dube T 2015 Assessing the aftermath of the fast track land reform programme in Zimbabwe on land-use and land-cover changes *Transactions of the Royal Society of South Africa* **70** 181–6
- Mathieu R et al 2013 Toward structural assessment of semi-arid African savannahs and woodlands: the potential of multitemporal polarimetric RADARSAT-2 fine beam images *Remote Sens. Environ.* **138** 215–31
- Mathieu R, Main R, Roy D P, Naidoo L and Yang H 2019 The effect of surface fire in savannah systems in the kruger national park (KNP), South Africa, on the backscatter of c-band sentinel-1 images *Fire* **2** 1–24
- Mayes M, Mustard J, Melillo J, Neill C and Nyadzi G 2017 Going beyond the green: senesced vegetation material predicts basal area and biomass in remote sensing of tree cover conditions in an African tropical dry forest (miombo woodland) landscape *Environ. Res. Lett.* **12** 085004
- Mayr M J, Vanselow K A and Samimi C 2018 Fire regimes at the arid fringe: a 16-year remote sensing perspective (2000–2016) on the controls of fire activity in Namibia from spatial predictive models *Ecol. Indic.* **91** 324–37
- Mbow C, Brandt M, Ouedraogo I, de Leeuw J and Marshall M 2015 What four decades of earth observation tell us about land degradation in the Sahel? *Remote Sensing* **7** 4048–67
- McCarthy J, Gumbrecht T and McCarthy T S 2005 Ecoregion classification in the Okavango Delta, Botswana from multitemporal remote sensing *Int. J. Remote Sens.* **26** 4339–57
- McKenzie J E, Brennan S E, Ryan R E, Thomson H J, Johnston R V and Thomas J 2019 Defining the criteria for including studies and how they will be grouped for the synthesis *Cochrane Handbook for Systematic Reviews of Interventions* **23** 33–65
- Méndez V E, Gliessman S R and Gilbert G S 2007 Tree biodiversity in farmer cooperatives of a shade coffee landscape in western El Salvador. *Agriculture Ecosystems & Environment* **119** 145–59
- Mendivelso H A, Camarero J J, Gutiérrez E and Zuidema P A 2014 Time-dependent effects of climate and drought on tree growth in a Neotropical dry forest: short-term tolerance versus long-term sensitivity *Agric. For. Meteorol.* **188** 13–23
- Metternicht G, Zinck J A, Blanco P D and del Valle H F 2010 Remote sensing of land degradation: experiences from Latin America and the Caribbean *J. Environ. Qual.* **39** 42–61
- Meyer T, D'Odorico P, Okin G S, Shugart H H, Caylor K K, O'Donnell F C, Bhattachan A and Dintwe K 2014 An analysis of structure: biomass structure relationships for characteristic species of the western Kalahari, Botswana *African Journal of Ecology* **52** 20–9
- Miles L, Newton A C, DeFries R S, Ravilious C, May I, Blyth S, Kapos V and Gordon J E 2006 A global overview of the conservation status of tropical dry forests *Journal of Biogeography* **33** 491–505
- Mitchard E T and Flintrop C M 2013 Woody encroachment and forest degradation in sub-Saharan Africa's woodlands and savannas 1982–2006 *Philosophical Transactions of the Royal Society B: Biological Sciences* **368** 20120406
- Mitchard E T, Saatchi S S, Woodhouse I H, Nangendo G, Ribeiro N S, Williams M, Ryan C M, Lewis S L, Feldpausch T R and Meir P 2009 Using satellite radar backscatter to predict above-ground woody biomass: a consistent relationship across four different African landscapes *Geophys. Res. Lett.* **36** L23401
- Moher D, Liberati A, Tetzlaff J, Altman D G and Prisma Group 2009 Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement *PLoS Med.* **6** e1000097
- Mooney H A, Bullock S H and Medina E 1995 Introduction ed S H Bullock, H A Mooney and E Medina *Seasonally Dry Tropical Forests*. (Cambridge: Cambridge University Press)
- Moore C E et al 2016 Reviews and syntheses: Australian vegetation phenology: new insights from satellite remote sensing and digital repeat photography *Biogeosciences* **13** 5085–102
- Muraoka H and Koizumi H 2009 Satellite Ecology (SATECO)—linking ecology, remote sensing and micrometeorology, from plot to regional scale, for the study of ecosystem structure and function *Journal of Plant Research* **122** 3–20
- Murphy P G and Lugo A E 1986 Ecology of tropical dry forest *Annual Review of Ecology and Systematics* **17** 67–88
- Mutanga O, Van Aardt J and Kumar L 2009 Imaging spectroscopy (Hyperspectral remote sensing) in Southern Africa: an overview *S. Afr. J. Sci.* **105** 193–8
- Mutanga O, Dube T and Ahmed F 2016 Progress in remote sensing: vegetation monitoring in South Africa *South African Geographical Journal* **98** 461–71
- Mutanga O and Rugege D 2006 Integrating remote sensing and spatial statistics to model herbaceous biomass distribution in a tropical savanna *Int. J. Remote Sens.* **27** 3499–514
- Naidoo Laven, Mathieu Renaud, Main Russell, Wessels Konrad and Asner Gregory P 2016 L-band Synthetic Aperture Radar imagery performs better than optical datasets at retrieving woody fractional cover in deciduous, dry savannahs *International Journal of Applied Earth Observation and Geoinformation* **52** 54–64
- Nanni A S, Sloan S, Aide T M, Graesser J, Edwards D and Grau H R 2019 The neotropical reforestation hotspots: a biophysical and socioeconomic typology of contemporary forest expansion *Global Environ. Change* **54** 148–59
- Narine L L, Popescu S C and Malambo L 2019 Synergy of ICESat-2 and Landsat for mapping forest aboveground biomass with deep learning *Remote Sensing* **11** 1503
- O'Connor Tim G, Puttick James R and Hoffman M Timm 2014 Bush encroachment in southern Africa: changes and causes *African Journal of Range & Forage Science* **31** 67–88
- Olson D M et al 2001 Terrestrial ecoregions of the world: a new map of life on earth a new global map of terrestrial ecoregions provides an innovative tool for conserving biodiversity *BioScience* **51** 933–8
- Olsson L, Eklundh L and Ardö J 2005 A recent greening of the Sahel—trends, patterns and potential causes *J. Arid. Environ.* **63** 556–66
- Owe M., de Jeu R. and Walker J. 2001 A methodology for surface soil moisture and vegetation optical depth retrieval using the microwave polarization difference index *IEEE Transactions on Geoscience and Remote Sensing* **39** 1643–54
- Paruelo J M, Teixeira M, Staiano L, Mastrángelo M, Amdan L and Gallego F 2016 An integrative index of Ecosystem Services provision based on remotely sensed data *Ecol. Indic.* **71** 145–54
- Pennington R T, Lehmann C E and Rowland L M 2018 Tropical savannas and dry forests *Current Biology* **28** R541–5
- Pennington R T and Ratter J A (ed) 2006 *Neotropical Savannas And Seasonally Dry Forests: Plant Diversity, Biogeography, And Conservation*. (Boca Raton, Florida, USA: CRC press)

- Pereira Júnior A C, Oliveira S L, Pereira J M and Turkman M A A 2014 Modelling fire frequency in a Cerrado savanna protected area *PLoS One* **9** e102380
- Petheram L, Campbell B M, Marunda C T, Tiveau D and Shackleton S 2006 The wealth of the dry forests: can sound forest management contribute to the millennium development goals in Sub-Saharan Africa? CIFOR Livelihood Brief No.5. Bogor, Indonesia, Center for International Forestry Research (CIFOR) (<https://cgspace.cgiar.org/handle/10568/19577>)
- Portillo C 2010 Assessing the Conservation status of Neotropical Dry forests using Geographic Information Systems and Optical Remote Sensing (<https://doi.org/10.7939/R36F06>)
- Portillo-Quintero C A, Sanchez A M, Valbuena C A, Gonzalez Y Y and Larreal J T 2012 Forest cover and deforestation patterns in the Northern Andes (Lake Maracaibo Basin): a synoptic assessment using MODIS and Landsat imagery *Appl. Geogr.* **35** 152–63
- Portillo-Quintero C A and Sánchez-Azofeifa G A 2010 Extent and conservation of tropical dry forests in the Americas *Biological Conservation* **143** 144–55
- Pricope N G and Binford M W 2012 A spatio-temporal analysis of fire recurrence and extent for semi-arid savanna ecosystems in southern Africa using moderate-resolution satellite imagery *J. Environ. Manage.* **100** 72–85
- Pricope N G, Gaughan A E, All J D, Binford M W and Rutina L P 2015 Spatio-temporal analysis of vegetation dynamics in relation to shifting inundation and fire regimes: disentangling environmental variability from land management decisions in a southern african transboundary watershed *Land* **4** 627–55
- Putz F E and Redford K H 2010 The importance of defining ‘forest’: tropical forest degradation, deforestation, long-term phase shifts, and further transitions *Biotropica* **42** 10–20
- Quesada M, Sanchez-Azofeifa G. A, Alvarez-Anorve M, Stoner K. E, Avila-Cabadilla L, Calvo-Alvarado J, Castillo A, Espirito-Santo M. M, Fagundes M, Fernandes G. W and Gamon J 2009 Succession and management of tropical dry forests in the Americas: Review and new perspectives *Forest Ecology and Management* **258** 1014–24
- Quijas S, Romero-Duque L P, Trilleras J M, Conti G, Kolb M, Brignone E and Dellafiore C 2019 Linking biodiversity, ecosystem services, and beneficiaries of tropical dry forests of Latin America: review and new perspectives *Ecosystem Services* **36** 100909
- Rahmoune R, Ferrazzoli P, Kerr Y H and Richaume P 2013 SMOS level 2 retrieval algorithm over forests: description and generation of global maps *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **6** 1430–9
- Reiche J et al 2016 Combining satellite data for better tropical forest monitoring *Nat. Clim. Change* **6** 120–2
- Reynolds J, Wesson K, Desbiez A L, Ochoa-Quintero J M and Leimgruber P 2016 Using remote sensing and random forest to assess the conservation status of critical Cerrado habitats in Mato Grosso do Sul, Brazil *Land* **5** 12
- Ringrose S, Chipanshi A C, Matheson W, Chanda R, Motoma L, Magole I and Jellema A 2002 Climate- and human-induced woody vegetation changes in Botswana and their implications for human adaptation *Environmental Management* **30** 98–109
- Rodríguez J P, Nassar J M, Rodríguez-Clark K M, Zager I, Portillo-Quintero C A, Carrasquel F and Zambrano S 2008 Tropical dry forests in Venezuela: assessing status, threats and future prospects *Environ. Conserv.* **35** 311–8
- Rouse J W 1974 Monitoring the vernal advancement of retrogradation of natural vegetation. NASA/GSFC, type III, final report (1974 Greenbelt, MD College Station, Texas **77843** 371
- Roy David P, Huang Haiyan, Boschetti Luigi, Giglio Louis, Yan Lin, Zhang Hankui H. and Li Zhongbin 2019 Landsat-8 and Sentinel-2 burned area mapping - A combined sensor multi-temporal change detection approach *Remote Sensing of Environment* **231** 111254
- Ryan Casey M., Pritchard Rose, McNicol Iain, Owen Matthew, Fisher Janet A. and Lehmann Caroline 2016 Ecosystem services from southern African woodlands and their future under global change *Philosophical Transactions of the Royal Society B: Biological Sciences* **371** 20150312
- Sader S A, Waide R B, Lawrence W T and Joyce A T 1989 Tropical forest biomass and successional age class relationships to a vegetation index derived from Landsat TM data *Remote Sens. Environ.* **28** 143–98
- Schmidhuber Jürgen 2015 Deep learning in neural networks: An overview *Neural Networks* **61** 85–117
- Sánchez-Azofeifa A, Portillo-Quintero C, Wilson-Fernandes G, Stoner K and Shimizu T 2013 The policy process for land use/cover change and forest degradation in the semi-arid Latin American/Caribbean region: perspectives and opportunities. A literature review prepared for the Inter-American Development Bank *A Literature Review Prepared for the Inter-American Development Bank*
- Sánchez-Azofeifa G A et al 2005 Research priorities for Neotropical dry forests 1 *Biotropica: The Journal of Biology and Conservation* **37** 477–85 (<https://www.jstor.org/stable/30043216>)
- Sánchez-Azofeifa G A, Castro K L, Rivard B, Kalascka M R and Harriss R C 2003 Remote sensing research priorities in tropical dry forest environments *Biotropica* **35** 134–42
- Santos J R, Lacruz M S P, Araujo L S and Keil M 2002 Savanna and tropical rainforest biomass estimation and spatialization using JERS-1 data *Int. J. Remote Sens.* **23** 1217–29
- Scholes R J, Frost P G H and Tian Y 2004 Canopy structure in savannas along a moisture gradient on Kalahari sands *Global Change Biol.* **10** 292–302
- Schröder Jobst Michael, Ávila Rodríguez Lina Paola and Günter Sven 2021 Research trends: Tropical dry forests: The neglected research agenda? *Forest Policy and Economics* **122** 102333
- Schultz M, Shapiro A, Clevers J G, Beech C and Herold M 2018 Forest cover and vegetation degradation detection in the Kavango Zambezi transfrontier conservation area using BFAST monitor *Remote Sensing* **10** 1850
- Shafaei M A., Salem M A.-M, Ebied H M, Al-Berry M N and Tolba M F 2018 Deep Learning for Satellite Image Classification *Deep Learning for Satellite Image Classification* (AISI 2018: Springer, Cham.) pp 383–91
- Shao Z, Zhang L and Wang L 2017 Stacked sparse autoencoder modeling using the synergy of airborne LiDAR and satellite optical and SAR data to map forest above-ground biomass *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **10** 5569–82
- Shao Z, Zhang L and Wang L 2017 Stacked sparse autoencoder modeling using the synergy of airborne LiDAR and satellite optical and SAR data to map forest above-ground biomass *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing* **10** 5569–82
- Shelestov A, Lavreniuk M, Kussul N, Novikov A and Skakun S 2017 Exploring google earth engine platform for big data processing: classification of multi-temporal satellite imagery for crop mapping *Frontiers in Earth Science* **5** 17
- Shi J, Jackson T, Tao J, Du J, Bindlish R, Lu L and Chen K S 2008 Microwave vegetation indices for short vegetation covers from satellite passive microwave sensor AMSR-E *Remote Sens. Environ.* **112** 4285–300
- Shimada M 2018 *Imaging from Spaceborne and Airborne SARs, Calibration, and Applications* (Boca Raton, Florida, USA: CRC Press)
- Silva J M N, Pereira J M C, Cabral A I, Sa A C L, Vasconcelos M J P, Mota B and Gregoire J M 2003 An estimate of the area burned in southern Africa during the 2000 dry season using SPOT-VEGETATION satellite data *Journal of Geophysical Research-Atmospheres* **108** NO. D13, 8498
- Smith W K et al 2019 Remote sensing of dryland ecosystem structure and function: progress, challenges, and opportunities *Remote Sens. Environ.* **233** 111401

- Stan K and Sanchez-Azofeifa A 2019 Deforestation and secondary growth in Costa Rica along the path of development *Regional Environmental Change* **19** 587–97
- Stan K and Sanchez-Azofeifa A 2019 Tropical dry forest diversity, climatic response, and resilience in a changing climate *Forests* **10** 443
- Staver A C, Archibald S and Levin S A 2011 The global extent and determinants of savanna and forest as alternative biome states *Science* **334** 230–2
- Stoner K E, Timm R M, Frankie G W, Mata A and Vinson S B 2004 *Tropical dry-Forest Mammals of Palo Verde. Biodiversity Conservation in Costa Rica. Learning the Lessons in a Seasonal Dry Forest*. (Berkeley, Los Angeles: University of California Press) 48–66
- Sunderland T et al 2015 Global dry forests: a prologue *International Forestry Review* **17** 1–9
- Suresh H S, Dattaraja H S, Mondal N and Sukumar R 2011 Seasonally dry tropical forests in Southern India. An analysis of floristic composition, structure, and dynamics in Mudumalai Wildlife Sanctuary *The Ecology and Conservation of seasonally Dry Forests in Asia* 37–58
- Symeonakis E, Higginbottom T P, Petroulaki K and Rabe A 2018a Optimisation of savannah land cover characterisation with optical and SAR data *Remote Sensing* **10** 499
- Symeonakis E, Korkofigkas A, Vamvoukakis G, Stamou G and Arnau-Rosalen E 2020 Deep learning monitoring of woody vegetation density in a South African Savannah region *International Archives of the Photogrammetry, Remote Sensing and Spatial Information Sciences- ISPRS Archives* **43** 1645–9
- Tian F, Brandt M, Liu Yi Y, Verger A, Tagesson T, Diouf A A, Rasmussen K, Mbow C, Wang Y and Fensholt R 2016 Remote sensing of vegetation dynamics in drylands: Evaluating vegetation optical depth (VOD) using AVHRR NDVI and in situ green biomass data over West African Sahel *Remote Sensing of Environment* **177** 265–76
- Tian F, Brandt M, Liu Y Y, Rasmussen K and Fensholt R 2017 Mapping gains and losses in woody vegetation across global tropical drylands *Global Change Biol.* **23** 1748–60
- Toth C and Józków G 2016 Remote sensing platforms and sensors: a survey *ISPRS J. Photogramm. Remote Sens.* **115** 22–36
- Trejo I and Dirzo R 2000 Deforestation of seasonally dry tropical forest: a national and local analysis in Mexico *Biological Conservation* **94** 133–42
- Trier Ø D, Salberg A B, Kermit M, Rudjord Ø, Gobakken T, Næsset E and Aarsten D 2018 Tree species classification in Norway from airborne hyperspectral and airborne laser scanning data *European Journal of Remote Sensing* **51** 336–51
- Tsalyuk M, Kelly M and Getz W M 2017 Improving the prediction of African savanna vegetation variables using time series of MODIS products *ISPRS J. Photogramm. Remote Sens.* **131** 77–91
- Tucker C J 1979 Red and photographic infrared linear combinations for monitoring vegetation *Remote Sens. Environ.* **8** 127–50
- Urban M, Berger C, Mudau T E, Heckel K, Truckenbrodt J, Onyango Odipo V, Smit I P and Schullius C 2018 Surface moisture and vegetation cover analysis for drought monitoring in the Southern Kruger National Park using sentinel-1, sentinel-2, and landsat-8 *Remote Sensing* **10** 1482
- Valentini R et al 2014 A full greenhouse gases budget of Africa: synthesis, uncertainties, and vulnerabilities *Biogeosciences* **11** 381–407
- Van Bommel F P J, Heitkönig I M A, Epema G F, Ringrose S, Bonyongo C and Veenendaal E M 2006 Remotely sensed habitat indicators for predicting distribution of impala (*Aepyceros melampus*) in the Okavango Delta, Botswana *Journal of Tropical Ecology* **22** 101–10
- Van Marle M J E, Van Der Werf G R, De Jeu R A M and Liu Y Y 2016 Annual South American forest loss estimates based on passive microwave remote sensing (1990–2010) *Biogeosciences* **13** 609–24
- Veldkamp A and Lambin E F 2001 Predicting land-use change *Agriculture, Ecosystems & Environment* **85** 6
- Venter Z S, Scott S L, Desmet P G and Hoffman M T 2020 Application of Landsat-derived vegetation trends over South Africa: potential for monitoring land degradation and restoration *Ecol. Indic.* **113** 106206
- Verbesselt J, Somers B, van Aardt J, Jonckheere I and Coppin P 2006 Monitoring herbaceous biomass and water content with SPOT VEGETATION time-series to improve fire risk assessment in savanna ecosystems *Remote Sens. Environ.* **101** 399–414
- Verbesselt J, Somers B, Lhermitte S, Jonckheere I, van Aardt J and Coppin P 2007 Monitoring herbaceous fuel moisture content with SPOT VEGETATION time-series for fire risk prediction in savanna ecosystems *Remote Sens. Environ.* **108** 357–68
- Verlinden A and Laamanen R 2006a Long term fire scar monitoring with remote sensing in northern Namibia: relations between fire frequency, rainfall, land cover, fire management and trees *Environ. Monit. Assess.* **112** 231–53
- Verlinden A and Laamanen R 2006b Modeling woody vegetation resources using Landsat TM imagery in northern Namibia *Southern African Forestry Journal* **207** 27–39
- Walter H and Burnett J H 1971 *Ecology of Tropical and Subtropical Vegetation*. 539 (Edinburgh: Oliver and Boyd) xviii + 539
- Wang L, Jia M, Yin D and Tian J 2019 A review of remote sensing for mangrove forests: 1956–2018 *Remote Sens. Environ.* **231** 111223
- Wang Y, Day J L and Davis F W 1998 Sensitivity of modeled C- and L-band radar backscatter to ground surface parameters in loblolly pine forest *Remote Sens. Environ.* **66** 331–42
- Wessels K J, Prince S D, Zambatis N, Macfadyen S, Frost P E and Van Zyl D 2006 Relationship between herbaceous biomass and 1-km(2) Advanced Very High Resolution Radiometer (AVHRR) NDVI in Kruger National Park, South Africa *Int. J. Remote Sens.* **27** 951–73
- Westinga E, Beltran A P R, De Bie C A and van Gils H A 2020 A novel approach to optimize hierarchical vegetation mapping from hyper-temporal NDVI imagery, demonstrated at national level for Namibia *Int. J. Appl. Earth Obs. Geoinf.* **91** 102152
- White B L A 2019 Satellite detection of wildland fires in South America *Floresta* **49** 851–8
- Whitecross M A, Witkowski E T F and Archibald S 2017 Assessing the frequency and drivers of early-greening in broad-leaved woodlands along a latitudinal gradient in southern Africa *Austral Ecology* **42** 341–53
- Williams M, Hill T C and Ryan C M 2013 Using biomass distributions to determine probability and intensity of tropical forest disturbance *Plant Ecology & Diversity* **6** 87–99
- Wingate V R, Phinn S R, Kuhn N and Scarth P 2018 Estimating aboveground woody biomass change in Kalahari woodland: combining field, radar, and optical data sets *Int. J. Remote Sens.* **39** 577–606
- Xue J and Su B 2017 Significant remote sensing vegetation indices: a review of developments and applications *Journal of Sensors* **2017**
- Yang J and Prince S 2000 Remote sensing of savanna vegetation changes in Eastern Zambia 1972–1989 *Int. J. Remote Sens.* **21** 301–22
- Zhou B, Okin G S and Zhang J 2020 Leveraging Google Earth Engine (GEE) and machine learning algorithms to incorporate *in situ* measurement from different times for rangelands monitoring *Remote Sens. Environ.* **236** 111521
- Zhu X X, Tuia D, Mou L, Xia G S, Zhang L, Xu F and Fraundorfer F 2017 Deep learning in remote sensing: a comprehensive review and list of resources *IEEE Geoscience and Remote Sensing Magazine* **5** 8–36
- Zhu Z 2017 Change detection using landsat time series: a review of frequencies, preprocessing, algorithms, and applications *ISPRS J. Photogramm. Remote Sens.* **130** 370–84
- Zhu Z and Woodcock C E 2012 Object-based cloud and cloud shadow detection in Landsat imagery *Remote Sens. Environ.* **118** 83–94