

Density functionals with spin-density accuracy for open shells

Accepted Manuscript: This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination, and proofreading process, which may lead to differences between this version and the Version of Record.

Cite as: J. Chem. Phys. (in press) (2021); <https://doi.org/10.1063/5.0071991>

Submitted: 18 September 2021 • Accepted: 24 February 2022 • Accepted Manuscript Online: 24 February 2022

Timothy Callow, Benjamin Pearce and  Nikitas Gidopoulos



View Online



Export Citation



CrossMark

ARTICLES YOU MAY BE INTERESTED IN

[A range-separated generalized Kohn–Sham method including a long-range nonlocal random phase approximation correlation potential](#)

The Journal of Chemical Physics **153**, 244118 (2020); <https://doi.org/10.1063/5.0031310>

[Assessing locally range-separated hybrid functionals from a gradient expansion of the exchange energy density](#)


The Journal of Chemical Physics **154**, 214101 (2021); <https://doi.org/10.1063/5.0047628>

[Software for the frontiers of quantum chemistry: An overview of developments in the Q-Chem 5 package](#)

The Journal of Chemical Physics **155**, 084801 (2021); <https://doi.org/10.1063/5.0055522>

Webinar

Interfaces: how they make
or break a nanodevice



March 29th – Register now



Zurich
Instruments



J. Chem. Phys. (in press) (2021); <https://doi.org/10.1063/5.0071991>

(c) 2022 Author(s).

Density functionals with spin-density accuracy for open shells

Timothy Callow,^{1,2,3,4} Benjamin Pearce,¹ and Nikitas Gidopoulos¹

¹*Department of Physics, Durham University, South Road, Durham, DH1 3LE, UK*

²*Max-Planck-Institut für Mikrostrukturphysik, Weinberg 2, D-06120 Halle, Germany*

³*Center for Advanced Systems Understanding (CASUS), D-02826 Görlitz, Germany*

⁴*Helmholtz-Zentrum Dresden-Rossendorf, D-01328 Dresden, Germany*

(*Electronic mail: nikitas.gidopoulos@durham.ac.uk)

(*Electronic mail: b.j.pearce@durham.ac.uk)

(*Electronic mail: t.callow@hzdr.de)

(Dated: 18 February 2022)

Electrons in zero external magnetic field can be studied with the Kohn–Sham (KS) scheme of either density functional theory (DFT) or spin-DFT (SDFT). The latter is normally used for open shell systems because its approximations appear to model better the exchange and correlation (xc) functional, but also because so far the application of DFT implied a closed-shell-like approximation. In the first part of this Communication, we show that correcting this error for open shells allows the approximate DFT xc functionals to become as accurate as those in SDFT. In the second part, we consider the behaviour of SDFT for zero magnetic field. We show that the KS equations of SDFT emerge as the generalized KS equations of DFT in this limit, thus establishing a so-far unknown link between the two theories.

Keywords: local density approximation, implicit density functionals, semi-local density approximation

I. INTRODUCTION

Density functional theory (DFT)^{1,2} and spin-density functional theory (SDFT)^{3,4} are two reformulations of the many-electron problem. Their computational advantage is replacing the solution of Schrödinger’s equation for the multi-dimensional ground state of the physical electronic system of interest by the solution of a much less demanding equation that yields the ground-state (total) density ρ (DFT) or the spin-density ($\rho^\uparrow, \rho^\downarrow$) (SDFT) of the real system.⁵

Both theories can in principle be applied equally well to study electronic systems when the external magnetic field, B , vanishes ($B = 0$). Then, obviously, the predicted values for any observable quantity must be the same between the two, formally exact, theories. Specifically, the ground-state (total) density and energy of the electronic system must be the same in both theories.

Even though for $B = 0$ both exact theories can equally well describe an electronic system, this changes when density functional approximations (DFAs) are introduced. The flexibility of two basic variables in SDFT is thought to offer a better modelling of the exchange–correlation (xc) energy, $E_{xc}[\rho^\uparrow, \rho^\downarrow]$, compared with just a single variable in DFT, $E_{xc}[\rho]$. As Parr and Yang write⁶, in local and semi-local density functional approximations “the exchange–correlation energy of the electrons is approximated locally by results for the homogeneous spin-compensated electron gas. Such a procedure is not appropriate for systems with unpaired electrons, like open-shell molecules. A better description for such systems will be obtained through the use of the exchange–correlation energy of the homogeneous spin-polarised electron gas”. As a result, open shell systems are in practice usually treated in SDFT rather than DFT.

The first aim of this Communication is to revisit the above widely-held view, namely that approximations in SDFT are inherently superior to those of DFT. As it will transpire, this

view is a misconception, perhaps abetted by a qualitative error in the standard treatment of open-shells with KS-DFT approximations. We shall explain the source of this error and show that correcting for it yields KS-DFT results that are (almost) as accurate as the corresponding KS-SDFT results.

Although this result goes against current wisdom, its corollary is positive for the DFT literature in general. Take for example the xc derivative discontinuity, the shift of the KS-DFT xc potential when the number of electrons crosses an integer number of particles from N to $N + 1$. Obviously, either N or $N + 1$ is an odd number, and in the analysis of the derivative discontinuity it is assumed that it is possible to calculate the xc potential for an odd number of electrons accurately, without qualitative errors. With the present work, this assumption is justified, even for approximate xc potentials⁷.

In (S)DFT the value of any observable quantity becomes a functional of the (spin) density. So far, little is known about the functional dependence of most observable quantities, except the total energy and identically the total density (DFT) or the spin-density (SDFT). For open shell systems and for $B = 0$, the virtual KS systems of DFT and SDFT are not the same (despite sharing the same total density), and the single-particle KS equations of exact SDFT do not reduce to those of exact DFT. In DFT the (unknown) density functional for the spin density is required to obtain the physical spin-density.

The second aim of this Communication is to investigate the link between exact DFT and SDFT when $B = 0$. In general, a pair of external potentials is necessary for the formulation of SDFT (conjugate to the pair of spin-densities), whereas when $B = 0$, there is only one external potential. So, either SDFT becomes singular in this limit, or the KS-SDFT equations must reduce to a set of (yet unknown) single-particle equations of DFT. In section IV of this Communication, we show that when $B = 0$ the KS-SDFT equations reduce to the equations of generalized KS-DFT (GKS-DFT), establishing a link between DFT and SDFT. In addition, the spin-density of

the GKS-DFT system emerges as the unknown density functional for the physical (real) spin-density.

For simplicity, we shall discuss the special case of SDFT for a collinear magnetisation density $m = -\mu_B(\rho^\uparrow - \rho^\downarrow)$. We shall also restrict to densities that are pure-state v -representable (both for the interacting and for the KS system⁸), with ground states having an integer number of spin-up and spin-down electrons⁹. Finally, we consider that the KS states are Slater determinants.

We return now to the first aim of our Communication. For $B = 0$, the main error in modelling open shell electronic systems as unpolarised arises in the exchange energy functional, $E_x[\rho]$ (DFT) vs $E_x[\rho^\uparrow, \rho^\downarrow]$ (SDFT). When a local or semi-local approximate exchange energy expression is employed, depending explicitly only on the density ρ and its derivatives, it is essentially assumed that the spin-up and spin-down densities are equal to each other and to one half of the total density. For open shell systems, this amounts to mixing partly the spin-up ρ^\uparrow with the spin-down ρ^\downarrow densities in the exchange energy expression:

$$E_x[\rho] \simeq E_x[\rho/2, \rho/2], \quad \rho = \rho^\uparrow + \rho^\downarrow. \quad (1)$$

This mixing of the spin-densities in the exchange energy leads to a spurious error that we call the ghost exchange energy error, in analogy to the ‘ghost interaction’ error of Ref.¹⁰. We define it by the difference of the exchange energies with mixed spin-densities from the reference exchange energy where the spin-densities are separate:

$$G_x = E_x[\rho/2, \rho/2] - E_x[\rho^\uparrow, \rho^\downarrow]. \quad (2)$$

In DFT, the ghost exchange error is not caused by the approximate or exact expression for the exchange energy but by the mixing of two spin-densities when an open shell system is treated as if it were closed-shell, considering half its electrons as spin-up and half as spin-down.

In the following sections II-III, we show that open shell systems in zero magnetic field can be modelled within DFT, without any mixing of spin-up and spin-down KS orbitals and avoiding the ghost-exchange error. Then, we explore the link between exact DFT and SDFT in the limit $B = 0$ in section IV. Finally, in section V, we discuss the famous spin-symmetry dilemma¹¹ in the context of our work.

Before proceeding, we note that there are various extensions of KS-DFT targeted for open-shell systems. For example, we mention the restricted open-shell KS (ROKS) method^{12,13}, KS-DFT with complex, spin-restricted KS orbitals¹⁴ and the recent progress in ensemble DFT for degenerate/excited states¹⁵⁻²⁰. Methods with multi-determinant KS reference states, or using ensemble DFT, are indeed intrinsically advantageous for open-shell systems. Our approach is restricted to single-determinant reference states and cannot be expected to compete in accuracy. Nevertheless, we maintain the practical advantage of KS-DFT, by employing its standard and popular approximations, while our aim has been to understand better the basic theory and to reveal the missing link between KS-DFT and KS-SDFT.

II. THE XC ENERGY AS AN IMPLICIT DENSITY FUNCTIONAL

To continue, we consider the exact exchange energy in KS-DFT. It is given by the Fock expression in terms of the spin-orbitals of the KS Slater determinant,

$$E_x[\rho^\uparrow, \rho^\downarrow] = -\frac{1}{2} \iint d\mathbf{r} d\mathbf{x} \left\{ \frac{|\rho^\uparrow(\mathbf{r}, \mathbf{x})|^2}{|\mathbf{r} - \mathbf{x}|} + \frac{|\rho^\downarrow(\mathbf{r}, \mathbf{x})|^2}{|\mathbf{r} - \mathbf{x}|} \right\}. \quad (3)$$

where $\rho^\sigma(\mathbf{r}, \mathbf{x})$, $\sigma = \uparrow, \downarrow$, is the spin- σ , one-body reduced density matrix of KS orbitals.

The exact exchange energy functional separates in two disjoint terms,

$$E_x[\rho^\uparrow, \rho^\downarrow] = E_x[\rho^\uparrow, 0] + E_x[0, \rho^\downarrow], \quad (4)$$

in which the subsets of spin-up and spin-down KS orbitals do not mix. Hence, in order to avoid cross exchange effects between opposite spin-electrons also in approximations, the approximate exchange energy density functionals must also satisfy equality (4).

Since modelling the approximate exchange energy in terms of the total density and its derivatives, as in LDA and in semi-local DFAs, always violates equality (4), we have to model the exchange energy using the DFT-KS spin-density²¹. This modelling is still within DFT (not SDFT) since in DFT the KS spin-density (the spin-density of the KS determinant) is an implicit functional of the total density.

Equality (4) is satisfied by the local spin-density approximation (LSDA), and as far as we know all spin-dependent density functionals (DFs) for the exchange energy, approximate or the exact one. Therefore, we shall use expressions for the exchange energy which depend on the total density indirectly, or implicitly, via the KS spin-density (spin-density functionals, SDFs),

$$E_x^{\text{iDF}}[\rho] = E_x^{\text{SDF}}[\rho^\uparrow[\rho], \rho^\downarrow[\rho]], \quad (5)$$

where E_x^{SDF} must be the sum of two disjoint terms (4).

The acronyms iDF and SDF denote that the exchange energy on the lhs is an *implicit* density-functional and that the exchange energy expression on the rhs depends on the KS spin-density.

Eq. 5 not only (re)defines for any DFA a new, ghost-exchange-error free, x energy functional, but it is also a relation satisfied by the exact x energy functionals of DFT and SDFT. In other words, for exact exchange: $E_x[\rho] = E_x^{\text{iDF}}[\rho]$.

In order to keep the treatment of xc uniform, we decompose the correlation energy DF in an implicit DF, following (5), plus a remainder:

$$E_c[\rho] = E_c^{\text{iDF}}[\rho] + \Delta E_c[\rho], \quad (6)$$

$$E_c^{\text{iDF}}[\rho] = E_c^{\text{SDF}}[\rho^\uparrow[\rho], \rho^\downarrow[\rho]]. \quad (7)$$

$\Delta E_c[\rho] \neq 0$ in general, for open shell systems. In the following, we omit ΔE_c (which makes the rest of the treatment approximate) and write: $E_c[\rho] \approx E_c^{\text{iDF}}[\rho]$. In (semi)-local DFAs,

this redefinition of the correlation energy as an implicit DF yields a small accuracy gain, though this is typically a minor contribution compared to the ghost exchange error. The whole exchange-correlation (xc) density functional is thus written as

$$E_{xc}^{\text{iDF}}[\rho] \approx E_{xc}^{\text{SDF}}[\rho^\uparrow[\rho], \rho^\downarrow[\rho]]. \quad (8)$$

Hence, the total energy DF is now given by,

$$E_{\text{ven}}^{\text{iDF}}[\rho] = T_s[\rho] + \int d\mathbf{r} v_{\text{en}}(\mathbf{r})\rho(\mathbf{r}) + U[\rho] + E_{xc}^{\text{iDF}}[\rho], \quad (9)$$

where $U[\rho]$ is the Hartree energy and $v_{\text{en}}(\mathbf{r})$ is the external (electron-nuclear) potential of the interacting system.

As the xc energy functional is an implicit DF, the xc-potential, given by the functional derivative

$$v_{xc}[\rho](\mathbf{r}) = \frac{\delta E_{xc}^{\text{iDF}}[\rho]}{\delta \rho(\mathbf{r})}, \quad (10)$$

must be determined using the optimised effective potential (OEP) method^{22,23},

$$\int d\mathbf{r}' \sum_{\sigma} \chi^{\sigma}(\mathbf{r}, \mathbf{r}') v_{xc}[\rho](\mathbf{r}') = \int d\mathbf{r}' \sum_{\sigma} \chi^{\sigma}(\mathbf{r}, \mathbf{r}') v_{xc}^{\sigma}[\rho^\uparrow[\rho], \rho^\downarrow[\rho]](\mathbf{r}'), \quad (11)$$

with

$$v_{xc}^{\sigma}[\rho^\uparrow[\rho], \rho^\downarrow[\rho]](\mathbf{r}') = \left. \frac{\partial E_{xc}^{\text{SDF}}[\rho^\uparrow, \rho^\downarrow]}{\partial \rho^{\sigma}(\mathbf{r}')} \right|_{\substack{\rho^\uparrow = \rho^\uparrow[\rho] \\ \rho^\downarrow = \rho^\downarrow[\rho]}}, \quad (12)$$

$$\chi^{\sigma}(\mathbf{r}, \mathbf{r}') = -2 \sum_{i=1}^{N^{\sigma}} \sum_{a=N^{\sigma}+1}^{\infty} \frac{\phi_i(\mathbf{r})\phi_a(\mathbf{r})\phi_i(\mathbf{r}')\phi_a(\mathbf{r}')}{\varepsilon_a - \varepsilon_i}, \quad (13)$$

where $\chi^{\sigma}(\mathbf{r}, \mathbf{r}')$ is the spin- σ density-density response function²⁴.

The derivation of equations (8-11) constitutes our first important result. These equations determine the ghost-exchange-error-corrected xc energy and potential for open shell systems, in (semi)-local DFAs. From (11) we obtain the DFT-KS xc potential v_{xc} , which is used in the usual way during the self-consistent KS calculation. It is given as the weighted sum of the spin-dependent xc potentials v_{xc}^{σ} , with weighting factors the spin-dependent response functions χ^{σ} .

For closed shell systems, the solution of (11) reduces to the familiar KS xc functional derivative. For fully spin-polarised systems (eg spin up, $\rho^\uparrow = \rho, \rho^\downarrow = 0$), the spin-down response function vanishes, and the solution of (11) reduces to the spin-up xc potential, $v_{xc}[\rho] = v_{xc}^{\uparrow}[\rho, 0]$. Even for partially spin-polarised, open-shell systems, the spin-up (i.e. the majority spin) xc potential is still an accurate approximation to the full solution of (11). Firstly, this is because the latter must interpolate smoothly between the two limits just described, where $v_{xc} = v_{xc}^{\uparrow}$. Secondly, because the weighting of the spin- σ response function in (11) depends roughly on the inverse of the HOMO-LUMO gap, Δ^{σ} , of the spin- σ KS orbitals, and since $\Delta^{\uparrow} < \Delta^{\downarrow}$, the spin-up response function dominates. Another

	E_{LDA} (Ha)	E_{iLDA} (Ha)	E_{LSDA} (Ha)
Li	-7.388721	-7.398145	-7.398155
B	-24.43315	-24.44669	-24.44747
N	-54.12891	-54.14996	-54.15110
Na	-161.6491	-161.6571	-161.6572
Si ^a	-288.4640	-288.4905	-288.4910
LiH ⁺	-7.652062	-7.685603	-7.685608
O ₂ ^a	-149.6038	-149.6383	-149.6403
OH	-75.34813	-75.37077	-75.37208
NH ₄	-56.79800	-56.80389	-56.80404
Avg diff (mHa)	20.1	0.666	-

^a Triplet state

TABLE I. L(S)DA ground-state total energies calculated with: (i) standard LDA; (ii) implicit LDA (iLDA); and (iii) spin-LDA (LSDA). All states are doublets unless specified otherwise; auxiliary bases are uncontracted cc-pVTZ.

meaningful approximation for the solution is the weighted average, $v_{xc} \approx (\Delta^{\downarrow} v_{xc}^{\uparrow} + \Delta^{\uparrow} v_{xc}^{\downarrow}) / (\Delta^{\uparrow} + \Delta^{\downarrow})$. In the Supplementary Material we compare these two approximations with the full solution of the OEP equation (11).

III. RESULTS

We have implemented equations (5-11) in the Gaussian basis set code HIPPO²⁵. The solution of finite-basis OEP equations hides non-analytic behaviour²⁶, which leads to oscillations of the potential near the nuclei^{27,28}. We overcome these issues routinely following the method in Refs. 7 and 26 (please see Refs. 7 and 29 for details of the computational implementation). Unless otherwise stated, all results use cc-pVTZ bases for the KS orbitals and uncontracted cc-pVDZ auxiliary bases^{30,31} for the expansion of the ‘‘electron repulsion’’ or ‘‘screening’’ density^{7,32,33}.

Ground-state energies calculated with our iDF KS method compare favourably with those from SDFT-KS. This is demonstrated in Table I for some atoms and molecules at their equilibrium geometries, using the L(S)DA functional. We emphasise again that we seek only the ground-state energy in our approach, and thus where a state is referred to as ‘doublet’ this simply means

$$\Delta N = N^{\uparrow} - N^{\downarrow} = 1, \quad (14)$$

and likewise for triplet and so on. We also note the effect of the ghost exchange (and correlation) error, which is particularly pronounced for triplet states; we shall now explore some examples in which this error significantly affects results.

In Fig 1, we have calculated the energy dissociation curve for the H₂ molecule with the L(S)DA functional. As is well-known in the literature³⁴ and can be seen in this figure, the standard restricted solution yields a qualitatively incorrect dissociation curve; however, in our new method, once the bond is stretched enough the triplet state becomes lower in energy than the singlet, and the energy tends to the correct limit. This transition does not occur when the ghost exchange energy is




FIG. 1. Energy dissociation curves for the H_2 molecule. Top, iLDA energies for different values of ΔN ; bottom, comparison of LDA, spin LDA (LSDA), and iLDA minimum energies.

present: in this case the triplet energy is higher than the singlet at all bond distances.

We see a similar picture emerge for the stretched OH radical with the Perdew-Burke-Ernzerhof (PBE)³⁵ functional in Fig. 2. In this case, the energy dissociation curve is again qualitatively inaccurate for the solution contaminated with ghost-exchange errors; in our method which removes the ghost-exchange error, the energy correctly becomes equal to the energy of the two atoms in the infinitely stretched limit. In both these examples, the transition region is interesting: the unrestricted solution yields a smooth dissociation curve, whereas in our single-determinant method the transition is abrupt. Of course, our method does not correct the ubiquitous inability of typical semi-local functionals to capture static correlation effects³⁶. Naturally, using a spin-restricted formalism mandates constraints absent in an unrestricted scheme: for example, the HOMO and LUMO KS energy levels in a “doublet” state are by definition degenerate, though this is not typically the case for the corresponding energy levels in an unrestricted scheme.

IV. LIMIT OF SDFT-KS EQUATIONS FOR $B = 0$.

We now turn our attention to the limit of SDFT when $B = 0$ and compare with DFT. We start with the universal internal




FIG. 2. Energy dissociation curves for the OH radical. Top, implicit PBE (iPBE) energies for different values of ΔN ; bottom, comparison of PBE as total density functional (D-PBE), spin-PBE (SPBE), and iPBE minimum energies.

energy density functional,

$$F[\rho] = \min_{\Psi \rightarrow \rho} \langle \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle. \quad (15)$$

The minimising state Ψ_ρ depends on the total density ρ . When ρ is the ground state density of an interacting electronic system bound by a local potential v (i.e. when ρ is pure-state interacting v -representable), Ψ_ρ is the corresponding ground state. We remind the reader that we restricted our study to densities that are pure-state v -representable, both for the interacting and for the KS system⁸. The numbers of spin-up/down electrons N^\uparrow, N^\downarrow are good quantum numbers for Hamiltonians with collinear B , including $B = 0$. It follows the ground state Ψ_ρ can always be selected to have integer N^\uparrow, N^\downarrow . In DFT-KS theory, the KS reference state Φ_ρ is obtained from a similar minimisation to (15) for the noninteracting kinetic energy density functional, $T_s[\rho] = \min_{\Phi \rightarrow \rho} \langle \Phi | \hat{T} | \Phi \rangle$. Under the same assumptions, the KS ground state Φ_ρ can also be chosen to have integer N^\uparrow, N^\downarrow and be a Slater determinant.

To proceed, we follow Levy³⁷ and separate the minimisation in (15) into two separate minimisations with the same minimum,

$$F[\rho] = \min_{(\rho^\uparrow, \rho^\downarrow) \rightarrow \rho} \left[\min_{\Psi \rightarrow (\rho^\uparrow, \rho^\downarrow)} \langle \Psi | \hat{T} + \hat{V}_{ee} | \Psi \rangle \right]. \quad (16)$$

The inner minimisation, performed first, is over all states Ψ with spin-density $(\rho^\uparrow, \rho^\downarrow)$. The outer minimisation is over all spin-densities that add up to the total density ρ . The first minimisation inside the brackets on the rhs defines SDFT’s internal energy functional; we conclude that the DFT internal energy functional can be obtained from the SDFT functional with an extra optimisation over spin-densities with the same ρ (see also Appendix B in Ref. 11),

$$F[\rho] = \min_{(\rho^\uparrow, \rho^\downarrow) \rightarrow \rho} F[\rho^\uparrow, \rho^\downarrow]. \quad (17)$$

The two minimisations in (16) have the same overall minimum as the minimisation in (15). Therefore, the minimising spin-density, $(\rho_\rho^\uparrow, \rho_\rho^\downarrow)$, is equal to the spin-density of Ψ_ρ in (15) and hence equal to the spin-density of the real interacting system, even though the minimisation (17) is at fixed total density (not fixed spin-density), i.e. within DFT, not SDFT.

The minimisation in (17) is worth investigating further. We invoke SDFT’s KS system with spin-density $(\rho^\uparrow, \rho^\downarrow)$, to expand the internal energy functional in the usual way, $F[\rho^\uparrow, \rho^\downarrow] = T_s[\rho^\uparrow, \rho^\downarrow] + E_{xc}[\rho^\uparrow, \rho^\downarrow] + U[\rho^\uparrow + \rho^\downarrow]$. Dropping the superscript SDF from the functionals of SDFT, we obtain

$$F[\rho] = \min_{(\rho^\uparrow, \rho^\downarrow) \rightarrow \rho} \left\{ T_s[\rho^\uparrow, \rho^\downarrow] + E_{xc}[\rho^\uparrow, \rho^\downarrow] \right\} + U[\rho]. \quad (18)$$

Since every SDFT-KS system has a different spin-density, the minimisation effectively searches over all SDFT-KS systems with common total density ρ and returns that with the correct spin-density $(\rho_\rho^\uparrow, \rho_\rho^\downarrow)$. The KS Slater determinant state is $\Phi_{\rho_\rho^\uparrow, \rho_\rho^\downarrow}$. The minimising SDFT-KS system depends only on the total density ρ . The corresponding SDFT-KS potential (functional of the spin-density) at the specific spin-density $(\rho_\rho^\uparrow, \rho_\rho^\downarrow)$ is also an implicit functional of ρ . In the Supplementary Material we show how this unrestricted KS potential emerges directly from the minimisation in (18).

The SDFT-KS system that minimises (18) defines a new non-interacting system in DFT (when $B = 0$) represented by the state $\Phi_{\rho_\rho^\uparrow, \rho_\rho^\downarrow}$. We call it the generalised KS (GKS) system, as its derivation is analogous to the well-known GKS scheme in the literature³⁸. However, unlike typical GKS schemes, which employ a nonlocal potential term, the potential in the present DFT-GKS scheme is local and spin-dependent.

We conclude that the SDFT-KS equations reduce to the DFT-GKS equations for $B = 0$. The DFT-GKS Slater determinant state $\Phi_{\rho_\rho^\uparrow, \rho_\rho^\downarrow}$ gives not only the true total density but also the true (i.e. physical, or observable) spin-density. The elusive exact density functional for the spin-density is the spin-density of the DFT-GKS system.

V. THE SPIN-SYMMETRY DILEMMA

One of the oft-cited issues with SDFT in the absence of an external magnetic field is the spin-symmetry dilemma, so-called because SDFT approximate results yield either accurate total energies but with a poor (broken symmetry) prediction

for the spin-density, or an accurate prediction for the spin-density with poor total energies^{11,39,40}. The dilemma lies in that the spin-density is the key quantity and hence SDFT (exact) results are supposed to yield both the exact total energy and the exact spin-density. For example, H_2 should dissociate into two Hydrogen atoms with zero net magnetisation (difference of spin densities): indeed the lowest energy solution of local and semi-local approximations in SDFT gets the energy right, but it wrongly dissociates the molecule into two atoms of opposite spin (broken symmetry solution). There is also a self-consistent solution where the spin-densities are correctly equal to each other but the total energy is too high.

In their famous paper¹¹, Perdew, Savin and Burke “escape” the spin-symmetry dilemma, by postulating a “nearly exact alternative theory” where the spin-densities obtained from (semi)-local functionals “are not physical spin densities, but are only intermediate objects (like the Kohn-Sham orbitals)”. Instead, the on-top electron pair density $P(\mathbf{r}, \mathbf{r})$, alongside the total density, become the two fundamental variables of the new theory. As an aside, we note that $P(\mathbf{r}, \mathbf{r}')$ is itself a useful quantity and has been used to develop improved LSD approximations for multi-determinant reference states⁴¹. We share the principle of this paper, namely that the spin-density is no longer the basic quantity in the absence of a magnetic field; but in our approach, we do not consider any alternative theory: our (ghost-exchange-error free) theory yields the correct ground-state energy and the correct total density in the dissociation limit, and thus the spin-symmetry dilemma is weakened as a fundamental problem, since the KS spin-density is not expected equal to the observable spin-density.

However, we also propose that the DFT-GKS spin-density is the exact density functional of the observable spin-density; knowledge of the (exact) SDFT xc functional $E_{xc}[\rho^\uparrow, \rho^\downarrow]$ is necessary to obtain from the density the (exact) DFT-GKS spin-density. The value of the latter is the same as the SDFT-KS spin-density⁴² which, in local and semi-local approximations, demonstrates the spin-symmetry dilemma. We believe the explanation is that the approximate DFT-GKS spin-density is not a sufficiently accurate density functional for the real spin-density. According to Cohen, Sanchez and Yang^{34,36} on the fractional spin error, the crux of the problem is the degeneracy of different Slater determinants with the same energy and total density but different spin-densities. This kind of strong correlation cannot be captured with local or semi-local functionals.

VI. CONCLUSION

We have addressed two related conceptual or fundamental theoretical questions in DFT and spin-DFT. The first is the challenge how to apply the KS equations of DFT for open shells, avoiding a serious qualitative error, which we called the “ghost-exchange error”. Associated with this problem is the widespread belief that spin-DFT approximations are inherently more flexible and hence can model more accurately the xc energy for open shells than the corresponding approximations of DFT. In the literature this common view seemed

plausible, almost obvious. We demonstrate that actually density functionals can be as accurate as spin-density functionals, provided the ghost-exchange error is corrected.

For zero external magnetic field ($B=0$), both DFT and spin-DFT can obviously be applied to study open shell electronic systems and both exact theories should give identical results for any observable quantity. Then, intuitively, one would expect in the limit $B = 0$ the single particle KS equations of spin-DFT to reduce to a set of single-particle equations of DFT. We demonstrate this intuition is correct and that the SDFT equations reduce to the generalised KS equations of DFT.

SUPPLEMENTARY MATERIAL

In the Supplementary Material we compare the full solution of the OEP equation (11) with two simple approximations and we provide an alternative derivation of the DFT-GKS system.

ACKNOWLEDGMENTS

We thank N. Lathiotakis for help with the code HIPPO. We acknowledge enlightening discussions with K. Burke, E.K.U. Gross and N. Lathiotakis. N.I.G. acknowledges financial support from The Leverhulme Trust, through a Research Project Grant with number RPG-2016-005. B.P. acknowledges financial support from EPSRC with grant number EP/R513039/1. During the final stage of the project T.C. was funded by the Center for Advanced Systems Understanding (CASUS) which is financed by Germany's Federal Ministry of Education and Research (BMBF) and by the Saxon Ministry for Science, Culture and Tourism (SMWK) with tax funds on the basis of the budget approved by the Saxon State Parliament.

DATA AVAILABILITY

The data that support the findings of this study are available from the corresponding author upon reasonable request.

¹P. Hohenberg and W. Kohn, "Inhomogeneous electron gas," *Phys. Rev.* **136**, B864–B871 (1964).

²W. Kohn and L. J. Sham, "Self-consistent equations including exchange and correlation effects," *Phys. Rev.* **140**, A1133–A1138 (1965).

³U. von Barth and L. Hedin, "A local exchange-correlation potential for the spin polarized case. i," *Journal of Physics C: Solid State Physics* **5**, 1629 (1972).

⁴N. I. Gidopoulos, "Potential in spin-density-functional theory of non-collinear magnetism determined by the many-electron ground state," *Phys. Rev. B* **75**, 134408 (2007).

⁵For simplicity, we discuss the special case of SDFT for a collinear magnetisation density $m = -\mu_B(\rho^\uparrow - \rho^\downarrow)$; the generalisation to the noncollinear case is straightforward.

⁶R. G. Parr and W. Yang, "Density functional theory of atoms and molecules," (Oxford University Press, 1989) Chap. 8, pp. 173–174.

⁷T. J. Callow, B. J. Pearce, T. Pitts, N. N. Lathiotakis, M. J. P. Hodgson, and N. I. Gidopoulos, "Improving the exchange and correlation potential in density-functional approximations through constraints," *Faraday Discuss.* **224**, 126–144 (2020).

⁸P. R. T. Schipper, O. V. Gritsenko, and E. J. Baerends, "One - determinantal pure state versus ensemble kohn-sham solutions in the case of strong electron correlation: Ch_2 and c_2 ," *Theoretical Chemistry Accounts* **99**, 329–343 (1998).

⁹The interacting Hamiltonian and the KS Hamiltonian commute with the spin-up and spin-down number operators⁴.

¹⁰N. I. Gidopoulos, P. G. Papaconstantinou, and E. K. U. Gross, "Spurious interactions, and their correction, in the ensemble-kohn-sham scheme for excited states," *Phys. Rev. Lett.* **88**, 033003 (2002).

¹¹J. P. Perdew, A. Savin, and K. Burke, "Escaping the symmetry dilemma through a pair-density interpretation of spin-density functional theory," *Phys. Rev. A* **51**, 4531–4541 (1995).

¹²M. Filatov and S. Shaik, "Spin-restricted density functional approach to the open-shell problem," *Chemical Physics Letters* **288**, 689 – 697 (1998).

¹³M. Filatov and S. Shaik, "Application of spin-restricted open-shell kohn–sham method to atomic and molecular multiplet states," *J. Chem. Phys.* **110**, 116–125 (1999).

¹⁴J. Lee, L. W. Bertels, D. W. Small, and M. Head-Gordon, "Kohn-Sham Density Functional Theory with Complex, Spin-Restricted Orbitals: Accessing a New Class of Densities without the Symmetry Dilemma," *Phys. Rev. Lett.* **123**, 113001 (2019).

¹⁵T. Gould and S. Pittalis, "Hartree and exchange in ensemble density functional theory: Avoiding the nonuniqueness disaster," *Phys. Rev. Lett.* **119**, 243001 (2017).

¹⁶Z.-h. Yang, A. Pribram-Jones, K. Burke, and C. A. Ullrich, "Direct extraction of excitation energies from ensemble density-functional theory," *Phys. Rev. Lett.* **119**, 033003 (2017).

¹⁷T. Gould, S. Pittalis, J. Toulouse, E. Kraisler, and L. Kronik, "Asymptotic behavior of the hartree-exchange and correlation potentials in ensemble density functional theory," *Phys. Chem. Chem. Phys.* **21**, 19805–19815 (2019).

¹⁸T. Gould, G. Stefanucci, and S. Pittalis, "Ensemble density functional theory: Insight from the fluctuation-dissipation theorem," *Phys. Rev. Lett.* **125**, 233001 (2020).

¹⁹J. G. Brandenburg, K. Burke, E. Fromager, M. Gatti, S. Giarrusso, N. I. Gidopoulos, P. Gori-Giorgi, D. Gowland, T. Helgaker, M. J. P. Hodgson, L. Lacombe, G. Levi, P-F. Loos, N. T. Maitra, E. Maurina Morais, N. Mehta, F. Monti, M. R. Mulay, K. Pernal, L. Reining, P. Romaniello, M. R. Ryder, A. Savin, D. Sirbu, A. M. Teale, A. J. W. Thom, D. G. Truhlar, J. Wetherell, and W. Yang, "New approaches to study excited states in density functional theory: general discussion," *Faraday Discuss.* **224**, 483–508 (2020).

²⁰T. Gould and L. Kronik, "Ensemble generalized kohn–sham theory: The good, the bad, and the ugly," *The Journal of Chemical Physics* **154**, 094125 (2021).

²¹In DFT the KS spin-density need not coincide with the true spin-density.

²²R. T. Sharp and G. K. Horton, "A variational approach to the unipotential many-electron problem," *Phys. Rev.* **90**, 317–317 (1953).

²³J. D. Talman and W. F. Shadwick, "Optimized effective atomic central potential," *Phys. Rev. A* **14**, 36–40 (1976).

²⁴S. Ivanov, S. Hirata, and R. J. Bartlett, "Exact exchange treatment for molecules in finite-basis-set kohn-sham theory," *Phys. Rev. Lett.* **83**, 5455–5458 (1999).

²⁵For information, contact NL at lathiot@eie.gr.

²⁶N. I. Gidopoulos and N. N. Lathiotakis, "Nonanalyticity of the optimized effective potential with finite basis sets," *Phys. Rev. A* **85**, 052508 (2012).

²⁷V. N. Staroverov, G. E. Scuseria, and E. R. Davidson, "Optimized effective potentials yielding hartree–fock energies and densities," *J. Chem. Phys.* **124**, 141103 (2006).

²⁸A. Heßelmann, A. W. Götz, F. Della Sala, and A. Görling, "Numerically stable optimized effective potential method with balanced gaussian basis sets," *J. Chem. Phys.* **127**, 054102 (2007).

²⁹T. J. Callow, "Systematic routes to improved approximations in kohn-sham theory." Available at Durham E-Theses Online: <http://etheses.dur.ac.uk/13817/> (2020).

³⁰T. H. Dunning, "Gaussian basis sets for use in correlated molecular calculations. i. the atoms boron through neon and hydrogen," *J. Chem. Phys.* **90**, 1007–1023 (1989).

³¹D. E. Woon and T. H. Dunning, "Gaussian basis sets for use in correlated molecular calculations. iii. the atoms aluminum through argon,"

- J. Chem. Phys.* **98**, 1358–1371 (1993).
- ³²N. I. Gidopoulos and N. N. Lathiotakis, “Constraining density functional approximations to yield self-interaction free potentials,” *The Journal of Chemical Physics* **136**, 224109 (2012).
- ³³T. J. Callow, N. N. Lathiotakis, and N. I. Gidopoulos, “Density-inversion method for the kohn–sham potential: Role of the screening density,” *J. Chem. Phys.* **152**, 164114 (2020).
- ³⁴A. J. Cohen, P. Mori-Sánchez, and W. Yang, “Insights into current limitations of density functional theory,” *Science* **321**, 792–794 (2008).
- ³⁵J. P. Perdew, K. Burke, and M. Ernzerhof, “Generalized gradient approximation made simple,” *Phys. Rev. Lett.* **77**, 3865–3868 (1996).
- ³⁶A. J. Cohen, P. Mori-Sánchez, and W. Yang, “Fractional spins and static correlation error in density functional theory,” *J. Chem. Phys.* **129**, 121104 (2008).
- ³⁷M. Levy, “Universal variational functionals of electron densities, first-order density matrices, and natural spin-orbitals and solution of the v -representability problem,” *Proceedings of the National Academy of Sciences* **76**, 6062–6065 (1979).
- ³⁸A. Seidl, A. Görling, P. Vogl, J. A. Majewski, and M. Levy, “Generalized kohn–sham schemes and the band-gap problem,” *Phys. Rev. B* **53**, 3764–3774 (1996).
- ³⁹O. Gunnarsson and B. I. Lundqvist, “Exchange and correlation in atoms, molecules, and solids by the spin-density-functional formalism,” *Phys. Rev. B* **13**, 4274–4298 (1976).
- ⁴⁰A. Görling, “Symmetry in density-functional theory,” *Phys. Rev. A* **47**, 2783–2799 (1993).
- ⁴¹A. D. Becke, A. Savin, and H. Stoll, “Extension of the local-spin-density exchange-correlation approximation to multiplet states,” *Theoretica chimica acta* **91**, 147–156 (1995).
- ⁴²Hence SDFT does not have a singular limit at $B = 0$.

