

Double copy structure and the flat space limit of conformal correlators in even dimensions

Arthur E. Lipstein^{*}*Department of Mathematical Sciences, Durham University, Durham DH1 3LE, United Kingdom*Paul McFadden[†]*School of Mathematics, Statistics & Physics, Newcastle University, Newcastle NE1 7RU, United Kingdom*

(Received 24 April 2020; accepted 27 May 2020; published 8 June 2020)

We analyze the flat space limit of 3-point correlators in momentum space for general conformal field theories in even spacetime dimensions, and show they exhibit a double copy structure similar to that found in odd dimensions. In even dimensions, the situation is more complicated because correlators contain branch cuts and divergences which need to be renormalized. We describe the analytic continuation of momenta required to extract the flat space limit, and show that the flat space limit is encoded in the leading singularity of a 1-loop triangle integral which serves as a master integral for 3-point correlators in even dimensions. We then give a detailed analysis of the renormalized correlators in four dimensions where the flat space limit of stress tensor correlators is controlled by the coefficients in the trace anomaly.

DOI: [10.1103/PhysRevD.101.125006](https://doi.org/10.1103/PhysRevD.101.125006)

I. INTRODUCTION

Over the past few decades, the study of scattering amplitudes and conformal correlation functions has revealed remarkable new insight into the structure of quantum field theory and quantum gravity. For example, there is now considerable evidence that scattering amplitudes in quantum gravity can be computed from the correlation functions of a quantum field theory in one lower dimension. This holographic correspondence is best understood when the bulk geometry is anti-de Sitter [1], and conformal field theory (CFT) correlators in the boundary can be computed from Witten diagrams in the bulk. In the flat space limit, these correlators reduce to scattering amplitudes in one higher dimension [2]. Similar methods can also be applied to compute cosmological observables [3–5].

Since scattering amplitudes arise from the flat space limit of correlators, they are far simpler objects and many more tools are available to compute them. It is therefore of great interest to understand how to generalize these tools to correlators, and there has been important progress in this direction. For example, techniques analogous to Britto-Cachazo-Feng-Witten recursion [6] and unitarity methods

[7,8] for scattering amplitudes have been proposed for correlators [9–14]. Another remarkable property of scattering amplitudes is a set of relations connecting gauge to gravitational amplitudes known collectively as the double copy (see [15] for a recent review). Recently, analogous double copy relations were found for general conformal correlators in odd spacetime dimensions [16]. In particular, Euclidean 3-point correlators of stress tensors, conserved currents and marginal scalars were shown to reduce to gauge and gravitational scattering amplitudes in one higher dimension in the flat space limit. This was achieved by working in momentum space and taking the energy (defined as the sum of the magnitudes of the three momenta) to zero. In three dimensions, certain aspects of this double copy structure even extend beyond the flat space limit.

In odd dimensions, 3-point CFT correlators are rational functions of the momentum magnitudes which exhibit poles in the energy. The scattering amplitudes can then be read off from the coefficients of the most singular poles. In even dimensions, the situation is more subtle because the correlators contain branch cuts and need to be analytically continued before taking the flat space limit. Our strategy will be to analyze first the flat space limit of a certain 1-loop triangle integral. All the correlators we consider can then be constructed by applying differential operators to this master integral. (Note these correlators are nonperturbative, being fixed by conformal symmetry.) If the energy of each particle is taken to be positive, the master integral is nonsingular as the total energy tends to zero. To reach the flat space limit, we must instead analytically continue at least one of the energies to be negative before sending their

^{*}arthur.lipstein@durham.ac.uk[†]paul.l.mcfadden@ncl.ac.uk

Published by the American Physical Society under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/). Further distribution of this work must maintain attribution to the author(s) and the published article's title, journal citation, and DOI. Funded by SCOAP³.

sum to zero. This continuation involves crossing certain branch cuts giving rise to a new term with the desired singular behavior in the flat space limit. Interestingly, this new term is precisely the leading singularity of the 1-loop triangle integral computed decades ago by Cutkosky [17]. We present a more modern derivation of this result by first mapping the triangle integral to a box integral with a remarkable property known as dual conformal invariance, and then evaluating the leading singularity of this box integral by taking the global residue.

The flat space limit of correlators in general even dimensions can then be deduced by applying the appropriate differential operators to the master integral, and we discover the same double copy structure that we previously found in odd dimensions. In even dimensions there is one further complication coming from the fact that correlators are divergent and need to be renormalized. This renormalization has been worked out explicitly in four dimensions [18,19], and we carefully verify our general arguments in this case. We also find that the coefficients of the scattering amplitudes which arise in the flat space limit of stress tensor correlators are controlled by conformal anomalies, in agreement with general holographic expectations [20–22].

The structure of this paper is as follows. In Sec. II, we review some basic results about scattering amplitudes and conformal correlators in momentum space that will be relevant for this paper. In Sec. III we derive the flat space limit of the master integral after analytic continuation, and in Sec. IV we use this result to deduce the flat space limit of correlators of stress tensors, currents and marginal scalars in general even dimensions by applying certain differential operators. This reveals double copy structure similar to that previously found in odd dimensions. In Sec. VI, we specialize the discussion to four dimensions where the renormalized correlators have been explicitly computed and we verify the general arguments of the previous section. We also show how the anomaly coefficients parametrize the flat space limit. We present our conclusions and future directions in Sec. VII. In the Appendix we compute the leading singularity of the master integral.

II. FROM CORRELATORS TO AMPLITUDES

In this section, we review some results about momentum-space conformal correlators in d Euclidean dimensions [18,19,23], and their relation to scattering amplitudes in $(d+1)$ -dimensional Minkowski space, which for odd d were worked out in [16]. The tensor structure of correlators is first decomposed into a basis of transverse traceless tensors, where each component is multiplied by a scalar form factor.¹ For 3-point correlators,

¹We restrict here to the parity even sector; for the parity odd sector the basis of transverse traceless tensors is much larger due to the presence of the Levi-Civita tensor; see [23]. Recent work includes [24,25], and in momentum space, [26–28].

these form factors are functions purely of the momentum magnitudes,

$$p_i = +\sqrt{\mathbf{p}_i^2}, \quad i \in \{1, 2, 3\}, \quad (1)$$

since momentum conservation allows us to replace $\mathbf{p}_1 \cdot \mathbf{p}_2 = (p_3^2 - p_1^2 - p_2^2)/2$, etc. For physical kinematics, these magnitudes also obey the triangle inequalities $0 \leq p_i \leq p_j + p_k$.

If desired, the nontransverse traceless parts of correlators can be recovered from lower-point functions via the trace and transverse Ward identities. Here, since our interest is in scattering amplitudes, we will instead contract all indices with transverse polarization vectors $\epsilon_i = \epsilon(\mathbf{p}_i)$ satisfying

$$\epsilon_i \cdot \mathbf{p}_i = 0, \quad \epsilon_i \cdot \epsilon_i = 0. \quad (2)$$

Inserting this tensorial decomposition into the conformal Ward identities, one finds the form factors are given by specific linear combinations of triple- K integrals [23],

$$I_{\alpha\{\beta_1, \beta_2, \beta_3\}}(p_1, p_2, p_3) = \int_0^\infty dx x^\alpha \prod_{i=1}^3 p_i^{\beta_i} K_{\beta_i}(p_i x), \quad (3)$$

where K_{β_i} is a modified Bessel function of the second kind. The parity of the correlators is encoded in the tensor structures that multiply the triple- K integrals.

To connect with scattering amplitudes, we first lift to $(d+1)$ -dimensional Minkowski space by introducing the bulk null momenta and polarization vectors

$$p_i^\mu = (p_i, \mathbf{p}_i), \quad \epsilon_i^\mu = (0, \epsilon_i). \quad (4)$$

For gravitons, we write polarization tensors in terms of polarization vectors as $\epsilon_i^{\mu\nu} = \epsilon_i^\mu \epsilon_i^\nu$. Contractions of polarization vectors can then be lifted to their bulk counterparts by replacing $\epsilon_i \cdot \mathbf{p}_j \rightarrow \epsilon_i \cdot p_j$ and $\epsilon_i \cdot \epsilon_j \rightarrow \epsilon_i \cdot \epsilon_j$. However, while d -dimensional momentum is conserved, the bulk momentum is not since

$$\sum_{i=1}^3 p_i^\mu = (E, \mathbf{0}), \quad (5)$$

where the total bulk energy

$$E = p_1 + p_2 + p_3. \quad (6)$$

We are therefore interested in extracting the leading behavior of CFT correlators in the limit $E \rightarrow 0$ for which energy conservation is restored.

This limit is naturally regarded as a *flat space* limit, either of $(d+1)$ -dimensional anti-de Sitter space [2,11,29], or alternatively of $(d+1)$ -dimensional de Sitter space; see [3,5,16,30] or [31–33]. This follows since the leading

behavior as $E \rightarrow 0$ is governed by the asymptotic behavior of modes deep in the interior of anti-de Sitter (AdS), or equivalently at very early times in de Sitter space, where the effects of spacetime curvature can be neglected.

For CFTs in odd spacetime dimensions, the triple- K integrals feature half-integer indices and the form factors are simple rational functions of the momentum magnitudes. Taking the flat space limit is then simply a matter of extracting the leading behavior as $E \rightarrow 0$. The coefficients of the leading divergences are $(d+1)$ -dimensional flat space scattering amplitudes which exhibit double copy structure. In [16], we found that 3-point correlators of stress tensors and currents reduce to linear combinations of the following gauge and gravitational amplitudes, which are related to each other by a double copy:

$$\mathcal{A}_{EG} = (\mathcal{A}_{YM})^2, \quad (7)$$

$$\mathcal{A}_{\phi R^2}^{222} = \mathcal{A}_{F^3} \mathcal{A}_{YM}, \quad (8)$$

$$\mathcal{A}_{W^3} = (\mathcal{A}_{F^3})^2. \quad (9)$$

Here, \mathcal{A}_{EG} is the 3-graviton scattering amplitude for Einstein gravity, \mathcal{A}_{W^3} is that for Weyl-cubed gravity, while $\mathcal{A}_{\phi R^2}^{222}$ is the 3-graviton amplitude (indicated by the 222 superscript) for the curvature-squared theory of gravity coupled to scalars constructed in [34].² As indicated, these gravitational amplitudes are double copies of the gauge theory amplitudes

$$\begin{aligned} \mathcal{A}_{YM} &= \epsilon_1 \cdot \epsilon_2 \epsilon_3 \cdot p_1 + \text{cyclic}, \\ \mathcal{A}_{F^3} &= \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_3 \epsilon_3 \cdot p_1, \end{aligned} \quad (10)$$

where \mathcal{A}_{YM} is the 3-gluon Yang-Mills amplitude and \mathcal{A}_{F^3} is the corresponding amplitude in a higher-derivative gauge theory with an F^3 interaction constructed in [35]. It is also natural to consider 3-point CFT correlators involving marginal scalars. In [16], we found the correlator of two stress tensors and a marginal scalar reduces in the flat space limit to the amplitude

$$\mathcal{A}_{\phi R^2}^{220} = (\mathcal{A}_{\phi F^2})^2, \quad (11)$$

where

$$\mathcal{A}_{\phi F^2} = \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_1. \quad (12)$$

Here, $\mathcal{A}_{\phi R^2}^{220}$ is the scattering amplitude of two gravitons and a scalar (indicated by the superscript 220) in the ϕR^2

²In $d=4$, this theory reduces to a certain nonminimally coupled version of conformal gravity [35].

theory, which is a double copy of the amplitude $\mathcal{A}_{\phi F^2}$ for two gluons and a scalar in a Yang-Mills dilaton theory.

In even spacetime dimensions, a more detailed analysis is required in order to extract the flat space limit. The two issues are that, firstly, the form factors for CFTs in even dimensions diverge introducing the additional complication of regularization and renormalization; and secondly, the resulting renormalized form factors have a more complicated analytic structure containing branch cuts. As a result, the nature of the analytic continuation required to take the flat space limit $E \rightarrow 0$ must be carefully specified. This is the central question we address in this paper.

III. FLAT SPACE LIMIT OF THE MASTER INTEGRAL

As we will review later in Sec. IV C, for even-dimensional correlators all form factors can be obtained recursively starting from the triple- K integral $I_{1\{000\}}$. Our first task, therefore, is to evaluate the flat space limit of this master integral. We will discuss this from several points of view, but our basic strategy will be to analytically continue the momentum magnitude

$$p_3 = |p_3| e^{i\theta}, \quad 0 \leq \theta \leq \pi, \quad (13)$$

where the momenta are ordered so that p_3 is the largest magnitude. After continuing from $\theta = 0$ to $\theta = \pi$, the flat space limit $E = p_1 + p_2 + p_3 \rightarrow 0$ then corresponds to sending

$$|p_3| \rightarrow p_1 + p_2. \quad (14)$$

Noting that

$$K_\nu(e^{i\pi}x) = e^{-i\pi\nu} K_\nu(x) - i\pi I_\nu(x), \quad x \in \mathbb{R}^+, \quad \nu \in \mathbb{Z}, \quad (15)$$

where I_ν is a modified Bessel function of the first kind, we immediately obtain the following expression for the analytic continuation of $I_{1\{000\}}$:

$$\begin{aligned} I_{1\{000\}}(p_1, p_2, p_3) &= I_{1\{000\}}(p_1, p_2, |p_3|) \\ &\quad - i\pi \int_0^\infty dx x K_0(p_1 x) \\ &\quad \times K_0(p_2 x) I_0(|p_3| x). \end{aligned} \quad (16)$$

The first term on the right-hand side is simply the original triple- K integral and is finite in the flat space limit as we will see shortly. The second term can be evaluated using the formula [36]




FIG. 1. The momenta in the 3-point function form a triangle by momentum conservation, with angle ϕ_i appearing opposite the side of length $|p_i|$.

$$\begin{aligned} & \int_0^\infty dx x^{1+\mu} K_\nu(p_1 x) K_\nu(p_2 x) I_\mu(|p_3| x) \\ &= 2^{-2\mu-2} \sqrt{\frac{\pi}{2}} \Gamma(1+\mu+\nu) \Gamma(1+\mu-\nu) \frac{|c_{123}|^\mu}{\Delta^{2\mu+1}} \\ & \quad \times (\sin \phi_3)^{\mu+1/2} P_{\nu-1/2}^{-\mu-1/2}(\cos \phi_3), \end{aligned} \quad (17)$$

where

$$c_{123} = p_1 p_2 p_3, \quad (18)$$

the P_ν^μ are Legendre functions and Δ is the area of the triangle spanned by the momenta as depicted in Fig. 1. Moreover, using Heron's formula, the area can be written as

$$\Delta = \frac{1}{2} |p_i| |p_j| \sin \phi_k = \sqrt{J^2}/4, \quad (19)$$

where ϕ_i are the angles of the triangle in Fig. 1 and

$$J^2 = E(p_1 + p_2 - p_3)(p_1 - p_2 + p_3)(-p_1 + p_2 + p_3). \quad (20)$$

Notice the value of J^2 is the same at the start and end-point of our analytic continuation. The formula (17) is valid for $1 + \mu - |\nu| > 0$, so choosing $\mu = \nu = 0$ we find

$$I_{1\{000\}}(p_1, p_2, p_3) = I_{1\{000\}}(p_1, p_2, |p_3|) - \frac{i\pi\phi_3}{4\Delta}. \quad (21)$$

Hence, after analytic continuation to $\theta = \pi$, the master integral $I_{1\{000\}}$ acquires a new term which is simply the ratio of the angle ϕ_3 (opposite to the side p_3) to the area of the triangle. In the flat space limit, the angle $\phi_3 \rightarrow \pi$ and the area of the triangle vanishes according to

$$4\Delta = \sqrt{J^2} \rightarrow \sqrt{8|c_{123}|E}. \quad (22)$$

Hence, we find that

$$\lim_{E \rightarrow 0} I_{1\{000\}}(p_1, p_2, p_3) \rightarrow -\frac{i\pi^2}{\sqrt{8|c_{123}|E}} = \frac{\pi^2}{\sqrt{-J^2}}, \quad (23)$$

where the positive sign is taken in the square roots.

Further insight can be obtained by rederiving this result from a different point of view. First, we map the momenta to the complex plane according to

$$u = \frac{p_1^2}{p_3^2} = z\bar{z}, \quad v = \frac{p_2^2}{p_3^2} = (1-z)(1-\bar{z}). \quad (24)$$

Choosing $\Im(z) \geq 0$, we can invert to find

$$z = \frac{1}{2} \left(1 + u - v + \sqrt{(1+u-v)^2 - 4u} \right), \quad (25)$$

$$\bar{z} = \frac{1}{2} \left(1 + u - v - \sqrt{(1+u-v)^2 - 4u} \right). \quad (26)$$

For physical momentum configurations satisfying the triangle inequalities $p_i + p_j \geq p_k$, the quantity under the square root is negative (i.e., $-J^2/p_3^4 \leq 0$) meaning z and \bar{z} are complex conjugates. For such momenta, the master integral $I_{1\{000\}}$ is equivalent to a 1-loop triangle integral [23,37]

$$I_{1\{000\}}(p_1, p_2, p_3) = \frac{1}{4\pi^2} \int \frac{d^4\ell}{\ell^2(\ell+p_1)^2(\ell-p_3)^2}, \quad (27)$$

which can be evaluated in terms of z and \bar{z} as

$$I_{1\{000\}} = \frac{1}{2p_3^2(z-\bar{z})} \left[\text{Li}_2 z - \text{Li}_2 \bar{z} + \frac{1}{2} \ln(z\bar{z}) \ln\left(\frac{1-z}{1-\bar{z}}\right) \right], \quad (28)$$

where Li_2 is the dilogarithm. In fact, $I_{1\{000\}}$ is simply the Bloch-Wigner function [38,39] divided by

$$\sqrt{-J^2} = p_3^2(z-\bar{z}). \quad (29)$$

Geometrically, the Bloch-Wigner function expresses the volume of an ideal tetrahedron (i.e., with vertices at 0, 1, z and ∞) living in the hyperbolic 3-space spanned by the complex z -plane times the real line. Since $\sqrt{-J^2}$ is proportional to the area of the Euclidean triangle in Fig. 1, the master integral $I_{1\{000\}}$ is thus given by the ratio of these quantities.

Recall that J^2 is proportional to E via (20). Naively, we may therefore expect to reach the flat space limit by taking $J^2 \rightarrow 0$, or from (29), taking $z \rightarrow \bar{z}$. On the other hand, the Bloch-Wigner function has the property that all branch cuts in the logarithms and dilogarithms cancel, rendering $I_{1\{000\}}$ single-valued everywhere in the complex plane.³ As a result, we do not obtain a singularity corresponding to a flat space limit since the pole in $z-\bar{z}$ is canceled by the

³Single-valued polylogarithms of higher transcendality have been studied in, e.g., [40–42].




FIG. 2. (a) As we increase θ from 0 to π , z and \bar{z} move clockwise in the complex plane following the solid blue and orange dashed paths, respectively. Starting from generic complex conjugate initial values (corresponding to physical momentum configurations), they ultimately end up exchanging positions. In the process, z crosses the branch cut between $(1, \infty)$ while \bar{z} crosses the branch cut between $(-\infty, 0)$. (b) The trajectory of z and \bar{z} as we continue from a collinear initial configuration to one with $E = 0$. The flat space limit thus corresponds to bringing z and \bar{z} to a point on the real axis between 0 and 1 after crossing the cuts in the direction shown.

vanishing of the numerator in (28) [43]. Physically, the reason for this is that the energy of each particle is positive so taking the limit $z \rightarrow \bar{z}$ corresponds to taking the collinear limit $p_1 + p_2 - p_3 \rightarrow 0$. (Recall we ordered our momenta so $p_1 + p_2 \geq p_3$, hence this specific collinear limit is selected.)

In order to reach the flat space limit $E \rightarrow 0$, we must instead continue one of the energies to become negative, as described in (13), which sends

$$u = |u|e^{-2i\theta}, \quad v = |v|e^{-2i\theta}. \quad (30)$$

As θ ranges from zero to π , the trajectories of z and \bar{z} are then as plotted in Fig. 2. Starting from complex conjugate initial values, for $0 < \theta < \pi$, one finds z and \bar{z} are no longer complex conjugates meaning the branch cuts in the logarithms and dilogarithms no longer cancel. From (28), these cuts are located along the negative real axis, and along the positive real axis for values greater than unity. As we increase θ , we find z crosses the branch cut on the positive real axis, while \bar{z} crosses the branch cut on the negative real axis, both in a clockwise sense. Upon reaching $\theta = \pi$ their values are once again complex conjugates, but their final positions are now exchanged relative to their initial ones. While the exact shape of the trajectory depends on the initial values, the manner in which the respective cuts are crossed is always the same.

As a result of crossing these cuts, we acquire the following new contributions, whose signs are fixed by the direction in which the cuts are crossed:

$$\begin{aligned} \text{Li}_2(z) &\rightarrow \text{Li}_2(z) - 2\pi i \ln z, \\ \ln(1-z) &\rightarrow \ln(1-z) + 2i\pi, \\ \ln \bar{z} &\rightarrow \ln \bar{z} + 2i\pi. \end{aligned} \quad (31)$$

After analytic continuation, we therefore find

$$\begin{aligned} I_{1\{000\}} \Big|_{\theta=\pi} &= I_{1\{000\}} \Big|_{\theta=0} + \frac{1}{2p_3^2(z-\bar{z})} \\ &\times \left[i\pi \left(\ln \left(\frac{\bar{z}}{z} \right) + \ln \left(\frac{1-z}{1-\bar{z}} \right) \right) - 2\pi^2 \right], \end{aligned} \quad (32)$$

where the z and \bar{z} on the right-hand side refer to their final values at $\theta = \pi$. (For the first term however this distinction is immaterial since $I_{1\{000\}} \Big|_{\theta=0}$ is even under exchanging z and \bar{z} .) If we now send $z \rightarrow \bar{z}$ so that they collide on the real axis between zero and unity, without crossing any further cuts, we obtain the leading singular behavior

$$\begin{aligned} \lim_{E \rightarrow 0} I_{1\{000\}} \Big|_{\theta=\pi} &= -\lim_{z \rightarrow \bar{z}} \frac{\pi^2}{p_3^2(z-\bar{z})} \\ &= +\frac{\pi^2}{\sqrt{-J^2}} = -\frac{i\pi^2}{\sqrt{8|c_{123}|E}}. \end{aligned} \quad (33)$$

The positive sign in the penultimate equation follows because z and \bar{z} have exchanged positions relative to their initial values in (29). The value of $\sqrt{-J^2}$ is the same at $\theta = 0$ and $\theta = \pi$, and was fixed as the positive root $\sqrt{-J^2} = +i\sqrt{8|c_{123}|E}$ by our choice $\Im(z) \geq 0$ for $\theta = 0$.

We thus recover the same flat space limit as in (23) above, and the answer will clearly be the same for any analytic continuation so long as the cuts are traversed in the same manner before sending $z \rightarrow \bar{z}$. Had we continued in the opposite sense (i.e., with θ running from zero to $-\pi$) the flat space limit would take the opposite sign, but our real interest here is only in the momentum dependence. Continuing from $\theta = 0$ to π nevertheless seems more natural since this preserves $\Im(z) \geq 0$ for the first part of the trajectory in cases where z and \bar{z} are initially collinear, as illustrated in the right-hand panel of Fig. 2.

Remarkably, the analytic continuation of this same integral has been studied in the context of position-space 4-point functions in [29,44,45], where the general behavior is very similar to that found here. The basis for this

connection is explained in the Appendix, where we relate the 3-point master integral $I_{1\{000\}}$ to a dual conformal box integral. Taking the flat space limit of $I_{1\{000\}}$ then corresponds to computing the leading singularity of this box integral via its global residue. In fact, this connection between the flat space limit and the leading singularity is also visible at the level of the 3-point function. In [17], Cutkosky showed that the leading singularity of the triangle integral (28), obtained by putting all three propagators on shell, has the beautiful geometrical interpretation

$$\int d^4\ell \delta(\ell^2) \delta((\ell + p_1)^2) \delta((\ell - p_3)^2) = \frac{\pi}{8\Delta}, \quad (34)$$

where Δ is the area of the triangle (19). Including the factor of $(2\pi i)^3$ accompanying the delta functions, and the factor of $1/4\pi^2$ in (27), we again recover precisely (23).

IV. EVALUATING THE FLAT SPACE LIMIT OF CORRELATORS

Having analyzed the master integral, let us now discuss how to evaluate the flat space limit of correlators for general even dimensions $d = 2n \geq 4$. The relevant correlators for the double copy are $\langle JJJ \rangle$ and $\langle TTT \rangle$, as well as $\langle JJO \rangle$ and $\langle TTO \rangle$ for a marginal operator \mathcal{O} . As for the master integral, a number of different approaches can be taken. We discuss these in each of the following three subsections. While all approaches give the same result, they present different features of interest.

A. Asymptotic analysis

First we present a simple asymptotic formula for the flat space limit of a general triple- K integral. We start by analytically continuing the general triple- K integral (3). Using (13) and (15), we obtain

$$\begin{aligned} & I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1, p_2, p_3) \Big|_{\theta=\pi} \\ &= I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1, p_2, |p_3|) - i\pi p_1^{\beta_1} p_2^{\beta_2} p_3^{\beta_3} \\ & \quad \times \int_0^\infty dx x^\alpha K_{\beta_1}(p_1 x) K_{\beta_2}(p_2 x) I_{\beta_3}(|p_3| x), \end{aligned} \quad (35)$$

where the phase $e^{-i\pi\beta_3}$ from the continuation of the Bessel function cancels with that from the continuation of $p_3^{\beta_3}$. To evaluate the flat space limit of this analytically continued integral, we now consider the asymptotic behavior of its integrand. Physically, the flat space limit is reached by going to $p_i x \gg 1$, which corresponds to the deep interior of the bulk spacetime. Replacing the Bessel functions with their asymptotic behaviors,

$$K_\beta(p_i x) \rightarrow \sqrt{\frac{\pi}{2p_i x}} e^{-p_i x}, \quad I_\beta(p_i x) \rightarrow \sqrt{\frac{1}{2\pi p_i x}} e^{p_i x}, \quad (36)$$

we obtain the flat space limit

$$\begin{aligned} & \lim_{E \rightarrow 0} I_{\alpha\{\beta_1\beta_2\beta_3\}}(p_1, p_2, p_3) \Big|_{\theta=\pi} \\ &= -\frac{\pi^{3/2} \Gamma(\alpha - 1/2) \prod_{i=1}^3 p_i^{\beta_i - 1/2}}{\sqrt{8} E^{\alpha - 1/2}}. \end{aligned} \quad (37)$$

Note this result derives entirely from the KKI integral in (35), since the triple- K integral is finite for collinear configurations [23]. This asymptotic formula agrees with our result (23) for the flat space limit of $I_{1\{000\}}$, as well as the results in [16] for odd spacetime dimensions.

In even dimensions, however, we encounter divergent triple- K integrals and hence we must take into account the effects of regularization and renormalization. In general, a triple- K integral diverges whenever the indices satisfy [46]

$$\alpha + 1 \pm \beta_1 \pm \beta_2 \pm \beta_3 = -2n \quad (38)$$

for any (independent) choice of \pm signs and non-negative integer n . To regulate, one performs infinitesimal shifts of the operator and spacetime dimensions, and thus of the indices $\alpha, \{\beta_i\}$ parametrizing the triple- K integrals. The divergences can then be extracted and eliminated by the addition of covariant local counterterms, before removing the regulator to obtain the renormalized correlators.

For the correlators of interest here, one finds from the detailed analysis of [18,19] that the regulated form factors contain only *ultralocal* divergences, meaning they are analytic functions of the squared momenta.⁴ Terms of this form, and the corresponding counterterm contributions, cannot contribute any singular behavior in the flat space limit: this would require the appearance of factors of E raised to negative powers, which are not ultralocal. It therefore suffices to apply the continuation (35) to the regulated form factors and extract the leading behavior as $E \rightarrow 0$ using the asymptotic formula (37). The result is necessarily finite as the regulator is removed and all dimensions are restored to their physical values.⁵

B. Analytic continuation of the renormalized form factors in $d = 4$

Where the renormalized form factors are known explicitly, we can alternatively apply the analytic continuation

⁴Such singularities correspond to triple- K integrals with sign choice $(---)$ in (38). In fact, for $\langle TTT \rangle$, $\langle JJO \rangle$ and $\langle TTO \rangle$, one also encounters individual triple- K integrals with semilocal $(--+)$ or $(+-)$ divergences, however these either cancel with one another, or else are multiplied by vanishing coefficients. The regulated form factors then contain only ultralocal $(---)$ divergences. This is consistent with the absence of $(--+)$ or $(+-)$ type counterterms for these correlators.

⁵From the analysis of [46], the KKI integral in (35) is singular only when the sign of β_3 in (38) is $+$. However, as above, all singularities of this type either cancel or are multiplied by vanishing coefficients.

(13) directly to the renormalized form factors. In this approach we never encounter any divergences since we always work within the renormalized theory.

To illustrate this we consider the case $d = 4$, where all renormalized form factors have been evaluated in terms of differential operators acting on the finite master integral $I_{1\{000\}}$ [18,19]. These derivatives can be evaluated by making repeated use of the relation

$$p_1 \frac{\partial}{\partial p_1} I_{1\{000\}} = \frac{1}{J^2} \left[2p_1^2(p_1^2 - p_2^2 - p_3^2) I_{1\{000\}} - p_1^2 \ln p_1^2 + \frac{1}{2}(p_1^2 + p_2^2 - p_3^2) \ln p_2^2 + \frac{1}{2}(p_1^2 - p_2^2 + p_3^2) \ln p_3^2 \right]. \quad (39)$$

Ultimately, one finds that all renormalized form factors are given by a linear combination of the master integral $I_{1\{000\}}$ multiplied by some rational function of the squared momenta and $\sqrt{-J^2}$, plus logarithms of the momenta and renormalization group scale multiplied by similar rational functions, plus polynomials in the squared momenta. As we now discuss, the analytic continuation of all these terms is easily accomplished using the results of Sec. III.

The situation is clearest in the (z, \bar{z}) variables, where all factors of $\sqrt{-J^2}$ rationalize according to (29). The renormalized form factors A_n then take the form

$$p_3^\ell A_n = a_n^{(0)} p_3^2 I_{1\{000\}} + a_n^{(1)} \ln(z\bar{z}) + a_n^{(2)} \ln((1-z)(1-\bar{z})) + a_n^{(3)} \ln \frac{p_3^2}{\mu^2} + a_n^{(4)}, \quad (40)$$

where the factor of p_3^ℓ renders A_n dimensionless, and the $a_n^{(m)}$ are specific rational functions of z and \bar{z} . In particular, these rational functions may diverge as $\bar{z} \rightarrow z$. Although we acquire new terms from analytically continuing the logs, the leading behavior in the flat space limit in fact always comes from the continuation of the master integral $I_{1\{000\}}$. This can be understood as follows. The key point is that the renormalized form factors are finite in the collinear limit,⁶ which corresponds to sending $\bar{z} \rightarrow z$ while keeping $\theta = 0$. Since in the collinear limit [43]

$$\lim_{\bar{z} \rightarrow z} I_{1\{000\}} \Big|_{\theta=0} = -\frac{1}{2p_3^2} \left(\frac{\ln z}{1-z} + \frac{\ln(1-z)}{z} \right), \quad (41)$$

⁶In the regulated theory, the form factors are linear combinations of triple- K integrals, and triple- K integrals do not have collinear singularities [23]. The counterterm contributions are ultralocal and thus do not have collinear singularities either. The renormalized form factors are then finite for collinear configurations, as can be checked explicitly using the results of [18,19].

we see that if $a_n^{(0)}$ diverges as $(z - \bar{z})^{-k}$ for some k , then $a_n^{(1)}$ and $a_n^{(2)}$ must also diverge at this same order, so that

$$\begin{aligned} \chi_1 &= \lim_{\bar{z} \rightarrow z} \left(2a_n^{(1)} - \frac{1}{2(1-z)} a_n^{(0)} \right), \\ \chi_2 &= \lim_{\bar{z} \rightarrow z} \left(2a_n^{(2)} - \frac{1}{2z} a_n^{(0)} \right) \end{aligned} \quad (42)$$

are both finite. The remaining rational functions $a_n^{(3)}$ and $a_n^{(4)}$ are both subleading: $a_n^{(3)}$ is finite in the collinear limit since no cancellations are possible for this term, while any collinear divergences in $a_n^{(4)}$ are necessarily of order $(z - \bar{z})^{-k+2}$, since they must cancel against the polynomial terms that arise in $I_{1\{000\}}$ at subleading order [i.e., at order $(z - \bar{z})^2$ relative to the leading term shown in (41)].

After we analytically continue to $\theta = \pi$, the coefficient $a_n^{(0)}$ now acquires an additional factor of $(z - \bar{z})^{-1}$ from analytically continuing the master integral according to (33). The continuations of the log terms in (40) do not produce any additional divergences, however, and so overall the leading $(z - \bar{z})^{-k+1}$ behavior of the form factor is that associated with the master integral. Thus, to find the leading behavior of the renormalized form factors in the flat space limit, we actually only need to know the coefficient $a_n^{(0)}$ of the master integral. This is a substantial simplification.

Finally, let us remark that our approach is also readily applicable to the problem of continuing Euclidean CFT correlators to Lorentzian signature, as studied in [47–49]. One simply needs to analyze how z and \bar{z} move in the complex plane under Wick rotation, then evaluate the corresponding continuation of the master integral as discussed in Sec. III.

C. Extracting the dependence on the master integral in general dimensions

For general even dimensions above four, results are available for the regulated form factors, and the nature of all divergences and counterterms have been tabulated [18,19]. Once again, all triple- K integrals can be computed starting from the master integral $I_{1\{000\}}$ using the reduction scheme of [37]. To obtain the final renormalized form factors then requires a certain amount of additional case-by-case analysis of counterterm contributions. While this analysis is easy to perform in any specific case, it is difficult to write down general closed-form expressions. From the $d = 4$ discussion above, however, all we really need to know is the contribution to the renormalized form factors coming from the master integral $I_{1\{000\}}$, since this is the term that dominates in the flat space limit. This contribution is easily evaluated as we now explain.

TABLE I. Index-shifting operations generated by cyclic permutations of (43) (operations **1**, **2** and **3**), along with (44) and (45) (operations **4** and **5**). Through repeated use of these operations, all the regulated triple- K integrals in even-dimensional correlators can be reduced to $I_{0\{111\}}$.

| Operation | $\delta\alpha$ | $\delta\beta_1$ | $\delta\beta_2$ | $\delta\beta_3$ |
|-----------|----------------|-----------------|-----------------|-----------------|
| 1 | 1 | 1 | 0 | 0 |
| 2 | 1 | 0 | 1 | 0 |
| 3 | 1 | 0 | 0 | 1 |
| 4 | 2 | 0 | 0 | 0 |
| 5 | 1 | 1 | 1 | 1 |

Working in the regulated theory to avoid divergences, all triple- K integrals are first reduced to the integral $I_{0\{111\}}$. This can be achieved using the relations [37]

$$I_{\alpha+1\{\beta_1+1,\beta_2,\beta_3\}} = (2\beta_1 - p_1 \partial_{p_1}) I_{\alpha\{\beta_1\beta_2\beta_3\}}, \quad (43)$$

$$I_{\alpha+2\{\beta_1,\beta_2,\beta_3\}} = \left(\partial_{p_j}^2 + \frac{1-2\beta_j}{p_j} \partial_{p_j} \right) I_{\alpha\{\beta_1\beta_2\beta_3\}}, \quad (44)$$

$$I_{\alpha+1\{\beta_1+1,\beta_2+1,\beta_3+1\}} = \frac{1}{\alpha - \beta_t - 1} B_{\beta_1,\beta_2,\beta_3} I_{\alpha\{\beta_1\beta_2\beta_3\}}, \quad (45)$$

where in (44) one can choose any p_j from $j = 1, 2, 3$ and the operator in (45) is

$$B_{\beta_1,\beta_2,\beta_3} = p_1^2(2\beta_2 - p_2 \partial_{p_2})(2\beta_3 - p_3 \partial_{p_3}) + \text{cyclic}. \quad (46)$$

From these relations, we can construct the five index-shifting operations listed in Table I. The first three operations in this table follow from cyclic permutations of (43), while the fourth and fifth operation are (44) and (45), respectively.

From (4.2)–(4.7) of [37], the form of the dimensionally regulated integral $I_{0\{111\}}$ is

$$\begin{aligned} I_{0+ue\{1+v_1\epsilon, 1+v_2\epsilon, 1+v_3\epsilon\}} &= \frac{I^{(-2)}}{\epsilon^2} + \frac{I^{(-1)}}{\epsilon} + I^{(\text{scheme})} \\ &+ I^{(\text{scale-violating})} + I^{(\text{nonlocal})} \\ &+ O(\epsilon), \end{aligned} \quad (47)$$

where the indices have been shifted by an infinitesimal parameter ϵ times scheme-dependent constants (u, v_1, v_2, v_3). The divergent terms $I^{(-2)}$ and $I^{(-1)}$ are ultralocal and semilocal, respectively, and will ultimately be removed by subtracting counterterm contributions. The scheme-dependent term $I^{(\text{scheme})}$ contains logarithms of the individual momentum magnitudes and hence is semilocal, while the scale-violating piece $I^{(\text{scale-violating})}$ contains products of such logarithms and is nonlocal. The final scale-invariant, nonlocal piece $I^{(\text{nonlocal})}$ (also referred to as

$I_{0\{111\}}^{(\text{fin})}$ in [18,19]) encodes the dependence on the master integral we seek:

$$I^{(\text{nonlocal})} = \frac{J^2}{4} I_{1\{000\}}. \quad (48)$$

As in our discussion for $d = 4$, the finiteness of the renormalized form factors in the collinear limit ensures the leading contribution in the flat space limit comes solely from $I^{(\text{nonlocal})}$. Unlike $I_{1\{000\}}$, neither $I^{(\text{scheme})}$ or $I^{(\text{scale-violating})}$ acquire any additional divergences as $\bar{z} \rightarrow z$ after continuing to $\theta = \pi$. Since for $\theta = 0$ any divergences as $\bar{z} \rightarrow z$ must cancel, the leading divergence as $\bar{z} \rightarrow z$ for $\theta = \pi$ must then come from $I_{1\{000\}}$. For this reason, we can simply replace

$$I_{0\{111\}} \rightarrow \frac{J^2}{4} I_{1\{000\}} \quad (49)$$

for the purposes of computing the flat space limit of the renormalized form factors. All details of the regularization scheme and renormalization analysis can be safely neglected, since their contribution is subleading in the flat space limit.

In summary, the known expressions for the regulated form factors can be related to $I_{0\{111\}}$ using the operations (43)–(45) summarized in Table I, after which we substitute (49). The leading behavior in the flat space limit then corresponds to applying the same sequence of differential operators to the leading flat space behavior of $(J^2/4)I_{1\{000\}}$, evaluated using (23). In fact, one only needs to keep track of the leading contributions when evaluating these derivatives, which enables further simplification. For example, the flat space limit of $I_{2\{111\}}$ can be obtained by applying operation **5** in Table I, which simplifies to

$$\lim_{E \rightarrow 0} I_{2\{111\}} \propto (p_1^2 p_2 p_3 + \text{cyclic}) \partial_E^2 \frac{1}{\sqrt{c_{123}} E} \propto \frac{\sqrt{c_{123}}}{E^{3/2}}. \quad (50)$$

Here, we retained only the terms in (46) featuring derivatives, since their action is to generate more singular powers of E . For this same reason, the factor of $\sqrt{c_{123}}$ from the flat space limit of $I_{1\{000\}}$ can be moved outside the derivatives, which can then be replaced by derivatives with respect to E using the chain rule. Using this method, we find that the triple- K integrals we encounter all exhibit flat space behavior in agreement with the asymptotic formula (37).

V. GENERAL EVEN DIMENSIONS

In this section, we apply the procedure of Sec. IV C to compute the flat space limit of 3-point correlators of stress tensors, currents and marginal scalar operators, which we




FIG. 3. Reduction scheme for the regulated triple- K integrals appearing in $\langle JJJ \rangle$, where the numbered operations refer to Table I, and 5^{n-1} means applying operation 5 a total of $n-1$ times.

denote T , J , and \mathcal{O} , respectively. We show that correlators reduce to flat space scattering amplitudes in one higher dimension related by a double copy, extending the results of [16] to even dimensions. We take the spacetime dimension to be $d = 2n > 4$, postponing the analysis of $d = 4$ (where more detailed results for the renormalized form factors are available) to the following section.

A. $\langle JJJ \rangle$

First, we consider the 3-point correlator of conserved currents. This can be decomposed into form factors as follows:

$$\langle JJJ \rangle = A_1(p_1, p_2, p_3)\epsilon_1 \cdot p_2 \epsilon_2 \cdot p_3 \epsilon_3 \cdot p_1 + [A_2(p_1, p_2, p_3)\epsilon_1 \cdot \epsilon_2 \epsilon_3 \cdot p_1 + \text{cyclic}]. \quad (51)$$

Here, $\langle JJJ \rangle$ represents the correlator fully contracted with polarization vectors, and with color factors suppressed. We also strip off the overall delta function associated with momentum conservation. (Such correlators are denoted $\langle\langle \dots \rangle\rangle$ in [18,19].) From the conformal Ward identities, one finds [18]

$$\begin{aligned} A_1 &= C_1 I_{n+2\{n-1,n-1,n-1\}}, \\ A_2 &= C_1 I_{n+1\{n-1,n-1,n\}} + C_2 I_{n\{n-1,n-1,n-1\}}, \end{aligned} \quad (52)$$

where C_1 and C_2 are constants, and

$$C_2 = \#C_1 + \#C_{JJ}, \quad (53)$$

where C_{JJ} is the normalization of the 2-point function. The $\#$ represent specific dimension-dependent constants whose precise form is not important for reasons explained below.

The reduction to $I_{0\{111\}}$ and then to $I_{1\{000\}}$ via (49) is depicted in Fig. 3, where circled numbers correspond to operations in Table I. Applying these operations to (23) and using (53), we can read off the flat space limit,

$$\begin{aligned} \lim_{E \rightarrow 0} \langle JJJ \rangle &\propto c_{123}^{(d-3)/2} \left[\frac{C_1}{E^{(d+3)/2}} (\mathcal{A}_{F^3} + \mathcal{O}(E)) \right. \\ &\quad \left. + \frac{C_{JJ}}{E^{(d-1)/2}} (\mathcal{A}_{\text{YM}} + \mathcal{O}(E)) \right], \end{aligned} \quad (54)$$

where the gauge theory scattering amplitudes \mathcal{A}_{F^3} and \mathcal{A}_{YM} are given in (10). In this calculation, the contribution from

the first term in (53) is subleading, while the $\#$ in the second term proportional to C_{JJ} has been absorbed into the overall constant of proportionality in (54). Since C_1 is an arbitrary constant, we have additionally rescaled C_1 to eliminate any relative factors between the two terms in (54).

B. $\langle TTT \rangle$

The form factor decomposition for the stress tensor 3-point function is

$$\begin{aligned} \langle TTT \rangle &= A_1(p_1, p_2, p_3)(\epsilon_1 \cdot p_2 \epsilon_2 \cdot p_3 \epsilon_3 \cdot p_1)^2 \\ &\quad + (A_2(p_1, p_2, p_3)\epsilon_1 \cdot \epsilon_2 \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_3 (\epsilon_3 \cdot p_1)^2 + \text{cyclic}) \\ &\quad + (A_3(p_1, p_2, p_3)(\epsilon_1 \cdot \epsilon_2)^2 (p_1 \cdot \epsilon_3)^2 + \text{cyclic}) \\ &\quad + (A_4(p_1, p_2, p_3)\epsilon_1 \cdot \epsilon_3 \epsilon_2 \cdot \epsilon_3 \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_3 + \text{cyclic}) \\ &\quad + A_5(p_1, p_2, p_3)\epsilon_1 \cdot \epsilon_2 \epsilon_2 \cdot \epsilon_3 \epsilon_3 \cdot \epsilon_1, \end{aligned} \quad (55)$$

where $\langle TTT \rangle$ represents the correlator fully contracted with polarization vectors and with the delta function of momentum conservation stripped off. The form factors are [18]

$$\begin{aligned} A_1 &= C_1 I_{5+n\{n,n,n\}}, \\ A_2 &= 4C_1 I_{4+n\{n,n,n+1\}} + C_2 I_{3+n\{n,n,n\}}, \\ A_3 &= 2C_1 I_{3+n\{n,n,n+2\}} + C_2 I_{2+n\{n,n,n+1\}} + C_3 I_{n+1\{n,n,n\}}, \\ A_4 &= 8C_1 I_{3+n\{n+1,n+1,n\}} - 2C_2 I_{2+n\{n,n,n+1\}} \\ &\quad + C_4 I_{n+1\{n,n,n\}}, \\ A_5 &= 8C_1 I_{n+2\{n+1,n+1,n+1\}} + 2C_2 [I_{n+1\{n+1,n+1,n\}} \\ &\quad + I_{n+1\{n+1,n,n+1\}} + I_{n+1\{n,n+1,n+1\}}] \\ &\quad + C_5 I_{n-1\{n,n,n\}}, \end{aligned} \quad (56)$$

where C_1 to C_5 are constants (independent of those introduced in the previous subsection) which are related by

$$C_4 = 2C_3 + \#C_2, \quad C_5 = \#C_1 + \#C_2 + \#C_3. \quad (57)$$

Here, the $\#$ are specific dimension-dependent constants whose form is not of interest since the corresponding terms are subleading in the flat space limit. The factor of two in the first equation is however important in order to recover the Einstein gravity amplitude. We can likewise replace C_3 in terms of the normalization C_{TT} of the stress tensor 2-point function,

$$C_3 = \#C_1 + \#C_2 + \#C_{TT}, \quad (58)$$

where the terms proportional to C_1 and C_2 are also subleading in the flat space limit.

The reduction of the triple- K integrals to $I_{0\{111\}}$ is depicted in Fig. 4. Using (49), the flat space limit of these




FIG. 4. Reduction scheme for the regulated triple- K integrals appearing in $\langle TTT \rangle$, where the numbered operations refer to those in Table I.

integrals can then be deduced from that of $I_{1\{000\}}$. After rescaling the (theory-specific) constants C_1 and C_2 to absorb dimension-dependent constants, we obtain the flat space limit

$$\begin{aligned} \lim_{E \rightarrow 0} \langle TTT \rangle \propto c_{123}^{(d-1)/2} & \left[\frac{C_1}{E^{(d+9)/2}} (\mathcal{A}_{W^3} + \mathcal{O}(E)) \right. \\ & + \frac{C_2}{E^{(d+5)/2}} (\mathcal{A}_{\phi R^2}^{222} + \mathcal{O}(E)) \\ & \left. + \frac{C_{TT}}{E^{(d+1)/2}} (\mathcal{A}_{EG} + \mathcal{O}(E)) \right]. \end{aligned} \quad (59)$$

Remarkably, the gravitational amplitudes arising in the flat space limit of $\langle TTT \rangle$ are double copies of the gauge theory amplitudes arising in the flat space limit of $\langle JJJ \rangle$, as given in (9). This result takes the same form as in odd dimensions [16], as one would expect from the dimension-independent nature of the amplitudes themselves.

C. $\langle JJO \rangle$

For the case of two currents and a marginal scalar, the form factor decomposition is

$$\begin{aligned} \langle JJO \rangle = & -A_1(p_1, p_2, p_3) \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_1 \\ & + A_2(p_1, p_2, p_3) \epsilon_1 \cdot \epsilon_2. \end{aligned} \quad (60)$$

The form factors are [19]

$$\begin{aligned} A_1 = & C_1 I_{n+1\{n-1, n-1, n\}}, \\ A_2 = & C_1 I_{n\{n-1, n-1, n+1\}} + C_2 I_{n-1\{n-1, n-1, n\}}, \end{aligned} \quad (61)$$

where the constants C_1 and C_2 (which are once again independent from those defined in previous subsections) satisfy

$$C_2 = \#C_1. \quad (62)$$

Using the reduction to $I_{1\{000\}}$ in Fig. 5, we then obtain

$$\lim_{E \rightarrow 0} \langle JJO \rangle \propto c_{123}^{(d-3)/2} p_3 \frac{C_1}{E^{(d+1)/2}} (\mathcal{A}_{\phi F^2} + \mathcal{O}(E)), \quad (63)$$




FIG. 5. Reduction scheme for the regulated triple- K integrals appearing in $\langle JJO \rangle$.

where $\mathcal{A}_{\phi F^2}$ is the amplitude for two gluons and a scalar in (12).

D. $\langle TTO \rangle$

Finally, in the case of two stress tensors and a marginal scalar, we have

$$\begin{aligned} \langle TTO \rangle = & A_1(p_1, p_2, p_3) (\epsilon_1 \cdot p_2 \epsilon_2 \cdot p_1)^2 \\ & - A_2(p_1, p_2, p_3) \epsilon_1 \cdot \epsilon_2 \epsilon_1 \cdot p_2 \epsilon_2 \cdot p_1 \\ & + A_3(p_1, p_2, p_3) (\epsilon_1 \cdot \epsilon_2)^2, \end{aligned} \quad (64)$$

where the form factors are [19]

$$\begin{aligned} A_1 = & C_1 I_{3+n\{n, n, n\}}, \\ A_2 = & 4C_1 I_{2+n\{n, n, n+1\}} + C_3 I_{n+1\{n, n, n\}}, \\ A_3 = & 2C_1 I_{n+1\{n, n, n+2\}} + C_2 I_{n\{n, n, n+1\}} \\ & + C_3 I_{n-1\{n, n, n\}}, \end{aligned} \quad (65)$$

and the constants satisfy

$$C_2 = \#C_1, \quad C_3 = \#C_1. \quad (66)$$

Using the triple- K reduction in Fig. 6, the flat space limit is given by

$$\lim_{E \rightarrow 0} \langle TTO \rangle \propto c_{123}^{(d-1)/2} \frac{C_1}{E^{(d+5)/2}} (\mathcal{A}_{\phi R^2}^{220} + \mathcal{O}(E)), \quad (67)$$

where the amplitude $\mathcal{A}_{\phi R^2}^{220}$ for two gravitons and a scalar is a double copy of that for two gluons and a scalar arising in the flat space limit of $\langle JJO \rangle$, as given in (11).




FIG. 6. Reduction scheme for the regulated triple- K integrals appearing in $\langle TTO \rangle$. All numbered operations refer to those in Table I.

Once again, we find the same double copy structure for even-dimensional correlators as that obtained in [16] for odd-dimensional correlators.

VI. FOUR DIMENSIONS

We present the case of $d = 4$ separately since the complete renormalization analysis has been carried out in [18,19]. For $\langle TTT \rangle$ this allows us to parametrize the flat space limit in terms of the trace anomaly coefficients. We also briefly discuss the double copy structure for anomalies.

A. $\langle JJJ \rangle$

In $d = 4$, the renormalized form factors for $\langle JJJ \rangle$ are given by [18]

$$\begin{aligned}
A_1 &= C_1 I_{7\{222\}} + \dots, \\
A_2 &= 2 \left[a + c - 2C_1 p_3 \frac{\partial}{\partial p_3} \right] I_{5\{222\}}^{(\text{fin})} + \dots, \\
A_3 &= 2 \left[2c - (a + c + C_1) p_3 \frac{\partial}{\partial p_3} + C_1 p_3^2 \frac{\partial^2}{\partial p_3^2} \right] I_{3\{222\}}^{(\text{fin})} + \dots, \\
A_4 &= 4 \left[c - a + (a + c) p_3 \frac{\partial}{\partial p_3} + 2C_1 \left(8 - 4 \sum_{j=1}^3 p_j \frac{\partial}{\partial p_j} + p_1 p_2 \frac{\partial^2}{\partial p_1 \partial p_2} \right) \right] I_{3\{222\}}^{(\text{fin})} + \dots, \\
A_5 &= 2(a + c) \left[32 - 8 \sum_{j=1}^3 p_j \frac{\partial}{\partial p_j} + 2 \sum_{i < j} p_i p_j \frac{\partial^2}{\partial p_i \partial p_j} \right] I_{1\{222\}}^{(\text{fin})} - 8C_1 p_1^3 p_2^3 p_3^3 \frac{\partial^3}{\partial p_1 \partial p_2 \partial p_3} I_{1\{000\}} + \dots, \quad (69)
\end{aligned}$$

where once again the ellipses denote terms that are nonsingular in the flat space limit. Here, the triple- K integral $I_{7\{222\}}$ and the finite integrals $I_{5\{222\}}^{(\text{fin})}$, $I_{3\{222\}}^{(\text{fin})}$ and $I_{1\{222\}}^{(\text{fin})}$ can all be expressed as derivatives of the master integral, see (3.198)–(3.201) of [18]. The coefficients a and c are those entering the trace anomaly,

$$\langle T_\mu^\mu \rangle = aE_4 + cW^2, \quad (70)$$

where E_4 is the Euler density and W^2 is the square of the four-dimensional Weyl tensor. (Note that the Euler coefficient a is often defined with an additional minus sign to here.) To ensure the 2-point function is traceless, we work in a scheme where the $\square R$ contribution to the trace anomaly vanishes.

Using (23) and (43)–(49), we recover the flat space limit in (59) with⁷

$$C_2 \rightarrow a + c, \quad C_{TT} \rightarrow c. \quad (71)$$

⁷The exact relation from [18] is $C_{TT} = -2c$, but here as in (59) we omit such numerical coefficients.

$$\begin{aligned}
A_1 &= -C_1 p_1 p_2 p_3 \frac{\partial^3}{\partial p_1 \partial p_2 \partial p_3} I_{1\{000\}} + \dots, \\
A_2 &= C_1 p_1 p_2 p_3^2 \frac{\partial^2}{\partial p_1 \partial p_2} I_{1\{000\}} + 4C_{JJ} I_{2\{111\}}^{(\text{fin})} + \dots, \quad (68)
\end{aligned}$$

where the ellipses denote terms which are nonsingular in the flat space limit. We have additionally suppressed a factor relating to the color and the charge multiplying the 2-point normalization C_{JJ} . The finite integral $I_{2\{111\}}^{(\text{fin})}$ is given in (3.48) of [18]. Using (23) and (43)–(49), we recover the flat space limit in (54).

B. $\langle TTT \rangle$

The renormalized form factors for $\langle TTT \rangle$ are [18]

We can equivalently write this as

$$\lim_{E \rightarrow 0} \frac{E^{13/2}}{c_{123}^{3/2}} \langle TTT \rangle \propto C_1 \mathcal{A}_{W^3}, \quad (72)$$

$$\lim_{E \rightarrow 0} \frac{E^{9/2}}{c_{123}^{3/2}} \langle TTT \rangle|_{C_1=0} \propto (a + c) \mathcal{A}_{\phi R^2}^{222}, \quad (73)$$

$$\lim_{E \rightarrow 0} \frac{E^{5/2}}{c_{123}^{3/2}} \langle TTT \rangle|_{C_1=0, a+c=0} \propto c \mathcal{A}_{EG}. \quad (74)$$

The coefficients of the ϕR^2 and Einstein gravity amplitudes arising in the flat space limit are thus parametrized by the trace anomaly. This is natural from a holographic perspective since these anomaly coefficients are determined by bulk gravitational interactions [20–22]. Hence for Einstein gravity, for example, $a + c = 0$ and the $\mathcal{A}_{\phi R^2}^{222}$ contribution vanishes. Note that our results do not rely on holography and therefore also apply to conformal field theories without a holographic dual.

Finally, while our focus is on the transverse traceless parts of correlators, we note that double copy structure also arises in the trace part of this correlator as shown in [18]. From the trace Ward identity, this takes the form

$$\begin{aligned} & \langle T_\mu^\mu(\mathbf{p}_1)T(\mathbf{p}_2)T(\mathbf{p}_3) \rangle \\ & = 2\langle T(\mathbf{p}_2)T(-\mathbf{p}_2) \rangle + 2\langle T(\mathbf{p}_3)T(-\mathbf{p}_3) \rangle + \mathcal{A}, \end{aligned} \quad (75)$$

where we trace over the first two indices and contract the rest with polarization tensors. The anomalous contribution \mathcal{A} comes from functionally differentiating the trace anomaly (70). The part proportional to the Euler anomaly coefficient a is then a double copy of the chiral anomaly:

$$\begin{aligned} \mathcal{A}_{\text{Euler}} & = 40a\mathcal{A}_{\text{chiral}}^2, \\ \mathcal{A}_{\text{chiral}} & = \epsilon_{\mu_2\mu_3\mu_4\mu_5}\epsilon_2^{\mu_2}\epsilon_3^{\mu_3}p_2^{\mu_4}p_3^{\mu_5}, \end{aligned} \quad (76)$$

where the chiral anomaly arises in the transverse Ward identity for currents,

$$\begin{aligned} & \langle (p_1 \cdot J^a(\mathbf{p}_1))J^b(\mathbf{p}_2)J^c(\mathbf{p}_3) \rangle \\ & = gf^{adc}\langle J^d(\mathbf{p}_2)J^b(-\mathbf{p}_2) \rangle - gf^{abd}\langle J^d(\mathbf{p}_3)J^c(-\mathbf{p}_3) \rangle \\ & \quad + d^{abc}\mathcal{A}_{\text{chiral}}. \end{aligned} \quad (77)$$

In this identity, f^{abc} is the structure constant, g the gauge coupling, and d^{abc} is a group-theoretic factor depending on the matter content. The double copy (76) derives from the specific structure of type A anomalies (in the classification of [50]), and is not present for type B anomalies such as the Weyl-squared contribution to the trace anomaly.

C. $\langle JJO \rangle$

The renormalized form factors are given by [19]

$$\begin{aligned} A_1 & = C_1 \left(2 - p_3 \frac{\partial}{\partial p_3} \right) I_{2\{111\}}^{(\text{fin})} + \dots, \\ A_2 & = C_1 p_3^2 I_{2\{111\}}^{(\text{fin})} + \dots, \end{aligned} \quad (78)$$

where the omitted terms are nonsingular in the flat space limit. Using (23) and (43)–(49), we recover the flat space limit in (63).

D. $\langle TTO \rangle$

The renormalized form factors are given by [19]

$$\begin{aligned} A_1 & = C_1 \left(2 - p_1 \frac{\partial}{\partial p_1} \right) \left(2 - p_2 \frac{\partial}{\partial p_2} \right) \left(2 - p_3 \frac{\partial}{\partial p_3} \right) I_{2\{111\}}^{(\text{fin})} \\ & \quad + \dots, \\ A_2 & = 4C_1 \left(1 - p_3 \frac{\partial}{\partial p_3} \right) I_{3\{222\}}^{(\text{fin})} + \dots, \\ A_3 & = 2C_1 I_{3\{222\}}^{(\text{fin})} + \dots, \end{aligned} \quad (79)$$

where again the omitted terms are nonsingular in the flat space limit. Using (23) and (43)–(49), we recover the flat space limit in (67).

VII. CONCLUSION

In this paper, we extended to even spacetime dimensions our results for the double copy structure of momentum-space CFT correlators [16]. This double copy structure is inherited from the bulk scattering amplitudes that arise on taking the flat space limit of correlators. Our main achievement is to understand the analytic continuation required to reach the flat space limit. Analytically continuing the largest momentum magnitude, which is an energy from the bulk perspective, we arrive at configurations with $E = 0$ for which bulk energy conservation is restored. Analyzing the behavior of the master integral $I_{1\{000\}}$ under this continuation, the flat space limits of all renormalized correlators can then be constructed.

Prior to analytic continuation, the master integral can be expressed as a 1-loop triangle integral and evaluated in terms of a Bloch-Wigner function in suitable complex variables. Under analytic continuation, these complex variables follow a simple path in the complex plane which involves crossing two branch cuts. The resulting discontinuities produce a new term which supplies the necessary singular behavior in the flat space limit. This term can also be derived from the leading singularity of the 1-loop triangle integral, revealing interesting connections to dual conformal symmetry. The flat space limit of all other triple- K integrals, and that of the correlators themselves, can then be deduced by applying differential operators to $I_{1\{000\}}$.

The above discussion holds for 3-point correlators of stress tensors, currents, and marginal scalars of general CFTs in all even dimensions greater than two, and we verified it explicitly in four dimensions where the renormalized correlators have been fully evaluated. In this case, we showed that anomalies play an important role in the flat space limit. In particular, we found that the flat space limit of stress tensor correlators is controlled by conformal anomalies, in line with general holographic expectations.

It is remarkable that double copy structure plays such a ubiquitous role in correlation functions of general CFTs. It would be interesting to explore how this extends to higher-point correlators in momentum space. A general solution to the conformal Ward identities for n -point scalar correlators in momentum space was recently proposed in [51], so it would be interesting to extend this to tensorial correlators and understand how to systematically compute their flat space limit. Note that the general solution in [51] can be written as a 3-loop Feynman integral so it is conceivable that the flat space limit is encoded in the leading singularity of this integral. It may also be fruitful to look for double copy structure in the correlators derived from Witten diagrams of specific theories in the bulk such as bi-adjoint scalars, Yang-Mills, and Einstein gravity. KLT-like relations for inflationary graviton correlators have been explored in [52], and our results for double copy structure can likewise be applied to cosmology. In the future it would also be interesting to consider correlators of higher spin currents which would

probe higher spin interactions in the bulk [53]. Finally, it would be of interest to explore if the analytic continuation we used to reach the flat space limit can be adapted to continue CFT correlators from Euclidean to Lorentzian signature. This problem has recently been analyzed via other methods in [47–49], but our approach here seems particularly promising.

ACKNOWLEDGMENTS

A.L. is supported by the Royal Society as a Royal Society University Research Fellowship holder. P.M. is supported by the Science & Technologies Facilities Council through an Ernest Rutherford Fellowship, No. ST/P004326/2.

APPENDIX: LEADING SINGULARITY OF THE MASTER INTEGRAL

The master integral $I_{1\{000\}}$ for four-dimensional 3-point CFT correlators is a limit of the dual conformal box integral [54,55]. This box integral also plays a prominent role in the context of $\mathcal{N} = 4$ SYM [56,57]. The leading singularity can easily be computed by writing the box integral in coordinates which make the dual conformal symmetry manifest (the region momentum coordinates), then evaluating the global residue.

We begin by writing the box integral as

$$\Phi(u, v) = x_{13}^2 x_{24}^2 \int d^4 \mathbf{x}_5 \frac{1}{x_{15}^2 x_{25}^2 x_{35}^2 x_{45}^2}, \quad (\text{A1})$$

where $\mathbf{x}_{ij} = \mathbf{x}_i - \mathbf{x}_j$ and the region momentum coordinates are related to the external momenta by

$$\mathbf{x}_{12} = \mathbf{p}_1, \quad \mathbf{x}_{23} = \mathbf{p}_2, \quad \mathbf{x}_{34} = \mathbf{p}'_3, \quad \mathbf{x}_{41} = \mathbf{p}'_4. \quad (\text{A2})$$

The integral is invariant under translations and inversions $\mathbf{x}_i \rightarrow \mathbf{x}_i/x_i^2$, and therefore has conformal symmetry in region momentum space, known as dual conformal symmetry [58]. As a result, it depends only on the dual conformal cross-ratios

$$u = \frac{x_{12}^2 x_{34}^2}{x_{13}^2 x_{24}^2}, \quad v = \frac{x_{14}^2 x_{23}^2}{x_{13}^2 x_{24}^2}. \quad (\text{A3})$$

To recover a 3-point function in momentum space, we define

$$\mathbf{x}_{12} = \mathbf{p}_1, \quad \mathbf{x}_{23} = \mathbf{p}_2, \quad \mathbf{x}_{31} = \mathbf{p}_3, \quad (\text{A4})$$

where $\mathbf{p}_3 = \mathbf{p}'_3 + \mathbf{p}'_4$, and take the limit $\mathbf{x}_4 \rightarrow \infty$. We then recover

$$u = \frac{p_1^2}{p_3^2}, \quad v = \frac{p_2^2}{p_3^2}, \quad (\text{A5})$$




FIG. 7. Region momenta relating the massless box to the 3-point master integral $I_{1\{000\}}$. All external momenta are taken as ingoing.

and setting $l = \mathbf{x}_{51}$, we find

$$\begin{aligned} \lim_{x_4 \rightarrow \infty} \Phi(u, v) &= \int d^4 \mathbf{x}_5 \frac{x_{13}^2}{x_{15}^2 x_{25}^2 x_{35}^2} \\ &= \int d^4 \ell \frac{p_3^2}{\ell^2 (\ell + p_1)^2 (\ell - p_3)^2} \\ &= 4\pi^2 p_3^2 I_{1\{000\}}. \end{aligned} \quad (\text{A6})$$

The box and triangle integrals in region momentum space are depicted in Fig. 7.

An efficient method to evaluate the leading singularity of the box integral was devised in [59]. In place of x_5^μ , we change variables to the four new coordinates

$$P_i = x_{i5}^2, \quad i = 1, \dots, 4. \quad (\text{A7})$$

The Jacobian for this transformation is

$$\mathcal{J} = \det \left(\frac{\partial P_i}{\partial x_5^\mu} \right) = \det(-2x_{i5}^\mu). \quad (\text{A8})$$

As $\det(\mathbf{M}\mathbf{M}^T) = (\det \mathbf{M})^2$, taking $M_i^\mu = -2x_{i5}^\mu$ we have

$$\mathcal{J}^2 = \det(4x_{i5} \cdot x_{j5}) = 2^4 \det(x_{ij}^2 - x_{i5}^2 - x_{j5}^2), \quad (\text{A9})$$

since $\mathbf{x}_{ij} = \mathbf{x}_{i5} - \mathbf{x}_{j5}$ and the matrix is 4×4 . We now have

$$\int d^4 \mathbf{x}_5 \frac{1}{x_{15}^2 x_{25}^2 x_{35}^2 x_{45}^2} = \int \frac{d^4 P_i}{\mathcal{J} P_1 P_2 P_3 P_4}, \quad (\text{A10})$$

and the leading singularity is just the global residue

$$(2\pi i)^4 \frac{1}{\mathcal{J}} \Big|_{P_i=0} = 4\pi^4 \frac{1}{\sqrt{\det x_{ij}^2}} = \frac{4\pi^4}{x_{13}^2 x_{24}^2 (z - \bar{z})}. \quad (\text{A11})$$

The leading singularity of $I_{1\{000\}}$ is then

$$\frac{\pi^2}{p_3^2 (z - \bar{z})} = \frac{\pi^2}{\sqrt{-J^2}}, \quad (\text{A12})$$

which agrees with (23).

- [1] J. M. Maldacena, *Adv. Theor. Math. Phys.* **2**, 231 (1998).
- [2] J. Penedones, *J. High Energy Phys.* **03** (2011) 025.
- [3] J. M. Maldacena and G. L. Pimentel, *J. High Energy Phys.* **09** (2011) 045.
- [4] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **04** (2013) 047.
- [5] N. Arkani-Hamed, D. Baumann, H. Lee, and G. L. Pimentel, *J. High Energy Phys.* **04** (2020) 105.
- [6] R. Britto, F. Cachazo, B. Feng, and E. Witten, *Phys. Rev. Lett.* **94**, 181602 (2005).
- [7] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, *Nucl. Phys.* **B425**, 217 (1994).
- [8] Z. Bern, L. J. Dixon, D. C. Dunbar, and D. A. Kosower, *Nucl. Phys.* **B435**, 59 (1995).
- [9] S. Raju, *Phys. Rev. Lett.* **106**, 091601 (2011).
- [10] S. Raju, *Phys. Rev. D* **83**, 126002 (2011).
- [11] S. Raju, *Phys. Rev. D* **85**, 126009 (2012).
- [12] L. F. Alday and S. Caron-Huot, *J. High Energy Phys.* **12** (2018) 017.
- [13] L. Rastelli and X. Zhou, *J. High Energy Phys.* **04** (2018) 014.
- [14] X. Zhou, *J. High Energy Phys.* **05** (2019) 006.
- [15] Z. Bern, J. J. Carrasco, M. Chiodaroli, H. Johansson, and R. Roiban, [arXiv:1909.01358](https://arxiv.org/abs/1909.01358).
- [16] J. A. Farrow, A. E. Lipstein, and P. McFadden, *J. High Energy Phys.* **02** (2019) 130.
- [17] R. E. Cutkosky, *J. Math. Phys. (N.Y.)* **1**, 429 (1960).
- [18] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **11** (2018) 153.
- [19] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **11** (2018) 159.
- [20] M. Henningson and K. Skenderis, *J. High Energy Phys.* **07** (1998) 023.
- [21] S. Nojiri and S. D. Odintsov, *Int. J. Mod. Phys. A* **15**, 413 (2000).
- [22] F. Bugini and D. E. Diaz, *J. High Energy Phys.* **04** (2017) 122.
- [23] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **03** (2014) 111.
- [24] S. Giombi, S. Prakash, and X. Yin, *J. High Energy Phys.* **07** (2013) 105.
- [25] S. D. Chowdhury, J. R. David, and S. Prakash, *J. High Energy Phys.* **11** (2017) 171.
- [26] L. Bonora, A. D. Pereira, and B. L. de Souza, *J. High Energy Phys.* **06** (2015) 024.
- [27] L. Bonora and B. L. de Souza, in *Proceedings of the 18th Workshop on What Comes Beyond the Standard Models?, Bled, Slovenia, 2015* (2015), pp. 22–34.
- [28] J. P. B. Almeida, J. Mota-Manzano, and C. A. Valenzuela-Toledo, *J. Cosmol. Astropart. Phys.* **11** (2017) 015.
- [29] M. Gary, S. B. Giddings, and J. Penedones, *Phys. Rev. D* **80**, 085005 (2009).
- [30] N. Arkani-Hamed and J. Maldacena, [arXiv:1503.08043](https://arxiv.org/abs/1503.08043).
- [31] J. M. Maldacena, *J. High Energy Phys.* **05** (2003) 013.
- [32] P. McFadden and K. Skenderis, *J. Cosmol. Astropart. Phys.* **05** (2011) 013.
- [33] P. McFadden and K. Skenderis, *J. Cosmol. Astropart. Phys.* **06** (2011) 030.
- [34] J. Broedel and L. J. Dixon, *J. High Energy Phys.* **10** (2012) 091.
- [35] H. Johansson and J. Nohle, [arXiv:1707.02965](https://arxiv.org/abs/1707.02965).
- [36] A. Gervois and H. Navelet, *J. Math. Phys. (N.Y.)* **27**, 682 (1986).
- [37] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **02** (2016) 068.
- [38] D. Zagier, in *Proceedings of the Les Houches School of Physics: Frontiers in Number Theory, Physics and Geometry II: On Conformal Field Theories, Discrete Groups and Renormalization, Les Houches, France, 2003* (Springer-Verlag, Berlin, Heidelberg, 2007), pp. 3–65.
- [39] L. Lewin, *Polylogarithms and Associated Functions* (Noord-Holland, New York, 1981).
- [40] F. C. S. Brown, *Compt. Rend. Math.* **338**, 527 (2004).
- [41] L. J. Dixon, C. Duhr, and J. Pennington, *J. High Energy Phys.* **10** (2012) 074.
- [42] J. M. Drummond, *J. High Energy Phys.* **02** (2013) 092.
- [43] F. Chavez and C. Duhr, *J. High Energy Phys.* **11** (2012) 114.
- [44] J. Maldacena, D. Simmons-Duffin, and A. Zhiboedov, *J. High Energy Phys.* **01** (2017) 013.
- [45] C. Cardona, *J. High Energy Phys.* **11** (2019) 044.
- [46] A. Bzowski, P. McFadden, and K. Skenderis, *J. High Energy Phys.* **03** (2016) 066.
- [47] T. Bautista and H. Godazgar, *J. High Energy Phys.* **01** (2020) 142.
- [48] M. Gillioz, [arXiv:1909.00878](https://arxiv.org/abs/1909.00878).
- [49] N. Anand, Z. U. Khandker, and M. T. Walters, [arXiv:1911.02573](https://arxiv.org/abs/1911.02573).
- [50] S. Deser and A. Schwimmer, *Phys. Lett. B* **309**, 279 (1993).
- [51] A. Bzowski, P. McFadden, and K. Skenderis, *Phys. Rev. Lett.* **124**, 131602 (2020).
- [52] S. Y. Li, Y. Wang, and S. Zhou, *J. Cosmol. Astropart. Phys.* **12** (2018) 023.
- [53] M. A. Vasiliev, *Nucl. Phys.* **B862**, 341 (2012).
- [54] D. J. Broadhurst, *Phys. Lett. B* **307**, 132 (1993).
- [55] N. I. Usyukina and A. I. Davydychev, *Phys. Lett. B* **298**, 363 (1993).
- [56] A. Brandhuber, P. Heslop, and G. Travaglini, *Phys. Rev. D* **78**, 125005 (2008).
- [57] J. M. Drummond, J. Henn, G. P. Korchemsky, and E. Sokatchev, *Nucl. Phys.* **B828**, 317 (2010).
- [58] J. M. Drummond, J. Henn, V. A. Smirnov, and E. Sokatchev, *J. High Energy Phys.* **01** (2007) 064.
- [59] J. Drummond, C. Duhr, B. Eden, P. Heslop, J. Pennington, and V. A. Smirnov, *J. High Energy Phys.* **08** (2013) 133.