Accepted Manuscript

Victoria cabrera site: A middle stone age site at Olduvai gorge, Tanzania

José-Manuel Maíllo-Fernández, Juan Marín, Irene Solano-Megías, David Uribelarrea, David Martín-Perea, Julia Aramendi, Alicia Medialdea, Carlos Arteaga, Marta Pernas-Hernández, Agness Gidna, Sayantani Neogi, Eva Baudot, Carlos Narváez, Audax Mabulla

PII: S1040-6182(19)30365-9

DOI: https://doi.org/10.1016/j.quaint.2019.07.032

Reference: JQI 7960

To appear in: Quaternary International

Received Date: 18 April 2019

Revised Date: 24 July 2019

Accepted Date: 27 July 2019

Please cite this article as: Maíllo-Fernández, José.-Manuel., Marín, J., Solano-Megías, I., Uribelarrea, D., Martín-Perea, D., Aramendi, J., Medialdea, A., Arteaga, C., Pernas-Hernández, M., Gidna, A., Neogi, S., Baudot, E., Narváez, C., Mabulla, A., Victoria cabrera site: A middle stone age site at Olduvai gorge, Tanzania, *Quaternary International* (2019), doi: https://doi.org/10.1016/j.quaint.2019.07.032.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.



1 2

6 7

8

VICTORIA CABRERA SITE: A MIDDLE STONE AGE SITE AT OLDUVAI GORGE, TANZANIA.

José-Manuel Maíllo-Fernández^{1, 2*}, Juan Marín³, Irene Solano-Megías¹, David Uribelarrea^{2,4}, David
 Martín-Perea^{2,5}, Julia Aramendi^{2,6}, Alicia Medialdea⁷, Carlos Arteaga⁸, Marta Pernas-Hernández⁶, Agness
 Gidna⁹, Sayantani Neogi¹⁰, Eva Baudot⁸, Carlos Narváez⁸, Audax Mabulla⁹.

Abstract:

9 Olduvai Gorge (Tanzania) is a key site for the study and comprehension of human evolution in East 10 Africa. However, the origin of Homo sapiens and the Middle Stone Age have been poorly understood in 11 the Gorge thus far. In this study, we present the dating, taphonomic, technological and typological 12 analyses of the lithic industry and faunal remains excavated at the Victoria Cabrera Site (VCS) during the 13 2017 fieldwork season. The stratigraphic sequence of the site contains several levels of fluvial origin, 14 some of them with lithic and faunal remains. Most remains (lithics and bones) are affected by rounding 15 processes. Infrared stimulated luminescence (IRSL) dating has provided ages between 90-70 ka BP for 16 these layers that can be related to the Marine Isotope Stage 5a. Lithics are scarce and the dominant raw 17 materials are Naibor Soit quarzite and basalt, followed by phonolite, chert and quartz as the most 18 common raw materials. The lithic technology is based on the production of flakes obtained from prepared 19 cores, with the discoid method and, to a lesser extent the Levallois method being the most frequently 20 used. The retouched blanks are described as "substratum" or "domestic" tools (sidescrapers, notches or 21 denticulates). There is just one heavy duty piece and points are absent. The faunal assemblage includes 22 Alcelaphinae, Antilopinae and Equidae. The major parts of the unidentified faunal remains are size 3 23 ungulates, followed by carcasses of size 2 and size 4 species. We cannot relate lithic and bones because 24 no cut marks or percussion marks have been identified and carnivore action is scarce. In sum, VCS 25 represents the first accurately dated Middle Stone Age site, with lithic and faunal remains in Olduvai 26 Gorge. 27

Key words: Middle Stone Age, Lithic, Fauna, Olduvai, Tanzania

- 28 29 30
 - ¹ Dpt. Prehistory and Archaeology, UNED. Paseo Senda del Rey, 7, 28040 Madrid, Spain.
- 31 ² IDEA, The Institute of Human Evolution in Africa. C/ Covarrubias 36, 28010, Madrid, Spain.
- 32 ³ Muséum National d'Histoire Naturelle (MNHN). 57, rue Cuvier, 75005. Paris, France.
- ⁴ Dpto. de Geodinámica, Estratigrafía y Paleontología, UCM. C/ José Antonio Novais, 12, 28040 Madrid,
 Spain.
- ⁵ Dpto. de Paleobiología (MNCN). C/ José Gutiérrez Abascal, 2, 38006 Madrid, Spain.
- ⁶ Dpto. de Prehistoria, Historia Antigua y Arqueología, UCM. C/ Profesor Aranguren, s/n, 28040,
 Madrid, Spain.
- ⁷ Institute of Geography, University of Cologne. Albertus-Magnus-Platz, 50923 Cologne, Germany.
- ⁸ Laboratorio de Geografía Física, UAM. C/ Francisco Tomás y Valiente, 1, 28019, Madrid, Spain.
- 40 ⁹ National Museum of Tanzania. 5, Shaaban Robert St, Dar es Salaam, Tanzania.
- 41 ¹⁰ Department of Archaeology, Durham University. Lower Mount Joy, South Rd, Durham DH1 3LE, UK.
 - * Corresponding author: <u>ilmaillo@geo.uned.es</u> (José-Manuel Maíllo-Fernández)

45 46 1.

42 43

44

47

1. Introduction.

48 Olduvai Gorge has been known on merit as the Cradle of Humankind since 1959, when Mary and Louis 49 Leakey discovered the *Zinjanthropus* (*Paranthropus*) skull in the FLK site. Since that time, several 50 research teams and projects have devoted considerable effort to the paleoanthropological, cultural and 51 behavioural knowledge of the first figures of genus *Homo*: the Oldowan and the Acheulean. However, 52 little attention has been paid to upper beds in Olduvai Gorge, as well as to the origin of our species and its 53 material culture.

54

Olduvai Gorge is located in the South-eastern Serengeti (northern Tanzania), in the great East African Rift (Fig. 1). The erosion of the Olduvai River, cutting through a plain amidst a line of volcanoes in the South—e.g. Sadiman, Lemagrut, Olmoti or Ngorongoro—and lower-height metamorphic reliefs in the North—like Olongoidjo or Watumi—generated the gorge (Uribelarrea, 2014). The structural landscape remained stable until the Ngorongoro Collapse 2.2 Ma, whilst the plain was affected by external processes (weathering, and fluvial and aeolian sedimentation). Following the basin sedimentation, the

61 Olduvai River eroded the plain generating the gorge we know today, split into the Main and Side Gorges. 62 Throughout that process, an intense tectonic activity as well as further basin sedimentation took place, generally by volcanic processes during the Middle and Late Pleistocene. Thanks to their low-energy 63 64 sedimentary conditions, the availability of water sources and the primary sources of lithic raw material, 65 the spot was the perfect place for the occupation of the territory by hominin groups. Such an assertion is 66 contrasted with the more than 120 sites and the remains of four hominin species discovered (Paranthopus 67 boisei, Homo habilis, Homo ergaster/erectus and Homo sapiens). Nowadays, the gorge geology can be 68 split into seven beds (Hay, 1976).

69

70 Bed I (2.1Ma) in Olduvai Gorge has been well-known since the 1930s. Bed I deposits contain an 71 important amount of sites bearing hominin fossil remains and archaeological assemblages (including 72 fauna and lithic remains) that seem to belong to the first representatives of our genus (Homo) as well as to 73 other contemporary hominin species (Paranthropus boisei) (Leakey, 1971, Uribelarrea, 2014, 74 Domínguez-Rodrigo et al., 2007, 2010, Barboni et al., 2010). Oldowan and Acheulean industries have 75 been discovered in Bed II in sites (e.g. BK or SHK) in association with remains of Homo erectus (Ashley 76 et al., 2014, Domínguez-Rodrigo et al., 2013, Uribelarrea et al., 2014). Beds III and IV have provided 77 more than 40 Acheulean sites (Leakey and Roe, 1994). After that, the Masek Bed was deposited while the 78 great faults affect the gorge move (Leakey and Roe, 1994, Hay, 1976). This Bed represents the 79 sedimentary clogging of the basin. At this juncture, around 0.6 and 0.4 Ma, the first fault movement 80 happened, together with the movement of the five faults affecting the gorge. The result is the creation of 81 the Olbalbal depression-in the east of the gorge-and the creation of the Olduvai River itself, which 82 begins in Lake Ndutu and flows into the newly created depression. This marks the start of Olduvai Gorge 83 due to the exposition of the above-mentioned deposits on account of the erosive process (Hay, 1976, 84 Uribelarrea, 2014). The levels of Ndutu and Naisiusiu Beds will be deposited above them.

85

86 The Middle Stone Age (MSA) has been defined by the knapping of prepared cores, especially the 87 Levallois-type and, above all, by retouched pieces, particularly projectile points associated to Homo 88 sapiens (Clark, 1988; McBrearty and Books, 2000, Basell, 2013). Stratigraphically, the MSA in Olduvai 89 falls within the Ndutu Bed, which consists principally of tuffaceous, aeolian, fluvial and colluvial 90 deposits, which are deposited in the final stage of the gorge formation and, which cover the eroded areas 91 by the Olduvai River itself (Hay, 1976). Research on the human occupation of this Bed has been 92 discontinuous and biased in the three most relevant lines of research: chronology, human remains, and 93 archaeology.

94

95 One unsatisfactorily resolved issue for Bed Ndutu is its chronology. The initial approach was carried by 96 Hay, who considered that Upper Ndutu might be in the range of 60-32 ka BP, and Lower Ndutu in the 97 range of 60-400 ka BP (Hay, 1976: 152), although he used indirect methods to this end. Manega (1993) is 98 without a doubt, the researcher who has devoted the utmost efforts to discover the chronology of the 99 Ndutu Bed. According to his research, on the basis of the Single Cristal Laser Fusion (SCLF) technique, 100 Upper Ndutu might be located in a range between 210 ± 20 ka BP and 450 ± 40 ka BP and using amino 101 acids around 260-500 ka BP. Manega dates the start of Bed Naisiusiu in 42 ± 10 ka BP by using SCLF. 102 The latter runs counter to Skinner's dating for the same Bed, which was dated in 62 ± 5 ka BP (Skinner et 103 al., 2003)-and it is therefore, far older than Manega's proposal-by using ESR (electron spin 104 resonance).

105

Several Homo sapiens remains have been found in the Ndutu Bed. The first one is OH-11, a palate and 106 107 maxillary arch of Homo sp found on the surface in western DK and assigned to the Ndutu Bed (Hay, 1976, pp. 159, Leakey 1971, pp. 230), which presents a sturdy morphology for the modern standards 108 109 (Rightmire, 1980, pp. 227). The second one is OH-1968, a skull fragment partially in situ in Upper Ndutu 110 levels (Von Zieten, 2009), but whose origin and study are uncertain. The last one is OH-83, a partial 111 calvaria partially recovered in stratigraphy in the Upper Ndutu levels, close to the PLK site (Reiner et al., 2017) and identified as Homo sapiens. There is no direct dating for any of them, therefore they have been 112 113 placed in the chronological range proposed by Hay for Ndutu (Hay, 1976, pp. 152) with the uncertainties 114 that, as we have seen, it presents.

115

The archaeological study of the Ndutu Bed has also been intermittent over the last decades, with only three previous interventions. The first paper on MSA was carried out by Mary Leakey and colleagues, and in it, two sites with MSA industry are presented (Leakey et al., 1972). Both sites were not located on the paper, tough follow-up work placed them in the localities 4b and 26 (Hay, 1976, pp. 159) without further details of planimetry or stratigraphy, one of them belonging to Upper Ndutu and the other to

Lower Ndutu (Hay, 1976, pp. 28). The lithics of both sites were presented uniformly, and in there, the
presence of prepared cores (Levallois and discoid) and a few retouched tools was outlined. The most used
raw material was basalt, and there was a small quantity of phonolite and quartz. The assemblages,
although with due precaution, were linked to the Early MSA of Lake Eyasi (Leakey et al. 1972).

- The second project related to the MSA was directed by A. Mabulla, co-author in this study, from
 the University of Dar es Salaam in collaboration with the University of Florida between 1989 and 1990
 (Mabulla 1990). Several localities shedding light on lithic and fauna find-spots were explored—none of
 them in stratigraphy except for locality 26 (Ndutu type site according to Hay, 1976).
- 130

131 The third research project was led by M. Eren during the course of the 2013 field season and was the 132 most intense study conducted so far. They carried out a systematic survey between the second fault and 133 the Olbalbal depression, besides other selective surveys in the junction of both gorges, resulting in the 134 discovery of 72 find-spots which provided 521 MSA flaked pieces, as well as some faunal remains (Eren et al., 2014).

136

Even though the two first projects had already made it clear that the Ndutu Bed reported having
occupations belonging to MSA, certainly after the work of M. Eren's team, the extraordinary potential
which Olduvai Gorge has for the first modern humans in northern Tanzania became clear.

In this paper, we present the first results of Victoria Cabrera Site (VCS), which is the first site in stratigraphy with radiometric dating and several stratigraphic levels with lithic industry and faunal remains found in Olduvai Gorge.

145 2. Victoria Cabrera Site.

Victoria Cabrera Site (VCS) is located on the South side of Olduvai Gorge, at the junction between the
Main and the Side gorge. It is on the same hill as DS, 70 meters due East of it and over 600 meters from
FLK Zinj (Fig. 1). It was discovered by one of us (JMMF), along with Manuel Domínguez-Rodrigo in
the course of the 2016 survey work, and it has been dug in two seasons during February and July 2017. It
could not be excavated in 2018 as the site suffered an act of vandalism in which some "vandals" opened
the site without official permission, exposing the stratigraphic profiles.

The works carried out so far have consisted of opening a 24 x 2m linear trench on the hillside, as well as a
9 x 2m trench transversal to the former. Both of them have exposed a Lower Ndutu stratigraphy made up
of several levels, six of them containing lithic industry corresponding to the MSA and faunal remains.

157

Following the geologic order of formation, the following five stratigraphic levels have been identified at
VCS Trenches 1 and 2 (Fig. 2):

161 Level 5, silty tuffaceous claystone

162
163 Level 5 is a < 120 cm reddish brown sandy and silty tuffaceous claystone. This level can be interpreted as
164 an overbank deposit, formed in decantation facies, highly contaminated by tuffaceous deposits. In Trench
1, the uppermost section of the level contains a higher tuffaceous content and has been named Level 5a.

166167 Level 4, Very coarse to medium sand

Level 4 is a >30 cm thick greyish very coarse to medium parallel-stratified sand (20.50% very coarse
sand, 20.60% coarse sand, 21.30% medium sand). Level 4 is interpreted as a fluvial bedload level,
deposited in a constant and relatively high velocity regime. Similar deposits can be found currently in the
bank deposits of Olduvai River. Horizontal sets extend over 1 m, part of a fluvial bar, yet to be described
in detail. In Trench 1, Level 4 appears more massive, and no sedimentary structures are observed,
probably due to the degree of weathering suffered.

175

176 Level 3, Tuff breccia with sandy matrix (debris flow)

177

Level 3 is a yellowish matrix-supported breccia, composed by heterogeneous angular tuff clasts from 2
cm of diameter to blocks over 20 cm, and most probably eroded from upper Bed I tuffs. The matrix is
brownish yellow tuffaceous sand (77.38% sand) with a considerable amount of fines (10.08% silt,

181 12.60% clay). The level has an erosive base, eroding the uppermost section of underlying Levels 4 and 5.
182 Level 3 is lost eastwards due to present-day erosive processes. Size and angular shape of tuff blocks, with provenance most likely from Bed I marker tuffs which are seen eroded by these deposits in Trench 1, indicate very short transport.

186 Level 2, Sandy gravel

185

202

208 209

210

218

227

236

187 188 A greyish sandy gravel (66.90% gravel, 29.53% sand), very washed and clean (<5% silt and clay), lies 189 over a channelled erosional surface, pinching out towards the East and West in Trench 2 and towards the North and South in Trench 1 (Fig. 2). Trough cross-stratification (Gt lithofacies according to Miall, 1985) 190 191 is observed in the South Profile at Trench 2 (Fig. 2). Level 2 is interpreted as a fluvial bedload deposit. A 192 separate channel has been identified in Trench 1 and named Level 2a. In Trench 2, Level 2 contains some 193 Bed I tuff blocks, interpreted as topples generated from the erosion of the northern river margin. Lenses 194 of finer grained material are found in between bedding, suggesting a multi-episodic genesis. 195

196 Level 1, clayey tuffaceous siltstone197

The fluvial sequence ends with Level 1, which is a < 80 cm reddish brown sandy and clayey (24.55% fine sand, 21.88% clay) tuffaceous siltstone (22.39% silt), interpreted as another low-energy overbank deposit in the floodplain. Level 1 is eroded towards the East and is discomformably overlain by Holocene deposits.</p>

The whole sequence is overlain discomformably by Upper Ndutu aeolian sand bodies, and is being
eroded, forming a gully perpendicular to present-day Olduvai River, partially infilled with Ndutu, Bed I
and fluvial terrace deposits (Uribelarrea et al, this volume).

206207 3. Methodology.

3.1. Spatial Analysis.

Total stations were used to geo-reference all remains > 2 cm. Orientation patterns of bone and lithic remains were considered to reinforce the geological reconstruction of the site and to assess the impact of post-depositional events (e.g. wind, water flows, gravity, sedimentation) on the archaeological arrangement in each of the documented levels. We used compasses and clinometers to measure the horizontal and vertical orientations of each piece of remains with a longitudinal axis at least twice of its width (Voorhies 1969, Fiorillo 1991, Alcalá 1994, Howard 2007, Domínguez-Rodrigo and García-Pérez, 2013).

219 Orientation data were statistically treated using the "circular" package in the R free software 220 (http://www.r-project.org, Core R-Team, 2013). The uniformity of the different levels was statistically 221 evaluated by combining tests that can detect isotropy/anisotropy, where significance values (p) > 0.05222 indicate that the null hypothesis of isotropy cannot be rejected. Rayleigh test (Z) was used to evaluate 223 anisotropy (Fisher, 1995), and the omnibus Kuiper (V) and Watson (U^2) tests were used to determine if 224 the orientation of the archaeological assemblage was uniform, bimodal or polymodal (Fisher, 1995). 225 Statistical tests are accompanied by graphical plots. The spatial information was analyzed using ArcGIS 10.4.1 (Desktop, Engine), Rose.NET (Rosediagrams) and OpenStereo (Stereograms). 226

228 Orientation analyses were conducted for all VCS levels treating bones and stone tools separately as they 229 respond differently to post-depositional forces that can generate preferential orientation patterns (Frostick 230 and Reid, 1983, Schick, 1984). Since data tend to be scarce in all levels, measurements were bootstrapped 231 before performing statistical tests. A small number of permutations (replacement of 100 times) was 232 preferred as they provide meaningful statistical results without excessively distorting the original data. 233 Only in the case of the bone assemblage in level Tuff 2, the sample size was not large enough to provide 234 any clear results (4 pieces. The rest of the samples included at least ca. 10 elements. Ripley's K function 235 was used to describe how materials occur in any specific spatial area (Domínguez-Rodrigo et al., 2017).

237 3.2. Faunal Analysis.

Faunal remains were assigned to taxa, size category, anatomical element, side and portion when possible.
For the anatomical part representation analysis, carcasses were divided in anatomical areas: cranial (horn,

cranium, mandible, hyoid and teeth), axial (vertebra, ribs, pelvis and scapula) and appendicular (limb
bones). Four indexes have been used to estimate the taxonomic and anatomical abundance: number of
identified specimens (NISP); minimal number of element (MNE); standardized minimal animal units
(%MAU); minimal number of individual (MNI) (Grayson, 1984, Binford, 1984, Lyman, 1994). To
estimate the MNE, side, landmarks and age of each identifiable specimen were considered. The size of
the specimens was established according to the categories described by Bunn (1982): size 5 (over 1000
kg); size 4 (300-1000 kg); size 3 (70-300 kg); size 2 (18-100 kg); size 1 (>4.5-22 kg) (Brain, 1981).

248

249 The Spearman's rank order was used to test the correlation between %MAU and bone mineral density 250 (Lam et al., 1999), and to test a possible correlation between %MAU and standard food utility index 251 ((S)FUI). Following the proposal of Marean and Clerghorn (2003) we used high survival element. Length 252 distribution was examined in order to evaluate the impact of hydraulic processes, addressing also only 253 long bones separately, and finally long bones with green bone breakage (Domínguez-Rodrigo and 254 Martínez-Navarro, 2012, Organista et al., 2016). Bone surface modifications were examined using 10x40 255 hand lenses under a strong oblique light source. Cortical surfaces were evaluated and analyzed, and as a 256 result numerous types of modifications such as tooth marks, trampling, polished, fissures and rounding 257 were identified (Behrensmeyer, 1978, Binford, 1981, Capaldo, 1998, Fernández-Jalvo and Andrews, 258 2003, Domínguez-Rodrigo et al., 2009, Pickering and Egeland, 2006).

259

The description and quantification of the ostrich eggshells follows the criteria stated by Collins and Steele(2017) for the study of OES flakes.

262263 3.3. Lithic Analysis.

3.3.1. Taphonomic analysis: Knowing all the possible post-depositional processes undergone by an archaeological assemblage is essential in order to make correct interpretations. For the VCS lithic assemblage, we paid particular attention to the following alterations:

Rounding. It refers to the wear and tear of an artefact surface due to the brush of the lithic tool with other sedimentary particles, giving rise to breakages, wear, and micro-detachments (Pedraza, 1996, p. 210). It can happen because of several reasons, though it is linked to hydric phenomena where water converges with the carried load. A greater rounding could mean a greater transport. It is important to take into account the transported sediment as well (Shackley, 1978). We have classified the rounding into three categories: not very intense, and very intense.

275

276 Pseudo-retouch. It refers to a mechanical alteration affecting the surface of the lithic material. After 277 Sieveking (1976) it provokes an alternating and denticulated edge and the scars have variable sizes. Other 278 scholars note the random scars distribution of no fixed orientation or size (Tringham et al. 1974: 192) It is 279 usually provoked by the stone movement in the sediment because of trampling or hydraulic processes 280 (Stapert, 1976). The analysis of this kind of alteration has focused on the dispersion in the vertical 281 direction (Courtin and Villa, 1982, Hughes and Lampert, 1977, Trigham et al., 1974, Gifford-González et 282 al., 1985), on its relationship with the use-wear (Levi-Sala, 1986) and on the creation of pseudo-tools 283 (McBrearty et al., 1998). We have classified the pseudo-retouch into three categories: marginal, deep and 284 very deep.

286 3.3.2. Techno-typological analysis:

The term technology is understood as a conceptual approach to the material culture of the various human
groups (Inizan et al., 1996, p. 13). The ultimate goal of this sort of studies is to investigate human
behaviour through the study of technical, social or symbolic behaviours (Inizan et al. 1996, p. 14).

291

285

287

292 To carry out this kind of research, we analysed the *chaîne operatoire* understood as a sequence of acts, 293 gestures, and tools constituting a technical process with a roughly predictable series of steps (Karlin, 294 1992, p. 109; Karlin et al., 1992, Audouze and Karlin, 2017). From an empirical viewpoint, it is defined 295 as joint operations converging towards a single objective, whereas at the structural level it is defined as a 296 combination of technical factors, which materialize into technical schemes. Thus, the chaîne operatoire 297 only goes as far as explaining the use of a core/blanks when several blanks and cores have been exploited 298 by following the same method, and these are then grouped into operational schemes that make up the 299 lithic technical subsystem of a site (Audouze and Karlin, 2017).

The lithic industry studies pursue two different approaches: a quantitative approach comprising the data
 for each individual piece, and a qualitative approach focussed on the reconstruction of the operational
 schemes (Karlin, 1992). Unfortunately, the VCS lithic assemblage is very scarce to develop the second
 one.

306 In the former, raw materials, dimensions, support, cortex, accidents, fractures, alterations, platform type, 307 scar direction, section, morphology, technique and technological classification are recorded. In addition, 308 if the piece was modified this is also recorded as a retouched cutting edge, as well as the sort of retouch 309 and its typological classification. In addition, we analyse other parameters on the cores such as 310 morphology, method, type of blank (flake/blade/bladelet) and length and width of the last and biggest 311 blanks (after Geneste 1986, Maíllo-Fernández, 2003). Prepared knapping methods are described by 312 different protocols: Levallois (Boëda, 1994, Boëda et al., 1990) and Discoid (Boëda, 1993, 1995, 313 Terradas, 2003, Mourre, 2003).

314315 3.4. Chronology.

316

305

317 Three samples from levels 1, 4 and 5 of Trench 2 have been collected for luminescence dating to establish 318 a chronological record for this trench. Infrared luminescence (IRSL) on the feldspar extracts of these units 319 has been measured at the Institute of Geography of the University of Cologne. The potassium feldspar 320 fractions of size 180-250 µm have been used for the luminescence measurements and equivalent dose 321 estimation. post-IR-IRSL225 protocol (Thomsen et al., 2008). This protocol involves measurement of the 322 IRSL signal at 225°C after a low temperature (50°C) stimulation which minimizes the effect of 323 anomalous fading. This is the fading of the signal which can be significant over the burial time but would 324 be negligible over the timescale of the laboratory measurements and therefore, challenging to quantify 325 accurately. This can lead to underestimation of the age. Fading tests involving the measurement of the 326 IRSL signal after delays of different lengths in the laboratory were carried out to determine the impact of 327 fading (cf. Auclair et al., 2003). Also, dose recovery tests on artificially bleached and irradiated 328 subsamples were carried out to confirm the suitability of the chosen measurement protocol. 24-36 multi-329 grain aliquots of each sample were measured in order to get representative natural dose distributions. The 330 resulting data have been reduced to limit the scatter by removing individual outliers according to the 331 outlier definition of Tukey, 1977 (values outside 1.5 times the inter-quartile-range). Following this 332 criterion less than 10% of the measured dose values were excluded. Central Age Model (CAM, Galbraith 333 et al., 1999) from the reduced dose distribution was used to estimate the burial dose. Environmental dose 334 rates are based on the radionuclide activity concentration from high resolution gamma spectrometry 335 measured at the Radioisotopes Unit of the University of Seville. An internal potassium content of 336 12.5 \pm 0.5 % has been assumed for the feldspar grains of the three samples. A water content of 10 \pm 5 % has 337 been considered to cover the humidity conditions during the burial period. Total dose rates have been 338 calculated using DRAC v1.2 (Durcan et al., 2015). 339

340 **4. Results.**

4.1. Spatial Analysis.

342 4.1. S 343

341

344 The stereographic projections and omnibus tests performed on the arrangement of lithic and bony elements show a horizontal trend and anisotropic distribution in all VCS levels (Figs. A.1 and A.2). The 345 346 obtained p values < 0.05 indicate that there is not a uniform distribution along the stratigraphic sequence. 347 The Kuiper test of uniformity, the Watson goodness-of-fit test of circular uniformity and the Rayleigh test 348 of a general unimodal alternative suggest that the assemblage is largely oriented (Table 2). Thus, the 349 hydraulic flows detected in the geology of the site were sufficiently intense to preferentially orient 350 materials and cause a strong post-depositional disturbance. Most tests performed on the lithics and bones 351 found in VCS show very similar results and levels of significance; only the osseous assemblage in level 4 352 indicates that the null hypothesis of uniform distribution cannot be rejected, and that the unimodal 353 alternative is not valid in this case. This is also supported by the graphical output displayed in Fig. 2. The 354 Ripley's K function indicates a random disposition for lithics and faunal remains in distances smaller than 355 200 cm, except for faunal remains of level 3 that show a regular dispersion (Figs. A.3 and A.4).

357 4.2. Fauna.

356

358

A total of 148 faunal remains were discovered at VCS. Attending to the NISP, Levels 4 and 3 contain the
 major part of the materials, followed by Levels 2 and 2a. Levels 5a and 5 have five and three remains

361 respectively (Table 3). The identification rate at VCS is low (9.4%). Remains of Alcelaphinae have been 362 found in Level 3 and 4, of Antilopinae in Level 2 and some belonging to Equidae in Levels 2, 2a, 3 and 4. 363 In most cases it was not possible to identify the specific taxa. A fragment of Hippopotamidae tooth was 364 found in Level 2, a plastron fragment of *Testudinae* in Level 5a and two remains of *Rodentia* in Level 3. 365 A total of 35 ostrich eggshell flakes (OES) were recovered from Level 2a (NR=3) and Level 4 (NR=32). 366 The major part of the unidentified faunal remains belongs to size 3 ungulates, followed by size 2 and size 367 4 species (Table 3). Large animals (size 5) are only present in Levels 3 and 4. Small animals (size 1) are 368 scarcely represented in Level 2 (two fragments) (Tables A.1-A.4). The total MNE (n = 41) and the total 369 MNI (n = 11) present the same distribution as the NISP (Table 3).

370

381

371 The anatomical distribution of elements in not homogeneous; while Levels 2 and 3 show a predominance 372 of cranial elements (Level 2=50 %/Level 3=41.7 %), followed by axial (Level 2=33.3%/ Level 3=33.3%) 373 and appendicular elements (Level 2=16.7%/ Level 3=25%), Level 4 contains a predominance of axial elements (50%), followed by appendicular (35.7%) and I cranial element (14.3%). Level 2a has the same 374 375 proportion of cranial and appendicular elements (40 %), followed by axial bones (20%). Finally, in 376 Levels 5a and 5 only one cranial and one piece of axial remains were found (Tables A.2-A.4). These 377 anatomical representations indicate a predominance of low survival elements (58.2%) at VCS (Table A.5) 378 existed at the entire sample a low positive relationship between % MAU and bone mineral density 379 (r=0.22921; p=0.023189). This indicates a bias of the low dense portions of bones. The lack of faunal 380 refitting indicates as well intense post-depositional disturbance.

382 The anatomical distribution patterns observed at VCS, especially in Levels 3 and 4, is similar to the 383 scenario proposed by Behrensmeyer (1982) for fluvial contexts. He described three possible sources of 384 pathways for fluvial deposits characterized by specific anatomical profiles: Pathway A with high 385 proportions of denser and more compact elements (Teeth, mandibles, limb ends); Pathway B with the 386 presence of the more transportable bones (rib, vertebrae, scapulae, pelvis and skulls; Pathway C with all 387 bones available. VCS is probably the result of a combination between A and B, with the presence of the 388 more dense and compact elements and a predominance of the lighter and more transportable bones (axial 389 skeleton). Organista and colleagues (2017) suggest that an underrepresentation of shafts may indicate the 390 impact of water flow during sedimentation. At VCS epiphysis fragments were found only in Level 3, 391 including a complete metatarsal of an Equidae and a proximal epiphysis fragment of a metapodial that 392 belongs to a sized-2 ungulate. The statistical analysis performed on the (S)FUI and % MAU of high 393 survival elements indicates that there is not a correlation between the anatomical distribution and the 394 utility index (Table A.6). 395

Bone fragment lengths are differently distributed among levels. Specimens smaller than 20 mm are 396 397 poorly represented or absent in all levels. Specimens between 20-60 mm are abundant, especially in 398 Levels 2, 3, 4 and 2a. The amount of remains larger than 100 mm is remarkable in Level 5, but very low 399 in the rest of the levels at VCS (Fig. A.5). Some previous works relate the scarcity of small bone 400 fragments with water transport (Schick, 1984, Domínguez-Rodrigo and García-Pérez, 2013). A total of 401 five complete bones were found, in Level 4, including an astragalus, a thoracic vertebra, a first phalanx of 402 a sized-2 animal, and a cervical vertebra and a metatarsal of an Equidae. This data, alongside the 403 anatomical distribution patterns observed, reinforce the idea that water transport played an important role 404 in the final arrangement of site. A total of 32 remains present signs of green bone breakage, while only 8 405 bones show signs of dry fracture. In Levels 4 and 2a two bones fractured by the action of carnivores were 406 found. No bones show a combination of green and dry bone breakage. Green bone breakage is more 407 common among long bones (NR=27), however it was also identified in some flat bones (NR=5). The 408 absence of small bone fragments, together with the complete lack of faunal refitting, support the 409 hypothesis of VCS faunal arrangement as the result of water flow (Domínguez-Rodriguez et al., 2012, 410 Domínguez-Rodrigo and García-Pérez, Organista et al., 2016).

411

412 Considering the small sample size of the assemblage all remains were used to infer the taphonomical 413 history of VCS. No atrophic modifications were identified in any of the levels. Only four pieces bearing 414 tooth marks have been identified, two in Level 4, one in Level 3 and one in Level 2a. In Level 4 remains 415 of a sized-3 ungulate, including a vertebra that presents a crenulated edge over the spinous process and a 416 metacarpal affected by pitting, were found. Carnivore traces were also found in Level 4 (a sized-3 417 ungulate mandible with two pits) and in Level 2a (a femur that presents a pit on the diaphysis and over 418 the edge of the fracture). The identification of marks inflicted by carnivores at the site impedes from 419 ruling out their contribution to the assemblage. However, the low carnivore modification rate suggests that carnivore activity was, in any case, moderate. 420

422 Post-depositional modifications are common in all levels. The most abundant are rounding and polished 423 (Table A.7), that affect 42.6% and 33.3% of the bone remains at VCS, respectively. Although the 424 abrasion, polish and rounding could be produced by plant roots, soil chemistry or hyena gnawing (Domínguez-Rodrigo et al., 2007), the high percentages observed in all levels, and the presence of micro-425 426 abrasion, indicate the action of water flows, especially in Levels 2a and 2. In Level 2a a rib fragment of a 427 size-5 individual presents a complete black coloration with a high degree of mineralization. Besides, these 428 remains present a high grade of subaerial weathering and a type of fossilization completely different to 429 the rest of the materials of the level. This might indicate that during the formation of the level, some 430 faunal remains were incorporated by the erosion of ancient beds and that these remains do not belong to 431 the original assemblage in Level 2a.

432

421

The Ostrich eggshell sample shows conchoidal fractures (Kandel, 2004; Kandel & Conard, 2005), unlike
natural processes. Ostrich (*Struthio camelus*) could be identified in Level 4 (*sensu* Dauphin et al. 1996).
After sieving, three different refittings have been identified in Level 4. *Struthio sp.* is present in Level 2a,
but proper species could not be identified. All eggshells of Level 2a are affected by water abrasion and
moderate polishing; whereas 78,9% of the sample in Level 4 shows signs of microabrasion and moderate
polishing. No heating changes have been identified in any level.

439

440 Several terrestrial (Limicolaria martensiana catharia, Succinea baumanni, Gittenedouardia cf. 441 hanningtoni, Euonyma percivali and Haplohelix cf. lateaperta) and freshwater mollusc species 442 (Biomphalaria pfeifferi and Bulinus tropicus/truncatus) have been identified in VCS (Fig. A.6 and Table 443 A.8). Only two of them have previously been reported in Olduvai stratigraphy: Limicolaria martensiana 444 catharia and Bulinus tropicus/truncatus (Rowson, 2017: 10). The mixture of two mollusc communities 445 indicates that transport or reworking of the material must have occurred, although the distances travelled 446 could have been short. Most of the terrestrial species reflect an open, sparsely vegetated area analogous to 447 the contemporary habitat in the Olduvai area. Large shrubs or scattered trees would have been present, 448 and the area was probably at least as well vegetated as today (Rowson, 2017: 11).

449 450 4.3.

450 4.3. Lithic Analysis.451

Lithic industry has been recovered from six levels at VCS. The lithic assemblage is not very abundant; it is composed of 99 pieces, eight of which are retouched and 9 are cores. The small sample size of the different levels does not allow a technological study of the collection (Table 4). Notwithstanding, the lithic sample provides important qualitative information to frame the occupation by Anatomically Modern Humans (AMH) in Olduvai Gorge.

457458 4.3.1. Taphonomic results:

459

460 The number of pieces in each level is very scarce, which can distort the rounding and pseudo-retouch 461 results. In broad terms, lithic tools found in every level present some rounding alteration, which is 462 consistent with the fluvial formation of the site. The levels most affected by rounding are level 2 with the 463 34.4% and level 2a with 100% of the remains. Which is no surprise due to their high energy fluvial 464 origin. Levels 4 and 5 present rounding alteration on some part of the assemblages, 32% and 50% 465 respectively, although the intensity of the rounding was lower than on levels 2 and 2a (Fig. A.7a). This 466 can be attributed to the low energy fluvial nature of the site. Level 5a do not present modifications. 467 However, this could be related with the due to the small number of remains recovered.

468

Pseudo-retouch affects 20 to 44% of the pieces in a low-intense way (Fig. A.7b). We deem the fluvial agent to be responsible for most of the rounding and pseudo-retouch of the pieces, and we should bear in mind—except for Level 2a that the energy of the altering agent and the displacement of the parts does not seem very intense.

473

474 4.3.2. Techno-typological results:475

476 From a techno-typological point of view, only general comments can be made because of the sample size:477

478 The raw materials used in the site are naibor quartzite (56%), basalt (29%), phonolite (14%), quartz (1%)
479 and chert (1%) (Fig. A.8).

- Naibor quartzite comes from the Naibor Soit, a Precambrian inselberg 3,5 km North to the site. This
 quartzite is almost entirely made up of crystallized quartz, it looks very much like mineral quartz and
 responds to knapping in a similar way (Rubio-Jara et al., 2017, pp. 4).
- 484

The basalt comes from Lemagrut volcano, 10-12 km South to VCS. The acquisition of this raw material
may have been obtained in palaeochannels from Lemagrut or from the Side Gorge, as in the lower Beds
of Olduvai Gorge (Jones, 1994).

488

The closest phonolite source to Olduvai Gorge originates in the Engelosen, 7 km North to VCS. It is a hill
made up of "flow banded nepheline phonolite lava, slightly porphyric, dark greenish grey" (Hay, 1976,
pp.12). It comes in slabs and, despite being of good quality for knapping, it could be affected by flow
banding (Jones, 1994, pp. 257).

493

494 Chert is of the highest quality and easy to find in the current riverbed, but it is appear as cobbles of
495 irregular and small morphology, which makes it exceedingly difficult to knap prepared methods
496 (Levallois/discoid).

497

498 Quartz is pretty scarce in the studied collection and can be obtained in the Olduvai riverbed itself. 499

Level 2: Level 2 contains the largest lithic assemblage at VCS: 32 pieces were recovered from this level. Four of them are cores, and four are retouched. Quartzite is the most common raw material (63%), followed by basalt (31%), and phonolite (6%) (Table 4 and Table A.9). Blanks are divided into cortical flakes, a tested block of raw material or ordinary flakes. There are seven discoid flakes (three of them being cordal and three having a centripetal direction) and three Levallois flakes, which have been obtained using different methods: preferential, bipolar and centripetal (Fig. 3: 2-8).

507 There are four retouched pieces (Table A.15): two denticulates (one of them in basalt and the other one in 508 quartzite), a notch in quartzite (Fig. 3: 7), and a heavy-duty piece (core-axe) in basalt (Fig. 3: 1). 509

The cores are two centripetal (not discoid), one unifacial in basalt, one bifacial in quartzite, one polyhedral in basalt, and another unipolar core (single-striking platform) in basalt, also indicating that blanks were obtained from not-prepared cores (Table A.16).

Level 2a: this level has 16 pieces—10 of the in basalt and the remaining six in quartzite (Table 4 and Table A.10). There are no retouched blanks but there is a core. Over half are ordinary flakes (9/16), but the assemblage also includes two discoid flakes (cordal and centripetal), and a preferential Levallois flake (Fig. 4: 6-8). The core is polyhedral-type and it was knapped in quartzite (Table A.16).

519 The retouched piece is a denticulate-endscraper in quartzite on an ordinary flake (Fig. 4: 8 and Table
520 A.15).
521

Level 3: A total of 15 pieces have been found in this level: 11 of are quartzite blanks and four were
knapped in basalt (Table 4 and Table A.11). Besides, there is an ochre fragment without evidence of use.
There is only one discoid flake with centripetal direction. The remaining blanks are ordinary flakes
(8/16), chunks and undetermined pieces. The only core present at the level is a unifacial discoid core in
quartzite (Table A.16).

Level 4: 23 stone tools have been recovered from Level 4; 65% of them are made in quartz, followed by phonolite (22%). There is also a piece of basalt, a piece of quartz, and another of chert (4% each). Four of the pieces clearly come from prepared cores: a *desbordant* flake in quartz, a preferential Levallois flake, and a *récurrent* bipolar flake, the latter two were knapped in phonolite (Fig. 4: 9-11). There is also a cordal flake, probably from a discoid core also in phonolite (Table 4 and Table A.12).

533

527

534 In addition, one of the three cores are Levallois preferential in quartzite and the remaining two are
535 opportunistic-type, where the natural angle ratio has been used to extract some flakes (Table A.16).
536

537 There are two retouched pieces (Table A.15). One of them is a notch in quartzite on an ordinary flake and
538 the other is a *desbordant* flake retouched on a *déjeté* sidescraper in quartz (Fig. 4: 11, 12).

Level 5: This level contains 10 pieces: four of them in basalt, three in phonolite and three in quartzite
(Table 4 and Table A.13). Seven of them are ordinary flakes, although there are two Levallois flakes—
one being recurrent unipolar and undetermined-type, in basalt and phonolite, respectively (Fig. 4: 4, 5).
There are no cores, and one of these pieces, the Levallois bipolar one, is retouched in simple inverse
sidescraper (Table A.15).

Level 5a: this level only has three pieces, all of them in basalt and obtained by using discoid methods, as
there are two flakes with a centripetal direction, and a pseudo-Levallois point, typical of this sort of
exploitation (Fig. 4: 1-3, Table 4 and Table A.14). No piece presents retouch.

550 4.4 Chronology

558

571 572

573

582

551 552 The quartz fractions extracted from the three samples collected for luminescence dating were insufficient 553 for the measurement of the natural doses and the test required to provide robustness to the method. In 554 addition, the few measurements carried out on the quartz fractions show that their optically stimulated 555 signal (OSL) is not dominated by the fast component (necessary requirement for blue-OSL dating). Dose 556 recovery tests show an overestimation of the given dose by more than 50%. According to these 557 observations, K-feldspar has been chosen as the best dosimeter in this case.

Dose recovery tests from feldspar using the pIR-IR225 protocol derive in a recovered to given dose ratio of 1.1±0.1 for given doses of 200 Gy. The fading tests yielded an average fading rate per decade (g-value) of 1.2±0.8 %. Such low fading rates have been suggested to be an artifact of the laboratory measurements (Thiel et al., 2011) and therefore no fading correction has been applied.

564 Dose distributions are well defined with average over-dispersion values below 25%. The low scatter in
565 the distribution suggests that these samples are not affected by incomplete bleaching.
566

Total dose rates, estimated burial doses and derived ages are summarized in Table 1. Results show that
deposition of these sediments occurred between 70 and 90 ka BP. This can be related to the Marine
Isotope Stage (MIS) 5a, associated to the first warm phase of this MIS.

5. Discussion.

The archaeological remains from VCS are too scarce to hold a deep discussion about the role of the site in
the regional MSA. Still, its study allows the formulation of some working hypotheses related to its
stratigraphic position and the palaeo-environment conditions, as well as to its lithic industry.

578 The geological record at VCS, dating back to around 75-86 ka BP, is a Lower Ndutu fluvial terrace 579 deposit (Uribelarrea et al., in this volume), just prior to Upper Ndutu sedimentation, which can be 580 approximately dated 90 ka BP. This is a slightly older chronology than that suggested by Hay (1976) and 581 far from those proposed by Manega (1993).

Based upon chronology, we place VCS within MIS 5a (90-75 ka BP). Bearing in mind that Olduvai Gorge is located in a transition zone between eastern Africa and tropical Africa, and that climate changes are asynchronous and regional in Africa (Bloome et al. 2012, Sturchio et al., 1993., Scholz et al., 2007), we should be cautious when talking about paleoclimatic conditions with the analytical conditions we currently have. The fauna found in the site does not provide any paleoclimatic information. Only the study of gastropods could indicate a plant covert? similar to the current one.

590 Cobo-Sánchez et al. (2014) proved that anisotropy can be detected in autochthonous assemblages under 591 certain conditions (e.g. lacustrine environments) and where no transport processes can be detected. 592 Orientation patterns were used to detect the autochthony of VCS' archaeological assemblages, as isotropy 593 is associated with the lack of post-depositional transport processes. According to our results, the 594 assemblage was transported, as the remains are significantly oriented, as expected from a fluvial context. 595

Also, the analysis of faunal remains found at VCS indicates that the assemblage was mainly the result of
water flow transport. Besides, lithic remains show evidences of intense fluvial transport at levels 2 and
and low fluvial transport at level 3 and 4. This supports the interpretation of faunal anisotropy due to
hydraulic flow on an autochthonous assemblage. The absence of anthropogenic modification on any sense

600 does not permit to related directly faunal and lithic remains. However, the predominance of rounding 601 indicate that they were deposited and transported by fluvial processes, although it should have happened 602 separately. Carnivore activity, although secondary, was the main biotic agent. This conclusion is in line 603 with the skeletal part representation, predominated by the presence of the axial skeleton, and the absence 604 of small fragments of bones, refitting and long bone epiphyses. This pattern also corresponds to 605 assemblages affected by hydraulic processes.

606 607

608

609 The MSA in northern Tanzania is defined by a number of industries from very few sites. It was proposed 610 by Mehlman (1989) and is synthesized in the Table 5 and Fig. 5. From a chronological point of view, 611 VCS would be coetaneous at the beginning of the Kisele industry (Mehlman, 1989). This industry is 612 identified from Mumba Unit VI and levels 12-17 and 18-25 from Nasera rockshelter. Chronologically, it 613 is located between 73/63 ka BP for Mumba VIa-dated by using OSL- and 56 ka BP for level 17 of Nasera-dated by using U-Th-(Mehlman, 1989, Gliganic et al., 2012) although Mehlman puts it off 614 615 until 90 ka BP (Mehlman, 1989, p. 560). Kisele Industry is characterized by a smaller industry than the 616 one before (Sanzako from Mumba VIb), by a high presence of retouched points and bifacial pieces, as 617 well as by prepared cores methods (Levallois/discoid) and a low presence of heavy duty and bipolar 618 methods (Mehlman, 1989).

619

The industry prior to Kisele in the local sequence of the MSA is Sanzako, and it is only located in Mumba
Unit VIb (Mehlman, 1989), and by using Th-230 the dating result is 131,710 (+6,924 -6,026) and Pa-231
measures to 109,486 (+44,404 -23,020) (Mehlman 1989, Brauer and Mehlman, 1988). This industry is
characterized by prepared cores methods, especially discoid, a low proportion of bipolar methods, points
and bifacial points, and a higher presence of domestic-type pieces: scrapers, notches and denticulates
(Solano-Megías, 2018).

627 Although, the lithic industry found in VCS is very scarce and makes it impossible to compare it directly 628 with industries which are sort of synchronous. From a general point of view, VCS levels are characterised 629 by the flake production from prepared core methods (Levallois and particularly discoid), the absence of 630 bipolar methods and points of any type, and a limited presence of retouched blanks-being the existing 631 ones domestic-type (sidescrapers, denticulates and notches). Heavy duty is also present. Besides, the 632 overall percentage of retouched pieces is 8%, although in levels with retouched pieces (Levels 2a, 2, 4 633 and 5) it ranges from 6.25% to 17.3%. Thus, as a result of the data presented, VCS should be related to 634 the beginning of the Kisele industry. The site with the closest chronology to VCS is Unit VIa from 635 Mumba rockshelter, which approximately dates from 73-63 ka by using OSL (Mehlman, 1989, Gliganic 636 et al., 2012). However-and stressing again how flimsy the collection is-from a qualitative point of 637 view, VCS seems to be more akin to the oldest industry (Sanzako) than to its contemporary (Kisele), even 638 though this conclusion is tentative until the larger area of the site is excavated. Under no circumstances 639 can VCS be associated with the Sangoan/Early Middle Stone Age of Lake Eyasi, as was the case in the 640 past for Olduvai surface collections (Leakey et al, 1972, Eren et al. 2014). As a result, we consider the 641 above-ground evidence of MSA in Olduvai Gorge to correspond to several temporary moments rather 642 than to a number of Early MSA assemblages. 643

644 6. Conclusions.

645
646 Victoria Cabrera Site is the first site which clearly belongs to the Middle Stone Age with a stratigraphic
647 and chronological control in Olduvai Gorge. Six levels from the site—dating back between 75 to 86 ka
648 BP—contain lithic industry which, despite being small-numbered, is characteristic for prepared core
649 methods, especially discoid and Levallois, with some "domestic-type" retouched pieces (sidescrapers,
650 denticulates and notches), scarce heavy-duty, and a total absence of points. Lithic industry and fauna have
651 been found together, though the association between them is not clear yet.

652

The site corresponds to MIS 5a, a period of climatic favourable conditions. However, neither the paleoenvironment features of that time—as Olduvai Gorge is on the border between eastern Africa and tropical
Africa characteristics— nor the asynchronous and local climate changes in the area (Blome et al. 2012)
can be characterised in detail yet.

657

Only a few sites that belong to the MIS5 period have been found in East Africa (Mirazón-Lahr and Foley,
2016, pp. 223). One explanation might be related to the absence of investigation and dating of deposits.

The site of VCS adds to the list of other sites with levels corresponding to MIS5, such as Mumba, with
 the expectation that more knowledge about the human presence, patterns of settlement, mobility of human
 groups, and interaction with the ecosystem in Olduvai Gorge and northern Tanzania can be gained.

Acknowledgments.

To Victoria Cabrera-Valdés (1951-2004), In Memoriam.

669 This study has been conducted with the funding of the project HAR2015-64407-P MINECO/FEDER UE 670 of the Spanish Ministry of Science, Innovation and Universities, Fundación PALARQ and Instituto de 671 Patrimonio Cultural de España. Juan Marín is the beneficiary of an Erasmus Mundus Doctoral 672 scholarship for an International Doctorate in Quaternary and Prehistory. We want to thank COSTECH (Tanzania Commission for Science and Technology) for the permission to carry out the work and the 673 674 Department of Antiquities of Tanzania for its support. We want to thank to Museum of Olduvai, 675 especially to its director Ladislaus Kashaija Domician and Mr. Masaki, its curator. We want to thank to TOPPP team, especially Manuel Domínguez-Rodrigo, Enrique Baquedano, Fernando Díez-Martín and 676 677 his team, Elia Organista, Lucía Cobo, Manuel Santonja and his team, Alfredo Pérez, Joaquín Panera and 678 Susana Rubio. Also, we thank to José Yravedra and Mari Carmen Arriaza their support, advices and help. Finally, we want to thank the two anonymous reviewers for their suggestions and comments on the 679 680 manuscript and Andrew Whitehead for proofreading. 681

682683 References

684

697

701

664 665

666 667

668

685 Alcalá, L., 1994. Macromamíferos neógenos de la fosa Alfambra-Teruel. Instituto de Estudios
686 Turolenses, Zaragoza.
687

Auclair, M., Lamothe, M., and Huot, S., 2003. Measurement of anomalous fading for feldspar IRSL using
SAR. Radiat. Meas. 37, 4, 487-492. <u>https://doi.org/10.1016/S1350-4487(03)00018-0</u>

Audouze, F., Karlin, C., 2017. La chaîne opératoire a 70 ans: qu'en ont fait les préhistoriens français.
Journal of Lithic Studies 4 (2), 1-69 doi:10.2218/jls.v4i2.2539

Ashley, G.M., Bunn, H.T., Delaney, J., Barboni, D., Dominguez-Rodrigo, M., Mabulla, A., Gurtov, A.,
Baluyot, R., Beverly, E., Baquedano, E., 2014. Paleoclimatic and paleoenvironmental framework of FLK
North archaeological site, Olduvai Gorge, Tanzania. Quaternary International 322-323, 54-65.

Barboni, D., Ashley, G., Domínguez-Rodrigo, M., Bunn, H. T., Mabulla, A. Z., Baquedano, E., 2010.
Phytoliths infer locally dense and heterogeneous paleovegetation at FLK North and surrounding localities
during upper Bed I time, Olduvai Gorge, Tanzania. Quaternary Research 74, 344-354.

Basell, L., 2013. The Middle Stone Age of Eastern Africa. In: Mitchel, P., Lane, P. (Eds), *The Oxford Handbook of African Archaeology*. Oxford University Press, Oxford, pp. 387-401.

704
705 Behrensmeyer, A. K., 1982. Time resolution in fluvial vertebrate assemblages. Paleobiology 8(3), 211706 227.

Behrensmeyer, A.K. 1978. Taphonomic and ecologic information from bone weathering. Paleobiology 4, 150–162.

710

707

Binford, L.R. 1981. Bones: Ancient Men and Modern Myths. Academic Press, New York.

Blome, M.W., Cohen, A.S., Tryon, T.A., Brooks, A.S., Russell, J., 2012. The environmental context for
the origins of modern human diversity: a synthesis of regional variability in African climate 150,00030,000 years ago. Journal of Human Evolution 62, 563-592.

716

Boëda, E., 1995. Caractéristiques techniques des chaînes opératoires lithiques des niveaux micoquiens de
 Külna (Tchécoslovaquie). Actes du colloque Les industries à pointes foliacées d'Europe Centrale,

719 Miscolk 1991. Paléo, Supplément 1, 57-72.

Boëda F

720

733

736

Boëda, E., 1994. Le Concept Levallois: variabilité des méthodes. Monographie du C.R.A., Paris.

- 723 Boëda, E., 1993. Le débitage discoide et le débitage levallois récurrent centripéte. Bulletin de la Société
 724 Préhistorique Francaise 90 (6), 392-404.
 725
- Boëda, E., Geneste, J.M., Meignen, L., 1990. Identification de chaînes opératoires lithiques du
 Paléolithique ancien et moyen. Paléo 2, 43-80.
- Bower, J. R. F., Mabulla, A., 2013. Setting In: Evidence of Territorial Exclusion in the Late Middle Stone
 Age of Nothern Tanzania. Studies in the African Past 11, 7-24.
- 732 Brain, C. K., 1981. The Hunters or the Hunted?. University of Chicago Press, Chicago.
- Capaldo, S. D., 1998. Methods, marks, and models for inferring hominid and carnivore behavior. Journal
 of Human Evolution 35(3), 317-320.
- Clark J. D., 1988. The Middle Stone Age of East Africa and the Beginnings of Regional Identity. Journal of World Prehistory 2 (3), 235-303.
- Cobo-Sánchez, L., Aramendi, J., Domígnuez-Rodrigo, M., 2014. Orientation patterns of wildebeest bones
 on the lake Masek floodplain (Serengeti, Tanzania) and their relevance to interpret anisotropy in the
 Olduvai lacustrine floodplain. Quaternary International 322-323, 277-284.
- Collins, B., Steele, T. E., 2017. An often overlooked resource: Ostrich (*Struthio sp.*) eggshell in the
 archaeological record. Journal of Archaeological Science: Reports 13, 121-131.
- 746
 747 Core R-Team, 2013. A Language and Environment for Statistical Computing. R Foundtion for Statistical Computing, Vienna, Austria URL <u>http://www.Rproject.org/</u>.
 749
- Courtin, J., Villa, P., 1982. Une expérience de piétinement. Bulletin de la Société Préhistorique Francaise
 79 (4), 117-123.
- 752
 753 Dauphin, Y., Pickford, M., Senut, B., 1996. Microstructures des coquilles d'oeufs d'oiseaux fossiles de
 754 Namibie. Revue de Paléobiologie 15 (1), 225-241.
- Domínguez-Rodrigo, M., Cobo-Sánchez, L., Yravedra, J., Uribelarrea, D., Arriaza, C., Organista, E.,
 Baquedano, E., 2017. Fluvial spatial taphonomy: a new method for the study of post-depositional
 processes. Archaeological and Anthropological Sciences 10 (7), 1-21.
- Domínguez-Rodrigo, M., Martinez-Navarro, B., 2012. Taphonomic analysis of the early Pleistocene (2.4
 Ma) faunal assemblage from AL 894 (Hadar, Ethiopia). Journal of Human Evolution 62 (3), 315-327.
- Domínguez-Rodrigo, M., Mabulla, A., Bunn, H. T., Barba, R., Diez-Martín, F., Egeland, C. P., Espilez,
 E., Egeland, A., Yravedra, J., Sánchez, P., 2013a. Unravelling hominin behaviour at another
 anthropogenic site from Olduvai Gorge (Tanzania): new archaeological and taphonomic research at BK,
 Upper Bed II. Journal of Human Evolution 57, 260-283.
- 767
- Domínguez-Rodrigo, M., Garcia-Pérez, A., 2013b. Testing the accuracy of different A-axis types for
 measuring the orientation of bones in the archaeological and paleontological record. PLoS ONE 8(7),
 e68955.
- Domíngez-Rodrigo, M., Bunn, H., Pickering, T.R., Mabulla, A., Musiba, C.M., Baquedano, E., Ashley,
 G. M., Diez-Martin, F., Santonja, M., Uribelarrea, D., Barba, R., Yravedra, J., Barboni, D., Arriaza, C.,
 Gidna, A., 2012. Autochthony and orientation patterns in Olduvai Bed I: are examination of the status of
 post-depositional biasing of archaeological assemblages from FLK North (FLKN). Journal of
 Archaelogical Science 39, 2116-2127.
- 777

- Domínguez-Rodrigo, M., Bunn, H. T., Mabulla, A. Z., Baquedano, E., Pickering, E. T., 2010.
 Paleoecology and hominin behavior during Bed I at Olduvai Gorge (Tanzania). Quaternary Research 74, 301-303.
 Domínguez-Rodrigo, M., De Juana, S., Galán, A.B., Rodríguez, M., 2009. A new protocol to differentiate trampling marks from butchery cut marks. Journal of Archaeological Science 36 (12), 2643–2654
- 783 trampling marks
- 785 Domínguez-Rodrigo, M., Barba, R., Egeland, C. P., 2007. Desconstructing Olduvai: a taphonomic study
 786 of the Bed I sites. Springer, New York.
- 787

Durcan, J.A., King, G.E., Duller, G.A.T., 2015. DRAC: Dose rate and age calculator for trapped charge dating. Quaternary Geochronology 28, 54-61.

790

800

802

805

813

821

Fren, M., Durant, M.E., Prendergast, M., Mabulla, A.Z.P., 2014. Middle Stone Age archaeology at
Olduvai Gorge, Tanzania. Quaternary International 322-323, 292-313.

- Fernández-Jalvo, Y., Andrews, P., 2003. Experimental effects of water abrasion on bone fragments.
 Journal of Taphonomy 1, 147–163.
- Fiorillo, A.R., 1991. Taphonomy and depositional setting of Careless Creek Quarry (Judith River
 formation), Wheatland County, Montana, U.S.A. Palaeogeography, Palaeoclimatology, Palaeoecology
 81, 281–311.
- 801 Fisher, N.I., 1995. Statistical Analysis of Circular Data. Cambridge University Press, Cambridge.
- Frostick, L., Reid, I., 1983. Taphonomic significance of sub-aereal transport of vertebrate fossils on steep
 semi-arid slopes. Lethaia 16, 157-164.
- Galbraith, R.F., Roberts, R.G., Laslett, G.M., Yoshida, H., Olley, J.M., 1999. Optical dating of single and
 multiple grains of quartz from Jinmium rock shelter, Northern Australia: part 1, experimental design and
 statistical models. Archaeometry 41, 339-364.
- 810 Geneste, J-M. 1985. Analyse lithique d'industries moustériennes du Périgord: une approche
 811 technologique du comportement des groupes humains au Paléolithique moyen. Ph.D. Thesis, Université
 812 de Bordeaux I, France.
- 814 Gifford-González, D. P., Damrosch, D. B., Damrosch, D. R., Pryor, J., Thunen, R. L. 1985. The third dimension in site structure: an experiment in trampling and vertical dispersal. American Antiquity 50 (4), 803-818.
 817
- Gliganic, L. A., Jacobs, Z., Roberts, R.G., Domínguez-Rodrigo, M., Mabulla, A.Z.P., 2012. New ages for
 Middle and Later Stone Age deposits at Mumba rockshelter, Tanzania: Optically stimulated luminescence
 dating of quartz and feldspar grains. Journal of Human Evolution 62 (4), 533-547.
- Hay, R., 1987. Geology of the Laetoli are. In: Leakey, M. D., Harris, J. M. (Eds.), Laetoli: A Pliocene site
 in Northern Tanzania. Claredon Press, Oxford, pp. 23-47.
- Hay, R., 1976. The Geology of Olduvai Gorge. University of California Press, Berkeley.
- 826827 Howard, P., 2007. Archaeological survey and mapping. Taylor and Francis, London.
- 828
 829 Hughes, P. J., Lampert, R. J., 1977. Occupational disturbance and types of archaeological deposits.
 830 Journal of archaeological Science 4, 135-140.
 831
- 832 Inizan, M-L., Reduron, M., Roche, H., Tixier, J., 1995. Technologie de la Pierre taillée. C.R.E.P., Paris.

Jones, P. R., 1994. Results of experimental work in relation to the Stone industries of Olduvai Gorge. In:
Leakey, M. (Ed.), Olduvai Gorge. 5. Excavations in Beds III, IV and the Masek Beds, 1968–1979.
Cambridge University Press, Cambridge, pp. 254-298.

837

- Kandel, A.W., 2004. Modification of ostrich eggs by carnivores and its bearing on the interpretation of
 archaeological and paleontological finds. Journal of Archaeological Science 31, 377-391.
- Kandel, A.W., Conard, N.J., 2005. Production sequences of ostrich eggshell beads and settlement
 dynamics in the Geelbek Dunes of the Western Cape, South Africa. Journal of Archaeological Science
 32, 1711-1721.
- Karlin, C., 1992. Connaissances et savoir-faire: comment analyser un processus technique en Préhistoire.
 Introduction. In: R. Mora, X. Terradas, A. Parpal, C. Plana (Eds.), Tecnología y Cadenas operativas
 líticas. Treballs d'Arqueología, Barcelona, pp. 99-124.
- 848

865

844

- Karlin, C., Bodu, P., Pelegrin, J., 1992. Processus techniques et chaînes opératoires: comment les préhistoriens s'appropient un concept élaboré par les ethnologues. In : Balfet, P. (Ed.), Observer l'action technique: des chaînes opératoires, pour quoi faire? CNRS, Paris, pp.101-117.
- Lam, Y. M., Chen, X., Pearson, O.M., 1999. Intertaxonomic variability in patterns of bone density and
 the differential representation of bovid, cervid, and equid elements in the archaeological record. American
 Antiquity 64 (2): 343–362.
- 856 857 Leskey M. Roe
- Leakey, M., Roe, D. A., 1994. Olduvai Gorge. 5. Excavations in Beds III, IV and the Masek Beds, 1968–
 1979. Cambridge University Press, Cambridge.
- Leakey, M., Hay, R., Thurber, D., Protsch, R., Berger, R., 1972. Stratigraphy, archaeology, and age of the
 Ndutu and Naisiusu Beds, Olduvai Gorge, Tanzania. World Archaeology 3, 328-341.
- Leakey, M. D., 1971. Olduvai Gorge vol 3. Excavations in Beds I & II, 1960-1963. Cambridge
 University Press, Cambridge.
- Levi-Sala, I. 1986. Use wear and post-depositional surface modifications: a word of caution. Journal of
 Archaeological Science 13, 229-244.
- Mabulla, A., 1990. Preliminary report on an archaeological survey of the Ndutu Beds, Olduvai Gorge,
 Tanzania. Nyame Akuma 33, 20-25.
- 872 Maíllo-Fernández, J. M. 2003. La transición Paleolítico Medio-Superior en Cantabria: análisis
 873 tecnológico de la industria lítica de Cueva Morín. Ph.D. Thesis, UNED, Spain.
 874
- Manega, P., 1993. Geochronology, Geochemistry, and Isotopic Study of the Plio-pleistocene Hominid
 Sites and the Ngorongoro Volcanic Highland in Northern Tanzania. Ph.D. Thesis, University of Colorado
 at Boulder, USA.
- Marean, C.W., Cleghorn, N., 2003. Large mammal skeletal element transport. Applying foraging theory
 in a complex taphonomic system. Journal of Taphonomy 1, 15–42
- Miall, A. D., 1985. Architectural-element analysis: a new method of facies analysis applied to fluvial
 deposits. Earth-Science Reviews 22(4), 261-308.
- McBrearty, S., Brooks, A., 2000. The revolution that wasn't: A new interpretation of the origin of modern
 human behavior. Journal of Human Evolution 39 (5), 453-563.
- McBrearty, S., Bishop. L. Plummer, T., Dewar, R., Conard, N., 1998. Tools underfoot: human trampling
 as an agent of lithic artifacts edge damage. American Antiquity 63 (1), 108-129.
- Melhman, M. J., 1989. Late Quaternary Archaeological Sequences in Northern Tanzania. Ph. D. Thesis,
 University of Illinois, USA.
- Mirazón-Lahr, M., Foley, R. A., 2016. Human Evolution in Late Quaternary Eastern Africa. In: Jones, S.
 C., Stewart, B. A. (Eds.), Africa from MIS 6-2: Population Dynamics and Paleoenvironments. Springer, New York, pp. 215-231.
- 897

884

- Mourre, V., 2003. Discoïde ou pas Discoïde ? Réflexions sur la pertinence des critères techniques
 définissant le débitage Discoïde. In: Peresani, M. (Ed.), Discoid lithic technology. Advances and
 implications. BAR International Series 1120, Oxford, pp. 19-32.
- 901

905

902 Organista, E., Domínguez-Rodrígo, M., Egeland, C.P., Uribelarrea, D., Mabulla, A., Baquedano, E.,
903 2016. Did *Homo erectus* kill a *Pelorovis* herd at BK (Olduvai Gorge)? A taphonomic study of BK5.
904 Archaeological and Anthropological Science, 8, 601-624.

Organista, E., Domínguez-Rodrigo, M., Yravedra, J., Uribelarrea, J., Arriaza, M. C., Ortega, M.C.,
Mabulla, A., Gidna, A., Baquedano, E., 2017. Biotic and abiotic processes affecting the formation of BK
Level 4c (Bed II, Olduvai Gorge) and their bearing on hominin behavior at the site. Palaeogeography,
Palaeoclimatology, Palaeoecology 488, 59-75.

910

911 Pedraza, J., 1996. Geomorfología. Principios, Métodos y Aplicaciones. Editorial Rueda, Madrid.
 912

Pickering, T. R., Egeland, C. P., 2006. Experimental patterns of hammerstone percussion damage on
bones and zooarchaeological inferences of carcass processing intensity by humans. Journal of
Archaeological Science 33, 459–469

Reiner, W. B., Masao, F., Sholts, S. B., Songita, A. V., Stanistreet, I., Stollhofen, H., Taylor, R. E.,
Hlusko, L. J., 2017. OH 83: A new early modern human fossil cranium from the Ndutu beds of Olduvai
Gorge, Tanzania. American Journal of Physical Anthropology 164 (3), 533-545.

921 Rightmire, G. P., 1980. Middle Pleistocene hominids from Olduvai Gorge, northern Tanzania. American
922 Journal of Physical Anthropology 53, 225–241.
923

- 924 Rowson, B., 2017. Terrestrial and freshwater Mollusca (Gastropoda Pulmonata) from a Middle Stone
 925 Age excavation in Olduvai Gorge, Tanzania. Unpublished report, 12 pages.
- Rubio-Jara, S., Panera, J., Snatonja, M., Pérez-González, A., Yravedra, J., Domínguez-Rodrigo, M.,
 Bello, P., Rojas, R., Mabulla, A., Baquedano, E., 2017. Site function and lithic technology in the
 Acheulean technocomplex: a case study from Thiongo Korongo (TK), Bed II, Olduvai Gorge, Tanzania.
 Boreas. DOI 10.1111/bor.12275
- 932 Schick, K., 1984. Towards the interpretation of early archaeological sites: Experiments in site formation
 933 processes. Ph.D. Thesis, University of California, USA.
- 934

926

931

Scholz, Ch., A., Johnson, T. C., Cohen, A. S., King, J. W., Peck, J. A., Overpeck, J. T., Talbot, M. R.,
Brown, E. T., Kalindekafe, L., Amoako, P. Y. O., Lyons, R. P., Shanahan, T. M., Castañeda, I. S., Heil,
C. W., Forman, S. L., McHargue, L. R., Beuning, K. R., Gomez, J., Pierson, J., 2007. East African
megadroughts between 135 and 75 thousand years ago and bearing on early-modern human origins.
PNAS 104 (42), 16416-16421

- 941 Skinner, A., Kay, R., Masao, F., Blackwell, B., 2003. Dating the Naisiusiu Beds, Olduvai Gorge, by
 942 electron spin resonance. Quaternary Science Reviews 22, 1361-1366
 943
- 944 Shackley, M. L., 1978. The behavior of artefacts as sedimentary particles. Archaeometry 20 (1), 55-61.
- 945

946 Solano-Megías, I., 2018. El tecnocomplejo Sanzako: estudio tecnotipológico del nivel VI-B de Mumba
947 (Tanzania). MA Thesis. University Rovira I Virgili, Spain.

948

949 Stapert, D., 1976. Some natural surface modifications on flint in the Netherlands. Palaeohistoria XVIII, 7950 41.
951

Sturchio, N. C., Dunkley, P. N., Smith, M., 1993. Climate-driven variations in geotermal activity in the
 northern Kenya rift valley. Nature 362: 233-234.

954

955 Terradas, X. 2003. Discoid flaking method: Conception and technological variability. In: Peresani, M.
956 (Ed.), Discoid lithic technology. Advances and implications. BAR International Series 1120, Oxford, pp.
957 19-31.

- Thiel, C., Buylaert, J.-P., Murray, A.S., Terhorst, B., Hofer, I., Tsukamoto, S., Frechen, M., 2011.
 Luminescence dating of the Stratzing loess profile (Austria) testing the potential of an elevated temperature post-IR IRSL protocol. Quat. Int. 234, 23-31.
- 963 Thomsen, K.J., Murray, A.S., Jain, M., and Bøtter-Jensen, L., 2008. Laboratory fading rates of various
 964 luminescence signals from feldspar-rich sediment extracts. Radiat. Meas. 43, 9, 1474-1486.
 965 https://doi.org/10.1016/j.radmeas.2008.06.002.
 966
- 967 Tringham, R. Cooper. G. Odell G., Voytex, B., Whitman, A., 1974. Experimentation in the formation of
 968 edge damage: a new approach to lithic analysis. Journal of Field Archaeology 1, 171-196.

- 971 Tukey, J.W., 1977. Exploratory Data Analysis. Addison Wesley, Reading, Mass.972
- 973 Uribelarrea del Val, D., Martín-Perea, D. M., Marín, J., Solano-Megías, I., Barba Egido, R., Baquedano,
 974 E., Mabulla, A., Domínguez-Rodrigo, M., Gidna, A., Maíllo-Fernández, J. M. This volumen.
 975 Geomorphological reconstruction of VCS, a Middle Stone Age site in the terrace system of Lower Ndutu,
 976 Olduvai Gorge (Tanzania). Quaternary International.
 977
- 978 Uribelarrea, D., 2014. Geología y geomorfología de Olduvai y Laetoli. In: Domínguez, Rodrigo, M.;
 979 Baquedano; E., (Eds), La cuna de la Humanidad, vol II. Museo Arqueológico Regional, Madrid, pp 120-135.
- Uribelarrea, D., Domínguez-Rodrigo, M. Pérez-González, A., Vegas-Salamanca, J., Baquedano, E.,
 Mabulla, A. Z. P., Musiba C., Barboni, D. and Cobo-Sánchez, L., 2014. Geo-archaeological and
 geometrically corrected reconstruction of the 1.84 Ma FLK Zinj paleolandscape at Olduvai Gorge,
 Tanzania. Quaternary International 322-323:7-31
- 987 Von Zieten, R. R. R., 2009. Preliminary report on a new Olduvai hominid from 1968–an "anatomically988 modern-man" from Upper Ndutu Bed. Retrieved from www1.uni-hamburg.de/helmut-ziegert.
 989
- 990 Voorhies, M., 1969. Taphonomy and population dynamics of an early Pliocene vertebrate fauna, Knox
 991 County, Nebraska. Contributions to Geology Special Paper No. 1, University of Wyoming, Laramie.
 992
- Woodcock, N.H., 1977. Specification of fabric shapes using an eigen value method. Geological Society
 of America Bulletin 88 1231–1236.

ACCEPTED MANUSCRIPT
CAPTIONS
CALITONS
Table 1: Summary of luminescence dating from VCS including the depth and water content used for calculation of the environmental dose rate. A 5% error has been added to the assumed water content. Estimated equivalent doses derived from IRSL measurements and corresponding burial ages, expressed in kiloyears (ka).
Table 2. Omnibus statistical tasts applied to the different archaeological levels of VCS, divided by bones
and lithics
Table 3. NISP, MNE and MNI by taxa and weights size category at archaeological levels of VCS.
Table 4. Blanks' chaîne opératoire inventory by levels at VCS.
Table 5. Northern Tanzania MSA's sites cited in the text.
Figure 1. a) Map of Africa; b) Map of Tanzania; c) Map of Olduvai Gorge; d) Location of Victoria Cabrera site (VCS). ©google maps.
Figure 2. Figure 2. 1. Victoria Cabrera Site Trenches 1 and 2 stratigraphic profiles. 2. Schematic stratigraphical correlation between Victoria Cabrera Site Trench 1, Trench 2 South Profile and Trench 2 West Profile. 3. South Profile picture.
Figure 3. Lithics from Level 2: 1. Heavy-duty (core-axe) on basalt, 2-4. Cordal flakes on quartzite, 5. Centripetal discoid flake on basalt, 6. Recurrent Levallois flake on basalt, 7. Preferential Levallois flake on quartzite (notch), 8. Bipolar Levallois flake on quartzite.
 Figura 4. Lithics from VCS: 1-3: Level 5a; 4-5: level 5; 6-8: level 2a; 9-12: Level 4. 1. Centripetal discoid flake on basalt, 2. Pseudolevallois point on basalt, 3. Centripetal discoid flake on basalt, 4. Unipolar Recurrent Levallois on basalt, 5. Ordinary flake on basalt, 6. Preferential Levallois flake on basalt, 7. Centripetal discoid flake on basalt., 8. Ordinary flake on Quartzite (denticulate), 9. Preferential Levallois flake on basalt, 10. Bipolar levallois flake on basalt, 11. Desbordant flake on quarzite (sidescraper), 12. Ordinary flake on quartzite (notch). Figure 5. Map of Northern Tanzania with the major sites cited in text (source NASA).

Appendix A.

VICTORIA CABRERA SITE: A MIDDLE STONE AGE SITE AT OLDUVAI GORGE, TANZANIA.

José-Manuel Maíllo-Fernández^{1, 2}, Juan Marín³, Irene Solano-Megías¹, David Uribelarrea^{2,4}, David Martín-Perea^{2,5}, Julia Aramendi^{2,6}, Alicia Medialdea⁷, Carlos Arteaga⁸, Marta Pernas-Hernández⁶, Agness Gidna⁹, Neogi Sayantani¹⁰, Eva Baudot⁸, Carlos Narváez⁸, Audax Mabulla⁹.

¹ Dpt. Prehistory and Archaeology, UNED, Spain.

² IDEA, The Institute of Human Evolution in Africa, Madrid, Spain.

³ Muséum National d'Histoire Naturelle (MNHN), France.

⁴ Dpto. de Geodinámica, Estratigrafía y Paleontología, UCM. Spain.

⁵ Dpto. de Paleobiología (MNCN), Madrid, Spain.

⁶ Dpt. Geographisches Institut, University of Kölh, Germany.

- ⁷ Dpto. de Prehistoria, Historia Antigua y Arqueología, UCM, Spain.
- ⁸ Laboratorio de Geografía Física, UAM, Spain.
- ⁹ National Museum of Tanzania, Tanzania.

¹⁰ Department of Archaeology, Durham University, UK.



Figure A.1. a) Rose diagram showing lithic orientation; b) Stereogram showing the azimuth orientation of lithic specimens with a longitudinal axis from BK5. The first eigenvalue comprising most of the inertia is presented here, as well as the confidence interval of the mean trend/plunge orientation (in red); c) Woodcock diagram with von Mises distribution K.



Figure A.2. a) Rose diagram showing bones orientation; b) Stereogram showing the azimuth orientation of lithic specimens with a longitudinal axis from BK5. The first eigenvalue comprising most of the inertia is presented here, as well as the confidence interval of the mean trend/plunge orientation (in red); c) Woodcock diagram with von Mises distribution K.



Figure A.3. Ripley's K function for lithic assemblages.



Figure A.4. Ripley's K function for bone assemblages.

NISP (MNE)	Size 5			
	Level 2	Level 3	Level 4	Level 2a
Isolated teeth	1 (-)			
Rib	1 (1)	2 (1)		1 (1)
Flat			1 (-)	

Table A.1. NISP and MNE by anatomical element for Size 5 animals.

NISP (MNE)	Size 4				
	Level 2	Level 3	Level 4	Level 2a	Level 5
Cranium	1 (1)				
Mandible		-2			V
Isolated teeth		6 (-)		1 (-)) 7
Vertebrae		1 (1)	2 (2)		
Rib			1 (1)	S	
Metacarpal				1 (1)	
Tibiae		1 (1)	X		
Metatarsal		1 (1)			
Long	1 (-)	2 (-)	1 (-)	Y	
Flat	3 (-)		2 (-)	1 (-)	1 (-)

Table A.2. NISP and MNE by anatomical element for Size 4 animals.

NISP (MNE)	Size 3				
	Level 2	Level 3	Level 4	Level 2a	Level 5
Cranium		3 (1)	1 (1)		1(1)
Mandible	1 (1)	1 (1)	1 (1)	1 (1)	
Isolated teeth					1 (-)
Rib		2(1)	1 (1)		1 (1)
Scapula		1 (1)			
Соха		1 (1)			
Radius		1 (1)		A	
Ulna		1 (1)			
Metacarpal	1 (1)		2 (1)		
Femur					1(1)
Long	2 (-)	3 (-)	4 (-)	2 (-)	
Flat	1 (-)	4 (-)	4 (-)	1 (-)	

Table A.3. NISP and MNE by anatomical element for Size 3 animals.

NISP (MNE)	Size 2			
	Level 3	Level 4	Level 2a	Level 5a
Vertebrae		1 (1)		
Rib				1 (1)
Scapula	1 (1)	1 (1)		
Соха		1 (1)		
Radius			1 (1)	
Metacarpal	1 (1)	1 (1)		
Femur	1 (1)			
Astragalus		1 (1)		
Phalanx		1 (1)		
Long	6 (-)	2 (-)	2 (-)	

Table A.4. NISP and MNE by anatomical element for Size 2 animals.

NISP / %NISP	Level 2	Level 3	Level 4	Level 2a	Level 5a	Level 5
Axial cranial skeleton	3/50%	10/41.7%	2/14.3%	2/40%	2/50%	1/50%
Axial postcraneal skeleton	2/33.3%	8/33.3%	7/50%	1/20%	1/25%	1/50%
Appendicular bones	1/16.7%	6/25%	5/35.7%	2/40%	1/25%	0

Table A.5. NISP and % NISP distribution by anatomicaL segment.

Level	Size category	Spearman's r	р
Level 2	Size 3	-0.73259	0.10714
Level 3	Size 4	-0.25198	0.64286
	Size 3	-0.53611	0.19048
Level 4	Size 4	-0.51437	0.28571
	Size 3	-0.57735	0.25
Level 2a	Size 4	-0.57735	0.25
	Size 3	-0.15121	0.7381

Table A.6. Spearman's statistics correlation between high survival elements by level and size category and (S)FUI.



Figure A.5 %NISP distribution in function of length sizes.

NR	1	1	1	1	Laural Era	
%NR	Level 2	Level 3	Level 4	Level 2a	vei za Level 5a	
Dounding	16/26	21/45	20/50	10/19	2/4	0/3
Rounding	69.5%	46.7%	40%	77%	50%	0
Deliched	13/26	19/45	10/50	9/19	3/5	0/3
Polisned	56.5%	42.2%	20%	69.2%	60%	0
Ficcure	2/26	12/45	5/50	8/19	0/5	0/3
rissure	8.7%	26.6%	10%	61.5%	0	0
Concretions	5/26	5/45	5/50	1/19	2/5	1/3
Concretions	21.7%	11.1%	10%	7.6%	40%	33.3%
Trampling	0/26	7/45	2/50	0/19	0/5	0/3
rrampling	0	15.5%	4%	0	0	0

Table A.7. NISP (NISP%) by archaeological level and postdepositional modification.

		NMI					
	Level 2	Level 2a	Level 4	Level 5	Level 5a	NMI	%
pfeifferi	2		107			109	60
tropicus/truncatus			10		1	11	6
	2		117	~	1	120	66
				S			
				Y			
		1					
martensiana catharia	1	2	46	3	1	53	29
baumanni			5			5	3
cf. hanningtoni			2			2	1
percivali			1			1	1
cf. lateaperta			1			1	1
	1	2	55	3	1	62	34
		2	172	2	2	102	100
	pfeifferi tropicus/truncatus martensiana catharia baumanni cf. hanningtoni percivali cf. lateaperta	pfeifferi 2 tropicus/truncatus 2 2 martensiana catharia 1 baumanni cf. hanningtoni percivali cf. lateaperta 1 3	pfeifferi2tropicus/truncatus2222112martensiana catharia122baumanni1cf. hanningtoni2percivali2cf. lateaperta11232	pfeifferi 2 107 tropicus/truncatus 10 2 117 2 117 1 2 5 5 3 2	pfeifferi 2 107 tropicus/truncatus 10 2 117 2 117 2 117 2 117 10 1 2 117 10 1 2 117 1 2 1 2 1 2 46 3 baumanni 5 cf. hanningtoni 2 1 2 1 2 1 2 55 3 3 2	Image: state of the state	Image: state of the state

Table A.8. Material identified, minimum numbers of individuals, and percentage occurrence.



Figure A.6. Representative shells from VCS: **A**, *Biomphalaria pfeifferi*; **B**, *Bulinus truncatus/tropicus*; **C**, *Succinea baumanni*; **D**, *Gittenedouardia* cf. *hanningtoni*; **E**, *Euonyma percivali*; **F**, *Limicolaria martensiana catharia*; **G**, *?Haplohelix* cf. *lateaperta*. All scalebars 5 mm. All shown to scale, except for F (shown at 0.5x scale) and enlargements of sculpture for B and G (Rowson 2017).



VCS - Rounding

VCS - Pseudo-retouch



Figure A.7. Lithic alterations at VCS: a) rounding, b) pseudo-retouch.



	Basalt	Phonolite	Quartzite (naibor)	Total
Tested cobble.	1			1
Cortical flake with no cortical platform.	1			1
Cordal flake			3	3
Desbordant flake			1	1
Striking platform preparation flake		1		1
Ordinary flake	1	1	6	8
Ordinary flake with cortex	2		1	3
Centripetal flake (discoid)	1		2	3
Cortical centripetal flake (discoid)			1	1
Levallois preferential			1	1
Levallois recurrent bipolar			1	1
Levallois (centripetal récurrent)	1			1
Core	1	Š.	3	4
Chunck	1	7	1	2
Hammer	1			1
Total	10	2	20	32

Table A.9: Lithic inventory by raw materials at Level 2.

	Basalt	Quartzite (naibor)	Total
Cordal flake	1		1
Striking platform preparation flake	1		1
Ordinary flake	4	4	8
Ordinary flake with cortex	1		1
Centripetal flake (discoid)	1		1
Levallois preferential	1	1	2
Blade	1		1
Core		1	1
Total	10	6	16

Table A.10: Lithic inventory by raw materials at Level 2a.

	Phonolite	Ochre	Quartzite (naibor)	Total
Cobble (fragment)	1			1
Ordinary flake	1		7	8
Centripetal flake (discoid)			1	1
Core			1	1
Chunck	1		2	3
Indet.	1			1
Other		1		1
Total	4	1	11	16

Table A.11: Lithic inventory by raw materials at Level 3.

	Basalt	Phonolite	Quartz	Quartzite (naibor)	Chert	Total
Cordal Flake		1				1
Desbordant flake			1			1
Striking platform preparation flake				1		1
Ordinary flake		2		6	ľ	8
Ordinary flake with cortex	1			1	1	3
Levallois Preferential		1				1
Levallois (récurrent bipolar)		1		S		1
Core				3		3
Chunck				1		1
Chip				3		3
Total	1	5	1	15	1	23

Table A.12: Lithic inventory by raw materials at Level 4:

	Basalt	Phonolite	Quartzite (naibor)	Total
Ordinary flake	2	2	2	6
Ordinary flake with cortex	1			1
Levallois (récurrent unipolar)	1			1
Levallois (indet.)		1		1
Total	4	3	3	10

Table A.13: Lithic inventory by raw materials at Level 5.

	Basalt	Total
Pseudolevallois point	1	1
Centripetal flake (discoid):	2	2
Total	3	3

Table A.14: Lithic inventory by raw materials at Level 5a.

	Quartzite (naibor)	Phonolite	Basalt	Quartz	Total
Level 2					
Denticulate	1		1		2
Heavy Duty-core axe			1		1
Notch	1				1
Level 2a					
Denticulate-endscraper	1				1
Level 4					
Notch	1				1
Sidescraper (dejeté)				1	1
Level 5					
Sidescraper inverse simple			1		1
Total	4		3	1	8

Table A.15: Retouched blanks inventory by levels and raw materials.

	Quartzite (Naibor)	Basalt	Total	
Level 2				
Centripetal		1	1	
Centripetal (bifacial)	1		1	
Poliédrico	1		1	
Unipolar	1		1	
Level 2a				
poliédrico	1		1) ´
Level 3				
Discoid	1		1	
(unifacial)				
Level 4				
Levallois preferential	1		1	
Oportunistic	2		2	
Total	8	1	9	

Table A.16: Core inventory by levels and raw materials.

Table 1: Summary of luminescence dating from VCS including the depth and water content used for calculation of the environmental dose rate. A 5% error has been added to the assumed water content. Estimated equivalent doses derived from IRSL measurements and corresponding burial ages, expressed in kiloyears (ka).

Sample	Lab. Code Luminescence	Lab. Code Gamma spectrometry	Trench 2 Level	Depth (m)	w (%)	Environmental dose rate (Gy/ka)	Equivalent dose (Gy)	Age (ka)
VCS-L1	C-L4535	2915	1	0.6	10	4.44 ± 0.21	334.4 ± 17.2	75.3 ± 5.2
VCS-L4	C-L4536	2916	4	0.8	10	4.26 ± 0.21	365.4 ± 17.7	85.8 ± 5.9
VCS-L5	C-L4537	2917	5	2.0	10	3.97 ± 0.20	342.3 ± 13.4	86.2 ± 5.5

	Lev	vel 2	Lev	vel 2a	Leve	Level 3 Level 4 Lev		Level 5	
	Bones	Lithics	Lithics Bones Lithics Bones Lithics		Bones	Lithics	Lithics		
	V: 13.8988	V: 13.738	V: 12.7429	V: 11.996	V: 11.332	V: 10.1176	V: 9.2738	V: 9.1716	V: 11.8716
Kuiper	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01
	U ² : 11.2854	U ² : 11.7863	U ² : 10.165	U ² : 9.5455	U ² : 4.9172	U ² : 4.2042	U ² : 3.6202	U ² : 3.8968	U ² : 8.2133
Watson	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01	p<0.01
	Z: 0.2144	Z: 0.3436	Z: 0.2581	Z: 0.296	Z: 0.0749	Z: 0.129	Z: 0.2144	Z: 0.1469	Z: 0.2894
Rayleigh	p=0	p=0	p=0	p=0	p=0	p=0	p=0.21	p=0	p=0

Table 2. Omnibus statistical tests applied to the different archaeological levels of VCS, divided by bones and lithics.

ACER.

Taxa/Size	LEVE	L 2		LEVE	L 2a		LEVE	L 3		LEVE	L 4		LEVE	L 5		LEVE	L 5a	
	NISP	MNE	MNI	NISP	MNE	MNI	NISP	MNE	MNI	NISP	MNE	MNI	NISP	MNE	MNI	NISP	MNE	MNI
Alcelaphini							1	1	1	1	1	1						
Antilophini	1	1	1								R							
Equidae	1	1	1	1	1	1	3	2	1	1	1	1						
Hippopotamidae	1	1	1															
Testudinae																1	1	1
Rodentia				1	1	1	2	2	1									
Size 5	1	1		1	1		2	1	\mathcal{O}									
Size 4	4			2	1		8	2		5	2		1					
Size 3	4	1		4	1		18			15	6		2	2				
Size 2	3			3	1		9	Y		9	6					1	1	
Size 1	2						$\overline{\mathbf{A}}$											
Indeterminate	9			7			2			19						1		
Total	28	5	3	21	6	2	46	8	3	59	16	2	3	2	0	3	2	1

Table 3. NISP, MNE and MNI by taxa and weights size category at archaeological levels of VCS.

	Level 2	Level 2a	Level 3	Level 4	Level 5	Level 5a	Total
Phase 0: test and raw material preparation.		1				1	
Cobble			1				1
Tested cobble	1						1
Cortical flake with no cortical platform	1						1
Phase 1: Preparation						A ST	
Desbordant flake	1			1			2
Striking platform preparation flake	1	1		1	C		3
Pseudolevallois point						1	1
Cordal flake	3	1		1			5
Phase 2: Exploitation					2		
Ordinary flake	11	9	8	11	7		46
Centripetal flake (discoid)	4	1	1			2	8
Levallois (Preferential)	1	2		1			4
Levallois (unipolar)					1		1
Levallois (bipolar)	1			1			2
Levallois (centripetal)	1						1
Levallois (indet.)					1		1
Blade		1					1
Phase 4. Abandon			Y				
Core	4	1	1	3			9
Phase 5. Divers							
Chunck	2		3	1			6
Hammer	1						1
Indet.			1				1
Chip				3			3
Other			1				1
	Y						
Total	32	16	16	23	9	3	99

Table 4. Blanks' chaîne opératoire inventory by levels at VCS.

Industry	Site	Techno-Typological features	Туре	Year	Chronology (Ka BP)	Bibliography	
Njarasan	Eyasi Shore	Heavy Duty and scrapers Discoidal method	Surface	1935 and 1977	± 200	Mehlman, 1989	
Ngaloban	Laetoli	Scrapers, heavy Duty, points (uni/bifacial) Discoidal method	Surface/pit	1998-2003	120 130-100 240-500	Hay, 1987 Manega, 1993 Manega, 1993	
Sanzako	Mumba VI-B	Scrapers, bifacial pieces, Heavy Duty Discoid method	Excavation	1938	131-109	Mehlman, 1989	
Kisele	Mumba VI-A Nasera 18-25 Nasera 12-17	Smaller industry Scrapers, points, bifacial pieces Discoid/Levallois methods	Excavation	1938 1975-76 1975-76	90-56	Mehlman, 1989 Gliganic et al., 2012	
Loiyangalanian	Loiyangalani	Scrapers, denticulates Discoid/Bipolar methods	Excavation	1985 and 2001-2005	64	Bower & Mabulla, 2013	
Figure 5. Map of N	lorthern Tanzania v	vith the major sites cited in text (source NASA).					













CONFLITC OF INTERESTS:

The authors declare that they have no conflict of interest.