Twisted handaxes in Middle Pleistocene Britain and their implications for regional-scale cultural variation and the deep history of Acheulean hominin groups.

Mark White<sup>1</sup>, Nick Ashton<sup>2</sup> and David Bridgland<sup>3</sup>

Department of Archaeology, Durham University, South Road, Durham DH1 3LE.
 British Museum (Frank's House), 38-46 Orsman Road, London N1 5QJ
 Department of Geography, Durham University, South Road, Durham DH1 3LE.

## Introduction

Two decades ago, White (1998a) suggested that Acheulean assemblages with high frequencies of twisted ovate handaxes all belonged to late Marine Isotope Stage [MIS] 11 or early MIS 10. This was not the first time this highly distinctive type had been singled out for special attention. Evans (1872, 520) recorded several early discoveries of twisted handaxes at Hoxne and Santon Downham, noting that while they were 'by no means uncommon' they were probably accidental. Spurrell (1883) likewise thought twisted edges were a defect, left uncorrected due to raw material constraints. This view was not shared by Smith and Dewey (1913, 1914), who regarded the rare twisted forms from Swanscombe and Dartford as technologically advanced. Despite coming from the highest, and therefore oldest, terrace in the Lower Thames, they assigned them to the culturally-evolved St Acheul II stage, equivalent to the 'Upper Acheulean' described by Victor Commont (1908) in the Somme Valley. Henri Breuil included the latter material in his Acheulean IV, the penultimate phase of the Acheulean before, according to him, it evolved into the Micoquian (Acheulean VI and VII; Breuil and Kelley 1954). Collins (1969) similarly saw twists as a technically advanced trait, using them as the basis for his youngest 'Elveden' stage of the Acheulean, which he dated, on the basis of typology, to the mid-Rissian (i.e. the penultimate glacial) period. Around the same time, Roe (1968) identified a large group of assemblages (his Ovate Tradition, Sub-Group VI) with above-average frequencies of twisted handaxes, but the lack of secure dating and compressed chronological framework, which recognised too few climatic cycles, again meant that he was unable to link them by anything other than artefact form. More mindful of the spurious use of typological dating, Roe leaned towards a

functional explanation in print (Roe 1981), although privately he always suspected they were of the same age (Derek Roe, personal communication to MJW, 1998).

By the late 1990s, important advances in lithostratigraphy (e.g. Bridgland 1994, Antoine et al 2000) and biostratigraphy (e.g. Keen 1990; Schreve 1997, 2001a, b), in particular, had made it possible to correlate the fragmentary terrestrial record with the more complete oxygen isotope record from deep sea and terrestrial ice cores. This provided archaeologists with a new and expanded chronological framework that allowed the archaeological evidence to be arranged in a more coherent order. White (1998a) simply recognised that all of the better understood and dated sites with twisted handaxes belonged, according to several new and independent lines of evidence, to the same period, which he then believed to be late MIS 11/ early MIS10. He further suggested that they might represent an endemic cultural practice, perhaps one that developed during a period of high sea-level when Britain was cut off from Europe. He warned against uncritically using isolated twisted handaxes for dating purposes, following the prevailing near-consensus among British workers that any handaxe type could feasibly occur any place and any time, simply by chance: a statistical consequence of giving enough hominins enough time and enough stone.

White's (1998a) interpretation ran contrary to the dominant theoretical paradigm of the time, and contrary to his main thesis (White 1998b). Older ideas that handaxe shapes reflected the cultural norms of their makers, conscious or unconscious reflections of the groups who made them, had given way to more 'Processual' interpretations, which emphasised the impact of function, raw materials and resharpening on biface form. Similarly, the notion that handaxe shapes were the physical expression of 'mental templates', traditional and socially-resonant designs held in the mind's eye, was recast in more techno-functional terms as a 'mental construct' (Ashton and McNabb 1994). This involved a basic set of rules uniting all handaxes - bifacial working to form cutting edges, an idea of symmetry and distinct butts and tips – in which the realisation of a practical bifacial edge was more important than final form and in which any consistency in shape emerged from repetitive rhythms of making, rather than design (e.g., White 1998b).

Today twisted handaxes are one of a number of temporally-restricted types used by White and Bridgland (White et al. 2006, 2017; White, 2015; Bridgland and White 2014, 2015) to construct a new Lower Palaeolithic framework, based not on notional schemes of typological progression but on independent relative and absolute dating methods. This includes a consistent pattern in the timing of Clactonian and Levalloisian industries, as well as a number of handaxe assemblage types that belong to different interglacial cycles. In other words, Roe's (1981) hunch that the apparent lack of coherent cultural patterning was due to an inaccurate and inadequate chronological framework was correct; some of the variation in handaxe shape is culturally significant, and this has the potential to reveal much about hominin settlement, social networks and modes of social transmission during the Middle Pleistocene.

Here we return to our original temporal peculiarity, twisted handaxes, and explore more fully their distribution, significance and contribution to the deep history of Palaeolithic Britain. A map showing the sites discussed in this paper is presented in Figure 1. <FIGURE 1>

#### The Twisted Handaxe Knapping Method: what, how and why

Twisted handaxes are usually ovate or cordate forms, although they are defined not by the outline shape of the tool but by the profile of the edges, which display strong (usually reversed) ogee curves on all sides (Figure 2). In more mathematical terms, 'when viewed in the four orthogonal directions (left, right, distal and proximal)', twisted handaxes present a 'sinusoidal edge [with] four inflection points, i.e., the point on a curve at which the curvature changes sine and the curve changes from concave upwards to concave downwards or vice versa' (Gallotti et al. 2010, 310).

#### <FIGURE 2>

Twisted handaxes were created using a specific knapping strategy (Figure 2a). As described by White (1998a), this involved the division of the implement into four quarters or arcs, each of which was worked in the opposite direction to that adjacent to it, causing their edges to curve away from one another. Each quarter was brought into the correct knapping position by a series of rotations and inversions in the hand, the precise sequence of which was free to vary provided that contiguous quarters were preferentially worked in different directions and diagonally opposed quarters in the same direction (Figure 2b). Expressed in the terms employed by Gallotti et al. (2010), there has to be an inflection point on each edge at which the knapping direction reverses, causing the edges on either side to curve away from one another. The twist could presumably have been imposed upon a

handaxe at almost any stage of manufacture, although it was preserved or enhanced during finishing. Cognitively, the production of twisted edges demands the mental ability to conceive and impose wave forms onto handaxe edges, by keeping several opposing and future knapping operations in mind. The individual edge segments must continuously be conceived as parts of the greater whole, and work in harmony with each other to achieve what is, in our opinion, not an accident but a clear design template.

White also noted that 94% of twisted handaxes displayed z-twists rather than stwists, a likely indicator of handedness and brain laterality among Acheulean hominins. Unknown to White (1998a), Flaxman C.J. Spurrell (1883) had described the technique used to make twisted edge, and its implications for handedness, 115 years earlier.

Similar deviations from the more typical straight or zig-zagged bifacial edge can be produced accidentally, but such 'pseudo-twists' are usually found only on one edge and are typically curved at one end, rather than being properly sinusoidal. The assemblage from Broom is a case in point. Hosfield and colleagues reported that 3% (n=30) of handaxes showed a twisted profile, but that "in many cases the S-twists are not pronounced, and in no examples were they as diagnostic as the S and Z-twisted ovates discussed by White (1998a)" (Hosfield and Chambers 2009, 87; Hosfield and Green 2014). From the examples we have seen, we concur with this assessment: the Broom 'twists' represent unstructured deviations from straight, some with repeated sinusoidal waves on a single edge, rather than classic twisted handaxes. These pseudo-twists would thus appear to add to the already unusual set of practices seen at Broom, which include the use of Greensand chert (95% of handaxes), the production of asymmetrical handaxes (23%) and specific sharpening techniques (Hosfield and Green 2014; cf. Ashton and Hosfield 2009; Ashton et al 2011).

An obvious question is whether twisted handaxes served a specific function or represent a re-sharpened (but nonetheless formalised) expression of untwisted forms. Our analyses (Supplementary Data ) found little technological or morphometrical support for the idea that twisted edges emerged from regularised re-sharpening practices. Nor are twisted handaxes found in flint-poor regions where hominins might have been more economical in their use of stone resources and re-sharpened their tools more frequently: quite the contrary in fact. Twisted edges conceivably served a specific or auxiliary function, such as scraping (Walters 1996), although this is equally possible for many non-twisted forms. None

Twists 2018/ 5

of the existing examples are fresh enough for micro-wear analyses, and the suggestion remains untested (Walters 1996; cf. Keeley 1980).

In contrast, evidence that twisted edges were a deliberate design might be found at the primary-context site at Foxhall Road, Ipswich. Here, three 'twinned' pairs of untwisted handaxes, and a trio of near-identical twisted forms (Figure 3), were found together in a small cluster around a central focus, which Nina Layard, the excavator, thought was a hearth (White and Plunkett 2004). We interpret the pairs and the trio as the products of different individuals, each having their own style based around the common ovate form. They are witness to mental templates repeatedly imposed using skill and precision, derived from socially-acquired techniques and ideas, learnt as children and developed throughout life. If these groupings do not represent the distinctive products of specific individuals, then they were made by different people who closely copied each other, which would indicate that imitation (or overimitation: Shipton and Nielsen 2015) was a key factor in the social transmission of handaxes between contemporaries and across generations.

Regardless of whether twisted edges were a redundant design feature or a shared practical solution, the temporal and geographical patterning of twisted handaxes described below suggest that this was part of a long-lasting regional tradition.

# The Stratigraphical History of the Twisted Handaxe

White (1998a) suggested that assemblages containing relatively high frequencies of twisted handaxes all belonged to late MIS 11/early MIS 10, their occurrence at the top of the Swanscombe sequence suggesting they were present during the final part of the interglacial or interglacial complex (cf. Bridgland, 1994; Conway et al., 1996). Recent advances in our understanding of MIS 11, combined with new discoveries and a wider examination of the twisted-ovate phenomenon, suggests that this did not reveal the real story. MIS 11, which spanned approximately 60,000 years, encompassed two warm periods, MIS 11c (the Hoxnian *sensu stricto*) and MIS 11a, separated by a cold interval, MIS 11b (Ashton et al. 2008; White et al 2013). During the past 20 years, more MIS 11 sites have been correlated to the sub-stage level (Table 1). A greater number of sites can now also be confidently attributed to other Marine Isotope Stages and substages (e.g., Candy and Schreve 2007; Penkman et al. 2013), providing valuable comparative data from which to explore the

significance of the twisted ovate phenomenon (Table 2). Below we summarise the dating evidence for the key twisted-handaxe assemblages (see Figure 4).

<TABLE 1> <FIGURE 4>

# East Anglia and Hertfordshire

Barnham and Elveden are situated in the Breckland region of East Anglia, 3.5km south and 5km southwest of Thetford, respectively. The Barnham sequence consists of a deep glaciogenic channel filled with Anglian (MIS 12) till and glacial outwash, upon which a sequence of interglacial lacustrine, fluvial and colluvial deposits has been emplaced (Ashton et al. 1998). At Elveden there is a depression within the Chalk, mantled by Anglian till and filled by a similar sequence of aqueous and slope deposits to that at Barnham. The stratigraphy and environmental signatures of the two sequences led Ashton et al. (2005) to conclude they were part of the same evolving drainage network, spanning the late Anglian and early Hoxnian: late MIS 12 and MIS 11c (cf. Penkman et al. 2013; Ashton et al. 2016;). Twisted handaxes account for 35% of the total Elveden assemblage and 45% of the (metrically defined) ovates alone. Handaxes from Barnham are rare. Only three have been recovered in the recent excavations, but one of these is a twisted cordate similar to those found at Elveden. It was recovered in 2017, on the surface of the organic clays in the centre of the pit.

The sequence at Foxhall Road comprises a series of interglacial lacustrine deposits filling a basin formed in MIS 12 till and outwash, once again very similar to the records at Barnham and Elveden. These are overlain by fluvial sands and gravels indicative of a return to cold conditions (Allen et al. in press). The whole describes an immediate post-glacial landscape, one covered by a pock-marked till plain with disrupted drainage (cf. Magerud 1991), and containing interglacial lakes that gradually filled before being subsumed by an integrated fluvial system. Allen and White (2004) suggested that deposition had spanned the Anglian–Hoxnian, a proposal supported by recent OSL dates of 416±36 ka and 434±54 ka (Allen et al., in press). The twisted-ovate assemblage came from the deepest archaeological horizon, on the sloping Grey Clay deposits, 4m below the (1903) ground surface and 2m below cold-climate horizontal beds, part of a continuous sequence of lake beds that had been accumulating since Anglian deglaciation; we therefore correlate the twisted ovate assemblage with the Hoxnian, MIS 11c.

The artefacts from Hitchin were almost certainly in primary context, although their collection history has probably conflated two separate assemblages: a point-dominated assemblage with forms not dissimilar to those from the Swanscombe Middle Gravel, and an ovate assemblage with a high proportion of twisted ovates (37.5 %). With dedicated archival research or new fieldwork, it might prove possible to disentangle these assemblages. The sequence at Hitchin represents a stream-fed interglacial lake occupying a basin (kettle hole?) in outwash gravel and till (Boreham and Gibbard 1995). The lake later fragmented into small pools, within which shelly freshwater marl accumulated, and was ultimately incorporated into a maturing drainage system. Later infilling of the basin is represented by a series of poorly stratified silts, possibly laid down under cold conditions. Pollen from the lake beds and *Chara* marl showed accumulation throughout Ho I and Ho II of the Hoxnian, the absence of later pollen being either a result of weathering or cessation of organic sedimentation (Boreham and Gibbard 1995). The majority of the artefacts were found in the lake beds, the Chara marl and the brickearth at Jeeves's Pit and Ransom's Pit, towards the top of the lake sequence; according to West's pollen results from Jeeves's Pit, the Chara marl represents deposition during Hollc (West 1955). Boreham and Gibbard linked evidence for a reduced water table at Hitchin with similar phenomena across East Anglia, all of which appear to span Holl–III. Thus, both the pollen and hydrological sequence suggest that Hitchin spans a substantial length of the Hoxnian *sensu stricto*, MIS 11c.

Roe's Group VI also included the assemblage from Allington Hill, Cambridgeshire, on the edge of the Breckland, where 46% of the handaxes showed a marked twisted profile. This site is very poorly known, but artefacts came from gravel at 150ft (~48 m) O.D. Exposures seen by Hughes (1916) and the British Geological Survey (BGS) (Worssam and Taylor 1969) showed that the gravel was heavily festooned and contorted, and overlay what might have been a till. Hughes believed the contortions to have been caused by dissolution of the underlying Chalk. BGS mapping suggests that this gravel represents head deposits, evidence of 'drastic landscape changes during some early glacial episode' (Wymer 1999, 167). If the basal deposit is till, then this can only represent MIS12 in this part of East Anglia, so the deposits above most likely belong to the succeeding MIS 11 interglacial. However, for present purposes the assemblage remains undated and is excluded from the current discussion. The site is clearly a pressing target for further examination. Owing to similar uncertainties about context, assemblage integrity and age, we also exclude the possible MIS 11 assemblage at Santon Downham, Suffolk, which has 18% twisted handaxes but was assigned by Roe (1968) to his Group IV (mixed or intermediate).

# South of the Thames

The deposits at Barnfield Pit Swanscombe (= the skull site) record a more-or-less complete sequence through MIS 11 as well, perhaps, as the end of MIS 12 and the beginning of MIS 10 (Figure 5). In his review of the Swanscombe deposits and their significance, Bridgland (1994) adopted the stratigraphical scheme established by Bernard Conway during the 1968–1972 Waechter investigation of the site (Conway 1969, 1970, 1972, 1996; Conway and Waechter 1977). This recognised three divisions within the sequence, termed phases I, II and III, with the first two recording terminal MIS 12 and the whole of MIS 11c, a long-accepted attribution based on both lithostratgraphical and biostratigraphical evidence such as height, the presence of Rhenish molluscan fauna (which first appears in the Thames in Hollla of the Hoxnian interglacial) and a typical MIS 11 mammalian suite (Bridgland 1994; Schreve 2001b, White et al. 2013). Archaeologically, these deposits contain the much-disputed Clactonian industry (Phase I) and a point-dominated Acheulean assemblage attributed to Roe's Group II (Phase II). They have yielded no twisted handaxes, unlike contemporaneous East Anglian sites of similar age.

#### <FIGURE 5>

The age of the Phase III deposits is less clear-cut, although they certainly post-date the posttemperate stage of the Hoxnian (MIS 11c) interglacial. Evidence for deteriorating climatic conditions in the top of the Phase II sequence comes from the arrival of lemming, and the replacement of closed-canopy molluscs by open-ground species in the Upper Middle Gravel (Sutcliffe and Kowalski 1976; Kerney 1971). The overlying Phase III deposits, the Upper Sand and Upper Gravel, show periglacial features such as cryoturbation and small ice-wedge casts (Conway 1996); the latter also contains musk-ox, a clear indicator of arctic conditions. Conway interpreted these cold-climate levels as representing the MIS 10 glacial with the overlying Upper Loam deposits, which have been interpreted as interglacial estuarine deposits, therefore belonging to MIS 9. However, Schreve (2001b), having recognised MIS 9 deposits with a completely separate biostratigraphical signature in the lower-level Corbets Tey terrace at sites such as Purfleet (Schreve et al., 2002), attributed the Swanscombe Upper Loam to MIS 11; in which case, the cold episode must represent MIS 11b and the Upper Loam MIS 11a. There is further support for this climatic complexity within the usually disregarded palynological study from Barnfield Pit by Hubbard (1972, 1982; see, however, Turner, 1985).

The presence of a white-patinated twisted-ovate assemblage in the Swanscombe Upper Loam has been accepted for over 105 years, although Roe was unable to reconstruct a satisfactory sample (just 18 handaxes marked as Barnfield UL, four of which were twisted) and their rarity in the Stopes Collection has led to doubt about whether it ever really existed outside workmen's anecdotes (Wenban-Smith, pers. comm 2015). However, White et al.'s (2013) invertebrate-based correlation and historical reconstructions of various Swanscombe localities convincingly shows that the twisted ovates found by Newton (1901) at Dierden's Pit, and, by extension, those from at Rickson's Pit too, came from lateral equivalents of the Barnfield Upper Loam (Figure 4), which confirms the presence of a significant twisted component within assemblages from the Phase III deposits, temporally and morphologically distinct from the untwisted point-dominated industry ubiquitous throughout the area during Phase II times.

The Bowman's Lodge and Wansunt Pits revealed a sequence of ~15m of gravel (Dartford Heath Gravel, DHG) overlain by silts and clays (Wansunt Loam), at a height of ~42m O.D. The apparent ~10m height difference between Dartford Heath and the top of the Boyn Hill Terrace at Barnfield Pit, Swancombe, led some to attribute the Dartford Heath deposits to the (MIS 12) Black Park Terrace (this historical debate has been summarised by Bridgland 1994 and 2006). However, more recent archaeological evaluation at the Swan Valley Community School and the adjacent Sweyne County Primary School, to the west of Southfleet Road, Swanscombe, revealed a hitherto unrecorded occurrence of the Swanscombe sequence with a much thicker Upper Loam extending above 40 m O.D. This effectively settles the argument in favour of correlation of Swanscombe with Dartford Heath and the attribution of both to the Boyn Hill Formation (Wenban-Smith and Bridgland, 2001) of MIS 11 age.

This MIS 11 correlation can be more finely tuned by further comparison with Swanscombe. Dewey (1959) recorded a tributary channel feature beneath the DHG at Pearson's Pit, cut to a level comparable with the Swanscombe Lower Gravel. Newton (1895) reported molluscs recovered by Spurrell from the DHG at Dartford Brent (probably at TQ 555743; cf. Bridgland, 1994), including *Corbicula fluminalis, Bithynia tentaculata* and *Valvata piscinalis*. The first of these is a member of the distinctive assemblage known as the '*Rhenish* fauna', which appears in Lower Thames deposits between Hoxnian pollen zones Ho IIIa-IIIb of MIS 11c (White et al. 2013), and strongly supports both the correlation of the DHG with the Swanscombe Middle Gravel and the formation of a land-connection with Europe. Chandler and Leach (1912; Leach 1913) reported faunal remains from the lower sandy gravel of the DHG, including *Palaeoloxodon antiquus, Cervus elaphas, Equus ferus* and indeterminate rhinoceros, the first of which at least is an indicator of wooded interglacial conditions. We thus equate the basal channel and the main body of gravel at Dartford Heath with the Phase I and Phase II interglacial deposits at Swanscombe. Both thus represent MIS 11c, encompassing Ho I to Ho IIIb-IV.

Artefacts from within the body of the Dartford Heath Gravel are few and poorly contextualised, although where reported are similar to those from the Middle Gravel at Swanscombe (Wymer 1968). Primary-context twisted-ovate assemblages occur above the Dartford Heath Gravel at both Wansunt and Bowman's Lodge: in the Wansunt Loam at the former (White et al. 1995); in the Wansunt Loam and on the surface of the DHG at the latter (Tester 1951, 1953, 1975). They must be younger than the late temperate stage of the MIS 11c interglacial represented by the bulk of the DHG, and we equate them with MIS 11a, the same age, same height and same context lithology as the Upper Loam at Swanscombe.

Another major occurrence of twisted ovate handaxes, not considered by us before, is found at Limpsfield, Surrey, where several hundred handaxes, forming a surface assemblage exposed by a deep ploughing of terrace gravels 0.6m below the surface, were collected by A.M. Bell between 1883 and 1906 (Field et al., 1999; Bridgland, 2003). The gravel, on the interfluve between the catchments of the Eden–Medway, to the south, and the Darent, to the north, is at ~150 m O.D. It contains Hastings Beds material, leading to the conclusion that it was deposited by a formerly more extensive River Darent that drained the central Weald (Gossling 1940; Bridgland, 1999, 2003). Prestwich (1891) made a connection between Limpsfield and Dartford Heath, correlating the DHG with his 'Upper Valley Gravels' of the Darent, which he traced from the Darent–Eden watershed at Limpsfield, where he noted the occurrence of artefacts. Bridgland (2003) similarly concluded that the Limpsfield

gravel correlated with the Boyn Hill of the Thames, the presence of twisted ovates being among the several criteria he used to establish the connection. Nonetheless, this can be presumed to represent the final floodplain of a Darent draining from the Weald, prior to its subsequent beheading (capture) by the Medway system (cf. Wooldridge and Linton, 1955; Worssam, 1973; Bridgland, 2003), and is correlated here with MIS 11.

Twisted handaxes were also reported from Terrace B at Farnham, Surrey, the MIS 11 'step' in the terrace-staircase of the River Wey (Oakley 1939; Bridgland and White 2018). They were not apparently found in the other terraces, although further research is required to confirm and quantify the Farnham sequence. Only the Farnham Terrace A material in Table 2 is included in the present paper.

## <TABLE 2>

#### Twisted handaxes in MIS 11 and other interglacials

Twisted handaxes are rare (Table 3). From a sample of 4722 handaxes from 27 British sites (30 assemblages), only 302 (6.3%) show twisted edges (Table 2). But of these, 252, or 84%, occur in sites attributed to MIS 11. (Table 3 shows alternative calculations that exclude the Broom 'twists' and the entire Limpsfield sample, although even when the data are cleansed the pattern of MIS 11 predominance remains).

# <TABLE 3>

Twisted forms account for 21% of handaxes from all sites dated to MIS 11. At the sub-stage level, greater regional patterning becomes evident. Twisted handaxes make up 33% of MIS 11c assemblages in East Anglia and Hertfordshire, but just 1% of assemblages of this age from south of the Thames. Conversely, twisted handaxes constitute 22% of the five MIS 11a assemblages south of the Thames, but the solitary East Anglian site thus far firmly assigned to this sub-stage, Hoxne, has only 3%. The latter is little more than background variation and is based on Roe's older sample: the smaller but contextually more secure sample of MIS 11a handaxes (n=19) from the Wymer/Chicago Excavations has 0% (Singer et al 1993).

Pre-Anglian (MIS 13) contexts contain just 1% twisted handaxes, with over half of these coming from the Warren Hill 'fresh' assemblage, which is poorly contextualised and might possibly include later material. The percentage of twisted handaxes from sites correlated with the MIS 9 (Purfleet) interglacial is practically zero (0.1%, excluding Broom), with even the two from Furze Platt possibly being intrusive (naturally or through collector/museum error) from the higher (and nearby) Boyn Hill terrace. The combined percentage of twisted handaxes in assemblages belonging to MIS13 and MIS9 is just 1.4%, or 0.5% excluding Broom. Thus the twisted handaxes from MIS13 and MIS 9 contexts, where they occur at all, are merely background variation, and may be due to many factors, including chance production, intrusion of younger or older objects, collector error, or an idiosyncrasy that had little relevance to the wider group.

The MIS 11 assemblages from Swanscombe Middle Gravels, Dovercourt and Hoxne, which contain very low levels of classic twisted ovates, show 6–15% 'twisted-tips' (Roe 1968, Table VI), also seen in lower frequencies in Roe's typologically and stratigraphically mixed MIS 11 samples from Foxhall Road and Hitchin. This feature is not found in other interglacials. So, regardless of whether they made classic twisted ovates, hominins throughout MIS 11 appear to have been familiar with a technique for producing twisted edges that was not used in other periods. It is important to note here that all these assemblages fall into Roe's point-dominated tradition, a group which may owe some of its large-scale characteristics to local raw materials (White 1998b). It remains to be seen whether an MIS 11c ovate-dominated assemblage south of the Thames or a larger MIS 11a assemblage north of it would contain twisted handaxes, but nonetheless the influence of raw-material packages on handaxe form in Britain has been hotly disputed (see for example Wenban-Smith et al. 2001), and a more moderate view would see raw material selection as part of a cultural chaîne opératoire. Nodules were chosen by the knapper because they assisted the production of the culturally desired form; stone did not impose form onto a passive human instrument. There are also enough examples of straight-edged ovates from all relevant sites, which could have been twisted but were not, to suggest this is not a controlling factor.

Collector preferences may have created a bias towards fine and unusual forms such as twists, and may be affecting the very high proportion of twisted handaxes seen at Limpsfield, but the fact remains that they must have occurred in greater numbers at some sites to allow such biases to develop and leave us with the collections we have. Llewellyn Treacher, who was active around the Maidenhead area of the Middle Thames, found twisted ovates in pits located on the Boyn Hill Terrace, but almost none when exploring sites on the next terrace, the Lynch Hill, where ficron and cleaver assemblages occurred instead. This was not the expected pattern and caused interpretative problems (e.g. King and Oakley 1936). Similarly, twisted forms made up 10% of the handaxes recovered by Worthington Smith (1894) from the rather low-resolution secondary context sites at Stamford Hill and Leytonstone, situated on the Boyn Hill Terrace of the Lea, but there were none at Stoke Newington, where a primary-context assemblage was collected from sands and gravels of the Lynch Hill Terrace (Green et al. 2004). Smith is known to have been a comprehensive collector not prone to selection biases, other than those forced upon him by heavy items that he could not carry nor afford to transport.

Twisted handaxes are thus rare, highly distinctive, and tightly restricted in time and space.

# Twisted handaxes in global context

On a global scale, twisted-handaxe assemblages are extremely rare, an observation underlined by the fact that only one well-stratified and well-studied site (Gombore II, Melka Kunture, Ethiopia), plus a handful of surface collections, is known for the entire African Early Stone Age (Gallotti et al 2010). In Europe, twisted-handaxe assemblages have been widely reported only from northern France (Callow 1976), particularly from the complex terrace deposits and overlying loessic and colluvial sequences of the Somme.

The oldest are the two series (fresh and worn) from fluvial terrace gravel at Cagny la Garenne (Breuil 1934; Bordes 1956; Bourdier 1969), now dated to MIS12 (Antoine et al 2015). Callow's analysis of the worn series in the Musée de l'Homme and Bordeaux (n=>100) found it to be dominated by thick ovate handaxes (limandes and amydaloids), of which 48% were twisted (27% of the entire assemblage). Fewer twisted handaxes (13%) were recorded in the more refined (i.e. thinner) fresh series, none of which were pronounced.

Victor Commont recorded the presence of classic twisted ovates at the St Acheul type-site. 'Some' came from the *sable roux/brun* (Bed H) at Bultel and Tellier's Pit (*Atelier Commont*; Commont 1909, 47), while large numbers (116 out of 300) were reported from the *sable roux* (Bed D) at No. 54 Rue de Cagny (Commont 1908, 559). These deposits were described as clayey-sands and were positioned above the fluviatile sands and gravels at both localities, at the base of the loessic and slope deposits. At Rue de Cagny the base of the sable roux was associated with large angular flint and whole nodules, while at Atelier

Commont, an in situ knapping floor occurred in this position, underlain locally by a shelly silt. More recent work at St Acheul identified an MIS 11 tufa, locally preserved in the same stratigraphical position as Commont's sable roux (Antoine and Limondin-Lozouet 2004), and it would seem reasonable to infer that the twisted ovate assemblage from the latter is of a very similar age. The MIS 12/early MIS 11 from the underlying fluvial sands and gravels in the St Acheul region, however, contain a mixture of pointed and ovate assemblages, none notably twisted. Other undisputed MIS 11 levels in France, such as La Celle and St Pierre les Elbeuf IV in the Seine basin (Cliquet et al 2009; Limondin-Lozouet et al 2015) similarly contain no twisted ovates, again indicating regional and temporal complexity in handaxe form.

A potentially younger, MIS 9, occurrence is found in the primary-context assemblage from the sable roux at Cagny L'Epinette (Agache 1971; Callow 1976, 1986; Antoine et al 2015), on the lower L'Epinette Terrace. Here, Callow (1976) recorded an unusual assemblage containing a high proportion of thick ovates with unworked butts. Approximately one-quarter (n=65) had twisted edges, although these were not pronounced. We have not studied these handaxes, nor those from Cagny la Garenne, but wonder whether they conform more to the Broom variation than true twisted forms. Younger still might be the twisted-ovate handaxes from the Older Loess at St. Acheul and Mareuil (Callow 1986), although these could well represent older inclusions.

In summary, various forms of twisted handaxes are found in the Somme in deposits currently thought to belong to MIS 12, MIS 11 and MIS 9. Those belonging to MIS 11 show the pronounced twists on well-made refined forms, exactly as found in the British sample, but those from MIS12 and MIS9 are on often found on thick handaxes and are rarely pronounced. Despite an extensive (although by no means exhaustive) literature review and consultation with European colleagues (see acknowledgments), we know of no major concentrations of twisted handaxes in Iberia, central or southern France, Germany, Italy, Belgium or the Netherlands. At present, such forms appear to be restricted to Britain and the closest neighbouring region of France, although we would genuinely welcome all information to the contrary.

#### Handaxes and the Deep History of MIS 11 Britain

The twisted-ovate phenomenon in Britain occurred in different regions before and after MIS 11b, the Thames acting as the physical dividing line. The presence of twisted tips in pointdominated assemblages on the 'wrong' side of this physical and temporal divide suggests that some form of twisted technique was familiar to all MIS 11 hominin groups, but was variably expressed among individuals and their networks.

Given the estimated small size of archaic hominin populations, and the small social groups and regional landscapes they inhabited (Gamble 2002), we doubt whether these preferences and variations had any meaning outside of Gamble's (1999) intimate and effective networks, involving kith and kin and maybe familiar others from neighbouring groups. In other words, they may have been used for 'assertive' signalling about the identity of individuals within their local group and social networks, but not 'emblemic' signalling representing the identity of the group within or without it (cf. Wiessner 1983; Gamble 1999). For the purposes of Palaeolithic archaeologists, however, these amount to the same thing: signals emerging from living in groups.

The greatest distance between our MIS 11c occurrences north of the Thames is 98km, between Hitchin and Ipswich (Figure 1), and they all could be subsumed within a network ~50km radius. This provides an area ca 50% greater (5026 km2 vs 7853 km2: radii of 40 km vs 50 km) than Gamble's 'local hominin network' (1999, 2002), but that was based on maximum distances of raw-material transfers, which almost certainly underestimated mobility in flint-rich landscapes. Thus, the twisted-ovate phenomenon in East Anglia and adjacent regions north of the Thames could feasibly be the product of just one or two local hominin groups. The same is true south of the Thames in MIS 11a. Dartford and Swanscombe are just 9km apart, a distance that could have been covered in a few hours. Limpsfield and Farnham, although not correlated to the sub-stage level, are only 25km and 75km from Dartford, respectively. The distance between these two sites is 56km. Again, the area south of the Thames might have been home to only a few related local groups, perhaps just a few hundred individuals (the social brain hypothesis predicts a group size of ~150, e.g., Dunbar et al 2014; cf. Gamble 2002; Pettitt and White 2012).

The twisted ovate phenomenon might thus represent a fleeting event recording perhaps a few generations of biased transmission (perhaps influenced by the makers' status, reproductive success, homophily or conformism) before drift took handaxes in another direction. At some sites it is preceded or succeeded by assemblages with quite different characteristics: at Foxhall Road the twisted-ovate assemblage occurred beneath a horizon with only straight-edged pointed handaxes, and the same might be true of the material from Hitchin. In MIS 11a, a similar pattern is seen at Hoxne, but here untwisted ovates were replaced by points (Singer et al 1993). But at a wider scale, the appearance of such an unusual type north of the Thames during the earlier warm substage of MIS 11 and south of the Thames in the later one, when it is missing from all other interglacials and rare on a global scale, suggests that the two are connected not just by coincidence but by history. The practice was much longer-lived on a regional level, suggesting that despite a small demographic presence, geographical barriers and being the human settlement furthest northwest from Africa (cf. Lycett and von Cramon-Taubadel 2008), handaxe manufacturing skills and ideas were preserved through strong networks of social transmission and conservatism in NW Europe.

Ashton (2017) has suggested that during periods of stable environment local groups were able to persist in the landscape over multi-generational timescales, and once established they developed ways to deal with local circumstances that became embedded into social practices. So technological practices and handaxe shape 'preferences' might still emerge from local resources, but through choice and historical engagement: landscapes of habit (e.g. Gamble 1999) created landscapes of cultural tradition. Conversely, periods of climatic instability would have caused large-scale shifts in population, particularly within Northern Europe, where the southwards translocation and/or extinction of populations would have occurred. In Britain, we might thus expect to find our clearest signatures in our longest interglacials, such as MIS 11.

Britain was first re-colonised after the Anglian glaciation (MIS 12) by populations from Europe that did not habitually make handaxes, leaving an industry widely known as the Clactonian (White 2000). The source populations for the Clactonian have never been satisfactorily identified, but possibly derived from areas of central Europe, where handaxes are rare or absent (White and Schreve 2001; Ashton 2017). They persisted through the stable environments of the early and full temperate periods (Ho I and II) before being replaced by Acheulean populations in HoIIb-III, coinciding with a pan-European catastrophic event registered in the pollen record as a period of rapid deforestation (Ashton et al 2008, Ashton 2017). That the Acheulean appeared at a period of environmental instability when 'resident populations' in the affected areas may have begun to decline or be displaced, can hardly be co-incidence. The question of whether sea-level change might also have been implicated is unresolved. Britain was almost certainly connected to the continent during Hollb-III, when the freshwater 'Rhenish fauna' arrived, but its status during the main interglacial is uncertain; different sea-level reconstructions provide different answers (see White 2015; Pettitt and White 2012). It is equally uncertain whether these populations ever met or whether, as Ashton (2017) evocatively asks, the first Acheulean settlers would have found a landscape eerily littered with the tools, structures, hearths and meals of an earlier extinct people.

The first MIS 11c handaxe makers south of the Thames did not make twisted handaxes, while those who settled East Anglia and adjacent regions did. This probably represents different regional source populations from neighbouring areas of Europe, and/or the drowned landscapes of the North Sea and Channel basins. The pattern reverses after the cold interval of MIS 11b, when twisted handaxes are found only south of the Thames. It seems reasonable to infer that cold conditions would have pushed East Anglian populations south during MIS 11b, as the climate deteriorated and the Thames became more braided and less of a physical barrier, or that these same factors would have facilitated greater mobility and contact between the two areas. If this were the case, these populations survived into MIS 11a in the south, but a different population without twists must have entered East Anglia in MIS 11a. Alternatively, both MIS 11c populations might have moved east and south never to return, or died out entirely, in which case Britain was colonised in MIS 11a by new populations with their own handaxe traditions. One of these was probably related to the original twisted populations, who survived MIS 11b somewhere.

The pattern in neighbouring Europe is intriguing in this regard. On current dating estimates, twists of some form were made in the Somme during MIS12, MIS 11 and MIS 10/9, making this an obvious region in which to seek deep historical connections with Britain. It is equally interesting that, during this long period, classic twisted ovates are found in France only in MIS 11 contexts, those from sites which temporally sandwich this period being less-well-developed early and 'degenerating' late examples. These are tentative observations that require further research, but they provide potentially important information about the nature and resilience of regional traditions and populations. Another major implication is that the populations occupying the large area now submerged by the

North Sea and Channel were maintaining and following different sets of handaxe-shape traditions, perhaps two or more regional groups overlapping on this ephemeral land. This overlap might be expressed in the succession of assemblage types at Foxhall Road, Hitchin or Hoxne. That MIS 11b was potentially 'survivable', either in Britain or nearby, is also shown by the strong similarities between the fauna from MIS 11c and 11a (Schreve 2001b; Ashton et al 2008). Pan-European studies might help tease out some of these regional groups, although the cultural geography of mainland Europe is likely to be more complex than the British cul-de-sac, a northern sink zone with episodes of residency, abandonment and re-colonisation synchronised by the climatic rhythms of the Pleistocene (White 2015; Ashton 2017).

# Conclusions

Handaxes have been key to understanding archaic hominin culture and behaviour from the beginning of our discipline and Britain has a record that is extremely well suited to exploring these issues. It is a valid geographical entity, the NW uplands of the north European Plain, fringing the Atlantic and surrounded on two sides by basins that were periodically flooded. It was also occasionally glaciated and inhospitable. Consequently, it has a punctuated pattern of occupation with distinct periods of abandonment, colonisation, settlement and isolation, all of which can be synchronised to Pleistocene climatic fluctuations. Despite being a sink area at the limits of human occupation, accessible only at certain times and from certain directions, it provides insights into demographic patterns that can be isolated in far more detail than in more permanently occupied areas. It is also increasingly well dated and well researched.

Nearly two centuries of interpretation have created a (monochrome) tapestry of hominin behaviour in a landscape context, but British archaeologists had all but abandoned ideas that group-level patterns could be detected in the Acheulean, and ergo that handaxe shape contained a meaningful cultural signal. Paradoxically, it was partly the better documented record, which facilitated influential syntheses ranging from Breuil's to Wymer's and to Roe's, that exposed this lack of structure. To explain variation, archaeologists thus turned to more ecological or techno-functional explanations, attempting to find some behavioural order within the seemingly directionless and unstructured archaeological record, described, equally accurately, as both a 'bewildering variety' (Roe 1981) and a 'variable sameness' (Isaac 1977). This failure can now largely be attributed not to the absence of cultural patterns, but to the use of an incomplete Quaternary framework. The revolution brought about by correlation with the MIS record, with the greater number of climatic events thus recognisable in the terrestrial domain, eventually revealed these patterns. Now, using large datasets interpreted at different scales, we can suggest that handaxes indeed provide cultural signals.

The British handaxe record does not show linear or predictable patterns, and does not form the progressive developmental sequence expected by past cultural frameworks, but it is no longer a bewildering variety in time and space. Acheulean handaxe assemblages contain much variation, but also clear modal tendencies. Such variation is perhaps an inevitability in stone objects that were hand-made by people of all ages and skill-levels using inconsistent materials. Twisted handaxes and other time-locked forms (e.g. MIS 9 ficrons and cleavers), however, isolate clear modalities, distinctive and (often) highly accomplished forms that earlier workers would have called *fossile directeurs* (cultural markers) although we appreciate that experience has taught many colleagues to be wary of such things. We are equally aware that the persistence of local traditions over potentially vast time periods presents some scalar problems in terms of social transmission, although we are probably observing, in primary context assemblages, only a few hundred years. Handaxe assemblages thus capture hominin socio-politics at centennial scales, cross-generational histories of regional groups involving a moving roster of related individuals through time, with social transmission for particular forms biased by the success (or not) of real individuals and networks. We would further note that similar time scales are not deemed overly problematic when studying Upper Palaeolithic industries, such as the ~9000 year-long Aurignacian.

The next challenge is to understand better the social mechanisms at work, and to reexamine the European record in similar terms, using standardised methods and new analyses, acknowledging the possibility that only another bewildering variety might emerge. This was the reality faced by Paul Callow (1976), who failed to detect even a clear pointdominated/ovate-dominated division in northern France. As a potentially atypical but somewhat predictable case, Britain might just provide some of the empirical and theoretical tools necessary to unlock this complexity. Acknowledgements: We would like to thank Pascal Depaepe, Agnes Lamotte, Jean-Luc

Locht, Marie-Hélène Moncel, Marina Mosquera and Andreu Ollé for information on twisted

handaxes (or their absence) from neighbouring regions of Europe. We would also like to

thank three anonymous referees for their comments on the original version of this paper.

# Bibliography

- Agache, P. 1971. Information Archaeologiques: Circonscription de Nord et Picardie. *Gallia Prehistoire* **11**: 267-309
- Allen, P. & White, M.J. 2004. The geology of Foxhall Road and the surrounding area. In: M.J. White & S.J. Plunkett, *Miss Layard Excavates: a Palaeolithic site at Foxhall Road, Ipswich, 1903-1905*. Bristol: Western Academic and Specialist Press. pp 55-75.
- Allen, P., Bridgland, D.R, Haggart, A., Schwenninger, J-L, White, M.J., Wilson, D. & Wray, D. In press. Geological and Archaeological Re-Investigation of Foxhall Road, Ipswich, 2005, *Proceedings of the Geologists' Association*
- Antoine, P. & Limondin-Lozouet, N. 2004. Identification of MIS 11 Interglacial tufa deposit in the Somme valley (France): new results from the Saint-Acheul fluvial sequence. *Quaternaire* 15: 41-52
- Antoine, P., Lautridou, J.P., & Laurent, M., 2000. Long-Term Fluvial archives in NW France: response of the Seine and Somme rivers to tectonic movements, climatic variations and sea level changes. *Geomorphology* **33**: 183–207.
- Antoine, P, Auguste, P., Bahain, J-J., Chaussé, C., Falguëres, C., Galehb, B., Limondin-Lozouet, N., Locht, J-L., & Voinchet, P. 2010. Chronostratigraphy and palaeoenvironments of Acheulean occupations in Northern France (Somme, Seine and Yonne valleys). *Quaternary International* 223-224: 456–461.
- Antoine, P., Moncel, M-H., Locht, J-L., Limondin-Lozouet, N., Auguste, P., Stoetzel, E., Dabkowski, J., Voinchet, P., Bahain J-J., Falgueres, C. 2015. Dating the earliest human occupation of Western Europe: New evidence from the fluvial terrace system of the Somme basin (Northern France). *Quaternary International* **370**: 77–99.
- Ashton, N.M. 2017. Landscapes of habit and persistent places during MIS 11 in Europe. A return journey from Britain. In: C. Gamble, M. Pope, R. Scott, A. Shaw, & M. Bates (eds), *Crossing the Threshold*. London: Routledge.
- Ashton, N. M. & McNabb, J. 1994. Bifaces in perspective. In: N.M. Ashton & A. David (eds), *Stories in Stones*. London: Lithic Studies Society (Occasional Paper 4). pp 182-191.
- Ashton, N. & Hosfield, R. 2009. Mapping the human record in the British early Palaeolithic: evidence from the Solent River system. *Journal of Quaternary Science* **25**: 737-753.
- Ashton, N., Cook, J., Lewis, S. G. & Rose, J. 1992. *High Lodge. Excavations by G. de G. Sieveking 1962-68 and J. Cook 1988.* London: British Museum Press.
- Ashton, N., Lewis, S. G. & Parfitt, S. (eds). 1998. *Excavations at the Lower Palaeolithic site at East Farm, Barnham, 1989-1994.* London: British Museum Press
- Ashton, N., Lewis, S., Parfitt, S., Candy, I., Keen, D., Kemp, R., Penkman, K., Thomas, G., Whittaker, J.
  & White, M. J. 2005. Excavations at the Lower Palaeolithic site at Elveden, Suffolk, UK. *Proceedings of the Prehistoric Society* **71**: 1-61.
- Ashton, N., Lewis, S. G., Parfitt, S., Penkman, K. E. H. & Coope, G. R. 2008. New evidence for complex climate change in MIS 11 from Hoxne, UK. *Quaternary Science Reviews* **27**: 652-668.
- Ashton, N., Lewis, S. & Hosfield, R. 2011. Mapping the human record: population change during the later Middle Pleistocene. In: N. Ashton, S. Lewis & C. Stringer (eds), *The Ancient Human Occupation of Britain*. Amsterdam: Elsevier. pp 39-52.

- Ashton, N., Lewis, S. G., Parfitt, S. A., Davis, R. J., & Stringer, C. 2016. Handaxe and non-handaxe assemblages during Marine Isotope Stage 11 in northern Europe: recent investigations at Barnham, Suffolk, UK. *Journal of Quaternary Science* **31**: 837-843
- Boreham, S., & Gibbard P.L. 1995. Middle Pleistocene Hoxnian Stage interglacial deposits at Hitchin, Hertfordshire, England. *Proceedings of the Geologist's Association* **106**: 259-270.
- Bordes, F. 1956. Some observations on the Pleistocene succession in the Somme valley. *Proceedings* of the Prehistoric Society **22**:1-5.
- Bourdier, F. 1969. *Etude comparée des dépôts quaternaires des bassins de la Seine et de la Somme*. AGBP. Union internationale pour l'étude du quaternaire Congrès. pp169-220
- Breuil, H. 1934. De l'importance de la solifluction dans l'etude des terrains Quaternaires du nord de la France et des pays voisins. *Revue de Geographie physique et de Geologie dynamique* **7**: 269-332.
- Breuil, H., and Kelley, H. 1954. Le Paléolithique Ancien. *Bulletin de la Société Préhistorique de France* **51**: 9-26
- Bridgland, D. R. 1994. Quaternary of the Thames. London: Chapman and Hall.
- Bridgland, D. R. 1996. Dating of Lower Palaeolithic industries within the framework of the Lower Thames terrace sequence. In: N. Ashton & A. David (eds), *Stories in Stone*. London: Lithics Study Society (Occasional Paper 4). pp 28-40.
- Bridgland, D.R. 1999. 'Wealden rivers' north of the Thames: a provenance study based on gravel clast analysis. *Proceedings of the Geologists' Association* **110**: 133-148.
- Bridgland, D. R. 2003. The evolution of the River Medway, SE England, in the context of Quaternary palaeoclimate and the Palaeolithic occupation of NW Europe. *Proceedings of Geologists' Association* **114**: 23-48.
- Bridgland, D.R. 2006. The Middle and Upper Pleistocene sequence in the Lower Thames; a record of Milankovitch climatic fluctuation and early human occupation of southern Britain. Henry Stopes Memorial Lecture. *Proceedings of the Geologists' Association* **117**: 281–305.
- Bridgland, D.R. & White, M.J. 2014. Fluvial archives as a framework for the Lower and Middle Palaeolithic: patterns of British artefact distribution and potential chronological implications. *Boreas* **43**: 543–555.
- Bridgland, D.R., White, M.J.,2015. Chronological variations in handaxes: patterns detected from fluvial archives in NW Europe. *Journal of Quaternary Science* **30**: 623–638.
- Bridgland, D.R. and White, M.J. 2018. The Farnham River terrace staircase: an optimal record of the Thames Palaeolithic. *Earth Heritage* 49: 49-51
- Bridgland, D. R., Gibbard, P. L. & Preece, R. C. 1990. The Geology and Significance of the Interglacial Sediments at Little Oakley, Essex. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences* **328**, 307-339.
- Bridgland, D.R., D.H. Keen, D.C. Schreve and M.J. White. 1998. Quaternary drainage of the Kentish Stour. In: J.B. Murton, C.A. Whiteman, M.B. Bates, D.R. Bridgland, A.J. Long, M.B. Roberts and M.P. Waller, *The Quaternary of Kent and Sussex: Field Guide*. London: Quaternary Research Association. pp 39-41
- Bridgland, D. R., Allen, P., White, M. J., White, T. S. & Wenban-Smith, F. F. 2014. Wansunt Pit, Dartford Heath (TQ 513737). In: D. Bridgland, P. Allen, P. & T.S. White (eds), *The Quaternary of the Lower Thames and Eastern Essex: Field guide*. London: Quaternary Research Association. pp152-156.
- Callow, P. 1976. The Lower and Middle Palaeolithic of Britain and Adjacent Areas of Europe. Unpublished PhD Thesis, University of Cambridge.
- Callow, P. 1986. A Comparison of British and French Acheulian bifaces. In: S.N. Collcutt (ed.), *The Palaeolithic of Britain and its Nearest neighbours: Recent Trends*. Sheffield: University of Sheffield. pp 3-7.

- Candy, I. & Schreve, D. 2007. Land-sea correlation of Middle Pleistocene temperate sub-stages using high-precision uranium-series dating of tufa deposits from Southern England. *Quaternary Science Reviews* **26**: 1223-1235
- Chandler, R. H. & Leach, A. L. 1912. On the Dartford Heath Gravel and on a Palaeolithic implement factory. *Proceedings of the Geologist's Association* **23**: 102-111.
- Cliquet, D., Lautridou, J.-P., Antoine, P., Lamotte, A., Leroyer, M., Limondin-Lozouet, N., Mercier, N., 2009. La séquence loessique de Saint-Pierre-lès-Elbeuf (Normandie, France) : nouvelles données archéologiques, géochronologiques et paléontologiques. *Quaternaire* **20**: 321-343.
- Collins, D. 1969. Culture traditions and environment of early man. *Current Anthropology* **10**: 267-316.
- Commont, V. 1908. Les industries de l'ancien Saint- Acheul. L'Anthropologie XLIX: 527–572.
- Commont, V. 1909. Saint-Acheul et Montières. Notes de Geologie, de Paléontologie et de Préhistoire. Memoirs de la Société Géologique du Nord VI: 1-77
- Conway, B. 1969. Preliminary Geological investigation of Boyn Hill terrace deposits at Barnfield Pit, Swanscombe, Kent, during 1968. *Proceedings of the Royal Anthropological Institute of Great Britain and Ireland for 1968*: 59-61.
- Conway, B. 1970. Geological investigation of Boyn Hill terrace deposits at Barnfield Pit, Swanscombe, Kent, during 1969. *Proceedings of the Royal Anthropological Institute of Great Britain and Ireland for 1969:* 90-93.
- Conway, B., 1972. Geological investigation of Boyn Hill terrace deposits at Barnfield Pit, Swanscombe, Kent, during 1971. *Proceedings of the Royal Anthropological Institute of Great Britain and Ireland for 1971*: 80-85
- Conway, B., 1996. The geology outside the National Nature Reserve, 1968-72. In: B. Conway, J. McNabb & N. Ashton (eds), *Excavations at Barnfield Pit, Swanscombe, 1968-72*. London: British Museum Occasional Paper 94. pp. 67-88.
- Conway, B. & Waechter, J. d'A., 1977. Lower Thames and Medway valleys: Barnfield Pit, Swanscombe. In: ER. Shephard-Thorn & J.J. Wymer (eds.), *South East England and the Thames Valley. Guide Book for Excursion A5. X INQUA Congress*. Norwich: Geoabstracts. pp. 38-44.
- Dewey, H., 1959. *Palaeolithic Deposits of the Thames at Dartford Heath and Swanscombe*. Unpublished edited text of Henry Stopes Memorial Lecture, read to the Geologists' Association and available from archive.
- Dunbar, R.I.M., Gamble, C., and Gowlett, J.A.J. 2014. *From Lucy to Language: the Benchmark Papers*. Oxford University Press: Oxford.
- Evans, J. 1872. Ancient Stone Implements, Weapons and Ornaments of Great Britain. 1st Edition. London: Longmans & Co.
- Field, D., Nicolayson, P., & Cotton, J. 1999. The Palaeolithic sites at Limpsfield, Surrey: an analysis of the artefacts collected by A.M. Bell. *Surrey Archaeological Collections* **86**: 1-32.
- Gallotti, R, Collina, C, Raynal J\_P, Kieffer, G., Geraads, D., & Piperno, M. 2010. The Early Middle Pleistocene Site of Gombore II (Melka Kunture, Upper Awash, Ethiopia) and the Issue of Acheulean Bifacial Shaping Strategies. *African Archaeology Reviews* **27**:291–322
- Gamble, C. 1999. The Palaeolithic Societies of Europe. Cambridge: Cambridge University Press.
- Gamble, C. 2002. Early beginnings 500-35,000 years ago. In: P. Slack & R. Ward (eds), *The Peopling of Britain: the shaping of a human landscape. The Linacre Lectures 1999*. Oxford: Oxford University Press. pp 11-37.
- Gossling, F. 1940. A contribution to the Pleistocene history of the Upper Darent valley. *Proceedings of the Geologists' Association* **51**: 311-340.
- Green, C. P., Gibbard, P. L. & Bishop, B. J. 2004. Stoke Newington: geoarchaeology of the Palaeolithic 'floor'. *Proceedings of the Geologists' Association* **115**: 193-208.
- Green, C. P., Branch, N. P., Coope, G. R., Field, M. H., Keen, D. H., Wells, J. M., Schwenninger, J.-L., Preece, R. C., Schreve, D. C., Canti, M. G. & Gleed-Owen, C. 2006. Marine Isotope Stage 9 environments of fluvial deposits at Hackney, north London, UK. *Quaternary Science Reviews* 25, 89-113.

- Hardaker, T. 2012. Artefacts from the present land surface at the Palaeolithic site of Warren Hill, Suffolk, England. *Proceedings of the Geologists' Association* 123: 692-713
- Hosfield, R. & Chambers, J., 2009. Genuine Diversity? The Broom Biface Assemblage. *Proceedings of the Prehistoric Society* 75: 65–100
- Hosfield, R.T. & Green, C.H. 2014. *Quaternary History and Palaeolithic Archaeology in the Axe Valley at Broom, South West England*. Oxford: Oxbow.
- Hubbard, R. 1972. An interim report on the pollen record at Swanscombe. *Proceedings of the Royal Anthropological Institute of Great Britain and Ireland for 1971*: 79
- Hubbard, R. 1982. The environmental evidence from Swanscombe and its implications forPalaeolithic Archaeology. In: P.E. Leach (ed.), *Archaeology in Kent to AD 1500*. London: Council for British Archaeology Research Report 48. pp 3-7.
- Hughes, T.M. 1916. The Gravels of East Anglia. Cambridge: Cambridge University Press.
- Isaac, G. 1977. Olorgesailie: Archaeological Studies of a Middle Pleistocene Lake Basin in Kenya. Chicago: Chicago University Press.
- Keeley, L.H., 1980. Experimental Determination of Stone Tool Uses. Chicago: University of Chicago Press.
- Keen, D. H. 1990. Significance of the record provided by Pleistocene fluvial deposits and their included molluscan faunas for palaeo-environmental reconstruction and stratigraphy: a case study from the English Midlands. *Palaeogeography, Palaeoclimate, Palaeoeology* 80: 25-34.
- Kerney, M.P. 1971. Interglacial deposits at Barnfield Pit, Swanscombe, and their molluscan fauna. *Journal of the Geological Society of London* **127**: 69-86.
- King, W. B. R. & Oakley, K. P. 1936. The Pleistocene Succession in the Lower parts of the Thames Valley. *Proceedings of the Prehistoric Society* **2**: 52-76.
- Leach, A. L. 1913. On buried channels in the Dartford Heath Gravel: Together with the report of an excursion to Dartford Heath, May 17th, 1913. *Proceedings of the Geologists' Association* **24**: 337-344.
- Limondin-Lozouet, N., Antoine, P., Bahain, J.-J., Cliquet, D., Coutard, S., Dabkowski, J., Ghaleb, B., Locht, J.-L., Nicoud, E., Voinchet, P., 2015. North-West European MIS 11 malacological successions: a framework for the timing of Acheulean settlements. *Journal of Quaternary Science* **30**: 702–712.
- Lycett, S.J. & von Cramon-Taubadel, N. 2008. Acheulean variability and hominin dispersals: a modelbound approach. *Journal of Archaeological Science* **35**: 553-562

McNabb, J., Hosfield, R., Dearling, K., Barker, D., Strutt, K., Cole, J., Bates, M. & Toms, P. 2013. Recent Work at the Lower Palaeolithic Site of Corfe Mullen, Dorset, England. *Proceedings of the Prehistoric Society* **78**: 35–50.

- Mangerud, J. 1991. The Last Interglacial/Glacial cycle in northern Europe. In: L.C.K. Shane & E.J. Cushing (eds), *Quaternary Landscapes*. London: Bellhaven Press. pp 38-75.
- Newton, W.M., 1901. The occurrence in a very limited area of the rudest with the finer forms of worked stones. *Man* 1: 81-82.
- Oakley, K.P..1939. 'Geology and Palaeolithic studies', in K.P. Oakley, W.F. Rankine and A.W.G. Lowther (eds), *A Survey of the Prehistory of the Farnham District (Surrey)*. Guildford: Butler and Tanner. pp 3–58.
- Penkman, K. E. H., Preece, R. C., Bridgland, D. R., Keen, D. H., Meijer, T., Parfitt, S. A., White, T. S. & Collins, M. J. (2013). An aminostratigraphy for the British Quaternary based on Bithynia opercula. *Quaternary Science Reviews* **61**: 111-134.
- Pettitt, P. B. & White, M. J. (2012). *The British Palaeolithic: hominin societies on the edge of the Pleistocene world*. London: Routledge.
- Preece, R. C., Gowlett, J. A. J., Parfitt, S. A., Bridgland, D. R. & Lewis, S. G. 2006. Humans in the Hoxnian: habitat, context, and fire use at Beeches Pit, West Stow, Suffolk, UK. *Journal of Quaternary Science* **21**: 485-496.

- Prestwich, J. 1891. On the Age, Formation, and Successive Drift-Stages of the Valley of the Darent; with Remarks on the Palaeolithic Implements of the District, and on the Origin of its Chalk Escarpment. *Quarterly Journal of the Geological Society* **47**: 126-163.
- Roberts, M. B. and Parfitt, S.A. 1999. *Boxgrove; A Middle Pleistocene Hominid Site at Eartham Quarry, Boxgrove, West Sussex*. London: English Heritage.
- Roe, D. A. 1968. British Lower and Middle Palaeolithic handaxe groups. *Proceedings of the Prehistoric Society* **34**: 1-82.
- Roe, D. A. 1981. *The Lower and Middle Palaeolithic Periods in Britain*. London: Routledge and Kegan Paul.
- Shipton, C and Nielsen, M. 2015. The evolutionary origins of overimitation and shared intentionality. *Human Nature* **26**: 331-345
- Schreve, D. C. 2001a. Differentiation of the British late Middle Pleistocene interglacials: the evidence from mammalian biostratigraphy. *Quaternary Science Reviews* **20**: 1693-1705.
- Schreve, D. C. 2001b. Mammalian evidence from Middle Pleistocene fluvial sequences for complex environmental change at the oxygen isotope substage level. *Quaternary International* **79**: 65-74.
- Schreve, D.C., Bridgland, D.R., 2002. Correlation of English and German Middle Pleistocene fluvial sequences based on mammalian biostratigraphy. *Netherlands Journal of Geosciences* 81: 357– 373.
- Schreve, D. C., Bridgland, D. R., Allen, P., Blackford, J. J., Gleed-Owen, C. P., Griffiths, H. I., Keen, D. H.
  & White, M. J. 2002. Sedimentology, palaeontology and archaeology of late Middle Pleistocene River Thames terrace deposits at Purfleet, Essex, UK. *Quaternary Science Reviews* 21: 1423-1464.
- Singer, R., Gladfelter, B.G. and Wymer, J.J. 1993. *The Lower Palaeolithic Site at Hoxne, England*. Chicago: Chicago Press
- Smith, R.H. & Dewey, H., 1913. Stratification at Swanscombe: report on excavation made on behalf of the British Museum and H.M. Geological Survey. Archaeologia **64**: 177–204.
- Smith, R.H. & Dewey, H., 1914. The High Terrace of the Thames: report on excavations made on behalf of the British Museum and H.M. Geological Survey in 1913. Archaeologia **65**: 187–212.
- Smith, W. G. 1894. *Man the Primeval Savage: his haunts and relics from the hill-tops of Bedfordshire to Blackwall*. London: Stanford.
- Spurrell, F.C.J. 1883. Palaeolithic implements from West Kent. Archaeologia Cantiana 15: 89-103
- Sutcliffe, A.J. & Kowalski, K. 1976. Pleistocene Rodents of the British Isles. *Bulletin of the British Museum (Natural History) Geology* **27**: 31-147.
- Tester, P. J. 1951. Palaeolithic flint implements from the Bowman's Lodge Gravel Pit, Dartford Heath. *Archaeologia Cantiana* **63**: 122-134.
- Tester, P.J. 1953. The discovery of Acheulian implements in the deposits of the Dartford Heath terrace. *Archaeologia Cantiana* **66**: 72–76.
- Tester, P.J. 1975. Further consideration of the Bowman's Lodge Industry. *Archaeologia Cantiana* **91**: 29–39.
- Turner, C. 1985. Problems and pitfalls in the application of palynology to Pleistocene archaeological sites in WesternEurope. In: J. Renault-Miskovsky, M. Bui-Thi & M. Girard (eds), Palynologie Archéologique: Notes et monographies techniques 17. Paris: CNRS. pp. 347-373
- Walter, D. 1996. Twisted bifaces: an analysis of intentionality and functional efficiency of twisted bifaces in the British Lower Palaeolithic. Unpublished MA Thesis. University of London.
- Wenban-Smith, F. 2004. Handaxe typology and the Lower Palaeolithic cultural development: ficrons, cleavers and two giant handaxes from Cuxton. *Lithics* **25**: 11-21.
- Wenban-Smith, F.F. & Bridgland, D.R. 2001. Palaeolithic archaeology at the Swan Valley Community School, Swanscombe, Kent. *Proceedings of the Prehistoric Society* **67**: 219–225.

- Wenban-Smith, F.F., Gamble, C.S., and ApSimon, A. 2000. The Lower Palaeolithic site at Red Barns: bifacial technology, raw material quality and the organisation of archaic behaviour. *Proceedings of the Prehistoric Society* **66**: 209-256.
- West, R. G. 1955. The Pleistocene Vegetation and Geology of East Anglia. Cambridge: Cambridge University
- Westaway, R., Bridgland, D. & White, M. J. (2006). The Quaternary uplift History of central southern England: evidence from the terraces of the Solent River system and nearby raised beaches. *Quaternary Science Reviews* **25**: 2212-2250.
- White, M. J. 1998a. Twisted Ovate Bifaces in the British Lower Palaeolithic: Some Observations and Implications. In: N.M. Ashton, F. Healy and P. Pettitt (eds), *Stone Age Archaeology: Essays in honour of John Wymer*. Oxford: Oxbow Books. pp 98-104.
- White, M. J. 1998b. On the significance of Acheulean biface variability in Southern Britain. *Proceedings of the Prehistoric Society* **64**: 15-44.
- White, M. J. 2000. The Clactonian Question: On the interpretation of core-and-flake assemblages in the British Lower Palaeolithic. *Journal of World Prehistory* **14**, 1-63.
- White, M.J. 2015. Dancing to the Rhythms of the Biotidal Zone': Settlement history and culture history in Middle Pleistocene Britain. In: F. Coward, F. Wenban-Smith, R. Hosfield & M. Pope (eds), Society, Settlement and Cognition. Cambridge: Cambridge University Press. pp 154-173
- White, M.J. and Schreve, D.C. 2000. Island Britain-Peninsula Britain: palaeogeography, colonisation and the Lower Palaeolithic settlement of the British Isles. *Proceedings of the Prehistoric Society* 66: 1-28
- White, M. J. & Plunkett.S.J. 2004. *Miss Layard Excavates: a Palaeolithic site at Foxhall Road, Ipswich, 1903-1905*. Liverpool: WASP.
- White, M. J., Bridgland, D. R., Ashton, N., McNabb, J. & Berger, M. 1995. Wansunt Pit, Dartford Heath (TQ 513737). In: D.R. Bridgland, P. Allen & A. Haggart (eds.) *The Quaternary of the Lower Reaches of the Thames: Field Guide*. London: Quaternary Research Association. pp 117-128.
- White, M., Scott, R. & Ashton, N. 2006. The Early Middle Palaeolithic in Britain: archaeology, settlement history and human behaviour. *Journal of Quaternary Science* **21**: 525-542.
- White, M. J., Bridgland, D. R., Schreve, D. C., White, T. S. & Penkman, K. E. H. 2017. Well-dated fluvial sequences as templates for patterns of handaxe distribution: understanding the record of Acheulean activity in the Thames and its correlatives. *Quaternary International* http://dx.doi.org/ 10.1016/j.quaint.2017.03.049.
- White, T. S., Preece, R. C. & Whittaker, J. E. 2013. Molluscan and ostracod successions from Dierden's Pit, Swanscombe: insights into the fluvial history, sea-level record and human occupation of the Hoxnian Thames. *Quaternary Science Reviews* **70**: 73-90.
- Wiessner, P. 1983. Style and social information in Kalahari San Projectile Points. *American Antiquity* **48**: 253-276
- Wooldridge, S.W. & Linton, D.L. 1955. *Structure, surface and drainage in southeast England* (2nd edn). London: G. Phillip.
- Worssam, B. 1973. A new look at river capture and at the denudation history of the Weald. *Report of the institute of Geological Sciences* 73/17.
- Worssam, B. and Taylor, J. 1969. *Geology of the Country Around Cambridgeshire*. Memoir of the Geological Survey of Great Britain. London. HMSO
- Wymer, J. 1968. *Lower Palaeolithic archaeology in Britain as represented by the Thames Valley*. London: John Baker.
- Wymer, J. J. 1985. Palaeolithic Sites of East Anglia. Norwich: Geobooks.
- Wymer, J. J. 1999. *The Lower Palaeolithic Occupation of Britain*. Salisbury: Wessex Archaeology and English Heritage.

# **FIGURE & TABLE CAPTIONS**

Figure 1: Map of Britain showing main sites mentioned in the text and tables

Figure 2: The Twisted Handaxe Knapping Schema: A) Diagram showing the conceptual pattern of inflection points, and the edge configuration of a twisted handaxe, in this case describing a Z-twist (redrawn after Gallotti et al 2010). B) The series of rotations and inversions employed in the 'Classic Twisted Strategy' (White 1998a). 1: The first quarter is knapped; 2: The piece is turned over through the long axis, presenting the opposite margin and other face for knapping. This quarter is knapped; 3: The piece is rotated 180°, presenting the quarter diagonally opposite 2 for knapping; 4: The handaxe is inverted through the long axis once more, presenting the final quarter for knapping. Each quarter is knapped in an opposite direction to that adjacent to it, and in the same direction as that diagonally opposed it.

Figure 3: A trio of 'identical' twisted ovate handaxes from Foxhall Road. These were found by Nina Layard in 1903, lying side by side. Photographs are the pair of handaxes in Ipswich Museum (Accession Number 1920-76-35/Layard #48 & 1920-76-37/Layard #42) outline drawing belongs to the handaxe given to Sir John Evans and now housed in the Ashmolean Museum, Oxford (photographs: MJW).

Figure 4: Schematic correlation diagram for sites in the Thames Valley and East Anglia, showing the chronological distribution of different archaeological industries (based on correlations in Preece et al 2006, Pettitt and White 2012; White et al 2013). Blue shading = cold climate. Yellow shading = palaeo-environmental evidence for temperate climate.

Figure 5: The Sequence at Barnfield Pit, Swanscombe showing Phases and Deposits, Archaeology, Climate, Pollen Zones and MIS (after Bridgland 1994)

Table 1: Assemblages showing high frequencies of twisted handaxe assemblages, organised by region and age correlations

Table 2: Percentage of twisted ovate in 30 well-studied sites, for which it is possible to propose an age based on lithostratigraphical, biostratgraphical or chronometrical methods. Artefact typology has played no critical role in assigning ages. Data from White (1996 &

unpublished), Roe's (1968), Wymer (1968), Field et al. (1991); Hosfield and Green's (2014) They are organised according to probable age.

\*= Almost certainly belongs to or is derived from the underlying Grey Clays. Mixing of older material is also suspected at Furze Platt, while collection or contextual issues surround Barton Cliff, Warren Hill and Limpsfield. The twists from Broom are not considered to conform to the technological strategy seen in MIS 11.

Table 3: Summary data showing percentages of twisted handaxes in different marine isotope stages. Due to the atypical nature of the Broom twists, and the element of circularity in the dating of the Limpsfield assemblage (Bridgland 2003), figures are given both with and without these samples.

Region	Site (% twisted handaxes)	MIS stage or sub-	
		stage	
East Anglia & adjacent counties	Barnham, Suffolk (33%)	MIS 11c	
East Anglia & adjacent counties	Elveden, Suffolk (36-40%)	MIS 11c	
East Anglia & adjacent counties	Foxhall Road Grey Clays, Suffolk (39%)	MIS 11c	
East Anglia & adjacent counties	Hitchin, Hertsforshire (45% of ovate assemblage)	MIS 11c	
East Anglia & adjacent counties	Santon Downham, Suffolk (18%)	Post-MIS 12	
East Anglia & adjacent counties	Allington Hill, Cambridgeshire (46%)	Post-MIS 12	
South the Thames	Swanscombe Barnfield Upper Loam, Kent (22%)	MIS 11a	
South of the Thames	Swanscombe Rickson's Upper Gravel, Kent (16%)	MIS11a	
South of the Thames	Swanscombe Dierden's Stony Loam, Kent (16%)	MIS 11a	
South of the Thames	Wansunt Pit, Kent (28%)	MIS 11a	
South of the Thames	Bowman's Lodge Pit, Kent (31%)	MIS 11a	
South of the Thames	Limpsfield, Kent (54%)	MIS 11	
South of the Thames	Farnham Terrace B (nd)	MIS 11	

TABLE 1

Site	Probable Age	Roe Group	Region	% Twists	Total Sample	Number Twists	Reference for dating
Baker's Farm	MIS9	I	Thames	0	239	0	Bridgland 1994
Stoke Newington	MIS9	I	Thames	0	63	0	Green et al 2004, 2006
Wolvercote	MIS9	ш	Thames	0	47	0	Bridgland 1996
Cuxton	MIS9-8	1	Thames	0	183	0	Wenban-Smith 2004
Furze Platt	MIS9	1	Thames	0.4	461	2	Bridgland 1994
Broom	MIS9-8	IV	South-west	3	995	30	Hosfield and Green 2014
Hoxne	MIS11a	П	East Anglia	3.5	111	4	Ashton et al 2008
Swanscombe UL	MIS11a	VI	Thames	22	18	4	See text
Bowman's Lodge	MIS11a	VI	Thames	33	30	10	See text
Wansunt	MIS11a	VI	Thames	28	32	8	See text
Swanscombe Ricksons/Dierdens	MIS11a	VI	Thames	16	50	8	See text
Limpsfield	MIS11	VI	Medway- Darent	55	283	157	Field et al 1999
Barton Cliff	MIS11	IV	Solent	8	110	9	Westaway et al 2006
Dovercourt	MIS11	П	East Anglia	4	165	6	Bridgland et al 1990
Elveden	MIS11c	VI	East Anglia	36	74	27	See text
Barnham	MIS11c		East Anglia	33	3	1	See text
Foxhall Road Grey Clay	MIS11c	VI	East Anglia	39	18	7	See text
Foxhall Road Red Gravel	MIS11c	П	East Anglia	5	17	1*	See text
Hitchin Ovates	MIS11c	VI	East of England	38	24	9	See text
Swanscombe MG	MIS11c	п	Thames	0	159	0	See text
Chadwell St Mary	MIS11	П	East Anglia	1	99	1	Bridgland 1994
Warren Hill Fresh	MIS13	VII	East Anglia	1.6	642	10	
Warren Hill Worn	MIS13+	v	East Anglia	0	116	0	
High Lodge	MIS13	VII	East Anglia	3	68	2	Ashton et al 1992
Corfe Mullen	MIS13	VII	Southwest	7	45	3	McNabb et al 2013
Boxgrove	MIS13	VII	Southeast	0	81	0	Roberts and Parfitt 1999
Fordwich	MIS13+	v	Southeast	1	194	1	Bridgland et al 1998
Highlands Farm	MIS13/12	VII	Thames	0	200	0	Wymer 1999
Farnham A	MIS13+	v	South	0	39	0	Bridgland and White 2018
Caversham	MIS13/12	VI	Thames	3	156	2	Wymer 1999

Twisted handaxes as % of entire sample (n=4722)	Twisted handaxes as % of MIS 13 sample (n=1541)	Twisted handaxes as % of combined MIS 11 sample (n=1193)	Twisted handaxes as % of MIS 11c in East Anglia/Herts	Twisted handaxes as % of MIS 11a sites South of the Thames (n=130)	Twisted handaxes as % of MIS 9 sample (n=1988)
6.3 (3.2 excl. Limpsfield)	1	21 (10.4 excl. Limpfield)	33	23	1.6% (0.1% excl. Broom twists)
Number Twisted Handaxes	% from MIS 13	% from MIS 11	% from MIS 9		
N=302	5.9	83.4	10.5		
Excluding Broom twists	% from MIS 13	% from MIS 11	% from MIS 9		
N=270	6.6	93.3	0.7		
Excluding Limpsfield	% from MIS 13	% from MIS 11	% from MIS 9		
=145	8	69	22	]	
Excluding Broom twists and Limpsfield	% from MIS 13	% from MIS 11	% from MIS 9		
115	10.4	87.8	1.7		

Table 3:





В

Α













Phase and Units	Archaeology	Climate	<b>Pollen Zone (after Ashton et al 2008a)</b>	MIS
llle: Higher Loams		?		MICAO
IIId:Upper Gravel	Acheulean (derived)	Cold		
llic: Upper Loam	Acheulean (Roe's Group VI)	Temperate		MIS11a
llib: 'Upper Sands' channel		Cold		MIS11b
Illa: Soli ucted Clay		Cold		
llb: Upper Middle Gravel	Acheulean (Roe's Group II )	Cool	Hollib-IV	
lla: Lower Middle Gravel	Acheulean (Roe's Group II)	Temperate	Hollia-b	
1e:Weathered Lower Loam				MIS11c
1d: Lower Loam	Clactonian	Temperate	Ho∎b-c	
1c: Lower Gravel Midden	Clactonian	Temperate		
1b: Lower Gravel	Clactonian	Temperate	Hol	
1a: Basal Gravel		Cold/Temperate	   ?1an	F
 Thanet Sand				MIS12