1	Glacial geomorphological mapping:
2	a review of approaches and frameworks for best practice
3	
4	Benjamin M.P. Chandler ^{1*} , Harold Lovell ² , Clare M. Boston ² , Sven Lukas ³ , Iestyn D. Barr ⁴ ,
5	Ívar Örn Benediktsson ⁵ , Douglas I. Benn ⁶ , Chris D. Clark ⁷ , Christopher M. Darvill ⁸ ,
6	David J.A. Evans ⁹ , Marek W. Ewertowski ¹⁰ , David Loibl ¹¹ , Martin Margold ¹² , Jan-Christoph Otto ¹³ ,
7	David H. Roberts ⁹ , Chris R. Stokes ⁹ , Robert D. Storrar ¹⁴ , Arien P. Stroeven ^{15, 16}
8	
9	¹ School of Geography, Oueen Mary University of London, Mile End Road, London, E1 4NS, UK
10	² Department of Geography, University of Portsmouth, Portsmouth, UK
11	³ Department of Geology, Lund University, Lund, Sweden
12	⁴ School of Science and the Environment, Manchester Metropolitan University, Manchester, UK
13	⁵ Institute of Earth Sciences, University of Iceland, Reykjavík, Iceland
14	⁶ Department of Geography and Sustainable Development, University of St Andrews, St Andrews, UK
15	⁷ Department of Geography, University of Sheffield, Sheffield, UK
16	⁸ Geography, School of Environment, Education and Development, University of Manchester, Manchester, UK
17	⁹ Department of Geography, Durham University, Durham, UK
18	¹⁰ Faculty of Geographical and Geological Sciences, Adam Mickiewicz University, Poznań, Poland
19	¹¹ Department of Geography, Humboldt University of Berlin, Berlin, Germany
20	¹² Department of Physical Geography and Geoecology, Charles University, Prague, Czech Republic
21	¹³ Department of Geography and Geology, University of Salzburg, Salzburg, Austria
22	¹⁴ Department of the Natural and Built Environment, Sheffield Hallam University, Sheffield, UK
23	¹⁵ Geomorphology & Glaciology, Department of Physical Geography, Stockholm University, Stockholm, Sweden
24	¹⁶ Bolin Centre for Climate Research, Stockholm University, Stockholm, Sweden
25	
26	*Corresponding author. Email: b.m.p.chandler@qmul.ac.uk
27	
28	Abstract
29	
30	Geomorphological mapping is a well-established method for examining earth surface processes
31	and landscape evolution in a range of environmental contexts. In glacial research, it provides
32	crucial data for a wide range of process-oriented and palaeoglaciological reconstruction studies;
33	in the latter case providing an essential geomorphological framework for establishing glacial
34	chronologies. In recent decades, there have been significant developments in remote sensing
35	and Geographical Information Systems (GIS), with a plethora of high-quality remotely-sensed
36	datasets now (often freely) available. Most recently, the emergence of unmanned aerial vehicle
37	(UAV) technology has allowed sub-decimetre scale aerial images and Digital Elevation Models
38	(DEMs) to be obtained. Traditional field mapping methods still have an important role in

glacial geomorphology, particularly in cirque glacier, valley glacier and icefield/ice-cap outlet settings. Field mapping is also used in ice sheet settings, but often takes the form of necessarily highly-selective ground-truthing of remote mapping. Given the increasing abundance of datasets and methods available for mapping, effective approaches are necessary to enable assimilation of data and ensure robustness. In this contribution, we provide a review and assessment of the various glacial geomorphological methods and datasets currently available, with a focus on their applicability in particular glacial settings. We distinguish two overarching 'work streams' that recognise the different approaches typically used in mapping landforms produced by ice masses of different sizes: (i) mapping of ice sheet geomorphological imprints using a combined remote sensing approach, with some field checking (where feasible); and (ii) mapping of alpine and plateau-style ice mass (cirque glacier, valley glacier, icefield and ice-cap) geomorphological imprints using remote sensing and considerable field mapping. Key challenges to accurate and robust geomorphological mapping are highlighted, often necessitating compromises and pragmatic solutions. The importance of combining multiple datasets and/or mapping approaches is emphasised, akin to multi-proxy/-method approaches used in many Earth Science disciplines. Based on our review, we provide idealised frameworks and general recommendations to ensure best practice in future studies and aid in accuracy assessment, comparison and integration of geomorphological data. These will be of particular value where geomorphological data are incorporated in large compilations and subsequently used for palaeoglaciological reconstructions. Finally, we stress that robust interpretations of glacial landforms and landscapes invariably requires additional chronological and/or sedimentological evidence, and that such data should be collected as part of a coupled inductive-deductive approach.

Keywords: glacial geomorphology; geomorphological mapping; GIS; remote sensing; field mapping

- 76 1. Introduction
- 77

78 *1.1 Background and importance*

79

Mapping the spatial distribution of landforms and features through remote sensing and/or field-based approaches is a well-established method in Earth Sciences to examine earth surface processes and landscape evolution (e.g. Kronberg, 1984; Hubbard and Glasser, 2005; Smith et al., 2011). Moreover, geomorphological mapping is utilised in numerous applied settings, such as natural hazard assessment, environmental planning and civil engineering (e.g. Kienholz, 1977, Finke, 1980; Paron and Claessens, 2011; Marc and Hovius, 2015; Griffiths and Martin, 2017).

86

87 Two overarching traditions exist in geomorphological mapping: Firstly, the classical approach 88 involves mapping all geomorphological features in multiple thematic layers (e.g. landforms, breaks of 89 slope, slope angles, drainage), regardless of the range of different processes responsible for forming 90 the landscape. This approach to geomorphological mapping has been particularly widely used in 91 mainland Europe and has resulted in the creation of national legends to record holistic 92 geomorphological data that may be comparable across much larger areas or between studies (Demek, 93 1972; van Dorsser and Salomé, 1973; Leser and Stäblein, 1975; Klimaszewski, 1990; Schoeneich, 94 1993; Kneisel et al., 1998; Gustavsson et al., 2006; Raczkowska and Zwoliński, 2015). The second 95 approach involves more detailed, thematic geomorphological mapping commensurate with particular 96 research questions; for example, the map may have an emphasis on mass movements or glacial and 97 periglacial landforms and processes. Such a reductionist approach is helpful in ensuring a map is not 98 'cluttered' with less relevant data that may in turn make a multi-layered map unreadable (e.g. Kuhle, 99 1990; Robinson et al., 1995; Kraak and Ormeling, 2006). In recent years, the second approach has 100 become much more widespread due to increasing specialisation and thus forms the basis for this 101 review, which focuses on geomorphological mapping in glacial environments.

102

In glacial research, the production and analysis of geomorphological maps provide a wider contextand basis for various process-oriented and palaeoglaciological studies, including:

- 105
- (1) analysing glacial sediments and producing process-form models (e.g. Price, 1970; Benn,
 107 1994; Lukas, 2005; Benediktsson et al., 2016);
- (2) quantitatively capturing the pattern and characteristics ('metrics') of landforms to understand
 their formation and evolution (e.g. Spagnolo et al., 2014; Ojala et al., 2015; Ely et al., 2016a);
- (3) devising glacial landsystem models that can be used to elucidate former glaciation styles or
 inform engineering geology (e.g. Eyles, 1983; Evans et al., 1999; Evans, 2017; Bickerdike et al., 2018);

- (4) reconstructing the extent and dimensions of former or formerly more extensive ice masses
 (e.g. Dyke and Prest, 1987a; Kleman et al., 1997; Houmark-Nielsen and Kjær, 2003; Benn
 and Ballantyne, 2005; Glasser et al., 2008; Clark et al., 2012);
- (5) elucidating glacier and ice sheet dynamics, including advance/retreat cycles, flow
 patterns/velocities and thermal regime (e.g. Kjær et al., 2003; Kleman et al., 2008, 2010;
 Evans, 2011; Boston, 2012a; Hughes et al., 2014; Darvill et al., 2017);
- (6) identifying sampling locations for targeted numerical dating programmes and ensuring robust
 chronological frameworks (e.g. Owen et al., 2005; Barrell et al., 2011, 2013; Garcia et al.,
 2012; Kelley et al., 2014; Stroeven et al., 2014; Blomdin et al., 2016a; Gribenski et al., 2016,
 2017);
- (7) calculating palaeoclimatic variables for glaciated regions, namely palaeotemperature and
 palaeoprecipitation (e.g. Kerschner et al., 2000; Bakke et al., 2005; Stansell et al., 2007; Mills
 et al., 2012; Boston et al., 2015)
- (8) providing parameters to constrain and test numerical simulations of ice masses (e.g. Kleman
 et al., 2002; Napieralski et al., 2007a; Golledge et al., 2008; Stokes and Tarasov, 2010;
 Livingstone et al., 2015; Seguinot et al., 2016; Patton et al., 2017a).
- 129

130 Thus, accurate representation of glacial and associated landforms is crucial to producing 131 geomorphological maps of subsequent value in a wide range of glacial research. This is exemplified 132 in glacial geochronological investigations, where a targeted radiometric dating programme first 133 requires a clear geomorphological (and/or stratigraphic) framework and understanding of the 134 relationships and likely relative ages of different sediment-landform assemblages. In studies that 135 ignore this fundamental principle, it can be challenging to then reconcile any scattered or anomalous 136 numerical ages with a realistic geomorphological interpretation, as the samples have been obtained 137 without a clear genetic understanding of the landforms being sampled (see Boston et al., 2015, for 138 further discussion).

139

140 The analysis of geomorphological evidence has been employed in the study of glaciers and ice sheets 141 for over 150 years, with the techniques used in geomorphological mapping undergoing a number of 142 significant developments in that time. The earliest geomorphological investigations involved intensive 143 field surveys (e.g. Close, 1867; Penck and Brückner, 1901/1909; De Geer, 1910; Trotter, 1929; Caldenius, 1932; Raistrick, 1933), before greater efficiency was achieved through the development of 144 145 aerial photograph interpretation from the late 1950s onwards (e.g. Lueder, 1959; Price, 1963; Welch, 146 1967; Howarth, 1968; Prest et al., 1968; Sugden, 1970; Sissons, 1977a; Prest, 1983; Kronberg, 1984; 147 Mollard and Janes, 1984). Satellite imagery and digital elevation models (DEMs) have been in widespread usage since their development in the late 20th Century and have, in particular, helped 148 149 revolutionise our understanding of palaeo-ice sheets (e.g. Barents-Kara Ice Sheet: Winsborrow et al.,

150 2010; British Ice Sheet: Hughes et al., 2014; Cordilleran Ice Sheet: Kleman et al., 2010; 151 Fennoscandian Ice Sheet: Stroeven et al., 2016; Laurentide Ice Sheet: Margold et al., 2018; 152 Patagonian Ice Sheet: Glasser et al., 2008). In recent times, increasingly higher-resolution DEMs have 153 become available due to the adoption of Light Detection and Ranging (LiDAR) technology (e.g. 154 Salcher et al., 2010; Jónsson et al., 2014; Miller et al., 2014; Dowling et al., 2015; Hardt et al., 2015; 155 Putninš and Henriksen, 2017) and Unmanned Aerial Vehicles (UAVs) (e.g. Chandler et al., 2016a; 156 Evans et al., 2016a; Ewertowski et al., 2016; Tonkin et al., 2016; Ely et al., 2017). Aside from 157 improvements to remote sensing technologies, the last decade has seen a revolution in data 158 accessibility, with the proliferation of freely available imagery (e.g. Landsat data), freeware mapping 159 platforms (e.g. Google Earth) and open-source Geographical Information System (GIS) packages 160 (e.g. OGIS). As a result, tools for glacial geomorphological mapping are becoming increasingly 161 accessible, both practically and financially.

162

163 Field mapping remains a key component of the geomorphological mapping process, principally in the 164 context of manageable study areas relating to alpine- and plateau-style ice masses, i.e. cirque glaciers, valley glaciers, icefields and ice-caps (e.g. Bendle and Glasser, 2012; Boston, 2012a, b; Jónsson et al., 165 166 2014; Pearce et al., 2014; Gribenski et al., 2016; Lardeux et al., 2016; Chandler and Lukas, 2017). 167 This approach is also employed in ice sheet settings, but typically in the form of selective ground 168 checking of mapping from remotely-sensed data or focused mapping of regional sectors (e.g. Stokes 169 et al., 2013; Bendle et al., 2017a; Pearce et al., 2018). Frequently, field mapping is conducted in 170 tandem with sedimentological investigations (see Evans and Benn, 2004, for methods), providing a 171 means of testing preliminary interpretations and identifying problems for specific (and more detailed) 172 studies. This interlinked approach is particularly powerful and enables robust interpretations of 173 genetic processes, glaciation styles and/or glacier dynamics (e.g. Benn and Lukas, 2006; Evans, 2010; 174 Benediktsson et al., 2010, 2016; Gribenski et al., 2016). In this context, it is worth highlighting the 175 frequent use of the term 'sediment-landform assemblage' (or 'landform-sediment assemblage') as 176 opposed to 'landform' in glacial geomorphology, underlining the importance of studying both surface 177 form and internal composition (e.g. Evans, 2003a, 2017; Benn and Evans, 2010; Lukas et al., 2017).

178

Geomorphological mapping using a combination of field mapping and remotely-sensed data 179 180 interpretation (hereafter 'remote mapping'), or a number of remote sensing methods, permits a holistic 181 approach to mapping, wherein the advantages of each method/dataset can be combined to produce an 182 accurate map with robust genetic interpretations (e.g. Boston, 2012a, b; Darvill et al., 2014; Storrar 183 and Livingstone, 2017). As such, approaches are required that allow the accurate transfer and 184 assimilation of data from these various sources, particularly where data are transferred from analogue 185 (e.g. hard-copy aerial photographs) to digital format. Apart from a few recent exceptions for specific 186 locations (e.g. the Scottish Highlands: Boston, 2012a, b; Pearce et al., 2014), there has been limited

explicit discussion of the approaches used to integrate geomorphological data in map production (i.e. the relative contributions of different methods and/or datasets and their associated uncertainties), with many contributions simply stating that the maps were produced through fieldwork and/or remote sensing (e.g. Ballantyne, 1989; Lukas, 2007a; Evans et al., 2009a; McDougall, 2013). Given the diversity of scales, data sources and research questions inherent in glacial geomorphological research, and the increasing abundance of high-quality remotely-sensed datasets, finding the most cost- and time- effective approach is difficult, especially for researchers new to the field.

194

195 *1.2 Aims and scope*

196

197 In this contribution, we review the wide range of approaches and datasets available to practitioners 198 and students for geomorphological mapping in glacial environments. The main aims of this review are 199 to (i) synthesise scale-appropriate mapping approaches that are relevant to particular glacial settings, 200 (ii) devise frameworks that will help ensure best practice when mapping, and (iii) encourage clear 201 communication of details on mapping methods used in glacial geomorphological studies. This will 202 ensure transparency and aid data transferability against a background of growing demand to collate 203 geomorphological (and chronological) data in regional compilations (e.g. the BRITICE project: Clark 204 et al., 2004, 2018a; the DATED-1 database: Hughes et al., 2016). A further aim of this contribution is 205 to emphasise the continued and future importance of field mapping in geomorphological research, 206 despite the advent of very high-resolution remotely-sensed datasets in recent years (e.g. UAV-207 captured imagery).

208

209 The following two sections of this review focus on field mapping (Section 2) and remote mapping 210 (Section 3), respectively. We consider these methods in a broadly chronological order to provide 211 historical context and illustrate the evolution of geomorphological mapping in glacial environments. 212 Section 4 discusses the errors associated with each mapping method, an important issue that often 213 receives limited attention within geomorphological studies. Within this discussion, we highlight 214 approaches that can help manage and minimise residual errors. Subsequently, we review the mapping 215 methods used in particular glacial environments (Section 5) and synthesise frameworks to help ensure 216 best practice when mapping (Section 6).

217

For the purposes of this review, we distinguish two overarching 'work streams': (i) mapping of palaeo-ice sheet geomorphological imprints using a combined remote sensing approach, with some field checking (where feasible); and (ii) mapping of alpine- and plateau-style ice mass geomorphological imprints using a combination of remote sensing and considerable field mapping/checking. The second workstream incorporates a spatial continuum of glacier morphologies, namely cirque glaciers, valley glaciers, icefields and ice-caps (cf. Sugden and John, 1976; Benn and 224 Evans, 2010). The rationale for this subdivision is fourfold: Firstly, the approaches are governed by 225 the size of the (former) glacial systems and thus feasibility of using particular mapping methods in 226 certain settings (cf. Clark, 1997; Storrar et al., 2013). Secondly, there is a greater overlap of spatial 227 and temporal scales (i.e. more detailed records are preserved) in areas glaciated by smaller ice masses 228 that respond more rapidly to climate (cf. Lukas, 2005, 2012; Bradwell et al., 2013; Boston et al., 229 2015; Chandler et al., 2016b). Thirdly, the different mapping methodologies reflect the difficulties in 230 identifying vertical limits, thickness distribution and surface topography of palaeo-ice sheets (i.e. 231 emphasis often on mapping bed imprints) (cf. Stokes et al., 2015). Finally, the overarching methods 232 employed to map glacial landforms in alpine and plateau settings do not differ fundamentally with ice 233 mass morphology, i.e. most studies in these environments employ a combination of field mapping and 234 remote sensing. In Section 5.3, we also specifically consider geomorphological mapping in modern 235 glacial environments to highlight important issues relating to the temporal resolution of remotely-236 sensed data and landform preservation potential. We emphasise the importance of utilising multiple 237 datasets and/or mapping approaches in an iterative process in all glacial settings (multiple remotely-238 sensed datasets in the case of ice sheet-scale geomorphology) to increase accuracy and robustness, 239 akin to multi-proxy methodologies used in many Earth Science disciplines.

240

241 **2. Field mapping methods**

242

243 2.1 Background and applicability of field mapping

244

245 Traditionally, glacial geomorphological mapping has been undertaken through extensive field surveys, an approach that dates back to the late 19th Century and early 20th Century (e.g. Close, 1867; 246 Goodchild, 1875; Partsch, 1894; Sollas, 1896; Penck and Brückner, 1901/1909; Kendall, 1902; 247 248 Wright, 1912; Hollingworth, 1931; Caldenius, 1932). Field mapping involves traversing the study 249 area and recording pertinent landforms onto (enlarged) topographic base maps (Figure 1). Typically, 250 field mapping is conducted at cartographic scales of ~1: 10,000 (e.g. Leser and Stäblein, 1975; Rupke 251 and De Jong, 1983; Thorp, 1986; Ballantyne, 1989; Evans, 1990; Benn et al., 1992; Mitchell and 252 Riley, 2006; Rose and Smith, 2008; Boston, 2012a, b) or 1: 25,000 (e.g. Leser, 1983; Ballantyne, 2002, 2007a, b; Benn and Ballantyne, 2005; Lukas and Lukas, 2006). Occasionally, it is conducted at 253 254 even larger scales, such as 1: 1,000 to 1: 5,000, but this is most appropriate for small areas or project-255 specific purposes (e.g. Kienholz, 1977; Leser, 1983; Lukas et al., 2005; Coray, 2007; Graf, 2007; 256 Reinardy et al., 2013).

257

With improvements in technology, the widespread availability of remotely-sensed datasets, and a concomitant ease of access to high-quality printing facilities, alternative approaches to the traditional, *purely* field mapping method have also been employed, including (i) documenting sediment-landform 261 assemblages during extensive field campaigns both prior to and after commencing remote mapping 262 (e.g. Dyke et al., 1992; Krüger 1994; Lukas and Lukas, 2006; Kjær et al. 2008; Boston, 2012a, b; 263 Jónsson et al., 2014; Schomacker et al. 2014; Everest et al., 2017), (ii) mapping directly onto or 264 annotating print-outs of imagery (e.g. aerial photographs) in the field (e.g. Lovell, 2014), (iii) 265 recording the locations of individual landforms using a (handheld) Global Navigation Satellite System (GNSS) device (e.g. Bradwell et al., 2013; Brynjólfsson et al., 2014; Lovell, 2014; Małecki et al., 266 267 2018), or (iv) digitally mapping landforms in the field using a ruggedised tablet PC with built-in GNSS and GIS software (e.g. Finlayson et al., 2011; Pearce et al., 2014). These approaches to field 268 269 mapping are particularly useful where large-scale topographic maps are unavailable or obsolete.

270

271 Detailed field mapping is typically restricted to alpine- and plateau-style ice masses due to logistical 272 and financial constraints (Clark, 1997; Storrar et al., 2013). When conducted at the ice sheet scale, 273 field mapping is (or historically was) undertaken either as part of long-term campaigns by national 274 geological surveys in conjunction with surficial geology mapping programmes (e.g. Barrow et al., 275 1913; Flint et al., 1959; Krygowski, 1963; Campbell, 1967a, b; Hodgson et al., 1984; Klassen, 1993; 276 Priamonosov et al., 2000; Follestad and Bergstrøm, 2004) or necessarily highly-selective ground-277 truthing of remote mapping (e.g. Kleman et al., 1997, 2010; Golledge and Stoker, 2006; Stokes et al., 278 2013; Darvill et al., 2014; Stroeven et al., 2016; Pearce et al., 2018).

279

280 2.2 The field mapping process

281

282 Field mapping should ideally begin with systematic traverses of the study area – sometimes referred 283 to as a 'walk-over' (e.g. Demek, 1972; Otto and Smith, 2013) – to get a sense of the scale of the study 284 area and ensure that subtle features of importance, such as the location and orientation of ice-flow 285 directional indicators (e.g. flutes, striae, roches moutonnées and ice-moulded bedrock), are not 286 missed. In a palaeo-ice sheet context, mapping the location and orientation of striae in the field may 287 be of most interest as these can provide information on multiple (local) ice flow directions, of which 288 not all are recorded in the pattern of elongated bedforms mappable from remotely-sensed data (cf. 289 Kleman, 1990; Smith and Knight, 2011). Similarly, in a contemporary outlet glacier context, flutes are an important indicator of ice flow direction - sometimes of annual ice flow trajectories of glacier 290 291 margins (cf. Chandler et al., 2016a; Evans et al., 2017) – but due to their subtlety they may only be identifiable in the field (e.g. Jónsson et al. 2014). 292

293

Traversing should ideally start from higher ground, where an overview can be gained, and proceed by crossing a valley axis (or a cirque floor, for example) many times to enable the viewing and assessment of landforms from as many perspectives, angles and directions as possible (cf. Demek, 1972). In addition to systematic traverses, landform assemblages in, for example, individual 298 valleys/basins should ideally be viewed from a high vantage point in low light (e.g. Benn, 1990). 299 Depending on the location and orientation of landforms, it may be beneficial to see the same area 300 either (i) early in the morning or late in the afternoon/evening due to longer shadows, or (ii) both in 301 the morning and afternoon/evening due to the changing position of longer shadows. These procedures 302 ensure that apparent dimensions and orientations, which are influenced by perception under different 303 viewing angles and daylight conditions, can be taken into account in descriptions and interpretations. 304 This approach circumvents potential complications relating to subtle features that may only be visible 305 from one direction or certain angles.

306

307 The location of features should be recorded on field maps or imagery (e.g. aerial photograph) extracts 308 with reference to 'landmarks' that are clearly identifiable both in the field and on the base 309 maps/imagery, such as distinct changes in contour-line inflection, river bends, confluences, prominent 310 bedrock exposures and large ridges or mounds (Lukas and Lukas, 2006; Boston, 2012a, b). Where 311 geomorphological features are small, background relief is low and/or conspicuous reference points are 312 absent, a network of mapped reference points can be established by either taking a series of cross-313 bearings on prominent features using a compass (e.g. Benn, 1990) or by verifying locations using a 314 handheld GNSS (e.g. Lukas and Lukas, 2006; Boston, 2012a, b; Brynjólfsson et al., 2014; Jónsson et 315 al. 2014; Lovell, 2014; Pearce et al., 2014; van der Bilt et al., 2016). The latter is useful for recording 316 the location of point-data such as striae, erratic or glacially-transported boulders, and sediment 317 exposures (cf. Lukas and Lukas, 2006; Boston, 2012a, b; Pearce et al., 2014). Additional information 318 between known reference points can then be interpolated and marked on the geomorphological map.

319

Establishing the size of landforms and features and plotting them on the map as accurately as possible is of crucial importance, and in addition to the inflections of contours (which may mark the location and boundaries of prominent ridges, for example), the mapper may pace out and/or estimate lengths, heights and widths. For larger landforms, or those masked by forest, walking around the perimeter of landforms and establishing a GNSS-marked 'waypoint-trail' is a good first approximation.

325

The strategy outlined above offers a broad perspective on the overall landform pattern and ensures accurate representation of landforms on field maps. To ensure accurate genetic interpretation of individual landforms, and the landscape as a whole, this field mapping strategy should ideally form part of an iterative process of observation and interpretation whilst still in the field (see Section 2.3, below).

331

332 2.3 Interpreting glacial landforms

333

334 In the preceding section, we focused on the technical aspects of field mapping and the means of 335 recording glacial landforms. However, geomorphological mapping typically forms the foundation of 336 process-oriented and palaeoglaciological reconstruction studies (see Section 1.1) and should, 337 therefore, be embedded within a process of observation and interpretation. Definitive interpretation of 338 glacial landforms, and glacial landscapes as a whole, can rarely be made on the basis of surface 339 morphology alone. Additional strands of field evidence may become highly relevant, if not essential, 340 depending on the objectives of the individual project: sedimentological data are crucial to interpreting 341 processes of landform formation and glacier dynamics (e.g. Lukas, 2005; Benn and Lukas, 2006; 342 Benediktsson et al., 2010, 2016; Chandler et al., 2016a), whilst chronological data are fundamental to 343 robust palaeoglaciological reconstructions and related palaeoclimatic studies (e.g. Finlayson et al., 344 2011; Gribenski et al., 2016; Hughes et al., 2016; Stroeven et al., 2016; Bendle et al., 2017b; Darvill 345 et al., 2017). Moreover, time and resources are limited and pragmatism necessary. Thus, observations 346 must be targeted efficiently and effectively, in line with the research aims.

347

348 Much field-based research adopts an inductive approach, in which observations are collected and used 349 to argue towards a particular conclusion. This is a valid approach at the exploratory stage of research, 350 but deeper understanding of a landscape requires a more iterative process, in which data collection is 351 conducted within a framework of hypothesis generation and testing. For this reason, it is useful to 352 adopt a number of alternative working hypotheses (Chamberlin, 1897) that can be tested and gradually eliminated, following the principle of falsification (Popper, 1972). This process is best 353 354 conducted in the field while it is possible to make key observations to test an interpretation, especially 355 if the field site is remote and expensive to re-visit.

356

357 Following initial data collection, preliminary interpretations can be used to predict the outcome of 358 new observations, which can then be used to test and refine the interpretation. Well-framed 359 hypotheses allow an investigator to anticipate other characteristics of a glacial landscape, then to test 360 those predictions by targeted investigation of key localities (see Benn, 2006). For example, the 361 presence of a certain group of landforms (e.g. moraines trending downslope into a side valley) can be 362 used to formulate hypotheses (e.g. blockage of the side-valley by glacier ice), which in turn can be 363 used to predict the presence of other sediment-landform associations in a particular locality (e.g. 364 lacustrine sediments or shoreline terraces in the side-valley). Further detailed geomorphological mapping (and sedimentological analyses) in that area would then allow testing and falsification of the 365 366 alternative working hypotheses. Iterations of this process during field mapping enable an increasingly 367 detailed and robust understanding of the glacier system to be constructed. This coupled inductive-368 deductive approach is much more powerful than a purely inductive process: narratives that 'explain' a 369 set of observations can appear very persuasive, even self-evident, but there may be other narratives 370 that are also consistent with the same observations (cf. Popper, 1972).

371

372 Process-form models are useful tools in this inductive-deductive approach to landscape interpretation. 373 In particular, landsystem or facies models make explicit links between landscape components and 374 genetic processes, providing structure and context for data collection and interpretation (e.g. Eyles, 375 1983; Brodzikowski and van Loon, 1991; Evans, 2003a; Benn and Evans, 2010). At best, process-376 form models are not rigid templates or preconceived categories into which observations are forced, 377 but a flexible set of possibilities that can guide, shape and enrich investigations (Benn and Lukas, 378 2006). For example, preliminary remote mapping may reveal features that suggest former glacier 379 lobes may have surged (e.g. Lovell et al., 2012). Systematic study of sediment-landform assemblages, 380 sediment exposures and other evidence, with reference to modern analogues (e.g. Evans and Rea, 381 2003), allows this idea to be rigorously evaluated in a holistic context (e.g. Darvill et al., 2017). This 382 then opens up new avenues for research in a creative and open-ended process.

383

This inductive-deductive approach to interpreting glacial landscapes and events should be embedded as part of the geomorphological mapping process (see Section 6). When dealing with palaeo-ice sheets, such field-based investigations may be guided by (existing) remote mapping. In alpine and plateau-style ice mass settings, sedimentological and chronological investigations should ideally form an integral part of field surveys.

389

390 3. Remote mapping methods

391

392 In the following sections, we review the principal remote mapping approaches employed in glacial 393 geomorphological research, with analogue (or hard-copy) remote mapping (Section 3.1) and digital 394 remote mapping (Section 3.2) considered separately. We give an overview of a number of datasets 395 used for digital remote (i.e. GIS-based) mapping, namely satellite imagery (see Section 3.2.2.1), aerial 396 photographs (see Section 3.2.2.2), digital elevation models (see Section 3.2.2.3), freeware virtual 397 globes (see Section 3.2.2.4) and UAV-captured imagery (see Section 3.2.2.5). Each individual section 398 provides a brief outline of the historical background and development of the methods, and we discuss 399 the individual approaches in a broadly chronological order. Section 3.3 then provides an overview of 400 image processing techniques, highlighting that pragmatic solutions are often required.

401

We focus principally on remotely-sensed datasets relevant to terrestrial (onshore) glacial settings in the following sections, since submarine (bathymetric) datasets and mapping of submarine glacial landforms have been subject to recent reviews elsewhere (see Dowdeswell et al., 2016; Batchelor et al., 2017). Nevertheless, we do acknowledge that the emergence of geophysical techniques to investigate submarine (offshore) glacial geomorphology is a major development over the last two decades. Similarly, the emergence of geophysical datasets of sub-ice geomorphology in the last decade or so has been revolutionary, particularly in relation to subglacial bedforms (see Stokes, 2018).
Many of the issues we discuss in relation to mapping from DEMs are transferable to those
environments.

411

412 *3.1 Analogue remote mapping*

413

414 *3.1.1 Background and applicability of analogue remote mapping*

415 Geomorphological mapping from analogue (hard-copy) aerial photographs became a mainstream approach in glacial geomorphology in the 1960s and 1970s, with early proponents including, for 416 417 example, the Geological Survey of Canada (e.g. Craig, 1961, 1964; Prest et al., 1968) and UK-based researchers examining the Quaternary geomorphology of upland Britain (e.g. Price, 1961, 1963; 418 419 Sissons, 1967, 1977a, b, 1979a, b; Sugden, 1970) and contemporary glacial landsystems (e.g. Petrie 420 and Price, 1966; Price, 1966; Welch, 1966, 1967, 1968; Howarth, 1968; Howarth and Welch, 1969a, 421 b). The latter research on landsystems in Alaska and Iceland was particularly pioneering in that it 422 exploited a combination of aerial photograph interpretation, surveying techniques and early photogrammetry (see Evans, 2009, for further details). 423

424

425 Despite continued development of remote sensing technologies and the availability of digital aerial 426 photographs (see Section 3.2.2.2), analogue stereoscopic aerial photographs are still used for glacial 427 geomorphological mapping (e.g. Hättestrand, 1998; Benn and Ballantyne, 2005; Lukas et al., 2005; 428 Boston, 2012a, b; Evans and Orton, 2015). Additionally, the availability of high-quality 429 photogrammetric scanners means that archival, hard-copy aerial photographs can be scanned at high 430 resolutions, processed using digital photogrammetric methods and subsequently used for on-screen 431 digitisation (Section 3.2; e.g. Bennett et al., 2010; Jónsson et al., 2014). As with field mapping, the 432 interpretation of analogue aerial photographs is primarily used for mapping alpine- and plateau-style ice mass geomorphological imprints. Historically, analogue aerial photograph interpretation was 433 434 extensively used for mapping palaeo-ice sheet geomorphological imprints, particularly by the 435 Geological Survey of Canada, who combined aerial photograph interpretation with detailed ground 436 checking and helicopter-based surveys (e.g. Craig, 1961, 1964; Hodgson et al., 1984; Aylsworth and Shilts, 1989; Dyke et al., 1992; Klassen, 1993; Dyke and Hooper, 2001). This approach has largely 437 438 been superseded by satellite imagery and DEM interpretation in palaeo-ice sheet settings (see Section 5.1) but is applied in palaeo-ice sheet contexts for more detailed mapping of selected/complex areas 439 (e.g. Dyke, 1990; Kleman et al., 2010; Stokes et al., 2013; Storrar et al., 2013; Darvill et al., 2014; 440 441 Evans et al., 2014).

442

444 For glacial geomorphological mapping purposes, vertical panchromatic aerial photographs have 445 traditionally been employed, with pairs of photographs (stereopairs) viewed in stereo using a 446 stereoscope (with magnification) (e.g. Melander, 1975; Horsfield, 1983; Krüger, 1994; Kleman et al., 447 1997; Hättestrand, 1998; Evans and Twigg, 2002; Benn and Ballantyne, 2005; Lukas and Lukas, 448 2006; Boston, 2012a, b; Chandler and Lukas, 2017). During aerial surveys, longitudinally-449 overlapping photographs along the flight path (endlap $\geq 60\%$) are captured in a series of laterally-450 overlapping parallel strips (sidelap \geq 30%), with the two different viewing angles of the same area 451 resulting in the stereoscopic effect (due to the principle of parallax; see Lillesand et al., 2015, for further details). This form of aerial photograph interpretation has been demonstrated to be a 452 453 particularly valuable tool for determining the exact location, shape and planform of small features in 454 glaciated terrain (e.g. Ballantyne, 1989, 2002, 2007a, b; Bickerton and Matthews, 1992, 1993; Lukas 455 and Lukas, 2006; Boston, 2012a, b), provided the photographs are of appropriate scale, quality and 456 tonal contrast (cf. Benn, 1990; Benn et al., 1992).

457

458 Mapping from hard-copy aerial photographs is undertaken by drawing onto acetate sheets 459 (transparency films) whilst viewing the aerial photographs through a stereoscope, with the acetate 460 overlain on one photograph of a stereopair (Figure 2). Ideally, mapping should be conducted using a 461 super-fine pigment liner with a nib size of 0.05 mm to enable small features to be mapped. Even so, it 462 may still be necessary to compromise on the level of detail mapped; for example, meltwater channels between ice-marginal moraines have been left off maps in some studies due to map scale, with the 463 464 associated text describing chains of moraines interspersed with meltwater channels (e.g. Benn and 465 Ballantyne, 2005; Lukas, 2005).

466

467 Examining stereopairs from multiple sorties ('flight missions') in parallel or in combination with 468 digital aerial photographs may be beneficial and help alleviate issues such as localised cloud cover, 469 snow cover, poor tonal contrast, afforestation and anthropogenic developments (e.g. Horsfield, 1983; 470 Bennett, 1991; McDougall, 2001). Additionally, it is advantageous to examine stereopairs multiple 471 times – preferably before and after field mapping – to increase feature identification and improve the 472 accuracy of genetic interpretations (Lukas and Lukas, 2006; Sahlin and Glasser, 2008). When 473 conducting mapping over a large area with multiple stereopairs, examining stereopairs from a sortie 474 'out of sequence' (i.e. not mapping from consecutive pairs of photographs) may provide a means of 475 internal corroboration and ensure objectivity and robustness (Bennett, 1991).

476

In order to reduce geometric distortion, which increases towards the edges of aerial photographs due to the central perspective (Lillesand et al., 2015), it is advisable to keep the areas mapped onto the acetate as close as possible to the centre of one aerial photograph of a stereopair (Kronberg, 1984; Lukas, 2002, 2005a; Evans and Orton, 2015). These hand-drawn overlays can subsequently be scanned at high resolutions and then georeferenced and digitised using GIS and graphics software (see
also Section 3.3.1; e.g. Lukas and Lukas, 2006; Boston, 2012a, b).

483

484 *3.2 Digital remote mapping*

485

486 *3.2.1 Background and applicability of digital remote mapping*

487 The development of GIS software packages (e.g. commercial: ArcGIS; open source: OGIS) and the proliferation of digital imagery, particularly freely available satellite imagery, have undoubtedly been 488 489 the most significant developments in glacial geomorphological mapping. GIS packages have provided 490 platforms and tools for visualising, maintaining, manipulating and analysing vast quantities of 491 remotely-sensed and geomorphological data (cf. Gustavsson et al., 2006, 2008; Napieralski et al., 492 2007b). Their use in combination with digital imagery allows geomorphological features to be 493 mapped directly in GIS software (Figure 3), with individual vector layers created for each 494 geomorphological feature. Moreover, the availability of digital imagery enables the mapper to alter 495 the viewing scale instantaneously and switch between various datasets/types, allowing for a flexible 496 but systematic approach.

497

498 Digital mapping (on-screen digitisation) also provides georeferenced geomorphological data, which 499 has two important benefits: Firstly, these data can easily be used to derive landform metrics (e.g. 500 Hättestrand et al., 2004; Clark et al., 2009; Spagnolo et al., 2010, 2014; Storrar et al., 2014; Ojala et 501 al., 2015; Dowling et al., 2016; Ely et al., 2016a, 2017a); and, secondly, these data can be seamlessly 502 incorporated into wider, regional-scale GIS compilations (e.g. Bickerdike et al., 2016; Stroeven et al., 503 2016; Clark et al., 2018a). Additionally, digital remote mapping allows the user to record attribute 504 data (e.g. data source) tied to individual map (vector) layers, which can be useful for large 505 compilations of previously published mapping (e.g. Bickerdike et al., 2016; Clark et al., 2018a). Such 506 compendia help to circumvent issues relating to the often-fragmented nature of geomorphological 507 evidence (i.e. numerous spatially separate studies) and identify gaps in the mapping record. Once 508 assembled across large areas, they also enable evidence-based reconstructions of entire ice sheets and 509 regional ice sheet sectors (see Clark et al., 2004, 2018a). Indeed, the ongoing open access data 510 revolution in academia and the increasing publication/availability of mapping output (in the form of 511 GIS files; e.g. Finlayson et al., 2011; Darvill et al., 2014; Bickerdike et al., 2016; Bendle et al., 512 2017a), means that geomorphological mapping can have wider impact beyond individual local to 513 regional studies.

514

515 3.2.2 Datasets for digital remote mapping

516 There is now a plethora of remotely-sensed datasets covering a wide range of horizontal resolutions 517 $(10^{-2} \text{ to } 10^2 \text{ m})$, enabling the application of digital mapping (in some form) to all glacial settings. We 518 provide an overview of the principal datasets used in digital mapping below, with mapping 519 approaches in specific glacial settings reviewed in Section 5.

520

521 3.2.2.1 Satellite imagery. The development of satellite-based remote sensing in the 1970s and 522 subsequent advances in technology have revolutionised understanding of glaciated terrain, particularly 523 with respect to palaeo-ice sheet geomorphology and dynamics (see Section 5.1; Clark, 1997; Stokes, 524 2002; Stokes et al., 2015). The potential of satellite imagery was first demonstrated by the pioneering 525 work of Sugden (1978), Andrews and Miller (1979) and Punkari (1980), with the availability of large-526 area view (185 km x 185 km) Landsat Multi-Spectral Scanner (MSS) images affording a new 527 perspective of glaciated regions. These allowed a single analyst to systematically map ice sheet-scale 528 (1:45,000 to 1: 1,000,000) glacial geomorphology (e.g. Boulton and Clark, 1990a, b) in a way that 529 previously would have required the painstaking mosaicking of thousands of aerial photographs (e.g. 530 Prest et al., 1968). Since the 1980s, there has been an explosion in the use of satellite imagery for 531 glacial geomorphological mapping and there is now a profusion of datasets available (Table 1). 532 Importantly, many of these sensors capture multispectral data, which can enhance landform detection through image processing and the use of different band combinations (see Section 3.3.2). The uptake 533 534 of satellite imagery has coincided with improvements in the availability and spatial and spectral 535 resolution of satellite datasets globally, with Landsat (multispectral: 30 m; panchromatic: 15 m), 536 ASTER (15 m), Sentinel-2 (10 m) and SPOT (up to 1.5 m) images proving the most popular. More 537 recently, satellite sensor advancements have enabled the capture of satellite images with resolutions 538 comparable to aerial photographs (Figure 4; e.g. SPOT6/7, QuickBird, WorldView). Thus, these 539 datasets are also suitable for mapping typically smaller and/or complex glacial landforms produced by 540 cirque glaciers, valley glaciers and icefield/ice-cap outlets (e.g. Chandler et al., 2016a; Evans et al., 541 2016b; Ewertowski et al., 2016; Gribenski et al., 2016; Małecki et al., 2018).

542

543 In general, as better-resolution imagery has become more widely available at low to no cost, older, 544 coarser-resolution datasets (e.g. Landsat MSS: 60 m) have largely become obsolete. Nevertheless, 545 Landsat data (TM, ETM+ and OLI: 15 to 30 m) are still the standard data source for ice-sheet scale 546 mapping, with the uptake of high-resolution commercial satellite imagery still relatively slow in such 547 studies. This is primarily driven by the cost of purchasing high-resolution commercial datasets, 548 making freely-available imagery such as Landsat a valuable resource. In addition, archival satellite data afford time-series of multi-spectral images that may facilitate assessments of geomorphological 549 550 changes through time; for example, fluctuations in highly dynamic (surging or rapidly retreating) 551 glacial systems (e.g. Flink et al., 2015; Jamieson et al., 2015). Conversely, for smaller research areas 552 (e.g. for a single valley or foreland), high-resolution satellite imagery is becoming an increasingly 553 viable option, with prices for georeferenced and orthorectified products comparable to those for 554 digital aerial photographs (see Section 3.2.2.2). This also has the benefit of saving time on

photogrammetric processing, with many vendors providing consumers with various processing options. Consequently, on-demand, high-resolution (commercial) satellite imagery will inevitably come into widespread usage, where costs are not prohibitive. Alternatively, freeware virtual globes and web mapping services (e.g. *Bing Maps, Google Earth*) offer valuable resources for free visualisation of such high-resolution imagery (see Section 3.2.2.4).

560

561 3.2.2.2 Digital aerial photographs. With improvements in technology, high-resolution (ground 562 resolution <0.5 m per pixel) digital copies of aerial photographs have become widely available and used for glacial geomorphological mapping (e.g. Brown et al., 2011a; Bradwell et al., 2013; 563 Brynjólfsson et al., 2014; Jónsson et al., 2014; Pearce et al., 2014; Schomacker et al., 2014; Chandler 564 et al., 2016a; Evans et al., 2016c; Lardeux et al., 2016; Lønne, 2016; Allaart et al., 2018). Indeed, 565 566 digital aerial photographs, along with scanned copies of archival aerial photographs, are now more 567 widely used than hard-copy stereoscopic aerial photographs, particularly in modern glacial settings. 568 Additionally, the introduction of UAV technology in recent years has allowed sub-decimetre 569 resolution aerial photographs to be captured on demand (see Section 3.2.2.5). Aside from their 570 typically superior resolution/scale compared with hard-copy aerial photographs, a key advantage of 571 aerial photographs in digital format is the ability to produce orthorectified aerial photograph mosaics 572 (or 'orthophotographs') and DEMs with low root mean square errors (RMSEs <1 m; see Section 4.4), 573 when combined with ground control points collected using surveying equipment (e.g. Kjær et al. 574 2008; Bennett et al., 2010; Schomacker et al., 2014; Chandler et al., 2016b; Evans et al., 2017). These 575 photogrammetric products can then be used for on-screen digitisation and generation of georeferenced 576 geomorphological mapping (Figure 5), as outlined above.

577

Typically, digital aerial photographs are captured by commercial surveying companies (e.g. 578 579 Loftmyndir ehf, Iceland; Getmapping, UK), meaning that they may be expensive to purchase and 580 costs may be prohibitive for large study areas. This is in contrast to hard-copy (archival) aerial 581 photographs that are often freely available for viewing in national collections. Additionally, digital aerial photographs are not readily viewable in stereo with a standard desktop setup, although on-582 583 screen mapping in stereoscopic view is possible on workstations equipped with stereo display and 584 software such as BAE Systems SOCET SET (e.g. Kjær et al., 2008; Benediktsson et al., 2009). 585 However, this approach is not applicable to orthophotographs. An alternative approach is to visualise orthophotographs in 3D by draping them over a DEM (see Section 3.2.2.3) in GIS software such as 586 ESRI ArcScene or similar (Figure 6; e.g. Benediktsson et al., 2010; Jónsson et al., 2014; Schomacker 587 588 et al., 2014; van der Bilt et al., 2016). Three-dimensional assessment in ArcScene, parallel to mapping 589 in ArcMap, may aid in landform detection, delineation and interpretation.

590

591 3.2.2.3 Digital Elevation Models. Over the last ~15 years there has been increasing use of DEMs in 592 glacial geomorphology, particularly for mapping at the ice-sheet scale (e.g. Glasser and Jansson, 593 2008; Hughes et al., 2010; Ó Cofaigh et al., 2010; Evans et al., 2014, 2016d; Principato et al., 2016; 594 Ojala, 2016; Stokes et al., 2016a; Mäkinen et al., 2017; Norris et al., 2017). DEMs are raster-based 595 models of topography that record absolute elevation, with each pixel or grid cell in a DEM 596 representing the average height for the area it covers (Clark, 1997; Smith et al., 2006). Terrestrial 597 DEMs can be generated by a variety of means, including from surveyed contour data, directly from 598 stereo imagery (aerial photographs, satellite and UAV-captured imagery), or from air- and space-599 borne radar and LiDAR systems (Smith and Clark, 2005). An important recent development in this regard has been the Surface Extraction with TIN-based Search-space Minimization (SETSM) 600 601 algorithm for automated extraction of DEMs from stereo satellite imagery (Noh and Howat, 2015), 602 which has been used to generate the ArcticDEM dataset (https://www.pgc.umn.edu/data/arcticdem/). 603 However, SETSM DEMs may contain systematic vertical errors that require correction (e.g. Carrivick 604 et al., 2017; Storrar et al., 2017).

605

The majority of DEMs with national- to international-scale coverage (Table 2) typically have a 606 607 coarser spatial resolution than aerial photographs and satellite imagery and represent surface 608 elevations rather than surface reflectance. As a result, it may be difficult to identify glacial landforms 609 produced by relatively small ice masses (cirque glaciers, valley glaciers and icefield outlets), 610 precluding detailed mapping of their planforms (cf. Smith et al., 2006; Hughes et al., 2010; Brown et 611 al., 2011a; Boston, 2012a, b; Pearce et al., 2014). Conversely, these DEMs can be particularly 612 valuable for mapping glacial erosional features (e.g. glacial valleys, meltwater channels), as well as 613 major glacial depositional landforms produced by larger ice masses (e.g. Greenwood and Clark, 2008; 614 Heyman et al., 2008; Livingstone et al., 2008; Hughes et al., 2010; Morén et al., 2011; Barr and Clark, 615 2012; Stroeven et al., 2013; Turner et al., 2014a; Margold et al., 2015a; Blomdin et al., 2016a, b; Lindholm and Heyman, 2016; Mäkinen et al., 2017; Storrar and Livingstone, 2017). However, the 616 617 recent development of UAV (see section 3.2.2.5) and LiDAR technologies have allowed the 618 generation of very high resolution DEMs (<0.1 m), enabling the application of DEMs to map small 619 glacial landforms (e.g. Evans et al., 2016a; Ewertowski et al., 2016; Ely et al., 2017). We anticipate 620 national-scale LiDAR DEMs becoming widely-used in the future, with a number of nations recently 621 releasing or currently capturing/processing high horizontal resolution (≤ 2 m) LiDAR data (Table 2; 622 e.g. Dowling et al. 2013; Johnson et al. 2015).

623

Although the principal focus of this contribution is terrestrial/onshore glacial geomorphological mapping, it is worth highlighting here that the availability of spatially-extensive bathymetric charts, such as the General Bathymetric Chart of the Oceans (GEBCO) and International Bathymetric Chart of the Arctic Ocean (IBCAO: Jakobsson et al., 2012), and high-resolution, regional (often industry628 acquired) bathymetric data has been an important development in submarine/offshore glacial 629 geomorphological mapping. This has enabled the gridding of DEMs to map submarine glacial 630 geomorphological imprints (see Dowdeswell et al., 2016), markedly enhancing understanding of 631 palaeo-ice sheets in marine sectors (e.g. Ottesen et al., 2005, 2008a, 2016; Bradwell et al., 2008; 632 Winsborrow et al., 2010, 2012; Livingstone et al., 2012; Ó Cofaigh et al., 2013; Hodgson et al., 2014; 633 Stokes et al., 2014; Margold et al., 2015a, b; Greenwood et al., 2017) and modern tidewater (often 634 surging) glaciers (e.g. Ottesen and Dowdeswell, 2006; Ottesen et al., 2008b, 2017; Robinson and Dowdeswell, 2011; Dowdeswell and Vazquez, 2013; Flink et al., 2015; Streuff et al., 2015; Allaart et 635 636 al., 2018). In addition, recent years have seen the production of DEMs of sub-ice topography from 637 geophysical datasets (radar and seismics) at spatial resolutions suitable for identifying and mapping bedforms (see King et al., 2007, 2009, 2016a; Smith et al., 2007; Smith and Murray, 2009). This work 638 639 has advanced understanding of the evolution of bedforms beneath Antarctic ice streams, providing 640 important genetic links between the formation of landforms beneath modern ice sheets and those left-641 behind by palaeo-ice masses (Stokes, 2018). The interested reader is directed to recent reviews 642 (Livingstone et al., 2012; Ó Cofaigh, 2012; Stokes et al., 2015; Dowdeswell et al., 2016; Batchelor and Dowdeswell, 2017; Stokes, 2018) for further discussion on the importance of geophysical 643 644 evidence for understanding ice sheet extent and dynamics.

645

646 3.2.2.4 Freeware virtual globes. The advent of freeware virtual globes (e.g. Google Earth, NASA 647 Worldwind) and web mapping services (e.g. Bing Maps, Google Maps) have provided platforms for 648 free visualisation of imagery from various sources and low-cost mapping resources. A key benefit of 649 virtual globes is the ability to visualise imagery and terrain in 3D and from multiple viewing angles, 650 which may aid landform detection when used in conjunction with other datasets and software (e.g. 651 Heyman et al., 2008; Bendle et al., 2017a). Moreover, a number of virtual globes and web mapping 652 services have the ability to link with other freeware and open-source programmes; for example, free plugins are available to import Google Earth and Bing Maps imagery into the open-source GIS 653 software package QGIS. Thus, a mapper can combine freely available, often high-resolution (e.g. 654 SPOT6/7, QuickBird, WorldView), imagery and the capabilities of GIS technology without the 655 656 expense associated with commercial imagery and software (see Sections 3.2.2.1 and 3.2.2.2).

657

The most widely-used virtual globe is *Google Earth*, with a 'professional' version (*Google Earth Pro*) freely available since 2015 (see Mather et al., 2015, for a review). An increasing number of glacial geomorphological studies are noting the use of *Google Earth* (but not necessarily the imagery type) as a mapping tool (see Table 1), principally to cross-check mapping conducted from other imagery. However, some studies have also utilised the built-in digitising tools for mapping (e.g. Margold and Jansson, 2011; Margold et al., 2011; Fu et al., 2012). There is a compromise on the functionality of freeware virtual globes and digitisation tools are often not as flexible and/or user-friendly, but these 665 can be counteracted by the option to import imagery into GIS software. In the case of *Google Earth*, it 666 is also possible to export Keyhole Markup Language (KML) files that can be used for subsequent 667 analyses and map production in GIS software (following file conversion). Open access remotely-668 sensed datasets are also available through commercial GIS software, with high resolution satellite 669 imagery (e.g. GeoEye-1, SPOT-5, WorldView) available for mapping through the in-built 'World 670 Imagery' service in *ESRI ArcGIS* (e.g. Bendle et al., 2017a).

671

672 Despite the benefits, some caution is necessary when using freeware virtual globes as there may be 673 substantial errors in georeferencing of imagery, which users cannot account/correct for. Moreover, 674 dating of imagery is not necessarily clear or accurate (Mather et al., 2015; Wyshnytzky, 2017). The latter may not be a concern if mapping in a palaeoglaciological setting, whilst any georeferencing 675 676 errors may not be as significant if mapping broad patterns at the ice sheet scale. Conversely, errors 677 associated with freeware mapping may be significant when comparing imagery from different times 678 and/or when mapping in highly dynamic, contemporary glacial environments. Aside from these 679 potential issues, limitations are imposed by pre-processing of imagery, with no option to, for example, modify band combinations to enhance landform detection (see Section 3.3.2). 680

681

682 3.2.2.5 UAV-captured imagery. The recent emergence of UAV technology provides an alternative 683 method for the acquisition of very high-resolution (<0.1 m per pixel) geospatial data that circumvents 684 some of the issues associated with more established approaches, particularly in relation to temporal 685 resolution and the high-cost of acquiring commercial remotely-sensed data (see also Smith et al., 686 2016a). This method provides a rapid, flexible and relatively inexpensive (following the initial 687 acquisition of the UAV and associated software) means of acquiring up-to-date imagery at an 688 unprecedented spatial resolution and is becoming increasingly employed in glacial research (Figure 7; 689 Rippin et al., 2015; Ryan et al., 2015; Chandler et al., 2016b; Ewertowski et al., 2016; Tonkin et al., 690 2016; Westoby et al., 2016; Ely et al., 2017; Allaart et al., 2018). UAV-captured images are processed 691 using Structure-from-Motion (SfM) photogrammetry techniques, with Agisoft Photoscan being the 692 most common software in use at present (e.g. Chandler et al., 2016b; Evans et al., 2016a; Ely et al., 693 2017; Allaart et al., 2018). This methodology has enabled the production of sub-decimetre resolution 694 orthophotographs and DEMs with centimetre-scale error values (RMSEs <0.1 m; see Section 4.4) for 695 glacial geomorphological mapping (e.g. Evans et al., 2016a; Ely et al., 2017). Although surveying of ground control points is still preferable for processing UAV-captured imagery, a direct georeferencing 696 697 workflow (see Turner et al., 2014b, for further details) is capable of producing reliable geospatial 698 datasets from imagery captured using consumer-grade UAVs and cameras, without the need for 699 expensive survey equipment (see Carbonneau and Dietrich, 2017).

700

701 The use of UAVs will be valuable in future glacial geomorphological research due to their flexibility 702 and low-cost. In particular, UAVs open up the exciting possibility of undertaking repeat surveys at 703 high temporal (sub-annual to annual) resolutions in modern glacial settings (Immerzeel et al., 2014; 704 Chandler et al., 2016b; Ely et al., 2017). Multi-temporal UAV imagery will enable innovative 705 geomorphological studies on issues such as (i) the modification and preservation potential of 706 landforms over short timescales (Ely et al., 2017), (ii) the frequency of ice-marginal landform 707 formation, particularly debates on sub-annual to annual landform formation (Chandler et al., 2016b), 708 and (iii) changes in process-form regimes at contemporary ice-margins (Evans et al., 2016a).

709

710 Using UAVs to capture aerial imagery is not without challenges, particularly in relation to the 711 challenge of intersecting suitable weather conditions in modern glacial environments: many UAVs are 712 unable to fly in high windspeeds, whilst rain can infiltrate electrical components and create hazy 713 imagery (Ely et al., 2017). Flight times and areal coverage are also limited by battery life, with some 714 battery packs permitting as little as 10 minutes per flight. There are also legal considerations, with the 715 use of UAVs prohibited in some localities/countries or requiring licenses/permits. Moreover, there 716 may be restrictions on flying heights and UAVs may need to be flown in visual line of sight, further 717 limiting areal coverage. Nevertheless, we envisage UAV technology becoming more widespread and 718 a key tool in high-resolution glacial geomorphological investigations, especially if future 719 technological developments can increase the range of conditions in which UAVs can be flown. In 720 future, it is likely that UAV technology will be primarily used for investigating short-term changes 721 across relatively small areas.

722

723 *3.3 Image processing for mapping*

724

725 An important part of geomorphological mapping is processing remotely-sensed datasets in preparation 726 for mapping, but this is often given limited prominence in glacial geomorphological studies. 727 Crucially, processing of remotely-sensed data aids the identification of glacial landforms and ensures accurate transfer of geomorphological data from the imagery. In the sections below, we provide a 728 729 brief overview of image processing solutions for aerial photographs (Section 3.3.1), satellite imagery 730 (Section 3.3.2) and DEMs (Section 3.3.3). Reference is made to common processing techniques used 731 to remove distortion and displacement evident in aerially-captured imagery (see Campbell and Wynne 732 (2011) and Lillesand et al. (2015) for further details), but these are not discussed in detail for reasons 733 of brevity and clarity. However, a detailed workflow diagram outlining the potential procedures for a 734 range of scenarios (depending on data, resources and time) is available as Supplementary Material. 735 We emphasise that compromises and pragmatic solutions are necessary, particularly in the case of 736 aerial photographs, as the 'idealised' scenario is frequently not an option due to data limitations or 737 logistical constraints.

738

739 *3.3.1 Aerial photograph processing*

740 Aerial photographs contain varying degrees of distortion and displacement owing to their central (or 741 perspective) projection. Geometric distortion is related to radial lens distortion and refraction of light 742 rays in the atmosphere. Additional displacement occurs as a result of the deviation of the camera from 743 a vertical position (caused by roll, pitch and yaw of the aircraft), relief and curvature of the Earth. 744 Non-corrected aerial photographs are therefore characterised by relief displacement and scale 745 variations, which increase towards the edges of the photograph (see Campbell and Wynne (2011) and 746 Lillesand et al. (2015) for further details). Thus, it is necessary to apply geometric corrections to aerial 747 photographs before geomorphological mapping.

748

749 Ideally, aerial photographs would be corrected using stereoscopic (or conventional) photogrammetric 750 processing in software packages such as *Imagine Photogrammetry* (formerly *Leica Photogrammetry*) 751 Suite, or LPS). This approach involves the extraction of quantitative elevation data from stereoscopic 752 (overlapping) imagery to generate DEMs and orthorectified imagery (see also Section 3.2). Internal 753 and external parameters, along with the location of GCPs, are used to establish the relationship 754 between the position of the images and a ground coordinate system (e.g. Kjær et al., 2008; Bennett et 755 al., 2010). However, this approach may be impractical and unsuitable in many glacial settings; for 756 example, it is unrealistic to collect GCPs using (heavy) survey equipment (e.g. RTK-GPS) in former 757 plateau icefield and ice-cap settings due to their location (remote, upland environments) and the size 758 of the study area (and thus quantity of aerial photographs and GCPs required). Moreover, camera 759 calibration data (focal length, fiducial marks, principal point coordinates and lens distortion) are 760 frequently unavailable or incomplete for archive datasets, and the process is also not applicable to 761 acetate overlays. Thus, orthorectification of imagery – correction of geometric distortions in both the 762 horizontal (x and y coordinates) and vertical (z coordinates) dimensions - is typically precluded over 763 larger areas, although it may be possible to employ this approach for individual circue basins, valleys 764 and glacier forelands (e.g. Wilson, 2005; Bennett et al., 2010; Chandler et al., 2016a). Consequently, pragmatic solutions are required for georectification of imagery, i.e. the process of transforming and 765 766 projecting imagery to a (local) planar coordinate system. Several approaches have been used to 767 overcome this and we briefly outline these below in relation to analogue aerial photographs (Section 768 3.3.1.1) and digital aerial photographs (Section 3.3.1.2).

769

3.3.1.1 Analogue aerial photograph processing. A potential pragmatic solution to correcting analogue
 (hard-copy) aerial photographs is to georeference scanned copies of acetate overlays or the original
 aerial photographs to reference points on other forms of (coarser) georeferenced digital imagery (if
 available; e.g. DEMs, orthorectified radar images, satellite images). The scanned images can then be
 georectified and resampled using the georeferencing functions within GIS and remote sensing

programmes such as *ArcGIS* or *Erdas Imagine* (cf. Boston, 2012a, for further details). This approach is particularly useful where hard-copy aerial photographs are used in combination with (coarser) digital imagery. Using this procedure, georeferenced acetate overlays of Quaternary features in the Scottish Highlands have been produced with RMSE values ranging between 2.71 m and 7.82 m (Boston, 2012a), comparable to archival aerial photographs that have been processed using conventional photogrammetric techniques (e.g. Bennett et al., 2010).

781

782 The above georectification method works best if relatively small areas are mapped on one acetate. 783 This is because radial distortion increases towards the edges of aerial photographs and, as such, 784 presents a significant problem to matching reference points when large areas have been mapped. From 785 our experience, we estimate the maximum effective area that can be corrected without the danger of 786 mismatches at ~6 km². However, this figure depends on the terrain conditions and would have to be 787 smaller in high mountain areas where relief distortion is increased due to greater differences between 788 valleys and adjacent peaks (Lillesand et al., 2015). The mapped area could, conversely, be somewhat 789 larger in low-relief terrain as objects are roughly equally far away from the camera lens over larger 790 areas and thus subject to less distortion (Kronberg, 1984; Lillesand et al., 2015). The aforementioned 791 constraints might seem to make georectification from hard-copy aerial photographs a laborious 792 process, but this is counterbalanced by being able to record small landforms in great detail due to the 793 high-resolution 3D visualisation allowed by stereopairs.

794

795 3.3.1.2 Digital aerial photograph processing. Digital aerial photographs can be georeferenced within 796 GIS and remote sensing software following a similar process to that outlined in Section 3.3.1.1, i.e. 797 digital aerial photographs can be georeferenced to other forms of (coarser) georeferenced imagery. 798 Alternatively, SfM photogrammetry can be used to produce orthophotographs and DEMs from digital 799 aerial photographs, which partly circumvents issues relating to incomplete or absent camera 800 calibration data (e.g. Chandler et al., 2016a; Evans et al., 2016e, 2017; Tonkin et al., 2016; Midgley 801 and Tonkin, 2017; Mertes et al., 2017). SfM photogrammetry functions under the same basic 802 principles as stereoscopic photogrammetry but there are fundamental differences: the geometry of the 803 'scene', camera positions and orientation are solved automatically in an arbitrary 'image-space' 804 coordinate system without the need to specify either the 3D location of the camera or a network of 805 GCPs with known 'object-space' coordinates (cf. Westoby et al., 2012; Carrivick et al., 2016; Smith 806 et al., 2016a, for further details). However, positional data (GCPs) are still required to process the 807 digital photographs for geomorphological mapping, i.e. to assign the SfM models to an 'object-space' 808 coordinate system. Ideally, this should be conducted through ground control surveys (see above), but 809 a potential pragmatic solution is to utilise coordinate data from freeware virtual globes such as Bing 810 Maps (see also Supplementary Material). Position information ('object-space' coordinates) is

introduced after model production, with the benefit that errors in GCPs will not propagate in theDEM.

- 813
- 814 *3.3.2 Satellite imagery processing*

815 Satellite imagery products are typically available in georectified form as standard and therefore do not 816 require geometric correction prior to geomorphological mapping. With respect to high-resolution, 817 commercial satellite imagery (e.g. WorldView-4 captured imagery; 0.31 m GSD), these products are 818 often available for purchase as either georeferenced and orthorectified products (with consumers able 819 to define the processing technique used) at comparable prices to commercial aerial photographs, 820 thereby removing the need for photogrammetric processing. Alternatively, it is possible to purchase 821 less expensive 'ortho-ready' imagery and perform orthorectification (where DEM or GCP data are 822 available), thus providing greater end-user control on image processing (e.g. Chandler et al., 2016a; 823 Ewertowski et al., 2016).

824

825 Although satellite imagery does not typically require geometric correction for mapping, it is important 826 to consider the choice of band combinations when using multispectral satellite imagery (e.g. Landsat, 827 ASTER; Table 1). Since the detection of glacial landforms from optical satellite imagery relies on the 828 interaction of reflected radiation with topography, different combinations of spectral bands can be 829 employed to optimise landform identification (see Jansson and Glasser, 2005). Manipulating the order 830 of bands with different spectral wavelengths allows the generation of various visualisations, or false-831 colour composites, of the terrain. For example, specific band combinations may be particularly useful 832 for detecting moraine ridges (7, 5, 2 and 5, 4, 2), mega-scale glacial lineations (4, 5, 6) and meltwater 833 channels (4, 3, 2) from Landsat TM and ETM+ imagery (Jansson and Glasser, 2005; Lovell et al., 834 2011; Morén et al., 2011). This is principally due to the change in surface vegetation characteristics 835 (e.g. type, density and degree of development) between different landforms, between landforms and the surrounding terrain. For example, former meltwater channels typically appear as overly-wide 836 837 corridors (relative to any modern drainage) of lush green vegetation and stand out clearly as bright red 838 when using a near-infrared false-colour composites (bands 4, 3, 2: Landsat TM and ETM+), since the 839 green colour of surface vegetation is strongly reflected in near-infrared bands (band 4: Landsat TM 840 and ETM+). In addition to the manipulation of band combinations during the mapping process, it can 841 also be beneficial to use satellite image derivatives based on ratios of band combinations, such as 842 vegetation indices (see Walker et al., 1995) and semi-automated image classification techniques (e.g. 843 Smith et al., 2000, 2016b).

844

Aside from manipulating spectral band combinations, it may also be beneficial to use the higherresolution panchromatic band as a semi-transparent layer alongside the multispectral bands to aid landform detection (e.g. Morén et al., 2011; Stroeven et al., 2013; Lindholm and Heyman, 2016), or to 848 merge the pixel resolutions of the panchromatic and multispectral bands through pan-sharpening 849 techniques (e.g. Glasser and Jansson, 2008; Greenwood and Clark, 2008; Storrar et al., 2014; 850 Chandler et al., 2016a; Ewertowski et al., 2016). Pan-sharpening can be particularly valuable when it 851 is desirable to have both multispectral capabilities (e.g. different band combinations to differentiate 852 between features with varying surface characteristics) and higher-spatial resolutions to help determine 853 the extent and morphology of individual landforms.

854

855 3.3.3 Digital Elevation Model processing

856 Various processing techniques are available that can be beneficial when identifying and mapping 857 glacial landforms from DEMs (Bolch and Loibl, 2017). For mapping and analytical purposes, DEM data are typically converted into 'hillshaded relief models' (Figure 8), whereby different solar 858 859 illumination angles and azimuths are simulated within GIS software to produce the shaded DEMs. 860 This rendition provides a visually realistic representation of the land surface, with shadows improving 861 detection of surface features. Consequently, hillshaded relief models should be generated using a 862 variety of illumination azimuths (direction of light source) and angles (elevation of light source) to alleviate the issue of 'azimuth bias', the notion that some linear landforms are less visible when 863 864 shaded from certain azimuths (see Lidmar-Bergström et al., 1991; Smith and Clark, 2005). An illumination angle of 30° and azimuths set at orthogonal positions of 45° and 315° have been 865 866 suggested as optimal settings for visualisation (Smith and Clark, 2005; Hughes et al., 2010). Vertical 867 exaggeration of these products (e.g. three to four times) can also aid landform identification (e.g. 868 Hughes et al., 2010). Semi-transparent DEMs can be draped over shaded-relief images to accentuate topographic contrasts (Figure 9), or a semi-transparent satellite image can be draped over a DEM to 869 870 achieve both a multispectral and topographic assessment of a landscape (e.g. Jansson and Glasser, 871 2005). First- and second-order DEM-derivatives, including surface gradient (slope) and curvature, 872 have also been found to be useful for mapping (e.g. Smith and Clark, 2005; Evans, 2012; Storrar and 873 Livingstone, 2017).

874

875 **4. Assessment of mapping errors and uncertainties**

876

877 In this section, we provide an overview of the main errors and uncertainties associated with the 878 various geomorphological mapping methods introduced in the preceding sections. Consideration and 879 management of mapping errors should be an important part of glacial geomorphological mapping 880 studies, since any errors/uncertainties incorporated in the geomorphological map will propagate into 881 subsequent palaeoglaciological and palaeoclimatic reconstructions. This is of most relevance to small 882 ice masses (cirque glaciers, valley glaciers, outlet glaciers), e.g. metre-scale geolocation errors would 883 have significant implications for studies aiming to establish ice-margin retreat rates at the order of 884 tens of metres (e.g. Krüger, 1995; Lukas and Benn, 2006; Lukas, 2012; Bradwell et al., 2013;

Chandler et al., 2016b). Conversely, mapping errors might be negligible in the context of continentalscale ice sheet reconstructions (e.g. Hughes et al., 2014; Stroeven et al., 2016; Margold et al., 2018).

887

888 The overall 'quality' of any geomorphological map is a function of three interlinked factors: mapping 889 resolution, accuracy and precision. It is important to highlight that, irrespective of the mapping 890 method employed (field or remote-based), the accuracy and precision of the mapping reflects two 891 related factors: (i) the skill, philosophy and experience of the mapper; and (ii) the detectability of the 892 landforms (Smith and Wise, 2007; Otto and Smith, 2013; Hillier et al., 2015). Mapper philosophy 893 concerns issues such as how landforms are mapped (e.g. generalised mapping vs. mapping the 894 intricate details of individual landforms) and interpreted (e.g. differences in terminology and landform 895 classification), which will partly vary with background and training. The significance of the skill, 896 philosophy and experience in mapping is exemplified by the stark differences across boundaries of 897 British Geological Survey (BGS) map sheets that have been mapped by different surveyors, for 898 example (cf. Clark et al., 2004).

899

900 A key determinant of landform detectability is resolution, generally defined as the finest element that 901 can be distinguished during survey/observation (Lam and Quattrochi, 1992). In geomorphological 902 mapping it may be, for example, the smallest distinguishable landform that is visible from remotely-903 sensed data or that can be drawn on a field map. The accuracy of geomorphological mapping relates 904 to positional accuracy (i.e. difference between 'true' and mapped location of the landform), geometric 905 accuracy (i.e. difference between 'true' and mapped shape of the landform), and attribute accuracy 906 (i.e. deviation between 'true' and mapped landform types) (Smith et al., 2006). For spatial data, it is 907 usually not possible to obtain absolute 'true' data, due to limitations such as the 'resolution' of 908 remotely-sensed data and the accuracy of instruments/surveying equipment. Precision is often used to 909 express the reproducibility of surveys, which is controlled by random errors. These are errors that are 910 innate in the survey/observation process which cannot be removed (Butler et al., 1998). We now 911 outline the specific uncertainties associated with field mapping (Section 4.1), analogue remote 912 mapping (Section 4.2) and digital remote mapping (Section 4.3).

913

914 *4.1 Field mapping errors and uncertainty*

915

The correct positioning, orientation and scale of individual geomorphological features on field maps is dependent on the skill of the mapper and the ability to correctly interpret and record landforms. If a handheld GNSS device is used to locate landforms in the field, the positional accuracy is usually restricted to several metres and related to three factors: (i) the quality of the device (e.g. antenna, number of channels, ability to use more than one GNSS); (ii) the position of satellites; and (iii) the characteristics of the surrounding landscapes and space weather (solar activity can affect signal 922 quality). Higher accuracy (cm- or even mm-scale) can only be achieved when supplemented by 923 measurements using additional surveying (e.g. differential Global Positioning Systems (dGPS), real 924 time kinematic (RTK-) GPS or total station). Alongside positioning errors, the horizontal resolution 925 (and, consequently, accuracy) of the field map is related to line thickness on the field map (Knight et 926 al., 2011; Boston, 2012a, b; Otto and Smith, 2013). A pencil line has a thickness of between 0.20 and 927 0.50 mm on a field map; therefore, individual lines represent a thickness of between 2 and 5 m on 1: 928 10,000 scale maps, rendering the maps accurate to this level at best (Raisz, 1962; Robinson et al., 929 1995; Boston, 2012a). This necessitates some element of selection during field mapping of relatively 930 small landforms formed by alpine- and plateau-style ice masses, as not all the information that can be 931 seen in the field can be mapped, even at a large scale such as 1: 10,000. In terms of the vertical 932 accuracy of field maps, it should be recognised that the mapping is only as accurate as the resolution 933 of the source elevation data: if the topographic base map has contours at 10 m intervals, the mapping 934 has a vertical resolution, and thus accuracy, of 10 m at best, irrespective of the (perceived) skill of the 935 cartographer. As with positional accuracy, higher vertical accuracy necessitates the use of geodetic-936 grade surveying equipment.

937

938 4.2 Analogue remote mapping errors and uncertainty

939

940 Accurate detection and mapping of individual landforms from analogue (hard-copy) aerial 941 photographs is influenced by factors such as the scale or resolution of the photographs, shadow length 942 (shadows may obscure the 'true' planform or landforms altogether), the presence/absence of 943 vegetation, cloud cover and tonal contrast (photographs may appear 'flat', thus limiting landform 944 detection). The resolution of analogue remotely-sensed datasets is associated with scale, which results 945 from the altitude of the plane and camera lens focal length, and the optical resolution of the lens and 946 sensor (Wolf et al., 2013). It should also be recognised that the accuracy of the (non-rectified) 947 mapping, as with field mapping, is limited by the thickness of the pen used for drawing on the acetate 948 sheets. Super-fine pens typically have a nib size of 0.05–0.20 mm; thus, lines on an acetate overlay 949 typically represent thicknesses between ~ 1.25 m and 5.00 m at a common aerial photograph average 950 scale of 1: 25,000. Despite being particularly useful for detailed mapping of small features and 951 complex landform patterns, the level of accuracy achievable using this methodology is therefore 952 ~ 1.25 m at best. However, further errors will be introduced to the geomorphological mapping once 953 the raw, non-rectified acetates are georectified (see Section 3.3.1).

954

955 *4.3 Digital remote mapping errors and uncertainty*

956

A key influence on landform detectability from digital remotely-sensed data is the scale of the featurerelative to the resolution of the digital dataset, with a particular challenge being the mapping of

959 features with a scale close to or smaller than the resolution of the imagery. Conversely, mapping 960 exceptionally large ('mega-scale') glacial landforms can be challenging, depending on the remotely-961 sensed dataset employed (e.g. Greenwood and Kleman, 2010). Unlike analogue mapping (both in the 962 field and remotely), the thickness of digital lines is not typically a problem for digital mapping, so 963 landform detection and recording are fundamentally linked to spatial resolution. Spatial resolution of digital remotely-sensed data refers to the capability to distinguish between two objects, typically 964 965 expressed as either (i) pixel size or grid cell size or (ii) ground sampled distance. Pixel/grid size refers to the projected ground dimension of the smallest element of the digital image (Figure 10), whilst 966 967 ground sampled distance (GSD) refers to the ground distance between two measurements made by the 968 detector (the value of measurement is subsequently assigned to a pixel) (Figure 10; Duveiller and 969 Defourny, 2010). In practice, the spatial resolution of digital imagery is lower than the pixel size 970 (Figure 10).

971

972 Landform detectability from raster images (i.e. remotely-sensed data) can be considered with 973 reference to the Nyquist-Shannon sampling theorem, since they comprise discrete sampled values. 974 According to this theorem, the *intrinsic resolution* is twice the sampling distance of the measured 975 values, whereas the nominal resolution is twice the pixel/grid size (cf. Pipaud et al., 2015, and 976 references therein). The effective resolution and, consequently, the minimum landform 977 footprint/planform that can be unambiguously sampled are defined by the smaller of these two values 978 (cf. Pike, 1988). Where the Nyquist–Shannon criterion is not satisfied for either the intrinsic or 979 nominal resolution, landforms with footprints below the critical value may be visible but are rendered 980 ambiguously in digital imagery, i.e. their boundaries are not clearly definable and mappable (cf. 981 Cumming and Wong, 2005). Further factors that influence landform identification from digital 982 imagery include the strength of the landform signal relative to background terrain, and the azimuth 983 bias introduced by differences in the orientation of linear features and the illumination angle of the 984 sun (Smith and Wise, 2007), along with localised issues such as cloud cover, snow cover and 985 vegetation. The timing of data collection is also a key factor, particularly in the case of modern glacial 986 environments (see Section 5.3).

987

Aside from the factors outlined above, (raw) remotely-sensed data will contain distortion and/or 988 989 geometric artefacts of varying degrees. Distortions inherent in raw aerial photographs can be partially 990 or almost fully removed during georeferencing of acetate sheets or photogrammetric processing of 991 aerial photographs (see Section 3.3.1). Raw satellite imagery will contain biases related to attitude, 992 ephemeris and drift errors as well as displacements related to the relief, which, similarly to aerial 993 photographs, is more visible in mountainous areas than in lowland settings (Grodecki and Dial, 2003; 994 Shean et al., 2016). With respect to DEMs, some datasets captured using air- and space-borne radar 995 approaches may contain a number of artefacts (Clark, 1997; Figure 11), with geometric artefacts

996 particularly significant in upland settings. Geometric artefacts, such as foreshortening and layover, are 997 corrected during image processing by stretching high terrain into the correct position, which can result 998 in a smoothed region on steep slopes (Figure 12). In other parts of upland terrain, information will be 999 lost on the leeside of slopes, away from the sensor, where high ground prevents the radar beam from 1000 reaching the lower ground beneath it (Figure 11). Such issues can be alleviated, at least partly, by 1001 examining multiple complementary remotely-sensed datasets and mapping at a variety of scales.

1002

1003 4.4. Assessment and management of uncertainties

1004

1005 Due to the subjective nature of geomorphological mapping, assessing mapping precision is not an 1006 easy task. One possible approach is to compare results of mapping using different datasets/methods 1007 with a perceived more 'truthful' dataset (i.e. field-based survey) (Smith et al., 2006). The number, size 1008 and shape of mapped landforms in comparison with a 'true' dataset can be used as an approximation 1009 of mapping reliability. Precision and accuracy of the produced geomorphological map can also be 1010 estimated based on the quality of the source data. Most of the datasets are delivered with at least some 1011 assessment of uncertainties, often expressed as accuracy; for example, the SRTM DEM has a 1012 horizontal accuracy of ± 20 m and a vertical accuracy of ± 16 m (Rabus et al., 2003). Alternatively, 1013 some remotely-sensed datasets have an associated total root mean square error (RMSE), which 1014 indicates displacement between 'true' control points and corresponding points on the remotely-sensed 1015 data (Wolf et al., 2013). However, both are measures of the overall ('global') quality of the dataset. 1016 Thus, these errors may be deceptive because such 'global' measures ignore spatial patterns of errors 1017 and local terrain characteristics (cf. Lane et al., 2005; James et al., 2017). For example, DEM errors 1018 will typically be more pronounced on steep slopes, where even a small horizontal shift will incur large 1019 differences in elevation.

1020

1021 Ideally, remotely-sensed datasets should be evaluated independently by the mapper to establish their 1022 geolocation accuracy (accuracy of x, y and z coordinates). If feasible, surveys of ground control points 1023 (GCPs) should be conducted using geodetic-grade surveying equipment (e.g. RTK-GPS, total station). 1024 A sub-sample of this GCP dataset can be used for photogrammetric processing, and RMSE values 1025 calculated. Subsequently, the remaining GCPs (i.e. those not used for photogrammetric processing) 1026 can be used to perform a further quality check, by quantifying deviations from the coordinates of the GCPs and the corresponding points on the generated DEM (e.g. Carrivick et al., 2017). An additional 1027 1028 approach, in geomorphologically stable areas, is to compare the location of individual data points 1029 from the DEM (or raw point cloud) being used for mapping with those on a reference DEM (or raw 1030 point cloud) (e.g. King et al., 2016b; Carrivick et al., 2017; James et al., 2017; Midgley and Tonkin, 1031 2017; Mertes et al., 2017). Parameters such as the mean deviation, standard deviation and relative 1032 standard deviation between the two datasets can then be calculated to perform a quantitative

assessment of quality and accuracy of the DEM (e.g. King et al., 2016b; Mertes et al., 2017).
Performing these assessments may then facilitate correction of the processed datasets (e.g. Nuth and
Kääb, 2011; Carrivick et al., 2017; King et al., 2017).

1036

1037 To some extent, residual uncertainties relating to the skill, philosophy and experience of the mapper 1038 may be reduced by developing a set of clear criteria for identifying and mapping particular landforms 1039 (e.g. Barrell et al., 2011; Darvill et al., 2014; Bendle et al., 2017a; Lovell and Boston, 2017). That 1040 said, there are currently no 'agreed' genetic classification schemes for interpreting glacial sediment-1041 landform assemblages, despite the development of facies and landsystem models for particular glacial 1042 environments (e.g. Eyles, 1983; Brodzikowski and van Loon, 1991; Evans, 2003a; Benn and Evans, 2010). Indeed, terminologies are inconsistently used in glacial geomorphological research, as different 1043 1044 'schools' or traditions still exist. Thus, it is probably most appropriate to select a scheme that has been 1045 in frequent use in a given area (to enable ready comparison) or to develop one suited for a particular 1046 area or problem. Notwithstanding potential discrepancies relating to genetic classification or 1047 terminology, this will at least ensure transparency in future use and analysis of the geomorphological 1048 mapping.

1049

1050 Given the influence of the individual mapper on accuracy and precision, it may be beneficial and 1051 desirable for multiple mappers to complete (initially) independent field surveys and examination of 1052 remotely-sensed datasets to enhance reliability and reproducibility (cf. Hillier et al., 2015; Ewertowski 1053 et al., 2017). However, this approach would only be applicable in collaborative efforts and may be 1054 impractical due to various factors (e.g. study area size, data access restrictions). The level of detection 1055 of individual landforms might be improved by employing multiple methods to enhance landform 1056 detectability, whilst the genetic interpretation of landforms (landform classification) can be tested by 1057 detailed sedimentological investigations (see Section 2.3). Some uncertainties associated with the 1058 quality of data source (e.g. shadows, artefacts) can be alleviated, at least partly, by examining multiple 1059 complementary remotely-sensed datasets and mapping at a variety of scales.

1060

1061 **5. Scale-appropriate mapping approaches**

1062

The following sections place the main geomorphological mapping methods (see Sections 2 and 3) in the spatial and temporal context of the glacial settings in which they are commonly used, demonstrating that particular methods are employed depending on factors such as the size of the study area, former glacial system and landforms (Table 3). We focus on three broad glacial settings for the purposes of this discussion: palaeo-ice sheets (Section 5.1), alpine- and plateau-style ice masses (Section 5.2) and modern glacier forelands (Section 5.3). Although geomorphological mapping in modern glacial settings follows the same general procedures as in former alpine and plateau-style ice mass settings (see Section 6.2), specific consideration of contemporary glacier forelands is warranted
 due to important issues relating to the temporal resolution of remotely-sensed data and landform
 preservation potential, which are not as significant in palaeoglaciological settings.

1073

1074 5.1 Palaeo-ice sheet settings

1075

1076 The continental-scale of palaeo-ice sheets typically necessitates a mapping approach that enables 1077 systematic mapping of a large area in a time and cost-effective manner while still allowing accurate 1078 identification of landform assemblages at a variety of scales. However, the nature of the approach will 1079 differ depending on the aim of the investigation, as this fundamentally determines what needs to be 1080 mapped and how it should be mapped. Palaeo-ice sheet reconstructions have been produced at a range 1081 of scales, from entire ice sheets (e.g. Dyke and Prest, 1987a, b, c; Kleman et al., 1997, 2010; Boulton 1082 et al., 2001; Glasser et al., 2008; Clark et al., 2012; Livingstone et al., 2015) to regional/local sectors 1083 (e.g. Hättestrand, 1998; Jansson et al., 2003; Stokes and Clark, 2003; Ó Cofaigh et al., 2010; 1084 Astakhov et al., 2016; Darvill et al., 2017). Depending on the aim of the study, some investigations may focus specifically on mapping particular landforms. For example, studies of ice-sheet flow 1085 1086 patterns frequently focus on mapping subglacial bedforms, such as drumlins (e.g. Boulton and Clark, 1087 1990a, b; Kleman et al., 1997, 2010; Stokes and Clark, 2003; Hughes et al., 2010). Nonetheless, 1088 cartographic reduction is often still required to manage the volume of information, resulting in the grouping of similarly-orientated bedforms into flow-sets (occasionally termed fans or swarms) (e.g. 1089 1090 Jansson et al., 2002, 2003; De Angelis and Kleman, 2007; Greenwood and Clark, 2009a, b; Stokes et 1091 al., 2009; Hughes et al., 2014; Atkinson et al., 2016).

1092

1093 In many cases, studies incorporate all or most of the landforms across ice sheet scales to derive 1094 palaeoglaciological reconstructions (e.g. Kleman et al., 1997, 2010; Stroeven et al., 2016). The 1095 rationale for this is that glaciation styles and processes (e.g. ice-marginal, subglacial) can be inferred 1096 from particular combinations of landforms in landform assemblages (e.g. Clayton et al., 1985; Stokes 1097 and Clark, 1999; Evans, 2003b; Kleman et al., 2006; Evans et al., 2008, 2014; Darvill et al., 2017; 1098 Norris et al., 2018). Establishing relationships between landforms is therefore valuable, not only in 1099 understanding glaciation styles, but also in helping decipher the relative sequence of formation (e.g. 1100 Clark, 1993; Kleman and Borgström, 1996) that may lay the foundations for absolute dating. 1101 Typically, ice sheet investigations are focused on the spatial and temporal evolution of these various 1102 aspects, requiring the robust integration of geomorphological mapping with absolute dating 1103 techniques (see Stokes et al., 2015). For example, following pioneering palaeoglaciological studies of 1104 the Fennoscandian ice sheet (e.g. Kleman, 1990, 1992; Kleman and Stroeven, 1997; Kleman et al., 1105 1997), cosmogenic nuclide exposure dating offered a means to quantify dates and rates (e.g. Fabel et 1106 al., 2002, 2006; Stroeven et al., 2002a, b, 2006; Harbor et al., 2006). Such data are crucial to tune and

validate numerical models used to reconstruct evolving ice sheet limits, flow configurations and
subglacial processes (e.g. Boulton and Clark, 1990a, b; Näslund et al., 2003; Evans et al., 2009b;
Hubbard et al., 2009; Stokes and Tarasov, 2010; Kirchner et al., 2011; Livingstone et al., 2015; Stokes
et al., 2016b; Patton et al., 2017a, b).

1111

1112 5.1.1 Manual mapping of palaeo-ice sheet geomorphological imprints

1113 Satellite imagery and DEMs are the prevailing remotely-sensed datasets used for mapping ice sheet-1114 scale landforms, and these datasets have been at the forefront of key developments in the 1115 understanding of palaeo-ice sheets (cf. Stokes, 2002; Stokes et al., 2015). Notably, the use of satellite 1116 imagery resulted in the identification of hitherto-unrecognised mega-scale glacial lineations (MSGLs; 1117 Boulton and Clark, 1990a, b; Clark, 1993), which are now recognised as diagnostic geomorphological 1118 evidence of ice streams within palaeo-ice sheets (see Stokes and Clark, 1999, 2001, and references 1119 therein). This has allowed tangible links to be made between the behaviours of former Quaternary ice sheets and present-day ice sheets (e.g. King et al., 2009; Stokes and Tarasov, 2010; Stokes et al., 1120 1121 2016b). Aerial photograph interpretation and field mapping are also used in some studies (e.g. 1122 Hättestrand and Clark, 2006; Kleman et al., 2010; Darvill et al., 2014), but satellite imagery and 1123 DEMs are in wider usage for practical reasons (see also Section 3.2). In recent years, the development 1124 of LiDAR datasets has led to their increasing application for high resolution mapping of landforms 1125 formed by palaeo-ice sheets, particularly in Scandinavia (e.g. Dowling et al., 2015; Greenwood et al., 1126 2015; Ojala et al., 2015; Ojala, 2016; Mäkinen et al., 2017; Peterson et al., 2017). We expect this to be 1127 a major area of growth in future mapping studies of former ice sheets.

1128

1129 Mapping glacial landforms from remotely-sensed data typically involves manual on-screen 1130 digitisation using one of two main approaches: (i) creating polylines along the crestline or thalweg of 1131 landforms or (ii) digitising polygons that delineate the breaks of slope around landform margins (i.e. 1132 digitising the planform). The approach employed will depend on the requirements of the study; for 1133 example, flow-parallel bedforms (e.g. drumlins and MSGLs) have variously been mapped as polylines 1134 (e.g. Kleman et al., 1997, 2010; Stokes and Clark, 2003; De Angelis and Kleman, 2007; Storrar and 1135 Stokes, 2007; Livingstone et al., 2008; Brown et al., 2011b) and polygons (e.g. Hättestrand and 1136 Stroeven, 2002; Hättestrand et al., 2004; Hughes et al., 2010; Spagnolo et al., 2010, 2014; Stokes et 1137 al., 2013; Ely et al., 2016a; Bendle et al., 2017a) (Figure 13). The rationale behind mapping flow-1138 parallel bedforms as linear features is that dominant orientations of a population provide sufficient 1139 information when investigating ice sheet-scale flow patterns and organisation, although image 1140 resolution may also be a determining factor. Mapping polygons allows the extraction of individual 1141 landform metrics (e.g. elongation ratios) that can provide insights into subglacial processes (e.g. Ely 1142 et al., 2016a) and regional variations in ice sheet flow dynamics (e.g. Stokes and Clark, 2002, 2003; 1143 Hättestrand et al., 2004; Spagnolo et al., 2014), but it is far more time-consuming than digitising linear features. Increasingly, it is being recognised that the population metrics and spectral
characteristics of the subglacial bedform 'field' as a whole are most important for quantifying
bedforms and deciphering subglacial processes and conditions (see Hillier et al., 2013, 2016;
Spagnolo et al., 2017; Clark et al., 2018b; Ely et al., 2018; Stokes, 2018).

1148

1149 5.1.2 Automated mapping of palaeo-ice sheet geomorphological imprints

1150

1151 Comprehensive mapping of palaeo-ice sheet geomorphological imprints, and particularly of 1152 bedforms, typically entails the identification and mapping of large numbers (in some cases >10,000) 1153 of the same, or very, similar types of features (e.g. Clark et al., 2009; Kleman et al., 2010; Storrar et 1154 al., 2013). The manual digitisation of such large numbers of landforms is a time-consuming process. 1155 Consequently, semi-automated and automated mapping techniques are increasingly being applied to 1156 glacial geomorphology (e.g. Napieralski et al., 2007b; Saha et al., 2011; Maclachlan and Eyles, 2013; Eisank et al., 2014; Robb et al., 2015; Yu et al., 2015; Jorge and Brennand, 2017a, b), particularly 1157 1158 given that features of a single landform type (e.g. drumlins or MSGLs) will have fairly uniform 1159 characteristics (orientation, dimensions and morphology). Automated and semi-automated mapping 1160 techniques typically use either a pixel- or object-based approach (see Robb et al., 2015, and references 1161 therein). Thus far, automated and semi-automated approaches have primarily focused on mapping 1162 drumlins or MSGLs from medium- to high-resolution DEMs. Several methods have been used, 1163 including multi-resolution segmentation (MRS) algorithms (Eisank et al., 2014), a Curvature Based 1164 Relief Separation (CBRS) technique (Yu et al., 2015), Object Based Image Analysis (OBIA) (Saha et 1165 al., 2011; Robb et al., 2015), and clustering algorithms (Smith et al., 2016b).

1166

1167 Most recently, 2D discrete Fourier transformations have been applied to automatically quantify 1168 MSGLs (see Spagnolo et al., 2017). In contrast to traditional mapping approaches, this new method 1169 analyses the whole topography simultaneously to identify the wavelength and amplitude of periodic 1170 features (i.e. waves or ripples across the topography) without the need to manually digitise them. This 1171 automated approach is in its infancy but is likely to provide quantitative data that are useful for (i) 1172 testing and parameterising models of subglacial processes and landforms (e.g. Barchyn et al., 2016; 1173 Stokes, 2018) and (ii) facilitating comparison between subglacial bedforms and other bedforms (e.g. 1174 Fourrière et al., 2010; Kocurek et al., 2010; Murray et al., 2014).

1175

1176 5.2 Alpine and plateau glacial settings

1177

1178 Mapping the geomorphological imprints of former alpine- and plateau-style ice masses (cirque 1179 glaciers, valley glaciers, icefields and ice-caps) is particularly significant, since the geomorphological 1180 imprints of such discrete ice masses facilitate reconstructions of their three-dimensional form (extent, 1181 morphology and thickness). By contrast, establishing the vertical limits, thickness distribution and 1182 surface topography of palaeo-ice sheets is challenging (cf. Stokes et al., 2015). Importantly, three-1183 dimensional glacier reconstructions permit the calculation of palaeoclimatic boundary conditions for 1184 glaciated regions (e.g. Kerschner et al., 2000; Bakke et al., 2005; Stansell et al., 2007; Mills et al., 1185 2012; Boston et al., 2015), data that cannot be obtained from point-source palaeoenvironmental 1186 records in distal settings (e.g. lacustrine archives). Empirical palaeoclimatic data derived from glacier 1187 reconstructions are important for three reasons. Firstly, these data facilitate analyses of wind patterns 1188 across loci of former glaciers and, in a wider context, regional precipitation gradients and atmospheric 1189 circulation patterns (e.g. Ballantyne, 2007a, b). Secondly, the data allow glaciodynamic conditions 1190 reconstructed from sediment-landform assemblages (e.g. moraines) to be directly linked to climatic 1191 regimes, thereby providing insights into glacier-climate interactions at long-term timescales (e.g. 1192 Benn and Lukas, 2006; Lukas, 2007a). Finally, independent, empirical information on climatic 1193 boundary conditions is fundamental to parameterising and testing numerical models used to simulate 1194 past glacier-climate interactions (e.g. Golledge et al., 2008). Thus, the geomorphological records of 1195 alpine and plateau-style ice masses are powerful proxies for understanding the interactions of such ice 1196 masses with climate.

1197

1198 Alpine- and plateau-style ice masses encompass a broad spatial spectrum of glacier morphologies (cf. 1199 Sugden and John, 1976; Benn and Evans, 2010), but geomorphological mapping of glacial landforms 1200 in alpine and plateau settings generally follows a similar approach that combines remote sensing and 1201 considerable field mapping/checking (Figure 14; e.g. Federici et al., 2003, 2017; Bakke et al., 2005; 1202 Lukas and Lukas, 2006; Reuther et al., 2007; Hyatt, 2010; Bendle and Glasser, 2012; Pearce et al., 1203 2014; Borsellino et al., 2017). Hence, alpine- and plateau-style ice masses are considered collectively 1204 here. The similarities in mapping approaches across a wider range of spatial scales partly reflect the 1205 fact that, in both alpine and plateau settings, the majority of (preserved) glacial landforms are 1206 confined to spatially- and/or topographically-restricted areas (e.g. glaciated valleys), i.e. glacial 1207 landforms relating to plateau-style ice masses (i.e. plateau icefields, ice-caps) are dominantly formed 1208 by outlet glaciers. Conversely, an important component of mapping in upland environments is often 1209 assessing any glacial geomorphological evidence for connections between supposed valley glaciers 1210 and plateau surfaces/rounded summits, i.e. alpine vs. plateau styles of glaciation (e.g. McDougall, 1211 2001; Boston et al., 2015). The recognition of any plateau-based ice has significant implications for 1212 studies aiming to assess glacier dynamics and regional palaeoclimate (see Rea et al., 1999; Boston, 1213 2012a, and references therein). Consequently, it is important to utilise a versatile geomorphological 1214 mapping approach in alpine and plateau settings that allows mapping of glacial landforms at a wide range of spatial scales and potentially across very large areas (>500 km²), whilst also providing 1215 1216 sufficiently high resolution imagery to map planforms of individual, small landforms (e.g. moraines).

1218 5.2.1 Remote mapping of alpine and plateau settings

1219

1220 Glacial geomorphological mapping from remotely-sensed data in alpine and plateau ice mass settings 1221 typically involves interpretation of either analogue or digital aerial photographs (see Sections 3.1 and 1222 3.2.2.2; e.g. Bickerton and Matthews, 1993; Boston, 2012a; Finlayson et al., 2011; Lukas, 2012; 1223 Izagirre et al., 2018). This reflects the superior resolution required to map in detail the frequently 1224 smaller glacial landforms produced by alpine and plateau-style ice masses, by contrast to the coarser 1225 resolution satellite imagery and DEMs predominantly used in ice sheet settings (see Section 5.1). The 1226 use of analogue (hard-copy) and digital aerial photographs varies in alpine and plateau settings, 1227 depending on data availability and the preference of individual mappers. For example, hard-copy, 1228 panchromatic aerial photographs have been widely used in conjunction with stereoscopes (see Section 1229 3.1) for mapping Younger Dryas glacial landforms in Scotland, owing to their excellent tonal contrast 1230 (e.g. Benn and Ballantyne, 2005; Lukas and Lukas, 2006; Boston, 2012a, b). Indeed, depending on the 1231 environment and quality/resolution of available remotely-sensed imagery, panchromatic, stereoscopic 1232 aerial photographs can provide the most accurate approach (in terms of landform identification), with 1233 photographs of this format having superior tonal contrast than their digital (colour) counterparts: 1234 digital colour aerial photographs may appear 'flat' (i.e. shadows are absent or less pronounced) 1235 making it more difficult to pick out subtle features, particularly in the absence of SOCET SET stereo 1236 display software and equipment (see Section 3.2.2.2). Nevertheless, mapping from digital aerial 1237 photographs has the advantage of providing georeferenced data and avoiding the duplication of effort, 1238 with hand-drawing on acetate overlays necessitating subsequent digitisation (see Sections 3.1 and 1239 3.2). Although panchromatic aerial photographs are invariably older, temporality usually presents no 1240 issue in palaeoglaciological (non-glacierised) settings, with the critical factor being image quality.

1241

1242 Irrespective of the type of aerial photographs used for geomorphological mapping, georectification is 1243 required to ensure accurate depiction of glacial landforms on the final maps (Section 3.3). This is 1244 important for minimising potential geospatial errors that will propagate into any subsequent glacier 1245 reconstructions and analyses of glacier-climate interactions. Ideally, georectification would involve 1246 stereoscopic photogrammetry, as discussed in Section 3.3, but this approach is impractical for larger 1247 ice masses (i.e. plateau icefields and plateau ice-caps). Thus, it is necessary to apply the pragmatic 1248 solutions described in Section 3.3.1.1, namely georectifying the aerial photographs or acetate overlays 1249 to other (coarser) georeferenced digital imagery or topographic data. Conversely, geomorphological 1250 studies at the scale of individual cirque basins, valley glaciers or glacier forelands would be 1251 appropriate for topographic surveys and hence stereoscopic photogrammetry, provided (i) the 1252 accessibility of the study area permits the use of surveying equipment and (ii) camera calibration data 1253 are available (see Section 3.3).

1254

1255 In some locations, coarse to medium resolution satellite imagery may be sufficient to map the 1256 geomorphological imprint of former or formerly more extensive icefields and ice-caps (Figure 15; e.g. 1257 Glasser et al., 2005; Heyman et al., 2008; Barr and Clark, 2009, 2012; Morén et al., 2011; 1258 Hochreuther et al., 2015; Loibl et al., 2015; Gribenski et al., 2016). However, these coarse remotely-1259 sensed datasets may only allow for mapping of broad landform arrangements and patterns, rather than 1260 the intricate details of individual landforms, and preclude mapping of small features (cf. Barr and 1261 Clark, 2012; Fu et al., 2012; Stroeven et al., 2013; Blomdin et al., 2016b). The emergence of high-1262 resolution (commercial) satellite imagery may result in more widespread use of satellite imagery for 1263 mapping in alpine and plateau settings, although the benefits of increased resolution may be 1264 counteracted by prohibitive costs for large study areas (see Section 3.2.2.1).

- 1265
- 1266 1267

266 5.2.2 Field mapping in alpine and plateau settings

1268 Detailed field mapping, following the procedures outlined in Section 2.2, has been widely applied as 1269 part of geomorphological studies focused on alpine- and plateau-style ice masses (e.g. Benn, 1992; 1270 Federici et al., 2003, 2017; Lukas, 2007a; Reuther et al., 2007; Boston, 2012a; Małecki et al., 2018). 1271 Although field mapping is widely used in such settings, many studies do not explicitly report whether 1272 this entails field mapping *sensu stricto* (i.e. the procedure outlined in Section 2.2), or verification of 1273 landforms mapped from remotely-sensed data by direct ground observations ('ground truthing'). We 1274 reaffirm the points raised in Sections 2.2 and 2.3 that, whenever possible, field mapping should be 1275 combined with remote mapping in cirque glacier, valley glacier, icefield and ice-cap settings in order 1276 to identify subtle glacial landforms and test interpretations of ambiguous features. While we advocate 1277 the application of detailed field mapping, we recognise that logistical and/or financial issues may 1278 preclude this and that ground truthing (of selected areas) may only be possible. Nevertheless, some 1279 form of field survey is important in alpine and plateau settings to (i) circumvent potential issues with 1280 the quality/resolution of remotely-sensed data (e.g. poor tonal contrast) and (ii) arrive at definitive of 1281 glacial landforms and landscapes (see also Section 2.3)

1282

1283 5.3 Modern glacial settings

1284

Many contemporary glacier forelands are rapidly evolving and new landscapes are emerging. This is largely due to changes resulting from the current retreat of ice masses and exposure of previouslyglaciated terrain, leading to destabilisation of some landforms (e.g. Krüger and Kjær, 2000; Kjær and Krüger, 2001; Lukas et al., 2005; Lukas, 2011), erosion by changing meltwater routes and remoulding or complete obliteration of extant landforms in areas following a glacier re-advance or surge (e.g. Evans et al., 1999; Evans and Twigg, 2002; Evans, 2003b; Evans and Rea, 2003; Benediktsson et al., 2008). Glaciofluvial processes on active temperate glacier forelands (e.g. Iceland) often make these 1292 environments unfavourable for preservation of (small) landforms (e.g. Evans and Twigg, 2002; 1293 Evans, 2003b, Kirkbride and Winkler, 2012; Evans and Orton, 2015; Evans et al., 2016a). In addition, 1294 de-icing and sediment re-working processes prevalent in many modern glacial environments (e.g. 1295 Iceland, Svalbard) typically result in substantial ice-marginal landscape modification and topographic 1296 inversion (e.g. Etzelmüller et al., 1996; Krüger and Kjær, 2000; Kjær and Krüger, 2001; Lukas et al., 1297 2005; Schomacker, 2008; Bennett and Evans, 2012; Ewertowski and Tomczyk, 2015). Anthropogenic 1298 activity can also have considerable implications for glacial systems (Jamieson et al., 2015; Evans et 1299 al., 2016b). The rapidity, ubiquity and efficacy of these censoring processes (cf. Kirkbride and 1300 Winkler, 2012, for further details) in contemporary glacial environments should be key considerations 1301 in geomorphological mapping studies; in particular, the recognition that ice-cored features mapped at 1302 a given interval in time are not the 'final' geomorphological products (cf. Krüger and Kjær, 2000; 1303 Kjær and Krüger, 2001; Everest and Bradwell, 2003; Lukas et al., 2005, 2007; Lukas, 2007b).

1304

1305 In addition to landform preservation potential, spatial and temporal scales will be key determinants in 1306 the approaches used in mapping of ice-marginal landscapes, with studies in such settings often 1307 focused on the formation of small features (<3 m in height) on recent, short-term timescales (0-30 1308 years) (e.g. Beedle et al., 2009; Lukas, 2012; Bradwell et al., 2013; Reinardy et al., 2013; Chandler et 1309 al., 2016b) and/or evolution of the glacier foreland over a given time period (e.g. Bennett et al., 2010; 1310 Bennett and Evans, 2012; Ewertowski, 2014; Jamieson et al., 2015; Chandler et al., 2016a, b; Evans et 1311 al., 2016a). Thus, the approach to geomorphological mapping discussed in Section 5.2 require some 1312 modification, as discussed below. It is also worth noting that geomorphological mapping usually 1313 forms part of process-oriented studies in modern glacial settings (Figure 16), often with the intention 1314 of providing modern analogues for palaeo-ice masses and their geomorphological imprints (e.g. Evans 1315 et al., 1999; Evans, 2011; Schomacker et al., 2014; Benediktsson et al., 2016).

1316

1317 Geophysical surveying methods can also strengthen links between modern and ancient landform 1318 records through surveying of the internal architecture of landforms that can be directly linked to 1319 depositional processes, as well as glaciological and climatic conditions (e.g. Bennett et al., 2004; 1320 Benediktsson et al., 2009, 2010; Lukas and Sass, 2011; Midgley et al., 2013, 2018). Recent advances 1321 in geophysical imaging of sub-ice geomorphology has allowed links to be made between modern and 1322 palaeo-ice sheets (see Section 3.2.2.3), and we expect this to a growth area going forward (see also 1323 Stokes, 2018). More broadly, geophysical methods can also be used to identify the extent of buried 1324 ice, allowing an assessment of the geomorphological stability of contemporary glacier forelands (e.g. 1325 Everest and Bradwell, 2003).

1326

1327 5.3.1 Remote mapping of modern glacial settings
1328 The spatial resolution of remotely-sensed data is of critical importance in modern glacial settings: 1329 spatial resolutions commensurate with the size of the landforms being mapped and the scope of the 1330 research are required. Typically, aerial photographs or satellite imagery with GSDs of <0.5 m are used 1331 in modern glacial settings to enable mapping of small features (e.g. Benediktsson et al., 2010; Lukas, 1332 2012; Bradwell et al., 2013; Brynjólfsson et al., 2014; Lovell, 2014; Schomacker et al., 2014; 1333 Chandler et al., 2016a; Ewertowski et al., 2016; Lovell et al., 2018). LiDAR or UAV-derived DEMs 1334 are also becoming increasingly used for mapping in modern glacial environments (e.g. Brynjólfsson et 1335 al., 2014, 2016; Jónsson et al. 2014, 2016; Benediktsson et al., 2016; Chandler et al., 2016a; 1336 Ewertowski et al., 2016; Everest et al., 2017; Lovell et al., 2018). Despite the high-resolution of the 1337 imagery, some compromise on the level of detail may be necessary, such as deciding on a maximum 1338 mapping scale (e.g. 1:500-1:1000, Schomacker et al., 2014) to prevent too detailed mapping or by 1339 simplifying the mapping of certain features. In studies of low-amplitude (annual) moraines, the 1340 crestlines rather than the planforms are typically mapped, reflecting a combination of image 1341 resolution and data requirements: annual moraine sequences are often used to calculate ice-margin 1342 retreat rates and mapping of crestlines is sufficient detail for this purpose (Figure 17; Krüger, 1995; 1343 Beedle et al., 2009; Lukas, 2012; Bradwell et al., 2013; Chandler et al., 2016a, b). Moreover, this 1344 approach can actually 'normalise' the data for subsequent analyses, removing the variability of, for 1345 example, moraine-base widths that result from gravitational processes during or after moraine 1346 formation.

1347

1348 The temporality (both month and year) of imagery takes on greater significance in modern glacial 1349 environments. Depending on the purpose of the research, either the most recent high resolution 1350 remotely-sensed dataset available or a series of images from a number of intervals during a given time 1351 period are commonly required (e.g. Benediktsson et al., 2010; Bennett et al., 2010; Bradwell et al., 1352 2013; Reinardy et al., 2013; Chandler et al., 2016a; Evans et al., 2016b; Ewertowski et al., 2016). In 1353 exceptional circumstances, the research may require an annual temporal resolution; for example, 1354 aerial photographs are commonly captured annually at the beginning and end of the ablation season in 1355 many forelands of the European Alps (cf. Lukas, 2012; Zemp et al., 2015). The burgeoning use of 1356 UAVs provides very high-resolution imagery (<0.1 m GSD) of contemporary glacier forelands and 1357 the option to capture up-to-date imagery during every visit to the site, circumventing issues relating to 1358 temporal resolution. This approach is likely to come into greater usage for studies examining short-1359 term ice-marginal landscape evolution and preservation potential.

1360

Photogrammetric image processing (see Section 3.3) is arguably of most importance in contemporary glacial environments, particularly where the purpose of the mapping is to investigate small variations of the order of metres to tens of metres at short-term (0–30 years) timescales (cf. Evans, 2009). However, such constraints are not necessarily applicable where broader landsystem mapping is

1365 conducted (e.g. Evans, 2009; Evans and Orton, 2015; Evans et al., 2016a). Ideally, digital aerial 1366 photographs should be processed using stereoscopic photogrammetry techniques using GCPs 1367 collected during topographic surveys to enable the production of DEMs and orthorectified imagery 1368 with low error values (RMSEs <2 m; see Section 3.3). It is preferable to survey GCPs and capture 1369 imagery contemporaneously, with surveyed GCPs appearing in the captured aerial imagery (e.g. 1370 Evans and Twigg, 2002; Evans et al., 2006, 2012; Schomacker et al., 2014), but imagery often pre-1371 dates the geomorphological investigations and topographic surveys (e.g. Bennett et al., 2010; 1372 Bradwell et al., 2013; Chandler et al., 2016b). Alternatively, the digital aerial photographs could be 1373 processed using SfM photogrammetry methods (see Section 3.3.1.2).

1374

1375 5.3.2 Field mapping in modern glacial settings

The rapidly-changing nature of modern glacier forelands presents a number of challenges when using topographic base maps (see Section 2). Firstly, in relation to spatial limitations, topographic maps available in many settings (typically at scales of 1: 25,000 or 1: 50,000) may offer insufficient spatial resolution for mapping: (i) the relief of the small geomorphological features ubiquitous in contemporary glacial environments is often less than the contour intervals depicted on the maps; and (ii) many forelands, such as those of southeast Iceland, have limited elevation changes across the foreland (cf. Evans and Twigg, 2002; Evans et al., 2016a).

1383

1384 Publicly-available topographic maps are rarely updated frequently enough to be useful for mapping 1385 the often rapid (annual to decadal-scale) changes exhibited by modern glacier margins and proglacial 1386 landscapes. Instead, it is desirable to undertake geodetic-grade surveying (i.e. using an RTK-GPS) of 1387 landforms and measurement of high-resolution topographic profiles, where conditions allow a safe 1388 approach towards the glacier margin (e.g. Benediktsson et al., 2008; Bradwell et al., 2013). Indeed, 1389 conducting detailed surveying with geodetic-grade equipment is essential for quantifying small 1390 changes in ice-marginal/proglacial landscapes (e.g. Schomacker and Kjær, 2008; Ewertowski and 1391 Tomczyk, 2015; Korsgaard et al., 2015) and obtaining metre-scale ice-margin retreat rates from the 1392 geomorphological record (e.g. Bradwell et al., 2013; Chandler et al., 2016a). This level of detail and 1393 accuracy may be unnecessary for some glacial geomorphological studies (e.g. those focused on the 1394 overall glacial landsystem), and annotation of aerial photograph extracts may be sufficient. There are 1395 still potential temporal limitations with these approaches, namely (a) limitations imposed by the 1396 date/year of image capture when mapping on print-outs and (b) difficulties with correlating survey 1397 data with imagery, depending on the time difference and rapidity of landscape changes. In localities 1398 where (parts of) the ice-marginal/proglacial landscape cannot be satisfactorily or safely traversed, 1399 imagery and elevation control from remotely-sensed sources will be necessary (e.g. Evans et al., 1400 2016e).

- 1402 **6. Frameworks for best practice**
- 1403

Based on our review of the various mapping approaches, we here synthesise *idealised* frameworks for mapping palaeo-ice sheet geomorphological imprints (Section 6.1) and alpine and plateau-style ice mass (cirque glaciers, valley glaciers, ice-fields and ice-caps) geomorphological imprints (Section 6.2). The aim is to provide frameworks for best practice in glacial geomorphological mapping, ensuring robust and systematic geomorphological mapping programmes. The templates outlined can be modified as necessary, depending on the study area size and project scope, along with the datasets, software and time available.

1411

1412 Before outlining the idealised frameworks, we offer four general recommendations for undertaking 1413 and reporting glacial geomorphological mapping that are applicable at all scales of investigation:

- 1414
- (1) The methods, datasets and equipment employed in mapping should be clearly stated,including the resolution and format of remotely-sensed data.
- (2) Any processing methods and imagery rectification errors (RMSEs) should be reported, as
 well as mapping uncertainties (both in terms of the location of the landforms and their
 identification/classification). Where remotely-sensed datasets are obtained as pre-processed,
 georeferenced products, this should also be stated.
- (3) Establishing and reporting criteria for identifying and mapping different landforms is
 desirable (e.g. Barrell et al., 2011; Darvill et al., 2014; Bendle et al., 2017a; Lovell and
 Boston, 2017). As a minimum, this could take the form of a brief definition of the mapped
 landform (e.g. Storrar and Livingstone, 2017).
- (4) GIS software (e.g. *ArcGIS*, *QGIS*) should be used for geomorphological mapping and
 digitisation to provide georeferenced geomorphological data.
- 1427

Following the above general recommendations will provide transparency about how the mapping was compiled and what considerations were made during the process, aiding accuracy assessment, comparison and integration of geomorphological data. This is particularly valuable for the incorporation of the geomorphological mapping in large compilations (Bickerdike et al., 2016; Stroeven et al., 2016; Clark et al., 2017a) and subsequently using the data for palaeoglaciological reconstruction and/or testing numerical ice sheet models (Stokes et al., 2015; Margold et al., 2018).

1434

In relation to software (recommendation 4), some practitioners may prefer to use graphics software
packages (e.g. *Adobe Illustrator*, *Canvas X*, *CorelDRAW*) for producing final glacial
geomorphological maps (e.g. Brynjólfsson et al., 2014; Darvill et al., 2014; Chandler et al., 2016a;
Bendle et al., 2017a; Norris et al., 2017). Such graphics software can provide greater functionality

than current GIS packages for fine adjustments of the final cartographic design. However, any modification in graphics software should be kept to a minimum in order to avoid compromising the transferability of the data for other users (e.g. as shapefiles), with the focus instead on adjustments to the map symbology and ensuring optimal map presentation.

- 1443
- 1444 6.1 Palaeo-ice sheet geomorphological imprints
- 1445

1446 For mapping of palaeo-ice sheet geomorphological imprints we recommend the use of multiple 1447 remotely-sensed datasets in a synergistic and systematic process, subject to data availability and 1448 coverage (Figure 18). As a minimum, remote-sensing investigations should involve reconnaissance-1449 level mapping using multiple remotely-sensed datasets to establish the most suitable dataset (e.g. 1450 Stokes et al., 2016a). However, mapping often benefits from utilising a range of imagery types and 1451 resolutions, enabling the advantages of each respective method/dataset to be integrated to produce an 1452 accurate geomorphological map (see below). At the outset of the mapping, a decision should be made 1453 on the level of mapping detail required for particular landforms (i.e. polyline or polygon mapping), in line with the aims and requirements of the study (see Section 5.1.1). 1454

1455

1456 Initially, mapping should involve an assessment of the study area using remotely-sensed data in 1457 conjunction with existing maps and literature to identify gaps in the mapping record and localities for 1458 focused mapping. Following this reconnaissance stage, the mapper may proceed with mapping from 1459 both DEMs and satellite imagery, adding increasing levels of detail with increasingly higher 1460 resolution datasets. Recommended techniques for processing the satellite images and DEMs are 1461 outlined in Sections 3.3.2 and 3.3.3, including the generation of false-colour composites with different 1462 spectral band combinations to aid landform identification (e.g. Jansson and Glasser, 2005; Lovell et 1463 al., 2011; Storrar and Livingstone, 2017).

1464

1465 DEMs may provide a superior source of imagery as they directly record the shape of landforms, rather 1466 than the interaction of reflected radiation and topography, and therefore allow for more accurate and 1467 intuitive mapping. For example, DEMs are often particularly useful for identifying and mapping meltwater channels (e.g. Greenwood et al., 2007; Storrar and Livingstone, 2017). Specific features 1468 1469 may also only be identifiable on satellite imagery, such as low-relief corridors of glaciofluvial 1470 deposits, due to their distinctive spectral signatures (e.g. Storrar and Livingstone, 2017). Moreover, the typically superior resolution of satellite imagery may enhance landform detectability and allow for 1471 1472 more detailed mapping. Many glacial landforms are also clearly distinguishable in one or more sets of 1473 remotely-sensed data (or through using a combination of datasets).

To ensure that all landforms are mapped from remotely-sensed data, the datasets should be viewed at a variety of scales and mapping conducted through multiple passes of the area, enabling the addition of increasing levels of detail to and/or refinement of initial mapping with each pass (Norris et al., 2017). It may be advantageous to perform a final check at a small cartographic scale (e.g. 1:500,000) to ensure there are no errors in the mapping, such as duplication of landforms at image overlaps (e.g. De Angelis, 2007). The mapping should be iterative, with repeated consultations of various remotelysensed datasets throughout the process recommended.

1482

1483 In this contribution, we have focused on the use of satellite imagery and DEMs for mapping palaeo-1484 ice sheet geomorphological imprints, since these are the most widely used for practical reasons. 1485 However, aerial photograph interpretation and fieldwork should not be abandoned altogether in 1486 palaeo-ice sheet settings. Aerial photographs, where available, can be used to add further detail and 1487 refine the mapping, whilst fieldwork enables ground-truthing of remote mapping (e.g. Hättestrand and 1488 Clark, 2006; Kleman et al., 2010; Darvill et al., 2014; Evans et al., 2014). Furthermore, mapping from 1489 satellite imagery and DEMs can direct fieldwork, highlighting areas for sedimentological and 1490 stratigraphic investigations. Such studies can provide invaluable data on landform genesis, subglacial 1491 processes and ice dynamics (e.g. Livingstone et al., 2010; Evans et al., 2015; Spagnolo et al., 2016; 1492 Phillips et al., 2017; Norris et al., 2018). Remote mapping of palaeo-ice sheet geomorphology also 1493 guides targeted dating for chronological investigations and should be an essential first phase in such 1494 studies (e.g. Darvill et al., 2014, 2015).

1495

1496 6.2 Alpine and plateau-style ice mass geomorphological imprints

1497

1498 Our idealised framework for mapping alpine and plateau-style ice mass geomorphological imprints is 1499 an iterative process involving several consultations of remotely-sensed data and field mapping 1500 (Figures 19 and 20). This methodology provides a robust approach to mapping that has been broadly 1501 used in previous studies (e.g. Benn and Ballantyne, 2005; Lukas and Lukas, 2006; Kjær et al., 2008; 1502 Boston, 2012a, b; Brynjólfsson et al., 2014; Jónsson et al., 2014; Pearce et al., 2014; Schomacker et 1503 al., 2014; Chandler et al., 2016a; Chandler and Lukas, 2017). This framework is also applicable to 1504 modern glacial settings as the overarching methods do not differ fundamentally, but practitioners 1505 should be aware of issues relating to the temporal resolution of remotely-sensed data (see Section 1506 5.3).

1507

In the initial preparatory stage, the mapper should consult topographic, geological and extant geomorphological maps (where available), and ideally undertake mapping of the study area using remotely-sensed data, at least at a reconnaissance level. This essential phase familiarises the mapper with the study area prior to fieldwork and enables the identification of significant areas for targeted, detailed field mapping (or ground verification) and sedimentological investigations of specific landforms. Conversely, the reconnaissance investigations may also clarify which areas are less important for a field visit and aid route planning. Importantly, this enables a systematic approach to mapping, and is particularly important in previously-unmapped areas (e.g. Boston, 2012a, b). During the initial stage, it may also be desirable to establish a legend/mapping system in readiness for subsequent field mapping (Otto and Smith, 2013).

1518

1519 Following the preparatory/reconnaissance stage, detailed field mapping, or at a minimum some 1520 ground verification, should ideally be conducted to avoid overlooking (subtle) landforms and 1521 misinterpreting others. Depending on the nature of the project and accessibility limitations, ground 1522 verification may be done during a single (and relatively short) field visit (e.g. Lukas, 2012; Chandler 1523 et al., 2016a), whilst detailed field mapping would usually require longer field visits or even repeated, 1524 long-term field campaigns (e.g. Kjær et al. 2008; Boston, 2012a, b; Schomacker et al., 2014; Evans et 1525 al., 2016a). During field surveys, consultation of initial remote mapping helps to ensure accurate 1526 representation of landforms on field maps and allows verification of all features identified remotely 1527 (e.g. Boston, 2012a, b; Pearce et al., 2014).

1528

1529 Following field mapping, which may be an intermittent and ongoing process in the case of large study 1530 areas and long-term research projects, it is ideal to finalise the geomorphological mapping using high-1531 resolution imagery (i.e. aerial photographs, satellite imagery, LiDAR DEMs, UAV-derived imagery). 1532 This allows complex patterns of landforms, such as Scottish 'hummocky moraine' (e.g. Lukas and 1533 Lukas, 2006; Boston, 2012b), crevasse-squeeze ridges (e.g. Kjær et al., 2008), drumlin fields (e.g. 1534 Benediktsson et al., 2016), and sawtooth 'annual' moraines (e.g. Chandler et al., 2016a; Evans et al., 1535 2016a), to be mapped with high spatial accuracy, following landform identification and interpretation 1536 in the field. Again, during this stage, previous mapping from DEMs and field maps should be 1537 consulted. As highlighted in the scale-appropriate examples, the procurement of remotely-sensed data 1538 with appropriate spatial and temporal resolution is important (see Sections 5.2 and 5.3).

1539

Depending on the type of imagery used (hard-copy or digital), the rectification of imagery/overlays may precede or follow aerial photograph mapping: where digital format aerial photographs are used, rectification will be undertaken before mapping (Figure 19), whilst acetate overlays will be corrected *after* mapping from hard-copy aerial photographs (Figure 20) (see also Supplementary Material). Subsequently, acetate overlays can be checked against digital imagery (if available) before being digitised, either in a GIS software package (e.g. *ArcMap, QGIS*) or in a graphics software package.

1546

In our view, geomorphological mapping in cirque glacier, valley glacier and icefield/ice-cap outletsettings should not be reliant solely on the morphological characteristics of features and should ideally

1549 be combined with detailed sedimentological investigations of available exposures as part of an 1550 inductive-deductive process, using standard procedures (cf. Evans and Benn, 2004; Lukas et al., 2013, 1551 and references therein). This reflects the fact that these glacier systems occupy more manageable 1552 study areas and therefore sedimentological analyses can be more readily applied. By combining 1553 geomorphological mapping and sedimentology, issues relating to equifinality (Chorley, 1962; Möller 1554 and Dowling, 2018) will be avoided, which is important when attempting to establish the wider 1555 palaeoglaciological and palaeoclimatic significance of the geomorphological evidence (cf. Benn and 1556 Lukas, 2006). This multi-proxy, process-form approach ensures accurate genetic interpretations on 1557 geomorphological maps.

1558

1559 **7. Conclusions**

1560

Geomorphological mapping forms the basis of a wide range of process-oriented, glacial chronological and palaeoglaciological studies. As such, it is imperative that effective approaches are used to ensure robust assimilation of data and that errors and uncertainties are explicitly reported. This is particularly the case where field mapping and analogue data are transferred to digital format and combined with digital remotely-sensed data.

1566

1567 In general, specific methods and datasets are often applied to particular glacial settings: (i) a mixture 1568 of satellite imagery (e.g. Landsat) and DEMs (e.g. ASTER GDEM, SRTM) are typically used for 1569 mapping in palaeo-ice sheet settings; and (ii) a combination of aerial photographs and field mapping 1570 are widely employed for mapping alpine and plateau-style ice mass geomorphological imprints. 1571 Increasingly, UAV-captured aerial imagery and high resolution DEMs (derived from UAV-captured 1572 imagery and LiDAR) are being utilised for mapping of modern glacial environments and are likely to 1573 be a growth area in future geomorphological mapping studies, enabling high resolution, multi-1574 temporal remotely-sensed datasets to be obtained at relatively low cost. The usage of particular 1575 methods reflects the spatial and temporal resolution of remotely-sensed datasets, along with the 1576 practicality of their application (both in terms of time and finance).

1577

In this contribution, we have highlighted that compromises and pragmatic solutions are often necessary in glacial geomorphological mapping, particularly with respect to processing techniques and the level of mapping detail. For example, detailed GNSS surveys using geodetic-grade equipment are desirable for photogrammetric processing of aerial photographs, but this is impractical for the large areas covered by icefields, ice-caps and ice sheets. Thus, pragmatic approaches may be used, such as georeferencing analogue-derived mapping to existing (coarser) georeferenced datasets (e.g. satellite imagery, DEMs or orthophotographs). In relation to the level of mapping detail, it is often

necessary to map particular landforms as linear features (e.g. subglacial bedforms, moraines) or define
a maximum scale during mapping, due to image resolution and/or study requirements.

1587

1588 Based on our review, we have outlined idealised frameworks and general recommendations to ensure 1589 best practice in future studies. In particular, we emphasise the importance of utilising multiple 1590 datasets or mapping approaches in synergy, akin to multi-proxy/-method approaches used in many 1591 Earth Science disciplines; multiple remotely-sensed datasets in the case of ice sheet-scale 1592 geomorphology and a combination of remote sensing and field mapping for circu glaciers to ice-1593 caps. Further key recommendations are the clear reporting of (i) the methods, datasets and equipment 1594 employed in mapping, (ii) any processing methods employed and imagery rectification errors 1595 (RMSEs) associated with imagery, along with mapping uncertainties, (iii) the criteria for identifying 1596 and mapping different landforms. We also recommend that mapping is conducted in GIS software to 1597 provide georeferenced geomorphological data. Finally, we advocate sedimentological investigations 1598 of available exposures as part of an inductive-deductive process during fieldwork to ensure accurate 1599 genetic interpretations of the geomorphological record as part of a holistic approach. Following these 1600 recommendations will aid in comparison, integration and accuracy assessment of geomorphological 1601 data, particularly where geomorphological data are incorporated in large compilations and 1602 subsequently used for palaeoglaciological reconstruction.

1603

1604 Acknowledgements

1605

1606 We are grateful to numerous colleagues for informal discussions that have directly or indirectly 1607 helped shape this paper. Alex Clayton is thanked for kindly supplying the UAV imagery and DEM for 1608 the Skálafellsjökull foreland, whilst Jon Merritt is thanked for providing CMB and SL with access to 1609 aerial photographs at the British Geological Survey in Edinburgh. We are also grateful to Jacob 1610 Bendle, Natacha Gribenski and Sophie Norris for kindly providing figures for inclusion in this contribution. The NEXTMap Great BritainTM data for Ben More Coigach was licensed to BMPC by 1611 1612 the NERC Earth Observation Data Centre under a Demonstration Use License Agreement. CMB and 1613 HL obtained access to aerial photographs and NEXTMap Great BritainTM data through NERC Earth 1614 Observation Data Centre whilst in receipt of NERC Algorithm studentships NE/G52368X/1 (CMB) 1615 and NE/I528050/1 (HL). This contribution was written whilst BMPC was in receipt of a Queen Mary 1616 Natural and Environmental Science Studentship, which is gratefully acknowledged. We thank Richard Waller and an anonymous reviewer for constructive comments that helped improve the clarity of this 1617 1618 contribution, along with Ian Candy for editorial handling.

1619

1620 **References**

- Alexanderson, H., Adrielsson, L., Hjort, C., Möller, P., Antonov, O., Eriksson, S., Pavlov, M., 2002.
 Depositional history of the North Taymyr ice-marginal zone, Siberia—a landsystem
 approach. J. Quat. Sci. 17(4), 361-382.
- Allaart, L., Friis, N., Ingólfsson, Ó., Håkansson, L., Noormets, R., Farnsworth, W.R., Mertes, J.,
 Schomacker, A., 2018. Drumlins in the Nordenskiöldbreen forefield, Svalbard. Gff, in press.
 doi: 10.1080/11035897.2018.1466832
- Andrews, J.T., Miller, G.H., 1979. Glacial erosion and ice sheet divides, northeastern Laurentide Ice
 Sheet, on the basis of the distribution of limestone erratics. Geology 7(12), 592–596.
- Astakhov, V., Shkatova, V., Zastrozhnov, A., Chuyko, M., 2016. Glaciomorphological Map of the
 Russian Federation. Quat. Int. 420, 4–14.
- Atkinson, N., Pawley, S., Utting, D.J., 2016. Flow-pattern evolution of the Laurentide and Cordilleran
 ice sheets across west-central Alberta, Canada: implications for ice sheet growth, retreat and
 dynamics during the last glacial cycle. J. Quat. Sci. 31(7), 753–768.
- Aylsworth, J.M., Shilts, W.W., 1989. Glacial features around the Keewatin Ice Divide: Districts of
 Mackenzie and Keewatin. Geological Survey of Canada paper 88-24.
- Bakke, J., Dahl, S.O., Paasche, Ø., Løvlie, R., Nesje, A., 2005. Glacier fluctuations, equilibrium-line
 altitudes and palaeoclimate in Lyngen, northern Norway, during the Lateglacial and
 Holocene. The Holocene 15, 518–540.
- Ballantyne, C.K., 1989. The Loch Lomond Readvance on the Isle of Skye, Scotland: glacier
 reconstruction and palaeoclimatic implications. J. Quat. Sci. 4, 95–108.
- Ballantyne, C.K., 2002. The Loch Lomond Readvance on the Isle of Mull, Scotland: glacier
 reconstruction and palaeoclimatic implications. J. Quat. Sci. 17, 759–771.
- Ballantyne, C.K., 2007a. The Loch Lomond Readvance on north Arran, Scotland: glacier
 reconstruction and palaeoclimatic implications. J. Quat. Sci. 22, 343–359.
- Ballantyne, C.K., 2007b. Loch Lomond Stadial glacier on North Harris, Outer Hebrides, North-West
 Scotland: glacier reconstruction and palaeoclimatic implications. Quat. Sci. Rev. 26, 3134–
 3149.
- Barchyn, T.E., Dowling, T.P.F., Stokes, C.R., Hugenholtz, C.H., 2016. Subglacial bed form
 morphology controlled by ice speed and sediment thickness. Geophys. Res. Lett. 43, 7572–
 7580.
- Barr, I.D., Clark, C.D., 2009. Distribution and pattern of moraines in Far NE Russia reveal former
 glacial extent. J. Maps 5, 186–193.
- 1654 Barr, I.D., Clark, C.D., 2012. An updated moraine map of Far NE Russia. J. Maps 8, 431–436.
- Barrell, D.J.A., Andersen, B.G., Denton, G.H. and Lyttle, B.S., 2011. Glacial geomorphology of the
 central South Island, New Zealand. GNS Science Monograph 27. GNS Science, Lower Hutt,
 81 pp + map (5 sheets).
- Barrell, D.J.A., Andersen, B.G., Denton, G.H. and Lyttle, B.S., 2013. Glacial geomorphology of the
 central South Island, New Zealand digital data. GNS Science Monograph 27a. GNS
 Science, 17 pp (GIS digital data files + explanatory notes).
- Barrow, G., Hinxman, L.W., Cunningham Craig, E.H., 1913. The Geology of Upper Strathspey,
 Gaick, and the Forest of Atholl (Explanation of Sheet 64). Memoirs of the Geological Survey,
 HMSO, Edinburgh, Scotland.
- Batchelor, C., Dowdeswell, J.A., Ottesen, D., 2017. Submarine Glacial Landforms. In: Micallef, A.,
 Krastel, S., Savini, A. (Eds.), Submarine Geomorphology. Springer, Cham, Switzerland: pp. 207–234.
- Beedle, M.J., Menounos, B., Luckman, B.H., Wheate, R., 2009. Annual push moraines as climate
 proxy. Geophys. Res. Lett. 36, L20501.
- Bendle, J.M., Glasser, N.F., 2012. Palaeoclimatic reconstruction from Lateglacial (Younger Dryas
 Chronozone) cirque glaciers in Snowdonia, North Wales. Proceedings of the Geologists'
 Association 123, 130–145.
- Bendle, J.M., Thorndycraft, V.R., Palmer, A.P., 2017a. The glacial geomorphology of the Lago
 Buenos Aires and Lago Pueyrredón ice lobes of central Patagonia. J. Maps 13, 654–673.
- Bendle, J.M., Palmer, A.P., Thorndycraft, V.R., Matthews, I.P., 2017b. High-resolution chronology
 for deglaciation of the Patagonian Ice Sheet at Lago Buenos Aires (46.5° S) revealed through
 varve chronology and Bayesian age modelling. Quat. Sci. Rev. 177, 314-339.

- Benediktsson, Í.Ö., Möller, P., Ingólfsson, Ó., van der Meer, J.J.M., Kjær, K.H., Krüger, J., 2008.
 Instantaneous end moraine and sediment wedge formation during the 1890 glacier surge of Brúarjökull, Iceland. Quat. Sci. Rev. 27, 209–234.
- Benediktsson, Í.Ö., Ingólfsson, Ó., Schomacker, A., Kjaer, K.H., 2009. Formation of submarginal and
 proglacial end moraines: implications of ice-flow mechanism during the 1963-64 surge of
 Brúarjökull, Iceland. Boreas 38, 440–457.
- Benediktsson, Í.Ö., Schomacker, A., Lokrantz, H., Ingólfsson, Ó., 2010. The 1890 surge end moraine
 at Eyjabakkajökull, Iceland: a re-assessment of a classic glaciotectonic locality. Quat. Sci.
 Rev. 29, 484–506.
- Benediktsson, Í.Ö., Schomacker, A., Johnson, M.D., Geiger, A.J., Ingólfsson, Ó., Guðmundsdóttir,
 E.R. 2015. Architecture and structural evolution of an early Little Ice Age terminal moraine at
 the surge-type glacier Múlajökull, Iceland. J. Geophys. Res.: Earth Surf. 120, 1895–1910.
- Benediktsson, Í.Ö., Jónsson, S.A., Schomacker, A., Johnson, M.D., Ingólfsson, Ó., Zoet, L., Iverson,
 N.R., Stötter, J., 2016. Progressive formation of modern drumlins at Múlajökull, Iceland:
 stratigraphical and morphological evidence. Boreas, 45, 567–583.
- Benn, D.I., 1990. Scottish Lateglacial moraines: debris supply, genesis and significance. Unpublished
 PhD thesis, University of St Andrews.
- Benn, D.I., 1994. Fluted moraine formation and till genesis below a temperate valley glacier:
 Slettmarkbreen, Jotunheimen, southern Norway. Sedimentology 41, 279–292.
- Benn, D.I. 2006. Interpreting glacial sediments. In: Knight, P. (Ed.) Glacier Science and Environmental Change. Oxford, Blackwell, pp. 434-439.
- Benn, D.I., Ballantyne, C.K., 2005. Palaeoclimatic reconstruction from Loch Lomond Readvance
 glaciers in the West Drumochter Hills, Scotland. J. Quat. Sci. 20, 577–592.
- Benn, D.I., Evans, D.J.A., 2010. Glaciers and Glaciation (2nd Edition). Hodder Education, London, 802 pp.
- Benn, D.I., Lukas, S., 2006. Younger Dryas glacial landsystems in North West Scotland: an
 assessment of modern analogues and palaeoclimatic implications. Quat. Sci. Rev. 25, 2390–
 2408.
- Benn, D.I., Lowe, J.J., Walker, M.J.C., 1992. Glacier response to climatic change during the Loch
 Lomond Stadial and early Flandrian: geomorphological and palynological evidence from the
 Isle of Skye, Scotland. J. Quat. Sci. 7, 125–144.
- Bennett, G.L., Evans, D.J.A., 2012. Glacier retreat and landform production on an overdeepened
 glacier foreland: the debris-charged glacial landsystem at Kvíárjökull, Iceland. Earth Surf.
 Process. Land. 37, 1584–1602.
- Bennett, G.L., Evans, D.J.A., Carbonneau, P., Twigg, D.R., 2010. Evolution of a debris-charged
 glacier landsystem, Kvíárjökull, Iceland. J. Maps 6, 40–67.
- Bennett, M.R., 1991. Scottish "hummocky moraine": its implications for the deglaciation of the North
 West Highlands during the Younger Dryas or Loch Lomond Stadial. University of Edinburgh,
 Unpublished PhD thesis, 362 pp.
- Bennett, M.R., Huddart, D., Waller, R.I., Cassidy, N., Tomio, A., Zukowskyj, P., Midgley, N.G.,
 Cook, S.J., Gonzalez, S., Glasser, N.F., 2004. Sedimentary and tectonic architecture of a large
 push moraine: a case study from Hagafellsjökull-Eystri, Iceland. Sedi. Geol. 172, 269–292.
- Bickerdike, H.L., Evans, D.J.A., Ó Cofaigh, C., Stokes, C.R., 2016. The glacial geomorphology of the
 Loch Lomond Stadial in Britain: a map and geographic information system resource of
 published evidence. J. Maps 12(5), 1178–1186.
- Bickerdike, H.L., Ó Cofaigh, C., Evans, D.J.A., Stokes, C.R., 2018. Glacial landsystems, retreat
 dynamics and controls on Loch Lomond Stadial (Younger Dryas) glaciation in Britain.
 Boreas 47, 202–224.
- Bickerton, R.W., Matthews, J.A., 1992. On the accuracy of lichenometric dates: an assessment based
 on the 'Little Ice Age' moraine sequence of Nigardsbreen, southern Norway. The Holocene 2,
 227–237.
- Bickerton, R.W., Matthews, J.A., 1993. 'Little ice age' variations of outlet glaciers from the
 jostedalsbreen ice-cap, Southern Norway: A regional lichenometric-dating study of icemarginal moraine sequences and their climatic significance. J. Quat. Sci. 8, 45–66.

- Blomdin, R., Stroeven, A.P., Harbor, J.M., Lifton, N.A., Heyman, J., Gribenski, N., Petrakov, D.A.,
 Caffee, M.W., Ivanov, M.N., Hättestrand, C., Rogozhina, I., Usubaliev, R., 2016a. Evaluating
 the timing of former glacier expansions in the Tian Shan: A key step towards robust spatial
 correlations. Quat. Sci. Rev. 153, 78–96.
- Blomdin, R., Heyman, J., Stroeven, A.P., Hättestrand, C., Harbor, J.M., Gribenski, N., Jansson, K.N.,
 Petrakov, D.A., Ivanov, M.N., Alexander, O., Rudoy, A.N., Walther, M., 2016b. Glacial
 geomorphology of the Altai and Western Sayan Mountains, Central Asia. J. Maps 12(1), 123–
 136.
- Bolch, T., Loibl, D., 2017. GIS for Glaciers and Glacial Landforms. Reference Module in Earth
 Systems and Environmental Sciences.
- Borsellino, R., Shulmeister, J., Winkler, S., 2017. Glacial geomorphology of the Barbizon & Butler
 Downs, Rangitata Valley, South Island, New Zealand. J. Maps 13, 502–510.
- Boston, C.M., 2012a. A Lateglacial plateau icefield in the Monadhliath Mountains, Scotland:
 reconstruction, dynamics and palaeoclimatic implications. Unpublished PhD thesis, Queen
 Mary University of London, 295 pp.
- Boston, C.M., 2012b. A glacial geomorphological map of the Monadhliath Mountains, Central
 Scottish Highlands. J. Maps 8(4), 437–444.
- Boston, C.M., Lukas, S., Carr, S.J. 2015. A Younger Dryas plateau icefield in the Monadhliath,
 Scotland, and implications for regional palaeoclimate. Quat. Sci. Rev. 108, 139–162.
- Boulton, G.S., Clark, C.D., 1990a. A highly mobile Laurentide ice sheet revealed by satellite images
 of glacial lineations. Nature 346(6287), 813–817.
- Boulton, G.S., Clark, C.D., 1990b. The Laurentide ice sheet through the last glacial cycle: the topology of drift lineations as a key to the dynamic behaviour of former ice sheets. Earth Environ. Sci. Trans. R. Soc. Edinb. 81(4), 327–347.
- Boulton, G.S., Dongelmans, P., Punkari, M., Broadgate, M., 2001. Palaeoglaciology of an ice sheet
 through a glacial cycle: the European ice sheet through the Weichselian. Quat. Sci. Rev.
 20(4), 591–625.
- Bradwell, T., Stoker, M.S., Golledge, N.R., Wilson, C.K., Merritt, J.W., Long, D., Everest, J.D.,
 Hestvik, O.B., Stevenson, A.G., Hubbard, A.L., Finlayson, A.G., Mathers, H.E., 2008. The
 northern sector of the last British Ice Sheet: maximum extent and demise. Earth-Sci. Rev. 88,
 207–226.
- Bradwell, T., Sigurðsson, O., Everest, J., 2013. Recent, very rapid retreat of a temperate glacier in SE
 Iceland. Boreas 42, 959–973.
- 1764 Brodzikowski, K., van Loon, A.J., 1991. Glacigenic Sediments. Elsevier, Amsterdam, 674 pp.
- Brown, V.H., Evans, D.J.A., Evans, I.S., 2011a. The Glacial Geomorphology and Surficial Geology of the South-West English Lake District. J. Maps 7(1), 221–243.
- Brown, V.H., Stokes, C.R., Ó Cofaigh, C., 2011b. The Glacial Geomorphology of the North-West
 sector of the Laurentide Ice Sheet. J. Maps 7, 409–428.
- Brynjólfsson, S., Schomacker, A., Ingólfsson, Ó., 2014. Geomorphology and the Little Ice Age extent
 of the Drangajökull ice cap, NW Iceland, with focus on its three surge-type outlets.
 Geomorphology 213, 292–304.
- Brynjólfsson, S., Schomacker, A., Korsgaard, N.J., Ingólfsson, Ó. 2016. Surges of outlet glacier from
 the Drangajökull ice cap, northwest Iceland. Earth Planet. Sci. Lett. 450, 140–151.
- Butler, J., Lane, S., Chandler, J., 1998. Assessment of DEM quality for characterizing surface
 roughness using close range digital photogrammetry. The Photogramm. Rec. 16(92), 271–
 291.
- 1777 Caldenius, C.C., 1932. Las glaciaciones cuaternarias en la Patagonia y Tierra del Fuego. Geogr. Ann.
 1778 14, 1–164.
- 1779 Campbell, J.B., Wynne, R.H., 2011. Introduction to remote sensing (5th Edition). Taylor and Francis,
 1780 London, 667 pp.
- Campbell, R.B., 1967a. Geology of Glenlyon map-area, Yukon Territory. Geological Survey of
 Canada, Memoir 352, 92 pp.
- 1783 Campbell, R.B., 1967b. Surficial Geology, Glenlyon, Yukon Territory. Geological Survey of Canada,
 1784 "A" Series Map 1222A, 1: 253,440.

- Carbonneau, P.E., Dietrich, J.T., 2017. Cost-effective non-metric photogrammetry from consumer grade sUAS: implications for direct georeferencing of structure from motion
 photogrammetry. Earth Surf. Proc. Land. 42, 473–486.
- Carrivick, J.L., Smith, M.W., Quincey, D.J., 2016. Structure from Motion in the Geosciences. New
 Analytical Methods in Earth and Environmental Science Series. John Wiley & Sons,
 Chichester, UK, 208 pp.
- Carrivick, J.L., Yde, J., Russell, A.J., Quincey, D.J., Ingeman-Nielsen, T., Mallalieu, J., 2017. Ice margin and meltwater dynamics during the mid-Holocene in the Kangerlussuaq area of west
 Greenland. Boreas 46, 369–387.
- 1794 Chamberlin, T.C., 1897/1965. The method of multiple working hypotheses. Science 148, 745–759.
- Chandler, B.M.P., Lukas, S., 2017. Reconstruction of Loch Lomond Stadial (Younger Dryas) glaciers
 on Ben More Coigach, NW Scotland, and implications for reconstructing palaeoclimate using
 small ice masses. J. Quat. Sci. 32(4), 475–492.
- 1798 Chandler, B.M.P., Evans, D.J.A., Roberts, D.H., Ewertowski, M.W., Clayton, A.I., 2016a. Glacial
 1799 geomorphology of the Skálafellsjökull foreland, Iceland: A case study of 'annual' moraines.
 1800 J. Maps 12(5), 904–916.
- 1801 Chandler, B.M.P., Evans, D.J.A., Roberts, D.H., 2016b. Characteristics of recessional moraines at a
 1802 temperate glacier in SE Iceland: Insights into patterns, rates and drivers of glacier retreat.
 1803 Quat. Sci. Rev. 135, 171–205.
- 1804 Chorley, R.J., 1962. Geomorphology and General Systems Theory. United States Geological Survey,
 1805 Professional Paper 500B.
- 1806 Clark, C.D., 1993. Mega-scale glacial lineations and cross-cutting ice-flow landforms. Earth Surf.
 1807 Proc. Land. 18(1), 1–29.
- 1808 Clark, C.D. 1997. Reconstructing the evolutionary dynamics of former ice sheets using multitemporal evidence, remote sensing and GIS. Quat. Sci. Rev. 16, 1067–1092.
- 1810 Clark, C.D., Stokes, C.R., 2001. Extent and basal characteristics of the M'Clintock Channel Ice
 1811 Stream. Quat. Int. 86, 81–101.
- 1812 Clark, C.D., Knight, J.K., Gray, J.T., 2000. Geomorphological reconstruction of the Labrador sector
 1813 of the Laurentide Ice Sheet. Quat. Sci. Rev. 19(13), 1343–1366.
- 1814 Clark, C.D., Evans, D.J.A., Khatwa, A., Bradwell, T., Jordan, C.J., Marsh, S.H., Mitchell, W.A.,
 1815 Bateman, M.D., 2004. Map and GIS database of glacial landforms and features related to the
 1816 last British Ice Sheet. Boreas 33, 359–375.
- 1817 Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Spagnolo, M., Ng, F.S.L., 2009. Size and shape
 1818 characteristics of drumlins, derived from a large sample, and associated scaling laws. Quat.
 1819 Sci. Rev. 28, 677–692.
- Clark, C.D., Hughes, A.L.C., Greenwood, S.L., Jordan, C., Sejrup, H.P., 2012. Pattern and timing of
 retreat of the last British-Irish Ice Sheet. Quat. Sci. Rev. 44, 112–146.
- 1822 Clark, C.D., Ely, J.C., Greenwood, S.L., Hughes, A.L.C., Meehan, R., Barr, I.D., Bateman, M.D.,
 1823 Bradwell, T., Doole, J., Evans, D.J.A., Jordan, C.J., Monteys, X., Pellicer, X.M., Sheehy, M.,
 1824 2018a. BRITICE Glacial Map, version 2: a map and GIS database of glacial landforms of the
 1825 last British–Irish Ice Sheet. Boreas, 47, 11–27.
- Clark, C.D., Ely, J.C., Spagnolo, M., Hahn, U., Hughes, A.L.C., Stokes, C.R., 2018b. Spatial
 organization of drumlins. Earth Surf. Proc. Land., 43, 499–513.
- 1828 Clayton, L., Teller, J.T., Attig, J.W., 1985. Surging of the southwestern part of the Laurentide Ice
 1829 Sheet. Boreas 14(3), 235–241.
- 1830 Close, M.H. 1867. Notes on the general glaciation of Ireland. J. R. Geogr. Soc. Lond. Dublin 1, 207–
 1831 242.
- 1832 Coray, S., 2007. Glazialsedimentologische Untersuchungen an Flutes im Vorfeld des
 1833 Findelengletschers. Unpublished MSc Thesis, Universität Bern, 116 pp.
- 1834 Coronato, A., Seppälä, M., Ponce, J.F., Rabassa, J., 2009. Glacial geomorphology of the Pleistocene
 1835 lake Fagnano ice lobe, Tierra del Fuego, southern South America. Geomorphology 112(1),
 1836 67–81.
- 1837 Craig, B.G., 1961. Surficial geology of northern District of Keewatin, Northwest Territories.
 1838 Geological Survey of Canada, Map 7-1961, Scale 1: 1,013,760.

- 1839 Craig, B.G., 1964. Surficial geology of east-central District of Mackenzie. Geological Survey of
 Canada, Bulletin, 99.
- 1841 Cumming, I.G., Wong, F.H., 2005. Digital Processing of Synthetic Aperture Radar Data: Algorithms
 1842 and Implementation. Artech House, Boston.
- da Rosa, K.K., Vieira, R., Fernandez, G.B., Simoes, F.L., Simoes, J.C., 2011. Formas glaciales y
 procesos glaciológicos del glaciar templado Wanda, Shetland del Sur. Investig. Geogr. 43, 3–
 16.
- da Rosa, K.K., 2013a. The landforms and pattern of deglaciation of the Dragon glacier, King George
 Island, South Shetlands, Antarctica. Rev. Geogr. (Recife) 30(2), 1–16.
- da Rosa, K.K., Vieira, R., Júnior, C.W.M., de Souza Júnior, E., Simões, J.C., 2013b. Compilation of geomorphological map for reconstructing the deglaciation of ice-free areas in the Martel Inlet, King George Island, Antarctica. Rev. Bras. Geomorfol. 14(2), 181–187.
- 1851 Darvill, C.M., Stokes, C.R., Bentley, M.R., Lovell, H., 2014. A glacial geomorphological map of the
 1852 southernmost ice lobes of Patagonia: the Bahía Inútil San Sebastián, Magellan, Otway,
 1853 Skyring and Río Gallegos lobes. J. Maps 10(3), 500–520.
- Darvill, C.M., Bentley, M.J., Stokes, C.R., Hein, A.S., Rodés, Á., 2015. Extensive MIS 3 glaciation in
 southernmost Patagonia revealed by cosmogenic nuclide dating of outwash sediments. Earth
 Planet. Sci. Lett. 429, 157–169.
- Darvill, C.M., Stokes, C.R., Bentley, M.J., Evans, D.J.A., Lovell, H., 2017. Dynamics of former ice
 lobes of the southernmost Patagonian Ice Sheet based on a glacial landsystems approach. J.
 Quat. Sci. 32, 857–876.
- 1860 De Angelis, H., 2007. Glacial geomorphology of the east-central Canadian Arctic. J. Maps 3, 323–
 1861 341.
- 1862 De Angelis, H. Kleman, J., 2007. Palaeo-ice streams in the Foxe/Baffin sector of the Laurentide Ice
 1863 Sheet. Quat. Sci. Rev. 26(9), 1313–1331.
- 1864 De Geer, G. 1910. Den svenska Setsbergsexkursionen 1910 för deltagare i den 11:te internationella
 1865 geologkonferensen i Stockholm. Ymer 30, 305-310.
- 1866 Demek, J., 1972. Manual of detailed geomorphological mapping. IUG, Prague, 344 pp.
- 1867 Dowdeswell, J.A., Vásquez, M., 2013. Submarine landforms in the fjords of southern Chile:
 1868 implications for glacimarine processes and sedimentation in a mild glacier-influenced
 1869 environment. Quat. Sci. Rev. 64, 1–19.
- 1870 Dowdeswell, J.A., Canals, M., Jakobsson, M., Todd, B. J., Dowdeswell, E.K., Hogan, K.A., 2016.
 1871 Atlas of Submarine Glacial landforms: Modern, Quaternary and Ancient. Geological Society, 1872 London, Memoirs, No. 46.
- 1873 Dowling, T.P.F., Alexanderson, H., Möller, P., 2013. The new high-resolution LiDAR digital height
 1874 model ('Ny Nationell Höjdmodell') and its application to Swedish Quaternary
 1875 geomorphology. GFF 135(2), 145–151.
- 1876 Dowling, T.P.F., Spagnolo, M., Möller, P., 2015. Morphometry and core type of streamlined
 1877 bedforms in southern Sweden from high resolution LiDAR. Geomorphology 236, 54–63.
- 1878 Dowling, T.P.F., Möller, P., Spagnolo, M., 2016. Rapid subglacial streamlined bedform formation at a
 1879 calving bay margin. J. Quat. Sci. 31, 879–892.
- Dunstone, R.B., 2014. Testing the groove-ploughing theory for mega-scale glacial lineaton (MSGL)
 formation, using a large dataset of their morphology. Unpublished MSc thesis, University of
 Durham, 127 pp. [Available at Durham e-theses: http://etheses.dur.ac.uk/9457/]
- 1883 Duveiller, G., Defourny, P., 2010. A conceptual framework to define the spatial resolution
 1884 requirements for agricultural monitoring using remote sensing. Remote Sens. Environ.
 1885 114(11), 2637-2650.
- 1886 Dyke, A.S. 1990. Quaternary geology of the Frances Lake map area, Yukon and Northwest
 1887 Territories. Geological Survey of Canada, Memoir 426.
- 1888 Dyke, A.S., Hooper, J.M.G., 2001. Deglaciation of northwest Baffin Island, Nunavut. Geological
 1889 Survey of Canada Map 1999A, Scale 1: 500,000.
- 1890 Dyke, A.S., Prest, V.K., 1987a. Late Wisconsinan and Holocene History of the Laurentide Ice Sheet.
 1891 Géogr. Phys. Quat. 41, 237–263.
- 1892 Dyke, A.S., Prest, V.K., 1987b. Late Wisconsinan and Holocene retreat of the Laurentide Ice Sheet,
 1893 Map 1702A. Geological Survey of Canada, Ottawa.

- 1894 Dyke, A.S., Prest, V.K., 1987c. Paleogeography of northern North America, 18 000 5 000 years ago,
 1895 Map 1703A. Geological Survey of Canada, Ottawa.
- 1896 Dyke, A.S., Morris, T.F., Green, D.E.C., England, J., 1992. Quaternary geology of Prince of Wales
 1897 Island, arctic Canada. Geological Survey of Canada Memoir 433.
- 1898 Eisank, C., Smith, M., Hillier, J., 2014. Assessment of multiresolution segmentation for delimiting
 1899 drumlins in digital elevation models. Geomorphology 214, 452–464.
- Ely, J.C., Clark, C.D., Spagnolo, M., Stokes, C.R., Greenwood, S.L., Hughes, A.L.C., Dunlop, P.,
 Hess, D., 2016a. Do subglacial bedforms comprise a size and shape continuum?
 Geomorphology 257, 108–119.
- Ely, J.C., Gribble, E.A., Clark, C.D., 2016b. The glacial geomorphology of the western cordilleran ice
 sheet and Ahklun ice cap, Southern Alaska. J. Maps 12(Sup. 1), 415–424.
- Ely, J.C., Graham, C., Barr, I.D., Rea, B.R., Spagnolo, M., Evans, J., 2017. Using UAV acquired photography and structure from motion techniques for studying glacier landforms: application to the glacial flutes at Isfallsglaciären. Earth Surf. Proc. Land. 42(6), 877–888.
- Ely, J.C., Clark, C.D., Spagnolo, M., Hughes, A.L., Stokes, C.R., 2018. Using the size and position of
 drumlins to understand how they grow, interact and evolve. Earth Surf. Proc. Land. 43(5),
 1073–1087.
- Ercolano, B., Coronato, A., Tiberi, P., Corbella, H., Marderwald, G., 2016. Glacial geomorphology of
 the tableland east of the Andes between the Coyle and Gallegos river valleys, Patagonia,
 Argentina. J. Maps 12(1), 304–313.
- 1914 Espinoza, J.M., Glacial geomorphology and paleoglacial behaviour estimation in the Sierra Baguales
 1915 (50°S): Paleoclimatic factors that controlled glacier variations within the Pleistocene –
 1916 Holocene regional context. Unpublished PhD thesis, Universidad de Chile.
- Etzelmüller, B., Hagen, J.O., Vatne, G., Ødegård, R.S., Sollid, J.L., 1996. Glacier debris accumulation
 and sediment deformation influenced by permafrost: examples from Svalbard. Ann. Glaciol.
 22, 53–62.
- 1920 Evans, D.J.A. (Ed.), 2003a. Glacial Landsystems. Arnold, London, 532 pp.
- Evans, D.J.A., 2003b. Ice-Marginal Terrestrial Landsystems: Active Temperate Glacier Margins. In:
 Evans, D.J.A. (Ed.), Glacial Landsystems. Arnold, London, pp. 12–43.
- Evans, D.J.A., 2009. Glacial Geomorphology at Glasgow. Scottish Geographical Journal 125, 285–
 320.
- Evans, D.J.A., 2010. Controlled moraine development and debris transport pathways in polythermal
 plateau icefields: examples from Tungnafellsjökull, Iceland. Earth Surf. Proc. Land. 35,
 1430–1444.
- Evans, D.J.A., 2011. Glacial landsystems of Satujökull, Iceland: A modern analogue for glacial landsystem overprinting by mountain icecaps. Geomorphology 129, 225–237.
- Evans, D.J.A., 2017. Chapter 4 Conceptual glacial ground models: British and Irish case studies. In:
 Griffiths, J.S., Martin, C.J. (Eds.), Engineering Geology and Geomorphology of Glaciated
 and Periglaciated Terrains: Engineering Group Working Party Report. Engineering Geology
 Special Publication 28. The Geological Society, London, pp. 369–500.
- Evans, D.J.A., Benn, D.I. (Eds.), 2004. A Practical Guide to the Study of Glacial Sediments. Arnold,
 London, 266 pp.
- Evans, D.J.A., Orton, C., 2015. Heinabergsjökull and Skalafellsjökull, Iceland: Active Temperate
 Piedmont Lobe and Outwash Head Glacial Landsystem. J. Maps 11(3), 415–431.
- Evans, D.J.A., Rea, B.R., 2003. Surging Glacier Landsystem. In: Evans, D.J.A. (Ed.), Glacial
 Landsystems. Arnold, London, pp. 259–288.
- Evans, D.J.A., Twigg, D.R., 2002. The active temperate glacial landsystem: a model based on Breiðamerkurjökull and Fjallsjökull, Iceland. Quat. Sci. Rev. 21, 2143–2177.
- Evans, D.J.A., Lemmen, D.S., Rea, B.R., 1999. Glacial landsystems of the southwest Laurentide Ice
 Sheet: modern Icelandic analogues. J. Quat. Sci. 14, 673–691.
- Evans, D.J.A., Twigg, D.R., Shand, M., 2006. Surficial geology and geomorphology of the bórisjökull
 plateau icefield, west-central Iceland. J. Maps 2, 17–29.
- Evans, D.J.A., Clark, C.D., Rea B.R., 2008. Landform and sediment imprints of fast glacier flow in the southwest Laurentide Ice Sheet. J. Quat. Sci. 23(3), 249–272.

- Evans, D.J.A., Twigg, D.R., Rea, B.R., Orton, C., 2009a. Surging glacier landsystem of Tungnaárjökull, Iceland. J. Maps 5, 134–151.
- Evans, D.J.A., Livingstone, S.J., Vieli, A., Ó Cofaigh, C., 2009b. The palaeoglaciology of the central sector of the British and Irish Ice Sheet: reconciling glacial geomorphology and preliminary ice sheet modelling. Quat. Sci. Rev. 28, 739–757.
- Evans, D.J.A., Strzelecki, M., Milledge, D.G., Orton, C., 2012. Hørbyebreen polythermal glacial
 landsystem, Svalbard. J. Maps 8, 146–156.
- Evans, D.J.A., Young, N.J.P., Ó Cofaigh, C., 2014. Glacial geomorphology of terrestrial-terminating
 fast flow lobes/ice stream margins in the southwest Laurentide Ice Sheet. Geomorphology
 204, 86–113.
- Evans, D.J.A., Roberts, D.H., Cofaigh, C.Ó., 2015. Drumlin sedimentology in a hard-bed, lowland
 setting, Connemara, western Ireland: implications for subglacial bedform generation in areas
 of sparse till cover. J. Quat. Sci. 30, 537-557.
- Evans, D.J.A., Ewertowski, M., Orton, C., 2016a. Fláajökull (north lobe), Iceland: active temperate
 piedmont lobe glacial landsystem. J. Maps 12(5), 777–789.
- Evans, D.J.A., Ewertowski, M., Jamieson, S.S.R., Orton, C., 2016b. Surficial geology and geomorphology of the Kumtor Gold Mine, Kyrgyzstan: human impacts on mountain glacier landsystems. J. Maps 12(5), 757–769.
- Evans, D.J.A., Ewertowski, M., Orton, C., 2016c. Eiríksjökull plateau icefield landsystem, Iceland. J.
 Maps 12(5), 747–756.
- Evans, D.J.A., Storrar, R.D., Rea, B.R., 2016d. Crevasse-squeeze ridge corridors: Diagnostic features
 of late-stage palaeo-ice stream activity. Geomorphology 258, 40–50.
- Evans, D.J.A., Ewertowski, M., Orton, C., Harris, C., Guðmundsson, S., 2016e. Snæfellsjökull
 volcano-centred ice cap landsystem, West Iceland. J. Maps 12(5) 1128–1137.
- Evans, D.J.A., Ewertowski, M., Orton, C., 2017. Skaftafellsjökull, Iceland: glacial geomorphology
 recording glacier recession since the Little Ice Age. J. Maps 13, 358–368.
- Evans, I.S., 1990. Cartographic techniques in geomorphology. In: Goudie, A.S. (Ed.),
 Geomorphological techniques (2nd Edition). Routledge, London, pp. 97–108.
- Evans I.S., 2012, Geomorphometry and landform mapping: What is a landform? Geomorphology 137,
 94–106.
- Everest, J.D., Bradwell, T., 2003. Buried glacier ice in southern Iceland and its wider significance.
 Geomorphology 52, 347–358.
- Everest, J., Bradwell, T., Jones, L., Hughes, L., 2017. The geomorphology of Svínafellsjökull and
 Virkisjökull-Falljökull glacier forelands, southeast Iceland. J. Maps 13, 936–945.
- Ewertowski, M.W., 2014. Recent transformations in the high-Arctic glacier landsystem, Ragnarbreen,
 Svalbard. Geogr. Ann. 96A(3), 265–285.
- Ewertowski, M.W., Tomczyk, A.M., 2015. Quantification of the ice-cored moraines' short-term
 dynamics in the high-Arctic glaciers Ebbabreen and Ragnarbreen, Petuniabukta, Svalbard.
 Geomorphology 234, 211–227.
- Ewertowski, M.W., Evans, D.J.A., Roberts, D.H., Tomczyk, A.M., 2016. Glacial geomorphology of
 the terrestrial margins of the tidewater glacier, Nordenskiöldbreen, Svalbard. J. Maps 12(Sup.
 1), 476–487.
- Ewertowski M.W., Kijowski A., Szuman I., Tomczyk A.M., Kasprzak L., 2017. Low-altitude remote
 sensing and GIS-based analysis of cropmarks: classification of past thermal-contraction-crack
 polygons in central western Poland. Geomorphology 293B, 418–432.
- Eyles, N. (Ed.), 1983. Glacial Geology: An Introduction for Engineers and Earth Scientists.
 Permagon, Oxford, 409 pp.
- Fabel, D., Stroeven, A.P., Harbor, J., Kleman, J., Elmore, D., Fink, D., 2002. Landscape preservation
 under Fennoscandian ice sheets determined from in situ produced 10Be and 26Al. Earth
 Planet. Sci. Lett. 201, 397–406.
- Fabel, D., Fink, D., Fredin, O., Harbor, J., Land, M., Stroeven, A.P., 2006. Exposure ages from relict
 lateral moraines overridden by the Fennoscandian ice sheet. Quat. Res. 65, 136–146.
- Federici, P.R., Pappalardo, M., Ribolini, A., 2003. Geomorphological Map of the Maritime Alps
 Natural Park and surroundings (Argentera Massif, Italy), 1: 25000 scale. Selca, Florence.

- Federici, P.R., Ribolini, A., Spagnolo, M., 2017. Glacial history of the Maritime Alps from the Last
 Glacial Maximum to the Little Ice Age. Geological Society, London, Special Publications
 433(1), 137–159.
- Finke, L., 1980. Anforderungen aus der Planungspraxis an ein geomorphologisches Kartenwerk. In:
 Barsch, D., Liedtke, H. (Eds.), Methoden und Anwendbarkeit geomorphologischer
 Detailkarten. Freie Universität Berlin, Berlin; pp. 75–81.
- Finlayson, A., Merritt, J., Browne, M., Merritt, J., McMillan, A., Whitbread, K., 2010. Ice sheet
 advance, dynamics, and decay configurations: evidence from west central Scotland. Quat. Sci.
 Rev. 29, 969–988.
- Finlayson, A.G., Golledge, N., Bradwell, T., Fabel, D., 2011. Evolution of a Lateglacial mountain
 icecap in northern Scotland. Boreas 40, 536–554.
- Flink, A.E., Noormets, R., Kirchner, N., Benn, D.I., Luckman, A., Lovell, H., 2015. The evolution of
 a submarine landform record following recent and multiple surges of Tunabreen glacier,
 Svalbard. Quat. Sci. Rev. 108, 37–50.
- Flint, R.F., Colton, R.B., Goldthwait, R.P., Willman, H.B., 1959. Glacial map of the United States
 east of the Rocky Mountains. Geological Society of America.
- Follestad, B., Bergstrøm, B., 2004. Otta 1718-IV. Quaternary geology map with description.
 Norwegian Geological Survey. 1:50,000.
- Fourrière, A., Claudin, P., Andreotti, B., 2010. Bedforms in a turbulent stream: Formation of ripples
 by primary linear instability and of dunes by nonlinear pattern coarsening, J. Fluid Mech. 649,
 287–328.
- Fu, P., Heyman, J., Hättestrand, C., Stroeven, A.P., Harbor, J.M., 2012. Glacial geomorphology of the
 Shaluli Shan area, southeastern Tibetan Plateau. J. Maps 8, 48–55.
- Garcia, J.L., Kaplan, M.R., Hall, B.L., Schaefer, J.M., Vega, R.M., Schwartz, R., Finkel, R., 2012
 Glacier expansion in southern Patagonia throughout the Antarctic cold reversal. Geology 40, 859–862
- Glasser, N.F., Jansson, K.N., 2005. Fast-flowing outlet glaciers of the last glacial maximum
 Patagonian Icefield. Quat. Res. 63(2), 206–211.
- 2030 Glasser, N., Jansson, K., 2008. The Glacial Map of southern South America. J. Maps 4, 175–196.
- Glasser, N.F., Jansson, K.N., Harrison, S., Rivera, A., 2005. Geomorphological evidence for
 variations of the North Patagonian Icefield during the Holocene. Geomorphology 71(3-4),
 263–277.
- 2034Glasser, N.F., Jansson, K.N., Harrison, S., Kleman, J., 2008. The glacial geomorphology and2035Pleistocene history of South America between 38 S and 56 S. Quat. Sci. Rev. 27(3), 365–390.
- Golledge, N.R., Stoker, M.S., 2006. A palaeo-ice-stream of the British Ice Sheet in eastern Scotland.
 Boreas 35, 231–243.
- 2038 Golledge, N.R., Hubbard, A., Sugden, D.E., 2008. High-resolution numerical simulation of Younger
 2039 Dryas glaciation in Scotland. Quat. Sci. Rev. 27, 888–904.
- 2040Goodchild, J.G., 1875. The glacial phenomena of the Eden Valley and the western part of the2041Yorkshire-dale District. Quart. J. Geol. Soc. Lond. 31, 55–99.
- Graf, A., 2007. Genese alpiner Seitenmoränen am Beispiel des Findelengletschers bei Zermatt (VS).
 Unpublished MSc Thesis, Universität Bern, 126 pp.
- 2044 Greenwood, S.L., Clark, C.D., 2008. Subglacial bedforms of the Irish Ice Sheet. J. Maps 4, 332–357.
- Greenwood, S.L., Clark, C.D., 2009a. Reconstructing the last Irish Ice Sheet 1: changing flow
 geometries and ice flow dynamics deciphered from the glacial landform record. Quat. Sci.
 Rev. 28, 3085–3100.
- Greenwood, S.L., Clark, C.D., 2009b. Reconstructing the last Irish Ice Sheet 2: a geomorphologically driven model of ice sheet growth, retreat and dynamics. Quat. Sci. Rev. 28(27), 3101–3123.
- Greenwood, S.L., Kleman, J., 2010. Glacial landforms of extreme size in the Keewatin sector of the
 Laurentide Ice Sheet. Quat. Sci. Rev. 29(15), 1894–1910.
- Greenwood, S.L., Clark, C.D., Hughes, A.L.C., 2007. Formalising an inversion methodology for
 reconstructing ice-sheet retreat patterns from meltwater channels: application to the British
 Ice Sheet. J. Quat. Sci. 22, 637–645.

- Greenwood, S.L., Clason, C.C., Mikko, H., Nyberg, J., Peterson, G., Smith, C.A., 2015. Integrated
 use of lidar and multibeam bathymetry reveals onset of ice streaming in the northern Bothnian
 Sea. GFF 137, 284–292
- 2058Greenwood, S.L., Clason, C.C., Nyberg, J., Jakobsson, M., Holmlund, P., 2017. The Bothnian Sea ice2059stream: early Holocene retreat dynamics of the south-central Fennoscandian Ice Sheet. Boreas206046, 346–362.
- Gribenski, N., Jansson, K.N., Lukas, S., Stroeven, A.P., Harbor, J.M., Blomdin, R., Ivanov, M.N.,
 Heyman, J., Petrakov, D.A., Rudoy, A., Clifton, T., Lifton, N.A., Caffee, M.W., 2016.
 Complex patterns of glacier advances during the late glacial in the Chagan Uzun Valley,
 Russian Altai. Quat. Sci. Rev. 149, 288–305.
- Gribenski, N., Lukas, S., Stroeven, A.P., Jansson, K.N., Harbor, J.M., Blomdin, R., Ivanov, M.N.,
 Heyman, J., Petrakov, D.A., Rudoy, A., Clifton, T., Lifton, N.A., Caffee, M.W., 2017. Reply
 to comment received from J. Herget et al. regarding "Complex patterns of glacier advances
 during the late glacial in the Chagan Uzun Valley, Russian Altai" by Gribenski et al. (2016),
 Quaternary Science Reviews 149, 288–305. Quat. Sci. Rev. 168, 219–221.
- 2070 Griffiths, J.S., Martin, C.J. (Eds.), 2017. Engineering Geology and Geomorphology of Glaciated and
 2071 Periglaciated Terrains: Engineering Group Working Party Report. Engineering Geology
 2072 Special Publication 28. The Geological Society, London, 953 pp.
- 2073 Grodecki, J., Dial, G., 2003. Block Adjustment of High-Resolution Satellite Images Described by
 2074 Rational Polynomials. Photogramm. Eng. Remote Sens. 69(1), 59–68.
- Gustavsson, M., Kolstrup, E., Seijmonsbergen, A.C., 2006. A new symbol-and-GIS based detailed
 geomorphological mapping system: Renewal of a scientific discipline for understanding
 landscape development. Geomorphology 77(1–2), 90–111.
- 2078 Gustavsson, M., Seijmonsbergen, A.C., Kolstrup, E., 2008. Structure and contents of a new geomorphological GIS database linked to a geomorphological map With an example from Liden, central Sweden. Geomorphology 95, 335–349.
- Harbor, J., Stroeven, A.P., Fabel, D., Clarhäll, A., Kleman, J., Li, Y.K., Elmore, D., Fink, D., 2006.
 Cosmogenic nuclide evidence for minimal erosion across two subglacial sliding boundaries of
 the late glacial Fennoscandian ice sheet. Geomorphology 75, 90–99.
- Hardt, J., Hebenstreit, R., Lüthgens, C., Böse, M., 2015. High-resolution mapping of ice-marginal
 landforms in the Barnim region, northeast Germany. Geomorphology 250, 41–52.
- Hättestrand, C., 1998. The glacial geomorphology of central and northern Sweden. Sver. Geol.
 Unders. Ca 85, 1–47.
- Hättestrand, C., Clark, C.D., 2006. The glacial geomorphology of Kola Peninsula and adjacent areas
 in the Murmansk Region, Russia. J. Maps 2, 30–42.
- Hättestrand, C., Stroeven, A.P., 2002. A relict landscape in the centre of Fennoscandian glaciation:
 Geomorphological evidence of minimal Quaternary glacial erosion. Geomorphology 44(1),
 127–143.
- Hättestrand, C., Kosche, S., Näslund, J.-O., Fabel, D., Stroeven, A.P., 2004. Drumlin formation time evidence from northern and central Sweden. Geogr. Ann. 86A, 155–167.
- Heiser, P.A., Roush, J.J., 2001. Pleistocene glaciations in Chukotka, Russia: moraine mapping using satellite synthetic aperture radar (SAR) imagery. Quat. Sci. Rev. 20(1), 393–404.
- Hess, D.P., Briner, J.P., 2009. Geospatial analysis of controls on subglacial bedform morphometry in the New York Drumlin Field – implications for Laurentide Ice Sheet dynamics. Earth Surf.
 Proc. Land. 34. 1126–1135.
- Heyman, J., Hättestrand, C., Stroeven, A.P., 2008. Glacial geomorphology of the Bayan Har sector of
 the NE Tibetan Plateau. J. Maps 4(1), 42–62.
- Hillier, J.K., Smith, M.J., Clark, C.D., Stokes, C.R., Spagnolo, M., 2013. Subglacial bedforms reveal an exponential size–frequency distribution. Geomorphology 190, 82–91.
- Hillier, J.K., Smith, M.J., Armugam, R., Barr, I., Boston, C.M., Clark, C.D., Ely, J., Frankl, A.,
 Greenwood, S.L., Gosselin, L., Hättestrand, C., Hogan, K., Hughes, A.L.C., Livingstone, S.J.,
 Lovell, H., McHenry, M., Munoz, Y., Pellicer, X.M., Pellitero, R., Robb, C., Roberson, S.,
 Ruther, D., Spagnolo, M., Standell, M., Stokes, C.R., Storrar, R., Tate, N.J., Wooldridge, K.,
 Maps 11, 719–729.

- Hillier, J.K., Kougioumtzoglou, I.A., Stokes, C.R., Smith, M.J., Clark, C.D., Spagnolo, M.S., 2016.
 Exploring Explanations of Subglacial Bedform Sizes Using Statistical Models. PLoS One 11, e0159489.
- Hochreuther, P., Loibl, D., Wernicke, J., Zhu, H., Grießinger, J., Bräuning, A., 2015. Ages of major
 Little Ice Age glacier fluctuations on the southeast Tibetan Plateau derived from tree-ringbased moraine dating. Palaeogeogr., Palaeoclimatol., Palaeoecol. 422, 1–10.
- Hodgson, D.A., Vincent, J.S., Fyles, J.G., 1984. Quaternary geology of central Melville Island,
 Northwest Territories. Geological Survey of Canada paper 83-16.
- Hodgson, D.A., Graham, A.G.C., Griffiths, H.J., Roberts, S.J., Ó Cofaigh, C., Bentley, M.J., Evans,
 D.J.A., 2014. Glacial history of sub-Antarctic South Georgia based on the submarine
 geomorphology of its fjords. Quat. Sci. Rev. 89, 129–147.
- Hollingworth, S.E., 1931. The glaciation of western Edenside and adjoining areas and the drumlins of
 Edenside and the Solway Basin. Quart. J. Geol. Soc. Lond. 87, 281–359.
- Horsfield, B.R., 1983. The deglaciation pattern of the western Grampians of Scotland. Unpublished
 PhD thesis, University of East Anglia.
- Houmark-Nielsen, M., Kjær, K.H. 2003. Southwest Scandinavia, 40-15 kyr BP: Palaeogeography and
 environmental change. J. Quat. Sci. 18, 769–786.
- Howarth, P.J., 1968. Geomorphological and Glaciological Studies, Eastern Breiðamerkurjökull,
 Iceland, Unpublished PhD thesis, University of Glasgow.
- Howarth, P.J., Welch, R., 1969a. Breiðamerkurjökull, South-east Iceland, August 1945, 1:30,000
 scale map. University of Glasgow.
- Howarth, P.J., Welch, R., 1969b. Breiðamerkurjökull, South-east Iceland, August 1965, 1:30,000
 scale map. University of Glasgow.
- Hubbard, A., Bradwell, T., Golledge, N., Hall, A., Patton, H., Sugden, D., Cooper, R., Stoker, M.,
 2009. Dynamic cycles, ice streams and their impact on the extent, chronology and
 deglaciation of the British–Irish ice sheet. Quat. Sci. Rev. 28, 758–776.
- Hubbard, B., Glasser, N.F., 2005. Field techniques in glaciology and glacial geomorphology. John
 Wiley and Sons, Chichester, 400 pp.
- Hughes, A.L.C., Clark, C.D., Jordan, C.J., 2010. Subglacial bedforms of the last British Ice Sheet. J.
 Maps 6, 543–563.
- Hughes, A.L.C., Clark, C.D., Jordan, C.J., 2014. Flow-pattern evolution of the last British Ice Sheet.
 Quat. Sci. Rev. 89, 148–168.
- Hughes, A.L.C., Gyllencreutz, R., Lohne, Ø.S., Mangerud, J., Svendsen, J.I., 2016. The last Eurasian
 ice sheets a chronological database and time-slice reconstruction, DATED-1. Boreas 45, 145.
- Hyatt, O.M., 2010. Insights into New Zealand Glacial Processes from studies of glacial geomorphology and sedimentology in Rakaia and other South Island Valleys. Unpublished PhD thesis, University of Canterbury, 251 pp.
- Immerzeel, W.W., Kraaijenbrink, P.D.A., Shea, J.M., Shrestha, A.B., Pellicciotti, F., Bierkens,
 M.F.P., de Jong, S.M., 2014. High-resolution monitoring of Himalayan glacier dynamics
 using unmanned aerial vehicles. Remote Sens. Environ. 150, 93–103.
- Izagirre, E., Darvill, C.M., Rada, C., Aravena, J.C., 2018. Glacial geomorphology of the Marinelli and
 Pigafetta glaciers, Cordillera Darwin Icefield, southernmost Chile. J. Maps 14, 269–281.
- Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J.A., Forbes, S., Fridman, B., Hodnesdal, H.,
 Noormets, R., Pedersen, R., Rebesco, M., Schenke, H.W., Zarayskaya, Y., Accettella, D.,
 Armstrong, A., Anderson, R.M., Bienhoff, P., Camerlenghi, A., Church, I., Edwards, M.,
 Gardner, J.V., Hall, J.K., Hell, B., Hestvik, O., Kristoffersen, Y., Marcussen, C., Mohammad,
 R., Mosher, D., Nghiem, S.V., Pedrosa, M.T., Travaglini, P.G., Weatherall, P., 2012. The
 International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0. Geophys. Res.
 Lett. 39, L12609.
- James, M.R., Robson, S., Smith, M.W., 2017. 3-D uncertainty-based topographic change detection
 with structure-from-motion photogrammetry: precision maps for ground control and directly
 georeferenced surveys. Earth Surf. Process. Land. 42, 1769–1788.
- Jamieson, S.S.R., Ewertowski, M.W., Evans, D.J.A., 2015. Rapid advance of two mountain glaciers
 in response to mine-related debris loading, J. Geophys. Res.: Earth Surf. 120, 1418–1435.

- 2165Jansson, K.N., Glasser, N.F., 2005. Using Landsat 7 ETM+ imagery and Digital Terrain Models for2166mapping glacial lineaments on former ice sheet beds. Int. J. Remote Sens. 26(18), 3931–3941.
- Jansson, K.N., Kleman, J., Marchant, D.R., 2002. The succession of ice-flow patterns in north-central
 Québec-Labrador, Canada. Quat. Sci. Rev. 21(4), 503–523.
- Jansson, K.N., Stroeven, A.P., Kleman, J., 2003. Configuration and timing of Ungava Bay ice
 streams, Labrador–Ungava, Canada. Boreas 32(1), 256–262.
- 2171Johnson, M.D., Fredin, O., Ojala, A.E.K., Peterson, G. 2015. Unravelling Scandinavian2172geomorphology: the LiDAR revolution. GFF 137, 245–251.
- Jónsson, S.A., Schomacker, A., Benediktsson, Í.Ö., Ingólfsson, Ó., Johnson, M.D., 2014. The drumlin
 field and the geomorphology of the Múlajökull surge-type glacier, central Iceland.
 Geomorphology 207, 213–220.
- Jónsson, S.A., Benediktsson, Í.Ö., Ingólfsson, Ó., Schomacker, A., Bergsdóttir, H.L., Jacobsson,
 W.R., Linderson, H. 2016. Submarginal drumlin formation and late Holocene history of
 Fláajökull, southeast Iceland. Ann. Glaciol. 57, 128–141.
- Jorge, M.G., Brennand, T.A., 2017a. Measuring (subglacial) bedform orientation, length, and
 longitudinal asymmetry–Method assessment. PloS one, 12(3), e0174312.
- Jorge, M.G., Brennand, T.A., 2017b. Semi-automated extraction of longitudinal subglacial bedforms
 from digital terrain models–Two new methods. Geomorphology, 288, 148–163.
- Juyal, N., Thakkar, P.S., Sundriyal, Y.P., 2011. Geomorphic evidence of glaciations around Mount
 Kailash (Inner Kora): implication to past climate. Current Science 100(4), 535-541.
- Kassab, C., Wang, J., Harbor, J., 2013. Glacial geomorphology of the Dalijia Shan region, northeastern Tibetan Plateau. J. Maps 9(1), 98–105.
- Kelley, S.E., Kaplan, M.R., Schaefer, J.M., Andersen, B.G., Barrell, D.J.A., Putnam, A.E., Denton,
 G.H., Schwartz, R., Finkel, R.C., Doughty, A.M., 2014. High-precision 10Be chronology of
 moraines in the Southern Alps indicates synchronous cooling in Antarctica and New Zealand
 42,000 years ago. Earth Planet. Sci. Lett. 405, 194–206.
- Kendall, P.F. 1902. A system of glacier lakes in the Cleveland Hills. Quart. J. Geol. Soc. Lond. 58, 471–571.
- Kerschner, H., Kaser, G., Sailer, R., 2000. Alpine Younger Dryas glaciers as palaeo-precipitation
 gauges. Ann. Glaciol. 31, 80–84.
- Kienholz, H., 1977. Kombinierte geomorphologische Gefahrenkarte 1:10,000 von Grindelwald.
 Geographica Bernensia, G4: 1–204.
- King, E.C., Woodward, J., Smith, A. M., 2007. Seismic and radar observations of subglacial bed
 forms beneath the onset zone of Rutford Ice Stream, Antarctica. J. Glaciol. 53(183), 665–672.
- King, E.C., Hindmarsh, R.C., Stokes, C.R., 2009. Formation of mega-scale glacial lineations observed
 beneath a West Antarctic ice stream. Nat. Geosci. 2(8), 585–588.
- King, E.C., Pritchard, H.D., Smith, A.M., 2016a. Subglacial landforms beneath Rutford Ice Stream,
 Antarctica: detailed bed topography from ice-penetrating radar. Earth Syst. Sci. Data 8(1),
 151–158.
- King, O., Hambrey, M.J., Irvine-Fynn, T.D.L., Holt, T.O., 2016b. The structural, geometric and volumetric changes of a polythermal Arctic glacier during a surge cycle: Comfortlessbreen, Svalbard. Earth Surf. Process. Land. 41, 162–177.
- King, O., Quincey, D.J., Carrivick, J.L., Rowan, A.V., 2017. Spatial variability in mass loss of glaciers in the Everest region, central Himalayas, between 2000 and 2015. The Cryosphere 11, 407–426.
- Kirchner, N., Greve, R., Stroeven, A.P., Heyman, J., 2011. Paleoglaciological reconstructions for the
 Tibetan Plateau during the last glacial cycle: evaluating numerical ice sheet simulations
 driven by GCM-ensembles. Quat. Sci. Rev. 30, 248–267.
- Kirkbride, M.P., Winkler, S., 2012. Correlation of Late Quaternary moraines: impact of climate
 variability, glacier response, and chronological resolution. Quat. Sci. Rev. 46, 1–29.
- Kjær, K.H., Krüger, J., 2001. The final phase of dead-ice moraine development: processes and sediment architecture, Kotlujokull, Iceland. Sedimentology 48, 935–952.
- Kjær, K.H., Houmark-Nielsen, M., Richardt, N. 2003. Ice-flow patterns and dispersal of erratics at the
 southwestern margin of the last Scandinavian Ice Sheet: signature of paleo-ice streams.
 Boreas 32, 130-148.

- Kjær, K.H., Korsgaard, N.J., Schomacker, A., 2008. Impact of multiple glacier surges a
 geomorphological map from Brúarjökull, East Iceland. J. Maps 4, 5–20.
- Kłapyta, P., 2013. Application of Schmidt hammer relative age dating to Late Pleistocene moraines
 and rock glaciers in the Western Tatra Mountains, Slovakia. Catena 111, 104–121.
- Klassen, R.A., 1993. Quaternary geology and glacial history of Bylot Island, Northwest Territories.
 Geological Survey of Canada Memoir 429.
- Kleman, J., 1990. On the use of glacial striae for reconstruction of palaeo-ice sheet flow patterns.
 Geogr. Ann. 72A, 217–236.
- Kleman, J., 1992. The Palimpsest Glacial Landscape in Northwestern Sweden. Late Weichselian
 Deglaciation Landforms and Traces of Older West-Centered Ice Sheets. Geogr. Ann. 74A(4),
 305–325.
- Kleman, J., Borgström, I., 1996. Reconstruction of Palaeo-Ice Sheets: The Use of Geomorphological
 Data. Earth Surf. Process. Land. 21, 893–909.
- Kleman, J. Stroeven, A.P., 1997. Preglacial surface remnants and Quaternary glacial regimes in northwestern Sweden. Geomorphology 19(1–2), 35–54.
- Kleman, J., Hättestrand, C., Borgström, I., Stroeven, A., 1997. Fennoscandian palaeoglaciology
 reconstructed using a glacial geological inversion model. J. Glaciol. 43(144), 283–299.
- Kleman, J., Fastook, J., Stroeven, A.P., 2002. Geologically and geomorphologically constrained numerical model of Laurentide Ice Sheet inception and build-up. Quat. Int. 95-96, 87–98.
- Kleman, J., Hättestrand, C., Stroeven, A.P., Jansson, K.N., De Angelis, H., Borgström, I., 2006.
 Reconstruction of palaeo-ice sheets; inversion of their glacial geomorphological record. In:
 Knight, P.G. (Ed.), Glacier Science and Environmental Change. Blackwell, Oxford, pp. 192–199.
- Kleman, J., Stroeven, A.P., Lundqvist, J., 2008. Patterns of Quaternary ice sheet erosion and deposition in Fennoscandia. Geomorphology 97, 73–90.
- Kleman, J., Jansson, K., De Angelis, H., Stroeven, A.P., Hättestrand, C., Alm, G., Glasser, N., 2010.
 North American Ice Sheet build-up during the last glacial cycle, 115–21kyr. Quat. Sci. Rev.
 2247 29, 2036–2051.
- 2248 Klimaszewski, M., 1990. Thirty years of detailed geomorphological mapping. Geogr. Pol. 58, 11–18.
- Kneisel, C., Lehmkuhl, F., Winkler, S., Tressel, E., Schröder, H., 1998. Legende für geomorphologische Kartierungen in Hochgebirgen (GMK Hochgebirge). Trier. Geogr. Stud.
 18, 1–24.
- Knight, J., Mitchell, W., Rose, J., 2011. Geomorphological Field Mapping. In: Smith, M.J., Paron, P.,
 Griffiths, J. (Eds.), Geomorphological Mapping: Methods and Applications. Developments in
 Earth Surface Processes 15. Elsevier, London, pp. 151–188.
- Kocurek, G., Ewing, R.C., Mohrig, D., 2010. How do bedform patterns arise? New views on the role
 of bedform interactions within a set of boundary conditions. Earth Surf. Process. Land. 35,
 51–63.
- Korsgaard, N.J., Schomacker, A., Benediktsson, Í.Ö., Larsen, N.K., Ingólfsson, Ó., Kjær, K.H., 2015.
 Spatial distribution of erosion and deposition during a glacier surge: Brúarjökull, Iceland.
 Geomorphology 250, 258–270.
- Kraak, M.–J., Oremling, F.J., 2010. Cartography: visualization of spatial data (3rd Edition). Routledge,
 London.
- Kronberg, P., 1984. Photogeologie. Eine Einfhrung in die Grundlagen und Methoden der
 geologischen Auswertung von Luftbildern. Enke, Stuttgart.
- Krüger, J., 1994. Glacial processes, sediments, landforms and stratigraphy in the terminus region of
 Mýrdalsjökull, Iceland. Folia Geographica Danica 21, 1–233.
- Krüger, J., 1995. Origin, chronology and climatological significance of annual moraine ridges at Mýrdalsjökull, Iceland. The Holocene 5, 420–427.
- Krüger, J., Kjær, K.H., 2000. De-icing progression of ice-cored moraines in a humid, subpolar climate, Kötlujökull, Iceland. The Holocene 10, 737–747.
- Krygowski B. (Ed.), 1963. Mapa Geomorfologiczna Niziny Wielkopolsko-Kujawskiej
 [Geomorphological Map of Wielkopolska-Kujawy lowland]. 1:300 000 Map. Adam
 Mickiewicz University.

- Kuhle, M., 1990. Quantificational reductionism as a risk in geography instanced by the 1:25,000
 Geomorphological Map of the Federal Republic of Germany. Geogr. Pol. 58, 41–54.
- Lam, N.S.N., Quattrochi, D.A., 1992. On the issues of scale, resolution, and fractal analysis in the mapping sciences. The Prof. Geogr. 44(1), 88–98.
- Lane, S.N., Reid, S.C., Westaway, R.M., Hicks, D.M., 2005. Remotely Sensed Topographic Data for River Channel Research: The Identification, Explanation and Management of Error. In: R.E.J.
 Kelly, Drake, N.A., Barr, S.L. (Eds.), Spatial Modelling of the Terrestrial Environment. John Wiley & Sons, Ltd, Chichester, UK, pp. 113–136.
- Lardeux, P., Glasser, N., Holt, T., Hubbard, B., 2015. Glaciological and geomorphological map of
 Glacier Noir and Glacier Blanc, French Alps. J. Maps 12, 582–596.
- Leser, H., 1983. Anwendung und Auswertung geomorphologischer Kartierungen und Karten. Mater.
 Physiogeogr. 5, 5–13.
- Leser, H., Stäblein, G., 1975. Geomorphologische Kartierung Richtlinien zur Herstellung
 geomorphologischer Karten 1: 25,000. Berliner Geographische Abhandlungen Sonderheft, 1–
 33.
- Levy, L.B., Larsen, N.K., Davidson, T.A., Strunk, A., Olsen, J., Jeppesen, E., 2017. Contrasting evidence of Holocene ice margin retreat, south-western Greenland. J. Quat. Sci. 32, 604–616.
- Li, Y., Li, Y., Chen, Y., Lu, X., 2016. Presumed Little Ice Age glacial extent in the eastern Tian Shan, China. J. Maps 12(1), 71–78.
- Lidmar-Bergström, K., Elvhage, C., Ringberg, B., 1991. Landforms in Skane, south Sweden.
 Geografiska Annaler 73A, 61–91.
- Lifton, N., Beel, C., Hättestrand, C., Kassab, C., Rogozhina, I., Heermance, R., Oskin, M., Burbank,
 D., Blomdin, R., Gribenski, N., Caffee, M., 2014. Constraints on the late Quaternary glacial
 history of the Inylchek and Sary-Dzaz valleys from in situ cosmogenic 10Be and 26Al,
 eastern Kyrgyz Tian Shan. Quat. Sci. Rev. 101, 77–90.
- Lillesand, T.M., Kiefer, R.W., Chipman, J.W., 2015. Remote Sensing and Image Interpretation (7th
 Edition). John Wiley & Sons, Hoboken, USA, 768 pp.
- Lindholm, M.S., Heyman, J., 2016. Glacial geomorphology of the Maidika region, Tibetan Plateau. J.
 Maps 12(5), 797–803.
- Livingstone, S.J., Ó Cofaigh, C., Evans, D.J.A., 2008. The glacial geomorphology of the central sector of the British-Irish Ice Sheet. J. Maps 4, 358–377.
- Livingstone, S.J., Evans, D.J.A., Ó Cofaigh, C., 2010. Re-advance of Scottish ice into the Solway
 Lowlands (Cumbria, UK) during the Main Late Devensian deglaciation. Quat. Sci. Rev. 29,
 2544-2570.
- Livingstone, S.J., Ó Cofaigh, C., Stokes, C.R., Hillenbrand, C.-D., Vieli, A., Jamieson, S.S.R., 2012.
 Antarctic palaeo-ice streams. Earth-Sci. Rev. 111, 90–128.
- Livingstone, S.J., Storrar, R.D., Hillier, J.K., Stokes, C.R., Clark, C.D., Tarasov, L., 2015. An ice sheet scale comparison of eskers with modelled subglacial drainage routes. Geomorphology
 246, 104–112.
- Loibl, D., Hochreuther, P., Schulte, P., Hülle, D., Zhu H, Bräuning, A., Lehmkuhl, F., 2015. Toward a
 late Holocene glacial chronology for the eastern Nyainqêntanglha Range, southeastern Tibet.
 Quat. Sci. Rev. 107, 243–259.
- Lønne, I., 2016. A new concept for glacial geological investigations of surges, based on High-Arctic
 examples (Svalbard). Quat. Sci. Rev., 132, 74-100.
- Lovell, H., 2014. On the ice-sediment-landform associations of surging glaciers on Svalbard.
 Unpublished PhD thesis, Queen Mary University of London, 312 pp.
- Lovell, H., Boston, C.M., 2017. Glacitectonic composite ridge systems and surge-type glaciers: an
 updated correlation based on Svalbard, Norway. arktos 3, 2.
- Lovell, H., Stokes, C.R., Bentley, M.J., 2011. A glacial geomorphological map of the Seno Skyring Seno Otway-Strait of Magellan region, southernmost Patagonia. J. Maps 7, 318–339.
- Lovell, H., Stokes, C.R., Bentley, M.J., Benn, D.I., 2012. Evidence for rapid ice flow and proglacial
 lake evolution around the central Strait of Magellan region, southernmost Patagonia. J. Quat.
 Sci. 27, 625–638.

- Lovell, H., Benn, D.I., Lukas, S., Spagnolo, M., Cook, S.J., Swift, D.A., Clark, C.D., Yde, J.C.,
 Watts, T.P., 2018. Geomorphological investigation of multiphase glacitectonic composite
 ridge systems in Svalbard. Geomorphology 300, 176–188.
- Lueder, D.R., 1959. Aerial Photographic Interpretation Principles and Applications. McGraw-Hill,
 New York.
- Lukas, S., 2002. Geomorphological evidence for the pattern of deglaciation around the Drumochter
 Pass, Central Grampian Highlands, Scotland. Unpublished MSc thesis, Ruhr-University of
 Bochum, Germany, 115 pp.
- Lukas, S., 2005. A test of the englacial thrusting hypothesis of 'hummocky' moraine formation case
 studies from the north-west Highlands, Scotland. Boreas 34, 287–307.
- Lukas, S., 2006. Morphostratigraphic principles in glacier reconstruction -a perspective from the
 British Younger Dryas. Prog. Phys. Geogr. 30, 719–736.
- Lukas, S., 2007a. Early-Holocene glacier fluctuations in Krundalen, south central Norway: palaeo glacier dynamics and palaeoclimate. The Holocene 17, 585–598.
- Lukas, S., 2007b. 'A test of the englacial thrusting hypothesis of "hummocky" moraine formation:
 case studies from the northwest Highlands, Scotland': Reply to comments. Boreas 36, 108–
 113.
- Lukas, S., 2011. Ice-cored moraines. In: Singh, V., Singh, P. Haritashya, U.K. (Eds.), Encyclopedia of
 Snow, Ice and Glaciers. Springer, Heidelberg, pp. 616–619.
- Lukas, S., 2012. Processes of annual moraine formation at a temperate alpine valley glacier: insights
 into glacier dynamics and climatic controls. Boreas 41(3), 463–480.
- Lukas, S., Benn, D.I., 2006. Retreat dynamics of Younger Dryas glacier in the far NW Scottish
 Highlands reconstructed from moraine sequences. Scott. Geogr. J. 122, 308–325.
- Lukas, S., Lukas, T., 2006. A glacial geological and geomorphological map of the far NW Highlands,
 Scotland. Parts 1 and 2. J. Maps 2, 43–56, 56–58.
- Lukas, S., Sass, O., 2011. The formation of Alpine lateral moraines inferred from sedimentology and radar reflection patterns: a case study from Gornergletscher, Switzerland. Geological Society, London, Special Publications 354, 77–92.
- Lukas, S., Nicholson, L.I., Ross, F.H., Humlum, O., 2005. Formation, meltout processes and landscape alteration of High-Arctic ice-cored moraines—examples from Nordenskiöld Land, Central Spitsbergen. Polar Geogr. 29, 157–187.
- Lukas, S., Nicholson, L.I., Humlum, O., 2007. Comment on Lønne and Lyså (2005): "Deglaciation dynamics following the Little Ice Age on Svalbard: Implications for shaping of landscapes at high latitudes", Geomorphology 72, 300–319. Geomorphology 84, 145–149.
- Lukas, S., Benn, D.I., Boston, C.M., Brook, M.S., Coray, S., Evans, D.J.A., Graf, A., KellererPirklbauer-Eulenstein, A., Kirkbride, M.P., Krabbendam, M., Lovell, H., Machiedo, M.,
 Mills, S.C., Nye, K., Reinardy, B.T.I., Ross, F.H., Signer, M., 2013. Clast shape analysis and
 clast transport paths in glacial environments: A critical review of methods and the role of
 lithology. Earth-Sci. Rev. 121, 96–116.
- Lukas, S., Preusser, F., Evans, D.J.A., Boston, C.M., Lovell, H., 2017. Chapter 2 The Quaternary.
 In: Griffiths, J.S., Martin, C.J. (Eds.), Engineering Geology and Geomorphology of Glaciated and Periglaciated Terrains: Engineering Group Working Party Report. Engineering Geology Special Publication 28. The Geological Society, London, pp. 31–57.
- Maclachlan, J.C., Eyles, C.H., 2013. Quantitative geomorphological analysis of drumlins in the
 Peterborough drumlin field, Ontario, Canada. Geogr. Ann. 95A, 125–144.
- Mäkinen, J., Kajuutti, K., Palmu, J., Ojala, A., Ahokangas, E., 2017. Triangular-shaped landforms
 reveal subglacial drainage routes in SW Finland. Quat. Sci. Rev. 164, 37–53.
- Małecki, J., Lovell, H., Ewertowski, W., Górski, Ł., Kurczba, T., Latos, B., Miara, M., Piniarska, D.,
 Płocieniczak, J., Sowada, T., Spiralski, M., Warczachowska, A., Rabatel, A., 2018. The
 glacial landsystem of a tropical glacier: Charquini Sur, Bolivian Andes. Earth Surf. Proc.
 Land., in press. doi: 10.1002/esp.4417
- Marc, O., Hovius, N., 2015. Amalgamation in landslide maps: effects and automatic detection. Nat.
 Hazards Earth Syst. Sci. 15, 723–733.
- Margold, M., Jansson, K.N., 2011. Glacial geomorphology and glacial lakes of central Transbaikalia,
 Siberia, Russia. J. Maps 7, 18–30.

- Margold, M., Jansson, K.N., 2012. Evaluation of data sources for mapping glacial meltwater features.
 Int. J. Remote Sens. 33, 2355–2377.
- Margold, M., Jansson, K.N., Kleman, J., Stroeven, A.P., 2011. Glacial meltwater landforms of central
 British Columbia. J. Maps 7, 486–506.
- Margold, M., Stokes, C.R., Clark, C.D., Kleman, J., 2015a. Ice streams in the Laurentide Ice Sheet: a
 new mapping inventory. J. Maps 11, 380–395.
- Margold, M., Stokes, C.R., Clark, C.D., 2015b. Ice streams in the Laurentide Ice Sheet: Identification,
 characteristics and comparison to modern ice sheets. Earth-Sci. Rev. 143, 117–146.
- Margold, M., Stokes, C.R., Clark, C.D., 2018. Reconciling records of ice streaming and ice margin
 retreat to produce a palaeogeographic reconstruction of the deglaciation of the Laurentide Ice
 Sheet. Quat. Sci. Rev. 189, 1–30.
- Mather, A.E., Mills, S., Stokes, M., Fyfe, R., 2015. Ten years on: what can Google Earth offer the
 Geoscience community? Geology Today 31(6), 216–221.
- May, J.H., Zech, J., Zech, R., Preusser, F., Argollo, J., Kubik, P.W., Veit, H., 2011. Reconstruction of
 a complex late Quaternary glacial landscape in the Cordillera de Cochabamba (Bolivia) based
 on a morphostratigraphic and multiple dating approach. Quat. Res. 76(1), 106–118.
- McDougall, D.A., 2001. The geomorphological impact of Loch Lomond (Younger Dryas) Stadial
 plateau icefields in the central Lake District, northwest England. J. Quat. Sci. 16, 531–543.
- McDougall, D.A., 2013. Glaciation style and the geomorphological record: evidence for Younger
 Dryas glaciers in the eastern Lake District, northwest England. Quat. Sci. Rev. 73, 48–58.
- McHenry, M., Dunlop, P., 2016. The subglacial imprint of the last Newfoundland Ice Sheet, Canada.
 J. Maps 12(3), 462–483.
- Melander, O., 1975. Geomorfologiska kartbladet 29 I Kebnekaise. Statens naturvårdsverk. 78 pp.,
 map scale 1:250,000.
- Mertes, J.R., Gulley, J.D., Benn, D.I., Thompson, S.S., Nicholson, L.I., 2017. Using structure-from motion to create glacier DEMs and orthoimagery from historical terrestrial and oblique aerial
 imagery. Earth Surf. Process. Land. 42, 2350–2364.
- Midgley, N.G., Tonkin, T.N., 2017. Reconstruction of former glacier surface topography from archive
 oblique aerial images. Geomorphology 282, 18–26.
- Midgley, N.G., Cook, S.J., Graham, D.J., Tonkin, T.N., 2013. Origin, evolution and dynamic context
 of a Neoglacial lateral–frontal moraine at Austre Lovénbreen, Svalbard. Geomorphology 198,
 96–106.
- Midgley, N.G., Tonkin, T.N., Graham, D.J., Cook, S.J., 2018. Evolution of high-Arctic glacial
 landforms during deglaciation. Geomorphology 311, 63–75.
- Miller, H., Cotterill, C.J., Bradwell, T., 2014. Glacial and paraglacial history of the Troutbeck Valley,
 Cumbria, UK: integrating airborne LiDAR, multibeam bathymetry, and geological field
 mapping. Proc. Geol. Assoc. 125, 31–40.
- Mills, S.C., Grab, S.W., Rea, B.R., Carr, S.J., Farrow, A., 2012. Shifting westerlies and precipitation
 patterns during the Late Pleistocene in southern Africa determined using glacier
 reconstruction and mass balance modelling. Quaternary Science Reviews 55, 145-159.
- Mitchell, W.A., Riley, J.M., 2006. Drumlin map of the Western Pennines and southern Vale of Eden,
 Northern England, UK. J. Maps 2, 10–16.
- Mollard, J.D., Janes, J.R., 1984. Airphoto Interpretation and the Canadian Landscape. Energy, Mines
 and Resources Canada, Ottawa.
- Möller, P., Dowling, T.P.F., 2016. Streamlined subglacial bedforms on the Närke plain, south-central
 Sweden Areal distribution, morphometrics, internal architecture and formation. Quat. Sci.
 Rev. 146, 182–215.
- Möller, P., Dowling, T.P.F., 2018. Equifinality in glacial geomorphology: instability theory examined
 via ribbed moraine and drumlins in Sweden. GFF, in press.
- Morén, B., Heyman, J., Stroeven, A.P., 2011. Glacial geomorphology of the central Tibetan Plateau.
 J. Maps 7, 115–125.
- Murray, A.B., Goldstein, E.B., Coco, G., 2014. The shape of patterns to come: from initial formation
 to long-term evolution. Earth Surf. Process. Land. 39, 62–70.

- Napieralski, J., Hubbard, A., Li, Y.K., Harbor, J., Stroeven, A.P., Kleman, J., Alm, G., Jansson, K.N.,
 2436 2007a. Towards a GIS assessment of numerical ice sheet model performance using
 2437 geomorphological data. J. Glaciol. 53 (180), 71–83.
- Napieralski, J., Harbor, J., Li, Y., 2007b. Glacial geomorphology and geographic information
 systems. Earth-Sci. Rev. 85, 1–22.
- Näslund, J.O., Rodhe, L., Fastook, J.L., Holmlund, P., 2003. New ways of studying ice sheet flow directions and glacial erosion by computer modelling examples from Fennoscandia. Quat.
 Sci. Rev. 22, 245–258.
- Noh, M.J., Howat, I.M., 2015. Automated stereo-photogrammetric DEM generation at high latitudes:
 Surface Extraction from TIN-Based Search Minimization (SETSM) validation and
 demonstration over glaciated regions. GIScience Remote Sens. 52(2), 198–217.
- Norris, S.L., Margold, M., Froese, D.G., 2017. Glacial landforms of northwest Saskatchewan. J. Maps
 13, 600–607.
- Norris, S.L., Evans, D.J.A., Cofaigh, C.Ó., 2018. Geomorphology and till architecture of terrestrial palaeo-ice streams of the southwest Laurentide Ice Sheet: A borehole stratigraphic approach.
 Quat. Sci. Rev. 186, 186–214.
- Nuth, C., Kääb, A., 2011. Co-registration and bias corrections of satellite elevation data sets for quantifying glacier thickness change. The Cryosphere 5, 271–290.

- Ó Cofaigh, C., 2012. Ice sheets viewed from the ocean: the contribution of marine science to understanding modern and past ice sheets. Phil. Trans. R. Soc. A 370(1980), 5512–5539.
- Ó Cofaigh, C., Evans, D.J.A., Smith, I.R., 2010. Large-scale reorganization and sedimentation of terrestrial ice streams during late Wisconsinan Laurentide Ice Sheet deglaciation. Geol. Soc.
 Am. Bull. 122, 743–756.
- Ó Cofaigh, C., Dowdeswell, J.A., Jennings, A.E., Hogan, K.A., Kilfeather, A., Hiemstra, J.F., Noormets, R.M., Evans, J., McCarthy, D.J., Andrews, J.T., Lloyd, J.M., Moros, M., 2013. An extensive and dynamic ice sheet on the West Greenland shelf during the last glacial cycle. Geology 41 (2), 219–222.
- Ojala, A.E.K., 2016. Appearance of De Geer moraines in southern and western Finland —
 Implications for reconstructing glacier retreat dynamics. Geomorphology 255, 16–25.
- Ojala, A.E.K., Putkinen, N., Palmu, J.P., Nenonen, K., 2015. Characterization of De Geer moraines in
 Finland based on LiDAR DEM mapping. GFF 137(4), 304–318.
- Orkhonselenge, A., 2016. Glacial Geomorphology of Mt. Munkh Saridag in the Khuvsgul Mountain
 Range, Northern Mongolia. Géomorphologie: relief, processus, environnement 22(4), 389 398.
- Ottesen, D., Dowdeswell, J.A., 2006. Assemblages of submarine landforms produced by tidewater
 glaciers in Svalbard. J. Geophys. Res. 111, F01016.
- Ottesen, D., Dowdeswell, J.A., Rise, L., 2005. Submarine landforms and the reconstruction of fast-flowing ice streams within a large Quaternary ice sheet: the 2500 km long Norwegian-Svalbard margin (57 degrees 80 degrees N). Geol. Soc. Am. Bull. 117 (7–8), 1033–1050.
- Ottesen, D., Dowdeswell, J.A., 2006. Assemblages of submarine landforms produced by tidewater
 glaciers in Svalbard. J. Geophys. Res. 111, F01016.
- Ottesen, D., Stokes, C.R., Rise, L., Olsen, L., 2008a. Ice-sheet dynamics and ice streaming along the
 coastal parts of northern Norway. Quat. Sci. Rev. 27, 922–940
- Ottesen, D., Dowdeswell, J.A., Benn, D.I., Kristensen, L., Christiansen, H.H., Christensen, O.,
 Hansen, L., Lebesbye, E., Forwick, M., Vorren, T.O., 2008b. Submarine landforms
 characteristic of glacier surges in two Spitsbergen fjords. Quat. Sci. Rev. 27, 1583–1599.
- Ottesen, D., Stokes, C.R., Bøe, R., Rise, L., Longva, O., Thorsnes, T., Olesen, O., Bugge, T., Lepland,
 A., Hestvik, O.B., 2016. Landform assemblages and sedimentary processes along the
 Norwegian Channel Ice Stream. Sediment. Geol. 338, 115–137.
- 2485Ottesen, D., Dowdeswell, J.A., Bellec, V.K., Bjarnadóttir, L.R., 2017. The geomorphic imprint of2486glacier surges into open-marine waters: Examples from eastern Svalbard. Mar. Geol. 392, 1–248729.

- Otto, J-C., Smith, M.J., 2013. Section 2.6: Geomorphological mapping. In: Clarke, L. (Ed.),
 Geomorphological Techniques (Online Edition). British Society for Geomorphology, London.
 ISSN: 2047-0371.
- Owen, L.A., Finkel, R.C., Barnard, P.L., Haizhou, M., Asahi, K., Caffee, M.W., Derbyshire, E., 2005.
 Climatic and topographic controls on the style and timing of Late Quaternary glaciation throughout Tibet and the Himalaya defined by 10Be cosmogenic radionuclide surface exposure dating. Quat. Sci. Rev. 24(12), 1391–1411.
- Paron, P., Claessens, L., 2011. Makers and Users of Geomorphological Maps. In: Smith, M.J., Paron,
 P., Griffiths, J.S. (Eds.), Geomorphological Mapping: Methods and applications.
 Developments in Earth Surface Processes, Volume 15. Elsevier, Oxford; pp. 75–106.
- Partsch, J., 1894. Die Vergletscherung des Riesengebirges zur Eiszeit. Forschungen zur Deutschen
 Landes- und Volkskunde VIII/2, 103–194.
- Patton, H., Hubbard, A., Andreassen, K., Auriac, A., Whitehouse, P.L., Stroeven, A.P., Shackleton,
 C., Winsborrow, M., Heyman, J., Hall, A.M., 2017a. Deglaciation of the Eurasian ice sheet
 complex. Quat. Sci. Rev. 169, 148–172.
- Patton, H., Hubbard, A., Bradwell, T., Schomacker, A., 2017b. The configuration, sensitivity and rapid retreat of the Late Weichselian Icelandic ice sheet. Earth-Sci. Rev. 166, 223–245.
- Pearce, D., Rea, B.R., Bradwell, T., McDougall, D., 2014. Glacial geomorphology of the Tweedsmuir
 Hills, Central Southern Uplands, Scotland. J. Maps 10(3), 457–465.
- Pearce, D.M., Mair, D.W.F., Rea, B.R., Lea, J.M., Schofield, J.E., Kamenos, N., Schoenrock, K.,
 2018. The glacial geomorphology of upper Godthåbsfjord (Nuup Kangerlua) in southwest
 Greenland. J. Maps 14, 45–55.
- 2510 Penck, A., Brückner, E., 1901/1909. Die Alpen im Eiszeitalter. Tauchnitz, Leipzig.
- Peterson, G., Johnson, M.D., Smith, C.A., 2017. Glacial geomorphology of the south Swedish uplands
 focus on the spatial distribution of hummock tracts. J. Maps 13, 534–544.
- Petrie, G., Price, R.J., 1966. Photogrammetric measurements of the ice wastage and morphological changes near the Casement Glacier, Alaska. Can. J. Earth Sci. 3, 827–840.
- Phillips, E., Evans, D.J.A., Atkinson, N., Kendall, A., 2017. Structural architecture and glacitectonic
 evolution of the Mud Buttes cupola hill complex, southern Alberta, Canada. Quat. Sci. Rev.
 164, 110-139.
- Pike, R.J., 1988. The geometric signature: quantifying landslide-terrain types from digital elevation models. Math. Geol. 20, 491–511.
- Pipaud, I., Loibl, D., Lehmkuhl, F., 2015. Evaluation of TanDEM-X elevation data for geomorphological mapping and interpretation in high mountain environments A case study from SE Tibet, China. Geomorphology 246, 232–254.
- 2523 Popper, K.R., 1972. Objective Knowledge. Oxford University Press, Oxford.
- 2524 Prest, V.K., 1983. Canada's Heritage of Glacial Features. GSC Miscellaneous Report 28.
- Prest, V.K., Grant, D.R., Rampton, V.N. 1968. Glacial map of Canada. Geological Survey of Canada,
 Map 1253A.
- Priamonosov, A.P., Kuznetsova, E.J., Abaturova, I.V., 2000. National geological map of the Russian
 Federation, Map of Quaternary Formations, Polar-Ural series, map sheet Q-41-XII Kharp.
 Ministry of Natural Resources of the Russian Federation. 1:200,000.
- Price, R.J., 1961. The Deglaciation of the Upper Tweed Basin. Unpublished PhD thesis, University of
 Edinburgh.
- Price, R.J., 1963. A glacial meltwater drainage system in Peebleshire, Scotland. Scott. Geogr. Mag.
 79, 133–141.
- 2534 Price, R.J., 1966. Eskers near the Casement Glacier, Alaska. Geogr. Ann. 48, 111–125.
- 2535 Price, R.J., 1970. Moraines at Fjallsjökull, Iceland. Arctic Alp. Res. 2, 27–42.
- Principato, S.M., Moyer, A.N., Hampsch, A.G., Ipsen, H.A., 2016. Using GIS and streamlined
 landforms to interpret palaeo-ice flow in northern Iceland. Boreas 45, 470–482.
- Punkari, M., 1980. The ice lobes of the Scandinavian ice sheet during the deglaciation in Finland.
 Boreas 9(4), 307–310.
- Punkari, M., 1995. Glacial flow systems in the zone of confluence between the Scandinavian and
 Novaja Zemlya Ice Sheets. Quat. Sci. Rev. 14(6), 589–603.

- Putniņš, A., Henriksen, M., 2017. Reconstructing the flow pattern evolution in inner region of the
 Fennoscandian Ice Sheet by glacial landforms from Gausdal Vestfjell area, south-central
 Norway. Quat. Sci. Rev. 163, 56–71.
- Rabus, B., Eineder, M., Roth, A., Bamler, R., 2003. The shuttle radar topography mission—a new class of digital elevation models acquired by spaceborne radar. ISPRS J. Photogramm.
 Remote Sens. 57(4), 241-262.
- Rączkowska Z., Zwoliński Z., 2015. Digital geomorphological map of Poland. Geogr. Pol. 88(2),
 205–210.
- Raistrick, A., 1933. The glacial and post-glacial periods in West Yorkshire. Proc. Geol. Assoc. 44, 263–269.
- 2552 Raisz, E.J., 1962. Principles of cartography. McGraw-Hill, New York, 315 pp.
- Rea, B.R., Whalley, W.B., Dixon, T.S., Gordon, J.E., 1999. Plateau icefields as contributing areas to
 valley glaciers and the potential impact on reconstructed ELAs: a case study from the Lyngen
 Alps, North Norway. Ann. Glaciol. 28, 97–102.
- Reinardy, B.T.I., Leighton, I., Marx, P.J., 2013. Glacier thermal regime linked to processes of annual
 moraine formation at Midtdalsbreen, southern Norway. Boreas 42(4), 896–911.
- Reuther, A.U., Urdea, P., Geiger, C., Ivy-Ochs, S., Niller, H.P., Kubik, P.W., Heine, K., 2007. Late
 Pleistocene glacial chronology of the Pietrele Valley, Retezat Mountains, Southern
 Carpathians constrained by 10Be exposure ages and pedological investigations. Quat. Int.
 164, 151–169.
- Rippin, D.M., Pomfret, A., King, N., 2015. High resolution mapping of supra-glacial drainage pathways reveals link between microchannel drainage density, surface roughness and surface reflectance. Earth Surf. Proc. Land. 40(10), 1279–1290.
- Robb, C., Willis, I., Arnold, N., Guðmundsson, S., 2015. A semi-automated method for mapping
 glacial geomorphology tested at Breiðamerkurjökull, Iceland. Remote Sens. Environ. 163,
 80–90.
- Robinson, A.H., Morrison, J.L., Muehrcke, P.C., Kimerling, A.J., Guptill, S.C., 1995. Elements of
 cartography (6th Edition). Wiley & Sons, Chichester, 674 pp.
- Robinson, P., Dowdeswell, J.A., 2011. Submarine landforms and the behavior of a surging ice cap
 since the last glacial maximum: The open-marine setting of eastern Austfonna, Svalbard. Mar.
 Geol. 286(1), 82–94.
- Rose, J., Smith, M.J., 2008. Glacial geomorphological maps of the Glasgow region, western central
 Scotland. J. Maps 4, 399–416.
- Rossi, C., Rodriguez Gonzalez, F., Fritz, T., Yague-Martinez, N., Eineder, M., 2012. TanDEM-X
 calibrated raw DEM generation. ISPRS J. Photogramm. Remote Sens. 73, 12–20.
- Rupke, J., De Jong, M.G.G., 1983. Slope collapse destroying ice-marginal topography in the Walgau
 (Vorarlberg, Austria)- an example of the application of a 1:10000 geomorphological mapping
 system. Mater. Physiogeogr. 5, 33–41.
- Ryan, J.C., Hubbard, A.L., Todd, J., Carr, J.R., Box, J.E., Christoffersen, P., Holt, T.O., Snooke, N.,
 2015. Repeat UAV photogrammetry to assess calving front dynamics at a large outlet glacier
 draining the Greenland Ice Sheet. The Cryosphere 9, 1–11.
- Sagredo, E.A., Moreno, P.I., Villa-Martínez, R., Kaplan, M.R., Kubik, P.W., Stern, C.R., 2011.
 Fluctuations of the Última Esperanza ice lobe (52 S), Chilean Patagonia, during the last
 glacial maximum and termination 1. Geomorphology 125(1), 92–108.
- Saha, K., Wells, N.A., Munro-Stasiuk, M., 2011. An object-oriented approach to automated landform mapping: A case study of drumlins. Computers & Geosciences 37(9), 1324–1336.
- Sahlin, E.A.U., Glasser, N.F., 2008. A geomorphological map of Cadair Idris, Wales. J. Maps 4, 299–314.
- Salcher, B.C., Hinsch, R., Wagreich, M., 2010. High-resolution mapping of glacial landforms in the
 North Alpine Foreland, Austria. Geomorphology 122, 283–293.
- Schoeneich, P., 1993. Cartographie géomorphologique en Suisse, Institut de Géogrphie Lausanne,
 Travaux et recherches, Lausanne, pp. 1–13.
- Schomacker, A., 2008. What controls dead-ice melting under different climate conditions? A
 discussion. Earth-Sci. Rev. 90, 103–113.

- Schomacker, A., Kjaer, K.H., 2008. Quantification of dead-ice melting in ice-cored moraines at the
 high-Arctic glacier Holmströmbreen, Svalbard. Boreas 37, 211–225.
- Schomacker, A., Benediktsson, Í.Ö., Ingólfsson, Ó., 2014. The Eyjabakkajökull glacial landsystem,
 Iceland: Geomorphic impact of multiple surges. Geomorphology 218, 98–107.
- Seguinot, J., Rogozhina, I., Stroeven, A.P., Margold, M., Kleman, J., 2016. Numerical simulations of
 the Cordilleran ice sheet through the last glacial cycle. The Cryosphere 10, 639-664.
- Shean, D.E., Alexandrov, O., Moratto, Z.M., Smith, B.E., Joughin, I.R., Porter, C., Morin, P., 2016.
 An automated, open-source pipeline for mass production of digital elevation models (DEMs)
 from very-high-resolution commercial stereo satellite imagery. ISPRS J. Photogramm.
 Remote Sens. 116, 101–117.
- 2606 Sissons, J.B., 1967. The Evolution of Scotland's Scenery. Oliver & Boyd, Edinburgh, 259 pp.
- Sissons, J.B., 1977a. The Loch Lomond Readvance in the northern mainland of Scotland. In: Gray,
 J.M., Lowe, J.J. (Eds.), Studies in the Scottish Lateglacial Environment. Permagon Press,
 Oxford, pp. 45–60.
- Sissons, J.B., 1977b. The Loch Lomond Readvance in southern Skye and some palaeoclimatic
 implications. Scott. J. Geol. 13, 23–36.
- Sissons, J.B., 1979a. The limit of the Loch Lomond Advance in Glen Roy and vicinity. Scott. J. Geol.
 15, 31–42.
- Sissons, J.B., 1979b. The Loch Lomond advance in the Cairngorm Mountains. Scott. Geogr. Mag. 95,
 66–82.
- Smith, A.M., Murray, T., 2009. Bedform topography and basal conditions beneath a fast-flowing
 West Antarctic ice stream. Quat. Sci. Rev. 28(7), 584–596.
- Smith, A.M., Murray, T., Nicholls, K.W., Makinson, K., Ađalgeirsdóttir, G., Behar, A.E., Vaughan,
 D. G., 2007. Rapid erosion, drumlin formation, and changing hydrology beneath an Antarctic ice stream. Geology 35(2), 127–130.
- Smith, G.R., Woodward, J.C., Heywood, D.I., Gibbard, P.L., 2000. Interpreting Pleistocene glacial
 features from SPOT HRV data using fuzzy techniques. Computers & Geosciences 26(4),
 479–490.
- Smith, M.J., Clark, C.D., 2005. Methods and visulisation of digital elevation models for landform
 mapping. Earth Surf. Process. Land. 30, 885–900.
- Smith, M.J., Knight, J., 2011. Palaeoglaciology of the last Irish ice sheet reconstructed from striae
 evidence. Quat. Sci. Rev. 30(1-2), 147–160.
- Smith, M.J., Wise, S.M., 2007. Problems of bias in mapping linear landforms from satellite imagery.
 Int. J. Appl. Earth Obs. Geoinform. 9(1), 65–78.
- Smith, M.J., Rose, J., Booth, S., 2006. Geomorphological mapping of glacial landforms from remotely sensed data: An evaluation of the principal data sources and an assessment of their quality. Geomorphology 76, 148–165.
- Smith, M.J., Griffiths, J., Paron, P. (Eds.)., 2011. Geomorphological Mapping: Methods and
 Applications. Developments in Earth Surface Processes, Volume 15. Elsevier, Oxford, 610
 pp.
- Smith, M.J., Anders, N.S., Keesstra, S.D., 2016b. CLustre: semi-automated lineament clustering for
 palaeo-glacial reconstruction. Earth Surf. Process. Land. 41(3), 364–377.
- Smith, M.W., Carrivick, J.L., Quincey, D.J., 2016a. Structure from motion photogrammetry in physical geography. Prog. Phys. Geogr. 40(2), 247–275.
- Sollas, W.J., 1896. A map to show the distribution of eskers in Ireland. The Sci. Trans. R. Dublin Soc.
 5, 785–822.
- Spagnolo, M., Clark, C.D., Hughes, A.L.C., Dunlop, P., Stokes, C.R., 2010. The planar shape of drumlins. Sediment. Geol. 232, 119–129.
- Spagnolo, M., Clark, C.D., Ely, J.C., Stokes, C.R., Anderson, J.B., Andreassen, K., Graham, A.G.C.,
 King, E.C., 2014. Size, shape and spatial arrangement of megascale glacial lineations from a
 large and diverse dataset. Earth Surf. Process. Land. 39, 1432–1448.
- Spagnolo, M., Phillips, E., Piotrowski, J.A., Rea, B.R., Clark, C.D., Stokes, C.R., Carr, S.J., Ely, J.C.,
 Ribolini, A., Wysota, W., Szuman, I., 2016. Ice stream motion facilitated by a shallowdeforming and accreting bed. Nat. Commun. 7, 10723.

- Spagnolo, M., Bartholomaus, T.C., Clark, C.D., Stokes, C.R., Atkinson, N., Dowdeswell, J.A., Ely,
 J.C., Graham, A.G.C., Hogan, K.A., King, E.C., Larter, R.D., Livingstone, S.J., Pritchard,
 H.D., 2017. The periodic topography of ice stream beds: Insights from the Fourier spectra of
 mega-scale glacial lineations. J. Geophys. Res.: Earth Surf. 122, 1355–1373.
- Stansell, N.D., Polissar, P.J., Abbott, M.B., 2007. Last glacial maximum equilibrium-line altitude and
 paleo-temperature reconstructions for the Cordillera de Mérida, Venezuelan Andes. Quat.
 Res. 67, 115–127.
- 2657Stokes, C.R., 2002. Identification and mapping of palaeo-ice stream geomorphology from satellite2658imagery: implications for ice stream functioning and ice sheet dynamics. Int. J. Remote Sens.265923(8), 1557–1563.
- Stokes, C.R., 2018. Geomorphology under ice streams: moving from form to process. Earth Surf.
 Process. Land., 43, 85–123.
- Stokes, C.R., Clark, C.D., 1999. Geomorphological criteria for identifying Pleistocene ice streams.
 Ann. Glaciol. 28(1), 67–74.
- 2664 Stokes, C.R., Clark, C.D., 2001. Palaeo-ice streams. Quat. Sci. Rev. 20(13), 1437–1457.
- Stokes, C.R., Clark, C.D., 2002. Are long subglacial bedforms indicative of fast ice flow? Boreas, 31(3), 239–249.
- Stokes, C.R., Clark, C.D. 2003. The Dubawnt Lake palaeo-ice stream: evidence for dynamic ice sheet
 behaviour on the Canadian Shield and insights regarding the controls on ice-stream location
 and vigour. Boreas 32, 263–279.
- Stokes, C.R., Tarasov, L., 2010. Ice streaming in the Laurentide Ice Sheet: A first comparison
 between data-calibrated numerical model output and geological evidence. Geophys. Res. Lett.
 37, L01501.
- 2673Stokes, C.R., Clark, C.D., Storrar, R., 2009. Major changes in ice stream dynamics during2674deglaciation of the north-western margin of the Laurentide Ice Sheet. Quat. Sci. Rev. 28, 721–2675738.
- Stokes, C.R., Spagnolo, M., Clark, C.D., Ó Cofaigh, C., Lian, O.B., Dunstone, R.B., 2013. Formation
 of mega-scale glacial lineations on the Dubawnt Lake Ice Stream bed: 1. size, shape and
 spacing from a large remote sensing dataset. Quat. Sci. Rev. 77, 190–209.
- Stokes, C.R., Corner, G.D., Winsborrow, M.C.M., Husum, K., Andreassen, K., 2014. Asynchronous
 response of marine-terminating outlet glaciers during deglaciation of the Fennoscandian Ice
 Sheet. Geology 42, 455–458.
- Stokes, C.R., Tarasov, L., Blomdin, R., Cronin, T.M., Fisher, T.G., Gyllencreutz, R., Hättestrand, C.,
 Heyman, J., Hindmarsh, R.C.A., Hughes, A.L.C., Jakobsson, M., Kirchner, N., Livingstone,
 S.J., Margold, M., Murton, J.B., Noormets, R., Peltier, W.R., Peteet, D.M., Piper, D.J.W.,
 Preusser, F., Renssen, H., Roberts, D.H., Roche, D.M., Saint-Ange, F., Stroeven, A.P., Teller,
 J.T., 2015. On the reconstruction of palaeo-ice sheets: Recent advances and future challenges.
 Quat. Sci. Rev. 125, 15–49.
- Stokes, C.R., Margold, M., Creyts, T.T., 2016a. Ribbed bedforms on palaeo-ice stream beds resemble
 regular patterns of basal shear stress ('traction ribs') inferred from modern ice streams. J.
 Glaciol. 62, 696–713.
- 2691Stokes, C.R., Margold, M., Clark, C.D., Tarasov, L., 2016b. Ice stream activity scaled to ice sheet2692volume during Laurentide Ice Sheet deglaciation. Nature 530(7590), 322–326.
- Storrar, R.D., Livingstone, S.J., 2017. Glacial geomorphology of the northern Kivalliq region,
 Nunavut, Canada, with an emphasis on meltwater drainage systems. J. Maps 13, 153–164.
- Storrar, R., Stokes, C.R., 2007. A Glacial geomorphological map of Victoria Island, Canadian Arctic.
 J. Maps 3, 191–210.
- Storrar, R.D., Stokes, C.R., Evans, D.J.A., 2013. A map of large Canadian eskers from Landsat satellite imagery. J. Maps 9, 456–473.
- Storrar, R.D., Stokes, C.R., Evans, D.J.A., 2014. Morphometry and pattern of a large sample
 (>20,000) of Canadian eskers and implications for subglacial drainage beneath ice sheets.
 Quat. Sci. Rev. 105, 1–25.
- Storrar, R.D., Jones, A., Evans, D.J.A., 2017. Small-scale topographically-controlled glacier flow
 switching in an expanding proglacial lake at Breiðamerkurjökull, SE Iceland. J. Glaciol.
 63(240), 745–750.

- Streuff, K., Forwick, M., Szczuciński, W., Andreassen, K., Ó Cofaigh, C., 2015. Submarine landform
 assemblages and sedimentary processes related to glacier surging in Kongsfjorden, Svalbard.
 arktos 1(1), 14.
- Stroeven, A.P., Fabel, D., Harbor, J., Hättestrand, C., Kleman, J., 2002a. Quantifying the erosional impact of the Fennoscandian ice sheet in the Torneträsk-Narvik corridor, northern Sweden, based on cosmogenic radionuclide data. Geogr. Ann. 84A, 275–287.
- Stroeven, A.P., Fabel, D., Hättestrand, C., Harbor, J., 2002b. A relict landscape in the centre of
 Fennoscandian glaciation: Cosmogenic radionuclide evidence of tors preserved through
 multiple glacial cycles. Geomorphology 44, 145–154.
- Stroeven, A.P., Harbor, J., Fabel, D., Kleman, J., Hättestrand, C., Elmore, D., Fink, D., Fredin, O.,
 2006. Slow, patchy landscape evolution in northern Sweden despite repeated ice-sheet
 glaciation. GSA Special Paper 398, 387–396.
- Stroeven, A.P., Hättestrand, C., Heyman, J., Kleman, J., Morén, B.M., 2013. Glacial geomorphology of the Tian Shan. J. Maps 9(4), 505–512.
- Stroeven, A.P. Fabel, D., Margold, M., Clague, J.J., Xu, S., 2014. Investigating absolute chronologies
 of glacial advances in the NW sector of the Cordilleran Ice Sheet with terrestrial in situ
 cosmogenic nuclides. Quat. Sci. Rev. 92, 429–443.
- Stroeven, A.P., Hättestrand, C., Kleman, J., Heyman, J., Fabel, D., Fredin, O., Goodfellow, B.W.,
 Harbor, J.M., Jansen, J.D., Olsen, L., Caffee, M.W., Fink, D., Lundqvist, J., Rosqvist, G.C.,
 Strömberg, B., Jansson, K.N., 2016. Deglaciation of Fennoscandia. Quat. Sci. Rev. 147, 91–
 121.
- Sugden, D.E., 1970. Landforms of Deglaciation in the Cairngorm Mountains, Scotland. Trans. Inst.
 Br. Geogr. 51, 201–219.
- 2728 Sugden, D.E., 1978. Glacial erosion by the Laurentide Ice Sheet. J. Glaciol. 20, 367–391.
- 2729 Sugden, D.E., John, B.S., 1976. Glaciers and Landscape. Arnold, London, 376 pp.
- Thorp, P.W., 1986. A mountain icefield of Loch Lomond Stadial age, western Grampians, Scotland.
 Boreas 15, 83–97.
- Tonkin, T.N., Midgley, N.G., Cook, S.J., Graham, D.J., 2016. Ice-cored moraine degradation mapped
 and quantified using an unmanned aerial vehicle: A case study from a polythermal glacier in
 Svalbard. Geomorphology 258, 1–10.
- Trommelen, M.S., Ross, M., 2010. Subglacial landforms in northern Manitoba, Canada, based on remote sensing data. J. Maps 2010, 618–638.
- Trommelen, M.S., Ross, M., 2014. Distribution and type of sticky spots at the centre of a deglacial
 streamlined lobe in northeastern Manitoba, Canada. Boreas 43, 557–576.
- Trotter, F.M., 1929. The glaciation of the eastern Edenside, the Alston Block and the Carlisle Plain.
 Quart. J. Geol. Soc. Lond. 88, 549–607.
- Turner, A.J., Woodward, J., Stokes, C.R., Ó Cofaigh, C., Dunning, S., 2014a. Glacial geomorphology
 of the Great Glen Region of Scotland. J. Maps 10, 159–178.
- Turner, D., Lucieer, A., Wallace, L., 2014b. Direct georeferencing of ultrahigh-resolution UAV
 imagery. IEEE Trans. Geosci. Remote Sens. 52(5), 2738–2745.
- van der Bilt, W.G.M., Bakke, J., Balascio, N.L., 2016. Mapping sediment–landform assemblages to
 constrain lacustrine sedimentation in a glacier-fed lake catchment in northwest Spitsbergen. J.
 Maps 12(5), 985–993.
- 2748 Van Dorsser, J.J., Salomé, A.I., 1973. Different methods of detailed geomorphological mapping.
 2749 K.N.A.G. Geografisch Tijdschrift VII: 71–74.
- Walker, D.A., Auerbach, N.A., Shippert, M.M., 1995. NDVI, biomass, and landscape evolution of
 glaciated terrain in northern Alaska. Polar Record, 31(177), 169-178.
- Welch, R., 1966. A comparison of aerial films in the study of the Breiðamerkur glacier area, Iceland,
 The Photogramm. Rec. 5, 289–306.
- Welch, R., 1967. The Application of Aerial Photography to the Study of a Glacial Area.
 Breiðamerkur, Iceland, Unpublished PhD thesis, University of Glasgow.
- Welch, R., 1968. Color aerial photography applied to the study of a glacial area. In: Smith, J.T. (Ed.),
 Manual of Color Aerial Photography. American Society of Photogrammetry and Remote
 Sensing, Falls Church, VA, pp. 400–401.

- Westoby, M.J., Brasington, J., Glasser, N.F., Hambrey, M.J., Reynolds, J.M., 2012. 'Structure-fromMotion' photogrammetry: A low-cost, effective tool for geoscience applications.
 Geomorphology 179, 300–314.
- Westoby, M.J., Glasser, N.F., Hambrey, M.J., Brasington, J., Reynolds, J.M., Hassan, M.A., 2014.
 Reconstructing historic Glacial Lake Outburst Floods through numerical modelling and geomorphological assessment: Extreme events in the Himalaya. Earth Surf. Process. Land.
 39(12), 1675–1692.
- Westoby, M.J., Dunning, S.A., Woodward, J., Hein, A.S., Marrero, S.M., Winter, K., Sugden, D.E.,
 2767 2016. Interannual surface evolution of an Antarctic blue-ice moraine using multi-temporal
 DEMs. Earth Surf. Dyn. 4, 515–529.
- Wilson, S.B., 2005. Morphological analysis and mapping of Loch Lomond Stadial moraines using
 digital photogrammetry and geographical information systems. Unpublished PhD thesis,
 University of Glasgow, 360 pp.
- Winsborrow, M.C.M., Andreassen, K., Corner, G.D., Laberg, J.S., 2010. Deglaciation of a marinebased ice sheet: Late Weichselian palaeo-ice dynamics and retreat in the southern Barents Sea
 reconstructed from onshore and offshore glacial geomorphology. Quat. Sci. Rev. 29(3–4),
 424–442.
- Winsborrow, M.C.M., Stokes, C.R., Andreassen, K., 2012. Ice stream flow switching during deglaciation of the southwestern Barents Sea. Geol. Soc. Am. Bull. 124(3–4), 275–290.
- Wolf, P.R., DeWitt, B.A., Wilkinson, B.E., 2013. Elements of Photogrammetry with Application in
 GIS, Fourth Edition. McGraw-Hill Education.
- 2780 Wright, W.B., 1912. The drumlin topography of south Donegal. Geol. Mag. 9, 153–159.
- Wyshnytzky, C.E., 2017. On the mechanisms of minor moraine formation in high-mountain
 environments of the European Alps. Unpublished PhD thesis, Queen Mary University of
 London, 329 pp.
- Yu, P., Eyles, N., Sookhan, S., 2015. Automated drumlin shape and volume estimation using high
 resolution LiDAR imagery (Curvature Based Relief Separation): A test from the Wadena
 Drumlin Field, Minnesota. Geomorphology 246, 589–601.
- Zasadni, J., Kłapyta, P., 2016. From valley to marginal glaciation in alpine-type relief: Lateglacial
 glacier advances in the Pięć Stawów Polskich/Roztoka Valley, High Tatra Mountains, Poland.
 Geomorphology 253, 406–424.
- Zech, R., Abramowski, U., Glaser, B., Sosin, P., Kubik, P.W. and Zech, W., 2005. Late Quaternary
 glacial and climate history of the Pamir Mountains derived from cosmogenic 10Be exposure
 ages. Quat. Res. 64(2), 212–220.
- 2793 Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S.U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstrøm, A.P., Anderson, B., Bajracharya, S., Baroni, C., Braun, L.N., 2794 2795 Cáceres, B.E., Casassa, G., Cobos, G., Dávila, L.R., Delgado Granados, H., Demuth, M.N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J.O., Holmlund, P., 2796 2797 Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V.V., Portocarrero, C.A., Prinz, R., Sangewar, C.V., Severskiy, I., Sigurðsson, O., Soruco, A., Usubaliev, R., Vincent, C., 2015. 2798 2799 Historically unprecedented global glacier decline in the early 21st century. J. Glaciology 61, 745-762. 2800
- 2801 2802
- 2803
- 2804
- 2805
- 2806
- 2807
- 2808
- 2809

- 2810 2811 2812 2813 2814 2815 2816 2817 2818 **Figure captions** 2819 2820 Figure 1. Digitised versions of two geomorphological maps drawn in the field for (A) Coire Easgainn 2821 and (B) Glen Odhar in the Monadhliath, Central Scottish Highlands. These field maps were used in 2822 the production of a 1:57,500 geomorphological map for the entire region (Boston, 2012a, b). 2823 2824 Figure 2. The aerial photograph overlay-mapping process using an example from the mountain Arkle, 2825 NW Scotland. (A) aerial photograph at an average scale of ~1:25,000 (extract from photo 38 88 087; 2826 ©RCAHMS 1988); (B) scan of original overlay mapped through a stereoscope from (A) (see Section
- 2.2.2 for method description), focusing on moraines, fluted moraines and the approximate upper limit
 of scree slopes as seen from the aerial photograph; (C) compiled, rectified geomorphological map,
 incorporating moraines and fluted moraines from (B) and additional data from field mapping, such as
 the exact upper limits of scree slopes, orientation of striae, solifluction lobes and mountaintop detritus,
 for example. For description and interpretation of the geomorphology, see Lukas (2006).
- 2832

Figure 3. Satellite image (A) and geomorphological mapping (B) showing suites of moraines formed by the Lago General Carrera–Buenos Aires ice lobe of the former Patagonian Ice Sheet, located to the east of the present-day Northern Patagonian Icefield. A combination of remotely-sensed datasets and field mapping were used to circumvent issues of localised cloud cover, as visible in A. Where areas were obscured, SPOT-5 and DigitalGlobe images available in *Google Earth* were used. Note the extensive outwash development between moraine suites. The geomorphological map extract is taken from Bendle et al. (2017a).

2840

Figure 4. Comparison of WorldView-2 satellite imagery (June 2012, European Space Imaging) with
digital colour aerial photographs (2006, Loftmyndir ehf) for the Skálafellsjökull foreland, SE Iceland.
A. Panchromatic satellite image (0.5 m ground sampled distance, GSD). B. Multispectral satellite
image (2.0 m GSD). C. Pansharpened three-band natural colour satellite image (0.5 m GSD). D.
Digital colour aerial photographs (0.41 m GSD). The satellite imagery is of sufficient resolution to
allow mapping of small-scale (<2 m in height) annual moraines (see Chandler et al., 2016a, b).

2847

Figure 5. Geomorphology of the Finsterwalderbreen foreland, Svalbard, mapped in the field and from
a digital aerial photograph captured in 2004. Photograph provided by the NERC Earth Observation
Data Centre. Modified from Lovell et al. (2018).

2851

Figure 6. Views at various points along the length of the 1890 surge end moraine at Eyjabakkajökull,
Iceland, visualised in *ESRI ArcScene* (Benediktsson et al., 2010). Aerial orthophotographs from 2008
are draped over a 3 m grid DEM with 1.5x vertical exaggeration.

2855

Figure 7. High-resolution geomorphological mapping of part of the Fláajökull foreland, Iceland,
based on UAV-derived imagery (Evans et al., 2016a). A 1:350 scale version of this map is freely
available for download from *Journal of Maps*: http://dx.doi.org/10.1080/17445647.2015.1073185.

Figure 8. Geomorphological mapping of lineations in northwest Saskatchewan, Canada, from SRTM imagery (modified from Norris et al., 2017). (A) and (B) Densely spaced drumlins displaying low length to width ratios. (C) and (D) Highly elongated fluting orientated NE–SW east of the Grizzly Bear Hills. Geomorphological map extracts in (C) and (D) show lineations (black lines), eskers (red lines) and meltwater channels (dashed blue lines).

2865

Figure 9. Examples of landforms in relief-shaded DEMs. Red indicates higher elevations and blue lower elevations. (A) Lineations in N Canada shown in 16 m resolution CDED data. (B) De Geer moraines in SW Finland shown in 2 m resolution LiDAR data. (C) Lineations of the Dubawnt Lake Ice Stream shown in 5 m resolution ArcticDEM mosaic data. (D) Esker-fed ice-contact outwash fan in SW Finland shown in 2 m resolution LiDAR data. See Table 2 for DEM data sources.

2871

Figure 10. Conceptual diagrams illustrating the distinction between ground sampled distance (B and E) and pixel size (C and F). The ground distances between two measurements by the detector (i.e. the ground sampled distances) are 30 m and 50 m in (B) and (E), respectively. These ground sample distances are then assigned to pixels in the resulting 30 x 30 m (C) and 50 x 50 m (F) digital images. Note, resultant images may fail to represent accurately the shape of the objects (upper row) or even may fail to reproduce them (lower row), even where the size of the object is the same or larger than the sampling distance.

2879

Figure 11. Geometric artefacts that may be present in space- and air-borne radar captured imagery, resulting from the effects of relief. (A) **Foreshortening**, occurring where the slope of the local terrain is less than the incidence angle (γ). The facing slope, a - b, becomes compressed to $a_1 - b_1$ in the resulting image. (B) **Layover**, occurring in steep terrain when the slope angle is greater than the

- incidence angle. As a mountain-top, *b*, is closer to the sensor than the base, *a*, this causes layover in the imagery (an incorrect positioning of b_1 relative to a_1). (C) **Radar shadow** in areas of rugged terrain as the illumination is from an oblique source. No data is recorded for the region $b_1 - d_1$. (D) In regions of varying topography, a **combination of artefacts** may be present: points *b* and *c* will be impacted by layover and will be positioned incorrectly relative to *a*; no data will be recorded for the region between *c* and *d* due to radar shadow; foreshortening occurs at slope facet d - e; further radar shadow occurs at e - f; and foreshortening at *f* and *g*. After Clark (1997).
- 2891
- Figure 12. Extracts from hillshaded relief models of Ben More Coigach, NW Scottish Highlands, showing the effect of geometric artefacts on the models. The hillshades were generated with azimuths of 45° (A) and 315° (B). Stretching of upland terrain during processing of the DEM data results in blurred regions on the hillshaded relief models. NEXTMap DSM from Intermap Technologies Inc. provided by NERC via the NERC Earth Observation Data Centre.
- 2897

Figure 13. Example mapping of subglacial bedforms from the Strait of Magellan, Patagonia (A–C), and the Dubawnt Lake Ice Stream (D–F). The bedforms are mapped as polylines along landform crests in B and E, and mapped as polygons delineating lower-break-of-slope in C and F. The Dubawnt Lake Ice Stream polylines (Stokes and Clark, 2003) and polygons (Dunstone, 2014) were mapped by different mappers at different times, which may account for small inconsistencies. For further details on the bedform examples from the Strait of Magellan, see Lovell et al. (2011) and Darvill et al. (2014).

2905

Figure 14. Geomorphological mapping of Coire Easgainn, Monadhliath, Scotland, using a
combination of NEXTMap DSMs, analogue aerial photographs and field mapping. Modified from
Boston (2012a, b).

2909

Figure 15. Examples of landforms in icefield and valley glacier settings mapped on medium to coarse resolution imagery. Landforms observed in the Chagan Uzun Valley, Russian Altai, displayed on (A) SPOT image and (B) Landsat 7 ETM+ image. (C) Associated geomorphological map extract from Gribenski et al. (2016). Moraines in the Anadyr Lowlands, Far NE Russia, displayed on (D) semitransparent shaded ViewFinder Panorama (VFP) DEM data (NE solar azimuth) draped over the raw VFP DEM. (E) Associated mapping of moraines (black polygons) from Barr and Clark (2012).

2916

Figure 16. Geomorphological mapping (A) from the Múlajökull foreland, Iceland, completed as part
of a process-oriented study examining the internal architecture and structural evolution of a Little Ice
Age terminal moraine at this surge-type glacier (Benediktsson et al., 2015). The mapping was

combined with sedimentological investigations (B) to produce a process-form model of moraineformation and evolution (C).

2922

2923 Figure 17. Geomorphological mapping of the foreland of Skálafellsjökull, an active temperate outlet 2924 of Vatnajökull, SE Iceland. (A) Digital aerial photographs (2006; 0.41 m GSD; Loftmyndir ehf), pan-2925 sharpened WorldView-2 multi-spectral satellite imagery (2012; 0.5 m GSD; European Space 2926 Imaging), a UAV-derived DEM (2013; 0.09 m GSD) and field mapping were employed to produce 2927 the mapping extract (B). A compromise on the level of detail was made, with annual moraines 2928 mapped along crestlines due to image resolution and map readability. This mapping detail was 2929 sufficient for calculating crest-to-crest moraine spacing (ice-margin retreat rates) shown in (C), which 2930 was the principal purpose of the study. Modified from Chandler et al. (2016a, b).

2931

Figure 18. Idealised workflow for mapping palaeo-ice sheet geomorphology. Some pathways in the workflow are optional (grey dashed lines) depending on data availability and the feasibility and applicability of particular methods. Note, where analogue (hard-copy) aerial photographs are used for mapping, processing of acetate overlays would be undertaken *after* mapping from the aerial photographs. Further details on image processing are shown on the processing workflow available as supplementary material.

2938

2939 Figure 19. Idealised workflow for mapping alpine- and plateau-style ice mass geomorphology. In this 2940 scenario, digital remotely-sensed datasets are used and this necessitates image processing before 2941 mapping is undertaken. Ideally, GNSS surveys would be conducted in order to process digital aerial 2942 photographs, as depicted in the workflow. Some pathways are optional (grey dashed lines) depending 2943 on data availability and the feasibility and applicability of particular methods. Although 2944 sedimentology is shown as 'optional', it is highly desirable to undertake sedimentological 2945 investigations, wherever possible. Alternative image processing solutions are available and readers 2946 should consult with the detailed processing workflow which is available as supplementary material.

2947

Figure 20. Idealised workflow for mapping alpine- and plateau-style ice mass geomorphology. In this scenario, analogue (hard-copy) aerial photographs are used and this necessitates image processing *after* mapping is undertaken. Some pathways are optional (grey dashed lines) depending on data availability and the feasibility and applicability of particular methods. Although sedimentology is shown as 'optional', it is highly desirable to undertake sedimentological investigations, wherever possible. Alternative image processing solutions are available and readers should consult with the detailed processing workflow which is available as supplementary material.

Satellite	Sensor	Temporal coverage	Spectral bands	Spatial resolution (m)	Source	Example studies
Landsat 1–5	MSS	1972-	4	80	USGS Earth Explorer	Clark and Stokes (2001); Stokes and Clark (2002,
		2013			(earthexplorer.usgs.gov)	2003); Jansson et al. (2003); see also Clark (1997, Table 1)
Landsat 4–5	TM	1982-	1	120	Global Land Cover Facility	Punkari (1995); Alexanderson et al. (2002); De
		2013	6	30	(landcover.org)	Angelis (2007); Storrar et al. (2013); Orkhonselenge (2016)
Landsat 7	ETM+	1999–	1	60		Kassab et al (2013); Stroeven et al. (2013); Darvill et
			6	30		al. (2014); Blomdin et al. (2016b); Ely et al. (2016b);
			1	15		Ercolano et al. (2016); Lindholm and Heyman (2016); Storrar and Livingstone (2017); see also Clark (1997, Table 1)
Landsat 8	OLI/TIRS	2013-	2	100		Espinoza (2016); Carrivick et al. (2017); Storrar and
			8	30		Livingstone (2017)
			1	15		
Terra	ASTER	2000-	5	90	LP DAAC	Glasser and Jansson (2005, 2008); Glasser et al.
			6	20	(LPDAAC.usgs.gov)	(2005); Lovell et al. (2011); Sagredo et al. (2011);
			5	15		Darvill et al (2014); Ercolano et al. (2016)
ERS 1	SAR	1991–	1	30	European Space Agency	Clark et al. (2000); Clark and Stokes (2001); Heiser
		2000			(earth.esa.int)	and Roush (2001); see also Clark (1997, Table 1)
SPOT 1–3	HRV	1986–	3	20	Airbus Defence and Space	Smith et al. (2000); Coronato et al. (2009)
		2009	1	20	(intelligence-airbusds.com)	
			1	10		
SPOT 4	HRVIR	1998–	1	10		Trommelen and Ross (2010, 2014); Ercolano et al.
		2013	3	20		(2016) [viewed in Google Earth [™]]; McHenry and
			1	20		Dunlop (2016); Principato et al. (2016)
SPOT 5	HRG/HRS	2002-	1	2.5, 5		Trommelen and Ross (2010, 2014); Ercolano et al.
		2015	3	10		(2016) [viewed in Google Earth [™]]; McHenry and
			1	20		Dunlop (2016); Principato et al. (2016); Bendle et al. (2017a)
SPOT 6–7	NAOMI	2012-	1	1.5		Gribenski et al. (2016)
			4	6		
CORONA/ARGON/LANYARD	КН1-КН6	1959– 1972	1	1.8–140	USGS Earth Explorer (earthexplorer.usgs.gov)	Alexanderson et al (2002); Zech et al. (2005); Lifton et al (2014)
IKONOS	HRG	1999_	1	1	DigitalGlobe	Iuval et al. (2011): Kłanyta (2013): Zasadni and
11201005	into	2015	4	4	(digitalglobe com)	K_{i} Kanvta (2016)
		2015		•	(angitungiobe.com)	15mp Jun (2010)

Table 1. Primary satellite imagery types used in glacial geomorphological mapping and example applications. Broadly ordered in terms of spatial resolution.

COSMO-Skymed	SAR	2008-	1/3/15/16/20	1	e-GEOS (e-geos.it)	da Rosa et al. (2013a)	
Quickbird	HRG	2001-	1	0.61	DigitalGlobe	da Rosa et al. (2011, 2013b); May et al (2011);	
		2014	4	2.44	(digitalglobe.com)	Lovell et al. (2011)	
GeoEye-1		2008-	1	0.46		Westoby et al. (2014)	
			4	1.84	European Space Imaging		
WorldView-2		2009-	1	0.46	(euspaceimaging.com)	Jamieson et al. (2015); Chandler et al. (2016a);	
			8	1.84		Evans et al. (2016e); Ewertowski et al (2016)	
Google Earth [™] (specific image details not given)	n/a	n/a	n/a	n/a	Google Earth	Margold and Jansson (2011); Margold et al. (2011); Kassab et al (2013); Stroeven et al. (2013); Darvill et al (2014); Blomdin et al. (2016b); Evans et al. (2016d); Li et al (2016); Lindholm and Heyman (2016); Orkhonselenge (2016)	
Dataset	Coverage	Spatial resolution (m)	RMSE or CE90 (m)				
--	----------------------	--	------------------	------------	---	---	--
			Vertical	Horizontal	Data source(s)	Example studies	
SRTM ¹	Global	~90 (3 arc-second) ~30 (1 arc-second)	~5–13	-	Global Land Cover Facility (landcover.org) USGS Earth Resources and Science Center (eros.usgs.gov)	Glasser and Jansson (2008); Barr and Clark (2009); Ó Cofaigh et al. (2010); Morén et al. (2011); Stroeven et al. (2013); Darvill et al. (2014); Evans et al. (2014, 2016d); Trommelen and Ross (2014); Stokes et al. (2016a); Ely et al. (2016b); Lindholm and Heyman (2016)	
ASTER GDEM (V2)	Global	~30 (1 arc-second)	~8.7	-	LP DAAC Global Data Explorer (gdex.cr.usgs.gov/gdex) NASA Reverb (reverb.echo.nasa.gov/reverb)	Barr and Clark (2012); Blomdin et al. (2016a, b); Lindholm and Heyman (2016)	
Canadian Digital Elevation Dataset (CDED)	Canada	~20 (0.75 arc-second)	-	-	Natural Resources Canada (geogratis.gc.ca)	Margold et al. (2011, 2015a); Evans et al. (2016c); Storrar and Livingstone (2017)	
USGS National Elevation Dataset (NED) ²	US	~30 (1 arc-second) ~10 (1/3 arc-second)	~2.4	-	US Geological Survey (ned.usgs.gov)	Hess and Briner (2009); Margold et al. (2015a); Ely et al. (2016a)	
TanDEM-X	Global	~12 (0.4 arc-second)	<10	<10	German Aerospace Center (DLR) (tandemx-science.dlr.de)	Pipaud et al. (2015)	
NEXTMap Britain TM	UK	5	~1	2.5	NERC Earth Observation Data Centre ³ (ceda.ac.uk)	Livingstone et al. (2008); Finlayson et al. (2010, 2011); Hughes et al. (2010); Brown et al. (2011a); Boston (2012a, b); Pearce et al. (2014); Turner et al. (2014a)	
ArcticDEM	Arctic	2	2.0	3.8	Polar Geospatial Center (pgc.umn.edu/data/arcticdem)	Levy et al. (2017)	
Maanmittauslaitos LiDAR DEM	Finland	2	~0.3	-	National Land Survey of Finland (maanmittauslaitos.fi)	Ojala et al. (2015); Ojala (2016); Mäkinen et al. (2017)	
Ny Nationell Höjdmodell	Sweden	2	~0.1	-	Lantmäteriet (lantmateriet.se)	Dowling et al. (2015, 2016); Greenwood et al. (2015); Möller and Dowling (2016); Peterson et al. (2017)	
Environment Agency LiDAR DEM	UK (partial)	2, 1, 0.5 and 0.25	0.05 - 0.15	0.4	DEFRA Environment Data (environment.data.gov.uk)	Miller et al. (2014)	
Iceland Met Office and Institute of Earth Sciences, University of Iceland, LiDAR DEM ⁴	Iceland (partial)	<5	<0.5	-	Iceland Meteorological Office (en.vedur.is)	Brynjólfsson et al. (2014, 2016); Benediktsson et al. (2016); Jónsson et al. (2016)	

Table 2. Examples of DEM datasets with national- to international-coverage that have been employed in glacial geomorphological map production.

¹ SRTM data was only freely available with a spatial resolution of ~90 m (3 arc-seconds) outside of the United States until late 2015 when the highest resolution data were thereafter made available globally (see <u>http://www2.jpl.nasa.gov/srtm/</u>)

² The USGS NED dataset has been superseded by the 3D Elevation Program (3DEP), with this data available as seamless 1/3 arc-second, 1 arc-second and 2 arc-second DEMs (see <u>https://nationalmap.gov/3DEP/3dep_prodserv.html</u>)

³ NEXTMap BritainTM data is freely available to NERC staff and NERC-funded researchers, though subsets can be applied for by non-NERC-funded researchers under a Demonstrator User License Agreement (DULA)

 4 The Icelandic LiDAR DEM data are available at 5 m resolution, but it is possible to derive higher-resolution DEMs (e.g. 2 m) from the point clouds using denser interpolation.

Table 3. Summary of the glacial settings where the main geomorphological mapping methods and remotely-sensed data types are *most* appropriate. \checkmark = the method/dataset is appropriate and should be used (where the dataset is available). • = the method is applicable in certain cases, depending on factors such as the resolution of the *specific* dataset, the size of the study area and landforms, and the accessibility of the study area.

Glacial setting	DEMs	Coarse satellite imagery	LiDAR DEMs	High-resolution satellite imagery	Aerial photographs	UAV imagery	Field mapping
Ice sheets	✓	\checkmark	\checkmark				
Ice sheet sectors/lobes	\checkmark	\checkmark	\checkmark	•	•		•
Ice-caps	•	•	•	\checkmark	✓		✓
Icefields			•	\checkmark	✓		✓
Valley (outlet) glaciers			•	\checkmark	✓	•	\checkmark
Cirque glaciers			•	\checkmark	✓	•	\checkmark
Modern glacier forelands			•	\checkmark	✓	\checkmark	\checkmark