

Critical Vertices and Edges in H -free Graphs^{*}

Daniël Paulusma^{1**}, Christophe Picouleau², and Bernard Ries³

¹ Durham University, Durham, UK, daniel.paulusma@durham.ac.uk

² CNAM, Laboratoire CEDRIC, Paris, France, christophe.picouleau@cnam.fr

³ University of Fribourg, Department of Informatics, Fribourg, Switzerland,
bernard.ries@unifr.ch

Abstract. A vertex or edge in a graph is critical if its deletion reduces the chromatic number of the graph by one. We consider the problems of deciding whether a graph has a critical vertex or edge, respectively. We give a complexity dichotomy for both problems restricted to H -free graphs, that is, graphs with no induced subgraph isomorphic to H . Moreover, we show that an edge is critical if and only if its contraction reduces the chromatic number by one. Hence, we also obtain a complexity dichotomy for the problem of deciding if a graph has an edge whose contraction reduces the chromatic number by one.

Keywords. edge contraction, vertex deletion, chromatic number.

1 Introduction

For a positive integer k , a k -colouring of a graph $G = (V, E)$ is a mapping $c : V \rightarrow \{1, 2, \dots, k\}$ such that no two end-vertices of an edge are coloured alike, that is, $c(u) \neq c(v)$ if $uv \in E$. The *chromatic number* $\chi(G)$ of a graph G is the smallest integer k for which G has a k -colouring. The well-known COLOURING problem is to decide if $\chi(G) \leq k$ for a given graph G and integer k . If k is not part of the input, then we call this problem k -COLOURING instead. Lovász [15] proved that 3-COLOURING is NP-complete.

Due to its computational hardness, the COLOURING problem has been well studied for special graph classes. We refer to the survey [11] for an overview of the results on COLOURING restricted to graph classes characterized by one or two forbidden induced subgraphs. In particular, Král', Kratochvíl, Tuza, and Woeginger [14] classified COLOURING for H -free graphs, that is, graphs that do not contain a single graph H as an induced subgraph. To explain their result we need the following notation. For a graph F , we write $F \subseteq_i G$ to denote that F is an induced subgraph of a graph G . The *disjoint union* of two graphs G_1 and G_2 is the graph $G_1 + G_2$, which has vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$. We write rG for the disjoint union of r copies of G . The graphs K_r , P_r and C_r denote the complete graph, path and cycle on r vertices, respectively. We can now state the theorem of Král' et al.

Theorem 1 ([14]). *Let H be a graph. If $H \subseteq_i P_4$ or $H \subseteq_i P_1 + P_3$, then COLOURING restricted to H -free graphs is polynomial-time solvable, otherwise it is NP-complete.*

For a vertex u or edge e in a graph G , we let $G - u$ and $G - e$ be the graph obtained from G by deleting u or e , respectively. Note that such an operation may

^{*} Results in this paper appeared in extended abstracts in the proceedings of ISCO 2016 [21] and LAGOS 2017 [20].

^{**} Author supported by the Leverhulme Trust (RPG-2016-258).

reduce the chromatic number of the graph by at most 1. We say that u or e is *critical* if $\chi(G - u) = \chi(G) - 1$ or $\chi(G - e) = \chi(G) - 1$, respectively. A graph is *vertex-critical* if every vertex is critical and *edge-critical* if every edge is critical.

If a problem is polynomial-time solvable, we can ask for a certifying algorithm. Such an algorithm provides a certificate that demonstrates the correctness of solutions. For the COLOURING problem, a yes-certificate is a k -colouring of the graph and a no-certificate can be a vertex-critical or edge-critical subgraph of G of size at most $f(k)$ for some function f depending only on k . To increase our understanding of the COLOURING problem and to obtain certifying algorithms that solve COLOURING for special graph classes, vertex-critical and edge-critical graphs have been studied intensively in the literature, see for instance [4–6, 8, 10, 12, 13, 16] for certifying algorithms for (subclasses of) H -free graphs and in particular P_r -free graphs.

In this paper we consider the problems CRITICAL VERTEX and CRITICAL EDGE, which are to decide if a graph has a critical vertex or critical edge, respectively. We also consider the edge contraction variant of these two problems. We let G/e denote the graph obtained from G after contracting $e = vw$, that is, after removing v and w and replacing them by a new vertex made adjacent to precisely those vertices adjacent to v or w in G (without creating multiple edges). Contracting an edge may reduce the chromatic number of the graph by at most 1. An edge e is *contraction-critical* if $\chi(G/e) = \chi(G) - 1$. This leads to the CONTRACTION-CRITICAL EDGE problem, which is to decide if a graph has a contraction-critical edge.

1.1 Our Results

We prove the following complexity dichotomies for CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE restricted to H -free graphs.

Theorem 2. *Let H be a graph. If a graph $H \subseteq_i P_4$ or of $H \subseteq_i P_1 + P_3$, then CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE restricted to H -free graphs are polynomial-time solvable, otherwise they are NP-hard or co-NP-hard.*

We note that the classification in Theorem 2 coincides with the one in Theorem 1. The polynomial-time cases for CRITICAL VERTEX and CONTRACTION-CRITICAL EDGE can be obtained from Theorem 1. The reason for this is that a class of H -free graphs is not only closed under vertex deletions, but also under edge contractions whenever H is a *linear forest*, that is, a disjoint union of a set of paths (see Section 5 for further details). However, no class of H -free graphs is closed under edge deletion. We get around this issue by proving, in Section 2, that an edge is critical if and only if it is contraction-critical. Hence, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are equivalent.

The NP-hardness constructions of Theorem 1 cannot be used for proving the hard cases for CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE. Instead we construct new hardness reductions in Sections 3 and 4. In Section 3 we prove that the three problems are NP-hard for H -free graphs if H contains a claw or a cycle on three or more vertices. Note that we cannot prove membership in NP, as COLORING is NP-complete for the class of H -free graphs if $H \supseteq_i K_{1,3}$ or $H \supseteq_i C_r$ for some $r \geq 3$ due to Theorem 1. As such, it is not clear if there exists a certificate. In the remaining case H is a linear forest. In Section 4 we prove that the three problems are co-NP-hard even for $(C_5, 4P_1, 2P_1 + P_2, 2P_2)$ -free graphs (also here we cannot prove membership in NP, as COLOURING is NP-complete for $(C_5, 4P_1, 2P_1 + P_2, 2P_2)$ -free

graphs [14]). In Section 5 we combine the known cases with our new results from Sections 2–4 in order to prove Theorem 2.

1.2 Consequences

Our results have consequences for the computational complexity of two graph blocker problems. Let S be some fixed set of graph operations, and let π be some fixed graph parameter. Then, for a given graph G and integer $k \geq 0$, the S -BLOCKER(π) problem asks if G can be modified into a graph G' by using at most k operations from S so that $\pi(G') \leq \pi(G) - d$ for some given *threshold* $d \geq 0$. Over the last few years, the S -BLOCKER(π) problem has been well studied, see for instance [1–3, 7, 9, 19–23]. If S consists of a single operation that is either a vertex deletion or edge contraction, then S -BLOCKER(π) is called VERTEX DELETION BLOCKER(π) or CONTRACTION BLOCKER(π), respectively. By taking $d = k = 1$ and $\pi = \chi$ we obtain the problems CRITICAL VERTEX and CONTRACTION-CRITICAL EDGE, respectively. We showed in [20] how the results for CRITICAL VERTEX and CONTRACTION-CRITICAL EDGE can be extended with other results to get complexity dichotomies for VERTEX DELETION BLOCKER(χ) and CONTRACTION BLOCKER(χ) for H -free graphs.

1.3 Future Work

For some family of graphs $\{H_1, \dots, H_p\}$ and integer $p \geq 2$, a graph G is (H_1, \dots, H_p) -free if G is H -free for every $H \in \{H_1, \dots, H_p\}$. As a direction for future research we propose classifying the computational complexity of our three problems for (H_1, \dots, H_p) -free graphs for any $p \geq 2$. We note that such a classification for COLORING is still wide open even for $p = 2$ (see [11]). Hence, research in this direction might lead to an increased understanding of the complexity of the COLORING problem.

2 Equivalence

We prove the following result, which implies that the problems CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are equivalent.

Proposition 1. *An edge is critical if and only if it is contraction-critical.*

Proof. Let $e = uv$ be an edge in a graph G . First suppose that e is critical, so $\chi(G-e) = \chi(G) - 1$. Then u and v are colored alike in any coloring of $G - e$ that uses $\chi(G - e)$ colors. Hence, the graph G/e , obtained by contracting e in G , can also be colored with $\chi(G - e)$ colors. Indeed, we simply copy a $(\chi(G - e))$ -coloring of $G - e$ such that the new vertex in G/e is colored with the same color as u and v in $G - e$. Hence $\chi(G/e) = \chi(G - e) = \chi(G) - 1$, which means that e is contraction-critical.

Now suppose that e is contraction-critical, so $\chi(G/e) = \chi(G) - 1$. By copying a $\chi(G/e)$ -coloring of G/e such that u and v are colored with the same color as the new vertex in G/e , we obtain a coloring of $G - e$. So we can color $G - e$ with $\chi(G/e)$ colors as well. Hence $\chi(G - e) = \chi(G/e) = \chi(G) - 1$, which means that e is critical. \square

3 Forbidding Claws or Cycles

The *claw* is the 4-vertex star $K_{1,3}$ on vertices a, b, c, d and edges ab, ac and ad . In this section we prove that the problems CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are NP-hard for H -free graphs whenever the graph H contains a claw or a cycle on at least three vertices.

A subset K of vertices in G is a *clique* if any two vertices in K are adjacent to each other. Let \mathcal{G} be a graph class with the following property: if $G \in \mathcal{G}$, then so are $2G$ and $G + K_r$ for any $r \geq 1$. We call such a graph class *clique-proof*. We will prove that several classes of H -free graphs are clique-proof. For such classes of H -free graphs we can apply the following theorem.

Theorem 3. *If COLORING is NP-complete for a clique-proof graph class \mathcal{G} , then CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are NP-hard for \mathcal{G} .*

Proof. By Proposition 1 it suffices to prove the theorem for CRITICAL VERTEX and CONTRACTION-CRITICAL EDGE. We first consider the CONTRACTION-CRITICAL EDGE problem. Let \mathcal{G} be a graph class that is clique-proof. From a given graph $G \in \mathcal{G}$ and integer $\ell \geq 1$ we construct the graph $G' = 2G + K_{\ell+1}$. Note that $G' \in \mathcal{G}$ by definition and that $\chi(G') = \max\{\chi(G), \ell + 1\}$. We first prove that $\chi(G) \leq \ell$ if and only if G' contains a contraction-critical edge.

Suppose that $\chi(G) \leq \ell$. Then $\chi(G') = \chi(K_{\ell+1}) = \ell + 1$. In G' we contract an edge of the $K_{\ell+1}$. This yields the graph $G^* = 2G + K_{\ell}$, which has chromatic number $\chi(G^*) = \ell$, as $\chi(K_{\ell}) = \ell$ and $\chi(G) \leq \ell$. As $\chi(G') = \ell + 1$, this means that $\chi(G^*) = \chi(G') - 1$. Hence G' contains a contraction-critical edge.

Now suppose that G' contains a contraction-critical edge. Let G^* be the resulting graph after contracting this edge. Then $\chi(G^*) = \chi(G') - 1$. As contracting an edge in one of the two copies of G in G' does not lower the chromatic number of G' , the contracted edge must be in the $K_{\ell+1}$, that is, $G^* = 2G + K_{\ell}$. Since this has lowered the chromatic number, we conclude that $\chi(G') = \chi(K_{\ell+1}) = \ell + 1$ and $\chi(G^*) = \chi(2G + K_{\ell}) = \max\{\chi(G), \ell\} = \ell$. The latter equality implies that $\chi(G) \leq \ell$.

From the above we conclude that CONTRACTION-CRITICAL EDGE is NP-hard. We can prove that CRITICAL VERTEX is NP-hard by using the same arguments. \square

We also need a result of Maffray and Preissmann as a lemma.

Lemma 1 ([17]). *The 3-COLORING problem is NP-complete for C_3 -free graphs.*

We are now ready to prove the main result of this section.

Theorem 4. *Let H be a graph such that $H \supseteq_i K_{1,3}$ or $H \supseteq_i C_r$ for some $r \geq 3$. Then the problems CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are NP-hard for H -free graphs.*

Proof. By Proposition 1 it suffices to prove the theorem for CRITICAL VERTEX and CONTRACTION-CRITICAL EDGE. If for a graph H , these problems are NP-hard for the class of H -free graphs, then they are also NP-hard on the class of H' -free graphs for any graph H' with $H' \supseteq_i H$. Therefore we only have to prove the theorem for the cases where $H = K_{1,3}$ or $H = C_r$ for every $r \geq 3$. If $H = K_{1,3}$ or $H = C_r$ for some $r \geq 4$, then the class of H -free graphs is clique-proof. Hence, in these cases NP-hardness immediately follows from Theorems 1 and 3.




Fig. 1. The Grötzsch graph.

Suppose $H = C_3$. We reduce from 3-COLORING restricted to C_3 -free graphs. This problem is NP-complete by Lemma 1. Let G be a C_3 -free graph that is an instance of 3-COLORING. We obtain an instance of CRITICAL VERTEX or CONTRACTION-CRITICAL EDGE as follows. Take the disjoint union of two copies of G and the Grötzsch graph F (see Figure 1), which is known to be 4-colorable but not 3-colorable (see [24]). Call the resulting graph G' , so $G' = 2G + F$. As G and F are C_3 -free, G' is C_3 -free. We claim that G is 3-colorable if and only if G' has a critical vertex if and only if G' has a contraction-critical edge. This can be proven via similar arguments as used in the proof of Theorem 3, with F playing the role of $K_{\ell+1}$. \square

4 Forbidding Linear Forests

In this section we prove our second hardness result needed to show Theorem 2. We first introduce some additional terminology.

Let G be a graph. The graph \overline{G} denotes the *complement* of G , that is, the graph with vertex set $V(G)$ and an edge between two vertices u and v if and only if u and v are not adjacent in G . A *clique cover* of a graph G is a set \mathcal{K} of cliques in G , such that each vertex of G belongs to exactly one clique of \mathcal{K} . The *clique covering number* $\sigma(G)$ is the size of a smallest clique cover of G . Note that $\chi(G) = \sigma(\overline{G})$. The size of a largest clique in a graph G is denoted by $\omega(G)$.

The hardness construction in the proof of our next result uses clique covers. Král et al. [14] proved that COLORING is NP-hard for $(C_5, 4P_1, P_1 + 2P_2, 2P_2)$ -free graphs. This does not give us hardness for CRITICAL VERTEX or CRITICAL EDGE, but we can use some elements of their construction. For instance, we reduce from a similar NP-complete problem, namely the NP-complete problem MONOTONE 1-IN-3-SAT, which is defined as follows.¹ Let Φ be a formula in conjunctive normal form, that is, Φ consists of a clause set C , where each clause in C consists of literals from a variable set X . For an instance Φ of MONOTONE 1-IN-3-SAT it holds that each clause in C consists of three distinct positive literals from X and each variable of X occurs in exactly three clauses (observe that this implies that $|C| = |X|$). The question is whether Φ has a truth assignment, such that each clause is satisfied by exactly one variable. In

¹ The MONOTONE 1-IN-3 SAT problem is also known as the POSITIVE 1-IN-3 SAT problem, and in the literature, monotonicity may also refer to the property that every clause consists of either only positive literals or only negative literals.

that case we say that Φ is *1-satisfiable*. Moore and Robson proved that the problem is NP-complete.

Lemma 2 ([18]). MONOTONE 1-IN-3-SAT is NP-complete.

We are now ready to prove the main result of this section.

Theorem 5. The problems CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE are co-NP-hard for $(C_5, 4P_1, 2P_1 + P_2, 2P_2)$ -free graphs.

Proof. By Proposition 1 it suffices to consider CRITICAL VERTEX and CRITICAL EDGE. We will first consider CRITICAL VERTEX and show that the equivalent problem whether a graph has a vertex whose deletion reduces the clique covering number by one is co-NP-hard for $(C_5, K_4, \overline{2P_1 + P_2}, C_4)$ -free graphs. We call such a vertex *critical* as well. The complement of a $(C_5, K_4, \overline{2P_1 + P_2}, C_4)$ -free graph is $(C_5, 4P_1, 2P_1 + P_2, 2P_2)$ -free. Hence by proving this co-NP-hardness result we will have proven the theorem for CRITICAL VERTEX.

As mentioned, we reduce from MONOTONE 1-IN-3-SAT, which is NP-complete due to Lemma 2. Given an instance Φ of MONOTONE 1-IN-3-SAT with clause set C and variable set X , we construct a graph $G = (V, E)$ as follows. For every clause $c \in C$, the clause gadget $G_c = (V_c, E_c)$ is a path of length 5. For $c = (x, y, z)$, we let three pairwise non-adjacent vertices $c(x), c(y), c(z)$ of G_c correspond to the three variables x, y, z . We denote the other two vertices of G_c by a_1^c and a_2^c , such that $G_c = c(x)a_1^c c(y)a_2^c c(z)$. For each variable $x_h \in X$ we let the variable gadget Q_{x_h} consist of the triangle $c(x_h)c'(x_h)c''(x_h)c(x_h)$, where c, c', c'' are the three clauses containing x_h . See Figure 2 for an illustration of the construction. We observe that $|V(G)| = 5n$ and that G is $(C_5, K_4, \overline{2P_1 + P_2}, C_4)$ -free with $\omega(G) = 3$.




Fig. 2. The clause gadget G_c and three variable gadgets Q_x, Q_y and Q_z .

In order to prove co-NP-hardness we first need to deduce a number of properties of our gadget. We do this via a number of claims.

Claim 1. There exists a minimum clique cover of G , in which each vertex a_i^c ($c \in C$, $1 \leq i \leq 2$) is covered by a clique of size 2.

We prove Claim 1 as follows. If for some $c \in C$, vertex a_1^c or a_2^c forms a 1-vertex clique in \mathcal{K} , then we add $c(x)$ or $c(z)$, respectively, to it. This proves Claim 1.

Now let \mathcal{K} be a minimum clique cover. By Claim 1, we may assume without loss of generality that each a_i^c is covered by a clique of size 2. Since the clause gadgets G_c are pairwise non-intersecting and isomorphic to P_5 , it takes at least three cliques to cover the vertices of every G_c . As the vertices a_i^c form an independent set, exactly $2n$ cliques

are needed to cover the $2n$ vertices a_i^c . As each a_i^c is covered by a clique of size 2, we also find that $2n$ vertices $c(x)$ are covered by these cliques. Since $\omega(G) = 3$, at least $n/3$ other cliques are necessary to cover the n remaining vertices $c(x)$. Hence, \mathcal{K} has size at least $\frac{7}{3}n$, that is,

$$\sigma(G) \geq \frac{7}{3}n.$$

We now prove three more claims.

Claim 2. Φ is 1-satisfiable if and only if $\sigma(G) = \frac{7}{3}n$.

We prove Claim 2 as follows. First suppose Φ is 1-satisfiable. We construct a clique cover \mathcal{K} in the following way. If x_h is true, then we let \mathcal{K} contain the triangle Q_{x_h} . Since each clause c contains exactly one true variable for each G_c , exactly one vertex of each G_c is covered by a variable gadget Q_{x_h} . Then \mathcal{K} contains two cliques of size 2 covering the four other vertices of G_c . Hence \mathcal{K} has size $\frac{7}{3}n$. As $\sigma(G) \geq \frac{7}{3}n$, this implies that $\sigma(G) = \frac{7}{3}n$.

Now suppose $\sigma(G) = \frac{7}{3}n$. Let \mathcal{K} be a minimum clique cover of G . By Claim 1, we may assume without loss of generality that each a_i^c is covered by a clique of size 2. Then at least $n/3$ other cliques are necessary to cover the vertices $c(x)$ that are not in a 2-vertex clique with a vertex a_i^c . Hence, as $\sigma(G) = \frac{7}{3}n$, these vertices are covered by exactly $n/3$ triangles, each one corresponding to one variable x (these are the only triangles in G). We assign the value true to a variable $x_h \in X$ if and only if its corresponding triangle Q_{x_h} is in the clique cover. Then, for each $c \in C$, exactly one variable is true, namely the one that corresponds to the unique vertex of G_c covered by a triangle. So Φ is 1-satisfiable. This completes the proof of Claim 2.

Claim 3. If G has a clique cover $\mathcal{K} = \{K_1, \dots, K_{\frac{7}{3}n}\}$, then each $K_i \in \mathcal{K}$ consists of either two or three vertices.

We prove Claim 3 as follows. As $\sigma(G) \geq \frac{7}{3}n$ and $|\mathcal{K}| = \frac{7}{3}n$, we find that \mathcal{K} is a minimum clique cover. With each $v \in V$, we associate a *weight* $w_v \geq 0$ as follows. For $K_i \in \mathcal{K}$ and $v \in K_i$, we define $w_v = 1/|K_i|$. Since $\omega(G) = 3$ we have $w_v \in \{\frac{1}{3}, \frac{1}{2}, 1\}$. So we have

$$\sum_{G_c} \sum_{v \in V_c} w_v = \sum_{v \in V} w_v = \sum_{i=1}^{\frac{7}{3}n} \sum_{v \in K_i} w_v = \frac{7}{3}n,$$

where the first equality holds, because the clause gadgets G_c are vertex-disjoint. We show that for every c we have $\sum_{v \in V_c} w_v \geq \frac{7}{3}$. Since every a_i^c has exactly two neighbours and these neighbours are not adjacent, we have $w_{a_i^c} \in \{\frac{1}{2}, 1\}$. If there exists an index i such that $w_{a_i^c} = 1$, then

$$\sum_{v \in V_c} w_v \geq 1 + \frac{1}{2} + 3 \times \frac{1}{3} = \frac{5}{2} > \frac{7}{3}.$$

Now, if a_i^c has weight $w_{a_i^c} = \frac{1}{2}$ for $i = 1, 2$, then a_i^c is covered by a clique of size 2 and the second vertex of this clique has weight $\frac{1}{2}$ as well by definition. Thus if $w_{a_i^c} = \frac{1}{2}$, at least two vertices of $c(x), c(y), c(z)$ have weight $\frac{1}{2}$. It follows that

$$\sum_{v \in V_c} w_v \geq 2 \times \frac{1}{2} + 2 \times \frac{1}{2} + \frac{1}{3} = \frac{7}{3}.$$

Hence $\sum_{v \in V_c} w_v = \frac{7}{3}$ if and only if each vertex of G_c is in a clique of size 2 or 3. Since $\sum_{G_c} \sum_{v \in V_c} w_v = \frac{7}{3}n$, we obtain $\sum_{v \in V_c} w_v = \frac{7}{3}$ for every $c \in C$. We conclude that each clique in \mathcal{K} is of size 2 or 3. This completes the proof of Claim 3.

Claim 4. If $\sigma(G) > \frac{7}{3}n$, then G has a minimum clique cover \mathcal{K} that contains a clique of size 1.

We prove Claim 4 as follows. Suppose $\sigma(G) > \frac{7}{3}n$. For contradiction, assume that every minimum clique cover of G has no cliques of size 1. Let \mathcal{K} be a minimum clique cover of G . By Claim 1, we may assume without loss of generality that each a_i^c is covered by a clique of size 2. Hence the remaining vertex $c(x)$ is covered by some clique $K_i \in \mathcal{K}$, such that either $K_i = \{c(x), c'(x)\}$ or $K_i = \{c(x), c'(x), c''(x)\}$.

If $K_i = \{c(x), c'(x)\}$, then $c''(x)$ is covered by some clique $K_j = \{c''(x), a\}$. However, then we can take $K_i = \{c(x), c'(x), c''(x)\}$ and $K_j = \{a\}$ to obtain a minimum clique cover with $|K_j| = 1$, a contradiction. Hence $K_i = \{c(x), c'(x), c''(x)\}$. As this holds for every G_c we find that $\sigma(G) = \frac{7}{3}n$, a contradiction. This completes the proof of Claim 4.

We claim that Φ is a 1-satisfiable if and only if G has no critical vertex. First suppose that Φ is 1-satisfiable. By Claims 2 and 3 we find that $\sigma(G) = \frac{7}{3}n$ and every clique in any minimum clique cover of G has size greater than 1. Hence, there is no vertex u of G with $\sigma(G - u) \leq \sigma(G) - 1$, that is, G has no critical vertex.

Now suppose that Φ is not 1-satisfiable. By Claims 2 and 4 we find that $\sigma(G) > \frac{7}{3}n$ and that there exists a minimum clique cover that contains a clique $\{u\}$ of size 1. This means that $\sigma(G - u) = \sigma(G) - 1$. So u is a critical vertex.

We are left to consider the CRITICAL EDGE problem. Just as for the CRITICAL VERTEX problem, we focus on the complementary problem. For CRITICAL EDGE, this is the problem of deciding if a graph has a non-edge whose addition to the graph (as an edge) reduces the clique covering number by one. We call such a non-edge *critical* as well and prove that this equivalent problem is co-NP-hard for $(C_5, K_4, 2P_1 + P_2, C_4)$ -free graphs. In order to do this we modify the previous construction by making the paths G_c isomorphic to P_9 . To be more precise, we let $G_c = c(x)a_1^c a_2^c a_3^c c(y)a_4^c a_5^c a_6^c c(z)$. Again the resulting graph G is $(C_4, C_5, K_4, 2P_1 + P_2)$ -free.

First suppose that Φ is 1-satisfiable. By using the same arguments as before we find that every clique in any minimum clique cover of G has size greater than 1. Hence, as G is K_4 -free, every clique in any minimum clique cover \mathcal{K} of G has size 2 or 3. This means that we can only merge two cliques of \mathcal{K} into one new clique via adding an edge if both cliques already have some edge in common. However, then the union of these two cliques induces a subgraph of G isomorphic to $2P_1 + P_2$. This is not possible, as G is $2P_1 + P_2$ -free. We conclude that G has no critical non-edge.

Now suppose that Φ is not 1-satisfiable. Then, by using the same arguments as before, we can prove that there exists a minimum clique cover \mathcal{K} that contains a clique $\{u\}$ of size 1. By the adjusted construction of G_c we find that u is adjacent to exactly one vertex of a 2-vertex clique $\{v, w\}$ of \mathcal{K} (in the previous construction with copies of P_5 instead of copies of P_9 , it could have happened that u is adjacent to only vertices that belong to distinct 3-vertex cliques of \mathcal{K} , namely when $u = a_i^c$ for some $1 \leq i \leq 2$ and $c \in C$). We may assume without loss of generality that u is adjacent to v but not to w . Then by adding the edge uw , which yields the graph $G + uw$, we merge the two cliques $\{u\}$ and $\{v, w\}$ into one new clique $\{u, v, w\}$. Hence we obtain $\sigma(G + uw) = \sigma(G) - 1$. So uw is a critical non-edge of G . This completes the proof of Theorem 5. \square

5 The Proof of Theorem 2

We are now ready to prove Theorem 2, which we restate below. In particular, the three problems are NP-hard for H -free graphs if H contains a claw or a cycle on three or more vertices and they are co-NP-hard if H contains an induced subgraph from $\{C_5, 4P_1, 2P_1 + P_2, 2P_2\}$.

Theorem 2. (restated) *If a graph $H \subseteq_i P_4$ or of $H \subseteq_i P_1 + P_3$, then CRITICAL VERTEX, CRITICAL EDGE and CONTRACTION-CRITICAL EDGE restricted to H -free graphs are polynomial-time solvable, otherwise they are NP-hard or co-NP-hard.*

Proof. Let $H \subseteq_i P_1 + P_3$ or $H \subseteq_i P_4$. Let G be an H -free graph. By Theorem 1 we can compute $\chi(G)$ in polynomial time. We note that any vertex deletion results in a graph that is H -free as well. Hence in order to solve CRITICAL VERTEX we can compute the chromatic number of $G - v$ for each vertex v in polynomial time and compare it with $\chi(G)$. As $(P_1 + P_3)$ -free graphs and P_4 -free graphs are closed under edge contraction as well, we can follow the same approach for solving CONTRACTION-CRITICAL EDGE. By Proposition 1 we obtain the same result for CRITICAL EDGE.

Now suppose that neither $H \subseteq_i P_1 + P_3$ nor $H \subseteq_i P_4$. If H has a cycle or an induced claw, then we use Theorem 4 to obtain NP-hardness. Assume not. Then H is a disjoint union of r paths for some $r \geq 1$. If $r \geq 4$ we use Theorem 5 to obtain co-NP-hardness. If $r = 3$ then either $H = 3P_1 \subseteq_i P_1 + P_3$, which is not possible, or $H \supseteq_i 2P_1 + P_2$ and we can apply Theorem 5 again to obtain co-NP-hardness. Suppose $r = 2$. If both paths contain an edge, then $2P_2 \subseteq_i H$. If at most one path has edges, then it must have at least four vertices, as otherwise $H \subseteq_i P_1 + P_3$. This means that $2P_1 + P_2 \subseteq_i H$. In both cases we apply Theorem 5 to obtain co-NP-hardness. If $r = 1$, then H is a path on at least five vertices, which means $2P_2 \subseteq_i H$. We apply Theorem 5 again to obtain co-NP-hardness. \square

Acknowledgments. We thank two anonymous reviewers for helpful comments on our paper.

References

1. C. Bazgan, C. Bentz, C. Picouleau and B. Ries, Blockers for the stability number and the chromatic number, *Graphs and Combinatorics* 31 (2015) 73–90.
2. C. Bazgan, S. Toubaline and Z. Tuza, The most vital nodes with respect to independent set and vertex cover, *Discrete Applied Mathematics* 159 (2011) 1933–1946.
3. C. Bentz, M.-C. Costa, D. de Werra, C. Picouleau and B. Ries, Weighted transversals and blockers for some optimization problems in graphs, *Progress in Combinatorial Optimization*, Wiley-ISTE, 2012.
4. D. Bruce, C.T. Hoàng, and J. Sawada, A certifying algorithm for 3-colorability of P_5 -free graphs, *Proc. ISAAC 2009, Lecture Notes in Computer Science* 5878 (2009), 595–604.
5. M. Chudnovsky, J. Goedgebeur, O. Schaudt and M. Zhong, Obstructions for three-coloring and list three-coloring H -free graphs, *Manuscript*, arXiv:1703.05684.
6. M. Chudnovsky, J. Goedgebeur, O. Schaudt and M. Zhong, Obstructions for three-coloring graphs without induced paths on six vertices, *Proc. SODA 2016*, 1774–1783.
7. M.-C. Costa, D. de Werra, C. Picouleau, Minimum d -blockers and d -transversals in graphs, *Journal of Combinatorial Optimization* 22 (2011) 857–872.
8. H.S. Dhaliwal, A.M. Hamel, C.T. Hoàng, F. Maffray, T.J.D. McConnell and S.A. Panait, On color-critical $(P_5, \overline{P_5})$ -free graphs, *Discrete Applied Mathematics* 216 (2017) 142–148.

9. O. Diner, D. Paulusma, C. Picouleau and B. Ries, Contraction blockers for graphs with forbidden induced paths, Proc. CIAC 2015, Lecture Notes in Computer Science 9079 (2015) 194–207.
10. J. Goedgebeur and O. Schaudt, Exhaustive generation of k -critical H -free graphs, Journal of Graph Theory 87 (2018) 188–207.
11. P.A. Golovach, M. Johnson, D. Paulusma and J. Song, A survey on the computational complexity of colouring graphs with forbidden subgraphs, Journal of Graph Theory 84 (2017) 331–363.
12. P. Hell and S. Huang, Complexity of coloring graphs without paths and cycles, Discrete Applied Mathematics 216 (2017) 211–232.
13. C.T. Hoàng, B. Moore, D. Recoskie, J. Sawada and M. Vatshelle, Constructions of k -critical P_5 -free graphs, Discrete Applied Mathematics 182 (2015) 91–98.
14. D. Král', J. Kratochvíl, Z. Tuza, and G.J. Woeginger, Complexity of coloring graphs without forbidden induced subgraphs, Proc. WG 2001, Lecture Notes in Computer Science 2204 (2001) 254–262.
15. L. Lovász, Coverings and coloring of hypergraphs, Proc. 4th Southeastern Conference on Combinatorics, Graph Theory, and Computing, Utilitas Mathematica (1973) 3–12.
16. F. Maffray and G. Morel, On 3-Colorable P_5 -free graphs, SIAM Journal on Discrete Mathematics 26 (2012) 1682–1708.
17. F. Maffray and M. Preissmann, On the NP-completeness of the k -colorability problem for triangle-free graphs, Discrete Mathematics 162 (1996) 313–317.
18. C. Moore, and J. M. Robson, Hard tiling problems with simple tiles, Discrete and Computational Geometry 26 (2001) 573–590.
19. F.M. Pajouh, V. Boginski and E. L. Pasiliao, Minimum vertex blocker clique problem, Networks 64 (2014) 48–64.
20. D. Paulusma, C. Picouleau and B. Ries, Reducing the chromatic number by vertex or edge deletions, Proc. LAGOS 2017, Electronic Notes in Discrete Mathematics 62 (2017) 243–248.
21. D. Paulusma, C. Picouleau and B. Ries, Reducing the clique and chromatic number via edge contractions and vertex deletions, Proc. ISCO 2016, Lecture Notes in Computer Science 9849 (2016) 38–49.
22. D. Paulusma, C. Picouleau and B. Ries, Blocking independent sets for H -free graphs via edge contractions and vertex deletions, Proc. TAMC 2017, Lecture Notes in Computer Science 10185 (2017) 470–483.
23. B. Ries, C. Bentz, C. Picouleau, D. de Werra, M.-C. Costa and R. Zenklusen, Blockers and transversals in some subclasses of bipartite graphs: when caterpillars are dancing on a grid, Discrete Mathematics 310 (2010) 132–146.
24. D. B. West, Introduction to Graph Theory, Prentice-Hall, 1996.