

Obesity, bone density relative to body weight and prevalent vertebral fracture at age 62 years: the Newcastle Thousand Families Study

H.A. Rudman ¹, F. Birrell ², M.S. Pearce ³, S.P. Tuck ², R.M.Francis ³, L. Treadgold ¹, K. Hind ^{2,4}

¹ Leeds Institute of Cardiovascular & Metabolic Medicine, University of Leeds, Leeds, United Kingdom.

² Institute of Cellular Medicine, Newcastle University, Newcastle Upon Tyne, United Kingdom.

³ Institute of Health and Society, Newcastle University, Sir James Spence Institute of Child Health, Royal Victoria Infirmary, Newcastle upon Tyne, United Kingdom.

⁴ Department of Sport and Exercise Sciences, Durham University, County Durham, United Kingdom.

For correspondence: Dr K Hind PhD, CCD. Assistant Professor of Sport and Exercise Sciences, and

Fellow of the Wolfson Research for Health and Wellbeing, Department of Sport and Exercise

Sciences, Durham University, 42 Old Elvet, Durham, DH1 3HN. United Kingdom.

Telephone: +44 191 334 6837

Email: karen.hind@durham.ac.uk

Abstract

Obesity is a global epidemic and there remains uncertainty over the effect of obesity on skeletal health, particularly in the context of osteoporosis. The aim of this study was to investigate associations of body mass index (BMI) and obesity with bone mineral density (BMD) and prevalent vertebral fracture (VF) in men and women aged 62 years. Three hundred and forty two men and women aged 62.5 ± 0.5 years from the Newcastle Thousand Families Study birth cohort, underwent DXA evaluations of femoral neck and lumbar spine BMD, and of the lateral spine for vertebral fracture assessment. The likelihood of prevalent VF was significantly increased in men when compared to women (OR = 2.7, $p < 0.001$, 95% CI 1.7– 4.4). As BMI increased in women, so did the likelihood of prevalent any-grade VF (OR = 1.09, $p = 0.006$, 95% CI 1.02 – 1.17). Compared to normal weight women, obese women were more likely to have at least one VF (OR = 2.65, $p = 0.025$, CI 1.13 – 6.20) and at least one grade 1 VF (OR = 4.39, $p = 0.005$, CI 1.57 – 12.28). Obese men were more likely to have a grade 2 and/or 3 VF compared to men of normal weight (OR = 3.36, $p = 0.032$, CI 1.11 – 10.16). In men and women, BMI was negatively associated with femoral neck BMD/ weight ($R = - 0.65$, $R = - 0.66$, $p < 0.001$) and lumbar spine BMD/ weight ($R = - 0.66$, $R = - 0.60$, $p < 0.001$). Obesity appears to be a risk factor for prevalent VF, and although absolute BMD is higher in obese individuals, the increase does not appear commensurate to their increase in body weight.

Keywords: *obesity; bone; fracture; DXA; fat.*

Introduction

1
2 Obesity is a global epidemic, with worldwide rates having more than doubled since 1980 (1).
3
4 Despite efforts to improve diet and exercise behaviours within the UK population, obesity
5 remains commonplace with 26% of adults documented as having an obese body mass index
6
7 (BMI) in the 2016 Health Survey for England (2). Our understanding of the implications of
8
9 obesity on cardiovascular health, diabetes and other common comorbidities is comprehensive
10
11 and well-publicised. However, there remains uncertainty over the effect of obesity on skeletal
12
13 health, particularly in the context of osteoporosis.
14
15
16
17

18
19 Osteoporosis is a systemic skeletal disorder characterised by compromised bone
20
21 strength, predisposing to an increased risk of fracture (3). Fractures occur when mechanical
22
23 loading exceeds the functional capacity of bone and those resulting from low trauma injuries
24
25 represent the clinical end point of osteoporosis. Of all osteoporotic fractures, those affecting
26
27 the vertebrae are the most common, but vertebral fractures are also the most poorly detected
28
29 and predicted (4, 5). Vertebral fractures are predominantly low trauma injuries, precipitated
30
31 by everyday activities such as bending to lift light objects or falling from standing (5),
32
33 although they can result from high impact forces to the spine (6). They are associated with
34
35 increased morbidity, reduced quality of life and a significant economic burden, costing
36
37 around €1,745 million in Europe in the year 2010 (7). While deformities can present
38
39 clinically as back pain, they are commonly asymptomatic and under-diagnosis is a worldwide
40
41 problem (5).
42
43
44
45
46

47
48 Understanding the risk factors and improving the identification of vertebral fractures
49
50 is important because even mild deformities are associated with an elevated risk of future
51
52 vertebral, and other osteoporotic fractures (8). Epidemiological research indicates that the
53
54 prevalence and number of vertebral fractures increases with age in both sexes (9), and recent
55
56
57
58
59
60
61
62
63
64
65

1 data show that in the United Kingdom, the rate of clinical vertebral fracture is greater in
2 women than in men over the age of 50 years (10).
3

4
5 Body weight is positively associated with bone mineral density (BMD), which is an
6 established measure of bone strength (11). A reduction in body weight results in bone loss
7 (12) and low BMI is an established risk factor for osteoporotic fractures, especially those at
8 the hip (13, 14). As such, low BMI is considered in fracture risk prediction algorithms
9 including FRAX and Garvan, with the highest gradient of risk seen at BMI levels less than 20
10 kg/m² (14). However, few studies have looked at BMD in relation to body weight (6) and
11 there continues to be uncertainty about the relationship between obesity and fracture risk.
12
13
14
15
16
17
18
19
20

21
22 Accumulating evidence indicates that the relationship between BMI and fracture
23 varies according to fracture site. Whereas the risk of hip fracture is known to be highest
24 among underweight individuals (14), fractures of the ankle, lower limb and upper arm have
25 been found to be more common in obese than non-obese or underweight postmenopausal
26 women (15, 16). Clinically diagnosed fractures of the vertebrae have been reported to be
27 lower in obese women (16), but it is known that a large proportion of vertebral fractures
28 remain clinically undetected (17, 18). This relationship is understudied in men, for whom
29 results have been discordant (19, 20).
30
31
32
33
34
35
36
37
38
39

40
41 The purpose of this study was to explore BMD relative to body weight, and the
42 prevalence of vertebral fracture, in a cohort of men and women according to BMI and BMI
43 category.
44
45
46
47
48
49

50 **Materials and Methods**

51 *Participants*

52
53 The research was conducted as a cross-sectional analysis of 342 participants from the
54 Newcastle Thousand Families Study (NTFS) birth cohort. The NTFS was initiated in 1947 by
55
56
57
58
59
60
61
62
63
64
65

1 Sir James Spence, when 1,142 children born in May and June of that year to mothers resident
2 in the city of Newcastle upon Tyne were recruited in response to the high infant mortality
3 rate and poverty levels. The birth cohort has been followed up at regular intervals with a
4 multidisciplinary research focus and the cohort has been described in detail at age 50 years
5 (21). The current study evaluated participants during the most recent NTFS follow up wave
6 in 2011. 37% of those presumed to still be alive at the end of the follow-up period returned
7 for a clinical examination having completed health and lifestyle questionnaires. A favourable
8 ethical opinion was obtained from the Sunderland Local Research Ethics Committee
9 (Reference 09/H0904/40) and all included study members gave their written consent.
10
11
12
13
14
15
16
17
18
19
20
21
22
23

24 *Clinical measurements*

25 Participants were assessed in light-weight clothing with shoes and jewellery removed. Body
26 weight was measured to the nearest 0.1kg using electronic calibrated scales (with heavy
27 clothing and shoes removed). Standing height was measured to the nearest 0.1 cm using a
28 stadiometer (with shoes removed). BMI (kg/m^2) was subsequently calculated [weight
29 (kg)/height(m)²]. BMI scores were categorised according to the WHO criteria as normal
30 weight (18.5-24.99 kg/m^2), overweight (25-29.99 kg/m^2) and obese (>30 kg/m^2) (22). One
31 female subject had been excluded from further analysis as the only participant calculated to
32 have an underweight BMI (<18.5 kg/m^2).
33
34
35
36
37
38
39
40
41
42
43
44

45 Fan beam dual energy X-ray absorptiometry (Lunar iDXA, GE Healthcare, Madison,
46 WI) was used to evaluate left femoral neck BMD, lumbar spine BMD (L1-L4) and prevalent
47 VF. Daily calibration and DXA quality control observations were recorded as per the
48 manufacturer's guidelines for the duration of the data collection, with no equipment drifts or
49 faults reported. Precision error for iDXA measurements are 0.4% coefficient of variation
50 (CV) for lumbar spine BMD and 0.9% CV for femoral neck BMD (23). Patient positioning
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 for the femoral neck BMD scan was assisted using the GE-dual femur positioning device
2 which allows both legs to be abducted and inwardly rotated 25°.
3

4
5 Vertebral fracture prevalence was assessed using a lateral vertebral scan and vertebral
6 fracture assessment (VFA) software (EnCore version 15.0). VFA, as described by the
7 International Society for Clinical Densitometry (ISCD), denotes the use of densitometric
8 spinal imaging for the purpose of identifying VFs (17). The method has been shown to be
9
10 equally as accurate as plain radiography (24) and is regarded as an effective method for
11 diagnosing and reporting VFs (17, 25). DXA scans were performed with participants in the
12 left lateral decubitus position, as recommended by the manufacturer (participants lay on their
13 left side with their knees and hips flexed at a 90° angle and arms flexed with both hands
14 joined together near the head). All scans began at the sacrum, targeting the T4-L4 segment of
15 the spine. Regions of interest and bone edge markers were verified or corrected where
16 required, by a certified clinical densitometrist (ISCD) with specific training in VFA.
17
18 Verification included the observation of correct placement of markers on the vertebral bodies
19 and differentiation between genuine VFs and deformities with other causes such as
20 degenerative disease. Measurements were compared with those of other vertebrae, allowing
21 the software to estimate the extent of any reduction in the anterior, middle or posterior
22 vertebral height. VFs were subsequently graded according to their severity as grade 1 (mild),
23 grade 2 (moderate) and grade 3 (severe) using the method set out by Genant et al (26). Grade
24 1 corresponds to a 20 to 25% reduction in any vertebral height; grade 2 a 25 to 40% reduction
25 in vertebral height and grade 3 represents a >40% reduction in vertebral height.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52

53 *FRAX scores*

54
55
56 In addition to age, sex and BMI, individual FRAX scores were included in the analyses (27) .
57
58 Information on previous fracture, parent hip fracture, current smoking, rheumatoid arthritis,
59
60
61
62
63
64
65

1 glucocorticoid use, secondary osteoporosis, and alcohol consumption, were self-recorded by
2 participants in the NTFS general health questionnaire. FRAX scores indicating the 10 year
3 probability of a major osteoporotic fracture were subsequently calculated from femoral neck
4 BMD and the aforementioned clinical risk factors.
5
6
7

8 9 10 11 *Statistical analysis*

12 The data were analysed using STATA software, version 12.0 (StataCorp, College Station,
13 TX). Ratios were created to express lumbar spine BMD as well as femoral neck BMD
14 relative to participant body weight as follows:
15
16
17
18
19

$$20 \quad [BMD / \text{body weight (kg)}] \times 100$$

21 Descriptive statistics including the mean and standard deviation (SD) were produced for all
22 measurements. Logistic regression was used to determine the associations of sex, BMI and
23 BMD / weight with the prevalence of VF, producing estimated odds ratios (OR). All models
24 were sex-specific and adjusted for lumbar spine BMD. Moderate and severe fractures (grade
25 2-3) are more reliable indicators of underlying disease than mild fractures (grade 1) when
26 using VFA (28). Hence the presence of a moderate or severe fracture was analysed
27 separately. Independent samples t-tests were used to determine whether differences in
28 anthropometric characteristics of those with and without fractures were significant.
29
30
31
32
33
34
35
36
37
38
39
40
41

42 Spearman's and Pearson's correlative analyses were used to examine associations between
43 BMI, BMD, FRAX scores, BMD/weight ratios and VF. One-way ANOVA models with post-
44 hoc Tukey analysis were used to determine significant differences in the BMD/weight ratios
45 of men and women of differing BMI categories. Statistical significance was set at $p = 0.05$.
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Results

The mean age of the cohort at the time of testing was 62.5 ± 0.5 years. There were fewer male (44% v 51%) and more female (49% v 56%) participants in the current sample compared to the original 1947 cohort ($p = 0.008$). More than half of the women were using hormone replacement therapy (HRT) (60%), as has previously been reported in U.K. cohorts (29). History of previously diagnosed fracture was reported by 15.8% of men and 16.8% of women. Six men and five women had previously had a clinically diagnosed vertebral fracture. Two men and one woman had previously suffered a hip fracture. Sex-specific anthropometric and bone density results are given in Table 1.

Table 1. Anthropometric and bone mineral density (BMD) summaries for men and women aged 62 years

	Women	Men	<i>p</i>
	n = 190	n = 152	
	(<i>mean</i> \pm <i>SD</i>)	(<i>mean</i> \pm <i>SD</i>)	
Body weight (kg)	73.8 ± 15.1	84.3 ± 15.2	< 0.001
Height (cm)	162.1 ± 6.9	173.5 ± 7.1	< 0.001
Body mass index (kg/m ²)	28.0 ± 5.3	28.0 ± 4.7	0.929
Femoral neck BMD (g/cm ²)	0.927 ± 0.143	0.944 ± 0.139	0.238
Lumbar spine (L1-L4) BMD (g/cm ²)	1.112 ± 0.180	1.204 ± 0.160	< 0.001
Femoral neck BMD / weight (g/cm ² : kg)	1.29 ± 0.27	1.15 ± 0.26	< 0.001
Lumbar spine BMD / weight (g/cm ² : kg)	1.55 ± 0.32	1.47 ± 0.30	0.015
FRAX Major Osteoporotic Fracture	8.2 ± 3.83	6.74 ± 3.83	< 0.001

Bone mineral density

Women had lower lumbar spine BMD than men, although men had lower lumbar spine BMD/ weight (Table 1). In men and women, BMI was negatively associated with femoral

1 neck BMD/ weight ($R = - 0.65$, $R = - 0.66$, $p < 0.001$) and lumbar spine BMD/ weight ($R = -$
2 0.66 , $R = - 0.60$, $p < 0.001$). Scatter graphs illustrating the association between BMD/weight
3 and BMI in men and women are shown in Figure 1.
4

5
6
7 Lumbar spine BMD/ weight and femoral neck BMD/ weight of men and women by
8
9 BMI category are given in Table 2. Both lumbar spine BMD/ weight ($F=37.78$, $p < 0.001$)
10 and femoral neck BMD/ weight ($F=34.93$, $p < 0.001$) varied across BMI category in men.
11
12 Lumbar spine BMD/ weight ($F=39.16$, $p < 0.001$) and femoral neck BMD/ weight ($F=56.44$,
13 $p < 0.001$) also varied across BMI category in women. Lumbar spine BMD/ weight and
14
15 femoral neck BMD/ weight were significantly lower in overweight and obese men and
16
17 women compared to those of normal weight (all $p < 0.001$). Lumbar spine and femoral neck
18
19 BMD/ weight were also lower in men and women who were obese compared to those who
20
21 were overweight (all $p < 0.001$).
22
23
24
25
26
27

28 29 30 31 *Sex-specific vertebral fracture prevalence* 32

33
34 The prevalence of any-grade VF was 48.0% in men and 25.3% in women. Men were more
35
36 than twice as likely to have at least one prevalent VF ($OR = 2.7$, $p < 0.001$, 95% CI 1.7– 4.4).
37

38
39 The prevalence of grade 1 vertebral deformities was higher in men compared to women
40
41 (38.2% v 19.5%). Men were also more than three times as likely to have at least one grade 2-
42
43 3 VF ($OR = 3.5$, $p < 0.001$, 95% CI 1.9 – 6.5). The prevalence of grade 2 and/or grade 3 VF
44
45 was 27.0% ($n = 41$) in men and 10.0% ($n = 19$) in women.
46

47
48
49 The prevalence of any-grade, grade 1 and grade 2 / 3 VFs in men and women at
50
51 each BMI category are given in Table 2. BMI was associated with an increased likelihood of
52
53 prevalent any-grade VF in women ($OR = 1.09$, $p = 0.006$, 95% CI 1.02 – 1.17). Compared to
54
55 normal weight women, obese women were more likely to have at least one VF ($OR = 2.65$, p
56
57 $= 0.025$) and at least one grade 1 VF ($OR = 4.39$, $p = 0.005$). No association between BMI
58
59
60
61
62
63
64
65

1 and grade 2 and/or 3 VF was identified in women (Table 3). Women with VFs had higher
2 weight ($p=0.002$) and BMI ($p<0.001$) than those without (Table 4). No anthropometric
3 differences were identified between women with and without grade 2 and/or 3 VFs.
4

5
6
7 In men, a higher BMI increased the likelihood of a prevalent grade 2 and/or 3 VF
8 (OR = 1.08, $p = 0.054$, 95% CI 1.00 – 1.17), and men in the obese category were more likely
9 to have a grade 2 and/or 3 VF (OR = 3.36, $p = 0.032$) (Table 3). No significant differences
10 were identified in the anthropometric characteristics of men with and without any grade VF
11 or grade 2 and/or 3 VF (Table 4).
12
13
14
15
16
17

18
19 In women, femoral neck BMD/ weight was negatively associated with prevalent
20 grade 1 vertebral deformity only (OR 0.11, $p=0.004$). There appeared to be a relationship
21 between LS BMD/ weight and grade 1 VF prevalence in women (OR 0.04, $p=0.080$),
22 although the association was not statistically significant. No association between lumbar
23 spine or femoral BMD/ weight and the likelihood of prevalent VF was identified in men.
24
25
26
27
28
29

30
31 In women, the number of prevalent VFs was associated with increasing body
32 weight ($R = 0.25$, $p < 0.001$), BMI ($R = 0.22$, $p < 0.001$), lumbar spine BMD ($R = 0.15$, $p =$
33 0.042) and femoral neck BMD/ weight ($R = - 0.15$, $p = 0.04$). The number of grade 1, but not
34 grade 2 / 3, deformities were positively associated with body weight ($R = 0.28$, $p < 0.001$),
35 BMI ($R = 0.26$, $p < 0.001$), lumbar spine BMD ($R = 0.18$, $p = 0.013$), and femoral neck
36 BMD/weight ($R = - 0.22$, $p = 0.003$). In men, the number of grade 1 vertebral deformities
37 were associated with higher body weight ($R = 0.17$, $p = 0.042$).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Table 2. Bone mineral density (BMD)/ weight and prevalent vertebral fracture across BMI categories in men and women aged 62 years.

	Women			Men		
	Normal weight n = 64	Overweight n = 68	Obese n = 58	Normal weight n = 35	Overweight n = 74	Obese n = 43
	BMD (mean ± SD)					
Femoral neck BMD			†			†
(g/cm ²)	0.911 ± 0.136	0.903 ± 0.145	0.973 ± 0.140	0.919 ± 0.151	0.959 ± 0.137	0.936 ± 0.133
Lumbar spine BMD	1.074 ± 0.168 [†]	1.073 ± 0.164 [†]	1.198 ± 0.185 [†]	1.166 ± 0.147	1.221 ± 0.173	1.206 ± 0.141
(g/cm ²)						
	BMD : Weight Ratio (mean ± SD)					
Femoral neck BMD / weight	1.50 ± 0.21*	1.27 ± 0.23*	1.09 ± 0.19*	1.36 ± 0.32*	1.17 ± 0.19*	0.93 ± 0.13*
(g/cm ² : kg)						
Lumbar spine BMD / weight	1.77 ± 0.25*	1.52 ± 0.30*	1.35 ± 0.23*	1.71 ± 0.29*	1.50 ± 0.26*	1.20 ± 0.16*
(g/cm ² : kg)						
	Number (%) with prevalent vertebral fracture (≥1)					
Any grade VF	12 (18.8%)	13 (19.1%)	23 (39.7%)	14 (40%)	37 (50%)	22 (51.2%)
Grade 1 VF	6 (9.4%)	11 (16.2%)	20 (34.5%)	12 (34.3%)	27 (36.5%)	19 (44.2%)
Grades 2 and/or 3 VF	8 (12.5%)	4 (5.9%)	7 (12.1%)	7 (20%)	18 (24.3%)	16 (37.2%)

15
16
17
18
19
28
38
59

† indicates significant variance between two groups i.e. 1 vs. 2 ($p < 0.05$).

* indicates significant variation between all groups i.e. 1 vs. 2, 1 vs. 3, 2 vs. 3 ($p < 0.05$).

60
61
62
63
64
65

16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
58
59
60
61
62
63
64
65

Table 3. Sex-specific odds ratios for the likelihood of prevalent vertebral fracture (VF) by BMI category

	Any grade VF		Grade 1		Grade 2 and/or 3	
	Women	Men	Women	Men	Women	Men
Odds ratio – Likelihood of prevalent VF vs. normal weight						
<i>OR (95% Confidence interval)</i>						
Overweight (25.0 – 29.99 kg/m ²)	1.0 (0.43 – 2.50)	1.49 (0.64 – 3.48)	1.9 (0.66 – 5.52)	0.99 (0.42 – 2.36)	0.44 (0.13 – 1.56)	1.45 (0.51 – 4.12)
Obese (30.0 kg/m ² or above)	2.65* (1.13 – 6.20)	1.91 (0.73 – 4.99)	4.39* (1.57 – 12.28)	1.59 (0.61 – 4.20)	0.93 (0.30 – 2.89)	3.36* (1.11 – 10.16)

* indicates result significant vs. normal weight reference group (p < 0.05).

15
16
17
18
19
20





58
59
60
61
62
63
64
65

Table 4. Descriptive results for men and women with and without any grade and grade 2 and/or 3 VFs.

	Women				Men			
	No VF n = 142	Any grade VF n = 48	No grade 2 or 3 VF n = 171	Grade 2 or 3 VF n = 19	No VF n = 78	Any grade VF n = 74	No grade 2 or 3 VF n = 111	Grade 2 or 3 VF n = 41
Height (cm)	162.2 ± 6.7	161.9 ± 7.6	162.3 ± 7.0	160.6 ± 6.5	173.0 ± 7.6	174.0 ± 6.7	173.5 ± 7.4	173.5 ± 6.5
Body weight (kg)	71.8 ± 13.8*	79.7 ± 17.2*	73.5 ± 14.9	76.7 ± 16.9	82.8 ± 15.6	86.0 ± 14.8	83.3 ± 15.08	87.1 ± 15.5
BMI (kg/m ²)	27.3 ± 4.7*	30.1 ± 6.3*	27.8 ± 5.0	29.2 ± 7.6	27.7 ± 5.0	28.4 ± 4.4	27.7 ± 4.71	28.9 ± 4.63
Femoral neck BMD (g/cm ²)	0.922 ± 0.144	0.945 ± 0.139	0.923 ± 0.142	0.967 ± 0.148	0.946 ± 0.155	0.942 ± 0.123	0.950 ± 0.146	0.929 ± 0.119
Lumbar spine BMD (g/cm ²)	1.100 ± 0.181	1.147 ± 0.176	1.111 ± 0.182	1.127 ± 0.163	1.201 ± 0.165	1.209 ± 0.156	1.207 ± 0.164	1.200 ± 0.151
FRAX Major Osteoporotic Fracture	8.25 ± 3.96	8.02 ± 3.46	8.19 ± 3.89	8.27 ± 3.27	7.04 ± 4.23	6.41 ± 3.34	6.78 ± 3.92	6.64 ± 3.64
Femoral neck BMD /weight (g/cm ² : kg)	1.31 ± 0.26	1.23 ± 0.29	1.29 ± 0.26	1.31 ± 0.32	1.18 ± 0.30	1.12 ± 0.22	1.17 ± 0.27	1.09 ± 0.24
Lumbar spine BMD /weight (g/cm ² : kg)	1.57 ± 0.31	1.50 ± 0.33	1.55 ± 0.32	1.54 ± 0.34	1.50 ± 0.32	1.44 ± 0.28	1.49 ± 0.31	1.41 ± 0.29

* indicates significant difference in mean values (p<0.05).

16
17
18
19
20
21
22 **Figure 1.** Bone mineral density (BMD)/ weight and BMI in men and women aged 62 years.



Discussion

The aim of this study was to investigate associations between BMI, BMD relative to body weight and prevalent VF in men and women aged 62 years. It is well-established that low BMI increases the risk of osteoporosis and fracture, but the skeletal risks associated with obesity are unclear. Our primary finding was that obese men and women have a greater likelihood of prevalent VF than their non-obese counterparts. Additionally, our results indicate that BMD is reduced in obese individuals, when considered relatively to body weight, which may contribute to VF formation in this group.

In agreement with reports elsewhere, the prevalence and number of VFs were greater in men than women (19). The overall prevalence of VF we observed was comparable to previous cohorts of men, although lower in women which may reflect an age-effect (25.3% vs. 37% by Laslett *et al.*) (15). The majority of VFs in women were grade 1 deformities, which were positively associated with BMI. Although grade 2/3 VFs reflect worsening architectural deterioration of bone (30, 31), grade 1 deformities have been proven to be important predictors of future VF (8). No associations between obesity and grade 2/3 VF were identified in women. However, we observed a relatively low number of grade 2/3 VFs in women and the study may have lacked the power to identify any potential relationship.

To date, there have been few studies examining risk factors for VF in older men, and findings have been conflicting (19, 20, 32-34). In the current study, we found that a high BMI increased the likelihood of prevalent grade 2/3 VF in men, with obese men three times more likely than normal weight men to have at least one grade 2/3 VF. Results from large, prospective cohorts examining obesity and VF in women have also varied (35-37). Compston *et al.* reported that high BMI was associated with lower rates of clinically reported incident VF (35). However, using radiographs, Tanaka *et al.* found a higher risk of incident VF in

1 overweight and obese women compared to normal weight women (37). We observed that
2 women with a higher BMI had an increased likelihood of any grade prevalent VF,
3
4 corroborating the findings of both Laslett *et al.* and Pirro *et al.* (11, 19), and that obese
5 women were more than twice as likely to have a prevalent VF. In agreement with previous
6
7 studies, we also found that the number of VFs were positively associated with BMI in women
8
9 (11, 19).
10
11
12

13
14 Women and men with an overweight BMI were not found to be significantly more
15
16 likely than normal weight counterparts to have prevalent VFs. This could indicate that the
17
18 fracture risk is greatest in underweight and obese individuals, whilst overweight individuals
19
20 are predisposed to VF to a lesser extent. At present this is speculative and longitudinal
21
22 analysis of VF incidence would be required to establish the gradient of risk across all four
23
24 BMI categories.
25
26
27
28

29
30 There have been a number of proposed explanations for the increased risk of VF with
31
32 obesity. It is possible that vertebral injuries sustained earlier in life, perhaps as a result of
33
34 occupational trauma, may become aggravated by the effects of obesity as age increases,
35
36 although this cannot be deduced from the current cross-sectional analysis. Postural stability is
37
38 reduced in obese individuals (38), which brings an increased risk of falling (39, 40), and
39
40 heavier individuals will generate greater forces on impact. VFs can also be low-trauma
41
42 injuries that develop insidiously as a result of sustained compressive forces on the vertebrae,
43
44 which are of the greater magnitude in the obese individual (41). In addition, obesity is
45
46 associated with total and visceral adiposity, and adipocyte-derived hormones and pro-
47
48 inflammatory cytokines can negatively impact bone metabolism (42).
49
50
51
52

53
54 The most pertinent explanation may be that the relationship between BMI and BMD
55
56 is non-linear, with increased in BMD in obese individuals not commensurate to increased
57
58
59
60
61
62
63
64
65

1 body mass. It is well established that BMD is higher in normal weight individuals, when
2 compared to underweight, conferring a protective advantage in the context of fracture risk
3
4 (14). However, our results would suggest this trend does not continue as BMI moves into the
5 overweight and obesity categories. We observed no significant variation in absolute BMD
6
7 between males of differing BMI group, and no increase between normal weight and
8
9 overweight women. Furthermore, when considered relative to body weight, we observed
10
11 that BMD was significantly lower in overweight and obese men and women. These results
12
13 add to a growing body of evidence highlighting that bone density does not increase
14
15 proportionally with body mass in obesity. Sornary Rendu *et al.* observed various measures of
16
17 bone strength to be relatively lower in obese postmenopausal women (43), whilst Beck et al
18
19 found that femoral BMD had scaled upwards in proportion with increases in lean tissue mass,
20
21 rather than increases in fat mass or total body mass (44). Lower femoral neck BMD/weight
22
23 was associated with an increased likelihood prevalent VF in women, but not men. We
24
25 propose that examining BMD relative to body weight is a useful measure when exploring
26
27 bone fragility in obese populations and further research on the clinical utility of this index is
28
29 warranted.
30
31
32
33
34
35
36
37
38
39

40 Previous studies validating the use of FRAX in predicting future VF reported that
41
42 femoral neck BMD, age and self reported fracture history were the main contributing factors
43
44 in generating 10-year risk score (45). The narrow age-range of this cohort, in combination
45
46 with adjustment for lumbar spine BMD in regression models, is therefore likely to have
47
48 limited the ability of the FRAX tool to discern high- from low-risk individuals. Further
49
50 analysis of the FRAX scoring in longitudinal studies will be required to further assess its role
51
52 in predicting VF.
53
54

55
56
57 There are a number of limitations to our study. First, we were unable to assess time
58
59 since the fracture or antecedent factors relating to vertebral fracture and are unable to make
60
61
62
63
64
65

1 any inferences about cause and effect relationships. Longitudinal investigations are needed
2 to address causality. Second, this cohort was, on average, overweight and therefore some
3 results may not be generalisable to populations with a lower prevalence of obesity. Finally,
4 caution should be taken when making inferences given that the study was performed among
5 members of a birth cohort born in Newcastle upon Tyne, UK who were aged 61 - 63 years.
6
7 However, this age group has particular public health importance because the risk of fracture
8 is increased compared to younger ages, and effective risk factor modification (e.g. exercise
9 and diet for fat loss) is still likely to be viable.

10
11 In conclusion, obesity appears to increase the likelihood of prevalent VF in both men
12 and women, and we observed an increased prevalence of VF in men from this cohort. BMD
13 was relatively low in obese individuals, when expressed relative to body weight, and
14 therefore although absolute BMD increases in obesity, this may not be commensurate to
15 increased body mass and fat mass. Given the rising global rates of obesity (1), the association
16 between BMD/weight and vertebral fracture risk warrants further and timely investigation.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Acknowledgements

1
2 All authors were involved with the preparation of the manuscript and interpretation of the
3
4 data. HR led the preparation of the manuscript and statistical analysis. KH led the concept,
5
6 MP and FB directed the data collection. MP is the lead for the Newcastle Thousand Families
7
8 Study. We would like to thank the previous research teams involved in the Newcastle
9
10 Thousand Families Study and the study members for taking part in the investigation. We are
11
12 grateful to previous funders for supporting the research and to JGW Patterson Foundation, the
13
14 NIHR BCC. Thanks also to Katherine Kirton and Emma Thompson for their excellent
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

References

1. Organization, W.H. *Obesity and overweight - factsheet*. [Online]. 2016. [Accessed]. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en/>
2. Barker, C. *Obesity statistics*. [Online]. 2018. [Accessed]. Available from: <https://researchbriefings.parliament.uk/ResearchBriefing/Summary/SN03336>
3. NIH Consensus Development Panel on Osteoporosis Prevention, D. and Therapy. Osteoporosis prevention, diagnosis, and therapy. *JAMA*. 2001, **285**(6), pp.785-795.
4. Ensrud, K.E. and Schousboe, J.T. Clinical practice. Vertebral fractures. *N Engl J Med*. 2011, **364**(17), pp.1634-1642.
5. Cummings, S.R. and Melton, L.J. Epidemiology and outcomes of osteoporotic fractures. *Lancet*. 2002, **359**(9319), pp.1761-1767.
6. Hind, K., Gannon, L., Brightmore, A. and Beck, B. Insights into relationships between body mass, composition and bone: findings in elite rugby players. *J Clin Densitom*. 2015, **18**(2), pp.172-178.
7. Hernlund, E., Svedbom, A., Ivergard, M., Compston, J., Cooper, C., Stenmark, J., McCloskey, E.V., Jonsson, B. and Kanis, J.A. Osteoporosis in the European Union: medical management, epidemiology and economic burden. A report prepared in collaboration with the International Osteoporosis Foundation (IOF) and the European Federation of Pharmaceutical Industry Associations (EFPIA). *Arch Osteoporos*. 2013, **8**, p.136.
8. Roux, C., Fechtenbaum, J., Kolta, S., Briot, K. and Girard, M. Mild prevalent and incident vertebral fractures are risk factors for new fractures. *Osteoporos Int*. 2007, **18**(12), pp.1617-1624.
9. European Prospective Osteoporosis Study, G., Felsenberg, D., Silman, A.J., Lunt, M., Armbrecht, G., Ismail, A.A., Finn, J.D., Cockerill, W.C., Banzer, D., Benevolenskaya, L.I., Bhalla, A., Bruges Armas, J., Cannata, J.B., Cooper, C., Dequeker, J., Eastell, R., Felsch, B., Gowin, W., Havelka, S., Hozzowski, K., Jajic, I., Janott, J., Johnell, O., Kanis, J.A., Kragl, G., Lopes Vaz, A., Lorenc, R., Lyritis, G., Masaryk, P., Matthis, C., Miazgowski, T., Parisi, G., Pols, H.A., Poor, G., Raspe, H.H., Reid, D.M., Reisinger, W., Schedit-Nave, C., Stepan, J.J., Todd, C.J., Weber, K., Woolf, A.D., Yershova, O.B., Reeve, J. and O'Neill, T.W. Incidence of vertebral fracture in europe: results from the European Prospective Osteoporosis Study (EPOS). *J Bone Miner Res*. 2002, **17**(4), pp.716-724.
10. Curtis, E.M., van der Velde, R., Moon, R.J., van den Bergh, J.P., Geusens, P., de Vries, F., van Staa, T.P., Cooper, C. and Harvey, N.C. Epidemiology of fractures in the United Kingdom 1988-2012: Variation with age, sex, geography, ethnicity and socioeconomic status. *Bone*. 2016, **87**, pp.19-26.
11. Pirro, M., Fabbriani, G., Leli, C., Callarelli, L., Manfredelli, M.R., Fioroni, C., Mannarino, M.R., Scarponi, A.M. and Mannarino, E. High weight or body mass index increase the risk of vertebral fractures in postmenopausal osteoporotic women. *J Bone Miner Metab*. 2010, **28**(1), pp.88-93.
12. Felson, D.T., Zhang, Y., Hannan, M.T. and Anderson, J.J. Effects of weight and body mass index on bone mineral density in men and women: the Framingham study. *J Bone Miner Res*. 1993, **8**(5), pp.567-573.
13. Johnell, O., Gullberg, B., Kanis, J.A., Allander, E., Elffors, L., Dequeker, J., Dilsen, G., Gennari, C., Lopes Vaz, A., Lyritis, G. and et al. Risk factors for hip fracture in

- 1 European women: the MEDOS Study. Mediterranean Osteoporosis Study. *J Bone*
2 *Miner Res.* 1995, **10**(11), pp.1802-1815.
- 3 14. De Laet, C., Kanis, J.A., Oden, A., Johanson, H., Johnell, O., Delmas, P., Eisman, J.A.,
4 Kroger, H., Fujiwara, S., Garnero, P., McCloskey, E.V., Mellstrom, D., Melton, L.J., 3rd,
5 Meunier, P.J., Pols, H.A., Reeve, J., Silman, A. and Tenenhouse, A. Body mass index as
6 a predictor of fracture risk: a meta-analysis. *Osteoporos Int.* 2005, **16**(11), pp.1330-
7 1338.
- 8 15. Armstrong, M.E., Cairns, B.J., Banks, E., Green, J., Reeves, G.K., Beral, V. and Million
9 Women Study, C. Different effects of age, adiposity and physical activity on the risk
10 of ankle, wrist and hip fractures in postmenopausal women. *Bone.* 2012, **50**(6),
11 pp.1394-1400.
- 12 16. Compston, J.E., Watts, N.B., Chapurlat, R., Cooper, C., Boonen, S., Greenspan, S.,
13 Pfeilschifter, J., Silverman, S., Diez-Perez, A., Lindsay, R., Saag, K.G., Netelenbos, J.C.,
14 Gehlbach, S., Hooven, F.H., Flahive, J., Adachi, J.D., Rossini, M., Lacroix, A.Z., Roux, C.,
15 Sambrook, P.N., Siris, E.S. and Glow, I. Obesity is not protective against fracture in
16 postmenopausal women: GLOW. *Am J Med.* 2011, **124**(11), pp.1043-1050.
- 17 17. Schousboe, J.T., Vokes, T., Broy, S.B., Ferrar, L., McKiernan, F., Roux, C. and Binkley,
18 N. Vertebral Fracture Assessment: the 2007 ISCD Official Positions. *J Clin Densitom.*
19 2008, **11**(1), pp.92-108.
- 20 18. Ensrud, K.E., Blackwell, T.L., Fink, H.A., Zhang, J., Cauley, J.A., Cawthon, P.M., Black,
21 D.M., Bauer, D.C., Curtis, J.R., Orwoll, E.S., Barrett-Connor, E., Kado, D.M., Marshall,
22 L.M., Shikany, J.M., Schousboe, J.T. and Osteoporotic Fractures in Men Research, G.
23 What Proportion of Incident Radiographic Vertebral Fractures in Older Men Is
24 Clinically Diagnosed and Vice Versa: A Prospective Study. *J Bone Miner Res.* 2016,
25 **31**(8), pp.1500-1503.
- 26 19. Laslett, L.L., Just Nee Foley, S.J., Quinn, S.J., Winzenberg, T.M. and Jones, G. Excess
27 body fat is associated with higher risk of vertebral deformities in older women but
28 not in men: a cross-sectional study. *Osteoporos Int.* 2012, **23**(1), pp.67-74.
- 29 20. Kanazawa, I., Yamaguchi, T., Yamamoto, M., Yamauchi, M., Yano, S. and Sugimoto, T.
30 Combination of obesity with hyperglycemia is a risk factor for the presence of
31 vertebral fractures in type 2 diabetic men. *Calcif Tissue Int.* 2008, **83**(5), pp.324-331.
- 32 21. Pearce, M.S., Unwin, N.C., Parker, L. and Craft, A.W. Cohort profile: the Newcastle
33 Thousand Families 1947 birth cohort. *Int J Epidemiol.* 2009, **38**(4), pp.932-937.
- 34 22. Organization, W.H. *WHO BMI classification*. [Online]. [Accessed 17/4/17]. Available
35 from: http://apps.who.int/bmi/index.jsp?introPage=intro_3.html.
- 36 23. Hind, K., Oldroyd, B. and Truscott, J.G. In vivo precision of the GE Lunar iDXA
37 densitometer for the measurement of total-body, lumbar spine, and femoral bone
38 mineral density in adults. *J Clin Densitom.* 2010, **13**(4), pp.413-417.
- 39 24. Diacinti, D., Del Fiacco, R., Pisani, D., Todde, F., Cattaruzza, M.S., Diacinti, D., Arima,
40 S., Romagnoli, E., Pepe, J., Cipriani, C. and Minisola, S. Diagnostic performance of
41 vertebral fracture assessment by the lunar iDXA scanner compared to conventional
42 radiography. *Calcif Tissue Int.* 2012, **91**(5), pp.335-342.
- 43 25. Genant, H.K., Li, J., Wu, C.Y. and Shepherd, J.A. Vertebral fractures in osteoporosis: a
44 new method for clinical assessment. *J Clin Densitom.* 2000, **3**(3), pp.281-290.
- 45 26. Genant, H.K., Wu, C.Y., van Kuijk, C. and Nevitt, M.C. Vertebral fracture assessment
46 using a semiquantitative technique. *J Bone Miner Res.* 1993, **8**(9), pp.1137-1148.
- 47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
27. Kanis, J.A., Johnell, O., Oden, A., Johansson, H. and McCloskey, E. FRAX and the assessment of fracture probability in men and women from the UK. *Osteoporos Int.* 2008, **19**(4), pp.385-397.
 28. Genant, H.K., Delmas, P.D., Chen, P., Jiang, Y., Eriksen, E.F., Dalsky, G.P., Marcus, R. and San Martin, J. Severity of vertebral fracture reflects deterioration of bone microarchitecture. *Osteoporos Int.* 2007, **18**(1), pp.69-76.
 29. Cumming, G.P., Currie, H., Morris, E., Moncur, R. and Lee, A.J. The need to do better - Are we still letting our patients down and at what cost? *Post Reprod Health.* 2015, **21**(2), pp.56-62.
 30. Sornay-Rendu, E., Cabrera-Bravo, J.L., Boutroy, S., Munoz, F. and Delmas, P.D. Severity of vertebral fractures is associated with alterations of cortical architecture in postmenopausal women. *J Bone Miner Res.* 2009, **24**(4), pp.737-743.
 31. Delmas, P.D., Genant, H.K., Crans, G.G., Stock, J.L., Wong, M., Siris, E. and Adachi, J.D. Severity of prevalent vertebral fractures and the risk of subsequent vertebral and nonvertebral fractures: results from the MORE trial. *Bone.* 2003, **33**(4), pp.522-532.
 32. Szulc, P., Varennes, A., Delmas, P.D., Goudable, J. and Chapurlat, R. Men with metabolic syndrome have lower bone mineral density but lower fracture risk--the MINOS study. *J Bone Miner Res.* 2010, **25**(6), pp.1446-1454.
 33. Hind, K., Pearce, M. and Birrell, F. Total and Visceral Adiposity Are Associated With Prevalent Vertebral Fracture in Women but Not Men at Age 62 Years: The Newcastle Thousand Families Study. *J Bone Miner Res.* 2017, **32**(5), pp.1109-1115.
 34. Scane, A.C., Francis, R.M., Sutcliffe, A.M., Francis, M.J., Rawlings, D.J. and Chapple, C.L. Case-control study of the pathogenesis and sequelae of symptomatic vertebral fractures in men. *Osteoporos Int.* 1999, **9**(1), pp.91-97.
 35. Compston, J.E., Flahive, J., Hosmer, D.W., Watts, N.B., Siris, E.S., Silverman, S., Saag, K.G., Roux, C., Rossini, M., Pfeilschifter, J., Nieves, J.W., Netelenbos, J.C., March, L., LaCroix, A.Z., Hooven, F.H., Greenspan, S.L., Gehlbach, S.H., Diez-Perez, A., Cooper, C., Chapurlat, R.D., Boonen, S., Anderson, F.A., Jr., Adami, S., Adachi, J.D. and Investigators, G. Relationship of weight, height, and body mass index with fracture risk at different sites in postmenopausal women: the Global Longitudinal study of Osteoporosis in Women (GLOW). *J Bone Miner Res.* 2014, **29**(2), pp.487-493.
 36. Holmberg, A.H., Johnell, O., Nilsson, P.M., Nilsson, J., Berglund, G. and Akesson, K. Risk factors for fragility fracture in middle age. A prospective population-based study of 33,000 men and women. *Osteoporos Int.* 2006, **17**(7), pp.1065-1077.
 37. Tanaka, S., Kuroda, T., Saito, M. and Shiraki, M. Overweight/obesity and underweight are both risk factors for osteoporotic fractures at different sites in Japanese postmenopausal women. *Osteoporos Int.* 2013, **24**(1), pp.69-76.
 38. Teasdale, N., Simoneau, M., Corbeil, P., Handrigan, G., Tremblay, A. and Hue, O. Obesity Alters Balance and Movement Control. *Current Obesity Reports.* 2013, **2**(3), pp.235-240.
 39. Hue, O., Simoneau, M., Marcotte, J., Berrigan, F., Dore, J., Marceau, P., Marceau, S., Tremblay, A. and Teasdale, N. Body weight is a strong predictor of postural stability. *Gait & Posture.* 2007, **26**(1), pp.32-38.
 40. Corbeil, P., Simoneau, M., Rancourt, D., Tremblay, A. and Teasdale, N. Increased risk for falling associated with obesity: mathematical modeling of postural control. *IEEE Trans Neural Syst Rehabil Eng.* 2001, **9**(2), pp.126-136.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
41. Pollintine, P., Luo, J., Offa-Jones, B., Dolan, P. and Adams, M.A. Bone creep can cause progressive vertebral deformity. *Bone*. 2009, **45**(3), pp.466-472.
 42. Zhao, L.J., Jiang, H., Papasian, C.J., Maulik, D., Drees, B., Hamilton, J. and Deng, H.W. Correlation of obesity and osteoporosis: effect of fat mass on the determination of osteoporosis. *J Bone Miner Res*. 2008, **23**(1), pp.17-29.
 43. Sornay-Rendu, E., Boutroy, S., Vilayphiou, N., Claustrat, B. and Chapurlat, R.D. In obese postmenopausal women, bone microarchitecture and strength are not commensurate to greater body weight: the Os des Femmes de Lyon (OFELY) study. *J Bone Miner Res*. 2013, **28**(7), pp.1679-1687.
 44. Beck, T.J., Petit, M.A., Wu, G., LeBoff, M.S., Cauley, J.A. and Chen, Z. Does obesity really make the femur stronger? BMD, geometry, and fracture incidence in the women's health initiative-observational study. *J Bone Miner Res*. 2009, **24**(8), pp.1369-1379.
 45. Donaldson, M.G., Palermo, L., Schousboe, J.T., Ensrud, K.E., Hochberg, M.C. and Cummings, S.R. FRAX and risk of vertebral fractures: the fracture intervention trial. *J Bone Miner Res*. 2009, **24**(11), pp.1793-1799.

