

www.ams.org

David Cimasoni, Anthony Conway, Kleopatra Zacharova Splitting numbers and signatures Proceedings of the American Mathematical Society DOI: 10.1090/proc/13156

Accepted Manuscript

This is a preliminary PDF of the author-produced manuscript that has been peer-reviewed and accepted for publication. It has not been copyedited, proofread, or finalized by AMS Production staff. Once the accepted manuscript has been copyedited, proofread, and finalized by AMS Production staff, the article will be published in electronic form as a "Recently Published Article" before being placed in an issue. That electronically published article will become the Version of Record.

This preliminary version is available to AMS members prior to publication of the Version of Record, and in limited cases it is also made accessible to everyone one year after the publication date of the Version of Record.

The Version of Record is accessible to everyone five years after publication in an issue.

SPLITTING NUMBERS AND SIGNATURES

DAVID CIMASONI, ANTHONY CONWAY, AND KLEOPATRA ZACHAROVA

ABSTRACT. The splitting number of a link is the minimal number of crossing changes between different components required to convert it into a split link. We obtain a lower bound on the splitting number in terms of the (multivariable) signature and nullity. Although very elementary and easy to compute, this bound turns out to be suprisingly efficient. In particular, it makes it a routine check to recover the splitting number of 129 out of the 130 prime links with at most 9 crossings. Also, we easily determine 16 of the 17 splitting numbers that were studied by Batson and Seed using Khovanov homology, and later computed by Cha, Friedl and Powell using a variety of techniques. Finally, we determine the splitting number of a large class of 2-bridge links which includes examples recently computed by Borodzik and Gorsky using a Heegaard Floer theoretical criterion.

1. INTRODUCTION

Any link $L = L_1 \cup \cdots \cup L_{\mu}$ in S^3 can be turned into the split union of its components by a sequence of crossing changes between different components. Following Batson and Seed [2] and Cha, Friedl and Powell [7], we call *splitting number* of L the minimal number of crossing changes in such a sequence, and denote it by sp(L). (Note that the same terminology is used in [1, 19, 23] for another invariant.) Obviously, upper bounds on sp(L) can be found by inspection of diagrams, so the difficulty in computing it is to find lower bounds.

As observed in [2] (see also [7, Lemma 2.1]), the linking numbers provide an elementary such bound. First note that $\operatorname{sp}(L)$ has the same parity as the total linking number $\sum_{i < j} \ell k(L_i, L_j)$. Furthermore, given a two component link $L_i \cup L_j$, let $b_{\ell k}(L_i, L_j)$ be equal to 0 if $L_i \cup L_j$ is split, to 2 if it is non-split but $\ell k(L_i, L_j)$ vanishes, and to $|\ell k(L_i, L_j)|$ otherwise. Then, one easily shows that

$$\sum_{i < j} b_{\ell k}(L_i, L_j) \le \operatorname{sp}(L) \,.$$

We shall call this bound the *linking number bound*.

In the same article, Batson and Seed defined a spectral sequence from the Khovanov homology of a link converging to the Khovanov homology of the corresponding split link, and used it to obtain a new lower bound on sp(L). Testing it on links up to 12 crossings, they found 17 examples where this *Batson-Seed bound* is strictly stronger than the linking number bound. This enabled them to compute the splitting number of 7 of these links, while the remaining ones were left undetermined.

In [7], Cha, Friedl and Powell introduced two new techniques for computing splitting numbers. The first one, based on *covering link calculus*, relies on the following observation. If a link L has an unknotted component L_i and can be split by $\alpha + \beta$ crossing changes, α involving L_i and β not involving L_i , then the preimage of $L \setminus L_i$ in the 2-fold cover of L branched

²⁰⁰⁰ Mathematics Subject Classification. 57M25.

over L_i bounds a smooth surface in the 4-ball whose Euler characteristic can be computed from α , β , and the slice genus of the components of L. (See [7, Theorem 3.2] for the precise statement.) This technique turns out to be very efficient. However, it can only be used on a case by case basis, as different links often require specific arguments (see in particular [7, Section 5.3]). Furthermore, proving that a link does not bound a given surface in the 4-ball (or more generally, in a rational homology 4-ball in case L_i is knotted) is by no means a trivial task. Therefore the authors often need very powerful tools and heavy computations to conclude, such as results of Casson-Gordon [5, 6], the Rasmussen s-invariant [22], or twisted Alexander polynomials [15].

Their second result, originally stated as [7, Theorem 4.2], was then strengthened by Borodzik, Friedl and Powell in [3, Corollary 4.3] (see also [16, Theorem 1.1]). It can be stated as follows. If the multivariable Alexander polynomial Δ_L does not vanish, then $\operatorname{sp}(L) \ge \mu - 1$ and if $\operatorname{sp}(L) = \mu - 1$, then

$$\Delta_L(t_1,\ldots,t_{\mu}) = \pm \prod_{i=1}^{\mu} \Delta_{L_i}(t_i) \cdot p(t_1,\ldots,t_{\mu}) \cdot p(t_1^{-1},\ldots,t_{\mu}^{-1}) \cdot \prod_{i=1}^{\mu} t_i^{r_i} \cdot \prod_{i=1}^{\mu} (1-t_i)^{s_i}$$

for some $p \in \mathbb{Z}[t_1^{\pm 1}, \ldots, t_{\mu}^{\pm 1}]$ and $r_i, s_i \in \mathbb{Z}$. This Alexander polynomial obstruction is powerful in showing that the splitting number of a μ -component link is strictly greater than $\mu - 1$, but it obviously cannot be used in other cases.

These two techniques (together with the linking number bound) allowed these authors to determine the splitting numbers of the 130 prime links with up to 9 crossings. To be more precise, for 7 of these links, Cha, Friedl and Powell relied on results of Kohn [17] on the unlinking number to exclude the value sp(L) = 1; nevertheless, these values can be recovered using the Alexander polynomial obstruction of [3]. Furthermore, these techniques were harnessed to compute the splitting numbers of all of the 17 links in the Batson-Seed list.

Finally, in the very recent preprint [4], Borodzik and Gorsky found a Heegaard Floer theoretical criterion for bounding the splitting number. As an application, they showed that for any positive a, the 2-bridge link with Conway normal form C(2a, 1, 2a) has splitting number 2a, even though the linking number of the two components vanishes.

In the present paper, we give a new lower bound on the splitting number of a link in terms of its multivariable signature and nullity, see Theorem 3.1 below. As a special case, we obtain a bound in terms of the (one-variable) Levine-Tristram signature and nullity (Corollary 3.4). Although very elementary and extremely easy to compute (see e.g. [10]), these bounds turn out to be remarkably powerful. Indeed, our bound is sharp for 127 out of the 130 prime links with up to 9 crossings, and two of the remaining splitting numbers can be determined with the linking number bound. Also, our method easily gives the splitting number of all but one of the 17 links in the Batson-Seed list. Finally, our bound easily implies the following generalization of [4, Theorem 7.12]: for any $n \ge 1$ and positive $a_1, \ldots, a_n, b_1, \ldots, b_{n-1}$, the splitting number of the 2-bridge link with Conway normal form $C(2a_1, b_1, 2a_2, b_2, \ldots, 2a_{n-1}, b_{n-1}, 2a_n)$ is equal to $a_1 + \cdots + a_n$ (Theorem 4.7).

Let us finally mention that our results hold for the computation of the splitting number of a *colored link*. In other words, our bound holds if $L = L_1 \cup \cdots \cup L_{\mu}$ denotes a partition of Linto sublinks, and sp(L) is the minimal number of crossing changes between components of different sublinks to obtain the split union of these links.



FIGURE 1. A clasp intersection crossed by a 1-cycle x.

This paper is organized as follows. In Section 2, we recall the definitions of the multivariable signature and nullity, together with a list of their properties. Section 3 contains our main result, its proof, as well as corollaries and remarks. Finally, Section 4 gathers several examples and applications.

Acknowledgments. The authors wish to thank Maciej Borodzik, Jae Choon Cha, Matthias Nagel, Sebastien Ott and Mark Powell for useful discussions. The first-named author was supported by the Swiss National Science Foundation. The second-named author was supported by the NCCR SwissMAP, funded by the Swiss National Science Foundation.

2. Multivariable signatures

Recall that the Levine-Tristram signature [20, 21, 24] of an oriented link L is the map

$$\sigma_L\colon S^1\setminus\{1\}\to\mathbb{Z}\,,$$

where $\sigma_L(\omega)$ is the signature of the Hermitian matrix $H(\omega) = (1 - \omega)A + (1 - \overline{\omega})A^T$ and A is any Seifert matrix for the oriented link L. The *nullity* of L is the map $\eta_L \colon S^1 \setminus \{1\} \to \mathbb{N}$ obtained by considering the nullity of $H(\omega)$. These invariants were extended to colored links using C-complexes in [11] (see references therein for previous versions of these extensions). We now briefly recall this construction.

A μ -colored link is an oriented link L in S^3 whose components are partitioned into μ sublinks $L_1 \cup \cdots \cup L_{\mu}$. Thus, a 1-colored link is simply an oriented link, while a μ -component μ -colored link is nothing but an ordered link. A *C*-complex [12] for a μ -colored link $L = L_1 \cup \cdots \cup L_{\mu}$ is a union $S = S_1 \cup \cdots \cup S_{\mu}$ of surfaces in S^3 which is connected, and such that:

- (i) for all i, S_i is a Seifert surface for the sublink L_i ;
- (*ii*) for all $i \neq j$, $S_i \cap S_j$ is either empty or a union of clasps (see Figure 1);
- (*iii*) for all i, j, k pairwise distinct, $S_i \cap S_j \cap S_k$ is empty.

The existence of a C-complex for an arbitrary colored link is fairly easy to establish, see [9, Lemma 1]. Note that in the case $\mu = 1$, a C-complex for the 1-colored link L is nothing but a connected Seifert surface for the oriented link L. Let us now define the corresponding generalization of the Seifert form.

Given a sequence $\varepsilon = (\varepsilon_1, \ldots, \varepsilon_{\mu})$ of signs ± 1 , let $i^{\varepsilon} \colon H_1(S) \to H_1(S^3 \setminus S)$ be defined as follows. Any homology class in $H_1(S)$ can be represented by an oriented cycle x which behaves as illustrated in Figure 1 whenever crossing a clasp. Then, define $i^{\varepsilon}([x])$ as the class of the 1-cycle obtained by pushing x in the ε_i -normal direction off S_i for $i = 1, \ldots, \mu$, and consider the bilinear form

$$\alpha^{\varepsilon} \colon H_1(S) \times H_1(S) \to \mathbb{Z}, \quad (x, y) \mapsto \ell k(i^{\varepsilon}(x), y),$$



FIGURE 2. The link C(4,3,2), and a natural C-complex for it.

where ℓk denotes the linking number. Fix a basis of $H_1(S)$ and denote by A^{ε} the matrix of α^{ε} . Note that for all ε , $A^{-\varepsilon}$ is equal to $(A^{\varepsilon})^T$. Using this fact, one easily checks that for any $\omega = (\omega_1, \ldots, \omega_{\mu})$ in the μ -dimensional torus \mathbb{T}^{μ} , the matrix

$$H(\omega) = \sum_{\varepsilon} \prod_{i=1}^{\mu} (1 - \overline{\omega}_i^{\varepsilon_i}) A^{\varepsilon}$$

is Hermitian. Since this matrix vanishes when one of the coordinates of ω is equal to 1, we restrict ourselves to the subset $\mathbb{T}^{\mu}_* = (S^1 \setminus \{1\})^{\mu}$ of \mathbb{T}^{μ} .

Definition. The multivariable signature and nullity of the μ -colored link L are the functions

$$\sigma_L, \eta_L \colon \mathbb{T}^{\mu}_* \to \mathbb{Z},$$

where $\sigma_L(\omega)$ is the signature of the Hermitian matrix $H(\omega)$ and $\eta_L(\omega)$ its nullity.

Note that in the case $\mu = 1$, one clearly recovers the Levine-Tristram signature and nullity. This multivariable generalization not only turns out to be well-defined (i.e. independent of the choice of the *C*-complex), but it also satisfies all the properties of the Levine-Tristram invariants, generalized from oriented links to colored links (see [11] and Proposition 2.2 below). Let us first illustrate this definition with an example.

Example 2.1. Figure 2 shows the 2-bridge link L with Conway normal form C(4,3,2), together with a natural C-complex S for it. A natural basis for $H_1(S)$ is given by the cycles α, β , which are also depicted on this figure. With respect to this basis, one gets $A^{++} = \begin{bmatrix} 0 & 0 \\ 0 & -2 \end{bmatrix}$ and $A^{+-} = \begin{bmatrix} -1 & 1 \\ 0 & -1 \end{bmatrix}$, leading in particular to $H(-1, -1) = 4 \begin{bmatrix} -2 & 1 \\ 1 & -6 \end{bmatrix}$. Therefore, we have $\sigma_L(-1, -1) = -2$.

Let us now collect some of the basic properties of these invariants, referring to Propositions 2.5, 2.8 and 2.13 of [11] for the easy proofs.

Proposition 2.2.

(i) Consider the $(\mu + \mu')$ -colored link $L \sqcup L'$ given by the split union of the μ and μ' -colored links L and L'. Then, for all $\omega \in \mathbb{T}^{\mu}_*$ and $\omega' \in \mathbb{T}^{\mu'}_*$,

$$\sigma_{L\sqcup L'}(\omega, \omega') = \sigma_L(\omega) + \sigma_{L'}(\omega') \quad and \quad \eta_{L\sqcup L'}(\omega, \omega') = \eta_L(\omega) + \eta_{L'}(\omega') + 1.$$

(ii) Given any colored link $L = L_1 \cup \cdots \cup L_{\mu}$, the Levine-Tristram signature and nullity of the underlying oriented link can be recovered from their multivariable analogue as follows: for all $\omega \in S^1 \setminus \{1\}$,

$$\sigma_L(\omega) = \sigma_L(\omega, \dots, \omega) - \sum_{i < j} \ell k(L_i, L_j) \quad and \quad \eta_L(\omega) = \eta_L(\omega, \dots, \omega).$$

5

(iii) Let $L = L_1 \cup \cdots \cup L_{\mu}$ be a colored link, and let L' be the colored link obtained from L by reversing the orientation of every component of the sublink L_1 . Then, for all $(\omega_1, \ldots, \omega_{\mu}) \in \mathbb{T}_*^{\mu}$,

$$\sigma_{L'}(\omega_1,\ldots,\omega_{\mu}) = \sigma_L(\omega_1^{-1},\omega_2,\ldots,\omega_{\mu}) \quad and \quad \eta_{L'}(\omega_1,\ldots,\omega_{\mu}) = \eta_L(\omega_1^{-1},\omega_2,\ldots,\omega_{\mu}) \,. \quad \Box$$

We now recall a deeper result, that requires some terminology. Let $L = L_1 \cup \cdots \cup L_\mu$ be a colored link and let X_L denote its exterior. The epimorphism $\pi_1(X_L) \to \mathbb{Z}^\mu$ given by $\gamma \mapsto (\ell k(\gamma, L_1), \ldots, \ell k(\gamma, L_\mu))$ induces a regular \mathbb{Z}^μ -covering $\widetilde{X}_L \to X_L$. The homology of \widetilde{X}_L is naturally a module over $\Lambda = \mathbb{Z}[t_1^{\pm 1}, \ldots, t_{\mu}^{\pm 1}]$, where t_i denotes the covering transformation corresponding to an oriented meridian of L_i . The Λ -module $H_1(\widetilde{X}_L)$ is called the *Alexander module* of the colored link L. We shall denote its rank by $\beta(L)$.

For $r \geq 0$, let $\Delta_L^{(r)} \in \Lambda$ denote the greatest common divisor of all $(m-r) \times (m-r)$ minors of an $m \times n$ presentation matrix of $H_1(\widetilde{X}_L)$. This polynomial, well-defined up to units of Λ , is called the r^{th} -Alexander polynomial of L. Finally, $\Delta_L^{(0)}$ is simply called the Alexander polynomial of L, and denoted by Δ_L . Note that the first non-vanishing Alexander polynomial is $\Delta_L^{(\beta(L))}$, that we shall denote by Δ_L^{tor} for obvious reasons.

Proposition 2.3.

- (i) The rank of the Alexander module of a colored link is the minimal value of its nullity: $\beta(L) = \min\{\eta_L(\omega) \mid \omega \in \mathbb{T}^{\mu}_*\}.$
- (ii) The signature is constant on the connected components of the complement in \mathbb{T}^{μ}_{*} of the zeroes of the Alexander polynomial.

Proof. To show the first point, let $E_r(L)$ denote the ideal of Λ generated by the $(m-r) \times (m-r)$ minors of an $m \times n$ presentation matrix of $H_1(\tilde{X}_L)$. Also, let Σ_r denote the set consisting of all $\omega \in \mathbb{T}^{\mu}_*$ such that $p(\omega) = 0$ for each $p \in E_{r-1}(L)$. Observe that the sets Σ_r form a decreasing sequence. By [11, Theorem 4.1], $\Sigma_r \setminus \Sigma_{r+1}$ consists of all ω such that $\eta_L(\omega) = r$. Therefore, if β denotes the minimal value of η_L , we have $\Sigma_r = \mathbb{T}^{\mu}_*$ for all $r \leq \beta$ and $\Sigma_{\beta+1} \neq \mathbb{T}^{\mu}_*$. Hence, β is equal to the maximal r such that $E_{r-1}(L) = 0$, which is nothing but the rank of the Alexander module. The second point is precisely [11, Corollary 4.2].

3. Main results

The splitting number $\operatorname{sp}(L)$ of a colored link $L = L_1 \cup \cdots \cup L_{\mu}$ is the minimal number of crossing changes between sublinks of different colors required to turn L into the corresponding split colored link $L_1 \sqcup \cdots \sqcup L_{\mu}$. If μ is equal to the number of components, which is the case to keep in mind, one recovers the splitting number discussed in the introduction. Our main result is the following inequality.

Theorem 3.1. If $L = L_1 \cup \cdots \cup L_{\mu}$ is a colored link, then

$$\left|\sigma_{L}(\omega_{1},\ldots,\omega_{\mu})-\sum_{i=1}^{\mu}\sigma_{L_{i}}(\omega_{i})\right|+\left|\mu-1-\eta_{L}(\omega_{1},\ldots,\omega_{\mu})+\sum_{i=1}^{\mu}\eta_{L_{i}}(\omega_{i})\right|\leq \operatorname{sp}(L)$$

for all $(\omega_{1},\ldots,\omega_{\mu})\in\mathbb{T}_{*}^{\mu}$.



FIGURE 3. A crossing change resulting in the addition of a clasp intersection.

Proof. Let L' be a colored link obtained from L by a single crossing change involving sublinks of different colors. Then, a C-complex S' for L' can be obtained from a C-complex S for Lby adding a clasp intersection, as illustrated in Figure 3. Since S is connected, it follows that $H_1(S') = H_1(S) \oplus \mathbb{Z}[\gamma]$ for some loop γ passing through the additional clasp. With respect to this choice of bases, the resulting Hermitian matrices can be written as

$$H'(\omega) = \begin{bmatrix} H(\omega) & z \\ \overline{z}^T & \lambda \end{bmatrix}$$

for some vector z and real number λ . It easily follows that

$$|\sigma_L(\omega) - \sigma_{L'}(\omega)| + |\eta_L(\omega) - \eta_{L'}(\omega)| = 1$$

for all $\omega \in \mathbb{T}^{\mu}_*$. Consequently if $L = L^{(0)}, L^{(1)}, \ldots, L^{(s)} = L_1 \sqcup \cdots \sqcup L_{\mu}$ is a splitting sequence which realizes the splitting number, then

$$sp(L) = \sum_{i=1}^{s} \left(|\sigma_{L^{(i-1)}}(\omega) - \sigma_{L^{(i)}}(\omega)| + |\eta_{L^{(i-1)}}(\omega) - \eta_{L^{(i)}}(\omega)| \right)$$

$$\geq |\sigma_{L}(\omega) - \sigma_{L^{(s)}}(\omega)| + |\eta_{L}(\omega) - \eta_{L^{(s)}}(\omega)| .$$

Since $L^{(s)}$ is the split union of its components, the first point of Proposition 2.2 gives

$$\sigma_{L^{(s)}}(\omega) = \sum_{i=1}^{\mu} \sigma_{L_i}(\omega_i) \quad \text{and} \quad \eta_{L^{(s)}}(\omega) = \sum_{i=1}^{\mu} \eta_{L_i}(\omega_i) + \mu - 1,$$

which completes the proof.

Remark 3.2. The proof of Theorem 3.1 has an interesting consequence. Namely, assume that $L = L^{(0)}, L^{(1)}, \ldots, L^{(s)} = L_1 \sqcup \cdots \sqcup L_{\mu}$ is a splitting sequence for which there exists $\omega \in \mathbb{T}^{\mu}_*$ and $\varepsilon = \pm 1$ such that the following holds: for each *i*, either $\sigma_{L^{(i+1)}}(\omega) = \sigma_{L^{(i)}}(\omega) + \varepsilon$, or $\eta_{L^{(i+1)}}(\omega) = \eta_{L^{(i)}}(\omega) + 1$. Then, this splitting sequence is minimal. This proof also shows that our bound has the same parity as the splitting number.

Remark 3.3. In the case of an ordered link L and for generic ω , Theorem 3.1 can also be proved by using 4-dimensional considerations. Indeed, a splitting sequence of length sprovides an immersed concordance in $S^3 \times [0,1]$ between L and the corresponding split link, with s transverse double points. Removing $B^3 \times [0,1]$, where B^3 is a 3-ball in S^3 that meets each component of L in a small unknotted arc untouched by the crossing changes, we obtain a collection of discs in B^4 with s double points, bounding the link $L \# \overline{L_1} \# \dots \# \overline{L_{\mu}}$. The result then follows from Theorem 7.2, Proposition 2.10 and Proposition 2.12 of [11].

As an immediately corollary of Theorem 3.1 and the second point of Proposition 2.2, we obtain the following lower bound for sp(L) in terms of the Levine-Tristram signature and nullity of L.

Corollary 3.4. If $L = L_1 \cup \cdots \cup L_{\mu}$ is a μ -component oriented link, then

$$\left|\sigma_{L}(\omega) + \sum_{i < j} \ell k(L_{i}, L_{j}) - \sum_{i=1}^{\mu} \sigma_{L_{i}}(\omega)\right| + \left|\mu - 1 - \eta_{L}(\omega) + \sum_{i=1}^{\mu} \eta_{L_{i}}(\omega)\right| \le \operatorname{sp}(L)$$

for all $\omega \in S^{1} \setminus \{1\}.$

Note that there are 2^{μ} choices of orientations for the components of L, which give $2^{\mu-1}$ lower bounds on $\operatorname{sp}(L)$. By the third point of Proposition 2.2, these correspond to the value of the multivariable invariants on the $2^{\mu-1}$ diagonals of $\mathbb{T}_{*}^{\mu} \simeq (0, 2\pi)^{\mu}$.

Our next corollary should be compared to [3, Corollary 4.3]. The first inequality is identical; in view of Proposition 2.3 (*ii*), the case $\beta(L) = 0$ of the second statement can be understood as the signature analogue of the Alexander polynomial obstruction.

Corollary 3.5. Let $L = L_1 \cup \cdots \cup L_\mu$ be a μ -colored link, $\beta(L)$ the rank of its Alexander module, and Δ_L^{tor} its first non-vanishing Alexander polynomial. Then, one has

$$\mu - 1 - \beta(L) \le \operatorname{sp}(L) \,.$$

Furthermore, if $\mu - 1 - \beta(L) = \operatorname{sp}(L)$, then

$$\sigma_L(\omega_1,\ldots,\omega_\mu) = \sum_{i=1}^{\mu} \sigma_{L_i}(\omega_i) \quad and \quad \eta_{L_1}(\omega_1) = \cdots = \eta_{L_\mu}(\omega_\mu) = 0$$

for all $\omega = (\omega_1, \dots, \omega_\mu) \in \mathbb{T}^\mu$ such that $\Delta_L^{\text{tor}}(\omega) \neq 0$.

Proof. By Theorem 3.1 and the first point of Proposition 2.3, we have the inequalities

$$sp(L) \ge \left| \sigma_L(\omega) - \sum_{i=1}^{\mu} \sigma_{L_i}(\omega_i) \right| + \left| \mu - 1 - \eta_L(\omega) + \sum_{i=1}^{\mu} \eta_{L_i}(\omega_i) \right|$$
$$\ge \mu - 1 - \eta_L(\omega) + \sum_{i=1}^{\mu} \eta_{L_i}(\omega_i)$$
$$\ge \mu - 1 - \eta_L(\omega)$$
$$\ge \mu - 1 - \beta(L) .$$

Let us now assume that $\Delta_L^{\text{tor}}(\omega) \neq 0$. Using the notations of the proof of Proposition 2.3, this implies that ω belongs to $\Sigma_{\beta(L)} \setminus \Sigma_{\beta(L)+1}$. By [11, Theorem 4.1], this means that $\eta_L(\omega) = \beta(L)$. The second statement now follows by setting equalities for each of the inequalities displayed above.

8

Remark 3.6. In the 2-component and $\beta(L) = 0$ case, one can also show Corollary 3.5 using other considerations. Indeed, it is known that if a 2-component link has splitting number one, then it consists of a *band clasping* of two knots (see the proof of Theorem 1 in [18]). A careful analysis of the situation using *C*-complexes then yields the result. As the Alexander polynomial may be computed via *C*-complexes [9], this alternative method also enables one to recover the Alexander obstruction in the 2-component case (and in particular [7, Theorem 4.2]).

Remark 3.7. As shown by Degtyarev, Florens and Lecuona [14], the multivariable signature behaves in a controlled way via the operation known as *splicing*, which generalizes the satellite, infection, and cabling operations. Therefore, Theorem 3.1 together with [14, Theorem 2.2] immediately yields a lower bound for the splitting number of the splice of two links in terms of the signature of its splice components. In particular, this gives a lower bound for the splitting number of the cabling of a link L (along one or several of its components) in terms of the signature of L and of the corresponding torus links. However, we have not been able to find compelling applications of this result, and therefore do not discuss it in more detail.

Finally note that all the techniques developed in this paragraph can also be used to obtain lower bounds on the unlinking number u(L) of a link L. For instance, if $L = L_1 \cup \cdots \cup L_{\mu}$ is a μ -component link, then one obtains

$$|\sigma_L(\omega_1,\ldots,\omega_\mu)| + |\mu - 1 - \eta_L(\omega_1,\ldots,\omega_\mu)| + \sum_{i < j} |\ell k(L_i,L_j)| \le u(L)$$

for all $(\omega_1, \ldots, \omega_\mu) \in \mathbb{T}^{\mu}_*$. Unfortunately, this bound is not very powerful so we shall not discuss it any further.

4. Examples and applications

In this section, we use our bound on three families of examples: the links with at most nine crossings, the links of the Batson-Seed list, and 2-bridge links. All the diagrams are taken from SnapPy [13].

As stated above, we first tested Theorem 3.1 on all 130 prime links with fewer than ten crossings, using the notations and data from LinkInfo [8]. In 125 cases, the Levine-Tristram bound of Corollary 3.4 is enough to recover the splitting number, while Corollary 3.5 recovers two additional cases. The three remaining links are L9a47 and L9n27 (for which the linking number bound is sharp) and L8a9 (whose splitting number can be recovered by the Alexander polynomial obstruction). Let us illustrate this with a couple of examples.



FIGURE 4. The link L9a29.

Example 4.1. The splitting number of the link L = L9a29 depicted in Figure 4 was shown to be 3 in [7, Section 4.2] by using the Alexander polynomial obstruction. Orienting L so that it consists of a right-handed trefoil L_1 and a trivial knot L_2 with linking number $\ell k(L_1, L_2) =$ -1, we have

 $\sigma_L(-1) = 5$, $\eta_L(-1) = 0$, $\sigma_{L_1}(-1) = 2$, $\eta_{L_1}(-1) = 0$, $\sigma_{L_2}(-1) = 0$, $\eta_{L_2}(-1) = 0$. It follows that the bound

$$|\sigma_L(-1) + \ell k(L_1, L_2) - \sigma_{L_1}(-1) - \sigma_{L_2}(-1)| + |2 - 1 - \eta_L(-1) + \eta_{L_1}(-1) + \eta_{L_2}(-1)| = 3$$
f Corollary 3.4 is sharp

of Corollary 3.4 is sharp.

Example 4.2. The splitting number of the link L = L9a24 depicted on the left hand side of Figure 5 was shown to be 3 in [7, Section 6] by using the Alexander polynomial obstruction. The lower bound of Corollary 3.4 gives 1, so it is not sufficient to conclude. However, Corollary 3.5 shows that the splitting number must be greater than one, and therefore equal to three due to the parity of the linking number. Indeed, one does not always have $\sigma_L(\omega_1, \omega_2) =$ $\sigma_{L_1}(\omega_1) + \sigma_{L_2}(\omega_2)$, as shown by the zero locus of Δ_L illustrated on the right hand side of Figure 5. (Recall Proposition 2.3 (ii).)

The exact same analysis shows that the splitting number of L9n17 is three.



FIGURE 5. The link L9a24 and the intersection of the zero locus of its Alexander polynomial with \mathbb{T}^2_* .

Next, we tested our bound on the 17 links of the Batson-Seed list, using Seifert matrix data kindly provided by J.C. Cha. Among these, there are seven 12-crossing links (namely, L12n1342, L12n1350, L12n1357, L12n1363, L12n1367, L12n1274 and L12n1404) for which both components are trefoils, and whose splitting number was shown to be equal to 3 by Batson and Seed. Cha, Friedl and Powell recovered these results via the Alexander obstruction. Applying Corollary 3.5 as in Example 4.2 immediately gives the same conclusion. Moreover, the splitting number of L12n1342, L12n1350, L12n1367 and L12n1274 can be recovered by using the Levine-Tristram signature and Corollary 3.4 alone. Let us illustrate this with one example.

Example 4.3. Orient the link L = L12n1367 (depicted in Figure 6) so that $\ell k(L_1, L_2) = 1$ and set $\omega = e^{\frac{i\pi}{3}}$. Using

$$\sigma_L(\omega) = 0, \quad \sigma_{L_1}(\omega) = 1, \quad \sigma_{L_2}(\omega) = -1, \quad \eta_L(\omega) = \eta_{L_1}(\omega) = \eta_{L_2}(\omega) = 1,$$

it follows that the bound

$$|\sigma_L(\omega) + \ell k(L_1, L_2) - \sigma_{L_1}(\omega) - \sigma_{L_2}(\omega)| + |2 - 1 - \eta_L(\omega) + \eta_{L_1}(\omega) + \eta_{L_2}(\omega)| = 3$$
 of Corollary 3.4 is sharp.



FIGURE 6. The link L12n1367.

For the remaining 10 links in this list, Batson and Seed could not determine whether the splitting number is 3 or 5. These links having two or three components, the Alexander polynomial obstruction cannot be applied. However, Cha, Friedl and Powell used various arguments based on covering link techniques to determine these values. As it turns out, our bound allows to easily determine these splitting number for all but one of them, namely L12n1321. Here are several of these examples.

Example 4.4. Consider the link L = L11a372 depicted in Figure 7, whose splitting number was shown to be 5 in [7, Section 5.2]. Orient L so that its trivial components L_1, L_2 satisfy $\ell k(L_1, L_2) = -1$. Since $\sigma_L(-1) = 5$ and $\eta_L(-1) = 0$, the bound

$$|\sigma_L(-1) + \ell k(L_1, L_2)| + |2 - 1 - \eta_L(-1)| = 5$$

given by the classical signature is enough to conclude.



FIGURE 7. The link L11a372.

Example 4.5. It took the whole Section 5.3 of [7] to show that the splitting number of the 3-component link L = L12a1622 depicted in Figure 8 is equal to 5. Orienting L so that its trivial components L_1, L_2, L_3 satisfy $\ell k(L_1, L_2) = 0$, $\ell k(L_1, L_3) = 0$, $\ell k(L_2, L_3) = 1$, and picking $\omega = e^{\frac{3\pi i}{4}}$, we have $\sigma_L(\omega) = -4$ and $\eta_L(\omega) = 0$. Hence, the bound

$$|\sigma_L(\omega) + \ell k(L_2, L_3)| + |3 - 1 - \eta_L(\omega)| = 5$$

of Corollary 3.4 immediately provides the desired splitting number.



FIGURE 8. The link L12a1622.

Example 4.6. Consider the link L = L12n1326 depicted in Figure 9. Cha-Friedl-Powell [7, Section 5.2] used the twisted Alexander polynomial to compute the slice genus of a covering link, and concluded that sp(L) = 3. Orienting L so that its trivial components L_1, L_2 satisfy $\ell k(L_1, L_2) = 1$, and picking $\omega = e^{\frac{\pi i}{5}}$ so that $\sigma_L(\omega) = 1$ and $\eta_L(\omega) = 0$, the bound

 $|\sigma_L(\omega) + \ell k(L_1, L_2)| + |2 - 1 - \eta_L(\omega)| = 3$

of Corollary 3.4 immediately provides the desired splitting number.



FIGURE 9. The link L12n1326.

In the very recent preprint [4], Borodzik and Gorsky used Heegaard Floer techniques to compute the splitting number of the 2-bridge link with Conway normal form C(2a, 1, 2a) (recall Figure 2 for an explanation of this notation). We conclude this article by showing how easily we can compute the splitting number of a much wider class of 2-bridge links.

Theorem 4.7. For any $n \ge 1$ and any positive integers $a_1, \ldots, a_n, b_1, \ldots, b_{n-1}$, the splitting number of the 2-bridge link $C(2a_1, b_1, 2a_2, b_2, \ldots, 2a_{n-1}, b_{n-1}, 2a_n)$ is equal to $a_1 + \cdots + a_n$.

Proof. Let us denote this link by L. Inspecting the standard diagram given by Conway's normal form, it is clear that $sp(L) \leq a_1 + \cdots + a_n =: s$. Furthermore, as exemplified in Figure 2, L admits a natural C-complex S made of two discs intersecting in s clasps, yielding a natural basis for $H_1(S)$ of cardinality s - 1. Computing the associated generalized Seifert matrices leads to

$$H(-1,-1) = 4 \begin{bmatrix} -2d_1 & 1 & & \\ 1 & -2d_2 & 1 & & \\ & 1 & \ddots & \ddots & \\ & & \ddots & & 1 \\ & & & 1 & -2d_{s-1} \end{bmatrix}$$

This is a pre-publication version of this article, which may differ from the final published version. Copyright restrictions may apply.

where the diagonal elements are positive integers whose precise value depends on the integers a_i and b_i , but will play no role here. Indeed, all the eigenvalues of a matrix of the form above are negative, so $\sigma_L(-1,-1) = 1 - s$ and $\eta_L(-1,-1) = 0$. Since L has $\mu = 2$ components, both unknotted, the result now follows from Theorem 3.1.

Note that this last result was obtained using the ordinary signature alone. Using the full power of Theorem 3.1, it is possible to determine the splitting number of many other 2-bridge links, such as all $C(a_1, \ldots, a_n)$ with $n \leq 3$. The first example of a 2-bridge link whose splitting number is unknown to us is L = C(4, 3, 1, 3): the linking number vanishes and our bound reads $\operatorname{sp}(L) \geq 2$, while the "obvious" splitting sequence has length 4.

References

- Colin C. Adams. Splitting versus unlinking. J. Knot Theory Ramifications, 5(3):295–299, 1996.
- [2] Joshua Batson and Cotton Seed. A link-splitting spectral sequence in Khovanov homology. Duke Math. J., 164(5):801-841, 2015.
- [3] Maciej Borodzik, Stefan Friedl, and Mark Powell. Blanchfield forms and Gordian distance. Journal of the Mathematical Society of Japan. to appear.
- Maciej Borodzik and Eugene Gorsky. Immersed concordances of links and Heegaard Floer homology. January 2016. http://arxiv.org/abs/1601.07507.
- [5] A. J. Casson and C. McA. Gordon. On slice knots in dimension three. In Algebraic and geometric topology (Proc. Sympos. Pure Math., Stanford Univ., Stanford, Calif., 1976), Part 2, Proc. Sympos. Pure Math., XXXII, pages 39–53. Amer. Math. Soc., Providence, R.I., 1978.
- [6] A. J. Casson and C. McA. Gordon. Cobordism of classical knots. In À la recherche de la topologie perdue, volume 62 of Progr. Math., pages 181–199. Birkhäuser Boston, Boston, MA, 1986. With an appendix by P. M. Gilmer.
- [7] Jae Choon Cha, Stefan Friedl, and Mark Powell. Splitting numbers of links. *Proceedings* of the Edinburgh Mathematical Society. to appear.
- [8] Jae Choon Cha and Charles Livingston. Linkinfo: Table of knot invariants. January 24, 2016. http://www.indiana.edu/~linkinfo.
- [9] David Cimasoni. A geometric construction of the Conway potential function. Comment. Math. Helv., 79(1):124-146, 2004.
- [10] David Cimasoni and Anthony Conway. Colored tangles and signatures. July 2015. http://arxiv.org/abs/1507.07818.
- [11] David Cimasoni and Vincent Florens. Generalized Seifert surfaces and signatures of colored links. Trans. Amer. Math. Soc., 360(3):1223–1264 (electronic), 2008.
- [12] D. Cooper. The universal abelian cover of a link. In Low-dimensional topology (Bangor, 1979), volume 48 of London Math. Soc. Lecture Note Ser., pages 51–66. Cambridge Univ. Press, Cambridge-New York, 1982.
- [13] Marc Culler, Nathan M. Dunfield, and Jeffrey R. Weeks. SnapPy, a computer program for studying the topology of 3-manifolds. Available at http://snappy.computop.org.
- [14] Alex Degtyarev, Vincent Florens, and Ana Lecuona. The signature of a splice. September 2014. http://arxiv.org/abs/1409.5873.
- [15] Chris Herald, Paul Kirk, and Charles Livingston. Metabelian representations, twisted Alexander polynomials, knot slicing, and mutation. Math. Z., 265(4):925–949, 2010.

- [16] Akio Kawauchi. The Alexander polynomials of immersed concordant links. Bol. Soc. Mat. Mex. (3), 20(2):559–578, 2014.
- [17] Peter Kohn. Unlinking two component links. Osaka J. Math., 30(4):741–752, 1993.
- [18] Hisako Kondo. Knots of unknotting number 1 and their Alexander polynomials. Osaka J. Math., 16(2):551–559, 1979.
- [19] Marc Lackenby. Elementary knot theory. to be published by the Clay Mathematics Institute, 2014.
- [20] J. Levine. Knot cobordism groups in codimension two. Comment. Math. Helv., 44:229– 244, 1969.
- [21] Kunio Murasugi. On a certain numerical invariant of link types. Trans. Amer. Math. Soc., 117:387–422, 1965.
- [22] Jacob Rasmussen. Khovanov homology and the slice genus. Invent. Math., 182(2):419– 447, 2010.
- [23] Ayaka Shimizu. The complete splitting number of a lassoed link. *Topology Appl.*, 159(4):959–965, 2012.
- [24] A. G. Tristram. Some cobordism invariants for links. Proc. Cambridge Philos. Soc., 66:251–264, 1969.

Université de Genève, Section de mathématiques, 2-4 rue du Lièvre, 1211 Genève 4, Switzerland

E-mail address: david.cimasoni@unige.ch

Université de Genève, Section de mathématiques, 2-4 rue du Lièvre, 1211 Genève 4, Switzerland

E-mail address: anthony.conway@unige.ch

Université de Genève, Section de mathématiques, 2-4 rue du Lièvre, 1211 Genève 4, Switzerland

 $E\text{-}mail\ address:\ \texttt{kleopatra.zacharova@etu.unige.ch}$