



Gum-feeder as environmental enrichment for zoo marmosets and tamarins

Journal:	<i>Zoo Biology</i>
Manuscript ID	ZOO-19-051.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Regaiolli, Barbara; Parco Natura Viva - Garda Zoological Park, Research Department Angelosante, C; Universita degli Studi di Bologna Dipartimento di Scienze Mediche Veterinarie, Veterinary Medical Sciences – DIMEVET University of Bologna Marliani, Giovanna; Universita degli Studi di Bologna Dipartimento di Scienze Mediche Veterinarie, Veterinary Medical Sciences – DIMEVET University of Bologna Accorsi, Pier; Universita degli Studi di Bologna Dipartimento di Scienze Mediche Veterinarie, Veterinary Medical Sciences – DIMEVET University of Bologna Vaglio, Stefano; University of Wolverhampton Faculty of Science and Engineering, Biology, Chemistry and Forensic Science; Durham University, Anthropology Spiezio, Caterina; Parco Natura Viva - Garda Zoological Park, Research Department;
Keywords:	zoo primate welfare, callitrichines, gum Arabic

SCHOLARONE™
Manuscripts

Research highlights

- Wooden gum feeders can improve the welfare of zoo marmosets and tamarins by stimulating their natural species-specific behaviors ~~and decreasing stress-related behavior, and helping them to cope with stressful contexts.~~
- Zoo callitrichines, managed by adequate enrichment programs, may show positive welfare.

For Peer Review

1 Regaiolli *et al.*

1

2
3
4 **Title:** Gum-feeder as environmental enrichment for zoo marmosets and tamarins

5
6
7
8 **Short running title:** Gum-feeder as enrichment for marmosets and tamarins

9
10
11
12 **Authors:** Regaiolli B.¹, Angelosante C.², Marliani G.², Accorsi P.A.², Vaglio S.^{1,3,4} & Spiezio C.¹

13
14
15
16
17 **Institutional affiliations:**

18
19 ¹Research and Conservation Department, Parco Natura Viva - Garda Zoological Park, Loc. Figara 40, 37012
20 Bussolengo, Italy

21
22
23 ²Dipartimento di Scienze Mediche Veterinarie - Università di Bologna - Via Tolara di Sopra 50, 40064
24 Ozzano Emilia (BO), Italy

25
26
27 ³Department of Biology, Chemistry and Forensic Science, University of Wolverhampton, Wulfruna Street,
28 WV1 1LY Wolverhampton, UK

29
30
31 ⁴Department of Anthropology & Behaviour, Ecology and Evolution Research (BEER) Centre, Durham
32 University, South Road, DH1 3LE Durham, UK

33
34
35
36
37 **Corresponding author:**

38 Dr Stefano Vaglio

39
40
41 S.Vaglio@wlv.ac.uk

42
43
44 +44 (0) 1902 323328

Regaiolli *et al.*

Abstract

Tamarins and marmosets are small-bodied social callitrichines. Wild callitrichines feed on exudates, such as sap and gum; particularly, marmosets are mainly gummivores, while tamarins consume gums only occasionally and opportunistically. Zoo marmosets and tamarins are usually provided with gum Arabic as an alternative to the exudates normally found in the wild. The aim of this study was to evaluate the effects of a gum feeder on the behavior and well-being of four zoo-managed callitrichines. We studied four cotton-top tamarins (*Saguinus oedipus*), four red-handed tamarins (*Saguinus midas*), two pygmy marmosets (*Cebuella pygmaea*) and three Geoffroy's marmosets (*Callithrix geoffroyi*) housed at Parco Natura Viva (Italy). We conducted the study over two different periods, a baseline (control, without the gum feeder) and then a gum-feeder (when the gum feeder was provided) period. We used continuous focal animal sampling to collect behavioral data, including durations of social and individual behaviors. We collected 240 minutes of observations per period per study subject, with a total of 3,120 minutes for all the subjects in the same period and of 6,240 minutes in both periods. We analyzed data by using non-parametric statistical tests. First, we found that the gum feeder promoted species-specific behaviors, such as exploration, and diminished self-directed behaviors, suggesting an enriching effect on tamarin and marmoset behavior. Moreover, in red-handed tamarins, the provision of the gum feeder reduced the performance of self-directed and abnormal behavior, specifically coprophagy. These results confirm that gum-feeders are effective foraging enrichment tools for zoo marmosets and tamarins.

Keywords: zoo primate welfare, callitrichines, gum Arabic

1 Introduction

Tamarins and marmosets are small-bodied New World monkeys belonging to the subfamily of Callitrichinae (Groves, 2001). They live in geographical areas ranging from Costa Rica to South Brazil, Bolivia and Paraguay, with the greatest variety of species found in the Amazonian region (Emmons, 1990; Buckner *et al.*, 2015). They are cooperative breeders and may form familial groups made up of a breeding pair, several siblings and other members helping with parental care of the offspring. The breeding pair is dominant over other group members (Emmons, 1990; Smuts *et al.*, 1986).

The diet of wild tamarins and marmosets is affected by seasonality and availability of food resources; they mainly feed on fruit, insects, small vertebrates, leaves and nectar (Rosenberger, 1992; Garber, 1993; Bairrão Ruivo, 2010). In addition, their natural diet includes plant exudates, such as gum and sap, which are rich in carbohydrates and minerals (Bairrão Ruivo, 2010; Power, 2010). Many plant gum exudates consist of β -linked complex polysaccharides which often also contain glycoproteins and proteoglycans (Gashua *et al.*, 2015) from which energy, water and minerals, particularly calcium, can be obtained after fermentation. Callitrichines require a high intake of calcium from their diet, as these species generally give birth to twins and plenty of milk is required during the nursing process (Garber, 1993; Kelly, 1993; Heymann and Smith, 1999; Pack *et al.*, 1999; Passamani and Rylands, 2000; Taylor and Vinyard, 2004). The benefits of gum feeding have been widely investigated in the wild. Therefore, supplementing the diet of zoo non-human primate species with plant gum exudates, might improve the diet and general well-being of these primate species in captive environments (Garber, 1993; Kelly, 1993; Heymann and Smith, 1999; Pack *et al.*, 1999; Passamani and Rylands, 2000; Taylor and Vinyard, 2004).

Plant gum exudates are an essential component of the marmoset diet, whilst it is less important for other callitrichines (Power and Oftedal, 1996; Bairrão Ruivo, 2010; Power, 2010). For this reason, marmosets show behavioral, morphological and metabolic adaptations to gum feeding. In particular, their dental adaptations allow them to gouge trees and thereby stimulate gum exudate production as part of the plants' response to wounding (Vinyard *et al.*, 2003; Vinyard *et al.*, 2004; Eng *et al.*, 2009; Burrows & Nash, 2010); while modifications of the gastro-intestinal trait permit the digestion of gums and other plant exudates (Coimbra-Filha and Mittermeier 1977; Heymann & Smith, 1999; Bairrão Ruivo, 2010). Tamarins tend to

Regaiolli *et al.*

1
2
3
4 28 feed on gum and sap only opportunistically, when this food is available in trees wounded by other animals,
5
6 29 with seasonal variation in terms of time investment for gum feeding (Garber, 1993; Power and Oftedal,
7
8 30 1996; Power, 2010).
9

10 31 Zoo tamarins and marmosets are generally provided with commercially available gum Arabic, as a
11
12 32 replacement for the various plant gum exudates that they consume in the wild (Goodrum *et al.*, 2000).
13
14 33 Furthermore, these plant gum exudates may be crucial biochemical digestive challenges for the normal
15
16 34 functioning of their digestive tract (Bairrão Ruivo, 2010). Wild tamarins show more frequent gum feeding
17
18 35 during the afternoon, as gum digestion is time consuming and would therefore be easier during the night
19
20 36 sleeping (Kelly, 1993; Heymann & Smith, 1999; Bairrão Ruivo, 2010). On the contrary, gummivore
21
22 37 marmosets are known to eat gum frequently throughout the daylight hours as their gastrointestinal system is
23
24 38 well-adapted to gum digestion (Kelly, 1993; Heymann & Smith, 1999; Bairrão Ruivo, 2010). Tamarins and
25
26 39 marmosets are usually attracted to novel objects, and vigilant and aware of what happens in their surrounding
27
28 40 environment (Menzel & Menzel, 1979). Promoting natural gum feeding behavior with specific devices might
29
30 41 be important to enhance the feeding strategy and husbandry of zoo-managed callitrichines in order to
31
32 42 improve their physical and mental well-being.
33
34

35 43 Giving zoo animals the opportunity to perform their species-specific behaviors represents one of the
36
37 44 primary goals of modern zoological gardens. In addition, ethological parameters have been proven to be a
38
39 45 valuable tool in assessing zoo animal welfare (Hill & Broom, 2009; Fontani *et al.*, 2014). The occurrence of
40
41 46 natural species-specific behaviors, such as exploratory behaviors, is considered an indicator of good welfare
42
43 47 status and enriched environment (Mench, 1994; Mellor & Beausoleil, 2015), while abnormal behaviors, such
44
45 48 as excessive inactivity, stereotypies and self-injurious behaviors, may indicate poor welfare or stressful
46
47 49 scenarios (Renner *et al.*, 2000; Manteca *et al.*, 2016). Moreover, self-directed behaviors such as self-
48
49 50 grooming and scratching, are usually benign activities that occur commonly in non-human primates
50
51 51 (Maestripieri *et al.*, 1992). However, in certain situations such as social tension and conflicting or frustrating
52
53 52 contexts, self-directed behaviors can be associated with uncertainty and anxiety and have been considered as
54
55 53 displacement activities (Troisi & Schino, 1987; Troisi, 2002; Lutz, 2014; Spiezio *et al.*, 2017). Thus, a
56
57 54 decrease in self-directed behaviors might be considered as a positive welfare indicator, although these
58
59 55 activities are normally included and well-represented in the species-specific behavioral repertoire of primate
60

1
2
3
4 56 species (Spiezio *et al.*, 2017; Leeds & Lukas, 2018). Similarly, coprophagy occurs in both captive and wild
5
6 57 non-human primates and may have an adaptive value in these species; however, in controlled environment
7
8 58 this behavior has been related to factors such as nutritional deficiency and medical problems but also
9
10 59 boredom and social stress (Krief *et al.*, 2004; Prates & Bicca-Marques 2005). Therefore, coprophagy has
11
12 60 been classified as an abnormal behavior and has been identified as a possible indicator of poor well-being
13
14 61 (*i.e.*, Prates & Bicca-Marques 2005; Lutz, 2018).
15
16

17 62 Environmental enrichment is a widespread practice among modern zoos and has been found to
18
19 63 promote the performance of species-specific behavioral repertoire and to address as well as prevent abnormal
20
21 64 behavior (Hosey *et al.*, 2013). Though some callitrichines may not require gum to reach their nutritional
22
23 65 needs in zoo settings, gum feeding may represent a behavioral necessity and could improve the diversity of
24
25 66 the behavioral repertoire as well as the welfare status of zoo marmosets and tamarins (Bairrão Ruivo, 2010).
26
27 67 In the current study, zoo tamarins and marmosets were provided with wooden drilled logs as gum-feeders.
28
29 68 Previous research investigating the effects of a similar gum-feeder on zoo marmosets and tamarins suggested
30
31 69 that this environmental enrichment could promote species-specific behaviors, reduce abnormal behaviors,
32
33 70 such as stereotypies and coprophagy, and decrease inactivity (McGrew *et al.*, 1986; Roberts *et al.*, 1999;
34
35 71 Taylor, 2002; Huber & Lewis, 2011). In particular, the provision of hanging feeder baskets and sticks
36
37 72 smeared with Acacia gum promoted feeding and foraging while decreasing stereotypic behavior, specifically
38
39 73 excessive coprophagy, in red-handed tamarins in zoo (Taylor, 2002). Similarly, common marmosets
40
41 74 (*Callithrix jacchus*) in different social housing conditions have been found to benefit from foraging
42
43 75 enrichments, specifically gum feeders, promoting natural behaviors and leading to a reduction in the time
44
45 76 spent performing stereotypic behavior such as pacing (McGrew *et al.*, 1986; Roberts *et al.*, 1999). In general,
46
47 77 providing zoo callitrichines with gum arabic in feeders that require specific feeding abilities has been found
48
49 78 to promote naturalistic behaviors and feeding strategies described in the wild. In particular, artificial gum
50
51 79 feeders are cheap and easy to build and can be helpful in enhancing the physical and mental well-being of
52
53 80 these species, which are well-disposed to work for gum, enhancing also the educative value of zoo exhibit
54
55 81 (McGrew *et al.*, 1986; Huber & Lewis, 2011).
56
57
58
59

60 82 The aims of this study were to:

- 1
2
3
4 83 • Assess the effects of the gum-feeder on the behavior of two tamarin species (*Saguinus*
5
6 84 *oedipus* and *Saguinus midas*) and two marmoset species (*Callithrix geoffroyi* and *Cebuella*
7
8 85 *pygmaea*) which were scarcely investigated in the past. On the basis of prior work by other
9
10 86 authors revealing that gum feeding represents a behavioral need and a digestive challenge for
11
12 87 the digestive system of marmosets and, to a lesser extent, tamarins (Heymann & Smith,
13
14 88 1999; Bairrão Ruivo, 2010; Hosey *et al.*, 2013; Roberts *et al.*, 1999; Taylor, 2002), we
15
16 89 predicted that the gum feeder would increase the performance of species-specific behaviors,
17
18 90 such as explorative and feeding behaviors and reduce inactivity and abnormal behavior.
19
20
21 91 • Compare the duration of gum Arabic feeding between tamarins and marmosets, particularly
22
23 92 the time spent feeding on gum between the morning and the afternoon within the two
24
25 93 groups, to identify the optimal time for gum provision. Since gum digestion is time
26
27 94 consuming and would therefore be easier during night sleeping, tamarins show gum feeding
28
29 95 more during the afternoon (Heymann & Smith, 1999; Bairrão Ruivo, 2010). Thus, we expect
30
31 96 that marmosets would perform gum feeding more than tamarins and during day time, while
32
33 97 tamarins would eat gum during afternoon hours rather than in the morning.
34
35
36 98
37

38 99 **Materials and Methods**

40 100 *Study subjects and area*

41
42 101 We studied eight tamarins, specifically four cotton-top tamarins and four red-handed tamarins; and
43
44 102 five marmosets, three Geoffroy's marmosets and a pair of pygmy marmosets. The study subjects belonged to
45
46 103 two different age-groups: juveniles, including individuals aged less than two years (approximate age of
47
48 104 sexual maturity, Tardif 1984; Ziegler *et al.* 1987; Abbott *et al.*, 2003; Tardif *et al.*, 2011) and adults,
49
50 105 including individuals aged more than two years (Abbott *et al.*, 2003; Tardiff *et al.*, 2008; Tardif *et al.*, 2011)
51
52
53 106 **(Table 1).**

54
55 107 All groups were housed at Parco Natura Viva-Garda Zoological Park (Bussolengo, Italy) in
56
57 108 separated enclosures in the Tropical Green House. Although enclosures were not adjacent to each other,
58
59 109 Geoffroy's marmosets, cotton-top tamarins and red-handed tamarins were in acoustic contact. Their
60
110 enclosures were made of an outdoor and an indoor area and each area was approximately 30 m² and

1
2
3
4 111 contained trees, branches and logs, ropes, wooden boxes, sheds and shelves; the indoor areas were heated
5
6 112 and provided with UV lamps. The pygmy marmoset enclosure was an 18 m² aviary.
7
8

9 113 During the study, the gum feeders were placed in the outdoor area of the enclosures. Tamarins and
10
11 114 marmosets were fed twice a day with fruits, multi-cereal pap, mealworms, gum Arabic, and occasionally
12
13 115 meat and eggs. Fresh water was available *ad libitum*. Manipulative, sensory or food-related devices were
14
15 116 provided daily as environmental enrichment. The study, which did not involve any invasive or stressful
16
17 117 techniques, was conducted in accordance with the EU Directive 2010/63/EU and the Italian legislative
18
19
20 118 decree 26/2014 for Animal Research.
21

22 119 *Procedure and data collection*

23
24
25 120 The study was made of two different periods, the baseline and the gum-feeder period. During the baseline,
26
27 121 gum Arabic was provided in bowls at the time of the afternoon meal following the daily routine husbandry.
28
29 122 In the gum-feeder period, gum Arabic was provided, using the new gum feeders, at the usual feeding times
30
31 123 over the morning and the afternoon. The gum-feeder consisted of a wooden drilled disc, with 10 to 15 holes
32
33 124 each, on both sides of the disc (**Figure 1**). The discs were hung with ropes on the enclosure trees and
34
35 125 branches, approximately 1.5 meters above the ground. The daily amount of gum Arabic was put in the holes
36
37 126 of the feeder. In particular, the amount of gum Arabic per subject was prepared using approximately 8 grams
38
39 127 of powder and 5 grams of water. The entire study consisted of 6,240 minutes (104 hours) of observation
40
41 128 divided in the baseline (52 hours) and in the gum-feeder period (52 hours) for all the subjects. Per period and
42
43
44 129 per subject, a total of 240 minutes of behavioral observations were carried out and two sessions per day per
45
46 130 monkey were run (one in the morning, one in the afternoon). The duration of the data collection sessions per
47
48 131 monkey differed between species based on the sample size. In particular, per period, data on each cotton-top
49
50 132 tamarin (N = 4) and on each red-handed tamarin (N = 4) were collected during eight 30-minute sessions. For
51
52 133 the marmosets, per period and per monkey, data were collected during six 40-minute sessions for Geoffroy's
53
54 134 marmosets (N = 3) and four 60-minute sessions for pygmy marmosets (N = 2) for an overall of 240 minutes
55
56 135 of observation for each subject within each period. Monkeys were observed in a prescribed sequence
57
58 136 following a specific design to avoid time-of-day bias in data collection. Feeding and enrichment times were
59
60 137 the same over the study period, one in the morning and one in the afternoon.

Regaiolli *et al.*

1
2
3
4 138 A continuous focal animal sampling method (Altmann, 1974) was used to collect durations of
5
6 139 normal and abnormal individual and social behaviors (**Table 2**). The time spent out-of-sight, here defined as
7
8 140 “not observed”, was also recorded. The red-handed tamarins were the only subjects to perform abnormal
9
10 141 behavior, specifically coprophagy, which was reported in three out of four monkeys, although for a very low
11
12 142 percentage of the total observation time (ranging from 0.13% to 1.67%). In order to assess the effect of the
13
14 143 feeder on the behavior of the red-handed tamarins, we created the category stress-related behaviors (SRB),
15
16 144 including coprophagy and self-directed behaviors. Although these behaviors are found in wild animals and
17
18 145 may be adaptive, they both have been related to stressful and conflict situations within controlled
19
20 146 environments (Troisi & Schino, 1987; Troisi, 2002; Prates & Bicca-Marques 2005; Lutz, 2014; Spiezio *et*
21
22 147 *al.*, 2017; Lutz, 2018). The gum-eating behavior was included in the category “feeding/foraging” as in the
23
24 148 first period gum Arabic was provided in bowls with other food; however, in order to compare the time spent
25
26 149 feeding on gum between marmosets and tamarins, the duration of gum feeding from the new feeder was also
27
28 150 collected (**Table 2**).
29
30

31 151 32 33 34 152 *Data analysis*

35
36 153 Data were analyzed using non-parametric statistic tests and significance level was set at $p < 0.05$ (Siegel &
37
38 154 Castellan, 1992). In particular, the Wilcoxon test was used to compare the behavior of the study subjects
39
40 155 between the two periods (baseline *vs.* gum-feeder), whereas the Mann-Whitney test was used to compare the
41
42 156 time spent feeding on gum between the study groups (marmosets *vs.* tamarins) in the gum-feeder period.
43
44 157 StatView version 5.0 (SAS Institute Inc.) was used for all the statistical analyses on behavioral data. In
45
46 158 addition, single-case analyses were used to test the effect of the gum-feeder on the behavior of each
47
48 159 individual, comparing the performance of SRBs between different periods within each individual of red-
49
50 160 handed tamarins (Fisch, 2001) performing stress-related behaviors, specifically abnormal behavior and self-
51
52 161 directed behavior. For the single-case analyses, Wilcoxon-Mann-Whitney test (Marx *et al.*, 2016) was used.
53
54 162 For all behavioral categories, medians and interquartile range (IQR) are reported in the manuscript and
55
56 163 tables.
57
58
59 164
60

165 **Results**

1
2
3
4 166
5
6 167 *Baseline vs. gum-feeder period*

7
8 168 We investigated the effects of the gum-feeder on the behavior of the study subjects comparing the time spent
9
10 169 in individual and social behaviors as well as “not observed” between the two periods (**Figure 2, Table 3**).

11
12
13 170 When considering each individual behavior, Wilcoxon tests revealed that feeding/foraging,
14
15 171 maintenance and self-directed were performed significantly more during the baseline than during the gum-
16
17 172 feeder period, whereas the opposite pattern was found for visual exploration. No other significant differences
18
19 173 were found (see **Figure 2** and **Table 3** for median, IQR and statistical values). On the other hand, within
20
21 174 social behaviors, interspecific behavior was performed more during the baseline than during the gum-feeder
22
23 175 period, whereas no other differences were found (see **Table 3** for median, IQR and statistical values).
24
25 176 Finally, Wilcoxon test revealed that “not observed” was performed significantly more during the baseline
26
27 177 than during the gum-feeder period ($z = -2.481, p = 0.013$) (**Table 3**).

28
29 178
30
31
32 179 *Effect of the gum-feeder on tamarins*

33
34 180 After evaluating the effects of the gum-feeder on all subjects, we focused on tamarins and marmosets
35
36 181 separately. Regarding tamarins, within individual behaviors, feeding/foraging and maintenance were
37
38 182 performed significantly more during the baseline than during the gum-feeder period, whereas no significant
39
40 183 differences were found in any other behavioral category (see **Table 3** for median, IQR and statistical values).

41
42 184
43
44 185 *Effects of the gum-feeder on marmosets*

45
46 186 Regarding marmosets, “not observed” was performed significantly more during baseline than during the
47
48 187 gum-feeder period ($z = -2.023, p = 0.043$), while no other significant differences were found between the two
49
50 188 periods (see **Table 3** for median, IQR and statistical values).

51
52 189
53
54
55 190 *Effects of the gum-feeder on stress-related behavior in the red-handed tamarins*

56
57 191 Three red-handed tamarins (CS, OB and Normann) showed the abnormal behavior ‘coprophagy’. In order to
58
59 192 test whether the gum-feeder positively affected the behavior as a measure of welfare of these individuals, a
60
193 single-case analysis was done to compare the performance of SRBs between different periods within each

Regaiolli *et al.*

1
2
3
4 194 individual. For CS, the median duration of SRB was 113.5 (50) seconds in the baseline and 74.5 (51.25)
5
6 195 seconds in the gum-feeder period; for OB, the median duration of SRB was 90 (70.25) seconds in the
7
8 196 baseline and 52 (72.75) seconds in the gum-feeder period; for Normann, the median duration of SRB was
9
10 197 101 (99.75) seconds in the baseline and 60 (70) seconds in the gum-feeder period. Wilcoxon-Mann-Whitney
11
12 198 tests revealed that SRB were performed significantly more during the baseline than the gum-feeder period in
13
14 199 CS ($p = 0.037$) but no significant differences were found for OB and Normann ($p > 0.05$).

20 201 *Gum-feeding in marmosets and tamarins*

21 202 Finally, we focused on the time spent feeding on gum Arabic- by the study subjects. The median time spent
22
23 203 feeding on gum Arabic was 42.5 (90.8) seconds for tamarins and 235 (255.5) seconds for marmosets.
24
25 204 Marmosets tended to spend more time eating gum Arabic than tamarins ($U = 7$, $p = 0.056$). When
26
27 205 considering gum feeding during the day within each group, the median time spent feeding on gum Arabic by
28
29 206 tamarins was 31.5 (91.8) seconds in the morning and 0 (20) seconds in the afternoon. In the case of
30
31 207 marmosets, the median time spent feeding on gum Arabic was 116 (207) seconds in the morning and 114
32
33 208 (281) seconds in the afternoon. Wilcoxon tests revealed no significant differences between morning and
34
35 209 afternoon in both tamarins ($Z = -1.572$, $p = 0.116$) and marmosets ($Z = -0.365$, $p = 0.715$) (**Figure 3**).

37 38 39 40 211 **Discussion**

41
42 212 This study aimed to investigate the effects of a gum-feeder, a wooden disc drilled with holes, on the behavior
43
44 213 of zoo tamarins and marmosets, to assess their welfare. Moreover, the study aimed to compare the time spent
45
46 214 feeding on gum Arabic between these species. First, each species interacted with the gum-feeders performing
47
48 215 a new behavior to obtain gum Arabic, confirming the enriching role on the animal daily routine and feeding
49
50 216 strategies of this device (McGrew *et al.*, 1986; Roberts *et al.*, 1999; Huber & Lewis, 2011). All the study
51
52 217 monkeys interacted with the gum-feeder and showed the behavior of eating Arabic gum from the holes. In
53
54 218 particular, marmosets and tamarins had to cling on to the wooden disc holding with one or both hands and
55
56 219 then retrieve gum directly with the mouth or grasping it with one hand. When retrieving the gum, marmosets
57
58 220 gouged and scraped the disc moving the head and mouth similarly as in the tree-gouging behavior reported in
59
60 221 the wild: indeed, the jaws were widely open around the gum in the hole, the upper jaw anchored on the

1
2
3
4 222 wooden disc whereas the lower jaws indented the area around the hole, favoring both gouging and scraping
5
6 223 movements (Rylands, 1984; Vinyard *et al.*, 2003; Vinyard *et al.*, 2004; Burrows & Nash, 2010; Thompson *et*
7
8 224 *al.*, 2014). This movement was less pronounced in tamarins as they simply retrieved the superficial gum
9
10 225 coming out the holes and consumed it.

12
13 226 First, we found that subjects were out-of-sight (“not observed”) more during the baseline than during
14
15 227 the gum-feeder period. This result seems to suggest that the gum-feeder could enhance the welfare of the
16
17 228 study subjects, engaging them in the performance of species-specific behaviors and increasing the time they
18
19 229 were visible to the public. This finding is consistent with previous research on gum-feeder provision in
20
21 230 marmosets, reporting an increased animal visibility (Kelly, 1993).

23 231 Regarding individual behaviors, when data on tamarins and marmosets were pooled together, we
24
25 232 found that feeding/foraging behaviors as well as maintenance behaviors were performed significantly less
26
27 233 during the gum-feeder period than during the baseline. Within each group (tamarins and marmosets), we
28
29 234 found the same patterns and significant differences for these categories in tamarins, whilst they did not differ
30
31 235 significantly in marmosets. Gum Arabic is a high-energy food source, rich in carbohydrates and minerals,
32
33 236 requiring longer time to be digested than other food items (Kelly, 1993; Power, 2010; Bairrão Ruivo, 2010).
34
35 237 It is possible that the provision of this food item in the new gum-feeder elicited tamarin and marmoset
36
37 238 interest, inducing them to eat gum Arabic before the rest of the food. Given the nutritional properties of the
38
39 239 gum (Power & Oftedal, 1996; Power, 2010), we speculate that a small amount of this food could be
40
41 240 sufficient for these small primates, leading to the reduction in the overall time spent feeding and foraging in
42
43 241 the second period. Also, it is possible that the interest toward the gum feeder as a novel device might supply
44
45 242 the callitrichines with the need of looking for food elsewhere in the enclosure (*e.g.* bowls, usual feeding
46
47 243 points). In other words, the study monkeys may have spent less time feeding and foraging in the second
48
49 244 period because they consumed gum Arabic first, and the same could extend to the decrease in maintenance
50
51 245 behavior during the second period. However, the presence of the gum feeder increased visual exploration,
52
53 246 which was performed significantly more during the gum-feeder period than during the baseline. This
54
55 247 behavior is particularly relevant in callitrichines as they are curious, attracted to novel objects and aware of
56
57 248 their surroundings (Menzel & Menzel, 1979). Such features of these small-bodied primates have been related
58
59 249 to improved vigilance and anti-predator behaviors (Caine, 1984). Therefore, the gum-feeder seemed to
60

Regaiolli *et al.*

1
2
3
4 250 encourage the performance of species-specific behaviors that are particularly important for the survival of
5
6 251 the species in the wild (Caine, 1984).

8 252 The presence of the gum-feeder also led to a significant decrease in self-directed behaviors, which
9
10 253 have been described as potential behavioral indicators of stress and anxiety in non-human primates, at least
11
12 254 in some situations (Maestriperi *et al.*, 1992; Leeds & Lukas, 2018). As our results underlined a decrease in
13
14 255 the performance of these behaviors, the presence of the gum-feeder seems to positively impact the behavior
15
16
17 256 of the study monkeys (Spiezio *et al.*, 2017; Leeds & Lukas, 2018). However, the study monkeys spent a
18
19 257 relatively low amount of time performing self-directed behaviors in both the baseline and the gum-feeder
20
21 258 periods (4% and 3% of the total observation time respectively) suggesting that this statistically significant
22
23 259 change may not necessarily be biologically important.

25 260 Regarding social behaviors, during the second period, a significant decrease of interspecific
26
27 261 behaviors, including interacting with or directing attention to humans was also reported. As visitors may be
28
29 262 distressing for the study subjects, this finding seems to highlight that the gum feeder might help tamarins and
30
31 263 marmosets to better cope with humans, promoting the performance of species-specific social behaviors and
32
33 264 discouraging possibly deleterious human-animal interactions. Together with the decrease in the time spent
34
35 265 out-of-sight in presence of the feeder, this result highlights a possible positive effect of gum-feeding devices
36
37
38 266 on callithricines' well-being in zoo environments.

40 267 Except for the decrease in feeding/foraging and maintenance behavior reported in the gum-feeder
41
42 268 period, no significant differences between periods were found in tamarins. The decrease in feeding/foraging
43
44 269 behaviors within tamarins seems consistent with the same result obtained when data from both species were
45
46 270 pooled together. Indeed, tamarins do not have as many adaptations for gum feeding and digestion as
47
48 271 marmosets (Heymann & Smith, 1999; Bairrão Ruivo, 2010; Hosey *et al.*, 2013; Roberts *et al.*, 1999; Taylor,
49
50 272 2002); therefore, consuming the gum before other food items due to the new presentation modality might
51
52 273 have reduced the tamarin appetite and the performance of other food-related behavior, but had no effect on
53
54 274 gummivore marmosets. However, the lack of significant differences in marmosets might be due to the small
55
56
57 275 sample size and thus further research is needed to better investigate this aspect.

58
59 276 On the other hand, marmosets were out-of-sight ("not observed") significantly more during the
60
277 baseline than the gum-feeder period, suggesting that for these species the presence of the drilled wooden

1
2
3
4 278 discs might help to increase the visibility of these species to zoo visitors. Inactive behaviors, such as
5
6 279 sleeping, were rather uncommon in the study subjects, as tamarins and marmosets tend to be very active
7
8 280 during the day. However, the study subjects had rest in the wooden nest boxes in the highest part of the
9
10 281 enclosure where the observer was not able to see them. Therefore, the behavioral category “not observed”
11
12 282 also included resting, which was otherwise not recorded in the study periods. For this reason, it is possible,
13
14 283 by decreasing the time spent out-of-sight, the presence of the gum-feeder also reduced inactive behavior of
15
16
17 284 the study subjects, especially within marmosets. This is consistent with findings by other authors on common
18
19 285 marmosets (*Callithrix jacchus*) (Roberts *et al.*, 1999).
20

21 286 When analyzing the behavior of each study subject during the two periods, we found that in three
22
23 287 red-handed tamarins, the gum-feeder had a positive effect as reduced SRBs, such as self-directed behaviors
24
25 288 and coprophagy. In addition, red-handed tamarins performed auto-coprophagy, which has been previously
26
27 289 related to physical or psychological stress in non-human primates (Anderson *et al.*, 1991; Krief *et al.*, 2004;
28
29 290 Prates & Bicca-Marques, 2005). One of the main effects of the gum-feeder was the reduction of SRBs,
30
31 291 including coprophagy. This was reported in other studies on gum-feeders in the red-handed tamarin,
32
33 292 suggesting that gum-related enrichment might be particularly relevant to prevent coprophagy in this species
34
35 293 (Taylor, 2002), and on laboratory common marmosets, where similar gum-feeding devices decreased
36
37
38 294 stereotypies (Roberts *et al.*, 1999).
39

40 295 Based on data collected during the second period, this study aimed at verifying the presence of
41
42 296 differences in gum-feeding between tamarins and marmosets and investigate whether the time spent feeding
43
44 297 on gum Arabic differed between morning and afternoon within each group. First, although no statistically
45
46 298 significant difference was found, marmosets tended to feed on gum more than tamarins. This finding agrees
47
48 299 with previous work on these species, as gummivore marmosets spend a large amount of time feeding on
49
50 300 gum, based on the importance of this food in their diet, whereas tamarins rely more on fruit and invertebrates
51
52 301 and cannot easily obtain and digest plant exudate (Kelly, 1993; Heymann & Smith, 1999; Bairrão Ruivo,
53
54 302 2010). However, it could not be excluded that the greater interaction of the marmosets with the feeder was
55
56 303 due to species differences in approaching the new enrichment devices, as previously suggested (Renner *et*
57
58 304 *al.*, 2000).
59
60

Regaiolli *et al.*

1
2
3
4 305 Similarly, no differences in time spent feeding on gum Arabic were found between mornings and
5
6 306 afternoons, neither within marmosets nor within tamarins. In the wild, marmosets feed on exudates
7
8 307 frequently throughout the daylight hours, as their gastrointestinal system is well-adapted to gum digestion
9
10 308 (Kelly, 1993; Heymann & Smith, 1999; Bairrão Ruivo, 2010). Therefore, the gum-feeding behavior of the
11
12 309 study Geoffroy's and pygmy marmosets resembles that reported in the wild, as the study subjects ate gum
13
14 310 Arabic to the same extent in both the morning and the afternoon. In the case of tamarins, these species are
15
16 311 known to eat exudates mainly during the afternoon, so that they can digest this food item overnight (Kelly,
17
18 312 1993; Heymann & Smith, 1999; Bairrão Ruivo, 2010). However, we collected data on the time spent eating
19
20 313 gum but not the amount of gum that was eaten. Again, providing gum in the new gum feeders might have
21
22 314 incentivized the consumption of this food item regardless of the time of the day.

23
24
25 315 Our findings seem to highlight positive effects of the gum-feeder on the behavior of zoo
26
27 316 callitrichines, as previously reported in the species considered in the current study as well as in other
28
29 317 species of marmosets and tamarins (McGrew *et al.*, 1986; Roberts *et al.*, 1999; Taylor, 2002; Huber &
30
31 318 Lewis, 2011). However, the small sample size and the relatively small amount of data collected in our study
32
33 319 hamper the ability to draw firm conclusions and future research is needed to investigate more deeply the
34
35 320 gum-feeding behavior of zoo callitrichines, focusing on larger samples and longer periods of time.

36 37 38 321 **Conclusion**

39
40 322 **Conclusion**
41
42 323 In conclusion, our findings suggest that providing zoo tamarins and marmosets with gum Arabic in a
43
44 324 naturalistic and innovative way, such as using a wooden gum feeder, might enhance the welfare of these
45
46 325 species by:

- 47
48 326 1) promoting the performance of species-specific behaviors, such as visual exploration and decreasing
49
50 327 self-directed behaviors, that can in some cases indicate a stressful situation of the animal;
- 51
52 328 2) improving the visibility to zoo visitors by decreasing the time spent out-of-sight and in interspecific
53
54 329 interactions;
- 55
56 330 3) reducing the performance of abnormal behaviors, as reported in the study of red-handed tamarins;
- 57
58 331 4) increasing the consumption of this crucial food item, at least in the first stages of the introduction of
59
60 332 the feeder.

Regaiolli *et al.*

15

1
2
3
4 333 Our results add to previous literature on gum-feeding devices in tamarins and marmosets, highlighting that
5
6 334 the wooden gum feeders can improve the welfare of these species in controlled environments by stimulating
7
8 335 natural behaviors and promoting naturalistic feeding strategies (McGrew *et al.*, 1986; Roberts *et al.*, 1999;
9
10 336 Taylor, 2002; Huber & Lewis, 2011).
11

12
13 337
14
15 338 **Acknowledgements**

16
17 339 We would like to thank Cesare Avesani Zaborra and Camillo Sandri for allowing this study to take place at
18
19 340 Parco Natura Viva and for supporting this project; Carolina Sammarini and Daniela Galietta, New-World
20
21 341 primate zookeepers, as well as Giuseppe Federico La Cauza and William Magnone, zoo veterinarians, for
22
23 342 their help during the study period; and Tim Baldwin for his invaluable suggestions on gum Arabic. Finally,
24
25 343 we thank two anonymous reviewers that helped us improve the manuscript with their constructive comments.
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 Regaiolli *et al.*

2

3

4

4 345

5

6 346

7

8 347

9

10 348

11

12 349

13

14 350

15

16 351

17

18 352

19

20 353

21

22 354

23

24 355

25

26 356

27

28 357

29

30 358

31

32 359

33

34 360

35

36 361

37

38 362

39

40 363

41

42 364

43

44 365

45

46 366

47

48 367

49

50 368

51

52 369

53

54 370

55

56 371

57

58 372

59

60

References

Abbott DH, Barnett DK, Colman RJ, Yamamoto ME, Schultz-Darken NJ. Aspects of common marmoset basic biology and life history important for biomedical research. *Comp Med.* 2003; 53:339–350.

Altman J. 1974. Observational study of behavior: sampling methods. *Behaviour* 49: 227-267.

Anderson, J. R., C. Combette, and J. J. Roeder. 1991. Integration of a tame adult female capuchin monkey (*Cebus apella*) into a captive group. *Prim. Rep* 31:87–94.

Bairrão Ruivo E. 2010. *EAZA Husbandry Guidelines for Callitrichidae, 2nd edition*. Beauval Zoo.

Buckner JC, Lynch Alfaro J, Rylands AB, Alfaro ME. 2015. Biogeography of the marmosets and tamarins (*Callitrichidae*). *Molecular Phylogenetics and Evolution*, 82: 413– 425.

Burrows AM, Nash LT. 2010. *The Evolution of Exudativory in Primates. Developments in Primatology: Progress and Prospects*. Springer, NY.

Caine, N.G. (1984) Visual scanning by tamarins: a description of the behavior and tests of two derived hypotheses. *Folia Primatol.* 43, 59–67

Coimbra-Filha AF, Mittermeier RA. 1977. Tree-gouging, exudate-eating, and the short-tusked condition in *Callithrix* and *Cebuella*. In: Kleiman DC, editor. *The biology and conservation of the Callitrichidae*. Washington DC: Smithsonian Institution Press, p 105-115.

Emmons LH. 1990. *Neotropical Rainforest Mammals: a Field Guide*. Chicago (US) and London (UK): The University of Chicago Press.

Regaiolli *et al.*

17

1
2
3
4 373 Eng CM, Ward SR, Vinyard CJ, Taylor AJ. 2009. The morphology of the masticatory apparatus facilitates
5
6 374 muscle force production at wide jaw gapes in tree-gouging common marmosets (*Callithrix jacchus*). *Journal*
7
8 375 *of Experimental Biology*. 212(24): 4040-55.

10 376
11
12 377 Fisch GS. 2001. Evaluating data from behavior analysis: Visual inspection or statistical models. *Behavioural*
13
14 378 *Processes*. 54: 137-154.

16
17 379
18
19 380 Fontani S., Vaglio S., Beghelli V., Mattioli M., Bacci S. & Accorsi PA. 2014 Fecal concentrations of
20
21 381 cortisol, testosterone, and progesterone in cotton-top tamarins housed in different zoological parks:
22
23 382 relationships among physiological data, environmental conditions and behavioral patterns. *Journal of*
24
25 383 *Applied Animal Welfare Science* 17: 228-252.

27 384
28
29 385 Garber PA. 1993. Feeding ecology and behavior of the genus *Saguinus*. In: Rylands AB, editor. *Marmosets*
30
31 386 *and Tamarins: Systematic, Behavior, and Ecology*. Oxford (England): Oxford University Press.

33
34 387
35
36 388 Gashua IB, Williams PA, Yadav MP, Baldwin TC (2015) Characterisation and molecular association of
37
38 389 Nigerian and Sudanese *Acacia* gum exudates. *Food Hydrocolloids* 51: 405-413.

39
40 390
41
42 391 Goodrum LJ, Patel A, Leykam JF, Kieliszewski MJ (2000) Gum arabic glycoprotein contains glycomodules
43
44 392 of both extensin and arabinogalactan-glycoproteins. *Phytochemistry* 54:99–106.

46 393
47
48 394 Groves C. 2001. *Primate taxonomy*. Washington DC (US): Smithsonian Institute Press.

50 395
51
52 396 Heymann EW, Smith AC. 1999. When to feed on gums: temporal patterns of gummivory in wild tamarins,
53
54 397 *Saguinus mystax* and *Saguinus fuscicollis* (*Callitrichinae*). *Zoo Biology* 18: 459-471.

56
57 398
58
59 399 Hill, SP, Broom DM. 2009. Measuring zoo animal welfare: theory and practice. *Zoo Biology* 28:531–544.

60
400

Regaiolli *et al.*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

401 Hosey G, Melfi V, Pankhurst S. 2013. *Zoo animals behavior, management and welfare*. Oxford (England):
402 Oxford University Press.
403
404 Huber, H. F. & Lewis, K. P. 2011. An assessment of gum-based environmental enrichment for captive
405 gummivorous primates. *Zoo Biology* 30: 71-78.
406
407 Kelly K. 1993. Environmental enrichment for captive wildlife through the simulation of gum feeding.
408 *Animal Welfare Information Center Newsletter* 4: 1-2, 5-10.
409
410 Krief, S., Jamart, A. and Hladik, C. M. 2004. On the possible adaptive value of coprophagy in free-ranging
411 chimpanzees. *Primates* 45: 141–145.
412
413 Leeds, A. & Lukas, K.E. (2018). Experimentally evaluating the function of self-directed behaviour in two
414 adult mandrills (*Mandrillus sphinx*). *Animal Welfare* 27(1): 81-86.
415
416 Lutz CK. 2014. Stereotypic behavior in non-human primates as a model for the human condition. *ILAR*
417 *Journal* 55: 284–296.
418
419 Lutz CK. 2018. A cross-species comparison of abnormal behavior in three species of singly-housed old
420 world monkeys. *Applied Animal Behavior Science* 199:52-58.
421
422 Maestriperi D, Schino G, Aureli F, Troisi A. 1992. A modest proposal: displacement activities as an
423 indicator of emotions in primates. *Animal Behaviour*, 44(5): 967-979.
424
425 Manteca X., Amat M., Salas M., Temple D. 2016. Animal-based indicators to assess welfare in zoo animals.
426 *CAB Reviews*, 11:1–10.

Regaiolli *et al.*

19

- 1
2
3
4 428 Marx, A.; Backes, C.; Meese, E.; Lenhof, H. P.; and Keller, A. (2016). EDISON-WMW: Exact Dynamic
5
6 429 Programming Solution of the Wilcoxon-Mann-Whitney Test. *Genomics Proteomics Bioinformatics*, 14(1):
7
8 430 55-61.
9
10 431
11
12 432 McGraw WC, Brennan JA, Russell C. 1986. An artificial gum tree for captive marmosets. *Zoo Biology* 5:
13
14 433 45-50.
15
16 434
17
18
19 435 Mellor, D. & Beausoleil, N. (2015) Extending the 'Five Domains' model for animal welfare assessment to
20
21 436 incorporate positive welfare states. *Animal Welfare* 24: 241-253.
22
23 437
24
25 438 Mench, J.A. (1994). Environmental enrichment and exploration. *Laboratory Animal*, 38-41.
26
27 439
28
29 440 Menzel, E. W., and Menzel, C. R. (1979). Cognitive, developmental and social aspects of responsiveness to
30
31 441 novel objects in a family group of marmosets (*Saguinus fuscicollis*). *Behaviour* 70: 251–279.
32
33 442
34
35 443 Pack KS, Henry O, Sabatier D. 1999. The insectivorous-frugivorous diet of the golden-handed tamarin
36
37 444 (*Saguinus midas midas*) in French Guiana. *Folia Primologica* 70: 1-7.
38
39 445
40
41 446 Passamani M, Rylands AB, 2000. Feeding behavior of Geoffroy's marmosets (*Callithrix geoffroyi*) in
42
43 447 Atlantic forest fragment of south-eastern Brazil. *Primates* 41: 27-38.
44
45 448
46
47 449 Power ML. 1996. The Other Side of Callitrichine Gummivory. In: Norconk MA, Rosenberger AL,
48
49 450 Garber PA, editors. *Adaptive Radiations of Neotropical Primates*. Boston, MA, Springer,
50
51 451
52
53 452 Power ML (2010) Nutritional and digestive challenges to being a gum feeding Primates. In: Burrows A, Nash
54
55 453 L, editors. *The evolution of exudativory in Primates*. New York: Springer. 25–44.
56
57
58
59 454
60

Regaiolli *et al.*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

455 Power ML, Oftedal OT (1996) Differences among captive callitrichids in the digestive responses to dietary
456 gum. *Am. J. Primatol* 40: 131–144.

457

458 Prates, HM, Bicca-Marques JC, 2005. Coprophagy in Captive Brown Capuchin Monkeys (*Cebus apella*),
459 *Neotropical Primates*, 13: 18-21.

460

461 Renner MJ, Feiner AJ, Orr MG, Delaney BA. Environmental enrichment for new world primates:
462 introducing food-irrelevant objects and direct and secondary effects. *J Appl Anim Welfare Sci* 2000;3(1):23 –
463 32.

464

465 Roberts RL, Roytburd LA, Newman JD. 1999. Puzzle feeders and gum-feeders as environmental enrichment
466 for common marmosets. *Contemporary Topics in Laboratories Animal Science*, 38(5): 27-31.

467

468 Rosenberger AL. 1992. Evolution of feeding niches in New World Monkeys. *American Journal of Physical*
469 *Anthropology* 88: 525–562.

470

471 Rylands AB (1984) Exudate-eating and tree-gouging by marmosets (Callitrichidae, Primates). In: Chavdwick
472 AC, Sutton SL editors. *Tropical Rain Forest: The Leeds Symposium Leeds Philosophical and Literary*
473 *Society*. 155–158.

474

475 Siegel S, Castellan N.J. 1992. *Statistica non parametrica*. Milano (IT). McGraw-Hill.

476

477 Smuts BB, Cheney DL, Seyfarth RM, Wrangham RW. 1986. *Primate Societies*. Chicago (US) and London
478 (UK): University of Chicago Press.

479

480 Spiezio C, Vaglio S, Scala C, Regaiolli B. 2017. Does positive reinforcement training affect the behaviour
481 and welfare of zoo animals? The case of the ring-tailed lemur (*Lemur catta*). *Applied Animal Behaviour*
482 *Science* 196: 91-99.

- 1
2
3
4 483
5
6 484 Tardif SD. 1984. Social influences on sexual maturation of female *Saguinus oedipus oedipus*. *Am J Primatol*
7
8 485 6(3): 199-209.
9
10 486
11
12 487 Tardif SD, Araujo A, Arruda MF, French JA, Sousa MBC, Yamamoto ME. 2008. Reproduction and aging in
13
14 488 marmosets and tamarins. In: Atsalis S, Margulis SW, Hof PR, editors. *Primate Reproductive Aging*.
15
16 489 *Interdisciplinary Topics in Gerontology*. vol 36. Basel: Karger;. pp. 29–48.
17
18
19 490
20
21 491 Tardif SD, Mansfield KG, Ratnam R, Ross CN, Ziegler TE. The marmoset as a model of aging and age-
22
23 492 related diseases. *ILAR J*. 2011;52: 54–65.
24
25 493
26
27 494 Taylor, T.D. 2002. Feeding Enrichment for Red-Handed Tamarins. *The Shape of Enrichment: A. Quarterly*
28
29 495 *Source of Ideas for Enrichment* 11(2): 1-3.
30
31 496
32
33 497 Taylor AB, Vinyard C. 2004. Comparative analysis of masseter fiber architecture in tree-gouging (*Callithrix*
34
35 498 *jacchus*) and non-gouging (*Saguinus oedipus*) callitrichids. *Journal of Morphology* 261: 276-284.
36
37
38 499
39
40 500 Thompson, C.L., M.M. Valença-Montenegro, L. César de Oliveira Melo, Y.B. Maranhão Valle, M.A.
41
42 501 Borstelmann de Oliveira, P.W. Lucas, and C.J. Vinyard (2014). Accessing Foods Can Exert Multiple
43
44 502 Distinct, and Potentially Competing, Selective Pressures on Feeding in Common Marmoset Monkeys.
45
46 503 *Journal of Zoology*, 294: 1-9.
47
48 504
49
50 505 Troisi A. 2002. Displacement activities as a behavioral measure of stress in non-human primates and human
51
52 506 subjects. *Stress* 5: 47-54.
53
54
55 507
56
57 508 Troisi A, Schino G. 1987. Environmental and social influences on autogrooming behaviour in a captive
58
59 509 group of Java monkeys. *Behaviour* 100: 292-302.
60
510

Regaiolli *et al.*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

511 Vinyard CJ, Lucas PW, Valença-Montenegro MM, Melo LCO, Valle YM, Monteiro da Cruz MAO. 2004.
512 Where the wild things are: Linking lab and field work in studying tree gouging in common marmosets
513 (*Callithrix jacchus*). *American Journal of Physical Anthropology* 38: 200–201.

514
515 Vinyard CJ, Wall CE, Williams SH, Hylander, WL. 2003. Comparative functional analysis of skull
516 morphology in tree-gouging primates. *American Journal of Physical Anthropology* 120:153–170.

517
518 Ziegler TE, Savage A, Scheffler G, Snowdon CT. 1987. The endocrinology of puberty and reproductive
519 functioning in female cotton-top tamarins (*Saguinus oedipus*) under varying social conditions. *Biol Reprod*
520 37(3): 618-27.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



188x176mm (150 x 150 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 1: The gum-feeder provided to marmosets and tamarins. In the gum-feeder period, tamarins and marmosets were provided with wooden drilled discs, hanged in the outdoor enclosure and filled with Arabic gum. The picture shows one of the study Geoffroy's marmoset interacting with the feeder.

188x176mm (150 x 150 DPI)

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2: Behaviors of the study marmosets and tamarins. The bars report the % total duration in seconds of individual and social behaviors as well as “not observed” in the baseline and in the gum-feeder. Asterisks indicate categories that differed significantly between periods (Wilcoxon test: $p < 0.05$).

336x146mm (150 x 150 DPI)

Species	Name	Sex	Age
	Mum	F	Adult
Cotton-top tamarin <i>(Saguinus oedipus)</i> <i>N = 4</i>	Franca	F	Juv
	Rubik	F	Juv
	Dad	M	Adult
	CS	F	Adult
Midas tamarin <i>(S. midas)</i> <i>N = 4</i>	OB	F	Adult
	Norman	M	Adult
	CC	M	Adult
	Mum	F	Adult
Geoffroy's marmoset <i>(Callithrix geoffroyi)</i> <i>N = 3</i>	Dad	M	Adult
	Sbiru	M	Juv
Pygmy marmoset <i>(Cebuella pigmea)</i> <i>N = 2</i>	Peace	F	Adult
	Love	M	Adult

Table 1: Tamarins and marmosets involved in the study. For each species the table reports the subject name, sex (F = female; M = male) and age class (Adult: > 2 years of age; Juv: < 2 years of age).

Individual behaviours

Abnormal behaviour	Coprophagy.
Alert	Being watchful.
Exploration/Play	Exploring the environment, hunting, manipulating enrichment devices, leaves, twigs and other objects found in the enclosure by sniffing, biting, chewing, gouging, handling, pouncing on, grappling with.
Feeding/Foraging	Eating food found in the enclosure, either in bowls or foraging on trees, ground and other substrates.
Gum-feeder*	Eating gum from the gum-feeders (second period) .
Locomotion	Moving around in the enclosure, walking, running or jumping along trees or walls of the enclosure
Maintenance	Peeing, defecating, drinking.
Scent-marking	Marking branches, shelves, ropes and other substrates with ano-genital, supra-pubic or sternal glands.
Self-grooming directed b.	Cleaning or licking the own hair or other parts of the body, scratching with the hands or with the legs.
Visual exploration	Looking around quietly.

Social behaviour ~~behavior~~

Affiliative b.	Allo-grooming (using the hands and/or mouth to clean the partner fur or other parts of the body), being in contact with conspecifics, nuzzling (rubbing or pushing gently with the nose and mouth), anogenital inspection (orienting the face against or toward anogenital region of partner or use hands or mouth to investigate anogenital region of partner), stealing food/objects from the hands or from the mouth.
Agonistic b.	Fighting (biting, clawing, and wrestling), attacking (lunging at or pouncing on partner aggressively), cuffing, chasing, threatening (staring with lower eyebrows, furl brow, tongue flicking), mounting between same-sex individuals.
Interspecific b.	Interacting with or directing attention toward individuals of different species, such as humans (visitors, zoo-keepers zookeepers) or Azara agoutis (only in the Geoffroy's marmoset enclosure).

Not visible

Not observed	The individual is not visible to the observer or it is not possible to identify the behaviour behavior being performed.
--------------	--

*This behavioural category was recorded only in the second period, when ~~Arabic~~ gum ~~arabic~~ was put in the gum feeder and it was possible to detect clearly that the subjects were feeding on that food item.

Table 2: Study ethogram. Individual and social behaviours collected in the study periods.

	All			Tamarins			Marmosets		
<i>Individual behavior</i>	Baseline	GF	Z and p-value	Baseline	GF	Z and p-value	Baseline	GF	Z and p-value
Abnormal	0 (0)	0 (9)	#	0 (60.8)	0 (79.5)	#	0 (0)	0 (0)	#
Alert	7 (17.5)	0 (4.5)	$z = -1.868$, $p = 0.062$	8 (15.8)	4 (9.5)	$z = -1.016$, $p = 0.310$	5 (18.5)	0 (0)	$z = -1.826$, $p = 0.068$
Expl/play	869 (1307.5)	663 (612)	$z = -1.922$, $p = 0.055$	974.5 (1569.5)	857.5 (1308.5)	$z = -1.400$, $p = 0.161$	321 (1509)	494 (723.5)	$z = -1.214$, $p = 0.225$
Feeding/Foraging	1219 (590.5)	826 (665.5)	$z = -2.481$, $p = 0.031^*$	1312.5 (635.3)	1233.5 (864.5)	$z = -2.100$, $p = 0.036^*$	1081 (653.5)	763 (485.5)	$z = -1.483$, $p = 0.138$
Locomotion	2920 (980.5)	2859 (1572.5)	$z = -0.384$, $p = 0.701$	2958.5 (890.3)	3340.5 (1155.8)	$z = -2.260$, $p = 0.208$	2920 (1199.5)	1622 (1432)	$z = -1.483$, $p = 0.138$
Maintenance	57 (70)	10 (32)	$z = -3.111$, $p = 0.002^*$	83 (42.8)	13 (30.5)	$z = -2.521$, $p = 0.012^*$	19 (11.5)	0 (22.5)	$z = -1.761$, $p = 0.078$
Scent-marking	38 (36.5)	22 (28.5)	$z = -1.049$, $p = 0.294$	27.5 (36)	28 (32.8)	$z = -0.491$, $p = 0.624$	49 (55.5)	19 (37)	$z = -0.944$, $p = 0.345$
Self-groomingdirected	420 (240.5)	392 (337)	$z = -2.201$, $p = 0.028^*$	464 (257.3)	437.5 (183.3)	$z = -1.260$, $p = 0.208$	384 (250.5)	208 (354)	$z = -1.753$, $p = 0.080$
Visual expl	6891 (2946.5)	8097 (1760)	$z = -2.201$, $p = 0.028^*$	6578 (3024.8)	7860.5 (2860.3)	$z = -1.680$, $p = 0.093$	7080 (2402.5)	8119 (2034)	$z = -1.483$, $p = 0.138$
<i>Social behavior</i>									
Affiliative	764 (632.5)	1331 (1180.5)	$z = -1.712$, $p = 0.087$	704.5 (753.5)	776.5 (798.8)	$z = -0.560$, $p = 0.575$	1110 (964)	1797 (1269.5)	$z = -1.753$, $p = 0.080$
Agonistic	0 (34.5)	8 (28.5)	$z = -0.533$, $p = 0.594$	9.5 (43.3)	7 (20.8)	$z = -0.280$, $p = 0.779$	0 (21.5)	8 (54.5)	$z = -1.604$, $p = 0.110$
Interspecific	0 (69)	0 (0)	$z = -2.023$, $p = 0.043^*$	0 (59.3)	0 (0)	$z = -1.342$, $p = 0.180$	38 (94)	0 (0)	$z = -1.604$, $p = 0.110$
<u>Not observed</u>	<u>956 (881.5)</u>	<u>207 (553.5)</u>	<u>$z = -2.481$</u> , <u>$p = 0.013^*$</u>	<u>724.5 (846.5)</u>	<u>172 (633)</u>	<u>$z = -1.540$</u> , <u>$p = 0.124$</u>	<u>1117 (740)</u>	<u>303 (615)</u>	<u>$z = -2.023$</u> , <u>$p = 0.043^*$</u>

*Significant difference between the two periods ($p < 0.05$).

1
2 **Table 3:** Individual behaviors, ~~and~~ social behaviors and “not observed” in the baseline and in the gum-feeder period. For each behavioral category,
3
4 the table reports the median (interquartile range) duration in seconds calculated in the baseline and in the gum-feeder period (GF), as well as the z
5
6 and p values of the Wilcoxon test performed to compare the two periods. All: data collected on different species pulled together; Tamarins: analysis
7
8 performed within the tamarin group ($N = 8$) (*Saguinus oedipus*, *Saguinus midas*); Marmosets: analysis performed within the marmoset group ($N =$
9
10
11 5) (*Callithrix geoffroyi*, *Cebuella pigmeapygmaea*).
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46

For Peer Review