

Conductance of ‘bare-bones’ tripodal molecular wires

Ross J. Davidson,^{a*} David C. Milan,^b Oday A. Al-Owaedi,^{c,d} Ali K. Ismael,^{c,e} Richard J. Nichols,^{b*} Simon J. Higgins,^b Colin J. Lambert,^c Dmitry S. Yufit,^a Andrew Beeby,^{a*}.

^a*Department of Chemistry, Durham University, South Rd, Durham, DH1 3LE, UK*

^b*Department of Chemistry, University of Liverpool, Crown St, Liverpool, L69 7ZD, UK*

^c*Department of Physics, University of Lancaster, Lancaster LA1 4YB, U.K.*

^d*Department of Laser Physics, Women Faculty of Science, Babylon University, Hilla, Iraq.*

^e*Department of Physics, College of Education for Pure Science, Tikrit University, Tikrit, Iraq*

ABSTRACT: Controlling the orientation of molecular conductors on the electrode surfaces is a critical factor in the development of single-molecule conductors. In the current study, we used the scanning tunnelling microscopy-based break junction (STM-BJ) technique to explore ‘bare-bones’ tripodal molecular wires, employing different anchor groups (AGs) at the ‘top’ and ‘bottom’ of the tripod. The triarylphosphine tris(4-(methylthio)phenyl)phosphine and its corresponding phosphine sulfide showed only a single high conductance feature in the resulting 1- and 2-dimensional conductance histograms, whereas analogous molecules with fewer than three thiomethyl AGs did not show clear conductance features. Thus, by systematic molecular modifications and with the aid of supporting DFT calculations, the binding geometry, with respect to the surface, was elucidated.

KEYWORDS: STM, single molecule conductance, DFT, tripodal, phosphine

Introduction

The field of single molecule conductance has almost exclusively involved the study of one-dimensional molecular wires, with one anchor group (AG), e.g. pyridine, thiol or amine, attached at each end.^{1, 2} This motif has been favoured as it provides the least-complicated example for examining structure-property relationships and conductance mechanisms. However, to afford greater spatial control relative to the substrate it is necessary to increase the number of AGs to give tripodal conductors. Increasing the number of AGs significantly enhances the strength of attachment of the molecule to the surface(s), thereby increasing the mechanical stability of the metal|molecule|metal junction.^{3, 4} Recent studies demonstrated the use of tripodal molecules consisting of three monodentate AGs (e.g., pyridine, thiophene) attached to a central carbon or silicon atom, bridged by a conductor such as phenylene or phenylene ethynylene, to give conductance values of 2×10^{-5} and $5 \pm 1 \times 10^{-4} G_0$ respectively.^{3, 5} ⁶ However, in each reported example, two or more AGs were attached to the central conjugated unit via sp^3 hybridised carbon or silicon atoms (by the use of a carbon atom, spirofluorene, or adamantane motif),^{5, 7-9} reducing the degree of conjugation between each of the AGs.

To examine the electrical conductance of a tripodal molecular wire in its simplest form, we examined a 'bare-bones' structure consisting a triaryl phosphine with AGs on each of the aryl groups. Ragaini demonstrated that 4,4',4''-phosphanetriyltribenzenethiol can simultaneously coordinate to metal complexes such as $Ni(CO)_3$ via the phosphine, and chemisorb to a gold surface via the thiols to give a trigonal pyramidal structure on the surface.¹⁰

Parameswaran *et. al* has previously demonstrated that diphenylphosphine moieties can function as mechanically and chemically stable AGs with a tendency to bind to under-coordinated gold atoms on the surface.¹¹ In addition to enhanced stability compared with corresponding dialkylphosphines, diphenylphosphines couple well to the σ -system of alkanes

resulting in relatively high conductances for $\text{Ph}_2\text{P}(\text{CH}_2)_n\text{PPh}_2$ junctions. Additionally, phosphines can also be readily oxidised by sulfur to give the corresponding phosphine sulfide. Fukazara demonstrated that dibenzophospholesulfides act as strong AGs and when bridged by 1,4-phenylene and biphenyl-4,4'-diyl groups, give conductance values of 5×10^{-4} and $5 \times 10^{-5} G_0$ respectively and favour conductance via the LUMO.¹²

Here we report conductance measurements and calculations on ‘bare-bones’ tripodal molecular wires based on triaryl-phosphines, -phosphine sulfides, -phosphine oxides and -selenides using scanning tunnelling microscopy (STM)-based techniques and we examine how conductance is affected by the number of available AGs in the tripods making contact with the single apex terminal group. The triarylphosphines and their P(V) derivatives are a good test case for this due to their simplicity and the fact that they can be easily modified in a modular fashion.

Experimental Methods

Synthetic work

Compound **3** was prepared by the literature methods,¹⁰ the synthesis of compounds **1**, **2**, **1-3=S**, **1-3=Se**, **2=O**, **3=O**, and **[3-Me]⁺** are given in the ESI.

Conductance measurements

All measurements were performed at room temperature and under ambient conditions with an Agilent (Keysight) STM running Picoscan 5.3.3. software. The STM-BJ setup used to perform all the measurements consists of a home-made scanner¹³ controlled by an Agilent Picoscan 5500 controller with a N9503A modified scanner and a multichannel current amplifier.

Samples were prepared by adsorption onto commercially-obtained gold-on-glass Au(111) substrates prepared with a chromium adhesion layer (Arrandee). The gold slides were flame annealed with a butane torch. The process involved heating the gold-on-glass slide in a dark room until it shows an orange glow, then allowing the slide to cool; this process is repeated as many times as necessary to obtain the desired Au(111) flat surface.¹⁴ Sample solutions were prepared by dissolving the amount of compound (a few milligrams) in 10 mL of mesitylene needed to achieve a 10^{-4} M concentration. Gold STM tips were fabricated from 0.25 mm Au wire (99.99 %) that was freshly mechanically cut for each STM experiment. Conductance experiment were carried out adding 100 μ L of the 10^{-4} M solution into the STM-BJ liquid cell, then the STM-BJ tip was repeatedly approached into contact and then withdrawn from the Au(111) surface. A current set point of 100 μ A was used to achieve a good gold contact with the surface, and the tip was then retracted 4 nm away from the surface at a rate of 20 nm s⁻¹. A bias voltage of 100 mV was applied between the sample and tip; current–distance traces were recorded during the approach to contact and the withdrawal process.

Results and Discussion

Synthesis. A family of triaryl phosphines and their oxidised derivatives were chosen as the tripods with aryl groups consisting of combinations of either thioanisole or phenyl groups. Thiomethyl AGs have been chosen due to their chemical stability and their proven record as contact groups in molecular junctions.¹⁵ Each of the phosphines were synthesised by carrying out halogen-lithium exchange on (4-bromophenyl)(methyl)sulfane, followed by reaction with PCl_3 , dichlorophenylphosphine or chlorodiphenylphosphine.^{10, 16} The phosphines (**1-3**) were then heated in the presence of either sulfur (to produce the respective phosphine sulfides **1=S**, **2=S** and **3=S**) or selenium (to produce the phosphine selenides **1=Se**, **2=Se** and **3=Se**). The

phosphine oxides **2=O** and **3=O** were prepared by reaction of 4-(methylthio)phenyllithium with phenylphosphinic dichloride or phosphoryl bromide respectively, to avoid complications from oxidation of thiomethyl groups that arose upon attempted H₂O₂ oxidation of **2** and **3**. To aid control experiments (*q.v.*), the methylphosphonium iodide salt of **3** (**[3-Me]⁺**) was synthesised using the standard techniques of reacting **3** with methyl iodide.




Chart 1. Triaryl phosphines and their derivatives used in this investigation

Molecular Conductance

The single-molecule conductances of tripodal compounds were determined using the STM-BJ method.¹⁷ Conductance values and break-off distances (95th percentile) are summarized in Table 1. Conductance histograms for the compounds (**2**, **3**, **3=S**, and **3=O**) are shown in Figure 1. Representative conductance traces with current plateaus, two-dimensional (2D) histograms; conductance values and break-off distances (95th percentile) are shown in the ESI.

Each of the three phosphine selenide compounds (**1=Se** – **3=Se**) showed a complete absence of molecular conductance peaks. In addition, in experiments with the phosphine selenides, the G_0 peaks in the histograms were significantly altered, with much-diminished peak heights, indicating that the gold surface had been chemically altered. The most likely cause of this alteration is ‘selenium poisoning’ resulting from the surface-induced decomposition of the phosphine selenide. In fact, in the present study, we examined the stability of these compounds by a ^1H and ^{31}P NMR study in CDCl_3 ; see ESI. This analysis shows that the compounds decayed over a number of hours, primarily to the corresponding phosphine oxide liberating the selenium.

Table 1. Conductance values and break-off distances (95th percentile) for **2**, **3**, **3=S** and **3=O**.

Molecule	Conductance (G_0)	Break-off distance (nm)
1	No peak	
2	3.46×10^{-5}	0.76
	5.88×10^{-4}	0.94
3	4.36×10^{-3}	0.76
1=S	No peak	
2=S	No peak	
3=S	1.23×10^{-3}	0.87
3=O	4.46×10^{-5}	0.91
	1.09×10^{-3}	0.76
2=O	No peak	
[3-Me]⁺	No peak	

In contrast, the **3** and **3=S** compounds showed single, well-defined conductance peaks, at $10^{-2.36}$ and $10^{-2.91} G_0$ respectively, similar in value to 1,4-bis(methylthio)benzene ($10^{-2.15} G_0$).¹⁸ However, replacement of a single thiomethyl group with a hydrogen resulted in two

conductance peaks being observed for compound **2** ($10^{-3.23}$ and $10^{-4.46}$ G_0), and no conductance peaks being observed for compound **2=S**.




Figure 1. Conductance histograms for the compounds **2**, **3**, **3=S** and **3=O**.

Further replacement of the thiomethyl groups with hydrogens (i.e., **1** and **1=S**) resulted in no molecular conductance peak being observed, suggesting that the conductance path is not simply that between one single thiomethyl group and the central donor group (either phosphorus or phosphorus sulfide). To further elucidate the nature of the central atom involvement in junction formation, we performed STM-BJ studies on the corresponding phosphine oxides (compounds **2=O** and **3=O**) and on the methylphosphonium iodide compound [**3-Me**]**I**. Compound **3=O** displayed conductance peaks at $10^{-2.96}$ and $10^{-4.35}$, while compound **2=O** displayed no conductance peak, as for the sulfide **2=S**. The presence of a second peak for **3=O** suggests that the phosphine oxide does not bind to the surface as strongly as the sulfide, allowing the formation of other contact geometries, possibly consisting of a thiomethyl-molecule-thiomethyl contacted arrangement based on the increased break-off distance for the low conductance band. Finally, [**3-Me**]⁺ showed no detectable conductance peak, which suggests that junction formation *via* thiomethyl groups at each contact does not contribute significantly

to the conductance peaks in these molecules. Rather, for the highest conductance mode to be reached, the central atom (P or P=S) must be able to act as a contact for one electrode, and all three thiomethyl groups must be available to make contact with the surface, giving the tripodal shape of the molecule. Indeed, previous work by Ragaini demonstrated that on a similar compound (4,4',4''-phosphanetriyltribenzenethiol),¹⁰ contact occurs via the aryl thiols leaving the central donor phosphorus atom available to act as a binding site able to coordinate to metal carbonyls.

To account for the low conductance mode that occurs when only two thiomethyl groups are attached, we propose a model in which one P-C₆H₄-SMe lies flat on the gold surface, allowing the lone pair in conjunction with the π -orbitals of the thioanisole to make contact with the surface. This arrangement would leave a single thioanisole raised perpendicular to the surface, providing a contact for the second electrode. In fact, similar behaviour has been observed by Su et al.,¹⁹ whereby the 1,1-bis(4-(methylthio)phenyl)silane system displayed two conductance peaks, which were an order of magnitude different. The low conductance peak corresponded to the end-to-end thiomethyl–thiomethyl contact mode, while the high conductance peak corresponded to attachment formed between the thiomethyl group at one gold contact and the central silicon at the other gold contact.

Theoretical calculations

Using the density functional code SIESTA⁴ the optimum geometries of the isolated molecules were obtained by relaxing the molecules until all forces on the atoms were less than 0.05 eV / Å. A double-zeta plus polarization orbital basis set^{20,21}, norm-conserving pseudopotentials, an energy cut-off of 250 Rydbergs defined the real space grid. Transmission functions were computed using and the local density approximation (LDA),²² a generalized gradient

approximations (GGA),²³ and using a van der Waals²⁴ functional and were found to be qualitatively similar in all cases,²⁵ (see Figure S45 in the SI).

The use of DFT to compute the ground state energy of various molecular junctions permits us to calculate binding energies to gold electrodes and optimal geometries. To avoid basis set superposition errors (BSSE)²⁶ we use the counterpoise correction²⁷ in which the binding energy of a molecule denoted *a* to an electrode *b* is expressed as:

$$\Delta E(ab) = E_{ab}^{ab} - E_a^{ab} - E_b^{ab} \quad (1)$$

In this expression, the total energy of the combined *a* and *b* system is E_{ab}^{ab} , while the total energies of isolated systems *a* and *b* are E_a^{ab} and E_b^{ab} respectively, maintaining identical basis sets (a,b) for the three energies.

Table 2. The binding energies $\Delta E(ab)$ of molecules to the Au(111) surface, calculated using a vdW functional. For comparison the corresponding binding energies obtained using GGA are shown. To obtain these results, we started from 252 different initial conditions for each molecule and allowed each to fully relax to a minimum energy. Then from these 252 different simulations, we chose the relaxed structure corresponding to the lowest energy. Figures S41-S44 in the SI show the various local energy minima obtained from this procedure.

The DFT calculations possible geometries involving two (compound 2). Based	System	$\Delta E(ab)$ (eV)GGA	$\Delta E(ab)$ (eV) vdW	predicted three for the system thiomethyl AGs on the binding energy
	2-A	- 0.57	-0.61	
	2-B	- 0.76	-0.79	
	2-C	- 0.82	-0.85	
	3	- 0.94	-1.02	

calculations shown in Table 2, the most probable geometry was shown to be 2-C (Figure 2),

since the order of binding energies between the molecule and gold electrode follows the trend $|E_{2-C}| > |E_{2-B}| > |E_{2-A}|$. For the structures with three thiomethyl anchor groups, our model shows one possible geometry for compounds **3**, **3=S** and **3=O** (Figure 3). The highest binding energy value is presented by compound **3** ($\Delta E = -1.02$ eV).




Figure 2. The relaxed geometries of all possible structures for **2**, **3**, **3=S** and **3=O**; where atoms are indicated by grey (carbon), white (hydrogen), yellow (sulfur), red (oxygen) and orange (Phosphorus).

To further explain the conductance behaviour we employed DFT-based transport code GOLLUM²⁸ to compute the transmission coefficient $T(E)$ for each relaxed junction geometry

(see Figure 3). This was achieved by first obtaining the corresponding Hamiltonian and overlap matrices with SIESTA, using a double-zeta polarized basis set. The optimized junction geometries as shown in Figure 3 confirm well that the thiomethyl-contacted compounds are not oriented normal to the electrode surface within the molecular junction. Rather, they are tilted within molecular junctions to accommodate the directionality of the lone pairs of electrons on the sulfur atoms that bind to the gold electrodes.^{3,4} From $T(E)$, the electrical conductance G was obtained using the Landauer formula.




Figure 3. The relaxed geometries of all molecular junctions; where atoms are indicated by grey (carbon), white (hydrogen), yellow (sulfur), red (oxygen) and orange (Phosphorus).

Figure 4 indicates that in all cases the Fermi level lies near the centre of the HOMO–LUMO gap, but shifted slightly towards the LUMO resonance, and therefore a LUMO-mediated electron tunnelling mechanism is anticipated. These results are consistent with our previous studies.^{1,2} The order of the calculated conductances is $G_3 > G_{3-S} > G_{3-O}$.




Figure 4. The calculated conductance as a function of Fermi energy for all molecular junctions, with molecules bound to Au (111) surfaces, obtained using a van der Waals functional^{20, 21}.

Table 3. The experimental (Exp. G/G_0) and calculated conductance values (Th. G/G_0) at $E_F - E_F^{DFT} = 0$ eV. Z^* is the experimental break-off distance. The calculated electrode separation in relaxed junctions (Z); $Z = d_{Au-Au} - 0.25$ nm, where 0.25 nm is the calculated centre-to-centre distance of the apex atoms of the two opposing gold electrodes when conductance = G_0 in the absence of a molecule. d_{Au-Au} is the calculated centre-to-centre distance of the apex atoms of the two opposing gold electrodes in relaxed junctions.

Molecule	Exp. G/G_0	Th. G/G_0	Z^* (nm)	Z (nm)	$d_{\text{Au-Au}}$ (nm)
3	4.36×10^{-3}	5.5×10^{-3}	0.76	0.27	0.52
3-S	1.23×10^{-3}	2.8×10^{-3}	0.87	0.39	0.64
3-O	1.09×10^{-3}	2.5×10^{-3}	0.76	0.26	0.51

Conclusions

In conclusion we have investigated the molecular conductance of a ‘bare-bones’ tripodal structure, consisting of either tris(4-(methylthio)phenyl)phosphine (**3**) or phosphine sulfide (**3=S**), and with both the thiomethyl and phosphorus/phosphine sulfide acting as surface binding groups. Using STM-BJ, we determined that both **3** and **3=S** are highly conductive, with only a single conductance peak.

Through the systematic substitution of thiomethyl groups with hydrogens, it was determined that three thiomethyl groups were required to achieve the single conductance mode, inferring that molecules bind to the surface via the thiomethyl groups, leaving the phosphorus (**3**) or phosphine sulfide (**3=S**) available to act as a ‘top’ contact for the molecular bridge. This was supported by DFT calculations, which showed that this geometry was the only stable geometry for the **3**, **3=S** and **3=O**. The calculations also indicate the Fermi-energy level lies in the vicinity of the middle of the HOMO-LUMO gap slightly closer to the LUMO.

Author Information

Corresponding Authors

*E-mail: ross.davidson@durham.ac.uk, andrew.beeby@durham.ac.uk and nichols@liv.ac.uk

Notes

The authors declare no competing financial interest.

Acknowledgements

AB and RJD also gratefully acknowledge the EPSRC (EP/K007785/1; EP/K007548/1) for funding this work. DCM, RJN, CJL and SJH also gratefully acknowledge the EPSRC (EP/M029522/1; EP/M005046/1; EP/M014169/1; EP/N017188/1; EP/M014452/1) for funding this work. This work support from the European Commission is provided by the FET Open project 767187 – QuIET and the Iraqi Ministry of Higher Education (SL-20). A.K.I. acknowledges financial support from Tikrit University (Iraq). This work is also supported by the EC FP7 ITN “MOLESCO” project no. 606728.



References

1. W. Hong, D. Z. Manrique, P. Moreno-García, M. Gulcur, A. Mishchenko, C. J. Lambert, M. R. Bryce and T. Wandlowski, *J. Am. Chem. Soc.*, 2012, **134**, 2292-2304.
2. V. Kaliginedi, A. V. Rudnev, P. Moreno-Garcia, M. Baghernejad, C. Huang, W. Hong and T. Wandlowski, *Phys. Chem. Chem. Phys.*, 2014, **16**, 23529-23539.
3. M. A. Karimi, S. G. Bahoosh, M. Valasek, M. Burkle, M. Mayor, F. Pauly and E. Scheer, *Nanoscale*, 2016, **8**, 10582-10590.
4. M. Valášek, M. Lindner and M. Mayor, *Beilstein J. Nanotechnol.*, 2016, **7**, 374-405.
5. Y. Ie, T. Hirose, H. Nakamura, M. Kiguchi, N. Takagi, M. Kawai and Y. Aso, *J. Am. Chem. Soc.*, 2011, **133**, 3014-3022.
6. L. Gerhard, K. Edelmann, J. Homberg, M. Valášek, S. G. Bahoosh, M. Lukas, F. Pauly, M. Mayor and W. Wulfhekel, *Nat. Commun.*, 2017, **8**, 14672.

7. M. Valášek and M. Mayor, *Chem. Eur. J.*, 2017, **23**, 1.
8. Y. Ie, K. Tanaka, A. Tashiro, S. K. Lee, H. R. Testai, R. Yamada, H. Tada and Y. Aso, *J. Phys. Chem. Lett.*, 2015, **6**, 3754-3759.
9. M. Lindner, M. Valášek, J. Homberg, K. Edelmann, L. Gerhard, W. Wulfhekel, O. Fuhr, T. Wächter, M. Zharnikov, V. Kolivoška, L. Pospíšil, G. Mészáros, M. Hromadová and M. Mayor, *Chem. Eur. J.*, 2016, **22**, 13218-13235.
10. F. Ragaini, L. Lunardi, D. Tomasoni and V. Guglielmi, *J. Organomet. Chem.*, 2004, **689**, 3621-3630.
11. R. Parameswaran, J. R. Widawsky, H. Vázquez, Y. S. Park, B. M. Boardman, C. Nuckolls, M. L. Steigerwald, M. S. Hybertsen and L. Venkataraman, *J. Phys. Chem. Lett.*, 2010, **1**, 2114-2119.
12. A. Fukazawa, M. Kiguchi, S. Tange, Y. Ichihashi, Q. Zhao, T. Takahashi, T. Konishi, K. Murakoshi, Y. Tsuji, A. Staykov, K. Yoshizawa and S. Yamaguchi, *Chem. Lett.*, 2011, **40**, 174-176.
13. R. J. Brooke, C. Jin, D. S. Szumski, R. J. Nichols, B.-W. Mao, K. S. Thygesen and W. Schwarzacher, *Nano Lett.*, 2015, **15**, 275-280.
14. W. Haiss, D. Lackey, J. K. Sass and K. H. Besocke, *J. Chem. Phys.*, 1991, **95**, 2193-2196.
15. Y. S. Park, A. C. Whalley, M. Kamenetska, M. L. Steigerwald, M. S. Hybertsen, C. Nuckolls and L. Venkataraman, *J. Am. Chem. Soc.*, 2007, **129**, 15768-15769.
16. O. Fuhr, A. Meredith and D. Fenske, *Dalton Trans.*, 2002, **22**, 4091-4094.
17. B. Xu and N. J. Tao, *Science*, 2003, **301**, 1221-1223.
18. Y. S. Park, J. R. Widawsky, M. Kamenetska, M. L. Steigerwald, M. S. Hybertsen, C. Nuckolls and L. Venkataraman, *J. Am. Chem. Soc.*, 2009, **131**, 10820-10821.

19. T. A. Su, J. R. Widawsky, H. Li, R. S. Klausen, J. L. Leighton, M. L. Steigerwald, L. Venkataraman and C. Nuckolls, *J. Am. Chem. Soc.*, 2013, **135**, 18331-18334.
20. W. Haiss, H. van Zalinge, S. J. Higgins, D. Bethell, H. Höbenreich, D. J. Schiffrin and R. J. Nichols, *J. Am. Chem. Soc.*, 2003, **125**, 15294-15295.
21. J. M. Soler, E. Artacho, J. D. Gale, A. Garcia, J. Junquera, P. Ordejon and D. Sanchez-Portal, *J. Phys. Condes. Matter*, 2002, **14**, 2745-2779.
22. W. Kohn and L. J. Sham, *Phys. Rev.*, 1965, **140**, A1133-A1138.
23. J. P. Perdew, K. Burke and M. Ernzerhof, *Phys. Rev. Lett.*, 1996, **77**, 3865-3868.
24. M. Dion, H. Rydberg, E. Schröder, D. C. Langreth and B. I. Lundqvist, *Phys. Rev. Lett.*, 2004, **92**, 246401.
25. A. K. Ismael, K. Wang, A. Vezzoli, M. K. Al-Khaykanee, H. E. Gallagher, I. M. Grace, C. J. Lambert, B. Xu, R. J. Nichols and S. J. Higgins, *Angew. Chem. Int. Ed.*, 2017, **56**, 15378-15382.
26. H. B. Jansen and P. Ros, *Chem. Phys. Lett.*, 1969, **3**, 140-143.
27. S. F. Boys and F. Bernardi, *Mol. Phys.*, 1970, **19**, 553-566.
28. J. Ferrer, C. J. Lambert, V. M. García-Suárez, D. Z. Manrique, D. Visontai, L. Oroszlany, R. Rodríguez-Ferradás, I. Grace, S. W. D. Bailey, K. Gillemot, S. Hatef and L. A. Algharagholy, *New J. Phys.*, 2014, **16**, 093029.

TABLE OF CONTENTS/ABSTRACTS GRAPHIC



When venturing into the field of tripodal single molecule conductors the geometry of the tripod with respect to the surface is a critical factor that determines their conductance. Here the authors examine the conductance behaviour of a simple tripodal conductor by systematic modifications of a triarylphosphine system.