

KHOVANOV HOMOTOPY CALCULATIONS USING FLOW CATEGORY CALCULUS

ANDREW LOBB, PATRICK ORSON, AND DIRK SCHÜTZ

ABSTRACT. The Lipshitz-Sarkar stable homotopy link invariant defines Steenrod squares on the Khovanov cohomology of a link. Lipshitz-Sarkar constructed an algorithm for computing the first two Steenrod squares. We develop a new algorithm which implements the flow category simplification techniques previously defined by the authors and Dan Jones. We give a purely combinatorial approach to calculating the second Steenrod square and Bockstein homomorphisms in Khovanov cohomology, and flow categories in general.

The new method has been implemented in a computer program by the third author and applied to large classes of knots and links. Several homotopy types not previously witnessed are observed, and more evidence is obtained that Khovanov stable homotopy types do not contain $\mathbb{C}P^2$ as a wedge summand. In fact, we are led by our calculations to formulate an even stronger conjecture in terms of $\mathbb{Z}/2$ summands of the cohomology.

1. INTRODUCTION

Framed flow categories were introduced by Cohen-Jones-Segal in [CJS95] as a way potentially to refine Floer homological invariants to space-level invariants. They were used by Lipshitz-Sarkar in [LS14a] to produce a stable homotopy type link invariant $\mathcal{X}_{Kh}(L)$ for links $L \subset S^3$. The cohomology of the spectrum $\mathcal{X}_{Kh}(L)$ is the Khovanov cohomology of the link L and so the spectrum can be used to define Steenrod square operations on the Khovanov cohomology of the link. The calculation of Steenrod squares becomes important in this context as a way to distinguish interesting stable homotopy types from those which are the suspension spectrum of a wedge of Moore spaces. Moreover, it was shown in [LS14b] that Steenrod squares in Khovanov cohomology can be used to refine the Rasmussen s -invariant of the smooth concordance class of a link, providing another important reason to be interested in their computation.

The main purpose of this paper is to describe a combinatorial algorithm for calculating Steenrod squares in the cohomology of the suspension spectrum associated to a general framed flow category based on the flow category simplifications developed in [JLS17, LOS18]. In particular this is a new method computing the Steenrod squares in Khovanov cohomology.

This combinatorial algorithm has been turned into a computer program by the third author. We present some of the more significant computations, including several stable homotopy type wedge summands not previously witnessed, and more evidence for the possibility that $\mathbb{C}P^2$ never appears as a wedge summand of the

The authors were partially supported by the EPSRC Grant EP/K00591X/1. PO was supported by a CIRGET postdoctoral fellowship.

Khovanov stable homotopy type of any link (see [LS14c, Question 5.2]). Guided by the calculations, we extend this idea to a conjecture about 2-torsion in Chang space wedge summands (see Conjecture 2.1).

1.1. 1-flow categories and the second Steenrod square. In [LS14c], Lipshitz-Sarkar already describe an algorithm for computing Steenrod squares from a framed flow category, but it is valid only in the specific context of flow categories arising directly from their construction in [LS14a] applied to a link $L \subset S^3$. Our algorithm is quite different from theirs, it works for any framed flow category, and indeed in the more general setting of what we call a *framed 1-flow category*. Any framed flow category restricts to one of these framed 1-flow categories, but a framed 1-flow category is a simpler object. In some sense it is nothing but a cochain complex with the minimal amount of extra data added so that one can define a combinatorial second Steenrod square on the cohomology (see Subsection 3.1 for precise details of this statement).

Put in terms of knot theory and the Lipshitz-Sarkar construction [LS14a], the Khovanov cochain complex of a link can only ‘see’ the information of the vertices and edges of the Khovanov cube of the link. The complete Lipshitz-Sarkar framed flow category can see all subcubes of the Khovanov cube of the link, and how they interrelate. The 1-flow category of a link is a collection of data which can see the information of how 2-dimensional faces of the Khovanov cube relate edges and vertices. However, this amount of information is sufficient if all you want is to compute the second Steenrod square.

1.2. Combinatorial arguments. While a link in S^3 is a geometric object, the input for Khovanov cohomology and for the Khovanov spectrum is the Khovanov cube, which can be viewed as a combinatorial object. We believe that there is some benefit to giving proofs and constructions purely combinatorially, particularly when a combinatorial input is involved. It is also possible that the combinatorics are amenable to generalizations and extensions in ways that non-combinatorial arguments may not be.

In the particular cases of the constructions considered in this paper (which arise originally from framings of embedded arcs and circles), it is a considerable effort to turn the topological data of a framed flow category into combinatorial data sufficient to make computations of the second Steenrod square. Working combinatorially from the start, this effort is sidestepped, leading to more elementary proofs.

1.3. Plan of the paper. We begin the paper by presenting in Section 2 some calculations made by the program implementing the algorithm due to the third author [Sch17]. We discuss what these calculations suggest for the conjectural picture of stable homotopy types realized by the Lipshitz-Sarkar construction.

For knots and links with more than 16 crossings the associated flow categories returned by the Lipshitz-Sarkar construction become very large, and we found that the new algorithm, in particular when combined with the results of [JLS15] on special diagrams, greatly speeds up the Steenrod square computations in these cases.

Section 3 is dedicated to a combinatorial construction and proof of the following theorem, stated here in abbreviated form:

Theorem 3.13. *Let (\mathcal{C}, s, f) be a framed 1-flow category. Then there is a well-defined linear map $\text{Sq}^2: H^*(\mathcal{C}; \mathbb{Z}/2) \rightarrow H^{*+2}(\mathcal{C}; \mathbb{Z}/2)$.*

What we mean by ‘well-defined’ in this context we postpone until later to make precise. Of course, the map defined in this theorem should (and does) agree with the second Steenrod square when computed in the context of the Khovanov homotopy type and its relatives, as we discuss at the end of Section 3.

In Sections 4 and 5 we turn to the algorithm. The algorithm makes heavy use of the *flow category moves* in [JLS17] and [LOS18], or rather their 1-flow category restrictions. The moves – called *handle slides*, *handle cancellation*, and *Whitney trick* – for modifying a framed flow category, are based on ideas from Morse theory. The moves do not change the stable homotopy type associated to the flow category, but we show in [LOS18, §6] that they can be used to simplify the flow category itself in an algorithmic way. To take advantage of this idea for our eventual Steenrod square algorithm, we describe explicitly in Section 4 how our flow category moves restrict to framed 1-flow categories.

In Section 5 we give a new combinatorial way to calculate the second Steenrod square. Subsection 5.2 gives a refinement of the auxiliary data used to compute the second Steenrod square, and Subsection 5.3 shows that this refinement suffices. Subsection 5.4 gives a treatment of the flow category moves in the contexts necessary for the algorithm, while Subsection 5.5 gives a complete description of the algorithm. The algorithm works by first simplifying the 1-flow category using the flow category moves, then applying the new combinatorial Steenrod square process to obtain the result.

2. CALCULATIONS

A computer program implementing the described algorithm has been developed by the third author, see [Sch17]. Both an integral and a mod 4 version have been written. The mod 4 version is noticeably faster when applied to diagrams with a larger number of crossings.

For more complicated diagrams such as for the torus knot $T_{7,4}$ the new program is faster than a previously written program of the third author. The new program is capable of doing calculations on the diagram for the torus knot $T_{6,5}$ given in Figure 1, although for different quantum degrees the time consumption varies greatly. As the algorithm still deals with a global approach to the diagram, the program has large memory requirements.

Note that the algorithm starts with objects of top degree in the normalization process. This can lead to a quite different performance depending on whether a knot or its mirror is considered! One may wonder whether a more balanced algorithm may lead to better results.

We denote by $\mathcal{X}_{Kh}^q(L)$ the stable homotopy type of the link L in quantum degree q , and we write $\tilde{\mathcal{X}}_{Kh}^q(L)$ for the reduced stable homotopy type, compare [LS14a, §8].

2.1. The torus link $T_{5,5}$. We used the diagram of Figure 1 to determine the Lipshitz-Sarkar stable homotopy types of $T_{5,5}$ for various quantum degrees. Figure 2 lists the Khovanov cohomology of $T_{5,5}$ which potentially contain non-trivial second




FIGURE 1. Glued diagrams for $T_{5,5}$ (left) and $T_{6,5}$ (right). For the definition of *glued diagram* see [JLS15].

Steenrod squares. For $q = 21$ we get the Chang¹ space $X(2\eta)$ which is already present in the stable homotopy type of $T_{3,3}$ and predicted by the stability result of Willis [Wil17]. Similarly, for $q = 23$ we get no non-trivial Steenrod squares, in line with the result we get for $T_{5,4}$ for $q = 19$.

$h \backslash q$	21	23	25	27	29	31	33
12						\mathbb{Z}	\mathbb{Z}^5
11						\mathbb{Z}_2	$\mathbb{Z} \oplus \mathbb{Z}_2^4$
10					\mathbb{Z}	$\mathbb{Z}^4 \oplus \mathbb{Z}_2$	\mathbb{Z}_2
9					\mathbb{Z}_4	\mathbb{Z}	
8			\mathbb{Z}	\mathbb{Z}^6	\mathbb{Z}^4		
7			\mathbb{Z}_2	$\mathbb{Z} \oplus \mathbb{Z}_2$	\mathbb{Z}		
6		\mathbb{Z}	\mathbb{Z}	0			
5		0	\mathbb{Z}	\mathbb{Z}			
4	\mathbb{Z}	\mathbb{Z}					
3	\mathbb{Z}_2	\mathbb{Z}					

FIGURE 2. The Khovanov cohomology of $T_{5,5}$ in quantum degrees from $q = 21$ to 33.

Both of these calculations are done rather fast. For $q = 25$ the calculation is still fairly efficient, but one can get the result even quicker by considering the reduced stable homotopy type.

The reduced Khovanov cohomology of $T_{5,5}$ is given in Figure 3, and calculations show that $\tilde{\mathcal{X}}_{Kh}^{26}(T_{5,5})$ is the central Baues-Hennes space $X(\eta 2\xi)$, while $\tilde{\mathcal{X}}_{Kh}^{32}(T_{5,5})$ is the wedge of a Chang space $X(2\eta)$ with Moore spaces. The central Baues-Hennes space $X(\eta 2\xi)$ can be realized as an appropriate suspension of $\mathbb{R}\mathbf{P}^5/\mathbb{R}\mathbf{P}^1$.

There is a cofibration sequence $\tilde{\mathcal{X}}_{Kh}^{q-1}(T_{5,5}) \hookrightarrow \mathcal{X}_{Kh}^q(T_{5,5}) \twoheadrightarrow \tilde{\mathcal{X}}_{Kh}^{q+1}(T_{5,5})$ by [LS14a, Thm.8.1]. From the resulting long exact sequence in cohomology and the naturality

¹See [Bau95, §11] for the definition of Chang space, and also for the definition of the various Baues-Hennes spaces.

$h \backslash q$	24	26	28	30	32	34
12					\mathbb{Z}	\mathbb{Z}^4
11					\mathbb{Z}_2	\mathbb{Z}^4
10				\mathbb{Z}	$\mathbb{Z}^3 \oplus \mathbb{Z}_2$	
9				\mathbb{Z}		
8		\mathbb{Z}	\mathbb{Z}^5			
7		\mathbb{Z}_2	\mathbb{Z}			
6	\mathbb{Z}	0				
5		\mathbb{Z}				

FIGURE 3. The reduced Khovanov cohomology of $T_{5,5}$ in quantum degrees from $q = 24$ to 34.

of Steenrod squares we get that both $\mathcal{X}_{Kh}^{25}(T_{5,5})$ and $\mathcal{X}_{Kh}^{27}(T_{5,5})$ contain $X(\eta 2\xi)$ and the rest are Moore spaces.

For $q \geq 29$ the calculations become very time and memory consuming, and we have not finished them. Particularly $q = 29$ would be interesting, as there is no $\mathbb{Z}/2$ summand in the Khovanov cohomology. One may expect a trivial second Steenrod square from degree 7 to 9 because of naturality with $T_{5,4}$, but a non-trivial second Steenrod square from degree 8 to 10 would lead to a previously unobserved Chang space.

2.2. The torus knot $T_{6,5}$. Similarly we can do calculations for $T_{6,5}$ using the right diagram of Figure 1. For $q = 29$ we also get a central Baues-Hennes space $X(\eta 2\xi)$, which should come from the $T_{5,5}$ result, although it does not fall into the stable range described by Willis [Wil17]. Again this Baues-Hennes space appears in the reduced stable homotopy type for $q = 30$, which means it also appears in the unreduced case for $q = 31$.

2.3. Knots with non-trivial Sq^3 . The disjoint union of three trefoils is known to contain the cyclic Baues-Hennes space $X(\xi^2 \eta_2, \text{id})$, see [LLS15] and [LOS18], giving the simplest example of a link with non-trivial Sq^3 . Recall that $Sq^1 Sq^2 = Sq^3$ by an Adem relation. Going through the prime knot tables of [Thi99], we found exactly one knot with at most 13 crossings that admits a non-trivial Sq^3 , namely $13n3663$ in quantum degree $q = 1$. The resulting Baues-Hennes space is again $X(\xi^2 \eta_2, \text{id})$. Figure 4 lists the prime knots with 14 crossings that exhibit this Baues-Hennes space as a summand. No other Baues-Hennes spaces have been found for prime knots with up to 14 crossings, although there could be Baues-Hennes spaces corresponding to ε -words among them.

2.4. Knots with Chang spaces involving 4-torsion. There are not that many prime knots with at most 16 crossings that have 4-torsion in their Khovanov cohomology. We have tested these knots for whether their stable homotopy type contains a Chang space involving 4-torsion. Those which do are listed in Figure 5. The knots have been mirrored so that the 4-torsion appears in a positive quantum degree. Note that all the occurring Chang spaces also contain 2-torsion, and the Chang space appears in quantum degree 13 or 25 (with one exception, where $q = 11$). The table only lists the summand in the stable homotopy type which

Knot	q	Knot	q
14n10164	-3	14n21627	-9
14n10510	-7	14n22180	3
14n18918	-7	14n22185	-1
14n18935	15	14n22589	-1
14n19177	-1	14n23524	-3
14n19265	-3	14n26039	-1
14n19315	9	14n26580	9

FIGURE 4. Prime knots with 14 crossings containing a cyclic Baues-Hennes space $X(\xi^2\eta_2, \text{id})$.

Knot	q	Chang space	Knot	q	Chang space	Knot	q	Chang space
15n46927	25	$X_{(2\eta_4)}$	16n300202	13	$X_{(4\eta_2)}$	16n785487	13	$X_{(2\eta_4)}$
15n49024	25	$X_{(2\eta_4)}$	16n300488	13	$X_{(2\eta_4)}$	16n785689	13	$X_{(2\eta_4)}$
15n53220	25	$X_{(2\eta_4)}$	16n306918	13	$X_{(2\eta_4)}$	16n787476	11	$X_{(2\eta_4)}$
15n131377	13	$X_{(2\eta_4)}$	16n324645	13	$X_{(2\eta_4)}$	16n795419	13	$X_{(2\eta_4)}$
15n136587	13	$X_{(2\eta_4)}$	16n327544	13	$X_{(4\eta_2)}$	16n796824	13	$X_{(2\eta_4)}$
15n136594	13	$X_{(2\eta_4)}$	16n332191	13	$X_{(2\eta_4)}$	16n796923	13	$X_{(2\eta_4)}$
15n137384	13	$X_{(2\eta_4)}$	16n334330	25	$X_{(2\eta_4)}$	16n807682	13	$X_{(2\eta_4)}$
15n137388	13	$X_{(2\eta_4)}$	16n336536	13	$X_{(2\eta_4)}$	16n814264	25	$X_{(2\eta_4)}$
15n141041	13	$X_{(2\eta_4)}$	16n365743	13	$X_{(4\eta_2)}$	16n830240	25	$X_{(2\eta_4)}$
15n166170	25	$X_{(2\eta_4)}$	16n696534	13	$X_{(2\eta_4)}$	16n831623	25	$X_{(2\eta_4)}$
16n224701	13	$X_{(4\eta_2)}$	16n696547	13	$X_{(2\eta_4)}$	16n836058	25	$X_{(2\eta_4)}$
16n224826	25	$X_{(2\eta_4)}$	16n697548	13	$X_{(2\eta_4)}$	16n838182	25	$X_{(2\eta_4)}$
16n228740	25	$X_{(2\eta_4)}$	16n711906	13	$X_{(2\eta_4)}$	16n839613	25	$X_{(2\eta_4)}$
16n240992	13	$X_{(4\eta_2)}$	16n763485	25	$X_{(2\eta_4)}$	16n881950	13	$X_{(4\eta_2)}$
16n265388	13	$X_{(4\eta_2)}$	16n781146	13	$X_{(2\eta_4)}$	16n992204	13	$X_{(4\eta_2)}$
16n281362	13	$X_{(4\eta_2)}$	16n784195	25	$X_{(2\eta_4)}$	16n992205	13	$X_{(4\eta_2)}$
16n287907	13	$X_{(4\eta_2)}$	16n784488	25	$X_{(2\eta_4)}$	16n994256	13	$X_{(4\eta_2)}$
16n289670	25	$X_{(2\eta_4)}$	16n784622	13	$X_{(2\eta_4)}$			

FIGURE 5. Prime knots with up to 16 crossings containing a Chang space involving 4-torsion.

contains the 4-torsion. None of the knots with 4-torsion in their Khovanov cohomology has more than one $\mathbb{Z}/4$ summand in their cohomology. On the basis of these calculations, we make the following tentative conjecture.

Conjecture 2.1. *If a Chang space occurs as a wedge summand of the Khovanov stable homotopy type of a link, then that Chang space has a $\mathbb{Z}/2$ summand in its cohomology.*

Note that, as $\mathbb{C}P^2$ has the stable homotopy type of the Chang space $X(\eta)$, this conjecture includes the statement that $\mathbb{C}P^2$ does not appear as a wedge summand of the Khovanov stable homotopy type of a link.

3. COMBINATORIAL STEENROD SQUARES

This section is in large part dedicated to the statement and combinatorial proof of Theorem 3.13. This theorem will not be surprising, given its topological motivations. The price paid for keeping the arguments in this section combinatorial is a somewhat lengthy although elementary proof.

We start in Subsection 3.1 giving the definition of our principal object of interest: a 1-flow category. Subsection 3.2 formalizes the auxiliary information needed to make a computation of the second Steenrod square, and does it in such a way as to be useful to us in giving the algorithm later. Finally Subsection 3.3 gives the proof of Theorem 3.13. We conclude with a few remarks concerning topological arguments.

3.1. 1-flow categories. In the combinatorial algorithm of [LS14c] for calculating the Steenrod squares on Khovanov cohomology, it was already observed that only a very restricted amount of the data of the flow category is actually required for the computation. In this paper we formalise this restricted amount of information in the following object.

Definition 3.1. A 1-flow category \mathcal{C} consists of a finite set $\text{Ob}(\mathcal{C})$, a function $|\cdot|: \text{Ob}(\mathcal{C}) \rightarrow \mathbb{Z}$ called the *grading*, and for each pair $a, b \in \text{Ob}(\mathcal{C})$ with $|a| - |b| = 1$ or 2 a *moduli space* $\mathcal{M}(a, b)$ which is a compact manifold of dimension $|a| - |b| - 1$, satisfying the following

- **Boundary Condition:** If $a, c \in \text{Ob}(\mathcal{C})$ with $|a| - |c| = 2$, then the boundary of $\mathcal{M}(a, b)$ is given by

$$\partial\mathcal{M}(a, c) = \coprod_{b \in \text{Ob}(\mathcal{C}), |b|=|a|-1} \mathcal{M}(b, c) \times \mathcal{M}(a, b)$$

If $|a| - |d| = 3$ we do not need a moduli space $\mathcal{M}(a, d)$. However, we define the boundary of $\mathcal{M}(a, d)$ as

$$\begin{aligned} \partial\mathcal{M}(a, d) = & \coprod_{b \in \text{Ob}(\mathcal{C}), |b|=|a|-1} \mathcal{M}(b, d) \times \mathcal{M}(a, b) \\ & \cup \coprod_{c \in \text{Ob}(\mathcal{C}), |c|=|a|-2} \mathcal{M}(c, d) \times \mathcal{M}(a, c) \end{aligned}$$

Notice that the two disjoint unions have a common subset, which is

$$\coprod_{\substack{(b, c) \in \text{Ob}(\mathcal{C}) \times \text{Ob}(\mathcal{C}) \\ |b|=|c|+1=|d|+2}} \mathcal{M}(c, d) \times \mathcal{M}(b, c) \times \mathcal{M}(a, b)$$

It is easy to see that $\partial\mathcal{M}(a, d)$ is a disjoint union of components which are homeomorphic to circles.

We caution that a 1-flow category is not a flow category. But one can usually think of a 1-flow category as a flow category where the higher dimensional moduli spaces are ignored (although not always, as we show in Example 3.19).

Definition 3.2. Let \mathcal{C} be a 1-flow category. A *sign assignment* s for \mathcal{C} is an assignment $s(P) \in \{0, 1\}$ for every point P in a 0-dimensional moduli space $\mathcal{M}(a, b)$ with the property that if $(P_1, Q_1) \in \mathcal{M}(b_1, c) \times \mathcal{M}(a, b_1)$ and $(P_2, Q_2) \in \mathcal{M}(b_2, c) \times \mathcal{M}(a, b_2)$ are the boundary of an interval component in $\mathcal{M}(a, c)$, then

$$s(P_1) + s(Q_1) + s(P_2) + s(Q_2) = 1.$$

For objects $a, b \in \text{Ob}(\mathcal{C})$ with $|a| = |b| + 1$ we define $[a : b] \in \mathbb{Z}$ as

$$[a : b] = \sum_{P \in \mathcal{M}(a, b)} (-1)^{s(P)}.$$

A *pre-framing* f of \mathcal{C} is an assignment $f(C) \in \{0, 1\}$ for every component $C \subset \mathcal{M}(a, c)$ of a 1-dimensional moduli space.

Remark 3.3. For the reader familiar with framed flow categories it is clear that the sign assignment leads to a framing of the 0-dimensional moduli spaces. We will therefore refer to points P with $s(P) = 0$ as *positively* framed, while points with $s(P) = 1$ are referred to as *negatively* framed.

The chain complex $C_*(\mathcal{C})$ is defined to have modules freely generated over \mathbb{Z} by the objects of \mathcal{C} , which are homologically graded by the grading function of \mathcal{C} . A sign assignment allows us to define a differential on this graded module by setting

$$\partial(a) = \sum [a : b]b$$

for $a \in \text{Ob}(\mathcal{C})$, where the sum is taken over all objects b with $|b| = |a| - 1$.

The purpose of a pre-framing is to give us enough information to define a cohomology operation combinatorially on the resulting cohomology with $\mathbb{Z}/2$ -coefficients. However, an arbitrary choice of pre-framing will not be sufficient for this purpose, and there must be a certain consistency in our choices as we now discuss.

Note that the connected components C of a 1-dimensional moduli space are either intervals or circles. A circle embedded in \mathbb{R}^n with $n \geq 4$ can be framed in two ways, one is trivial in the sense that it extends to a framing of the disc, and there is a non-trivial framing. We shall also refer to the trivial framing as the *standard framing*. Similarly, if an interval is embedded in $[0, \infty) \times \mathbb{R}^n$, with boundary embedded in $\{0\} \times \mathbb{R}^n$, and satisfying a transversality condition near the boundary, then there are two ways of framing it (provided a framing on the boundary is fixed and that this boundary framing is different at the two endpoints). Calling one of these two framings ‘standard’ or ‘non-standard’ requires consistent choices. Such choices were made in [LS14c, §3]. We will not repeat that somewhat technical discussion here, as we shall only need the combinatorial consequences.

When $f(C) = 0$ we will call this a trivial framing for C , and when $f(C) = 1$ we will call this a non-trivial framing. In order for this language to be meaningful, we need to add a consistency condition in terms of an induced framing on $\partial\mathcal{M}(a, d)$.

Let C be a component in $\partial\mathcal{M}(a, d)$. If C is a circle, which is either of the form $\{P\} \times S^1 \subset \mathcal{M}(c, d) \times \mathcal{M}(a, c)$ or $S^1 \times \{Q\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$, we define

$$\tilde{f}(C) = 1 + f(S^1) \in \mathbb{Z}/2,$$

where $f(S^1)$ is the framing value of the circle. If C is a union of intervals $\{P\} \times J \subset \mathcal{M}(c, d) \times \mathcal{M}(a, c)$ and $I \times \{Q\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$, we define

$$\tilde{f}(C) = \sum_{I \times \{Q\}} f(I) + \sum_{\{P\} \times J} (1 + s(P) + f(J)) \in \mathbb{Z}/2$$

where the first sum is over all intervals of the form $I \times \{Q\} \subset C$ and the second sum over all intervals of the form $\{P\} \times J$.

Definition 3.4. Let \mathcal{C} be a 1-flow category, s a sign assignment and f a pre-framing of \mathcal{C} . Then (\mathcal{C}, s, f) is called a *framed 1-flow category*, if the following *compatibility condition* is satisfied for f .

- Let $a, d \in \text{Ob}(\mathcal{C})$ satisfy $|a| = |d| + 3$. Then

$$\sum_C (1 + \tilde{f}(C)) = 0 \in \mathbb{Z}/2$$

where the sum is taken over all components in $\partial\mathcal{M}(a, d)$.

If f satisfies the condition with respect to s , we call f a *framing* of \mathcal{C} .

Remark 3.5. A framed flow category in the sense of [LS14a] determines a framed 1-flow category. The framing values for each interval component can be derived using the description in [LS14c], see also [JLS15]. We omit the slightly cumbersome derivation of this, but the reader is invited to check that the specific framing values given in [LS14c] and [JLS15] do indeed satisfy the compatibility condition.

3.2. Special graph structures.

Definition 3.6. Let V be a finite set. An *edge* is a subset $e \subset V$ with two elements. A *directed edge* is an edge e together with a function $d: \{e\} \rightarrow e$.

We can think of a directed edge e pointing towards $d(e)$, or write it as an ordered pair $e = (v, w)$ with $w = d(e)$.

Definition 3.7. A *special graph structure* $\Gamma = \Gamma(V, E, E', E'', L, s, f)$ consists of a set of vertices V , a function $s: V \rightarrow \{0, 1\}$, a set of edges E , a subset $E' \subset E$ together with a function $f: E' \rightarrow \{0, 1\}$, a subset $E'' \subset E - E'$ of directed edges, and a set of loops L . All sets are finite. It has to satisfy the following conditions:

- (1) Each vertex is contained in at least one edge, and in at most two edges. We denote the set of vertices contained in only one edge by ∂V and call these the boundary points.
- (2) If $v \in \partial V$, then the unique edge $e(v)$ with $v \in e(v)$ satisfies $e(v) \in E'$.
- (3) If $v \in V - \partial V$, then there is an edge $e_1 \in E'$ with $v \in e_1$, and an edge $e_2 \in E - E'$ with $v \in e_2$.
- (4) If $e \in E'$ and $e = \{v_1, v_2\}$, then $s(v_1) \neq s(v_2)$.
- (5) If $e \in E - E'$ and $e = \{v_1, v_2\}$, then $s(v_1) = s(v_2)$ if and only if $e \in E''$.

The edges E determine an equivalence relation on the vertex set V in the obvious way. By condition (1) the edges in an equivalence class either form an interval or a circle. The set of equivalence classes under this relation, taken together with the set L of loops, forms the set of *components* of Γ . Notice that L is a set independent from possible circles made from edges in E .

Example 3.8. Let (\mathcal{C}, s, f) be a framed 1-flow category and $a, c \in \text{Ob}(\mathcal{C})$ with $|a| - |c| = 2$. Define a special graph structure $\Gamma(a, c)$ as follows. The vertex set V consists of all points $(B, A) \in \mathcal{M}(b, c) \times \mathcal{M}(a, b)$ for some $b \in \text{Ob}(\mathcal{C})$ with $s(B, A) = s(B) + s(A)$. The edges $E' = E$ are determined by the intervals in $\mathcal{M}(a, c)$, with $f: E' \rightarrow \{0, 1\}$ coming from the framing of the 1-flow category. The set of loops L in $\Gamma(a, c)$ is given by the set of non-trivially framed circles in $\mathcal{M}(a, c)$.

Definition 3.9. Let $\Gamma = \Gamma(V, E, E', E'', L, s, f)$ be a special graph structure. For a circle component C of Γ which contains a vertex, let $F(C)$ be the sum of the framing values $f(e')$ where $e' \in E'$ is in C . Also, let $D(C) \in \mathbb{Z}/2$ be the number of oriented edges in C which point in a chosen given direction.

Lemma 3.10. *In the definition of $D(C)$, the value does not depend on the choice of given direction.*

Proof. Let C be a component and v a vertex in C . We need to show that the number of directed edges in C is even. First note that the total number of edges in C is even by Definition 3.7 (3). Let F'' be the directed edges in C and let F be the non-directed edges in C . Note that non-directed edges change the sign of their endpoints by Definition 3.7 (4) and (5), while directed edges do not, by Definition 3.7 (5). Since C is a circle, we have to have an even number of edges in F . It follows that the number of directed edges is also even. \square

Let (\mathcal{C}, s, f) be a framed 1-flow category. A cochain $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ can be identified with a function $\varphi: \text{Ob}_k(\mathcal{C}) \rightarrow \mathbb{Z}/2$, where $\text{Ob}_k(\mathcal{C}) \subset \text{Ob}(\mathcal{C})$ consists of those objects $c \in \text{Ob}(\mathcal{C})$ with $|c| = k$. For each cochain φ there exists a unique set $\{c_1, \dots, c_l\} \subset \text{Ob}(\mathcal{C})$ with $\varphi(c_i) = 1$ for all $i = 1, \dots, l$, and $\varphi(c) = 0$ if $c \neq c_i$ for all $i = 1, \dots, l$. We say that φ is *represented* by c_1, \dots, c_l . If φ is represented by exactly one c , we will simply write $c = \varphi$ by an abuse of notation.

Now suppose $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ is a cocycle represented by $c_1, \dots, c_l \in \text{Ob}(\mathcal{C})$. Let $b_1, \dots, b_m \in \text{Ob}(\mathcal{C})$ be all objects that have a non-empty 0-dimensional moduli space $\mathcal{M}(b_i, c_j)$ for some $j \in \{1, \dots, l\}$.

If we fix b_i , the union

$$\mathcal{M}(b_i, \varphi) := \bigcup_{j=1}^l \mathcal{M}(b_i, c_j)$$

is a set of even cardinality because φ is a cocycle with $\mathbb{Z}/2$ -coefficients.

Definition 3.11. Let (\mathcal{C}, s, f) be a framed 1-flow category and $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ be a cocycle. Writing b_1, \dots, b_m as above we may choose a partition of the elements of $\mathcal{M}(b_i, \varphi)$ into ordered pairs. If we make this choice for each $i = 1, \dots, m$, the overall choice is called a *combinatorial boundary matching* \mathcal{C} for φ .

Now suppose we have made a choice of combinatorial boundary matching \mathcal{C} for a cocycle $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ and let $a \in \text{Ob}(\mathcal{C})$ satisfy $|a| = k + 2$. Define a special graph structure $\Gamma_{\mathcal{C}}(a, \varphi)$ as follows. We begin with the disjoint union of the special graph structures $\Gamma(a, c_j)$, defined as in Example 3.8. Note that if $\mathcal{M}(a, c_i) = \emptyset$, we can also use the empty set for $\Gamma(a, c_j)$. Let (B_1, B_2) be a pair in \mathcal{C} . We then add an edge between (B_1, A) and (B_2, A) for all $A \in \mathcal{M}(a, b_i)$ with $B_k \in \mathcal{M}(b_i, c_{j_k})$. If $s(B_1) = s(B_2)$, this edge is directed towards B_2 , otherwise it is undirected. This defines $\Gamma_{\mathcal{C}}(a, \varphi)$.

Notice that by construction all vertices of $\Gamma_{\mathcal{C}}(a, \varphi)$ have valency two, so that the set of components C of $\Gamma_{\mathcal{C}}(a, \varphi)$ consists of a disjoint union of circles together with the loop set L .

Definition 3.12. Let (\mathcal{C}, s, f) be a framed 1-flow category and $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ a cocycle. Then define a cochain $\text{sq}^{\varphi}: C_{k+2}(\mathcal{C}; \mathbb{Z}/2) \rightarrow \mathbb{Z}/2$ by

$$\text{sq}^{\varphi}(a) := |L| + \sum_C (1 + F(C) + D(C)) \in \mathbb{Z}/2,$$

where L is the loop set of $\Gamma_{\mathcal{C}}(a, \varphi)$ and the sum is taken over all components C of $\Gamma_{\mathcal{C}}(a, \varphi)$ containing a vertex.

The following theorem states that we now have enough combinatorial information to define a Steenrod square for 1-flow categories.

Theorem 3.13. *Let (\mathcal{C}, s, f) be a framed 1-flow category and $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ a cocycle. Then sq^φ is a cycle and the cohomology class of sq^φ does not depend on the choice of combinatorial boundary matching for φ . Furthermore, there is a well defined linear map*

$$\text{Sq}^2: H^k(\mathcal{C}; \mathbb{Z}/2) \rightarrow H^{k+2}(\mathcal{C}; \mathbb{Z}/2); \quad z \mapsto [\text{sq}^\varphi],$$

where φ is any choice of cocycle representing the cohomology class z .

We will prove the theorem at the end of Subsection 3.3, once we have developed the necessary combinatorial machinery to do so.

Remark 3.14. The definition of sq is of course motivated by the formula given in [LS14c], and if the framed 1-flow category is induced from a framed flow category, the resulting linear map agrees with the second Steenrod square operation, compare also [JLS15, §5.1].


Example 3.15. Consider the 1-flow category \mathcal{C} consisting of four objects a, b_1, b_2, c and 0-dimensional moduli spaces

$$\begin{aligned} \mathcal{M}(a, b_1) &= \{X_1, X_2\}, & \mathcal{M}(a, b_2) &= \{Y_1, Y_2\} \\ \mathcal{M}(b_1, c) &= \{Z_1, Z_2\}, & \mathcal{M}(b_2, c) &= \{W_1, W_2\}. \end{aligned}$$

Framing Y_1 and Y_2 negatively, that is, $s(Y_1) = s(Y_2) = 1$, while all other points are framed positively, ensures that we can extend \mathcal{C} to a framed 1-flow category with four intervals between the eight endpoints (Z_i, X_j) and (W_k, Y_l) for $i, j, k, l = 1, 2$. The compatibility condition is trivially satisfied, so the framing values for these intervals can be arbitrary. Let us simply choose the standard framing for all of them.

Two possible choices are given by having intervals between (Z_i, X_j) and (W_i, Y_j) , or between (Z_i, X_j) and (W_j, Y_i) .

If we use the combinatorial matching (Z_1, Z_2) and (W_1, W_2) (note that our only choice is the order for both pairs), the resulting circles occur as follows.



The left side is what we get for connecting the points (Z_i, X_j) with (W_i, Y_j) , and the right side when we connect (Z_i, X_j) with (W_j, Y_i) . On the left we get for both circles $D(C) = 1$, so $\text{sq}^c(a) = 0$ which results in the trivial Steenrod square. On the right there is only one circle, with $D(C) = 0$, so $\text{sq}^c(a) = 1$, resulting in a non-trivial Steenrod square (provided there are no non-trivially framed circles in $\mathcal{M}(a, c)$). We remark that the second possibility arises for the disjoint union of two trefoil knots, compare [JLS17].


3.3. Combinatorial proof of Theorem 3.13.


Definition 3.16. For Γ a special graph structure with $\partial V = \emptyset$, define

$$o(\Gamma) := |L| + \sum_C (1 + F(C) + D(C)),$$

where the sum is taken over all components C of Γ which contain a vertex.
(With this notation Definition 3.12 can be written as $\text{sq}^\varphi(a) := o(\Gamma_C(a, \varphi))$.)

We now introduce a relation on special graph structures which is not going to change the value $o(\Gamma)$. The relation is described in terms of local moves on edges, listed below as (a) – (i). In the diagrams, the symbol ε stands for an element of $\mathbb{Z}/2$ and is the sign of a vertex. The numerical value above an edge (if any) indicates its framing value.





Note that the edges in (a), (b) and (d) are from $E - E'$, while the edges in (c) are from E' . The circles in (a), (c), (d) and (e) represent loops in L . Here (e) means that two such loops can be removed.

The equivalence relation \sim on special graph structures is the transitive closure of these local moves (a) – (i). Notice that the equivalence relation preserves the boundary points of a special graph structure.

Lemma 3.17. *Let Γ be a special graph structure with $\partial V = \emptyset$. Then $o(\Gamma)$ is invariant under \sim .*

Proof. We need to check this for all local moves (a) – (i). For (c) – (h) this is obvious. The remaining cases are similar, and we will only show (i).

Let v, w, x and y be vertices with $s(v) = s(w) = s(y) = s(x) + 1$ and $e_1, e_2 \in E''$ edges with $e_1 = (v, w)$ the directed edge and $e_2 = (x, y)$ the non-directed edge as in move (i). As $\partial V = \emptyset$, both edges are part of circle components C_1 and C_2 with $e_i \subset C_i$ for $i = 1, 2$. We can assume that $f(e) = 0$ for all $e \in E'$.

Assume that C_1 and C_2 are different components. Let $a \in \mathbb{Z}/2$ be the number of directed edges in C_1 that point in a different direction than e_1 . Also, let $b \in \mathbb{Z}/2$ be the number of directed edges in C_2 pointing in a chosen direction. The contribution of C_1 and C_2 to $o(\Gamma)$ is $a + b$.

After performing the move (i) we have a new special graph structure Γ' where the components C_1 and C_2 are replaced by one component C , and we have edges $e'_1 = (v, x)$ and $e'_2 = (w, y)$. Note that e'_2 is the directed edge, and there are a directed edges between v and w pointing in the same direction as e'_2 , and there are b directed edges between y and x pointing in the same direction as e'_2 . The contribution of C to $o(\Gamma')$ is therefore $1 + a + b + 1 = a + b$, as there are $a + b + 1$ directed edges pointing in the same direction as e'_2 in C (this includes e'_2). As all other components in Γ' are unchanged from Γ , we get $o(\Gamma) = o(\Gamma')$.

Now assume that $C_1 = C_2$. If we follow on from w in the direction of e_1 , we either reach x or y first. If we reach x first, then performing the (i) move leads to a graph Γ' where both edges $e'_1 = (v, x)$ and $e'_2 = (w, y)$ are again in the same component. Let $a \in \mathbb{Z}/2$ be the number of directed edges on the path between w and x which point in the same direction as e_1 , and let $b \in \mathbb{Z}/2$ be the number of directed edges on the path between y and v which point in the same direction as e_1 . The contribution of C_1 to $o(\Gamma)$ is then given by $1 + a + b + 1 = a + b$. Let C be the new component in Γ' containing e'_1 and e'_2 . The contribution of C to Γ' is also $a + b$, as there are $1 + b$ directed edges between w and v , starting with e'_2 , and there are a directed edges on the path between x and w pointing in the same direction

as e'_2 . Note that since $s(x) \neq s(w)$ there are indeed the same number of directed edges in a chosen direction between x and w , which in this case is a . Hence we get again $o(\Gamma) = o(\Gamma')$.

If we reach y first, then performing the (i) move leads to a graph Γ' where the edges $e'_1 = (v, x)$ and $e'_2 = (w, y)$ are now in different components. This case is now basically just the reverse of the first case where we changed from two circles to one. The same argument thus shows $o(\Gamma) = o(\Gamma')$. \square

Proof of Theorem 3.13. Let the cocycle $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ be represented by objects $c_1, \dots, c_m \in \text{Ob}(\mathcal{C})$ satisfying $|c_i| = k$ for all $i = 1, \dots, m$. We begin by showing that sq^φ is a cocycle. Let $u \in \text{Ob}(\mathcal{C})$ satisfy $|u| = k + 3$. Then

$$\partial\mathcal{M}(u, c_i) = \coprod_a \mathcal{M}(a, c_i) \times \mathcal{M}(u, a) \sqcup \coprod_b \mathcal{M}(b, c_i) \times \mathcal{M}(u, b)$$

where the first union is over all $a \in \text{Ob}(\mathcal{C})$ with $|a| = k + 2$ and the second union over all $b \in \text{Ob}(\mathcal{C})$ with $|b| = k + 1$. We associate a special graph structure $\Gamma(u, c_i)$ to this as follows. The vertices are given by

$$\coprod_{a,b} \mathcal{M}(b, c_i) \times \mathcal{M}(a, b) \times \mathcal{M}(u, a)$$

with $s(C, B, A) = s(C) + s(B) + s(A)$. As edges in E' we use $I \times \{A\} \subset \mathcal{M}(a, c_i) \times \mathcal{M}(u, a)$, where $I \subset \mathcal{M}(a, c_i)$ is an interval component. We set $f(I \times \{A\}) = f(I)$. The edges in $E - E'$ are given by the $\{C\} \times J \subset \mathcal{M}(b, c_i) \times \mathcal{M}(u, b)$ with J an interval component in $\mathcal{M}(u, b)$. Furthermore, if such an interval $\{C\} \times J$ satisfies $s(C) + f(J) = 0$, we add a loop in L . Finally, any non-trivially framed circle $\{C\} \times S^1 \subset \mathcal{M}(b, c_i) \times \mathcal{M}(u, b)$ and $S^1 \times \{A\} \subset \mathcal{M}(a, c_i) \times \mathcal{M}(u, a)$ gives rise to a loop in L . Note that $E'' = \emptyset$.

With this definition, we get $o(\Gamma(u, c_i)) = 0$, because loops and framings were chosen so that we get exactly the sum from the compatibility condition for $\partial\mathcal{M}(u, c_i)$.

Also form the special graph structure $\Gamma(u, \varphi)$ as the disjoint union of the $\Gamma(u, c_i)$. We want to replace edges coming from $\{C\} \times J$ with edges coming from the combinatorial boundary matching \mathcal{C} . Note that $C \in \mathcal{M}(b, c_i)$ is matched to some $C' \in \mathcal{M}(b, c_j)$, and we therefore also have an edge coming from $\{C'\} \times J$. We can then use a move (a) to replace these two edges with edges coming from the boundary matching. If $s(C) = s(C')$, we have $s(C) + f(J) = 0$ if and only if $s(C') + f(J) = 0$ so possible extra loops coming from this condition cancel each other. If $s(C) \neq s(C')$ this condition creates exactly one extra loop, which cancels the loop coming from move (a).

Replacing all the loops $\{C\} \times J$ this way leads to the special graph structure

$$\coprod_a \Gamma'_\mathcal{C}(a, \varphi) \times \mathcal{M}(u, a)$$

where $\Gamma'_C(a, \varphi) \times \{A\}$ only differs from $\Gamma_C(a, \varphi)$ in that the sign functions for vertices differ by $s(A)$. We thus get that $o(\Gamma'_C(a, \varphi) \times \{A\}) = o(\Gamma_C(a, \varphi))$. Furthermore,

$$\begin{aligned} 0 &= o\left(\prod_a \Gamma'_C(a, \varphi) \times \mathcal{M}(u, a)\right) \\ &= \sum_a o(\Gamma_C(a, \varphi)) \cdot [u : a] \\ &= \delta(\text{sq}^\varphi)(u). \end{aligned}$$

Therefore sq^φ is a cocycle.


To see that the cohomology class of sq^φ does not depend on the combinatorial boundary matching, first consider the order of a directed edge (C, C') with $C \in \mathcal{M}(b, c_i)$ and $C' \in \mathcal{M}(b, c_j)$ for some $b \in \text{Ob}(\mathcal{C})$ with $|b| = k + 1$. Then $o(\Gamma_{C'}(a, \varphi))$ differs from $o(\Gamma_C(a, \varphi))$ in that every occurrence of $(C, B) \in \mathcal{M}(b, c_j) \times \mathcal{M}(a, b)$ leads to a $+1$. In other words, $o(\Gamma_{C'}(a, \varphi)) + o(\Gamma_C(a, \varphi)) = [a : b] \in \mathbb{Z}/2$. This is true for every $a \in \text{Ob}(\mathcal{C})$ with $|a| = k + 2$, so this difference is compensated by adding $\delta(b)$ to sq^φ .

If we are given matchings (C_1, C_2) and (C_3, C_4) , we can replace this by matchings (C_1, C_3) and (C_2, C_4) which in the special graph structures correspond to moves (a) or (b) depending on signs of the points involved, again for all combinations with $A \in \mathcal{M}(a, b)$. So any possible circles arising from move (a) would again be compensated by $\delta(b)$. As we can go between any boundary matching using these steps, we see that the cohomology class does not depend on it.

If we are given two cocycles φ_1 and φ_2 we now show $\text{sq}^{\varphi_1 + \varphi_2} = \text{sq}^{\varphi_1} + \text{sq}^{\varphi_2}$. Taking respective combinatorial matchings \mathcal{C}_1 and \mathcal{C}_2 for φ_1 and φ_2 , we will have in general that some of the objects c_i are used for representing both φ_1 and φ_2 , so are not needed for $\varphi_1 + \varphi_2$. We can begin by letting \mathcal{C} be the disjoint union of \mathcal{C}_1 and \mathcal{C}_2 , and setting

$$\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2) = \Gamma_{\mathcal{C}_1}(a, \varphi_1) \sqcup \Gamma_{\mathcal{C}_2}(a, \varphi_2).$$

Let c_i be representing both φ_1 and φ_2 and assume that $I \subset \mathcal{M}(a, c_i)$ is an interval with endpoints (C, B) and (C', B') . Let C be matched to C_l by \mathcal{C}_l and C' be matched to C'_l by \mathcal{C}_l for $l = 1, 2$. In $\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2)$ the interval I occurs twice, and we can use the moves (a), (b) and (i) to change $\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2)$ without changing $o(\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2))$.



The square in the middle involving both intervals I contributes 0 to $o(\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2))$ and can be removed. Note that the directed edges may be in different places, but the square needs to have directed edges. We can now remove C and C' from \mathcal{C} and match C_1 with C_2 and C'_1 with C'_2 . After finitely many steps we get the right special

graph structure for $\text{sq}^{\varphi_1+\varphi_2}$, and as the value $o(\Gamma_{\mathcal{C}}(a, \varphi_1 + \varphi_2))$ never changed, we have $\text{sq}^{\varphi_1+\varphi_2} = \text{sq}^{\varphi_1} + \text{sq}^{\varphi_2}$.

It remains to show that sq^{φ} is a coboundary for φ a coboundary. So let $d \in \text{Ob}(\mathcal{C})$ satisfy $|d| = k-1$ and let c_1, \dots, c_m be the objects satisfying $|c_i| = k$ and $\mathcal{M}(c_i, d) \neq \emptyset$. Let us first assume that for each i , $\mathcal{M}(c_i, d)$ contains exactly one point.

If $a \in \text{Ob}(\mathcal{C})$ satisfies $|a| = k+2$, consider

$$\partial\mathcal{M}(a, d) = \coprod_b \mathcal{M}(b, d) \times \mathcal{M}(a, b) \sqcup \coprod_{c'} \mathcal{M}(c', d) \times \mathcal{M}(a, c')$$

where the first disjoint union is over all objects b with $|b| = k+1$ and the second over objects c' with $|c'| = k$. Note that we can restrict the second disjoint union to the objects c_1, \dots, c_m . Define a special graph structure $\Gamma(a, d)$ as follows. Vertices are given by


$$\coprod_{b,i} \mathcal{M}(c_i, d) \times \mathcal{M}(b, c_i) \times \mathcal{M}(a, b)$$

and we set $s(D, C, B) = s(D) + s(C) + s(B)$. As edges in E' we use $\{D\} \times I \subset \mathcal{M}(c_i, d) \times \mathcal{M}(a, c_i)$, where I is an interval component. We also set $f(\{D\} \times I) = s(D) + f(I)$. The edges in $E - E'$ are given by $J \times \{B\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$, where J is an interval component. For every such interval $J \times \{B\}$ we add a loop to L provided that $f(J) = 0$. Finally, any non-trivially framed circle $\{D\} \times S^1 \subset \mathcal{M}(c_i, d) \times \mathcal{M}(a, c_i)$ and $S^1 \times \{B\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$ gives rise to a loop in L . The set of directed edges $E'' = \emptyset$.

Note that $\Gamma(a, d)$ is very similar to the construction of $\Gamma(u, c_i)$ at the beginning of the proof, but the roles of E' and $E - E'$ are reversed. Nevertheless, we get again that $o(\Gamma(a, d)) = 0$ from the compatibility condition of $\partial\mathcal{M}(a, d)$.

Intervals $J \subset \mathcal{M}(b, d)$ have endpoints $(D_i, C) \in \mathcal{M}(c_i, d) \times \mathcal{M}(b, c_i)$ and $(D_j, C') \in \mathcal{M}(c_j, d) \times \mathcal{M}(b, c_j)$ for some i, j . As we assume that each $\mathcal{M}(c_i, d)$ has at most one point, we can use these intervals to define the combinatorial boundary matching \mathcal{C} . Namely, we match C and C' . If $s(C) = s(C')$, then $s(D_i) \neq s(D_j)$ and we choose the ordering (C, C') if and only if $s(D_i) = 0$.

Consider an edge $\{D_i\} \times I \subset \mathcal{M}(c_i, d) \times \mathcal{M}(a, c_i)$ in $\Gamma(a, d)$. There is a corresponding edge I in $\mathcal{M}_{\mathcal{C}}(a, \delta(d))$, but if $s(D_i) = 1$, the vertices have different sign, and the contribution to $o(\Gamma(a, d))$ and $o(\Gamma_{\mathcal{C}}(a, \delta(d)))$ differ by 1. However, we can perform a move of type (a) in $\Gamma(a, d)$ as follows:



The right graph has I as in $\Gamma_{\mathcal{C}}(a, \delta(d))$. Note that J and J' have as endpoints (D_j, C') and (D_k, C'') at the bottom. If $s(D_j) = 0 = s(D_k)$, then the blue directed edges are exactly as the boundary matching prescribes, and the contribution of $s(D_i) = 1$ corresponds to the 1 coming from the directed edge pointing with an orientation. If $s(D_k) = 1$, we have to do another such move to get the edge unoriented again. Repeating this throughout will show that the contributions of

$s(D_i)$ to $o(\Gamma(a, d))$ is the same as the contribution of the edges coming from the boundary matching via directed edges to $o(\Gamma_{\mathcal{C}}(a, \delta(d)))$.

There are still some contributions to $o(\Gamma(a, d))$ coming from loops which are not present in $o(\Gamma_{\mathcal{C}}(a, \delta(d)))$, namely coming from edges $J \times \{B\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$ with $f(J) = 0$ and non-trivially framed circles $S^1 \times \{B\} \subset \mathcal{M}(b, d) \times \mathcal{M}(a, b)$. But for each B we only need to check how many contributions from $\mathcal{M}(b, d)$ there are and compensate with $\delta(b)$. This shows that $\text{sq}^{\delta(d)}$ differs from 0 by a coboundary.

Finally, assume that a $\mathcal{M}(c_i, d)$ contains two different points X_1, X_2 . We can form $\Gamma(a, d)$ as before. Any interval $I \subset \mathcal{M}(a, c_i)$ leads to two edges $\{X_1\} \times I$ and $\{X_2\} \times I$ in $\Gamma(a, d)$ contributing $s(X_1) + s(X_2)$ to $o(\Gamma(a, d))$. We can replace these two edges by edges between their endpoints using a move (a) without changing $o(\Gamma(a, d))$. We can do this with any pairs of points in $\mathcal{M}(c_i, d)$ until there is only one point (or none) in each $\mathcal{M}(c_i, d)$ not paired this way. We can then use these remaining points to obtain a combinatorial boundary matching as in the previous case. The same argument as before shows that $\text{sq}^{\delta(d)}$ differs from 0 by a coboundary. \square

It is a great effort to turn the topological data of a framed flow category into the combinatorial data of a 1-flow category, and then to verify the formula for the second Steenrod square, given a combinatorial boundary matching. Nevertheless, it can be done and was carried in [LS14c] for the Lipshitz-Sarkar framed flow category and also in [JLS15] in general. Treating this (considerable) work as a black box, one is able to make the following remark.

Remark 3.18. Theorem 3.13 can alternatively be proved using the topological Steenrod square as follows. First we can truncate the framed 1-flow category \mathcal{C} so that all objects are in degree $k - 1$ to $k + 3$. This does not affect the cohomology groups $H^k(\mathcal{C}; \mathbb{Z}/2)$ and $H^{k+2}(\mathcal{C}; \mathbb{Z}/2)$. We now want to describe a framed flow category \mathcal{C}' which extends \mathcal{C} and so that the cohomology in degree k and $k + 2$ remains the same

We use the same objects for \mathcal{C}' as for \mathcal{C} , and it is straightforward to frame the 0- and 1-dimensional moduli spaces to extend \mathcal{C} . To get a 2-dimensional moduli space $\mathcal{M}(a, d)$ or $\mathcal{M}(b, e)$ where $|a| = k + 3 = |b| + 1 = |d| + 3 = |e| + 4$ note that compositions determine the boundary of these moduli spaces as a closed framed 1-manifold. The compatibility condition ensures that this framing is trivial, so we can choose any framed null-cobordism as the 2-dimensional moduli space.

For all pairs of objects a and e with $|a| = k + 3$ and $|e| = k - 1$, the possible compositions between a and e piece together a framed surface $S(a, e)$ which would be the boundary of the 3-dimensional moduli space $\mathcal{M}(a, e)$.

Recall that $\Omega_2^{fr} \cong \mathbb{Z}/2$, generated by the product of two non-trivially framed circles. If the framing of $S(a, e)$ is trivial we can extend $S(a, e)$ to a 3-dimensional framed moduli space $\mathcal{M}(a, e)$, and we are done.

If the framing is non-trivial, we add an extra object $c_{a,e}$ of grading $k + 1$ together with non-trivially framed circles as the moduli spaces $\mathcal{M}(a, c_{a,e})$ and $\mathcal{M}(c_{a,e}, e)$. This then adds a non-trivially framed component to $S(a, e)$, hence $S(a, e)$ becomes trivially framed overall. We are then able to choose a 3-dimensional framed moduli space $\mathcal{M}(a, e)$ with $\partial\mathcal{M}(a, e) = S(a, e)$, and the resulting framed flow category \mathcal{C}' has a well defined second Steenrod square from [JLS15]. Furthermore, the formula agrees exactly with the formula for \mathcal{C} .

As a neat application of Theorem 3.13, we finish this section with an example showing that not every framed 1-flow category extends to a framed flow category.

Example 3.19. Let \mathcal{C} have four objects a, b, d, e with $k + 3 = |a| = |b| + 1 = |d| + 3 = |e| + 4$ and let $\mathcal{M}(a, b)$ and $\mathcal{M}(d, e)$ both consist of two points with positive sign. Furthermore, let $\mathcal{M}(b, d)$ contain a non-trivially framed circle. This satisfies the compatibility condition and thus defines a framed 1-flow category. But notice that

$$\mathrm{Sq}^1 \mathrm{Sq}^2 \mathrm{Sq}^1(e) = a \neq 0$$

while $\mathrm{Sq}^2(e) = 0$. Therefore \mathcal{C} cannot be turned into a framed flow category, as the resulting space would violate the Adem relation $\mathrm{Sq}^1 \mathrm{Sq}^2 \mathrm{Sq}^1 = \mathrm{Sq}^2 \mathrm{Sq}^2$. Note that adding an object c with $|c| = k + 2$ as in Remark 3.18 establishes exactly this Adem relation.

4. MORSE MOVES IN 1-FLOW CATEGORIES

In [JLS17] and [LOS18] the authors, together with Dan Jones, defined a series of *flow category moves* – called *handle slides*, *handle cancellation*, and *Whitney trick* – for modifying and algorithmically simplifying a framed flow category. Section 4 is dedicated to an explicit description of the restrictions of these moves to 1-flow categories, with a mind to their application in Section 5 as part of the new algorithm for computing Steenrod squares.

4.1. Combinatorial handle slides. The handle slides of [LOS18] have obvious analogues in framed 1-flow categories.

Definition 4.1. Let (\mathcal{C}, s, f) be a framed 1-flow category, and let $x, y \in \mathrm{Ob}(\mathcal{C})$ satisfy $|x| = |y|$. For $\varepsilon \in \{0, 1\}$ let $(\mathcal{C}_s^\varepsilon, s', f')$ be the following framed 1-flow category:

- The objects of $\mathcal{C}_s^\varepsilon$ are in one-to-one correspondence with the objects of \mathcal{C} , and for $a \in \mathrm{Ob}(\mathcal{C})$ we write $a' \in \mathrm{Ob}(\mathcal{C}_s^\varepsilon)$ for the corresponding object. Furthermore, $|a'| = |a|$.
- The moduli spaces are given by

$$\begin{aligned} \mathcal{M}(x', b') &= \mathcal{M}(x, b) \sqcup \mathcal{M}(y, b) \times \{1\} \\ \mathcal{M}(a', y') &= \mathcal{M}(a, y) \sqcup \{0\} \times \mathcal{M}(a, x) \end{aligned}$$

If $|a| = |x| + 1 = |c| + 2$ we also have

$$\mathcal{M}(a', c') = \mathcal{M}(a, c) \sqcup \mathcal{M}(y, c) \times [0, 1] \times \mathcal{M}(a, x).$$

- The sign assignment s' is defined by

$$s'(X) = \begin{cases} s(X) & \text{if } X \in \mathcal{M}(a, b) \subset \mathcal{M}(a', b') \\ \varepsilon + s(X) & \text{if } X \in \mathcal{M}(y, b) \subset \mathcal{M}(x', b') \\ 1 + \varepsilon + s(X) & \text{if } X \in \mathcal{M}(a, x) \subset \mathcal{M}(a', y') \end{cases}$$

- The framing f' is defined by

$$f'(I) = \begin{cases} f(I) & \text{if } I \subset \mathcal{M}(a, b) \subset \mathcal{M}(a', b') \\ f(I) & \text{if } I \subset \mathcal{M}(y, b) \subset \mathcal{M}(x', b') \\ 1 + \varepsilon + f(I) & \text{if } I \subset \mathcal{M}(a, x) \subset \mathcal{M}(a', y') \\ 1 + \varepsilon + s(B) & \text{if } I = \{B\} \times [0, 1] \times \{A\} \subset \mathcal{M}(y, b) \times [0, 1] \times \mathcal{M}(a, x) \end{cases}$$

We also call the case $\varepsilon = 0$ a (+)-slide and the case $\varepsilon = 1$ a (-)-slide. We need to check that $(\mathcal{C}_S^\varepsilon, s', f')$ is indeed a framed 1-flow category.

Lemma 4.2. *Let (\mathcal{C}, s, f) be a framed 1-flow category, and let $x, y \in \text{Ob}(\mathcal{C})$ satisfying $|x| = |y|$. For $\varepsilon \in \{0, 1\}$ the collection $(\mathcal{C}_S^\varepsilon, s', f')$ is a framed 1-flow category.*

Proof. We begin by showing that the new moduli spaces behave well with respect to composition. First assume that $|x| = |b| + 2$ and consider

$$\begin{aligned} \partial\mathcal{M}(x', b') &= \partial\mathcal{M}(x, b) \sqcup \partial\mathcal{M}(y, b) \times \{1\} \\ &= \coprod_{a, |a|=|b|+1} \mathcal{M}(a, b) \times \mathcal{M}(x, a) \sqcup \coprod_{a, |a|=|b|+1} \mathcal{M}(a, b) \times \mathcal{M}(y, a) \times \{1\} \\ &= \coprod_{a, |a|=|b|+1} \mathcal{M}(a', b') \times \mathcal{M}(x', b') \end{aligned}$$

so we see that composition matches up with the topological boundary of the new moduli spaces. The case for $\mathcal{M}(a', y')$ where $|a| = |y| + 2$ is similar and will be omitted.

Now consider objects a, c with $|a| = |x| + 1 = |c| + 2$. Then

$$\begin{aligned} \partial\mathcal{M}(a', c') &= \partial\mathcal{M}(a, c) \sqcup \mathcal{M}(y, c) \times \{0, 1\} \times \mathcal{M}(a, x) \\ &= \coprod_{b \neq x, y} \mathcal{M}(b, c) \times \mathcal{M}(a, b) \sqcup \mathcal{M}(x, c) \times \mathcal{M}(a, x) \sqcup \mathcal{M}(y, c) \times \mathcal{M}(a, y) \\ &\quad \sqcup \mathcal{M}(y, c) \times \{0\} \times \mathcal{M}(a, x) \sqcup \mathcal{M}(y, c) \times \{1\} \times \mathcal{M}(a, x) \\ &= \coprod_{b \neq x, y} \mathcal{M}(b', c') \times \mathcal{M}(a', b') \\ &\quad \sqcup \mathcal{M}(x', c') \times \mathcal{M}(a', x') \sqcup \mathcal{M}(y', c') \times \mathcal{M}(a', y') \\ &= \coprod_{b, |b|=|c|+1} \mathcal{M}(b', c') \times \mathcal{M}(a', b'). \end{aligned}$$

It is easy to see that s' is a sign assignment. We need to check the compatibility condition for $\partial\mathcal{M}(a', d')$ for any $a', d' \in \text{Ob}(\mathcal{C})$ with $|a'| = |d'| + 3$. There are essentially four cases depending on $|a'| - |x'|$.

If $|a'| = |x'|$ we have $\partial\mathcal{M}(a', d') = \partial\mathcal{M}(a, d)$ with the same framings, unless $a' = x'$. In that case

$$\partial\mathcal{M}(x', d') = \partial\mathcal{M}(x, d) \sqcup \partial\mathcal{M}(y, d)$$

and the framing values in the relevant circle components are the same as for f .

If $|a'| = |x'| + 1$, every interval $J \subset \mathcal{M}(y, d)$ and $A \in \mathcal{M}(a, x)$ leads to two intervals $J \subset \mathcal{M}(y', d')$ and $J' \subset \mathcal{M}(x', d')$ and two points $A \in \mathcal{M}(a', x')$ and $A' \in \mathcal{M}(a', y')$. Let the endpoints of J be given by $(C, B) \in \mathcal{M}(c, d) \times \mathcal{M}(y, c)$ and $(\tilde{C}, \tilde{B}) \in \mathcal{M}(\tilde{c}, d) \times \mathcal{M}(y, \tilde{c})$. We then also get the intervals $I_{B, A} = \{B\} \times [0, 1] \times \{A\} \subset \mathcal{M}(a', c')$ and $I_{\tilde{B}, A} \subset \mathcal{M}(a', \tilde{c}')$.

We then get a component $C_{J,A}$ in $\partial\mathcal{M}(a', d')$ from the four intervals $J' \times \{A\}$, $J \times \{A'\}$, $\{C\} \times I_{B,A}$, and $\{\tilde{C}\} \times I_{\tilde{B},A}$. Then

$$\begin{aligned} \tilde{f}(C_{J,A}) &= f'(J') + f'(J) + 1 + s'(C) + f'(I_{B,A}) + 1 + s'(\tilde{C}) + f'(I_{\tilde{B},A}) \\ &= f(J) + f(J) + s(C) + s(\tilde{C}) + 1 + \varepsilon + s(B) + 1 + \varepsilon + s(\tilde{B}) \\ &= 1. \end{aligned}$$

This component therefore contributes nothing to the compatibility condition. Similarly, if $A \in \mathcal{M}(a, x)$ and $S \subset \mathcal{M}(y, d)$ a circle, we get two extra circles $S \times \{A'\}$ and $S' \times \{A\}$ which cancel each other in the compatibility condition. The remaining components of $\partial\mathcal{M}(a', d')$ are present in $\partial\mathcal{M}(a, d)$ with the same contributions.

The case $|a'| = |x'| + 2$ is similar to the previous case, we now get new components $C_{C,I}$ made up of four intervals for every $C \in \mathcal{M}(y, d)$ and every interval $I \subset \mathcal{M}(a, x)$. A straightforward check shows $\tilde{f}(C_{C,I}) = 1$ as in the previous case.

For $|a'| = |x'| + 3$ one can give an argument similar to the first case. We omit the details. \square

Lemma 4.3. *Let (\mathcal{C}, s, f) be a framed 1-flow category and $(\mathcal{C}_S^\varepsilon, s', f')$ be the framed 1-flow category obtained by a handle slide. Then there is an isomorphism*

$$\Phi: H^*(\mathcal{C}; \mathbb{Z}/2) \rightarrow H^*(\mathcal{C}_S^\varepsilon; \mathbb{Z}/2)$$

which commutes with Sq^2 .

Proof. From the definition we see that

$$[a' : y'] = [a : y] + (-1)^{\varepsilon+1}[a : x]$$

and

$$[x' : c'] = [x : c] + (-1)^\varepsilon[y : c]$$

and all other incidence numbers remain the same. It is now easy to check that we get an isomorphism of chain complexes $\Phi: C_*(\mathcal{C}_S^\varepsilon) \rightarrow C_*(\mathcal{C})$ which differs from the identity only in that $\Phi(x') = x + (-1)^\varepsilon y$. The induced map on cochain complexes $\Phi: C^*(\mathcal{C}; \mathbb{Z}/2) \rightarrow C^*(\mathcal{C}_S^\varepsilon; \mathbb{Z}/2)$ satisfies $\Phi(y) = x' + y'$.

Now let $\varphi \in C^n(\mathcal{C}; \mathbb{Z}/2)$ be a cocycle with $n = |x| = |y|$, and let $\varphi' = \Phi(\varphi)$ the corresponding cocycle in $C^n(\mathcal{C}_S^\varepsilon; \mathbb{Z}/2)$. If y is not among the objects representing φ , it is clear that the Steenrod square cannot change. Therefore assume that y is among the objects representing φ . If x is among the objects representing φ , then x' is not among the objects representing φ' and vice versa. Given an object a with $|a| = n + 2$, the moduli space $\mathcal{M}(a, x)$ either contributes to sq^φ or $\text{sq}^{\varphi'}$, depending on whether x is among the objects representing φ or not.

Let $\varepsilon = 1$, in which case the new intervals in $\mathcal{M}(a', y')$ coming from $\mathcal{M}(a, x)$ keep their framing value. Let \mathcal{C} be a combinatorial boundary matching for φ and assume first that x is among the objects representing φ . The elements of $\mathcal{M}(b, x)$ for objects b with $|b| = n + 1$ are matched by \mathcal{C} , possibly with elements of $\mathcal{M}(b, c)$ where $c \neq x$. As elements of $\mathcal{M}(b', x')$ they need no longer be matched, since x' is not representing φ' . But as elements of $\mathcal{M}(b', y')$ they need to be matched by a new combinatorial boundary matching \mathcal{C}' for φ' . So we can keep the same boundary matching, just interpreting it differently for the new moduli spaces. Since $\varepsilon = 1$, the signs remain the same, so orientations of the edges remain the same, and we get identical values for the Steenrod square.

If x is not among the objects representing φ , points in $\mathcal{M}(x, b)$ are not matched by \mathcal{C} , but a new boundary matching \mathcal{C}' for φ' needs to match these points both for $\mathcal{M}(b', x')$ and $\mathcal{M}(b', y')$. So we define a new combinatorial boundary matching \mathcal{C}' by keeping \mathcal{C} as a subset, and adding matchings $(B_1, B_2) \in \mathcal{M}(b', x') \times \mathcal{M}(b', y')$, where B_i are the copies of a $B \in \mathcal{M}(b, x)$ in the respective new moduli spaces. Note that $s(B_1) = s(B_2)$, so we get a directed edge.

Every interval I of $\mathcal{M}(a, x)$ now gives rise to intervals $I_1 \subset \mathcal{M}(a', x')$ and $I_2 \subset \mathcal{M}(a', y')$, and it is easy to see that $\Gamma_{\mathcal{C}'}(a', \varphi')$ is obtained from $\Gamma_{\mathcal{C}}(a, \varphi)$ by adding a square C_I for every such interval I . Furthermore, $F(C_I) = 0$ and $D(C_I) = 1$, so these new components do not contribute to the Steenrod square.

If $\varepsilon = 0$, then intervals in $\mathcal{M}(a, x)$ have different framing when considered as intervals in $\mathcal{M}(a', y')$. Also, points in $\mathcal{M}(b, x)$ change their sign when considered in $\mathcal{M}(b', y')$. Assume first that x is among the objects representing φ . Consider a combinatorial boundary matching \mathcal{C} for φ . We can assume that whenever an element $B \in \mathcal{M}(b, x)$ is matched to an element $B' \in \mathcal{M}(b, c_i)$ with $c_i \neq x$ and $s(B') = s(B)$, the matching is ordered as (B, B') . For the combinatorial boundary matching \mathcal{C}' we then keep the matching, although we now consider $B \in \mathcal{M}(b', y')$. Note that the sign has changed so the matching is no longer considered ordered. If $s(B') \neq s(B)$, we get $s'(B') = s'(B)$, and we choose the ordering (B, B') in \mathcal{C}' .

Now intervals in $\mathcal{M}(a, x)$ change their framing when considered as intervals in $\mathcal{M}(a', y')$, which can affect the value of $\text{sq}^{\varphi'}(a')$. Observe that the components in $\Gamma_{\mathcal{C}}(a, \varphi)$ differ from the components in $\Gamma_{\mathcal{C}'}(a', \varphi')$ only in that the framing values for intervals coming from $\mathcal{M}(a, x)$ are different, and that there are changes for those edges in $E - E'$ which have exactly one vertex (B, A) with B originally from $\mathcal{M}(b, x)$.

Now let C be a component of $\Gamma_{\mathcal{C}}(a, \varphi)$. We define a *plateau* of C of length k to be a sequence I_1, \dots, I_k of intervals in $\mathcal{M}(a, x)$ such that for $i = 1, \dots, k$ we have $\partial I_i = \{(B_i, A_i), (\tilde{B}_{i+1}, A_{i+1})\}$, where $B_i \in \mathcal{M}(b_i, x)$ for $i = 1, \dots, k$, $\tilde{B}_i \in \mathcal{M}(b_i, x)$ for $i = 2, \dots, k+1$, and $A_i \in \mathcal{M}(a, b_i)$ for $i = 1, \dots, k+1$. Furthermore, B_i and \tilde{B}_i are matched by \mathcal{C} for $i = 2, \dots, k$, but B_1 is matched to some $\tilde{B}_1 \in \mathcal{M}(b_1, c)$ with $c \neq x$ and \tilde{B}_{k+1} is matched to some $B_{k+1} \in \mathcal{M}(b_{k+1}, \tilde{c})$ with $\tilde{c} \neq x$. We can visualize a plateau as a subgraph of $\Gamma_{\mathcal{C}}(a, \varphi)$ as in Figure 6. It is easy to see that a




FIGURE 6. A plateau consisting of three intervals from $\mathcal{M}(a, x)$.

plateau of length k contributes an extra k to $F(C)$ and an extra 1 to $D(C)$ when C is considered a component of $\Gamma_{\mathcal{C}'}(a', \varphi')$. Furthermore, every interval of $\mathcal{M}(a, x)$ is contained in a plateau for some component C of $\Gamma_{\mathcal{C}}(a, \varphi)$.

So for every plateau of even length we get one extra summand in the calculation of $\text{sq}^{\varphi'}(a)$. This is also true if we consider a plateau of length 0 to be a component C which has two edges, one of which is from $\mathcal{M}(a, x)$.

Now notice that a plateau of even length has an odd number of edges involving a matching B, \tilde{B} with $B, \tilde{B} \in \mathcal{M}(b, x)$ and an element $A \in \mathcal{M}(a, b)$ for some object b .

If the plateau has odd length, we get an even number of such edges, and for every matching $B, \tilde{B} \in \mathcal{M}(b, x)$ and point $A \in \mathcal{M}(a, b)$, some object b , we get an edge contained in a plateau.

For b with $|b| = n + 1$ define $k_b \in \mathbb{Z}/2$ the number of matchings $(B, \tilde{B}) \in \mathcal{M}(b, x)^2$, and define a cochain $\psi \in C^{n+1}(\mathcal{C}'; \mathbb{Z}/2)$ by $\psi(b') = k_b$. It is now straightforward to check that

$$\text{sq}^{\varphi'}(a') = \text{sq}^{\varphi}(a) + \delta\psi(a') \in \mathbb{Z}/2$$

for all objects a with $|a| = n + 2$. Hence $\text{sq}^{\varphi'}$ differs from $\Phi(\text{sq}^{\varphi})$ by a coboundary.

If x is not among the objects contributing to φ , the argument is much closer to the case where $\varepsilon = 1$. As in that case, we get new square components C_I in $\Gamma_{\mathcal{C}'}(a', \varphi')$ for every interval $I \subset \mathcal{M}(a, x)$, but this time $F(C_I) = 1$ and $D(C_I) = 0$, so again these components do not contribute to $\text{sq}^{\varphi'}(a')$.

The cases where $|x| = n + 1$ or $n + 2$ are easier and we omit them. \square

4.2. Combinatorial Whitney trick. In [JLS17] the Whitney trick in a flow category was described. We now give the analogue for a 1-flow category.

Definition 4.4. Let (\mathcal{C}, s, f) be a framed 1-flow category, and let $x, y \in \text{Ob}(\mathcal{C})$ satisfy $|x| = |y| + 1$. Assume that $\mathcal{M}(x, y)$ contains two points P, M with $s(P) = 0$ and $s(M) = 1$. Then let (\mathcal{C}', s', f') be the framed 1-flow category given as follows:

- We have $\text{Ob}(\mathcal{C}') = \text{Ob}(\mathcal{C})$, and we write $a' \in \text{Ob}(\mathcal{C}')$ for $a \in \text{Ob}(\mathcal{C})$ with $|a'| = |a|$.
- The 0-dimensional moduli spaces are unchanged except for $\mathcal{M}(x', y') = \mathcal{M}(x, y) \setminus \{P, M\}$, furthermore $s'(A) = s(A)$ for every A in a 0-dimensional moduli space.
- The 1-dimensional moduli spaces are unchanged except for $\mathcal{M}(a', y')$ and $\mathcal{M}(x', b')$, which are obtained from $\mathcal{M}(a, y)$ and $\mathcal{M}(x, b)$ respectively, by identifying endpoints of intervals of the form $(P, A) \sim (M, A)$ for $\mathcal{M}(a, y)$, and endpoints $(B, P) \sim (B, M)$ for $\mathcal{M}(x, b)$.
- Each component C of $\mathcal{M}(a', y')$ is a quotient space of finitely many components C_1, \dots, C_k of $\mathcal{M}(a, y)$. The new framing value is given by

$$f'(C) = \sum_{i=1}^k f(C_i) \in \mathbb{Z}/2.$$

- Each component C of $\mathcal{M}(x', b')$ is a quotient space of finitely many components C_1, \dots, C_k of $\mathcal{M}(x, b)$. The new framing value is given by

$$f'(C) = \sum_{i=1}^k f(C_i) + \chi \in \mathbb{Z}/2$$

where χ is the number of gluings $(B, P) \sim (B, M)$ with $s(B) = 1$.

Remark 4.5. By [JLS17, Prop.3.4] one should expect that χ is the number of gluings $(B, P) \sim (B, M)$ with $s(B) = 0$, and [JLS17, Prop.3.3] suggests one should add 1 for every gluing that is taking place. This is however the result of an unfortunate framing convention used in [JLS17, §3.3]. If we mirror the roles of P and M in [JLS17, Figure 12] one gets a convention in line with Definition 4.4. The reader may want to check that this alternative convention also leads to a framing on \mathcal{C}' below.

Lemma 4.6. *The tuple (\mathcal{C}', s', f') is a framed 1-flow category. Furthermore, there is an isomorphism*

$$\Phi: H^*(\mathcal{C}; \mathbb{Z}/2) \rightarrow H^*(\mathcal{C}'; \mathbb{Z}/2)$$

which commutes with Sq^2 .

Proof. It is clear that s' is a sign assignment, so we need to check the compatibility condition for $\partial\mathcal{M}(a', d')$.

First consider the case where $|d'| = |y'|$. If $d' \neq y'$, the compatibility condition is clear. Consider $(P, B, A), (M, B, A) \in \partial\mathcal{M}(a, y)$. Then (P, B, A) is a boundary point of an interval $\{P\} \times I \subset \mathcal{M}(x, y) \times \mathcal{M}(a, x)$ and of an interval $J_1 \times \{A\} \subset \mathcal{M}(b, y) \times \mathcal{M}(a, b)$ for some object b with $|b| = |a| - 1$. Similarly, (M, B, A) is the boundary point of an interval $\{M\} \times I$ and of an interval $J_2 \times \{A\}$. The interval I has a second endpoint (B', A') , leading to two more points (P, B', A') and (M, B', A') , which also are endpoints of intervals $J_3 \times \{A'\}$ and $J_4 \times \{A'\}$, respectively. In $\partial\mathcal{M}(a', y')$ the intervals $\{P, M\} \times I$ are no longer present, and $J_1 \times \{A\}, J_2 \times \{A\}$ have been glued together, as well as $J_3 \times \{A'\}, J_4 \times \{A'\}$ which have also been glued together.

In $\partial\mathcal{M}(a, y)$ the two intervals $\{P, M\} \times I$ are either in the same or in different components. If they are in the same component, then $\partial\mathcal{M}(a', y')$ has one more component than $\partial\mathcal{M}(a, y)$, and if they are in different components, then $\partial\mathcal{M}(a', y')$ has one less component than $\mathcal{M}(a, y)$. But notice that $\{M\} \times I$ contributes an extra summand 1 to the compatibility condition, which offsets the change in number of components. Therefore the compatibility condition is also satisfied at $\partial\mathcal{M}(a', y')$.

The cases where $|a'| = |x'| + 1$ and $a' = x'$ are similar, and will be omitted.

To get the required isomorphism, we can simply use the identity on the cochain level. We also need to check that we get the same Steenrod square $\text{Sq}^2: H^k(\mathcal{C}'; \mathbb{Z}/2) \rightarrow H^{k+2}(\mathcal{C}'; \mathbb{Z}/2)$. If we consider the case $k = |y|$, and a cocycle φ which is non-zero on y , choosing the combinatorial matching to match P and M makes it clear that the same Steenrod square is obtained. If the cocycle is zero on y , nothing changes.

If $k = |y| - 1$ we need to be more careful. Let $z \in H^k(\mathcal{C}'; \mathbb{Z}/2) = H^k(\mathcal{C}; \mathbb{Z}/2)$ be represented by the cocycle φ . To determine $\text{sq}^\varphi(x)$ we need to look at $\mathcal{M}(x, \varphi)$ which contains various intervals with endpoints (B, M) and (B, P) . A combinatorial matching will match B to some B' , resulting in further endpoints (B', M) and (B', P) . In $\mathcal{M}(a', \varphi')$ the points (B, M) and (B, P) are identified, and so are (B', M) and (B', P) . There are now various cases to consider, depending on the signs of B and B' , but also on how the components in $\Gamma_{\mathcal{C}}(x, \varphi)$ and $\Gamma_{\mathcal{C}}(x', \varphi')$ are different. We will consider a slightly more general case of this in the proof of Lemma 5.8 from which it will follow that the Steenrod square is indeed the same. \square

4.3. Combinatorial handle cancellation. In view of [JLS17] the next flow category move to transfer to 1-flow categories is the handle cancellation. To save space we will only describe the most simple case of handle cancellation, but this case will be sufficient for our purposes later.

Definition 4.7. Let (\mathcal{C}, s, f) be a framed 1-flow category, and let $x, y \in \text{Ob}(\mathcal{C})$ satisfy $|x| = |y| + 1$. Assume that $\mathcal{M}(x, y)$ consists of exactly one point, and that $\mathcal{M}(x, b) = \emptyset$ for all $b \in \text{Ob}(\mathcal{C}) - \{y\}$ with $|b| = |y|$, and $\mathcal{M}(a, y) = \emptyset$ for all

$a \in \text{Ob}(\mathcal{C}) - \{x\}$ with $|a| = |x|$. Then let (\mathcal{C}', s', f') be the framed 1-flow category given as follows:

- We have $\text{Ob}(\mathcal{C}') = \text{Ob}(\mathcal{C}) - \{x, y\}$ with $|\cdot|' : \text{Ob}(\mathcal{C}') \rightarrow \mathbb{Z}$ given by restriction of $|\cdot|$.
- The moduli spaces $\mathcal{M}(a, b)$ of \mathcal{C}' agree with the moduli spaces of \mathcal{C} .
- Both s' and f' are the restrictions of s and f , respectively.

Lemma 4.8. *The tuple (\mathcal{C}', s', f') is a framed 1-flow category. Furthermore, there is an isomorphism*

$$\Phi : H^*(\mathcal{C}; \mathbb{Z}/2) \rightarrow H^*(\mathcal{C}'; \mathbb{Z}/2)$$

which commutes with Sq^2 .

Proof. This is much simpler than the previous cases. We will only consider the compatibility condition for objects a, b with $|a| = |x| + 1$ and $|b| = |y| - 1$. Because \mathcal{C} is a framed 1-flow category, we get $[a : x] = 0$ and $[y : b] = 0$ for all such $a, b \in \text{Ob}(\mathcal{C})$. Furthermore, any intervals I in $\mathcal{M}(a, y)$ and J in $\mathcal{M}(x, b)$ will have both endpoints going through x , respectively y . Each such pair of intervals (I, J) gives rise to a square C in $\partial\mathcal{M}(a, b)$ with edges given by the intervals $\{+\} \times I$, $\{-\} \times I$, $J \times \{+\}$ and $J \times \{-\}$. Clearly $\tilde{f}(C) = 1$, so C does not contribute to the compatibility condition.

Note that $\partial\mathcal{M}(a, b)$ in \mathcal{C}' is obtained from $\partial\mathcal{M}(a, b)$ in \mathcal{C} by removing these squares. Since \mathcal{C} satisfies the compatibility condition, so does \mathcal{C}' now. The isomorphism on cohomology is induced by inclusion $\Phi : C_*(\mathcal{C}') \rightarrow C_*(\mathcal{C})$ of objects, and it is easy to see that the Steenrod square commutes with this isomorphism. \square

5. COMPUTATIONAL ASPECTS: THE SMITH NORMAL FORM

In Section 5, we finally describe the new algorithm for computing Steenrod squares in a framed flow category. We begin with the definition of a particularly simple type of 1-flow category, which the algorithm achieves upon termination.

Definition 5.1. A framed 1-flow category \mathcal{C} is said to be in *primary Smith normal form*, if the following are satisfied.

- If b is an object, there is at most one non-empty 0-dimensional moduli space $\mathcal{M}(a, b)$, which contains p^k elements, all of which have the same framing, where $p \geq 2$ is a prime and $k \geq 1$.
- For each object a there exists at most one object b with $|a| = |b| + 1$ and $\mathcal{M}(a, b)$ non-empty.

The idea of the algorithm is as follows:

- (1) restrict from a framed flow category to a framed 1-flow category,
- (2) use flow category moves to algorithmically reduce the 1-flow category to primary Smith normal form,
- (3) instead of keeping track of the 1-dimensional moduli space information, we only keep track of relevant combinatorial data which will allow us to calculate the combinatorial Steenrod square.

Primary Smith normal form for flow categories was considered in [LOS18, §6], where it was shown that flow category moves can be used to move any flow category into

this simple form, so we already know that (2) is always possible. To do the right flow category moves we only need to look at the integral chain complex, but keeping track of the 1-dimensional moduli spaces can be computationally costly. As this section is computational in nature, we need to know the “relevant combinatorial data” in (3), and how we can keep track of it in a computationally efficient manner. For this, several tricks are used for reducing the amount of combinatorial data we have to carry with us.

5.1. Ideas to speed up the process. The Khovanov cohomology of a link has relatively few generators compared to the number of generators of the Khovanov cochain complex. Turning the Khovanov flow category of [LS14a] into a Smith normal form can be a time-consuming process. Furthermore, trying to keep track of the framed 1-dimensional moduli spaces is a difficult task as the numbers explode quickly. For example, for the diagram of the torus knot $T_{7,4}$ used to make Steenrod square calculations in [JLS15], the number of points in the 0-dimensional moduli spaces can exceed 10^{10000} , while the largest torsion coefficient is 4.

Large numbers do not have to represent a problem for a computer program, but points in the 0-dimensional moduli spaces combine to endpoints of intervals in 1-dimensional which can be combined in more diverse ways. These combinations may not behave well after applying handle slides.

We illustrate the key ideas for getting around this computational issue with an example.

Example 5.2. Assume a framed 1-flow category \mathcal{C} contains objects a, b_1, \dots, b_5, c as in Figure 7. If we assume that every 0-dimensional moduli space only contains points of the same sign, then there are 95 intervals in $\mathcal{M}(a, c)$. But we also need to know how their endpoints are paired. For example, of the 50 endpoints going through b_1 , 20 may be paired with endpoints going through b_3 , 17 with endpoints going through b_4 and the remaining 13 going through b_5 .

Performing a handle slide where we slide b_3 three times over b_1 with subsequent Whitney trick requires us to keep track of how these numbers change.

On the other hand, if c is part of a cocycle φ whose Steenrod square we want to compute, we have to choose a combinatorial boundary matching \mathcal{C} for φ , which will involve the objects b_1, \dots, b_5 . Clearly we can match the same four points in $\mathcal{M}(b_1, c)$ in the same way for every cocycle φ involving c , while the fifth point needs to be matched with another point in a $\mathcal{M}(b_1, c')$, where c' depends on φ . Similarly in $\mathcal{M}(b_3, c)$ and $\mathcal{M}(b_4, c)$ we can even match all the points in the same way for every such φ . So when we do the actual Steenrod square calculation we add 75 edges to $\Gamma(a, c)$ in a way that will work for any φ involving c . Let us call this new special graph structure $\tilde{\Gamma}(a, c)$, so that we have

$$\Gamma(a, c) \subset \tilde{\Gamma}(a, c) \subset \Gamma_{\mathcal{C}}(a, \varphi).$$

Furthermore, several of the components C in $\Gamma_{\mathcal{C}}(a, \varphi)$ may only involve edges already in $\tilde{\Gamma}(a, c)$. So their contribution to $\text{sq}^{\varphi}(a)$ can be pre-computed and stored with $\mathcal{M}(a, c)$.

Notice that there are still 40 endpoints in $\tilde{\Gamma}(a, c)$, ten of which are of the form (A_i, B_5) with $A_i \in \mathcal{M}(a, b_1)$ and $B_5 \in \mathcal{M}(b_1, c)$. We can also choose a matching for the ten points in $\mathcal{M}(a, b_1)$ (and which would be relevant for calculating the

Steenrod square of a cocycle involving b_1) and add five edges to $\tilde{\Gamma}(a, c)$ leading to a new special graph structure $\bar{\Gamma}(a, c)$. After doing this for the objects b_2 and b_5 (we




FIGURE 7. The 0-dimensional moduli spaces between a and c .

do not need to deal with b_3 and b_4 as there are no further endpoints left involving them), the resulting special graph structure $\bar{\Gamma}(a, c)$ has only two endpoints left, one going through b_2 and one going through b_5 .

We would like to calculate the Steenrod square of φ using the special graph structures $\bar{\Gamma}(a, c)$ by matching the remaining endpoints for varying c . An obvious problem is that $\bar{\Gamma}(a, c) \not\subset \Gamma_{\mathcal{C}}(a, c)$. This leads indeed to a difficulty, which requires us to define the *distortion* below. But it turns out that we can recover the Steenrod square from the $\bar{\Gamma}(a, c)$ directly, which will require much less information than the moduli spaces $\mathcal{M}(a, c)$.

5.2. Partial combinatorial matchings. For a framed 1-flow category \mathcal{C} with a combinatorial matching \mathcal{C} for a cocycle $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ represented by the objects c_1, \dots, c_l , we have the special graph structure $\Gamma_{\mathcal{C}}(a, \varphi)$ with assigned value $o(\Gamma_{\mathcal{C}}(a, \varphi)) \in \mathbb{Z}/2$ from which we can derive the second Steenrod square. For any object c (not necessarily the dual of a cocycle) we have $\Gamma(a, c)$ is defined for any object a with $|a| = |c| + 2$. We would like to generalize $\Gamma_{\mathcal{C}}(a, c)$ to this non-cocycle setting, taking advantage of a global combinatorial matching. Note that if for some j we have that $\mathcal{M}(b_i, c_j)$ has an odd number of elements, there is one element that needs to be matched with an element of some $\mathcal{M}(b_i, c_{j'})$ where $j' \neq j$. We therefore cannot do a global combinatorial matching which would work for all possible Steenrod squares, as different cocycles require different generators from the 1-flow category. But if $\mathcal{M}(b_i, c_j)$ has an even number of elements, we can match them all within, and use this matching whenever c_j is part of the dual of a cocycle.

Definition 5.3. Let \mathcal{C} be a framed 1-flow category. A *partial combinatorial matching* \mathcal{P} is a collection of pairs (A_1, A_2) where $A_1, A_2 \in \mathcal{M}(a, b)$ are different points where $|a| = |b| + 1$, and no point is paired up more than once.

Notice that pairs are ordered, so if A_1 and A_2 have the same sign, we have a direction from A_1 to A_2 once we use this in a combinatorial matching.

The set of partial combinatorial matchings can be partially ordered using inclusion, and a maximal partial combinatorial matching has the property that if $\mathcal{M}(a, b)$ has

an even number of points, all of them are paired up, and if there is an odd number of points in $\mathcal{M}(a, b)$, there is exactly one point which is not paired.

Definition 5.4. Let \mathcal{C} be a framed 1-flow category and \mathcal{P} a partial combinatorial matching. If $a, c \in \text{Ob}(\mathcal{C})$ are objects with $|a| = |c| + 2$, define a special graph structure $\Gamma_{\mathcal{P}}(a, c)$ as follows. The vertex set V and $s: V \rightarrow \{0, 1\}$ are the same as for $\Gamma(a, c)$, and the same holds for E' and $f: E' \rightarrow \{0, 1\}$. If $(B_1, A), (B_2, A) \in \mathcal{M}(b, c) \times \mathcal{M}(a, b)$, where $(B_1, B_2) \in \mathcal{P}$ we add an edge to $E - E'$ between the vertices $(B_1, A), (B_2, A)$ which is directed from (B_1, A) to (B_2, A) if $s(B_1) = s(B_2)$ and undirected otherwise.

If $(B, A_1), (B, A_2) \in \mathcal{M}(b, c) \times \mathcal{M}(a, b)$, where $(A_1, A_2) \in \mathcal{P}$ and B is not matched to any other point B' in \mathcal{P} , we add an edge to $E - E'$ between the vertices $(B, A_1), (B, A_2)$ which is directed from (B, A_1) to (B, A_2) if $s(A_1) = s(A_2)$ and undirected otherwise. Furthermore, for every new edge between (B, A_1) to (B, A_2) (directed or not) we add a loop to L if $s(B) = 1 \in \mathbb{Z}/2$.

Example 5.5. Consider the 1-flow category \mathcal{C} indicated in Figure 8. The superscript in the elements of the 0-dimensional moduli spaces indicate their sign.




FIGURE 8. The 1-flow category \mathcal{C} .

The moduli space $\mathcal{M}(a, c)$ consists of three intervals, which we can choose to connect the points (C_1^+, B_1^+) and (C_4^+, B_4^-) , (C_2^+, B_2^-) and (C_2^-, B_3^-) , (C_3^+, B_2^-) and (C_3^-, B_3^-) .

The partial matching pairs (C_2^+, C_3^+) and (C_2^-, C_3^-) . The special graph structure $\Gamma_{\mathcal{P}}(a, c)$ contains one interval with endpoints (C_1^+, B_1^+) and (C_4^+, B_4^-) , and a circle consisting of four edges, two of which are oriented. Note that these two oriented edges point in opposite directions on the circle.

If we change the moduli space $\mathcal{M}(a, c)$ by choosing to connect the points (C_1^+, B_1^+) and (C_2^+, B_2^-) , (C_4^+, B_4^-) and (C_2^-, B_3^-) , (C_3^+, B_2^-) and (C_3^-, B_3^-) , the graph will consist of one interval made up of five edges, two of which are oriented and again in opposite direction.

If we turn the 1-flow category upside down, we get two extra loops in $\Gamma_{\mathcal{P}}(c^*, a^*)$ coming from the fact that B_2^- and B_3^- are negatively framed (one loop from the edge between (B_2^-, C_2^+) and (B_2^-, C_3^+) , and one from the edge between (B_3^-, C_2^-) and (B_3^-, C_3^-)).

Note however that in this example c does not represent a cocycle, and a represents a coboundary, so this 1-flow category does not contain interesting Steenrod squares.

5.3. Steenrod squares using partial combinatorial matchings.

Definition 5.6. For Γ a special graph structure and C a component of Γ which does not contain any vertices of ∂V define $o(C) \in \mathbb{Z}/2$ as follows. If C is in L then $o(C) = 1$. If C contains a vertex then $o(C) := 1 + F(C) + D(C)$.

With this new notation, if for some Γ we have $\partial V = \emptyset$ then we may write $o(\Gamma) = \sum_C o(C)$ where the sum is over all components of Γ .

Now consider again an arbitrary framed 1-flow category \mathcal{C} . If a cocycle φ is represented by c_1, \dots, c_l we want to obtain values $\text{sq}^\varphi(a) \in \mathbb{Z}/2$ for objects $a \in \text{Ob}(\mathcal{C})$ using $\Gamma_{\mathcal{P}}(a, c_j)$ giving rise to the Steenrod square. Observe that the special graph structure $\Gamma_{\mathcal{P}}(a, c_j)$ has as boundary points the pairs (B, A) with neither B nor A matched by \mathcal{P} .

Denote by $\Gamma_{\mathcal{P}}(a, \varphi)$ the special graph structure given by the disjoint union of the $\Gamma_{\mathcal{P}}(a, c_j)$. If we extend \mathcal{P} to a combinatorial matching \mathcal{C} for φ by pairing those $B \in \mathcal{M}(b_i, c_j)$ which are not paired in \mathcal{P} , we can extend $\Gamma_{\mathcal{P}}(a, \varphi)$ to a special graph structure $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ by adding edges between vertices (B, A) and (B', A) , where (B, B') are matched in \mathcal{C} , but B, B' and A are not matched in \mathcal{P} .

Notice that $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ differs from $\Gamma_{\mathcal{C}}(a, \varphi)$ in that there are edges in the former between vertices (B, A) and (B, A') if B is not matched in \mathcal{P} , but A and A' are. Nevertheless, $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ has no boundary points, so we can assign each component C a value $o(C)$. Adding these values does not give the right value for the Steenrod square, so we need to make a further definition first.

Definition 5.7. Given a cocycle φ , with respect to $\mathbb{Z}/2$ -coefficients, represented by objects c_1, \dots, c_l , and an object $a \in \text{Ob}(\mathcal{C})$ with $|a| = |c_j| + 2$, let $b_1, \dots, b_m \in \text{Ob}(\mathcal{C})$ be the objects with $\mathcal{M}(b_i, c_j) \times \mathcal{M}(a, b_i)$ non-empty for at least one c_j , and let \mathcal{P} be a maximal partial combinatorial matching.

Let u_i be the number of moduli spaces $\mathcal{M}(b_i, c_j)$ for $j = 1, \dots, l$ which have an odd number of elements. Note that u_i is even as φ is a cocycle.

Also, let v_i be the number of pairings in \mathcal{P} involving elements of $\mathcal{M}(a, b_i)$ between points of the same sign.

Define the *distortion of \mathcal{P}* by

$$d_{\mathcal{P}}(a, \varphi) = \sum_i (u_i/2)v_i \in \mathbb{Z}/2.$$

Furthermore, define

$$\widetilde{\text{sq}}^\varphi(a) = \sum_C o(C) + d_{\mathcal{P}}(a, \varphi) \in \mathbb{Z}/2$$


where the sum is taken over the components of $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ for some extension \mathcal{C} of \mathcal{P} .

Lemma 5.8. *Let $\varphi \in C^k(\mathcal{C}; \mathbb{Z}/2)$ be a cocycle and $a \in \text{Ob}(\mathcal{C})$ with $|a| = k + 2$. Let \mathcal{P} be a maximal partial combinatorial matching and \mathcal{C} an extension of \mathcal{P} to a combinatorial matching for φ . Then $\text{sq}^\varphi(a) = \widetilde{\text{sq}}^\varphi(a) \in \mathbb{Z}/2$.*

Proof. Let $B \in \mathcal{M}(b_i, c_j)$ and $B' \in \mathcal{M}(b_i, c_k)$ be matched in \mathcal{C} , but not in \mathcal{P} . Also, let $A, A' \in \mathcal{M}(a, b_i)$ be matched in \mathcal{P} . The special graph structures $\bar{\Gamma}_{\mathcal{C}}(a, c)$ and $\Gamma_{\mathcal{C}}(a, \varphi)$ differ in that in the former there is an edge between (B, A) and (B, A') , and one between (B', A) and (B', A') , while in the latter there is an edge between (B, A) and (B', A) , and one between (B, A') and (B', A') . Also, if $s(B) = 1$, there will be an extra loop in L coming with every edge (B, A) to (B, A') .


We need to see how the sum over the components $\sum_C o(C)$ changes when we change these edges. We have to check four cases, depending on whether $s(B) = s(B')$ and $s(A) = s(A')$ or not. We note that the purpose of the distortion is to count the number of steps where $s(A) = s(A')$.

Let us begin with the case $s(A) \neq s(A')$ and $s(B) \neq s(B')$. In this case all four relevant edges are non-directed. We have the following picture



with the left-hand side a subgraph of $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ and the right-hand side a subgraph of $\Gamma_{\mathcal{C}}(a, \varphi)$. Note that we can assume all marked edges to be marked by 0 because of the local move (c).

In $\bar{\Gamma}_{\mathcal{C}}(a, \varphi)$ the endpoints X, Y, Z and W are connected somehow, and there are three ways of doing this. First, X and Y connect, and Z and W connect. This leads to the picture




If we assume that there are a directed edges on the outside path from Y to X pointing in a fixed direction on the component, and b directed edges on the outside path from Z to W pointing in a given direction, the contribution to the sum $o(C)$ over the components from these two components is $2 + 1 + a + b$, where 2 comes from the two components, the 1 comes from an extra loop in L , since $s(B) \neq s(B')$ means that one of those signs is 1, and so one of the edges comes with an extra loop. The summands a and b come from $D(C)$.

On the right we only get one component C and there will be no extra loop, so the contribution is $1 + D(C)$. But it is easy to see that $D(C) = a + b$, so both pictures contribute the same to $\sum_C o(C) \in \mathbb{Z}/2$.

The second case connects X with Z and Y with W , which is basically the same case as the previous one.

In the third case we connect X with W and Y with Z . We now get one component in the graph in both cases, but if we choose the orientation to be along the path from X to W , the orientation on the path from Y to Z changes.




Note that the assumption $s(A) \neq s(A')$ and $s(B) \neq s(B')$ implies that $s(Y) = s(W)$. Hence if there are a directed edges between Y and W pointing from Y to W , then there are $1 + a$ directed edges between Y and W pointing from W to Y . The contributions to $o(C)$ thus differ by 1, but since on the left we would get one extra loop, the total contribution to $\sum_C o(C)$ is the same again.

Now assume that $s(A) \neq s(A')$ and $s(B) = s(B')$. The differences to the previous case are that there are no extra loops (or two extra loops, which we can ignore) when $s(B) = 0$ (when $s(B) = 1$), and that the vertical edges on the right are directed, which we can assume to both point downwards.

Again we have three cases to check, and it turns out that again all contributions to $\sum_C o(C)$ are the same. For example, in the third case we have to count either both of the two extra directed edges, or none of them. As we count mod 2, this has no effect.

Now consider the assumption $s(A) = s(A')$ and $s(B) \neq s(B')$, leading again to an extra loop in $\bar{\Gamma}_C(a, c)$. This leads to the following picture.



An argument as above shows that for all three cases of connecting the endpoints X, Y, Z and W the contribution to $\sum_C o(C)$ this time around differs by 1. However, notice that since $s(A) = s(A')$ this pair contributes to v_i for the appropriate i . Also, since \mathcal{P} is maximal and B, B' are not matched by \mathcal{P} , this pair contributes 1 to $u_i/2$. Thus, taking the distortion into account will lead to the correct result.

The final case $s(A) = s(A')$ and $s(B) = s(B')$ is again similar, requiring again the distortion to give the right outcome. \square

Remark 5.9. Recall that we did not quite finish the proof that the Whitney trick does not affect the Steenrod square in the case where $k = |y| - 1$. In view of partial combinatorial matchings, we can think of $\text{sq}^c(x')$ as being obtained using a partial matching which combines (B, P) with (B, M) and (B', P) with (B', M) instead of (B, P) with (B', P) and (B, M) with (B', M) , so the above proof shows that the Steenrod square is the same for both ways. Notice that distortion plays no role as $s(P) \neq s(M)$.

Lemma 5.8 states that we can calculate the Steenrod square from the information given by $\Gamma_{\mathcal{P}}(a, \varphi)$, once we extend \mathcal{P} to a combinatorial matching \mathcal{C} for φ . If the 1-flow category is in primary Smith normal form, the values u_i are all 0. In the process of turning a 1-flow category into primary Smith normal form we need to keep track of the changes in distortions.

Instead of keeping track of the 1-dimensional moduli spaces $\mathcal{M}(a, c)$ to calculate the Steenrod square, we can now just keep track of the $\Gamma_{\mathcal{P}}(a, c)$. From the point of

view of a computer program, this is presumably not much of an improvement, but in fact we only need to keep track of the equivalence class of each $\Gamma_{\mathcal{P}}(a, c)$. This is going to be an improvement, as we can always find a rather simple representative of the special graph structure.

Recall that by definition, vertices in Γ have valency at most 2, so the connected components of Γ consist of intervals (with vertices) or circles.

Lemma 5.10. *Let Γ be a special graph structure. Then Γ is equivalent to a special graph structure Γ' which has at most one circle component and in which case this component is a loop in L . Furthermore, each interval component of Γ' either consist of a single edge $e \in E'$ with $f(e) = 0$, or of three edges e_1, e_2, e with $e_1, e_2 \in E'$, $e \in E''$, and $f(e_1) = 0 = f(e_2)$.*

Proof. The proof is by induction on the number of edges in Γ . If Γ has only one edge, this edge e will form an interval component. If $f(e) = 1$, we can use a move (c) to change the framing by adding a loop. If the set of loops L consists of more than one element, use moves (e) to reduce the number of loops. This finishes the induction start.

Now assume the result holds for all special graph structures with no more than n edges, $n \geq 1$, and let Γ be a special graph structure with $n + 1$ edges. If every edge is labelled with 0 or 1, each edge is its own interval component, and we can use moves (c) and (e) to bring the special graph structure into the desired form. Otherwise the set of unlabelled edges $E - E'$ is non-empty. Notice that the endpoints of $e \in E - E'$ are not in ∂V .

Assume there exists an undirected edge $e \in E - E'$. Then, up to (c) moves we can assume that e fits into the left-hand side of a move (g) or (h). In both cases, we can reduce the number of edges up to equivalence and invoke induction.

If all the edges in $E - E'$ are directed, there either exists a component which has more than one directed edge, or all the components are interval components with one or three edges. In the latter case we can use moves (c) to get all labels to be 0, and then reduce the number of loops with move (e) to get the desired form. If there exists a component with more than one directed edge, first use moves (d) to ensure that all directed edges in this component have the same direction. As there are no undirected, unlabelled edges by assumption, the left hand-side of move (f) fits into the component (after possibly another move (c)). Up to equivalence, we can again reduce the number of edges, and hence use induction. The result follows. \square

Definition 5.11. Let (Γ, I) consist of Γ a special graph structure and I an ordered pairing of the boundary points ∂V of Γ . Let Γ' be a special graph structure equivalent to Γ as in Lemma 5.10. Then I is also an ordered pairing of the boundary points of Γ' . Furthermore, assume that the boundary points of every interval component of Γ' are paired in I , and that if the interval component contains a directed edge then the direction of the edge agrees with the direction coming from the ordered pairing in I .

Given such a Γ' , define

$$o(\Gamma, I) = o(\Gamma''),$$

where Γ'' is the sub-special graph structure of Γ' where all interval components have been removed.

Remark 5.12. Our goal is to start with a framed 1-flow category and turn it into a framed 1-flow category in primary Smith normal form. But rather than keeping track of all the 1-dimensional moduli spaces, we want to keep track of equivalence classes of special graph structures $\Gamma_{\mathcal{P}}(a, c)$, where \mathcal{P} is a maximal partial combinatorial matching. By Lemma 5.10 and Definition 5.11 we only need to keep track of a pairing of the endpoints $I(a, c)$ of $\Gamma_{\mathcal{P}}(a, c)$ and the element $o(\Gamma_{\mathcal{P}}(a, c), I(a, c)) \in \mathbb{Z}/2$ for every pair of objects a, c with $|a| = |c| + 2$, in order to understand the equivalence class.

Notice that $I(a, c)$ is given by pairs $(B, A) \in \mathcal{M}(b, c) \times \mathcal{M}(a, b)$ with neither B nor A matched in \mathcal{P} . The ordered pairing $I(a, c)$ of boundary points is thus a collection of tuples $((B, A), (\bar{B}, \bar{A}))$. If $s(B, A) = s(\bar{B}, \bar{A})$ we interpret this order as a directed edge

$$\begin{array}{ccc} (B, A) & \longrightarrow & (\bar{B}, \bar{A}) \\ \circ & \longrightarrow & \circ \end{array}$$

while in the case $s(B, A) \neq s(\bar{B}, \bar{A})$ we interpret this as an undirected edge

$$\begin{array}{ccc} (B, A) & \text{---} & (\bar{B}, \bar{A}) \\ \circ & \text{---} & \circ \end{array}$$

In view of Lemma 5.10 the directed edge refers to actually three edges.

Turning a framed 1-flow category into primary Smith normal form is done by turning the underlying chain complex into primary Smith normal form using row and column operations, which on the 1-flow category level are done via handle slides. We need to know how \mathcal{P} , $I(a, c)$ and $o(\Gamma_{\mathcal{P}}(a, c), I(a, c))$ change during this process. Notice that changing the direction of one directed edge in $I(a, c)$ changes the value of $o(\Gamma_{\mathcal{P}}(a, c), I(a, c))$. We thus have to make a choice for the first 1-flow category, and carefully keep track of directions in subsequent pairings.

To simplify notation, we define

$$o(a, c) := o(\Gamma_{\mathcal{P}}(a, c), I(a, c)).$$

Lemma 5.13. *Let (\mathcal{C}, s, f) be a framed 1-flow category and \mathcal{P} a partial combinatorial matching. Let $x, y \in \text{Ob}(\mathcal{C})$ with $|x| = |y| + 1$ and assume $\mathcal{M}(x, y)$ contains two points P, M with $s(P) = 0$ and $s(M) = 1$. Assume further that either P, M are matched in \mathcal{P} or there exist $P', M' \in \mathcal{M}(x, y)$ with $s(P') = 0$ and $s(M') = 1$ such that $(P, P'), (M, M') \in \mathcal{P}$.*

Let (\mathcal{C}', s', f') be the framed 1-flow category obtained by performing the Whitney trick on P, M , and let \mathcal{P}' be the partial combinatorial matching obtained by removing the pairings involving P, M and adding (P', M') if P and M are not paired in \mathcal{P} . Then

$$o(a, y) = o(a', y') \quad \text{with } I(a, y) = I(a', y')$$

and

$$o(x, d) = o(x', d') \quad \text{with } I(x, d) = I(x', d')$$

for all $a, d \in \text{Ob}(\mathcal{C})$ with $|a| = |x| + 1$ and $|d| = |y| - 1$.

Proof. Notice that the $I(a, c)$ involve points which are not matched by \mathcal{P} , so we need not change them.

If P, M are matched, the result for (a, y) follows directly from move (g), if the vertices $(P, A), (M, A) \in \mathcal{M}(x, y) \times \mathcal{M}(a, x)$ are on different intervals, and from move

(h) if they are on the same interval. For (x, d) consider the vertices $(B, P), (B, M) \in \mathcal{M}(c, d) \times \mathcal{M}(x, c)$ for some object c . If B is not matched in \mathcal{P} and we get an edge between (B, P) and (B, M) in $\Gamma_{\mathcal{P}}(x, d)$ together with a loop if $s(B) = 1$. Using move (g) we get exactly $\Gamma_{\mathcal{P}}(x', d')$ by the definition of the Whitney trick. If B is matched to some B' , we get from $\Gamma_{\mathcal{P}}(x, d)$ to $\Gamma_{\mathcal{P}}(x', d')$ with a move (a).

Now assume P is matched to P' and M is matched to M' . For (a, y) we have to use a move (a) to get the desired result, while for (x, d) we have to consider again vertices (B, P) and (B, M) together with the cases whether B is matched or not. Again one checks that the special graph structures are equivalent. \square

Remark 5.14. Because of Lemma 5.13 performing a Whitney trick almost never changes the values $o(a, y)$ and $o(x, d)$. The only missing case is where one of P, M is not matched by \mathcal{P} . But if, say, P is not matched and $[x : y] > 0$ there is no reason to perform a Whitney trick involving P .

We will therefore try to ensure that if \mathcal{P} does not match a point $A \in \mathcal{M}(a, b)$, the sign of A agrees with the sign of $[a : b]$. This will require attention when the sign of $[a : b]$ changes after a handle slide, and the number of points in $\mathcal{M}(a, b)$ remains odd.

5.4. Effects of handle slides on special graph structures. Handle slides affect the special graph structures in a more significant way. In this section we describe how to change the partial combinatorial matchings during handle slides, and their effect on $o(a, c)$ and $I(a, c)$ during the handle slides.

Let \mathcal{C} be a framed 1-flow category and \mathcal{P} a partial combinatorial matching, where we assume that all points of a 0-dimensional moduli space $\mathcal{M}(a, b)$ have the same sign (this can be achieved using the Whitney trick). Let $\mathcal{C}_S^\varepsilon$ be the result of sliding an object x over y . We can define a partial combinatorial matching \mathcal{P}' for $\mathcal{C}_S^\varepsilon$ as follows. The inclusion $\mathcal{M}(a, b) \subset \mathcal{M}(a', b')$ for all objects a, b with $|a| = |b| + 1$ allows us to consider $\mathcal{P} \subset \mathcal{P}'$. If $Y \in \mathcal{M}(y, b)$, write $Y' \in \mathcal{M}(x', b')$ for the new point, and similarly write $X' \in \mathcal{M}(a', y')$ for $X \in \mathcal{M}(a, x)$. If $Y_1, Y_2 \in \mathcal{M}(y, b)$ are paired as $(Y_1, Y_2) \in \mathcal{P}$, we also pair $(Y'_1, Y'_2) \in \mathcal{P}'$. For $X_1, X_2 \in \mathcal{M}(a, x)$ paired as $(X_1, X_2) \in \mathcal{P}$, we pair them as $(X'_1, X'_2) \in \mathcal{P}'$ provided that $\varepsilon = 0$. If $\varepsilon = 1$ we pair them as $(X'_2, X'_1) \in \mathcal{P}'$. Note that we assume that X_1 and X_2 have the same sign, so the order matters.

Lemma 5.15. *Let \mathcal{C} be a framed 1-flow category with all points of a 0-dimensional moduli space having the same sign, and \mathcal{P} a partial combinatorial matching. Let \mathcal{C}' be the result of sliding an object x over y , and let \mathcal{P}' be the partial combinatorial matching for \mathcal{C}' described above. Then we have*

$$\Gamma_{\mathcal{P}'}(x', d') = \Gamma_{\mathcal{P}}(x, d) \sqcup \Gamma_{\mathcal{P}'}(y, d)$$

where $\Gamma_{\mathcal{P}'}(y, d)$ agrees with $\Gamma_{\mathcal{P}}(y, d)$ except that the sign function s' on vertices is given by $s'(X, Y) = \varepsilon + s(X, Y)$ for the vertices $(X, Y) \in \mathcal{M}(c', d') \times \mathcal{M}(y', c')$.

Furthermore,

$$\Gamma_{\mathcal{P}'}(b', d') = \Gamma_{\mathcal{P}}(b, d)$$

for all $b \in \text{Ob}(\mathcal{C})$ with $|b| = |x|$, $b \neq x$ and $|d| = |x| - 2$, and

$$\Gamma_{\mathcal{P}'}(a', c') \sim \Gamma_{\mathcal{P}}(a, c) \sqcup I_{\mathcal{P}}(a, c),$$

where $a, c \in \text{Ob}(\mathcal{C})$ with $|a| = |x| + 1 = |c| + 2$, and \sim stands for equivalence under moves (a)-(i). Here $I_{\mathcal{P}}(a, c)$ is a special graph structure containing one interval $I_{B,A}$ for every pair of points $(B, A) \in \mathcal{M}(y, c) \times \mathcal{M}(a, x)$ with neither B nor A matched by \mathcal{P} . The framing value of $I_{(B,A)}$ is given by $1 + \varepsilon + s(B)$.

Proof. We have $\mathcal{M}(x', d') = \mathcal{M}(x, d) \sqcup \mathcal{M}(y, d)$ and the signs on vertices $(X, Y) \in \mathcal{M}(c', d') \times \mathcal{M}(y', c')$ are given by $s'(X, Y) = \varepsilon + s(X, Y)$ with the ε coming from the new sign of $Y \in \mathcal{M}(y', c')$. The framings of intervals in $\mathcal{M}(y, d)$ do not change. Adding edges from the partial combinatorial matching do not change directions or add extra loops as the signs of $\mathcal{M}(c', d')$ do not change, and signs of $\mathcal{M}(y', c')$ change uniformly by ε . Hence $\Gamma_{\mathcal{P}'}(x', d') = \Gamma_{\mathcal{P}}(x, d) \sqcup \Gamma'_{\mathcal{P}}(y, d)$ as claimed.

We also get $\Gamma_{\mathcal{P}'}(b', d') = \Gamma_{\mathcal{P}}(b, d)$ for $b \neq x$ with $|b| = |x|$ as there is no change for the special graph structure.

Now for a, c with $|a| = |x| + 1 = |c| + 2$ we have

$$\mathcal{M}(a', c') = \mathcal{M}(a, c) \sqcup \mathcal{M}(y, c) \times [0, 1] \times \mathcal{M}(a, x)$$

which means we get a new interval for every pair $(B, A) \in \mathcal{M}(y, c) \times \mathcal{M}(a, x)$ with endpoints (B', A) and (B, A') , where $A' \in \mathcal{M}(a', y')$ and $B' \in \mathcal{M}(x', c')$ are new elements.

If B is matched to \bar{B} in \mathcal{P} , then B' is matched to \bar{B}' in \mathcal{P}' , and the intervals corresponding to (B, A) and (\bar{B}, A) form a circle component C together with two oriented edges coming from the partial combinatorial matching. Since $s(B) = s(\bar{B})$ by assumption, and B' and \bar{B}' are matched in the same order as B and \bar{B} , we get $o(C) = 1 + 1 = 0$, so we can drop C from $\Gamma_{\mathcal{P}'}(a', c')$.

Similarly, if B is not matched in \mathcal{P} , but A is matched to $\bar{A} \in \mathcal{M}(a, x)$, the intervals corresponding to (B, A) and (B, \bar{A}) form a circle component C together with two oriented edges from \mathcal{P}' . Notice that these oriented edges come with an extra loop if $s'(B)$ or $s'(B')$ equal 1. As $s'(B) + s'(B') = \varepsilon$, we get an extra loop if $\varepsilon = 1$. But notice that for $\varepsilon = 1$ the pair A' and \bar{A}' are directed opposite to the pair A and \bar{A} , so that this extra loop is cancelled with $o(C)$. So again this has no impact on $\Gamma_{\mathcal{P}'}(a', c')$. \square

Notice that even if \mathcal{P} is maximal, \mathcal{P}' as described above need not be maximal. But if \mathcal{P} is maximal, each $\mathcal{M}(x, b)$ contains at most one point which is not matched. So if $\mathcal{M}(x, c)$ and $\mathcal{M}(y, c)$ each have one point which is not matched in \mathcal{P} , these two points are not matched in \mathcal{P}' although they both are in $\mathcal{M}(x', c')$. Let \mathcal{P}'' be the partial combinatorial matching for \mathcal{C}' by matching these extra pairs in $\mathcal{M}(x', c')$ and $\mathcal{M}(a', y')$, where there is a similar situation. This matching is maximal, provided \mathcal{P} is maximal. If the signs of the points agree, we choose the ordering where the element from $\mathcal{M}(x, c)$, resp. $\mathcal{M}(a, y)$, is first.

Remark 5.16. Let $I(x, d)$, resp. $I(y, d)$, be an ordered pairing of the boundary points of $\Gamma_{\mathcal{P}}(x, d)$, resp. $\Gamma_{\mathcal{P}}(y, d)$. Then $I' = I(x, d) \sqcup I(y, d)$ is an ordered pairing of the boundary points of $\Gamma_{\mathcal{P}'}(x', d')$ and it is easy to see that

$$o(\Gamma_{\mathcal{P}'}(x', d'), I') = o(\Gamma_{\mathcal{P}}(x, d), I(x, d)) + o(\Gamma_{\mathcal{P}}(y, d), I(y, d)).$$

To determine $o(x', d')$, we have to consider the following special graph structure Γ_s . Take the disjoint union of edges determined by $I(x, d)$ and $I(y, d)$ (as described in Remark 5.12). For each object c with $|c| = |x| - 1$, there is at most one point

$A \in \mathcal{M}(x, c)$ and at most one point $A' \in \mathcal{M}(y, c)$ not matched by \mathcal{P} . If there exists a point $B \in \mathcal{M}(c, d)$ not matched by \mathcal{P} , we add an edge (possibly directed if $s(A) = s(A')$ as elements of $\mathcal{M}(x', c')$) between (B, A) and (B, A') , and also a loop if $s(B) = 1$. A choice of pairings for Γ_s gives us a pairing $I(x', c')$, and

$$o(x', c') = o(x, c) + o(y, c) + o(\Gamma_s, I(x, c)).$$

A similar problem appears in $\Gamma_{\mathcal{P}''}(a', c')$. First note that $I_{\mathcal{P}}(a, c)$ contains at most one interval, since we now assume that \mathcal{P} is maximal. Let us assume we get such an interval, which means that there is a $A \in \mathcal{M}(a, x)$ and a $B \in \mathcal{M}(y, c)$ with either of them not matched by \mathcal{P} . This interval $I_{(B, A)}$ then has endpoints (B', A) and (B, A') , where $B' \in \mathcal{M}(x', c')$ and $A' \in \mathcal{M}(a', y')$ are new points.

In order to understand the effect on $o(a', c')$, we need to construct a new special graph structure Γ_s as follows. It contains the edge $I_{(B, A)}$ labelled $1 + \varepsilon + s(B)$.

If there is a point $\tilde{A} \in \mathcal{M}(a, y)$ not matched by \mathcal{P} , then $\Gamma_{\mathcal{P}}(a, c)$ has an endpoint (B, \tilde{A}) . In $\Gamma_{\mathcal{P}''}(a', c')$ we then get a new edge between this point and (B, A') . Let (\hat{B}, \hat{A}) be the endpoint of $\Gamma_{\mathcal{P}}(a, c)$ paired with (B, \tilde{A}) by $I(a, c)$ and denote this pairing by I . Let C be the component of $\Gamma_{\mathcal{P}}(a, c)$ bounded by I . Up to equivalence in $\Gamma_{\mathcal{P}}(a, c)$ we can assume that $o(C, I) = 0$. We add C to Γ_s , together with the edge between (B, \tilde{A}) and (B, A') . The endpoints of Γ_s are now (\hat{B}, \hat{A}) and (B', A) .

If there is a point $\tilde{B} \in \mathcal{M}(x, c)$ not matched by \mathcal{P} , we have another endpoint (\tilde{B}, A) in $\Gamma_{\mathcal{P}}(a, c)$, which is paired with some (\check{B}, \check{A}) in $I(a, c)$, and we denote this pairing by I' . Let C' be the component of $\Gamma_{\mathcal{P}}(a, c)$ bounded by I' . Again we can assume that $o(C', I') = 0$. We need to distinguish the cases where (\tilde{B}, A) is (\hat{B}, \hat{A}) and where they are different.

In the first case $C = C'$ and $I = I'$, and we obtain Γ_s by adding an edge between $(\tilde{B}, A) = (\hat{B}, \hat{A})$ and (B', A) . The new pairing of endpoints in $\Gamma_{\mathcal{P}''}(a', c')$ is then given by $I(a', c') = I(a, c) \setminus I$.

In the second case we add C' to Γ_s together with an edge between (\tilde{B}, A) and (B', A) . The endpoints of Γ_s are then given by (\hat{B}, \hat{A}) and (\check{B}, \check{A}) . Choose a pairing of these points which we denote as I'' . The new pairing of endpoints in $\Gamma_{\mathcal{P}''}(a', c')$ is then given by $I(a', c') = (I(a, c) \setminus (I \cup I')) \cup I''$.

Depending on whether $\mathcal{M}(x, c)$ and $\mathcal{M}(a, y)$ are odd, we get various possibilities for the endpoints of Γ_s described above (including the empty set), and we now simply call the pairing I_s .

Lemma 5.17. *Let \mathcal{C} be a framed 1-flow category with all points of a 0-dimensional moduli space having the same sign, and \mathcal{P} a maximal partial combinatorial matching. Let \mathcal{C}' be the result of sliding an object x over y , and let \mathcal{P}'' be the partial combinatorial matching for \mathcal{C}' described above.*

Let a, c be objects in \mathcal{C} with $|a| = |x| + 1 = |c| + 2$ and assume that there exists $(B, A) \in \mathcal{M}(y, c) \times \mathcal{M}(a, x)$ with neither B nor A matched by \mathcal{P} . Then

$$o(a', c') = o(a, c) + o(\Gamma_s, I_s) + d(a, c),$$

where $d(a, c) \in \mathbb{Z}/2$ is 0 if $\mathcal{M}(a, y)$ has an even number of points, and if $\mathcal{M}(a, y)$ has an odd number of points then $d(a, c) = k \pmod{2}$, where $k \geq 0$ is the integer such that $\mathcal{M}(a, x)$ has $2k$ or $2k + 1$ elements.

Proof. If $\mathcal{M}(a, y)$ has an even number of points the equation $o(a', c') = o(a, c) + o(\Gamma_s, I_s)$ is clear from the discussion above. But if $\mathcal{M}(a, y)$ has an odd number of points, we need to consider the following. For every matched pair $A_1, A_2 \in \mathcal{M}(a, x)$ we have to remove an edge between (\tilde{B}, A_1) and (\tilde{B}, A_2) , and replace it by edges between (\tilde{B}, A_i) and (B', A_i) for $i = 1, 2$. Notice that we have to reintroduce the component corresponding to the intervals we get for B and A_i , which was removed in the last part of the proof of Lemma 5.15. As in the proof of Lemma 5.8 we get an extra summand of 1 in the contribution to $o(a', c')$. Since all the points in 0-dimensional moduli spaces have the same sign and \mathcal{P} is maximal, this is done k times. \square

We say that \mathcal{C}' is obtained from \mathcal{C} by a *double slide*, if it is the result of two handle slides of an object x over y of the same sign. For a double slide we can do different ways of matching points, and in particular we have to worry less about new interval components. So the points of $\mathcal{M}(y, c)$, resp. $\mathcal{M}(a, x)$, which are matched in \mathcal{P} now appear twice in $\mathcal{M}(x', c')$, resp. $\mathcal{M}(a', y')$, and we can keep this matching for \mathcal{P}' for both copies. But if $B \in \mathcal{M}(y, c)$ is not matched in \mathcal{P} , we get two copies $B', B'' \in \mathcal{M}(x', c')$ which we can match. Similarly, $A \in \mathcal{M}(a, x)$ leads to two points $A', A'' \in \mathcal{M}(a', y')$ that we can match. As both signs are the same, the order matters, and we choose a consistent order $(A', A'') \in \mathcal{P}'$. However, for $B \in \mathcal{M}(y, c)$ we choose $(B', B'') \in \mathcal{P}'$ if $s(B) = 0$, and $(B'', B') \in \mathcal{P}'$ if $s(B) = 1$.

Lemma 5.18. *Let \mathcal{C} be a framed 1-flow category and \mathcal{P} a partial combinatorial matching. Let \mathcal{C}' be the result of double sliding an object x over y . With the partial combinatorial matching \mathcal{P}' for \mathcal{C}' described above, we get*

$$\Gamma_{\mathcal{P}'}(x', d') = \Gamma_{\mathcal{P}}(x, d) \sqcup L_{\mathcal{P}}(y, d),$$

where $|d| = |x| - 2$, and $L_{\mathcal{P}}(y, d)$ is the special graph structure with empty vertex set, and one loop for every interval component in $\Gamma_{\mathcal{P}}(y, d)$. For $b \neq x$ with $|b| = |x|$ we get

$$\Gamma_{\mathcal{P}'}(b', d') = \Gamma_{\mathcal{P}}(b, d).$$

For $|a| = |x| + 1 = |c| + 2$ we get

$$\Gamma_{\mathcal{P}'}(a', c') = \Gamma_{\mathcal{P}}(a, c).$$

Proof. We get

$$\mathcal{M}(x', d') = \mathcal{M}(x, d) \sqcup \mathcal{M}(y, d) \sqcup \mathcal{M}(y, d)$$

and any non-trivially framed circle in $\mathcal{M}(y, d)$ appears twice and can be disregarded. Similarly, any circle component in $\Gamma_{\mathcal{P}}(y, d)$ appears twice in $\Gamma_{\mathcal{P}'}(x', d')$ and can be disregarded. If we have an interval with endpoints $(C, B) \in \mathcal{M}(c, d) \times \mathcal{M}(y, c)$ and $(\tilde{C}, \tilde{B}) \in \mathcal{M}(\tilde{c}, d) \times \mathcal{M}(y, \tilde{c})$, we now get two intervals with endpoints (C', B') , (\tilde{C}', \tilde{B}') , and (C', B'') , $(\tilde{C}', \tilde{B}'')$. These two intervals are connected via directed edges from the partial matchings of B', B'' and \tilde{B}', \tilde{B}'' , and by the choice of matchings in \mathcal{P}' we see that this component C satisfies $o(C) = 1$. We can therefore replace this component with a loop in $L_{\mathcal{P}}(y, d)$.

For $b \neq x$ with $|b| = |x|$ we get no change in the moduli spaces, so the result follows.

For $|a| = |x| + 1 = |c| + 2$ each pair $(B, A) \in \mathcal{M}(y, c) \times \mathcal{M}(a, x)$ leads to two intervals with endpoints (B', A) , (B, A') and (B'', A) , (B, A'') in $\mathcal{M}(a', c')$. It is easy to see that these intervals combine to a component C with $o(C) = 0$, and hence can be disregarded in $\Gamma_{\mathcal{P}'}(a', c')$. \square

Recall that we required that all points in a 0-dimensional moduli space in \mathcal{C} have the same sign, but after a handle slide this need not be the case for \mathcal{C}' anymore. We want to resolve this using Lemma 5.13, but we need that if a point $P \in \mathcal{M}(b, c)$ is not matched, then its sign agrees with the sign of $[b : c]$. Note that after a handle slide we may get that $[b' : c']$ has a different sign than $[b : c]$ although the same point is still not matched.

The next lemma shows that we can compensate this by a change in $I(a, c)$ and $I(b, d)$.

Lemma 5.19. *Let \mathcal{C} be a framed 1-flow category and \mathcal{P} a maximal partial combinatorial matching. Assume that $\mathcal{M}(b, c)$ is a 0-dimensional moduli space with an odd number of points, and let B be the point in $\mathcal{M}(b, c)$ not matched in \mathcal{P} . Assume that $s(B)$ is different from the sign of $[b : c]$. Then there is a maximal partial combinatorial matching \mathcal{P}' such that the unmatched point $B' \in \mathcal{M}(b, c)$ in \mathcal{P}' satisfies $s(B) \neq s(B')$, and the following holds.*

- (1) *Assume that d is an object with $|d| = |b| - 2$ such that $\Gamma_{\mathcal{P}}(b, d)$ has a boundary point (C, B) , that is, $\mathcal{M}(c, d)$ has an odd number of points. Let $I(b, d)$ be a choice of ordered pairings of boundary points of $\Gamma_{\mathcal{P}}(b, d)$ with $((C, B), (\bar{C}, \bar{B})) \in I(b, d)$ for some boundary point (\bar{C}, \bar{B}) . Let $I'(b, d)$ be the choice of ordered pairings of boundary points of $\Gamma_{\mathcal{P}'}(b, d)$ differing from $I(b, d)$ only as follows.*

$$\text{If } \begin{array}{ccc} (C, B) & \longrightarrow & (\bar{C}, \bar{B}) \\ \circ & & \circ \end{array} \in I(b, d), \text{ then } \begin{array}{ccc} (C, B') & \longrightarrow & (\bar{C}, \bar{B}) \\ \circ & & \circ \end{array} \in I'(b, d).$$

$$\text{If } \begin{array}{ccc} (C, B) & \longrightarrow & (\bar{C}, \bar{B}) \\ \circ & & \circ \end{array} \in I(b, d), \text{ then } \begin{array}{ccc} (C, B') & \longrightarrow & (\bar{C}, \bar{B}) \\ \circ & & \circ \end{array} \in I'(b, d).$$

Then

$$o(\Gamma_{\mathcal{P}}(b, d), I(b, d)) = o(\Gamma_{\mathcal{P}'}(b, d), I'(b, d)).$$

- (2) *Assume that a is an object with $|a| = |b| + 1$ such that $\Gamma_{\mathcal{P}}(a, c)$ has a boundary point (B, A) , that is, $\mathcal{M}(a, b)$ has an odd number of points. Let $I(a, c)$ be a choice of ordered pairings of boundary points of $\Gamma_{\mathcal{P}}(a, c)$ with $((B, A), (\bar{B}, \bar{A})) \in I(a, c)$ for some boundary point (\bar{B}, \bar{A}) . Let $I'(a, c)$ be the choice of ordered pairings of boundary points of $\Gamma_{\mathcal{P}'}(a, c)$ differing from $I(a, c)$ only as follows.*

$$\text{If } \begin{array}{ccc} (B, A) & \longrightarrow & (\bar{B}, \bar{A}) \\ \circ & & \circ \end{array} \in I(a, c), \text{ then } \begin{array}{ccc} (B', A) & \longrightarrow & (\bar{B}, \bar{A}) \\ \circ & & \circ \end{array} \in I'(a, c).$$


$$\text{If } \begin{array}{ccc} (B, A) & \longrightarrow & (\bar{B}, \bar{A}) \\ \circ & & \circ \end{array} \in I(a, c), \text{ then } \begin{array}{ccc} (B', A) & \longrightarrow & (\bar{B}, \bar{A}) \\ \circ & & \circ \end{array} \in I'(a, c).$$

Then


$$o(\Gamma_{\mathcal{P}}(a, c), I(a, c)) = o(\Gamma_{\mathcal{P}'}(a, c), I'(a, c)).$$

Proof. There is a matched pair (B', B'') in \mathcal{P} such that $s(B) \neq s(B') = s(B'')$, and we define \mathcal{P}' by matching (B, B'') , leaving B' unmatched, and keeping all other matchings from \mathcal{P} . Given a $C \in \mathcal{M}(c, d)$ we have intervals I, I', I'' in $\mathcal{M}(b, d)$ with endpoints $(C, B), (C, B'), (C, B'')$. If C is matched in \mathcal{P} , these endpoints are part of the same edges in both $\Gamma_{\mathcal{P}}(b, d)$ and $\Gamma_{\mathcal{P}'}(b, d)$. If C is not matched by \mathcal{P} , then (C, B) is connected to another endpoint (\bar{C}, \bar{B}) with either $s(C, B) = s(\bar{C}, \bar{B})$ or not.

Up to move-equivalence, the situation is as follows in $\Gamma_{\mathcal{P}}(b, d)$ and $\Gamma_{\mathcal{P}'}(b, d)$.



where the middle edge in the top row may be directed or not. Using a move (a) or (i), we see that the graph structure $\Gamma_{\mathcal{P}'}(b, d)$ is equivalent to



Note that the diagonal edge in the component with (C, B') is directed if and only if the original edge in the component of (C, B) is not directed. We thus get $o(\Gamma_{\mathcal{P}}(b, d), I(b, d)) = o(\Gamma_{\mathcal{P}'}(b, d), I'(b, d))$.

The proof that $o(\Gamma_{\mathcal{P}}(a, c), I(a, c)) = o(\Gamma_{\mathcal{P}'}(a, c), I'(a, c))$ is essentially the same and will be omitted. \square

Remark 5.20. Note that the possibility $\begin{matrix} (\bar{C}, \bar{B}) \\ \longrightarrow \end{matrix} \begin{matrix} (C, B) \\ \longrightarrow \end{matrix} \in I(b, d)$ is ruled out by requiring $((C, B), (\bar{C}, \bar{B})) \in I(b, d)$ in Lemma 5.19. If we would require $((\bar{C}, \bar{B}), (C, B)) \in I(b, d)$ we could get an analogous situation with the arrows reversed.

5.5. The algorithm. We may turn a general framed 1-flow category \mathcal{C} into a framed 1-flow category in primary Smith normal form as follows. We begin with objects of maximal degree, and move on to lower degrees. More formally, assume our 1-flow category satisfies the conditions

- (1_m) If b is an object with $|b| \geq m$, there is at most one non-empty 0-dimensional moduli space $\mathcal{M}(a, b)$, which contains p^k elements, all of which have the same framing, where $p \geq 2$ is a prime and $k \geq 1$.
- (2_m) For each object a with $|a| > m$ there exists at most one object b with $|a| = |b| + 1$ and $\mathcal{M}(a, b)$ non-empty.

If $m \geq |a|$ for all objects $a \in \text{Ob}(\mathcal{C})$, these conditions are trivially satisfied, and if $m \leq |a|$ for all objects $a \in \text{Ob}(\mathcal{C})$, these conditions are equivalent to \mathcal{C} being in primary Smith normal form.

Identifying the necessary handle slides to turn a framed 1-flow category satisfying (1_m) and (2_m) into one satisfying (1_{m-1}) and (2_{m-1}) is straightforward. We need

to turn the matrix of $\partial_m: C_m(\mathcal{C}) \rightarrow C_{m-1}(\mathcal{C})$ into Smith normal form using row and column operations, which determine the necessary handle slides.

With this in mind the algorithm proceeds as follows.

Step 1 Obtain a framed 1-flow category, for example using [LS14c] or [JLS15].

Step 2 Choose a maximal partial combinatorial matching \mathcal{P} , which means one groups the elements of 0-dimensional moduli spaces $\mathcal{M}(a, b)$ into pairs, possibly leaving one point unpaired if there is an odd number of points. This can be done so that the unpaired point has the same sign as $[a : b]$.

Step 3 For every two objects a, c with $|a| = |c| + 2$ choose an ordering $I(a, c)$ and determine $o(a, c)$.

Step 4 Perform handle slides according to the Smith normalization process, starting with objects of maximal degree. If we have a single slide of x over y , we obtain

$$o(a', y') = o(a, y) + o(a, x)$$

for objects a with $|a| = |x| + 2$. For objects b, d with $|b| = |x| + 1 = |d| + 2$ the values $o(b', d')$ are obtained from Lemma 5.17, and for objects e with $|e| = |x| - 2$ the values $o(y', e')$ are obtained from $o(x, e)$ and $o(y, e)$ according to Remark 5.16. Note that in the last two cases the special graph structure Γ_s needs to be determined. Also determine the new $I(b', d')$ and $I(y', e')$ according to Remark 5.16.

In the case of a double slide we only need to change $o(x', e')$ using Lemma 5.18. This involves counting the number of elements in $I(y, e)$.

Step 5 After each handle slide, check whether $[b' : c']$ has a different sign than $[b : c]$, in which case $I(a', c')$ and $I(b', c')$ need to be changed according to Lemma 5.19.

If two objects a', b' satisfy $[a' : b'] = \pm 1$ and $[a' : \tilde{b}'] = 0 = [\tilde{a}' : b']$ for all objects $\tilde{a}' \neq a'$ and $\tilde{b}' \neq b'$, remove a', b' from the list of objects.

Repeat Step 4 and 5 until the 1-flow category is in primary Smith normal form.

The information of 0-dimensional moduli spaces can be stored in an integer. After Step 5 the Whitney trick can be performed using Lemma 5.13, which changes neither $o(a, c)$ nor $I(a, c)$.

5.6. Computational complexity. The number of generators in the Khovanov chain complex is exponential in the number of crossings, in fact, slightly worse since the number of components in the smoothings of the diagram create extra generators. By working with diagrams that combine several crossings as in [JLS15] we can reduce the number of generators somewhat. As we work one quantum degree at a time, we can slightly improve this further, but the interesting quantum degrees still tend to have the most objects.

Standard methods for turning a matrix into Smith normal form use matrices of the form

$$G = \begin{pmatrix} p & q \\ -b/g & a/g \end{pmatrix} \in SL_2(\mathbb{Z})$$

where $g = ap + bq$, and $a, b \in \mathbb{Z}$ are in the same column of the original matrix. This amounts to an operation on two rows, but if $a = \pm 1$, we can choose $p = a$ and $q = 0$ to reduce this to an ordinary row operation. When working mod 4 this is practically not a restriction. Turning an $n \times n$ -matrix this way into Smith normal form requires $O(n^3)$ arithmetic operations [Sto98], with the computational

improvements described in [Sto98] not being available to us, as they use modular techniques which hide the relevant row and column operations.

In our situation this means we have $O(n^2)$ handle slides, with each handle slide giving rise to $O(n)$ arithmetic operations. When performing a handle slide of an object x over y , we also need to form the special graph structure Γ_s for all objects d with $|d| = |x| - 2$, which potentially involves all objects c with $|c| = |x| - 1$. This does not worsen the number of operations in terms of complexity. Note that n is not the total number of objects, but the maximum number of objects within a grading.

In practice, the boundary matrices are sparse, but the number of operations still seems to depend strongly on the link and not just the number of objects in the 1-flow category. For example, with the diagram used for $T_{5,5}$ in Figure 1 we get for $q = 25$ a 1-flow category with about 83,000 objects, with the resulting calculation done in less than two hours on an ordinary desktop computer. This required about 2.8 million handle slides. For $q = 27$ the 1-flow category has about 169,000 objects, and almost 17 million handle slides are needed. The calculation itself took two days. For $q = 29$ the 1-flow category has around 264,000 objects, and the calculation was abandoned after several weeks. On the other hand, there is a diagram for $T_{-16,3}$ giving rise to a 1-flow category with more than 1.2 million objects for $q = -59$, with the Smith normalization process only taking a few hours and requiring about 7.2 million handle slides.

REFERENCES

- [Bau95] H.J. Baues, *Homotopy types*, Handbook of algebraic topology, North-Holland, Amsterdam, 1995, pp. 1–72.
- [CJS95] R. Cohen, J.D.S. Jones, and G.B. Segal, *Floer’s infinite dimensional Morse theory and homotopy theory*, The Floer Memorial Volume, Prog. in Math., vol. 133, Birkhäuser Verlag, 1995, pp. 297–325.
- [JLS15] D. Jones, A. Lobb, and D. Schütz, *An sl_n stable homotopy type for matched diagrams*, ArXiv e-print 1506.07725 (2015).
- [JLS17] D. Jones, A. Lobb, and D. Schütz, *Morse moves in flow categories*, Indiana Univ. Math. J. **66** (2017), no. 5, 1603–1657.
- [LLS15] T. Lawson, R. Lipshitz, and S. Sarkar, *Khovanov homotopy type, Burnside category, and products*, ArXiv e-print 1505.00213 (2015).
- [LOS18] A. Lobb, P. Orson, and D. Schütz, *Framed cobordism and flow category moves*, ArXiv e-print 1605.02003, to appear in Algebr. Geom. Topol. (2018).
- [LS14a] R. Lipshitz and S. Sarkar, *A Khovanov stable homotopy type*, J. Amer. Math. Soc. **27** (2014), 983–1042.
- [LS14b] R. Lipshitz and S. Sarkar, *A refinement of Rasmussen’s s -invariant*, Duke Math. J. **163** (2014), no. 5, 923–952.
- [LS14c] R. Lipshitz and S. Sarkar, *A Steenrod square on Khovanov homology*, J. Topol. **7** (2014), no. 3, 817–848.
- [Sch17] D. Schütz, *XKnotJob*, software (2017), Available at www.maths.dur.ac.uk/~dma0ds/knotjob.html.
- [Sto98] A. Storjohann, *Computing Hermite and Smith normal forms of triangular integer matrices*, Linear Algebra Appl. **282** (1998), no. 1–3, 25–45.
- [Thi99] M. Thistlethwaite, *Knotscape*, software (1999), Available at <https://www.math.utk.edu/morwen/knotscape.html>.
- [Wil17] M. Willis, *Stabilization of the Khovanov homotopy type of torus links*, Int. Math. Res. Not. IMRN (2017), no. 11, 3350–3376.

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UK

E-mail address: `andrew.lobb@durham.ac.uk`

DEPARTMENT OF MATHEMATICS, BOSTON COLLEGE, USA

E-mail address: `patrick.orson@bc.edu`

DEPARTMENT OF MATHEMATICAL SCIENCES, DURHAM UNIVERSITY, UK

E-mail address: `dirk.schuetz@durham.ac.uk`