

Protocol for a feasibility randomised controlled trial of the use of Physical ACtivity monitors in an Exercise Referral Setting: The PACERS study.

Dr Jemma Hawkins¹ hawkinsj10@cardiff.ac.uk

Dr Michelle Edwards¹ edwardsm28@cardiff.ac.uk

Dr Joanna Charles² j.charles@bangor.ac.uk

Prof Russell Jago³ russ.jago@bristol.ac.uk

Dr Mark Kelson⁴ kelsonmj@cardiff.ac.uk

Dr Kelly Morgan¹ morgank22@cardiff.ac.uk

Prof Simon Murphy¹ murphys7@cardiff.ac.uk

Dr Emily Oliver⁵ emily.oliver@durham.ac.uk

Prof Sharon Simpson⁶ sharon.simpson@glasgow.ac.uk

Prof Rhiannon Tudor Edwards² r.t.edwards@bangor.ac.uk

Dr Graham Moore¹ mooreg@cardiff.ac.uk

1. Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), Cardiff University, Cardiff, CF10 3BD, UK
2. Centre for Health Economics and Medicines Evaluation, Bangor University, Bangor, LL57 2PZ, UK
3. Centre for Exercise, Nutrition and Health Sciences, University of Bristol, Bristol, BS8 1TZ, UK
4. Centre for Trials Research, Cardiff University, Cardiff, CF14 4YS, UK
5. School of Applied Social Sciences, Durham University, Durham, DH1 3HN, UK
6. MRC/CSO Social and Public Health Sciences Unit, University of Glasgow, Glasgow, G2 3QB, UK

Corresponding author: *Dr Jemma Hawkins*

27 **Abstract**

28 **Background:** Exercise referral schemes are recommended by the National Institute for
29 Clinical Excellence (NICE) for physical activity promotion among inactive patients with
30 health conditions or risk factors. Whilst there is evidence for the initial effectiveness and
31 cost-effectiveness of such schemes for increasing physical activity, evidence of long-term
32 effects is limited. Techniques such as goal setting, self-monitoring and personalised feedback
33 may support motivation for physical activity. Technologies such as activity monitoring
34 devices provide an opportunity to enhance delivery of motivational techniques. This paper
35 describes the PACERS study protocol, which aims to assess the feasibility and acceptability
36 of implementing an activity monitor within the existing Welsh National Exercise Referral
37 Scheme (NERS) and proposed evaluation methodology for a full-scale randomised controlled
38 trial.

39 **Methods/Design:** The PACERS study consists of a pilot randomised controlled trial, process
40 evaluation and exploratory economic analyses. Participants will be recruited from the generic
41 pathway of the Welsh NERS and will be randomly assigned to receive the intervention or
42 usual practice. Usual practice is a 16-week structured exercise programme, the intervention
43 consists of an accelerometry-based activity monitor (MyWellnessKey) and an associated web
44 platform (MyWellnessCloud). The primary outcomes are predefined progression criteria
45 assessing the acceptability and feasibility of the intervention and feasibility of the proposed
46 evaluation methodology. Postal questionnaires will be completed at baseline (time 0: T0), 16
47 weeks after T0 (T1), and 12 months after T0 (T2). Routinely collected data will also be
48 accessed at the same time points. A subsample of intervention participants and exercise
49 referral staff will be interviewed following initiation of intervention delivery and at the end of
50 the study.

Discussion: The PACERS study seeks to assess the feasibility of adding a novel motivational component to an existing effective intervention in order to enhance effects on physical activity and support longer-term maintenance. The study will provide insight into the acceptability of activity monitoring technologies to an exercise referral population and delivery staff. Data from this study will be used to determine whether and how to proceed to a full-scale trial of effectiveness of the intervention, including any necessary refinements to intervention implementation or the proposed evaluation methodology.

Trial registration:

ISRCTN85785652

Keywords:

Exercise referral, Physical activity, Autonomous motivation, Feasibility studies, Accelerometer/try, Physical activity monitors, Physical activity trackers, Costs, Economic evaluation

Background

Physical inactivity is a major cause of preventable illness with large costs to the National Health Service (NHS) [1]. Increasing activity at the population level and among at-risk groups is a public health priority [2, 3]. Physical activity interventions for at-risk groups often involve advice and/or signposting from primary care practitioners [4]. Exercise referral schemes (ERS) are one common model [5], usually involving referral to a community-based structured exercise program. In Wales, the 16-week National Exercise Referral Scheme (NERS) has been running since 2007. A previous effectiveness study of the scheme [6] showed that, at 12 months, NERS was associated with improvements in physical activity for

patients at risk of coronary heart disease, but not for those referred for anxiety and depression, despite an improvement in their mental health [7]. The evaluation also showed the base-case incremental cost-effectiveness ratio was £12,111 per quality adjusted life year (QALY), falling to £9741 if participants were to contribute £2 per session [7]. Qualitative data highlighted a need for post-intervention motivational support to maintain changes [7, 8]. Whilst there is evidence for effectiveness of ERS in increasing physical activity in the short-term [9-11] evidence of long-term effects is limited. The Department of Health's Quality Assurance Framework for Exercise Referral [12] highlights the need to understand how to support long-term maintenance of changes in physical activity.

On entry to an ERS, patients may be initially motivated by external sources such as GP advice to attend [13, 14]. However, sustained changes in physical activity are consistently associated with more internalised, or autonomous, motivation [15-17]. According to Self-Determination Theory [18], the development of autonomous motivation can be achieved through supporting psychological needs for autonomy (volitional and self-endorsed engagement), competence (personal mastery and effectiveness) and relatedness (meaningful interpersonal connections). Thus, developing ways to support these three needs should help to maintain changes in physical activity. Support for this notion is provided by the randomised controlled trial of the Welsh NERS which found increases in autonomous motivation after scheme exit. This increase explained almost half of the between group difference in physical activity six months later [19]. Integrating processes to further enhance and sustain autonomous motivation during and after involvement in an exercise referral scheme may lead to larger effects and longer-term maintenance of these. Existing evidence points to potential motivational effects of techniques such as goal setting, monitoring, and personalised feedback on progress towards goals [20, 21] which may support autonomous

motivation by enhancing patients' sense of mastery and competence and are recommended by NICE as core components of behaviour change interventions [22].

Technologies such as activity monitors, provide opportunities to enhance delivery of goal setting and feedback, allowing for more frequent, and automatic feedback on progress toward activity goals, tailored updating of goals based on achievement, and remote contact with intervention providers [23]. In addition to addressing psychological needs for competence, incorporation of social components may support motivation through promoting relatedness to other service users. Research on such technologies in exercise interventions suggests that use can be quickly integrated in participants' lives [24] and may increase physical activity levels [25-29], however overall the evidence is equivocal [23]. Furthermore, little is known about the acceptability of these technologies to ERS populations or if the benefits will remain once the initial novelty has ceased. Exercise referral patients are a diverse group with a range of ages and conditions. For example, although the average age of participants in the evaluation of the Welsh NERS was 52 years old, the overall ages ranged from 16 to 88. Thus, familiarity with technology and willingness to use it may differ within the group [30]. In addition, participant diversity in terms of socioeconomic status and geographic location may result in differences in access to high speed internet connections or the hardware required for engaging with some technologies (e.g. personal computer). Hence, prior to a trial of effectiveness, which may be undermined by difficulties integrating technologies into routine practice or facilitating uptake by patients, piloting is required to investigate these issues.

A preliminary investigation [31] tested a protocol for integrating activity monitoring devices (MyWellnessKey, Technogym) and a linked web platform in one local authority area of the Welsh NERS. The study showed potential for using the MyWellnessKey (MWK) devices in

the scheme; however, further work is required to understand the feasibility and acceptability of this on a larger scale with a demographically diverse population. In this paper we describe the protocol of the PACERS study, a pilot trial to assess the feasibility and acceptability of using the MWK activity monitors to promote maintenance of physical activity within NERS. The aim of the study is to evaluate the feasibility and acceptability of the intervention (the MWK) and its proposed evaluation methodology, in order to optimise design and delivery and evaluate whether a full scale randomised controlled trial of effectiveness is warranted and feasible.

Study aim

The primary aim of the study is to assess the feasibility and acceptability of implementing the MWK activity monitors within the Welsh NERS as well as the proposed evaluation methodology in order to optimise design and delivery for conducting a definitive evaluation trial.

Study objectives

The main objectives for this study are to investigate:

- a) the fidelity of delivery of the intervention and trial methodology including compliance with study invitation and randomisation processes;
- b) the acceptability of the intervention to participants in terms of its usability and likelihood of future use;
- c) whether randomisation is acceptable to 50% or more of the participants;
- d) the feasibility of recruiting 20% or more new NERS patients and retaining at least 80% of participants at 12-month follow-up (T2);

- e) contamination, by exploring whether less than 20% of control participants are ~~the~~ exposed to the intervention;
- f) the effect of the intervention on the main hypothesised change mechanism (autonomous motivation);
- g) the feasibility of collecting the primary, secondary and process outcome measures and economic evaluation methods.

Methods

Study design

The study design is an individually randomised pilot randomised controlled trial, plus a process evaluation and exploratory economic analyses, of implementing the MWK devices within Welsh NERS standard practice. Data will be collected at three time points: baseline (time 0 (T0)), at the end of the 16-week NERS programme (T1) and 12-months post-baseline (T2). Figure 1 shows the study flow diagram. The study was given favourable ethical opinion for conduct in the NHS on 1st December 2015 by the South East Scotland Research Ethics Committee 02 (REF: 189587).

Figure. 1 Flow diagram of the PACERS study design

Setting and participants

The study is being undertaken within the Welsh NERS across leisure centres in eight local authority areas in Wales, UK. The eight study sites were purposively selected to reflect a range of urbanisation and geography. Participants are eligible for the study if they; i) are referred into the NERS generic pathway (see Box 1), and ii) have the capacity to use the activity monitors (i.e. computer access and an email address).

Box 1. NERS Generic Pathway Criteria

For referral into the NERS generic pathway, patients must:

- be aged 16 years or above;
- be sedentary (defined as not moderately active for 3 times per week or deconditioned through age or inactivity);
- have at least one of the following:
 - Raised blood pressure 140/90,
 - BMI >28,
 - Cholesterol >5.0,
 - Controlled diabetes or impaired glucose intolerance,
 - Family history of heart disease or diabetes,
 - At risk of osteoporosis and/or musculoskeletal pain,
 - Mild arthritis or poor mobility,
 - Mild-moderate COPD, asthma, bronchitis, emphysema,
 - Mild anxiety, depression or stress,
 - Multiple sclerosis.

Recruitment

Participants will be recruited to the trial using opportunistic invites within the existing scheme structure. NERS exercise professionals will provide information about the study to all new generic pathway clients during their first consultation appointment on the scheme. Exercise professionals will transfer the contact details of clients who are eligible for and interested in joining the study to the research team using a secure electronic form. The research team will send a recruitment pack containing full informed consent materials and the baseline questionnaire to interested clients to be returned by post. Participants who return a signed consent form and completed baseline questionnaire will be entered into randomisation. Participants in the intervention group will be sent information about the process evaluation interviews following randomisation and will be asked to express an interest in taking part in

the interviews. From those who express an interest, participants will be selected to provide variation in local authority area, age, sex, and reason for referral. Where possible we will interview the same participants at 4-weeks and 12-months. Where not possible, additional participants matched by demographics (e.g. age and sex) will be recruited for 12-month interviews. All NERS staff involved in the study will be invited to participate in the process evaluation interviews. From those who express an interest, two staff members per local authority area will be selected.

Randomisation

After completion of baseline measures, study staff will randomly assign participants 1:1 to receive either the intervention (NERS plus MWK) or the control treatment (NERS standard practice) via a computer-generated random allocation sequence created by the South East Wales Trials Unit.

The Intervention

Box 2. Features of the MWK activity monitor and MyWellnessCloud web platform

- Real-time visual feedback via a screen on the activity monitor
- Detailed feedback on activity levels via a web platform to indicate progress towards goals, time spent in different activity intensities and calories burned
- Automatised goal setting via an algorithm which sets goals in a stepwise fashion such that forward progression is mastery-based
- Facilitation of social support for exercise via the web platform (through involvement in group challenges and remote communication with an exercise professional) and smartphone app (the option to share details about activity completed via social media)
- Free access to the web platform and smartphone application following discontinuation of use of the MWK via manual input or by linking the account to another monitoring device.

The intervention is an enhanced ERS that includes usual care (NERS standard practice) [6] plus an accelerometry-based activity monitor (MyWellnessKey; MWK). The MWK can be used for self-monitoring of physical activity levels in combination with a linked web platform (MyWellnessCloud) and smartphone application (see box 2). The MWK has been validated in terms of device accuracy at monitoring physical activity level and intensity [32, 33] and utility at fostering increased physical activity levels (high concurrent validity with ActiGraph accelerometer to detect physical activity in laboratory and free living environments) [34]. Intervention participants will be provided with a MWK to use for the remaining 12 weeks of their 16-week NERS programme after receiving it at their 4-week consultation and will be encouraged to use it for 36 weeks after they exit the scheme, up until their 12-month consultation when the device will be returned. In current practice conducting an 8-month telephone consultation to check clients' progress with exercise is an optional part of standard care. To encourage participants to maintain engagement in the study we have asked for the telephone consultation to take place with all intervention participants. Table 1 shows how the intervention will be implemented within the scheme.

It is anticipated that the intervention will enhance NERS through two key mechanisms; 1) goal setting and personalised feedback elements of the devices will support a sense of exercise mastery and perceived competence; 2) the web platform will provide a sense of relatedness to others via opportunities to communicate remotely with exercise professionals, other NERS clients and social media contacts. It is anticipated that these mechanisms will improve autonomous motivation for exercise, leading to greater maintenance of increases in physical activity, as depicted in the intervention logic model (see Figure 2).

Figure 2. PACERS logic model

Control treatment

Control participants will receive usual care which is NERS standard practice; a 16-week structured exercise programme which includes consultations with an exercise professional at the start, 4-weeks, on exiting the scheme (16-weeks) and at 12-month follow-up [6].

Primary outcome

The primary outcome will be the feasibility and acceptability of the intervention and its proposed evaluation methodology, to inform a decision on whether a full randomised controlled trial is warranted and feasible. This will be assessed against a set of predefined progression criteria related to recruitment and retention rates, exposure to the intervention in both intervention and control groups, and acceptability of the intervention, recruitment and randomisation processes to participants. The criteria were agreed by the Trial Steering Committee (TSC) and follow a traffic light assessment system (red=stop; amber=discuss with TSC whether there is enough evidence that sufficient improvements can be made to proceed to full trial without another feasibility assessment; green=proceed) using quantitative measures supported by qualitative data. The criteria, their measurement, and assessment criteria are summarised in Table 2. Qualitative data will provide insights into intervention and evaluation design features which need to be retained, or where metrics fall into the amber zone, modifications which may need to be made to improve feasibility and acceptability.

It is anticipated that in a full trial, the main outcome measure will be objectively measured physical activity using accelerometry. To examine the feasibility of collecting this data at follow-up in the NERS population, a sub sample of participants will be recruited to complete

the accelerometer measure at 16 months post-randomisation. Participants will wear a GT3X ActiGraph accelerometer around the waist for seven consecutive days during waking hours. Data will be processed to identify mean minutes of moderate to vigorous intensity activity per day and mean accelerometer counts per minute (volume of physical activity) using established processes [35].

Secondary outcomes

The effect of the intervention on the main hypothesised change mechanism (autonomous motivation) will be evaluated. Other secondary outcome measures will be piloted to estimate key trial parameters (e.g. standard deviation) to inform a future full trial.

Measures collected routinely in NERS

Data collected routinely within NERS will be obtained for use within the trial from T0, T1 and T2, as follows:

- Blood pressure and resting heart rate;
- Body Mass Index;
- Waist circumference;
- Self-reported physical activity (Scottish Physical Activity Questionnaire) [36];
- Health-related quality of life (EQ-5D-5L) [37];
- Fitness test (Chester fitness test) [38].

Measures included in PACERS study questionnaire

The following additional measures will be collected at all time-points, which in a full trial would be used to assess effectiveness of the added intervention component (MyWellnessKey):

- Autonomous Motivation (Behavioural Regulation in Exercise Questionnaire 3 (BREQ-3)) [39];
- Psychological need support (Intrinsic Motivation Inventory) [40];
- Anxiety and depression (Hospital Anxiety and Depression Scale (HADS)) [41].

Economic evaluation outcome measures

The PACERS study questionnaire will include an adapted Client Service Receipt Inventory (CSRI) based on the previous service use questionnaire used in the NERS evaluation [7] and examples in the DIRUM database (dirum.org) to capture client health and social care service use since the last time point (plus a four month retrospective period at baseline). Additional questions in the 12-month questionnaire will capture wider economic outcomes including current work status, days off work due to health problems and estimated income lost due to changes in work during the study period. Willingness to pay for the MWK will also be explored. Baseline demographic data on housing status and household income will also be collected in the PACERS study questionnaire for the purpose of the economic analysis.

Sample size

The proposed sample size for the study of 286 participants was calculated to allow the estimation of the feasibility proportions of adherence and retention to within at least plus or minus 8.2 percentage points using a 95% confidence interval, as well as to provide 80% power to detect an effect size of 0.4 at the 5% level on the main hypothesised mediator of autonomous motivation at 12-month follow-up, assuming 30% attrition [7]. The sample size was also planned to provide an indication of likely response rates, permit estimates of effect sizes of primary and secondary outcomes in advance of a larger trial, and allow exploration of

socio-demographic patterning in uptake and use of the MWKs in order to generate hypotheses regarding who the intervention might work for and why.

Data collection

Routinely collected data will be extracted from the NERS database at all T0, T1 and T2. The PACERS study questionnaire will be mailed to participants at all time-points. Telephone and email reminders will be made to non-responders. Semi-structured telephone interviews will be conducted with a sub-sample of intervention participants (n=20) following receipt of the intervention at 4-weeks and again at 12-months (T2) to explore feasibility and acceptability of the intervention and study methods. In addition, telephone interviews will be conducted with a sample of NERS exercise professionals at the same time points to explore feasibility and acceptability of implementing the intervention and study methods from a professional perspective. Figure 3 indicates the schedule of enrolment, interventions and assessments.

Figure 3. PACERS study schedule of enrolment, interventions, and assessments.

Process evaluation

A detailed process evaluation will examine the acceptability and feasibility of the intervention and evaluation methods, including intervention delivery and fidelity, potential contamination and contextual influences. Quantitative and qualitative data will be collected using a range of methods. Table 3 summarises the process evaluation methods.

Economic analysis methods

Data will be collected to estimate intervention costs and examine the feasibility of calculating cost-effectiveness alongside a definitive full pragmatic randomised trial. Health care service use will be costed using national unit costs [42, 43]. Both arms of the study will be costed,

revisiting and revising the costing methodology used in previous economic analysis of the Welsh NERS [44].

The additional costs of the intervention will consist of: the cost of the MWK; staff costs relating to the MWK (e.g. training, implementation and participant follow up support); the cost of the professional web cloud (e.g. licence fee) and additional staff interactions. These costs are in addition to the core programme costs (in both arms) including: NERS standard practice costs and participant contributions. Information about the additional staff resources required for the use of the MWK and professional web cloud will be derived from qualitative interviews with staff.

Data analysis

Quantitative analysis

The main outcomes in this feasibility study are related to the study progression criteria as outlined in Table 2. The methods of analysis for quantitative data collected for the process evaluation are summarised in Table 3. Analyses will be largely descriptive, with summary statistics being presented overall and also by key demographics. Evidence of whether the intervention could lead to behaviour change will be examined using regression analyses to quantify effects on autonomous motivation, using the Relative Autonomy Index derived from the BREQ-3.

To examine the direction of effect on physical activity Analysis of Covariance models [ANCOVA] will be used to estimate intervention effects on physical activity at 16 months. While likely non-significant due to limited power, this should be in the direction of a favourable intervention effect. Accelerometer data will be processed using standard procedures; periods of ≥ 60 minutes of zero counts will be recorded as “non-wear time” and

removed. Participants will be included in the analysis if they provide ≥ 3 valid days (i.e. 500 minutes of data between 6am and 11pm). Mean minutes of daily moderate to vigorous intensity activity will be estimated using a threshold value of ≥ 2020 counts per minute with minutes of light intensity physical activity estimated using thresholds of between 100 and 2019 counts per minute [35]. Sedentary time will be estimated based on a cut-point of less than 100 counts per minute; mean sedentary minutes per day will be derived.

Qualitative analysis

Qualitative data from interviews with exercise professionals and intervention participants will be transcribed verbatim and organised and coded into a thematic framework using NVivo 11 software. An approach to thematic analysis will be used that allows for both a deductive and inductive approach to data analysis [45]. Data will be initially coded using an a priori coding scheme of categories which align with the research questions as a means of organising the data for subsequent interpretation. An element of flexibility will be maintained to account for the emergence of any new and unexpected themes. The first three transcripts will be independently coded by two coders in order to develop a shared codebook via consensus. Any disagreements between coders will be discussed with a third coder. Divergence and convergence between interviews will be examined and comparisons made of the experiences of the intervention across and within areas (NERS clients and exercise professionals). We aim to develop a comprehensive understanding of the intervention acceptability, implementation and mechanisms of impact.

Economic Analysis

A pilot cost-consequence analysis will be conducted from a NHS and societal perspective. Response rates and level of completion of the measures will be reported using descriptive

statistics. Variables will be checked for out of range values before analysis begins. As data are expected to be skewed, non-parametric tests will be used to assess differences across groups or time points for the outcomes of QALYs (using the EQ-5D) and health and social care service use. We will bootstrap (5,000 replications) differences in cost and outcomes to produce a 95% confidence interval around these differences. Ceiling effects on the EQ-5D will also be assessed, assessing the proportion of participants that state “no problems” on all five dimensions on the EQ-5D questionnaire. QALY gains (using the EQ-5D) will be compared to those in similar samples from previous literature (where available).

A report on the data gathered about service use (from routinely collected data recorded by healthcare professionals delivering NERS) will explore if future studies could use this or a different method to the CSRI questionnaire used in the feasibility study. Descriptive statistics will be used to describe the amount participants are willing to pay for the MWK, both during the intervention and beyond. Response rates and level of completion of the questions exploring how best to capture productivity losses will be reported using descriptive statistics. Sub-group analyses will explore the effect on health related quality of life of socio-demographics (e.g. gender) and reason for referral. Sensitivity analysis will be conducted in accordance with NICE guidelines to vary the cost of the device [46], demonstrating what happened in the feasibility trial and how co-ordination may be varied in a future full-scale trial.

Serious adverse event reporting and monitoring

It is not anticipated that there will be any additional risks to participants over and above existing NERS standard practice for which standard operating procedures are in place covering referral into the scheme, provision of exercise instruction and support, and dealing

with adverse events. There are no serious adverse events expected to be related to the intervention. Any serious adverse event occurrence will be reported to the Chief Investigator within 48 hours of receiving notification. Assignment of causality will be made by the independent clinician member of the TSC.

Project management

A Trial Management Group is responsible for ensuring the appropriate, effective and timely implementation of the trial including monitoring adherence to standardised research protocols. The day-to-day operational management of the feasibility study is co-ordinated by a central project management team which meets weekly to monitor progress and any issues which may need relaying to the Trial Management Group. An independent TSC provides overall supervision for the trial and advice through its independent chair and also encompasses the role of Data Monitoring Committee.

Discussion

The PACERS feasibility trial aims to assess the feasibility and acceptability of implementing a novel motivational component, the MyWellnessKey, into the existing Welsh NERS. In addition, the trial also aims to determine the acceptability and feasibility of the proposed evaluation methodology for a definitive trial of the intervention for promoting long-term maintenance of physical activity. Whilst exercise referral approaches have been shown to be effective for increasing physical activity levels, evidence of long-term effects is limited [9, 10, 12] and so there is a need to better understand how to support long-term maintenance of physical activity [3]. The MWK intervention offers a potential mechanism for enhancing and sustaining autonomous motivation for physical activity via evidence-based techniques

including goal setting, self-monitoring and receiving personalised feedback on progress towards goals [20-22].

Findings from this study will determine whether progression to a full scale randomised controlled trial of effectiveness and cost-effectiveness is feasible and warranted, through the assessment of key progression criteria. The study will assess whether the outcomes being used are feasible and acceptable to use with the study population. Findings related to the acceptability and feasibility of implementing the intervention will inform potential refinement of the implementation processes where necessary. The findings will also allow refinement of the intervention logic model.

List of abbreviations

CSRI: Client Service Receipt Inventory; ERS: Exercise Referral Scheme; MWK: MyWellnessKey; MWC: MyWellnessCloud; NERS: National Exercise Referral Scheme; NHS: National Health Service; NICE: National Institute for Clinical Excellence; NRES: National Research Ethics Service; QALY: Quality Adjusted Life Year; REC: Research Ethics Committee; TSC: Trial Steering Committee.

Ethics approval and consent to participate

The PACERS study was given favourable ethical opinion for conduct in the NHS on 1st December 2015 by the South East Scotland Research Ethics Committee 02 (REF: 189587) and participants are required to provide written informed consent in order to participate in the study.

Consent for publication

Not applicable, this manuscript does not contain participant data.

Availability of data and material

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Funding

The PACERS study is funded by the Health and Care Research Wales Health Research Award (REF: HRA1019). The work was also undertaken with the support of The Centre for the Development and Evaluation of Complex Interventions for Public Health Improvement (DECIPHer), a UKCRC Public Health Research Centre of Excellence. Joint funding (MR/KO232331/1) from the British Heart Foundation, Cancer Research UK, Economic and Social Research Council, Medical Research Council, the Welsh Government and the Wellcome Trust, under the auspices of the UK Clinical Research Collaboration, is gratefully acknowledged. SS was supported by a MRC Strategic Award (MC-PC-13027, MC_UU_12017_14, & SPHSU14).

Authors' contributions

JH is the Principal Investigator and responsible for overall management of the PACERS study. She led the development of the study protocol. ME is the trial manager and responsible for coordinating the PACERS study. ME was involved in finalising the study protocol and implementing study processes. GM is the principal co-investigator and was involved in finalising the study protocol, in particular the process evaluation methods. The first draft of the manuscript was prepared by ME, JH and GM. EO and SS were involved in finalising the study protocol, in particular the qualitative methods. RJ and KM were involved in finalising

the study protocol, in particular the accelerometry methods. MK was involved in finalising the study protocol, in particular the statistical considerations. SM was involved in finalising the study protocol, in particular the trial design. JC and RTE were involved in finalising the study protocol, in particular the health economics analysis. All authors reviewed, contributed to and approved the final manuscript.

Acknowledgements

The authors would like to thank the following members of the Trial Steering Committee for their advice and guidance in the development of the study protocol and progression criteria: Professor Colin Greaves, Ms Annys Randell, Dr Nefyn Williams, Professor Monica Busse-Morris and Ms Yvonne Sylvestre. The support of the NERS National Co-ordinator, Regional Co-ordinators, Exercise Professionals and Public Health Wales colleagues in set up of the study and intervention is also gratefully acknowledged.

References

1. British Heart Foundation. The economic costs of physical inactivity. British Heart Foundation National Centre. Loughborough University: 2013.
2. Department of Health. Start Active, Stay Active: A report on physical activity from the four home countries. London: 2011.
3. NICE. Exercise referral schemes to promote physical activity. London: NICE Public Health Guidance 2014a.
4. Din NU, Moore G, Murphy S, Wilkinson C, Williams NH. Health professionals' perspectives on exercise referral and physical activity promotion in primary care: Findings from a process evaluation of the National Exercise Referral Scheme in Wales. *Health Education Journal*. 2014;74(6):743-57.
5. Sowden SL, Raine R. Running along parallel lines: how political reality impedes the evaluation of public health interventions. A case study of exercise referral schemes in England. *J Epidemiol Community Health*. 2008;62(9):835-41.
6. Murphy S, Raisanen L, Moore G, Edwards RT, Linck P, Williams N, et al. A pragmatic randomised controlled trial of the Welsh National Exercise Referral Scheme: protocol for trial and integrated economic and process evaluation. *BMC Public Health*. 2010;10:352.
7. Murphy SM, Edwards RT, Williams N, Raisanen L, Moore G, Linck P, et al. An evaluation of the effectiveness and cost effectiveness of the National Exercise Referral Scheme in Wales, UK: a randomised controlled trial of a public health policy initiative. *J Epidemiol Community Health*. 2012;66(8):745-53.
8. Moore GF, Raisanen L, Moore L, Ud Din N, Murphy SM. Mixed-method process evaluation of the Welsh national exercise referral scheme. *Health Education Journal*. 2013;113(6):476-501.
9. Pavey TG, Taylor AH, Fox KR, Hillsdon M, Anokye N, Campbell JL, et al. Effect of exercise referral schemes in primary care on physical activity and improving health outcomes: systematic review and meta-analysis. *BMJ*. 2011;343.
10. Campbell F, Holmes M, Everson-Hock E, Davis S, Buckley Woods H, Anokye N, et al. A systematic review and economic evaluation of exercise referral schemes in primary care: a short report. *Health Technol Assess*. 2015;19(60):1-110.

- 535 11. Williams NH, Hendry M, France B, Lewis R, Wilkinson C. Effectiveness of exercise-
536 referral schemes to promote physical activity in adults: systematic review. *The British*
537 *Journal of General Practice*. 2007;57(545):979-86.
- 538 12. Department of Health. Exercise Referral Systems: A National Quality Assurance
539 Framework. London: 2001.
- 540 13. Rouse PC, Ntoumanis N, Duda JL, Jolly K, Williams GC. In the beginning: role of
541 autonomy support on the motivation, mental health and intentions of participants entering an
542 exercise referral scheme. *Psychol Health*. 2011;26(6):729-49.
- 543 14. Markland D, Tobin VJ. Need support and behavioural regulations for exercise among
544 exercise referral scheme clients: The mediating role of psychological need satisfaction.
545 *Psychology of Sport and Exercise*. 2010;11(2):91-9.
- 546 15. Teixeira PJ, Silva MN, Mata J, Palmeira AL, Markland D. Motivation, self-
547 determination, and long-term weight control. *International Journal of Behavioral Nutrition*
548 *and Physical Activity*. 2012;9(1):1-13.
- 549 16. Teixeira PJ, Carraça EV, Markland D, Silva MN, Ryan RM. Exercise, physical
550 activity, and self-determination theory: A systematic review. *International Journal of*
551 *Behavioral Nutrition and Physical Activity*. 2012;9(1):1-30.
- 552 17. Fortier MS, Wiseman E, Sweet SN, O'Sullivan TL, Blanchard CM, Sigal RJ, et al. A
553 moderated mediation of motivation on physical activity in the context of the PAC
554 randomized control trial. *Psychology of Sport and exercise*. 2011;12(2):7-78.
- 555 18. Deci EL, Ryan RM. The 'what' and 'why' of goal pursuits: Human needs and the self-
556 determination of behavior. *Psychological Inquiry*. 2000;11(4):227-68.
- 557 19. Littlecott HJ, Moore GF, Moore L, Murphy S. Psychosocial mediators of change in
558 physical activity in the Welsh national exercise referral scheme: secondary analysis of a
559 randomised controlled trial. *International Journal of Behavioral Nutrition and Physical*
560 *Activity*. 2014;11(1):1-11.
- 561 20. O'Brien N, McDonald S, Araujo-Soares V, Lara J, Errington L, Godfrey A, et al. The
562 features of interventions associated with long-term effectiveness of physical activity
563 interventions in adults aged 55-70 years: a systematic review and meta-analysis. *Health*
564 *Psychol Rev*. 2015;9(4):417-33.
- 565 21. Shilts MK, Horowitz M, Townsend MS. Goal setting as a strategy for dietary and
566 physical activity behavior change: a review of the literature. *Am J Health Promot*.
567 2004;19(2):81-93.

- 568 22. NICE. Behaviour change: individual approaches. London: NICE Public Health
569 Guidance 49, 2014b.
- 570 23. Petit A, Cambon L. Exploratory study of the implications of research on the use of
571 smart connected devices for prevention: a scoping review. BMC Public Health. 2016;16:552.
- 572 24. Hunt K, McCann C, Gray CM, Mutrie N, Wyke S. "You've got to walk before you
573 run": positive evaluations of a walking program as part of a gender-sensitized, weight-
574 management program delivered to men through professional football clubs. Health Psychol.
575 2013;32(1):57-65.
- 576 25. Cupples M, Dean A, Tully MA, Taggart M, McCorkell G, . Using Pedometer Step-
577 Count Goals to Promote Physical Activity in Cardiac Rehabilitation: A Feasibility Study of a
578 Controlled Trial. International Journal of Physical Medicine and Rehabilitation. 2013; 1:157.
- 579 26. Vaes AW, Cheung A, Atakhorrami M, Groenen MT, Amft O, Franssen FM, et al.
580 Effect of 'activity monitor-based' counseling on physical activity and health-related outcomes
581 in patients with chronic diseases: A systematic review and meta-analysis. Ann Med.
582 2013;45(5-6):397-412.
- 583 27. Davies CA, Spence JC, Vandelanotte C, Caperchione CM, Mummery WK. Meta-
584 analysis of internet-delivered interventions to increase physical activity levels. International
585 Journal of Behavioral Nutrition and Physical Activity. 2012;9(1):1-13.
- 586 28. Kirwan M, Duncan JM, Vandelanotte C, Mummery KW. Using Smartphone
587 Technology to Monitor Physical Activity in the 10,000 Steps Program: A Matched Case
588 Control Trial. J Med Internet Res. 2012;14(2):e55.
- 589 29. Butryn ML, Arigo D, Raggio GA, Colasanti M, Forman EM. Enhancing physical
590 activity promotion in midlife women with technology-based self-monitoring and social
591 connectivity: A pilot study. J Health Psychol. 2014;21(8):1548-55.
- 592 30. Olson KE, O'Brien MA, Rogers WA, Charness N. Diffusion of Technology:
593 Frequency of Use for Younger and Older Adults. Ageing international. 2011;36(1):123-45.
- 594 31. Hawkins JL, Oliver EJ, Wyatt-Williams J, Scale E, van Woerden HC. Assessing the
595 Feasibility of Using Uniaxial Accelerometers With an Online Support Platform in the
596 Delivery of a Community-Based Exercise Referral Scheme. Journal of Primary Care &
597 Community Health. 2014;5(4):288-90.
- 598 32. Bergamin M, Ermolao A, Sieverdes JC, Zaccaria M, Zanuso S. Validation of the
599 Mywellness Key in Walking and Running Speeds. Journal of Sports Science & Medicine.
600 2012;11(1):57-63.

- 601 33. Sieverdes JC, Wickel EE, Hand GA, Bergamin M, Moran RR, Blair SN. Reliability
602 and validity of the Mywellness Key physical activity monitor. *Clin Epidemiol.* 2013;5:13-20.
- 603 34. Herrmann SD, Hart TL, Lee CD, Ainsworth BE. Evaluation of the MyWellness Key
604 accelerometer. *British Journal of Sports Medicine.* 2011;45(2):109-13.
- 605 35. Troiano RP, Berrigan D, Dodd KW, Masse LC, Tilert T, McDowell M. Physical
606 activity in the United States measured by accelerometer. *Med Sci Sports Exerc.*
607 2008;40(1):181-8.
- 608 36. Lowther M, Mutrie N, Loughlan C, McFarlane C. Development of a Scottish physical
609 activity questionnaire: a tool for use in physical activity interventions. *British Journal of*
610 *Sports Medicine.* 1999;33(4):244-9.
- 611 37. EuroQol--a new facility for the measurement of health-related quality of life. *Health*
612 *Policy.* 1990;16(3):199-208.
- 613 38. Sykes K. Chester step test; resource pack (Version 3). Cheshire, UK:: Chester College
614 of Higher Education, 1998.
- 615 39. Markland D, Tobin V. A modification of the Behavioral Regulation in Exercise
616 Questionnaire to include an assessment of amotivation. *Journal of Sport and Exercise*
617 *Psychology.* 2004;26:191-6.
- 618 40. McAuley E, Duncan T, Tammen VV. Psychometric properties of the Intrinsic
619 Motivation Inventory in a competitive sport setting: a confirmatory factor analysis. *Res Q*
620 *Exerc Sport.* 1989;60(1):48-58.
- 621 41. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr*
622 *Scand.* 1983;67(6):361-70.
- 623 42. Curtis L, Burns A. Unit Costs of Health and Social Care. Personal Social Services
624 Research Unit, 2016.
- 625 43. Department of Health. NHS Reference Costs. London: 2016.
- 626 44. Edwards RT, Linck P, Hounscome N, Raisanen L, Williams N, Moore L, et al. Cost-
627 effectiveness of a national exercise referral programme for primary care patients in Wales:
628 results of a randomised controlled trial. *BMC Public Health.* 2013;13(1):1021.
- 629 45. Fereday J, Muir -Cochrane E. Demonstrating rigor using thematic analysis: A hybrid
630 approach of inductive and deductive coding and theme development. *International Journal of*
631 *Qualitative Methods.* 2006;5(1):80-92.
- 632 46. Andronis L, Barton P, Bryan S. Sensitivity analysis in economic evaluation: an audit
633 of NICE current practice and a review of its use and value in decision-making. *Health*
634 *Technol Assess.* 2009;13(29):iii, ix-xi, 1-61.

635 **Figure titles and legends**

636 Figure 1. Flow diagram of the PACERS study design. (page 8)

637

638 Figure 2. PACERS logic model. (page 11)

639

640 Figure 3. PACERS study schedule of enrolment, interventions, and assessments. (page 14)

641 X = study participants, **X = intervention delivery staff**

Table 1. Implementation of the intervention components

Time-point	Exercise professionals	Intervention participants
At 4 week review appointment	Set up participants with a MWC account, configure initial activity goals on the MWK and demonstrate how to use the device and web platform.	Take the MWK home, sign into their MWC account on their home computer and connect their MWK to read data and charge it.
Over the study period (48Weeks)	Interact with participants to monitor and adjust their goals, send positive comments and set up group challenges through direct messaging via a linked website called Professional Cloud.	Use the device daily and connect the MWK to a computer at least twice per week to upload data to the MWC, receive feedback and charge the device. Manually enter information about activity that the device does not readily measure, i.e. swimming, weight training, cycling.
At 8 months from start	Telephone participants to check on their progress with exercising and remind them of the study and encourage use of the MWK, MWC and associated features.	Participants with a MWK continue to use it daily.
At 12 months from start	Exercise professionals will have a consultation with all	Hand the MWK back to the exercise professional.

	participants for usual NERS assessments and to collect the MWK.	
--	---	--

Table 2 Summary of progression criteria

Progression Criteria (PC)	Measures used	Assessment of whether criteria have been met
PC1. Feasibility to recruit a sufficient proportion of new NERS patients to participate in the trial, with appropriate retention rates to 12 month follow-up.	<ul style="list-style-type: none"> • The percentage of new NERS patients recruited to the trial, and retained at each subsequent follow-up. • Regression models will be used to identify predictors of loss to follow-up (e.g. demographics or baseline motivation). 	<ul style="list-style-type: none"> • If >20% of new NERS patients recruited = proceed; if <5% = full-scale trial unlikely to be feasible. If 5-20% the TSC will consider the feasibility of proceeding to a full-scale trial bearing in mind the data and feedback presented and representativeness of the recruited sample, and possible steps to increase the recruitment rate. • If >80% retained at 12-months = proceed, if <60% = full-scale trial unlikely to be feasible. If 60-80% the TSC will consider the feasibility of proceeding based on the available data and possible steps to increase retention.
PC2. Intervention and trial methodology delivered as intended	<ul style="list-style-type: none"> • Summary statistics for intervention fidelity measures overall and by area. • Compliance with study invite processes. • Compliance with randomisation processes. 	<ul style="list-style-type: none"> • The TSC will consider the data presented and make a judgement about whether the intervention and trial methodology were delivered as intended.
PC3. At least one of the two intervention	<ul style="list-style-type: none"> • Percentages of participants who report acceptability of the intervention components on four self-report 	<ul style="list-style-type: none"> • The TSC will consider the quantitative and qualitative data and make an overall judgement on whether the intervention is

components is acceptable to participants	<p>questions.</p> <ul style="list-style-type: none"> • Issues regarding acceptability of, and engagement with, the two intervention components explored in qualitative interviews with a sub-sample of intervention participants. 	acceptable.
PC4. Recruitment and randomisation processes acceptable to >50% of recruited participants	<ul style="list-style-type: none"> • Percentages of participants who report acceptability of the recruitment and randomisation processes on patient questionnaires. • Exploration of understanding and acceptability of recruitment and randomisation processes in qualitative interviews. 	<ul style="list-style-type: none"> • >50% of recruited participants report 'agree' or 'strongly agree' to questions about the acceptability of recruitment and randomisation processes. • The TSC will apply discretion in judging whether this criterion has been met, or could be addressed to improve acceptability in a full-scale trial.
PC5. < 20% of control group exposed to the intervention components	<ul style="list-style-type: none"> • The percentage of participants in intervention and control groups who report that they were provided with a MWK device or accessed the MWC web platform. 	<ul style="list-style-type: none"> • <20% of control participants report that they have used a MWK device during the study period. • <20% of control participants report that they have accessed the MWC during the study period.

Table 3. Summary of process evaluation methods

Fidelity/Feasibility/ Acceptability	Method of data collection	Aims to explore	Method of Analysis / Data to be presented	Participants	Time
Fidelity to trial methodology (PC2)	Audio recordings of NERS initial consultations with participants	The accuracy with which recruitment and consent processes were followed.	A summary score of adherence to the processes (range 0-4) will be calculated for each recording and presented overall and by area	Two participants per exercise professional	T0 (during NERS initial consultation)
Feasibility of implementing the intervention and trial methodology within routine NERS practice	Telephone interviews with NERS staff	Barriers/ facilitators, fit with local context, any adverse effects on usual NERS delivery, differences across settings, additional infrastructure or resources required for a full trial.	Thematic analysis	Two exercise professionals per area	After receipt of the intervention at 4-weeks and at T2
Acceptability of the trial methodology (PC4)	Telephone interviews with NERS staff and intervention participants	Understandings and acceptability of recruitment and randomisation processes.	Thematic analysis	Two exercise professionals per area, 20 intervention participants	After receipt of the intervention at 4-weeks and at T2
	Self-report questions on study questionnaire		Percentages of participants reporting acceptability of the randomisation process	All participants	T1

Acceptability of the intervention (PC3)	Telephone interviews with professionals and participants patients	Perceived acceptability of intervention components, barriers and facilitators in using the devices.	Thematic analysis	Two exercise professionals per area, 20 intervention participants	After receipt of the intervention at 4-weeks and at T2
	Self-report questions on study questionnaire	Frequency of use, ease of use, likelihood of future use.	Percentages of participants reporting that the intervention was easy to use, that they used it, and would do so in the future	All intervention participants	T1 and T2
Feasibility of collecting objective data on physical activity at long-term follow up	ActiGraph accelerometers	The feasibility of obtaining measures of physical activity over a 7 day period	A linear regression model controlling for age, gender, baseline self-reported physical activity and randomisation group will be fitted. Results will be expressed using regression coefficients, 95% confidence intervals, and standardised effect sizes.	100 participants	16 months post-randomisation
Contamination (PC5)	Self-report questions on study questionnaire on awareness of and exposure to intervention components	Assessment of contamination between trial arms.	Percentages of participants in intervention and control arms reporting exposure to the intervention will be presented alongside 95% confidence intervals.	All participants	T1 and T2