



Marine resource reliance in the human populations of the Atacama Desert, northern Chile – A view from prehistory

Charlotte L. King ^{a,*}, Andrew R. Millard ^b, Darren R. Gröcke ^c, Vivien G. Standen ^d, Bernardo T. Arriaza ^e, Siân E. Halcrow ^a

^a Department of Anatomy, University of Otago, Dunedin, Otago, New Zealand

^b Department of Archaeology, Durham University, Durham, United Kingdom

^c Department of Earth Sciences, Durham University, Durham, United Kingdom

^d Departamento de Antropología, Universidad de Tarapacá, Arica, Chile

^e Instituto de Alta Investigación, Universidad de Tarapacá, Arica, Chile



ARTICLE INFO

Article history:

Received 13 November 2017

Accepted 5 December 2017

Keywords:

Carbon
Nitrogen
South America
Dietary isotopes
FRUITS

ABSTRACT

The Atacama Desert is one of the most inhospitable terrestrial environments on Earth, yet the upwelling of the Humboldt Current off the coast has resulted in the presence of a rich marine biota. It is this marine environment which first enabled the human settlement of the northern Atacama Desert, and continues to form the basis of regional economies today. In this paper we explore how the desert has shaped human dietary choices throughout prehistory, using carbon and nitrogen isotope analysis of human bone collagen ($n = 80$) to reconstruct the diets of the inhabitants of the Arica region of the northern Atacama. This area is one of the driest parts of the desert, but has been generally understudied in terms of dietary adaptation. Statistical analysis using FRUITS has allowed deconvolution of isotopic signals to create dietary reconstructions and highlight the continued importance of marine resources throughout the archaeological sequence. Location also appears to have played a role in dietary choices, with inland sites having 10–20% less calories from marine foods than coastal sites. We also highlight evidence for the increasing importance of maize consumption, coinciding with contact with highland polities. In all periods apart from the earliest Archaic, however, there is significant variability between individuals in terms of dietary resource use. We conclude that marine resource use, and broad-spectrum economies persisted throughout prehistory. We interpret these results as reflecting a deliberate choice to retain dietary diversity as a buffer against resource instability.

© 2017 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

The Atacama Desert is one of the most extreme environments on Earth. As the driest hot desert in the world, life here is extremely marginal. Natural terrestrial resources are few, and freshwater availability is extremely limited (Williams et al., 2008). Human habitation of the desert, even today, is restricted to the valleys of the snowmelt-fed and seasonal rivers which traverse the Andean cordillera, and desert oases (Santoro et al., 2005). Yet, humans have occupied the desert for thousands of years (Arriaza et al., 2008). This is primarily due to the nearby marine environment. The Humboldt Current upwells off the north coast of Chile, bringing

with it nutrient-rich waters and sustaining a rich and complex marine ecosystem (Thiel et al., 2007). The presence of plentiful marine resources meant that habitation of the desert was possible even prior to the adoption of agriculture, after which the ability to manipulate the desert environment began (Santoro et al., 2017).

In order to evaluate how different societies and natural environments affected diet in the Atacama Desert, however, there is a need for palaeodietary data from multiple parts of the desert, and different time periods. Palaeodietary work already undertaken in the Atacama Desert suggests, relatively unsurprisingly, that human subsistence choices were constrained by the desert environment (Santana-Sagredo et al., 2015; Torres-Rouff et al., 2012). There is evidence from the incipient agricultural sites (dating from 1700 BCE) of coastal valleys of extreme northern Chile, such as Pampa Tamarugal and the Loa River Valley, that during the transition to agriculture, desert populations retained significant

* Corresponding author.

E-mail address: charlotte.king@otago.ac.nz (C.L. King).

marine-resource input into their diet (Bonilla et al., 2016; Pestle et al., 2015a; Santana-Sagredo et al., 2015; Torres-Rouff et al., 2012). This has been interpreted as reflecting the presence of trade networks (Pestle et al., 2015a, 2015b) to mitigate the instability of terrestrial crops through use of marine resources (Santana-Sagredo et al., 2015). While trade networks developed during the Formative Period (ca. 1700BCE – 450CE), these socio-economic interactions continued during all subsequent periods, allowing some areas to become focused on maize agriculture and camelid pastoralism (e.g. Pestle et al., 2016; Torres-Rouff et al., 2015). Other regions seemed to have retained broad-spectrum based subsistence despite being influenced by polities whose dietary focus was maize (Knudson et al., 2007), such as the Tiwanaku, Wari (ca. 450–900CE) and later Inka peoples (1450–1600CE). These polities whose homelands were in the Peruvian highlands, expanded and annexed surrounding areas in the Middle Period (Wari and Tiwanaku) and Late Period (Inka). While there has, in the past, been a strong focus on the impact of these external polities and their reliance on maize in the Atacama Desert, it is becoming increasingly recognised that this crop has played a variable role in the subsistence economies of the desert people (Cuellar, 2013; Tykot et al., 2006). Similarly, the role of different ecological niches in subsistence decisions is just beginning to be explored in the region (Zaro, 2007).

To date there have been relatively extensive isotopic studies of diet conducted in the San Pedro de Atacama and Tarapacá regions, but there is a dearth of data from the Arica region. These areas, while all located in the Atacama Desert, have quite different local environments, and therefore the potential for different subsistence choices in prehistory. For instance, the desert varies quite considerably in terms of freshwater availability and therefore suitability for human habitation and agriculture. In San Pedro de Atacama, in inland northern Chile, for example, archaeological sites are centred around the natural oases. Agriculture was possible in this area (Llagostera and Costa, 1999; Nuñez, 2007) and it was likely an important stopping point for camelid traders moving from the political centres of the Andean highlands (Hubbe et al., 2012; Kolata, 1991; Llagostera, 1996). In Peru there are fog ‘oases’ in the desert, providing moisture which increases floral variability (Beresford-Jones et al., 2015) and the potential for agricultural yields (Sandweiss et al., 1999). The desert in the Tarapacá and Arica regions, however is extremely dry, experiencing less than 0.6 mm of rain per year (Williams et al., 2008). Agriculture even today is possible in very restricted areas centering around the river valleys and inland oases.

As well as geographic variation in water availability, there has also been climatic variation in rainfall over time. The El Niño southern oscillation has serious effects on the amount of rainfall in any given year, and El Niño cycles have varied in intensity throughout prehistory (Gayo et al., 2012; Moseley and Keefer, 2008; Sandweiss et al., 2009). Around 500 BCE, for example, palaeoclimatic data suggest a change to El Niño regime, resulting in higher levels of humidity in the Atacama Desert (Gayo et al., 2012), and considerably expanding the agriculture land available. During this time period, archaeological evidence suggests the development of extensive field systems alongside villages such as Ramaditas, Guatacondo and Caserones (Uribe and Vidal, 2012; Vidal et al., 2012), located in harsh environments where today agriculture is not possible. In these sites, remains of *Zea mays*, *Phaseolus lunatus*, *P. vulgaris*, *Lagenaria*, *Arachis hypogaea*, *Chenopodium quinoa* and *Algarrobo* (*Prosopis*) pods have been identified (McRostie et al., 2017; Santoro et al., 2017). These plants are not endemic to the northern Atacama and it is likely that they were introduced from the Andes and the eastern lowlands. This research aims to assess both the impact of the arrival of these crops, and the marginality of the desert environment on resource choices.

2. Environmental context: Arica region

The Arica region (Fig. 1) has four snowmelt-fed rivers, the Lluta, San Jose, Chaca and Camarones. Further to the south it is more truly arid, with the Loa River and its tributaries providing some of the only freshwater (Santoro, 2012). Use of the terrestrial environment in the Arica region, however, is further complicated by the presence of high levels of contaminating heavy metals (arsenic, lithium and boron) in the rivers (Apata et al., 2017; Figueiroa et al., 2012). This means that, while there is freshwater, its use and consumption can have serious health implications (Arriaza et al., 2010; Swift et al., 2015).

Dietary adaptations in the Arica region have not yet been investigated using isotopic techniques, but instead inferred using archaeological evidence. Archaeological evidence suggests that marine resource consumption was fairly ubiquitous throughout prehistory, with marine faunal remains and material culture associated with fishing found even in interior valley sites (Table 1). From around 7000 BCE the area played host to a large Archaic period population of Chinchorro marine hunter-gatherers, who settled the coast and relied heavily on the ocean for all sustenance (Arriaza et al., 2008; Standen et al., 2017). From 1700 BCE, however, there appears to have been a change in the El Niño regime, resulting in depression of the fisheries, relocation of the coastal peoples into inland river valleys, and the beginnings of agriculture in the region (Grosjean et al., 2007; Moreno et al., 2009; Williams et al., 2008). The cultivation of the interior valleys will have provided the people of the region with the ability to produce staple terrestrial crops for the first time. Andean domesticates such as potato (*Solanum*), ullucu (*Ullucus*), and quinoa (*Chenopodium*) become common in the archaeological record (Pearsall, 2008). Later in prehistory the region begins to interact with highland Andean polities such as the Tiwanaku (Muñoz, 1983, 1995), eventually being annexed by the Inka Empire in the Late Period (Santoro et al., 2010). These cultures would have allowed resource trade with other parts of their empires, as well as having specific ideas surrounding diet. In particular, both the Tiwanaku people and the Inka Empire placed ceremonial importance on the consumption of maize (Goldstein, 2003; Staller, 2010). There is ethnographic and archaeological evidence that, despite the marginality of agricultural land, the area became significant for maize production under the Inka Empire (Murra, 1980; Santoro et al., 2010).

In this study we investigate the effect of the desert environment on prehistoric resource-use in the Arica region. We hypothesise that the marginality of the desert means that ancient populations are unlikely to have relied solely upon agricultural resources in any period. Instead a sustained reliance on marine resources is likely. The ocean has always been a plentiful source of food, leading to a maritime tradition which persists even into the present day. We therefore examined diet in coastal and near-coastal valley sites from both agricultural and pre-agricultural periods to examine whether or not dietary diversity, and marine resource use, is maintained throughout prehistory. We then compare the isotopic data to other lines of archaeological evidence relating to resource use to build a fuller picture of subsistence strategies in the region.

3. Archaeological context

The studied samples derive from archaeological sites in the northern Atacama Desert, close to the modern-day city of Arica (Fig. 1). Sites lie both on the coast, and in the near-coastal areas of the Azapa Valley. This region has a long human occupation history, but not all periods and cultures are present both inland and on the coast. For example, the Archaic period is not well-represented in the inland valleys. Conversely there is a lack of evidence for Middle




Fig. 1. Location of the study area and surrounding regions. Inset gives study site locations and occupation periods with the Azapa valley expanded for clarity.

Period occupation of the coast (Muñoz, 1982; Sutter, 2000). The known prehistoric cultural sequence is therefore briefly described here.

3.1. Archaic period

The first evidence for human settlement of the Arica region is between 9400–8200 BCE (Arriaza et al., 2008; Moreno et al., 2009), and corresponds with the presence of small communities of hunters, fishers, and collectors well adapted to exploitation of the marine ecosystem. Most bioarchaeological and archaeological evidence suggests that Chinchorro populations were almost completely reliant upon the marine environment for subsistence (Arriaza et al., 2017; Aufderheide et al., 1993). However, Holden (1994) and Reinhard et al. (2011) have both reported possible cases of potato starch and quinoa seeds in Chinchorro coprolites at Morro 1/6 site, suggesting the inhabitants harvested and consumed wild plants. By 5000 BCE these coastal populations developed extraordinary funerary practices including artificial mummification that lasted until ca. 1500–1000 BCE (Arriaza, 1995; Standen et al., 2014; Standen, 2003). These dates mark the end of the Chinchorro cultural tradition and also the transition towards food production in the coastal valleys of the Arica region (Grosjean et al., 2007; Santoro et al., 2017).

3.2. Formative period

Although fisher-gatherer populations continue to live on the coast during the following Formative Period, for example, at the site of Quiani (ca. 1600–1500 BCE), there is evidence that plants such as squash and gourds were being incorporated into mortuary rituals. Recent studies show there is a correlation between edible plants offered as grave goods in the prehistoric sites of the Arica region, the species identified in human dental calculus (as phytoliths and starch grains) and the frequency of dental caries (Arriaza et al., 2017). Archaeological evidence suggests that later Formative Period coastal groups like Faldas del Morro (800 BCE) had strong links to the inland valley sites and were incorporating agricultural products into their diet (e.g. (Belmonte, 1998; Erices, 1975).

There is, however, much that remains unknown regarding the development of early farming cultures in the Arica region. While

Formative coastal populations were incorporating agricultural products into their diets, it is probable that they maintained marine specialisation, with terrestrial products forming only a very small portion of dietary intake. Indeed, most coastal sites are dominated by objects associated with maritime subsistence: including hooks, harpoons, fishing lines and hook weights (Muñoz, 1993; Muñoz and Focacci, 1985). Inland sites also retain material culture associated with fishing, indicating that marine resources remained important (Santoro, 1980b). This is counter to traditional archaeological interpretations which have inferred that inland Formative sites are much more reliant upon agricultural resources.

The first farmers of the inland Azapa Valley sites have generally been associated with the Alto Ramírez cultural Phase, which according to Rivera (1994, 1975) corresponds with a migration from the complex Formative centres of the highlands, such as Wankarane and Pukara, down to the coastal valleys. This diffusionist model reflects archaeological paradigms which tended to try to establish cultural dependency between the marginal societies of the coastal valley in the Atacama Desert and the large power centers of the highlands. Although other studies (Muñoz, 2004; Núñez and Santoro, 2011) have highlighted the important and active role that local populations played during the Archaic-Formative transition, there must have been some exogenous contributions in this process. The first agricultural crops grown in the area are not endemic to the region and must have been brought in either through trade or movement of peoples or both. It is therefore possible that the inland valley populations of the Arica region will have had a diet based upon terrestrial resources and been buffered by exchange networks which extend to the highlands, as interpreted by Pestle et al. (2015a) further to the south.

3.3. Middle period

There is also debate in the archaeological literature over the extent to which the Arica region was involved in the expansion of highland polities (such as Tiwanaku and Wari) during the Middle Period. Traditionally the presence of Tiwanaku-style ceramics, particularly vessels (*queros*) used for the consumption of *chicha* (maize beer) has been used as evidence for these sites coinciding with the Middle Period (450–900 CE). However, there has been debate over the level to which groups were influenced by these

Table 1

Sites sampled in this study, geographic location, archaeologically-inferred subsistence regime and assigned time period with reasoning and key primary data sources.

	Archaeozoological/ botanical remains	Subsistence related material culture	Inferred subsistence strategy from archaeological evidence	Site dates (dating method)	Time period	Reason	Number of individuals sampled
Morro 1	Molluscs, crustaceans, seaweed, fish, marine mammals and birds, few camelid bones and skins (Standen, 2003). Plant fibre present in coprolites (Reinhard et al., 2011)	Harpoons, fishhooks, nets, ropes, 'chopes' (shellfish processing tools), spear, bows, darts, projectile points (Standen, 2003)	Marine hunter-gatherer	6226–1748 cal BCE (Allison et al., 1984) 3500 – 1600 cal BCE (Standen, 2003) (radiocarbon dates on burials)	Archaic	Radiocarbon dates	5
Az115	Curcubita, maize, tubers, camelids, dog, guinea pig, birds, fewer fish and marine mammal bones (Muñoz, 2017)	Darts, spear tips, slings	Broad-spectrum agriculturalists	100–600CE (radiocarbon dates on burials)	Late Formative	Radiocarbon dates. Although some recent dates place some burials in the Middle Period (Muñoz, 2017) there are no Tiwanaku cultural features	14
Az14	Shellfish, fish, guinea pig, maize, quinoa, manioc, camote (Santoro, 1980a)	Chopes, bows, wooden points, harpoons (context uncertain)(Santoro, 1980a)	Broad-spectrum agriculturalists	900–560BCE uncalibrated (Santoro, 1980b)	Formative	Radiocarbon dates	8
Az141	Maize, camote, squashes, coca	Wooden digging/tilling sticks	Agro-pastoralists	1044–1290 cal. CE (Korpisaari et al., 2014) 890 ± 100CE (Schiappacasse et al., 1991) 1018–1276CE (Sutter, 2005)	Middle Horizon – Late Intermediate	Both Tiwanaku and Cabuza pottery present	5
Az6	Maize, camote, squash, coca, dog, guinea pig, llama, shellfish/shells, birds (Focacci, 1990)	Wooden digging/tilling sticks, bows, arrows, quivers	Agro-pastoralists	380CE (Focacci, 1990) 890–1392 CE(Focacci, 1990) 850–1350CE (Korpisaari et al., 2014)	Middle Horizon – Late Intermediate	Both Tiwanaku and Cabuza ceramics and other items of material culture present	5
Az11	Camelids, dogs, guinea pigs, monkey, birds, rodents, maize, beans, squash, camote, manioc, potato, quinoa, molluscs, fish, freshwater snails	Arrowheads, darts, hoes, sticks, wooden shovels, harpoons, chopes, fishhooks	Agro-pastoralists	790–980CE (Muñoz and Focacci, 1985)	Middle Horizon	Radiocarbon dates but Maytas pottery (not Tiwanaku)	5
Az8	Maize, sweet potato, manioc, squash, beans, quinoa, camelids	Arrowheads, darts, hoes, sticks, wooden shovels	Agro-pastoralists	1150–1350CE (Rothhammer and Santoro, 2001) 910AD – 1120CE (Espouey et al., 1995) 500–890CE (Nuñez, 2007)	Late-Intermediate	Radiocarbon dates	13
Cam8	Maize, sweet potato, manioc, squash, beans, quinoa, camelids, dogs, fish, marine mammals (Muñoz, 1989)	Wooden shovels, harpoons, arrowheads	Marine subsistence complemented by agricultural crops	1050–1560CE (Thermoluminescence of potsherds) (Schiappacasse et al., 1991)	Late	Inka pottery, textiles	10
Cam9	Birds, sealions, sea turtle, fish, shellfish, camelids, maize, squash (Ulloa et al., 2000)	Harpoons, bows, darts, oars, fishhooks	Marine subsistence complemented by agricultural crops	1050–1560CE (Thermoluminescence of potsherds) (Schiappacasse et al., 1991)	Late	Inka textiles	15

polities. In the Arica region two cultural groups seem to have been present; Maytas and Cubuza, with Cabuza groups more influenced by the Tiwanaku. In this study we focus on the Cabuza sites as representative of the Middle Period. However, recent radiocarbon-dating of mortuary offerings in some of these Cabuza sites, has revealed them to belong to the Late-Intermediate Period (900–1450 CE), rather than the earlier Middle Period (Korpisaari et al., 2014). It is possible that Tiwanaku presence and influence in the Arica area was the result of political collapse in the Bolivian highlands, and displacement of peoples into the Northern Chilean valleys, rather than deliberate annexation during the height of Tiwanaku power (Korpisaari et al., 2014). Nonetheless, the importance of maize in the Tiwanaku culture is often inferred to have had an impact in the sites of the Arica region (Muñoz, 1983).

3.4. Later periods

The foundation of maize agriculture laid by the Middle Period polities is usually assumed to have been elaborated by cultural groups during the Late-Intermediate Period (900–1450 CE), culminating in the ritualised maize consumption that characterises the Late Period and associated Inka state (1450–1600 CE) (Goldstein, 2003; Staller, 2010). The cultivation and consumption of *chicha* underlay most social and ritual transactions during the Inka period, and was symbolically associated with imperial power (Bauer, 1996; Goldstein, 2003; Staller, 2010).

3.5. Chronology of sites used in this study

The dating of sites in the Arica region has been conducted somewhat haphazardly, and sometimes more recent radiocarbon dates directly contradict earlier dates (Korpisaari et al., 2014; Muñoz, 2017; Sutter, 2005). It was not possible to take radiocarbon samples for this project, we therefore use a combination of existing radiocarbon data and artefactual evidence for cultural affinity to assign sites used to archaeological phase. For instance, we consider sites with Tiwanaku pottery as representing the Middle Period, because their subsistence regime is likely to have been affected by Tiwanaku cultural values, similarly sites with Inka influence (i.e. the Camarones sites) are likely to have been influenced by Inka maize-reliance.

Table 1 gives the sites used in this study, their assigned phase and the rationale for that assignment. The focus of this isotopic study, however, is not chronologically constraining differences in subsistence choices, but instead understanding how the desert environment affected resource use during prehistory as a whole. Our analysis and interpretation does not look at change through time, but instead considers differences between inland and coastal sites, and sites with evidence for interaction with highland polities and those without. Diversity in resource use, or lack thereof, within and between sites will give us insight into resource-use choices regardless of chronological gaps and issues with the construction of cultural sequences.

4. Materials and methods

Analysis of stable carbon ($\delta^{13}\text{C}$) and nitrogen ($\delta^{15}\text{N}$) isotopic ratios from human collagen provide an insight into the diet of past human society. In particular, it is possible to differentiate marine from terrestrial resource use, the photosynthetic pathways of plant resources used, and the trophic level of food indicating meat consumption or the type of marine resources exploited (Ambrose and Norr, 1993; Schoeninger and DeNiro, 1984). Carbon isotopes are fractionated differently according to plant photosynthetic pathway (C_3 vs. C_4), with C_3 plants favouring fixation of lighter ^{12}C more than

C_4 plants do resulting in a more negative $\delta^{13}\text{C}$ value (Ambrose and Norr, 1993). In the Atacama Desert, crops such as quinoa, tubers and pulses (e.g. beans) are C_3 crops, and are differentiable from the C_4 crop, maize (which is less ^{12}C -enriched). Carbon isotopes may also be used to highlight marine resources, with marine carbonate concentrating ^{13}C , resulting in less negative $\delta^{13}\text{C}$ values in comparison to C_3 plants (Chisholm et al., 1982).

Nitrogen isotopic ratios vary with trophic level, and each step up the food chain increases $\delta^{15}\text{N}$ values between 2‰ and 6‰ (DeNiro and Epstein, 1981; O'Connell et al., 2012). This means that broadly speaking, in the absence of aquatic resource consumption, $\delta^{15}\text{N}$ values can be related to levels of terrestrial meat consumption. However, as marine food-chains tend to involve more steps, consumption of high trophic level marine resources results in especially high $\delta^{15}\text{N}$ values (Fry, 2006; Minagawa and Wada, 1984). $\delta^{15}\text{N}$ values in Atacama Desert terrestrial foodchains are also enriched in ^{15}N relative to other areas of the world (Díaz et al., 2016). This is a result of the aridity of the environment and resulting differences in nitrogen cycling (Ehleringer et al., 1992). Arid conditions result in the volatilisation of nitrogen compounds such as ammonia and this process preferentially involves the light isotope ^{14}N , thus concentrating ^{15}N in soils and foliage (Amundson et al., 2003), an enrichment which is carried through terrestrial food-chains (Gröcke et al., 1997; Hartman, 2011).

4.1. Materials

This study involved the sampling and isotopic analysis of adult individuals from the archaeological collections of the Museo Arqueológico San Miguel de Azapa (MASMA), Arica. The samples derive from the archaeological sites of the Azapa and Camarones river valleys, and the coastal sites of Arica (Fig. 1). Sites and numbers of individuals sampled are listed on Table 1, a complete list of individuals sampled is given in Table S1. MASMA enforces strict sampling regulations to preserve their collections. The sampling of bone was therefore restricted to 25 adult individuals from each phase. Collagen was assessed using Durham University Archaeology laboratory protocols and was considered to be of good quality if: C/N ratio = 2.9–3.6 and 35–50% carbon and 11–16% nitrogen (e.g. DeNiro, 1985). A number ($n = 13$) of individuals from the Archaic phase sampled had poor preservation of collagen, with the collagen extracted not passing standard quality control checks (see S1 Table). After excluding those with poor quality collagen the total analysed sample set comprised of 80 individuals. The poor preservation of collagen in the Archaic samples has been previously documented in this region (Aufderheide et al., 1993; Silva-Pinto et al., 2014), and persists despite repeat preparation procedures and addition of a filtration step to the protocol. Sampling was restricted to individuals who were not artificially mummified, ensuring that some bone was exposed and available for analysis. Only adults were sampled in this study to ensure all individuals were completely weaned and the isotopic signatures represent an adult diet. This work on adult diet, however, will form an important baseline for future work looking at infant diet and weaning patterns.

5. Methods

Skeletal sex estimation was undertaken by the authors [CK and VS] using the standards in Buikstra and Ubelaker (1994). Individuals were assigned to age groups (Young, Middle, Old) using standard scoring of the pubic symphysis (Brooks and Suchey, 1990) and auricular surface (Lovejoy et al., 1985), with epiphyseal fusion taken into account in the case of young adults (Buikstra and Ubelaker, 1994).

Samples of bone weighing between 100 and 200 mg were taken using a diamond cutting wheel. These were removed from already fragmented and disarticulated ribs to avoid unnecessary damage to any skeletal elements or mummified material. Adhering particulates were removed through surface abrasion with a diamond burr. Prior to collagen extraction a defatting procedure equivalent to those used in analysis of modern bone was employed (O'Connell et al., 2001). The samples were rinsed in deionised water then sonicated in test tubes containing a 2:1 methanol:chloroform solution for 2 hours, with the solvent changed every 30 minutes. Bone fragments were then prepared following a modified Longin (1971) method i.e. demineralised in 0.5 M HCl, gelatinised in a pH3 HCl solution at 75 °C overnight, centrifuged and decanted to remove particulates, then lyophilised.

Total organic carbon, total nitrogen content and stable isotope analysis of the samples was performed using a Costech Elemental Analyser (ECS 4010) connected to a Thermo Delta V Advantage isotope ratio mass spectrometer. Carbon isotope ratios were corrected for ^{17}O contribution and reported in standard delta (δ) notation in per mil (‰) relative to Vienna Pee Dee Belemnite (VPDB). Isotopic accuracy was monitored through routine analyses of in-house standards, which were stringently calibrated against international standards (e.g., USGS 40, USGS 24, IAEA 600, IAEA N1, IAEA N2): this provided a linear range for calibration in $\delta^{13}\text{C}$ between -46.7‰ and $+2.9\text{‰}$ and in $\delta^{15}\text{N}$ between -4.5‰ and $+20.4\text{‰}$. Analytical uncertainty in carbon and nitrogen isotope analysis was typically $\pm 0.1\text{‰}$ for replicate analyses of the international standards and typically $<0.2\text{‰}$ on replicate sample analysis (see S1 Table). Total organic carbon and nitrogen data were obtained as part of the isotopic analysis using an internal standard (Glutamic Acid, 40.82% C, 9.52% N).

The statistical software R (R core team, 2013) was used to script and visualise our data. In an effort to increase the reproducibility of research (Marwick, 2017), csv files and R scripts used in this analysis have been made available via a GitHub repository (https://github.com/DrCharlieKing/Atacama_Bulk). Values from modern terrestrial flora and fauna data were corrected for the Suess effect to pre-industrial levels (Long et al., 2005) using data from Francey et al. (1999). There is also a systematic offset between human collagen values and those of the diet consumed. When presenting human data with foodweb data this dietary offset is corrected for by shifting human values to those of their diet, to allow easier visualisation of possible dietary components. There is debate over the exact magnitude of this offset (O'Connell et al., 2012), but here we use controlled feeding data and consider the $\delta^{13}\text{C}$ diet-collagen offset as $4.8 \pm 0.5\text{‰}$, and $\delta^{15}\text{N}$ diet-collagen offset as $5.5 \pm 0.5\text{‰}$ (Fernandes et al., 2012; Froehle et al., 2010; Huelsemann et al., 2009).

In the Atacama Desert there are a number of resources that overlap substantially in terms of isotopic signatures, for example less negative $\delta^{13}\text{C}$ values may represent C₄ crop or marine resource consumption. Although analysis of dental enamel carbonate ($\delta^{13}\text{C}_{\text{apatite}}$) could potentially resolve these interpretive issues, sampling restrictions prevented this being undertaken as a part of this study. Instead, quantitative dietary reconstruction was undertaken using a Bayesian mixing model – Food Reconstruction Using Isotopic Transferred Signals (FRUITS) (Fernandes et al., 2014), in order to account to some extent for dietary routing and offsets.

FRUITS allows the estimation of the contribution of different food sources to the diet using measured isotopic values and food source data (Fernandes et al., 2014). It takes into account dietary routing (i.e. the preferential use of certain dietary components in specific tissues) and isotopic offsets, and allows the incorporation of *a priori* assumptions. In this study we used a weighted and concentration-dependant model, considering the input of both the

protein and energy components of the diet in the consumer isotopic values. The main FRUITS model used is given as Supplementary File 1. Mean values for each of the phases were used to model group diet, although we also present results from individuals to demonstrate the heterogeneity in diet within single sites. We considered the input of four dietary sources: C₃ plants, C₄ plants, terrestrial meat and marine meat (including shellfish, fish and marine mammals). Food source values were derived from previously conducted isotopic work in the Northern Atacama (Cadwallader et al., 2012; DeNiro and Hastorf, 1985; Szpak et al., 2012; Tieszen and Chapman, 1992) that is summarised in Andrade et al. (2015). Weighting of the model was based upon previous work detailing average nutrient contribution of the different food groups to human diet (Fernandes et al., 2015), with plant cereals protein: $10 \pm 2.5\text{ wtC \%}$; carbs/lipids: $90 \pm 2.5\text{ wtC \%}$, terrestrial meat sources: $30 \pm 2.5\text{ wtC \%}$; carbs/lipids: $70 \pm 2.5\text{ wtC \%}$ and marine foods: $35 \pm 5\text{ wtC \%}$; carbs/lipids: $65 \pm 5\text{ wtC \%}$. We also follow Fernandes et al. (2015, 2014), in using the *a priori* assumption that overall dietary protein intake will involve protein carbon contribution of between 5 and 45% (Otten et al., 2006). In addition, the Archaic Period reconstructions included the prior that C₄ resources form the least important component of diet, as there are no endemic C₄ resources in the Arica region, and therefore it is not until the Formative Period that we would expect these to become a possible important component of diet. Isotopic offsets were defined using the same controlled-feeding experiment data as used to plot our data ($\delta^{13}\text{C}$ diet-collagen offset = $4.8 \pm 0.5\text{‰}$, $\delta^{15}\text{N}$ diet-collagen offset = $5.5 \pm 0.5\text{‰}$) (Fernandes et al., 2012; Froehle et al., 2010; Huelsemann et al., 2009).

To test the robustness of dietary reconstructions alternative models using different dietary information and parameters were also run, and their results compared to our primary model. The first of the alternative dietary scenarios involved the inclusion of fertilised C₄ plant data (see Supplementary File 2). The fertilisation of maize with seabird guano was a common agricultural practice during the Late Intermediate Period (Julien, 1985), and results in elevated $\delta^{15}\text{N}$ values, such that fertilised maize overlaps isotopically with high trophic level marine resources (i.e. marine carnivores such as sealions). In our model we have considered guano fertilisation a




Fig. 2. Isotopic baseline data plotted alongside average human dietary values for each phase (with 95% confidence ellipses). Human isotopic values have been corrected for the diet-tissue offset (as described in-text). Datapoints represented by squares (rather than circles) are individuals representing the 'extremes' of dietary variation in each phase. These individuals have associated individual FRUITS dietary reconstructions (Fig. 4).




Fig. 3. FRUITS model output for each archaeological phase. Box and whisker plots (left) represent credibility intervals, with boxes representing a 68% credible interval, and whiskers a 95% credible interval. Horizontal lines represent the mean and median (dashed and continuous lines respectively). Probability distributions (right hand figures, y-axes) are given for each of the contribution estimates (x-axes).

possibility from the Middle Horizon onwards, as the earliest archaeological evidence for potential deliberate use of guano is from this period (Kelley et al., 1991; Muñoz and Focacci, 1985). The inclusion of this data in our model allows us to quantify the potential confounding effects of fertilization on our data. In addition, three other model types were run: 1) removing *a priori* assumptions (Supplementary File 3); 2) with offsets altered by $\pm 1\%$ (Supplementary Files 4 and 5); and 3) with food values altered by $\pm 1.5\%$ (Supplementary Files 6–9). Dietary reconstructions are considered robust if these changing parameters did not result in significant changes to dietary contribution estimates.

6. Results

The results of carbon and nitrogen isotopic analysis are given in Fig. 2, and are reported in full in Table S1.

6.1. FRUITS dietary estimates

Fig. 3 presents the dietary reconstructions generated by our main FRUITS model for each for each archaeological period. The estimated contributions of each food source, and uncertainties associated with these estimates are also given in Table 2.

As Fig. 2 and high standard deviations in Table 2 show, there are relatively high levels of dietary variability in each of the phases. We

Table 2

% Total caloric contribution to diet of each food source as reconstructed by FRUITS analysis. Errors given are 1SD.

Phase	C ₃ input	C ₄ input	Terrestrial meat input	Marine input
Archaic	32 \pm 10	10 \pm 5	17 \pm 7	41 \pm 6
Formative	31 \pm 21	24 \pm 16	30 \pm 18	15 \pm 10
Middle Horizon	20 \pm 14	46 \pm 13	13 \pm 9	21 \pm 13
Late-Intermediate	24 \pm 17	38 \pm 16	20 \pm 14	18 \pm 12
Late	16 \pm 11	39 \pm 12	9 \pm 7	36 \pm 9

acknowledge that the individuals grouped in archaeological phases may not be contemporaneous and therefore directly comparable due to the long occupation period of some sites. To fully describe the variation present, even within one site, we have generated FRUITS estimates of dietary contributions for individuals representing the extremes of diet in each phase (individuals marked as squares on Fig. 2). The Archaic Period is excluded from this individual analysis due to the small sample size and relative homogeneity of diet in this phase. These individual results are presented in Fig. 4 and Table 3.

6.2. Robustness and accuracy of dietary estimates

The dietary inputs calculated by the alternative models are given in supplementary tables S2–4. Each of the modelled dietary




Fig. 4. FRUITS model output for each archaeological phase. Boxes represent 68% credible intervals, and whiskers 95% credible intervals. Horizontal lines represent the mean and median (dashed and continuous lines respectively).

Table 3

% Total caloric contribution to diet of each food source for selected individuals, as reconstructed by FRUITS analysis. Errors given are 1SD.

Time period	Individual	C ₃ input	C ₄ input	Terrestrial meat input	Marine input
Formative	Az14 T10	9 ± 7	16 ± 9	7 ± 6	68 ± 9
	Az14 T76b	83 ± 6	3 ± 2	6 ± 5	8 ± 3
Middle	Az6 T19	6 ± 5	29 ± 13	3 ± 2	62 ± 13
	Az141 T33	37 ± 17	48 ± 12	12 ± 10	3 ± 2
Late Intermediate	Az8 T9	7 ± 6	51 ± 7	4 ± 3	38 ± 7
	Az11 T16	63 ± 16	9 ± 6	25 ± 14	3 ± 2
Late	Cam9 T59	8 ± 6	23 ± 10	3 ± 2	66 ± 11
	Cam8 TUW1	10 ± 8	46 ± 8	7 ± 4	37 ± 8

scenarios have similar probability distributions associated with different food sources (% input), although the actual percentages vary with the changing model parameters. Each dietary scenario has associated uncertainties ranging between 2 and 20%, with the majority of uncertainties being under 15%. These uncertainties are not large enough to change the overall interpretation of which resources are contributing the most to diet in each of our phases.

In most of the modelled dietary scenarios, despite the reconstructed % contributions to diet changing, the overall reconstruction of which resources contribute more or less to diet remains the same. For example, the Archaic estimates always show that marine food makes up the majority of the diet, with C₄ resources comprising only a small portion. In the Formative period it is terrestrial resources which make up the bulk of the diet, in particular C₃ plants. In the later phases C₄ plants form the foundation of the diet, although in the coastal Late Period sites marine resources form an equally important portion. Interestingly the inclusion of guano fertilised maize data (i.e. C₄ crops with significantly higher $\delta^{15}\text{N}$) from the Middle Period onwards does not significantly affect model outcomes (S4 table). Estimated dietary contributions in the guano-fertilised model differ by 2–6% from the main model presented in this paper.

The removal of prior assumptions from the model does significantly affect percent contribution estimates in the Archaic, with a

change to estimated C₄ contribution of 18%. Aside from large changes to the Archaic Period estimates when priors are removed, the majority of other models run change the percent contribution estimates by no more than 6%. This extreme change in the Archaic is due to the removal of the prior assumption that C₄ resources will comprise the smallest contribution to diet, due to lack of endemic C₄ plants. Other phases did not include this prior due to the presence of C₄ resources introduced from the highlands.

7. Discussion

The results of analysis using FRUITS clearly support the hypothesis that marine resources remained an important dietary component throughout the prehistory of the Arica region. In addition, they highlight that proximity to the coast, perhaps unsurprisingly, has the most significant impact on the level to which marine resources are consumed, though they are consumed in all sites regardless of time period or cultural affiliation. The arrival of maize and increasing importance of this resource from the Middle Period onwards is also visible isotopically. Here we discuss the persistence of marine resource use through time, the retention of broad-spectrum subsistence strategies and geographic constraints on resource use, and the possible influence of maize-reliant polities on subsistence in the region.

7.1. Continued consumption of marine resources despite the arrival of agricultural resources

The results of FRUITS analysis clearly indicate the continued consumption of marine resources throughout prehistory in the Arica region. In sites closest to the coast i.e. those from the Archaic and Late periods, marine resources provide around 40% of caloric input, although some individuals within those sites appear to have had a marine input closer to 60%. Most archaeological models assume the arrival of agriculture in the Formative Period involved an increase in terrestrial resource consumption, and this is borne out by our data which suggests that terrestrial resources do provide more caloric input during these periods. Marine resources, however, continue to comprise between 15 and 20% of the diet. This aligns well with isotopic research in other parts of the Atacama Desert which has shown the persistence of marine resource use in both coastal and inland sites during the Formative Period (Andrade et al., 2015; Pestle et al., 2015a; Santana-Sagredo et al., 2015).

In addition, within each agricultural archaeological period analysed in this study there remains variation in marine resource consumption, with some individuals obtaining up to 68% of their calories from the ocean (e.g. Az14 T10 in the Formative Period). This is a higher contribution even than the most extreme values from coastal sites in this study.

7.2. The importance of the environment – coastal vs. inland sites

In this sample over all periods and locations we see that resource decision-making is clearly affected by geographic availability, particularly relating to proximity to the coast. In our sample the sites located on the coast (i.e. the Archaic Morro sites and Late Period Camarones sites), unsurprisingly, consumed the most marine resources. There is no statistically significant difference in marine resource consumption between the Archaic and Late Period sites, but there are statistically significant differences between all inland (Formative, Middle and Late-Intermediate) sites and coastal (Archaic and Late) sites. This reflects the persistence of a marine tradition, which is present even today in the modern cities of the coastal desert.

Inland sites, however, have approximately 20% less caloric input from marine food sources than those at the coast. This is not necessarily as expected, as all sites analysed are within a day's walk from the coast, and marine resource gathering requires less energy input than rearing livestock or cultivating crops. There is, in addition, no correlation between absolute distance from the coast and marine input into the diet. Formative sites included in this study are the closest valley sites to the coast, yet display the least marine resource use. Middle Period sites analysed here are 10–15 km further from the coast, but use the most marine resources of all of the agricultural periods. Instead we consider it more likely that the inland sites, in general used less marine resources but the specific amounts used will have been dependent upon climatic conditions affecting the fisheries. Prevailing archaeological models, for example, predict that the initial move inland during the Formative Period was precipitated to changes to El Niño intensity which dramatically depressed the fisheries (de Bryson et al., 2001; Muñoz and Chacama, 2012). It is perhaps unsurprising then that we see the least use of marine resources in these earliest agricultural sites. Climatic oscillations during site occupation may also be responsible for the high levels of 'within site/phase' variation in isotopic results we see. Small-scale variations in El Niño cycles are likely to have caused differences in the availability of resources throughout prehistory. Individuals within the same site may have experienced quite different resource pressures, even generation to generation, if water availability changed or marine upwelling was affected. Thus

the variation in marine resource use may not have been personal choice or culturally-mediated, but instead environmentally-dictated.

7.3. The importance of maize and links to external polities

It has been generally assumed that maize played a more important role in subsistence from the Middle Period onwards, due to the presence of maize-reliant polities such as the Tiwanaku (Middle Period) and the Inka state (Late Period) (Goldstein, 2003). This does appear to be the case in the study sample, as C₄ contribution to the average diet increases dramatically between the Formative (24%) and Middle Horizon (46%), with C₄ contribution to diet remaining high throughout the subsequent archaeological phases. Even in coastal Late Period sites, where marine resources are likely to have been much easier to access, C₄ consumption provides a reasonable proportion of total caloric input (39%).

However, it should be noted that maize consumption in the Arica region never reaches the calculated levels of other areas annexed by or in contact with these same polities. For example, in the Ayacucho Valley of Peru, dietary isotope analysis has suggest that maize provided around 70% of dietary protein (Finucane, 2009). It is possible that the geographic location of the Azapa Valley, away from major centres of Tiwanaku and Inka control, meant that there was less cultural importance placed on maize consumption in this area (e.g. King et al., In press).

7.4. Linking isotopic and archaeological evidence for subsistence strategies

Overall the FRUITS reconstructions align well with archaeological evidence for subsistence. The dietary reconstructions presented here, however, also go some way to resolving archaeological questions about the importance of different resource types in each site. For example, archaeozoological and material culture evidence for marine resource consumption is present in all inland sites analysed (Focacci, 1990; Santoro, 1980a; Valenzuela et al., 2015), but it is difficult to know how important these resources were. Our results show that not only were marine resources present they also comprised an important part of the diet, although the input into individuals' diet was variable.

The variety of crop species and subsistence tools found even at inland agricultural sites may be used to infer that life in the Azapa Valley was never focussed on a single crop or subsistence strategy (Muñoz, 1987; Valenzuela et al., 2015). Our isotopic results support this interpretation, showing that the average diet during the Formative Period involved roughly equal proportions of C₃ and C₄ plants, terrestrial meat and marine resources. Even dietary outliers in this phase, who relied more upon one resource type than the others, still seem to have been consuming small proportions of other resource types. There is archaeological evidence for an increase in maize consumption in the Middle Horizon and Late-Intermediate Periods in that maize remains and material culture associated with *chicha* consumption become ubiquitous in sites (Muñoz and Focacci, 1985; Muñoz and Zalaquett, 2015). Our isotopic results corroborate the idea that maize becomes more important, as in both the Middle Horizon and Late-Intermediate periods C₄ resources are the most important dietary component. They are not, however, solely relied upon, and indeed there are some individuals for whom they form a much lesser proportion of the diet.

Archaeological research has previously questioned the importance of maize at Late Period coastal sites despite Inka cultural involvement. At sites such as Camarones maize cannot be grown locally and archaeobotanical evidence indicates that *chicha*

(traditionally a maize beer), appears to have been made primarily from C₃ crops rather than maize (Arriaza et al., 2016, 2017). Our research indicates that maize is likely to have formed an important component of the diet despite its lack of inclusion in *chicha*. In fact, the C₃ crops from which *chicha* was apparently made do not generally contribute much to diet. It is possible that the use of these crops in *chicha* was more an expression of local identity while under Inka rule, than an indication of lack of maize availability.

7.5. Variation in isotopic results, and diet in general, from each phase

Isotopic results from the inland Formative, Middle and Late-Intermediate Periods are extremely variable. Perhaps the most extreme example of this variation comes from Formative Period Az14, where individuals at one extreme appear to be consuming almost exclusively C₃ plants, while others seem to have a diet dominated by marine protein. It is tempting to interpret this isotopic variation within archaeological phases or single sites as evidence for heterogeneous diet between contemporary individuals. Unfortunately, however, the ambiguity of site dating in the Arica region means that individuals within the same site are not necessarily contemporaneous. Instead, this variation may indicate generational changes to diet, perhaps reflecting cyclical resource availability due to El Niño oscillations, or cultural change resulting in the changing popularity of some resources. Future work involving the tightening of site chronologies may reveal systematic temporal changes to diet within phases, that are currently not interpretable.

7.6. Differences between FRUITS dietary reconstructions and previous modelling in the region

One of the major differences between the dietary reconstructions generated in this study and those attempted previously (Aufderheide et al., 1993, 1994) is the percent input of marine resources estimated. The linear mixing models in these previous studies have predicted up to 80% of dietary protein was derived from marine resource consumption. We anticipate that this difference is partly due to our use of Bayesian rather than simple linear mixing modelling, but is also because our FRUITS model used metabolic experimental data to define the acceptable intake of dietary protein. As a result of this our dietary reconstructions seem to align better with archaeological evidence for subsistence than earlier estimates. For example, while marine resources undoubtedly formed the most significant portion of Archaic diet there is evidence from Archaic coprolites of C₃ plants and terrestrial meat (Reinhard et al., 2011). Our dietary reconstructions support this idea of terrestrial resource use, and reduce estimates of percent contribution of marine resources to what we consider more reasonable levels.

8. Conclusion

The isotopic results presented here paint a picture of deliberate retention of dietary diversity in the peoples of the Atacama Desert. The marine resource use traditions begun by the Archaic *Chinchorro* peoples persist throughout prehistory and into the modern day, with use of the rich marine environment buffering the desert populations against terrestrial resource insecurities. While the arrival of maize-reliant polities does seem to have had an impact on subsistence in later archaeological phases, agricultural crops are never fully relied upon. Instead, geographic constraints, such as proximity to the ocean, seem to have had a greater impact on overall diet. Our results also highlight dietary diversity both within

and between archaeological phases, potentially relating to changes to resource availability and El Niño cycles.

Author contribution

All coauthors made substantial contributions to, and have approved the submission of this manuscript. Lead author CLK was responsible for study design, with advice from SEH and ARM. CLK collected samples, with aid from BTA and VGS, who also provided contextual information and insight into the regional archaeology. CLK undertook sample prep, samples were run by DRG, who wrote the analytical methods section of the manuscript. The bulk of the manuscript was written by CLK, with editorial input and suggested rewrites from all other co-authors.

Acknowledgements

This work was funded by a Rutherford Postdoctoral Fellowship (Royal Society of New Zealand) awarded to CLK, and a Marsden Grant (UOO1413) awarded to SEH. Initial work was funded by a University of Otago Research Grant awarded to SEH. Both BTG and VGS are funded by El Fondo Nacional de Desarrollo Científico y Tecnológico (Fondecyt) grants. Bone and dental samples taken from this study were authorized by the Director of the San Miguel de Azapa Archaeological Museum of the University of Tarapacá, Arica, Chile; And the Council of National Monuments of Chile authorized the scientific publication of the results obtained (Ord. No. 04152). Susana Monsalve and Anita Flores provided help within the collections of the Museo San Miguel de Azapa. At Durham University Beth Upex and Steve Robertson (Archaeology Department) provided help during analysis.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.quascirev.2017.12.009>.

References

- Allison, M., Focacci, G., Arriaza, B.T., Standen, V.G., Rivera, M., Lowenstein, J., 1984. Chinchorro, momias de preparación complicada: métodos de momificación. *Chungará* 13, 155–173.
- Ambrose, S.H., Norr, L., 1993. Experimental evidence for the relationship of the carbon isotope ratios of whole diet and dietary protein to those of bone collagen and carbonate. In: Lambert, J.B., Grupe, G. (Eds.), *Prehistoric Human Bone: Archaeology at the Molecular Level*. Springer, Heidelberg, pp. 1–37.
- Amundson, R., Austin, A., Schuur, E., Yoo, K., Matzek, V., Kendall, C., Uebersax, A., Brenner, D., Baisden, W., 2003. Global patterns of the isotopic composition of soil and plant nitrogen. *Glob. Biogeochem. Cycles* 17, 1031.
- Andrade, P., Fernandes, R., Codjambassis, K., Urrea, J., Olguín, L., Rebolledo, S., Lira, F., Aravena, C., Berrios, M., 2015. Subsistence continuity linked to consumption of marine protein in the Formative period in the interfluvial coast of northern Chile: reassessing contacts with agropastoral groups from highlands. *Radio-carbon* 57, 679–688.
- Apata, M., Arriaza, B., Llop, E., Moraga, M., 2017. Human adaptation to arsenic in Andean populations of the Atacama Desert. *Am. J. Phys. Anthropol.* 163, 192–199.
- Arriaza, B.T., 1995. Chinchorro bioarchaeology: chronology and mummy seriation. *Lat. Amer. Antiq.* 6, 35–55.
- Arriaza, B.T., Amarasiqwariwdena, D., Cornejo, L., Standen, V., Byrne, S., Bartkus, L., Bandak, B., 2010. Exploring chronic arsenic poisoning in pre-Columbian Chilean mummies. *J. Archaeol. Sci.* 37, 1274–1278.
- Arriaza, B.T., Huamán, L., Villanueva, F., Tornero, R., Standen, V., Aravena, N., 2017. Estudio del cálculo dental en poblaciones arqueológicas del extremo norte de Chile. *Estud. Atacameños*. Early view.
- Arriaza, B.T., Ogalde, J., Chacama, J., Standen, V., Huaman, L., Villanueva, F., Aravena, N., Méndez-Quiros, P., Tapia, P., 2016. Microscopic analysis of botanical residues from Cerro Esmeralda burial in Northern Chile: state and death ritual implications. *Interciencia* 41, 844–850.
- Arriaza, B.T., Standen, V.G., Cassman, V., Santoro, C.M., 2008. Chinchorro culture: pioneers of the coast of the Atacama Desert. In: Silverman, H., Isbell, W. (Eds.), *The Handbook of South American Archaeology*. Springer, New York, pp. 45–58.
- Aufderheide, A.C., Kelley, M.A., Rivera, M., Gray, L., Tieszen, L.L., Iversen, E.,

- Krouse, H.R., Carevic, A., 1994. Contributions of chemical dietary reconstruction to the assessment of adaptation by ancient highland immigrants (Alto Ramírez) to coastal conditions at Pisagua, North Chile. *J. Archaeol. Sci.* 21, 515–524.
- Aufderheide, A.C., Muñoz, I., Arriaza, B., 1993. Seven Chinchorro mummies and the prehistory of northern Chile. *Am. J. Phys. Anthropol.* 91, 189–201.
- Bauer, B.S., 1996. Legitimization of the state in Inca myth and ritual. *Am. Anthropol.* 98, 327–337.
- Belmonte, E., 1998. Estudio comparativo de especies vegetales de contexto arqueológico (Museo Arqueológico San Miguel de Azapa). In: Informe Final de Investigación Proyecto N°. Universidad de Tarapacá, Arica, pp. 3740–3796.
- Beresford-Jones, D., Pullen, A.G., Whaley, O.Q., Moat, J., Chauca, G., Cadwallader, L., Arce, S., Orellana, A., Alarcón, C., Gorriti, M., 2015. Re-evaluating the resource potential of lomas fog oasis environments for Preceramic hunter-gatherers under past ENSO modes on the south coast of Peru. *Quat. Sci. Rev.* 129, 196–215.
- Bonilla, M.-Z., Drucker, D., Richardin, P., Silva-Pinto, V., Sepúlveda, M., Bocherens, H., 2016. Marine food consumption in coastal northern Chilean (Atacama Desert) populations during the Formative period: implications of isotopic evidence (C, N, S) for the Neolithic process in south central Andes. *J. Arch. Sci. Rep.* 6, 768–776.
- Brooks, S., Suchey, J.M., 1990. Skeletal age determination based on the os pubis: a comparison of the Acsádi-Nemeskéri and Suchey-Brooks methods. *Hum. Evol.* 5, 227–238.
- Buikstra, J.E., Ubelaker, D.H., 1994. Standards for Data Collection from Human Skeletal Remains. Arkansas Archeological Survey, Lafayette, AK.
- Cadwallader, L., Beresford-Jones, D.G., Whaley, O.Q., O'Connell, T.C., 2012. The signs of maize? A reconsideration of what $\delta^{13}\text{C}$ values say about palaeodiet in the Andean region. *Hum. Ecol.* 40, 487–509.
- Chisholm, B.S., Nelson, D.E., Schwarzc, H.P., 1982. Stable-carbon isotope ratios as a measure of marine versus terrestrial protein in ancient diets. *Science* 216, 1131–1132.
- Cuellar, A.M., 2013. The archaeology of food and social inequality in the Andes. *J. Archaeol. Res.* 21, 123–174.
- de Bryson, R., Luz, M., Bryson, R.U., Bryson, R.A., 2001. Paleoclimatic and material cultural perspective on the Formative Period of northern Chile. *Chungará* 33, 5–12.
- DeNiro, M., 1985. Postmortem preservation and alteration of *in vivo* bone collagen isotope ratios in relation to palaeodietary reconstruction. *Nature* 317, 806–809.
- DeNiro, M.J., Epstein, S., 1981. Influence of diet on the distribution of nitrogen isotopes in animals. *Geochim. Cosmochim. Acta* 45, 341–351.
- DeNiro, M.J., Hastorf, C.A., 1985. Alteration of $\delta^{15}\text{N}/\delta^{14}\text{N}$ and $\delta^{13}\text{C}/\delta^{12}\text{C}$ ratios of plant matter during the initial stages of diagenesis: studies utilizing archaeological specimens from Peru. *Geochim. Cosmochim. Acta* 49, 97–115.
- Díaz, F.P., Frugone, M., Gutiérrez, R.A., Latorre, C., 2016. Nitrogen cycling in an extreme hyperarid environment inferred from $\delta^{15}\text{N}$ analyses of plants, soils and herbivore diet. *Sci. Rep.* 6, 1–11.
- Ehleringer, J.R., Mooney, H.A., Rundel, P.W., Evans, R.D., Palma, B., Delatorre, J., 1992. Lack of nitrogen cycling in the Atacama Desert. *Nature* 359, 316–318.
- Erices, S., 1975. Evidencias de vegetales en tres cementerios prehispánicos, Arica - Chile. *Chungará* 5, 65–71.
- Espouyes, O., Uribe, M., Roman, A., Deza, A., 1995. Nuevos fechados por termoluminiscencia para la cerámica del período medio del Valle de Azapa (primera parte). *Hombre y Desierto* 9, 31–54.
- Fernandes, R., Grootes, P., Nadeau, M.J., Nehlich, O., 2015. Quantitative diet reconstruction of a Neolithic population using a Bayesian mixing model (FRUITS): the case study of Ostorf (Germany). *Am. J. Phys. Anthropol.* 158, 325–340.
- Fernandes, R., Millard, A.R., Brabec, M., Nadeau, M.-J., Grootes, P., 2014. Food reconstruction using isotopic transferred signals (FRUITS): a Bayesian model for diet reconstruction. *PLoS One* 9, e87436.
- Fernandes, R., Nadeau, M.-J., Grootes, P.M., 2012. Macronutrient-based model for dietary carbon routing in bone collagen and biapatite. *Archaeol. Anthropol. Sci.* 4, 291–301.
- Figueroa, L., Barton, S., Schull, W., Razmilic, B., Zumaeta, O., Young, A., Kamiya, Y., Hoskins, J., Ilgren, E., 2012. Environmental lithium exposure in the North of Chile—I. Natural water sources. *Biol. Trace Elem. Res.* 149, 280–290.
- Finucane, B.C., 2009. Maize and sociopolitical complexity in the Ayacucho Valley, Peru. *Curr. Anthropol.* 50, 535–545.
- Focacci, G., 1990. Excavaciones arqueológicas en el cementerio Az-6 Valle de Azapa. *Chungará* 69–123, 24/25.
- Francey, R., Allison, C., Etheridge, D., Trudinger, C., Enting, I., Leuenberger, M., Langenfelds, R., Michel, E., Steele, L., 1999. A 1000-year high precision record of $\delta^{13}\text{C}$ in atmospheric CO_2 . *Tellus B* 51, 170–193.
- Froehle, A.W., Kellner, C.M., Schoeninger, M.J., 2010. Effect of diet and protein source on carbon stable isotope ratios in collagen: follow up to Warinner and Tuross (2009). *J. Archaeol. Sci.* 37, 2662–2670.
- Fry, B., 2006. Stable Isotope Ecology. Springer, New York.
- Gayo, E., Latorre, C., Santoro, C., Maldonado, A., De Pol-Holz, R., 2012. Hydroclimate variability in the low-elevation Atacama Desert over the last 2500 yr. *Clim. Past* 8, 287–306.
- Goldstein, P., 2003. From stew-eaters to maize-drinkers. In: Bray, T. (Ed.), *The Archaeology and Politics of Food and Feasting in Early States and Empires*. Plenum Publishers, New York, pp. 143–172.
- Gröcke, D.R., Bocherens, H., Mariotti, A., 1997. Annual rainfall and nitrogen-isotope correlation in macropod collagen: application as a palaeoprecipitation indicator. *Earth. Planet. Sci. Lett.* 153, 279–285.
- Grosjean, M., Santoro, C.M., Thompson, L., Nunez, L., Standen, V.C., 2007. Mid-holocene climate and culture change in the south central Andes. In: Anderson, D.G., Maasch, K.A., Sandweiss, D.H. (Eds.), *Climate Change and Cultural Dynamics: a Global Perspective on Mid-Holocene Transitions*. Academic Press, London, pp. 51–115.
- Hartman, G., 2011. Are elevated $\delta^{15}\text{N}$ values in herbivores in hot and arid environments caused by diet or animal physiology? *Funct. Ecol.* 25, 122–131.
- Holden, T., 1994. Dietary evidence from the intestinal contents of ancient humans with particular reference to desiccated remains from northern Chile. In: Hether, J. (Ed.), *Tropical Archaeobotany. Applications and New Developments*. Routledge, London, pp. 65–85.
- Hubbe, M., Torres-Rouff, C., Neves, W.A., King, L.M., Da-Gloria, P., Costa, M.A., 2012. Dental health in northern Chile's Atacama oases: evaluating the Middle Horizon (AD 500–1000) impact on local diet. *Am. J. Phys. Anthropol.* 148, 62–72.
- Huelsmann, F., Flenker, U., Koehler, K., Schaenzer, W., 2009. Effect of a controlled dietary change on carbon and nitrogen stable isotope ratios of human hair. *Rapid Commun. Mass Spectrom.* 23, 2448–2454.
- Julien, C.J., 1985. Guano and resource control in sixteenth-century Arequipa. In: Masuda, C., Shimada, I., Morris, C. (Eds.), *Andean Ecology and Civilization*. University of Tokyo Press, Tokyo, pp. 185–231.
- Kelley, M., Levesque, D., Weidle, E., 1991. Contrasting patterns of dental disease in five early northern Chilean groups. In: Kelley, M., Larsen, C. (Eds.), *Advances in Dental Anthropology*. Wiley-Liss, New York, pp. 203–213.
- King, C.L., Halcrow, S.E., Millard, A.R., Grocke, D., Standen, V., Muñoz Ovalle, I., Arriaza, B.T., In press. El consumo de maíz en la prehistoria de Arica. *Chungará*.
- Knudson, K.J., Aufderheide, A.E., Buikstra, J.E., 2007. Seasonality and paleodiet in the Chiribaya polity of southern Peru. *J. Archaeol. Sci.* 34, 451–462.
- Kolata, A.L., 1991. The technology and organization of agricultural production in the Tiwanaku state. *Lat. Amer. Antiq.* 2, 99–125.
- Korpisari, A., Oinonen, M., Chacama, J., 2014. A reevaluation of the absolute chronology of Cabuza and related ceramic styles of the Azapa valley, northern Chile. *Lat. Amer. Antiq.* 25, 409–426.
- Llagostera, A., 1996. San Pedro de Atacama: nodo de complementariedad reticular. *Estudios y Debates Regionales Andinos* 96, 17–42.
- Llagostera, A., Costa, M.A., 1999. Patrones de asentamiento en la época agroalfarera de San Pedro de Atacama (norte de Chile). *Estud. Atacameños* 17, 175–206.
- Long, E.S., Sweitzer, R.A., Diefenbach, D.R., Ben-David, M., 2005. Controlling for anthropogenically induced atmospheric variation in stable carbon isotope studies. *Oecologia* 146, 148–156.
- Longin, R., 1971. New method of collagen extraction for radiocarbon dating. *Nature* 230, 241–242.
- Lovejoy, C.O., Meindl, R.S., Pryzbeck, T.R., Mensforth, R.P., 1985. Chronological metamorphosis of the auricular surface of the ilium: a new method for the determination of adult skeletal age at death. *Am. J. Phys. Anthropol.* 68, 15–28.
- Marwick, B., 2017. Computational reproducibility in archaeological research: basic principles and a case study of their implementation. *J. Archaeol. Method Th* 24, 424–450.
- Mcrostie, V.B., Gayo, E.M., Santoro, C.M., De Pol-Holz, R., Latorre, C., 2017. The pre-Columbian introduction and dispersal of Algarrobo (*Prosopis*, section *Algarobia*) in the Atacama Desert of northern Chile. *PLoS One* 12, e0181759.
- Minagawa, M., Wada, E., 1984. Stepwise enrichment of $\delta^{15}\text{N}$ along food chains: further evidence and the relation between $\delta^{15}\text{N}$ and animal age. *Geochim. Cosmochim. Acta* 48, 1135–1140.
- Moreno, A., Santoro, C.M., Latorre, C., 2009. Climate change and human occupation in the northernmost Chilean Altiplano over the last ca. 11 500 cal. a BP. *J. Quat. Sci.* 24, 373–382.
- Moseley, M.E., Keefer, D.K., 2008. Deadly deluges in the southern desert: modern and ancient El Niños in the oomre region of southern Peru. In: Sandweiss, D.H., Quilter, J. (Eds.), *El Niño, Catastrophe and Culture Change in Ancient America*. Harvard University Press, Washington DC, pp. 129–144.
- Muñoz, I., 1982. La Capilla No. 4: un asentamiento poblacional tardío en la costa de Arica. *Documentos de trabajo* (No. 2). Instituto de Antropología y Arqueología, Universidad de Tarapacá, Arica, Chile, pp. 95–125.
- Muñoz, I., 1983. El poblamiento Aldeano en el valle de Azapa y su vinculación con Tiwanaku (Arica-Chile): asentamientos Aldeanos en los valles costeros de Arica. *Documentos de Trabajo* (No. 3). Instituto de Antropología y Arqueología, Universidad de Tarapacá, Arica.
- Muñoz, I., 1987. Enteramientos en túmulos en el valle de Azapa: nuevas evidencias para definir la fase Alto Ramírez en el extremo norte de Chile. *Chungará* 19, 93–127.
- Muñoz, I., 1989. Perfil de la organización económico-social en la desembocadura del Río Camarones: períodos Intermedio Tardío e Inca. *Chungará* 22, 85–111.
- Muñoz, I., 1993. Spatial dimensions of complementary resource utilization at Achaca-2 and San Lorenzo, northern Chile. In: Aldenderfer, M. (Ed.), *Domestic Architecture, Ethnicity, and Complementarity in the South-Central Andes*. University of Iowa Press, Iowa City, pp. 94–102.
- Muñoz, I., 1995. Poblamiento humano y relaciones interculturales en el valle de Azapa: nuevos hallazgos en torno al período Formativo y Tiwanaku. *Diálogo Andino* 14, 241–276.
- Muñoz, I., 2004. El período Formativo en los valles del norte de Chile y sur de Perú: nuevas evidencias y comentarios. *Chungará* 36, 213–225.
- Muñoz, I., 2017. Espacios fúnebres, prácticas mortuorias y cronología en el cementerio Azapa-115: aproximaciones en torno a la estructura social de los agricultores prehispánicos del período Medio. *Estud. Atacameños* 54, 5–36.
- Muñoz, I., Chacama, J., 2012. Transformación del paisaje social en Arica, norte de Chile: de pescadores arcaicos a agricultores incipientes. *Estud. Atacameños* 44,

- 123–140.
- Muñoz, I., Focacci, G., 1985. San Lorenzo: testimonio de una comunidad de agricultores y pescadores postiwanaku en el valle de Azapa (Arica - Chile). Chungará 15, 7–30.
- Muñoz, I., Zalaquett, F., 2015. El paisaje en el ordenamiento territorial prehispánico durante el periodo Medio, valle de Azapa, norte de Chile. Revista de Geografía Norte Grande 60, 21–62.
- Murra, J.V., 1980. The Economic Organization of the Inka State. Jai Press, Greenwich.
- Núñez, L., 2007. Vida y Cultura en el oasis de San Pedro de Atacama. Editorial Universitaria, Santiago.
- Núñez, L., Santoro, C.M., 2011. El tránsito arcaico-formativo en la circumpuna y valles occidentales del centro sur Andino: hacia los cambios “Neolíticos”. Chungará 43, 487–530.
- O'Connell, T.C., Hedges, R.E., Healey, M., Simpson, A.H., 2001. Isotopic comparison of hair, nail and bone: modern analyses. *J. Archaeol. Sci.* 28, 1247–1255.
- O'Connell, T.C., Kneale, C.J., Tasevská, N., Kuhnle, G.G., 2012. The diet-body offset in human nitrogen isotopic values: a controlled dietary study. *Am. J. Phys. Anthropol.* 149, 426–434.
- Otten, J.J., Hellwig, J.P., Meyers, L.D., 2006. Dietary Reference Intakes: the Essential Guide to Nutrient Requirements. National Academies Press, Washington DC.
- Pearsall, D.M., 2008. Plant domestication and the shift to agriculture in the Andes. In: Silverman, H., Isbell, W. (Eds.), *The Handbook of South American Archaeology*. Springer, New York, pp. 105–120.
- Pestle, W.J., Torres-Rouff, C., Gallardo, F., Ballester, B., Clarot, A., 2015a. Mobility and exchange among marine hunter-gatherer and agropastoralist communities in the formative period Atacama Desert. *Curr. Anthropol.* 56, 121–133.
- Pestle, W.J., Torres-Rouff, C., Hubbe, M., 2016. Modeling diet in times of change: the case of Quitor, San Pedro de Atacama, Chile. *J. Archaeol. Sci. Rep.* 7, 82–93.
- Pestle, W.J., Torres-Rouff, C., Hubbe, M., Santana, F., Pimentel, G., Gallardo, F., Knudson, K.J., 2015b. Explorando la diversidad dietética en la prehistoria del Desierto de Atacama: un acercamiento a los patrones regionales. Chungará 47, 201–209.
- R core team, 2013. R: a Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria.
- Reinhard, K.J., Le-Roy-Toren, S., Arriaza, B., 2011. Where have all the plant foods gone? The search for refined dietary reconstruction from Chinchorro mummies. *Yearbook of Mummy Studies* 1, 139–151.
- Rivera, M., 1994. Hacia la complejidad social y política: el desarrollo Alto Ramírez del norte de Chile. *Diálogo Andino* 13, 9–38.
- Rivera, M.A., 1975. Una hipótesis sobre movimientos poblacionales altiplánicos y transatlánticos a las costas del norte de Chile. Chungará 5, 7–31.
- Rothhammer, F., Santoro, C.M., 2001. El desarrollo cultural en el Valle de Azapa, extremo norte de Chile y su vinculación con los desplazamientos poblacionales altiplánicos. *Lat. Am. Antiq.* 12, 59–66.
- Sandweiss, D.H., Maasch, K.A., Anderson, D.G., 1999. Transitions in the mid-Holocene. *Science* 283, 499–500.
- Sandweiss, D.H., Solís, R.S., Moseley, M.E., Keefer, D.K., Ortloff, C.R., 2009. Environmental change and economic development in coastal Peru between 5,800 and 3,600 years ago. *Proc. Natl. Acad. Sci.* 106, 1359–1363.
- Santana-Sagredo, F., Uribe, M., Herrera, M.J., Retamal, R., Flores, S., 2015. Dietary practices in ancient populations from northern Chile during the transition to agriculture (Tarapacá region, 1000 BC–AD 900). *Am. J. Phys. Anthropol.* 158, 751–758.
- Santoro, C., Williams, V., Valenzuela, D., Romero, A., Standen, V., 2010. An archaeological perspective on the Inka provincial administration of the south-central Andes. In: Malpass, M.A., Alconini, S. (Eds.), *Distant Provinces in the Inka Empire: toward a Deeper Understanding of Inka Imperialism*. University of Iowa Press, Iowa City, pp. 44–74.
- Santoro, C.M., 1980a. Estratigrafía y secuencia cultural funeraria fases: Azapa, Alto Ramírez y Tiwanaku (Arica-Chile). Chungará 6, 24–45.
- Santoro, C.M., 1980b. Fase Azapa transición del Arcaico al desarrollo agrario inicial en los valles bajos de Arica. Chungará 6, 45–56.
- Santoro, C.M., 2012. Rise and decline of Chinchorro sacred landscapes along the hyperarid coast of the Atacama Desert. Chungará 44, 637–653.
- Santoro, C.M., Arriaza, B.T., Standen, V.G., Marquet, P.A., 2005. People of the coastal Atacama Desert: living between sand dunes and waves of the Pacific Ocean. In: Veth, P., Smith, M., Hiscock, P. (Eds.), *Desert Peoples: Archaeological Perspectives*. Blackwell, Malden, pp. 243–260.
- Santoro, C.M., Capriles, J.M., Gayo, E.M., de Porras, M.E., Maldonado, A., Standen, V.G., Latorre, C., Castro, V., Angelo, D., McRostie, V., 2017. Continuities and discontinuities in the socio-environmental systems of the Atacama Desert during the last 13,000 years. *J. Anthropol. Archaeol.* 46, 28–39.
- Schiappacasse, V., Roman, A., Muñoz, I., Deza, A., Focacci, G., 1991. Cronología por termoluminiscencia de la cerámica del extremo norte de Chile: primera parte. In: *Actas Del XI Congreso Nacional De Arqueología Chilena. Museo Nacional de Historia Natural, Santiago*, pp. 43–60.
- Schoeninger, M.J., DeNiro, M.J., 1984. Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochim. Cosmochim. Acta* 48, 625–639.
- Silva-Pinto, V., Salazar-García, D., Muñoz, I., 2014. Una aproximación a la dieta consumida por las poblaciones constructoras de túmulos. In: Muñoz, I., Fernández Murillo, M.S. (Eds.), *Mil Años de Historia de los Constructores de Túmulos de los Valles Desérticos de Arica: Paisaje, Monumentos y Memoria*. Universidad de Tarapacá, Arica, pp. 89–102.
- Staller, J.E., 2010. *Maize Cobs and Cultures: History of Zea mays*. L. Springer, Heidelberg.
- Standen, V., Arriaza, B., Santoro, C., 2014. Chinchorro mortuary practices on infants: northern Chile Archaic period (BP 7000–3600). In: Thompson, J.L., Alfonso-Durruty, M.P., Crandall, J.J. (Eds.), *Tracing Childhood: Bioarchaeological Investigations of Early Lives in Antiquity*. University of Florida Press, Gainesville, pp. 58–74.
- Standen, V.G., 2003. Bienes funerarios del cementerio Chinchorro Morro 1: descripción, análisis e interpretación. Chungará 35, 175–207.
- Standen, V.G., Santoro, C.M., Arriaza, B., Coleman, D., 2017. Hunting, gathering, and fishing on the coast of the Atacama Desert: Chinchorro population mobility patterns inferred from strontium isotopes. *Geographia (Early View)*.
- Sutter, R.C., 2000. Prehistoric genetic and culture change: a bioarchaeological search for pre-Inka altiplano colonies in the coastal valleys of Moquegua, Peru, and Azapa, Chile. *Lat. Amer. Antiq.* 11, 43–70.
- Sutter, R.C., 2005. A bioarchaeological assessment of prehistoric ethnicity among early Late Intermediate period populations of the Azapa Valley, Chile. In: Reycraft, R. (Ed.), *Us and Them: Archaeology and Ethnicity in the Andes*. Cotsen Institute of Archaeology, Los Angeles, pp. 183–205.
- Swift, J., Cupper, M.L., Greig, A., Westaway, M.C., Carter, C., Santoro, C.M., Wood, R., Jacobsen, G.E., Bertuch, F., 2015. Skeletal arsenic of the pre-Columbian population of Caleta Vitor, northern Chile. *J. Archaeol. Sci.* 58, 31–45.
- Szpak, P., Millaire, J.-F., White, C.D., Longstaffe, F.J., 2012. Influence of seabird guano and camelid dung fertilization on the nitrogen isotopic composition of field-grown maize (*Zea mays*). *J. Archaeol. Sci.* 39, 3721–3740.
- Thiel, M., Macaya, E.C., Acuna, E., Arntz, W.E., Bastias, H., Brokordt, K., Camus, P.A., Castilla, J.C., Castro, L.R., Cortes, M., 2007. The Humboldt Current System of northern and central Chile: oceanographic processes, ecological interactions and socioeconomic feedback. *Oceanogr. Mar. Biol. Rev.* 45, 195–344.
- Tieszen, L.L., Chapman, M., 1992. Carbon and nitrogen isotopic status of the major marine and terrestrial resources in the Atacama desert of northern Chile. In: *Proceedings of the First World Congress on Mummy Studies*, vol. 2, pp. 409–426.
- Torres-Rouff, C., Pestle, W., Gallardo, F., 2012. Eating fish in the driest desert in the world: osteological and biogeochemical analyses of human skeletal remains from the San Salvador cemetery, North Chile. *Lat. Amer. Antiq.* 23, 51–69.
- Torres-Rouff, C., Knudson, K.J., Pestle, W.J., Stovel, E.M., 2015. Tiwanaku influence and social inequality: a bioarchaeological, biogeochemical, and contextual analysis of the Larache cemetery, San Pedro de Atacama, Northern Chile. *Am. J. Phys. Anthropol.* 158, 592–606.
- Tykot, R.H., Burger, R.L., Van Der Merwe, N.J., 2006. The importance of maize in initial period and early horizon Peru. In: Staller, J., Tykot, R., Benz, B. (Eds.), *Histories of Maize: Multidisciplinary Approaches to the Prehistory, Linguistics, Biogeography, Domestication, and Evolution of Maize*. Left Coast Press, Walnut Creek, CA, pp. 187–233.
- Ulloa, L., Standen, V.G., Gavilán, V., 2000. Estudio de una prenda textil asociada al Inca en la costa norte de Chile (Camarones 9): las "mantas" que envuelven los cuerpos. Chungará 32, 259–261.
- Uribe, M., Vidal, E., 2012. Sobre la secuencia cerámica del período Formativo de Tarapacá (900 AC-900 DC): estudios en Pircas, Caserones, Guatacondo y Ramaditas, norte de Chile. Chungará 44, 209–245.
- Valenzuela, D., Santoro, C.M., Capriles, J.M., Quinteros, M.J., Peredo, R., Gayo, E.M., Montt, I., Sepúlveda, M., 2015. Consumption of animals beyond diet in the Atacama Desert, northern Chile (13,000–410BP): comparing rock art motifs and archaeofaunal records. *J. Anthropol. Archaeol.* 40, 250–265.
- Vidal, A., García, M., Méndez-Quiros, P., 2012. Vida sedentaria y oportunismo: dos estrategias de producción agrícola durante el período Formativo en Tarapacá. In: *Actas del XIX Congreso Nacional de Arqueología Chilena*, vol. 1, pp. 183–192.
- Williams, A., Santoro, C.M., Smith, M.A., Latorre, C., 2008. The impact of ENSO in the Atacama Desert and Australian arid zone: exploratory time-series analysis of archaeological records. Chungará 40, 245–259.
- Zaro, G., 2007. Diversity specialists: coastal resource management and historical contingency in the Osmore desert of southern Peru. *Lat. Amer. Antiq.* 18, 161–179.