Published for SISSA by 🖄 Springer

RECEIVED: September 2, 2014 ACCEPTED: October 26, 2014 PUBLISHED: November 10, 2014

Two-loop helicity amplitudes for the production of two off-shell electroweak bosons in quark-antiquark collisions

Fabrizio Caola,^{*a*} Johannes M. Henn,^{*b*} Kirill Melnikov,^{*a*,1} Alexander V. Smirnov^{*c*} and Vladimir A. Smirnov^{*d*}

- ^aInstitute for Theoretical Particle Physics, KIT, Karlsruhe, Germany
- ^bInstitute for Advanced Study,
- Princeton, NJ 08540, U.S.A.
- ^cScientific Research Computing Center, Moscow State University, 119991 Moscow, Russia
- ^dSkobeltsyn Institute of Nuclear Physics of Moscow State University, 119991 Moscow, Russia

E-mail: fabrizio.caola@kit.edu, jmhenn@ias.edu, melnikov@pha.jhu.edu, asmirnov80@gmail.com, smirnov@theory.sinp.msu.ru

ABSTRACT: Knowledge of two-loop QCD amplitudes for processes $q\bar{q}' \rightarrow V_1V_2 \rightarrow (l_1\bar{l}_1)(l_2\bar{l}_2)$ is important for improving the theoretical description of four-lepton production in hadron collisions. In this paper we compute these helicity amplitudes for all intermediate vector bosons, $V_1V_2 = \gamma^*\gamma^*$, W^+W^- , ZZ, $W^{\pm}Z$, $W^{\pm}\gamma^*$, including off-shell effects and decays to leptons.

KEYWORDS: NLO Computations, Hadronic Colliders

ARXIV EPRINT: 1408.6409



¹On leave of absence from Department of Physics and Astronomy, Johns Hopkins University, Baltimore, MD, U.S.A.

Contents

T	Introduction	1
2	The setup of the computation	2
	2.1 $\gamma^* \gamma^*$ production	8
	2.2 ZZ production	9
	2.3 $Z\gamma^*$ production	9
	2.4 $W^+\gamma^*$ and for $W^-\gamma^*$ production	9
	2.5 W^+Z and for W^-Z production	9
	2.6 W^+W^- production	10
3	Calculation of the form factors and the amplitudes	10
4	Conclusions	15

1 Introduction

Production of vector boson pairs at the LHC is an interesting process for a variety of reasons. For definiteness, let us consider the W^+W^- final state. In that case, ATLAS and CMS collaborations have recently observed that the W^+W^- production cross section is about twenty percent higher than existing theoretical predictions [1-3]. This observation prompted speculations about the possibility to explain this excess by physics beyond the Standard Model [4, 5] and, at the same time, strongly emphasized the need to improve predictions for W^+W^- production within the Standard Model itself [6, 10]. In addition, production of W^+W^- pairs is an important process for studying anomalous couplings of electroweak gauge boson. Although current limits are already quite impressive [7-9], it is clear that studies of anomalous gauge boson couplings will intensify once the LHC Run II is underway. Making use of higher experimental precision will require improved modeling of W^+W^- production in the Standard Model. Finally, the $pp \to WW^*$ process with one W-boson on the mass shell and the other off the mass-shell, is an important background to Higgs boson production in $pp \to H \to WW$ channel. Better understanding of this background should allow improved measurements of Higgs boson couplings to W-bosons — including the anomalous ones — in the next run of the LHC. Similar arguments can be given for processes with other vector bosons in the final state.

It follows from these examples, that higher theoretical accuracy for vector-boson pair production in hadron collisions is essential. It can be achieved by extending existing computations of cross sections and kinematic distributions of $pp \rightarrow V_1V_2$ processes [6, 10–38] to next-to-next-to-leading order (NNLO) in perturbative QCD.¹ In particular, predictions

¹First NNLO QCD results for electroweak boson pair production have recently appeared, see e.g. refs. [39–42].

for fiducial volume cross sections, where kinematic restrictions on final state particles are taken into account exactly, are crucial. A NNLO QCD prediction for $pp \rightarrow V_1V_2$ needs three ingredients: *i*) real-emission matrix elements for $q\bar{q}' \rightarrow V_1V_2gg$, $qg \rightarrow V_1V_2qg$ and $q\bar{q} \rightarrow V_1V_2q\bar{q}$; *ii*) one-loop matrix elements for $q\bar{q} \rightarrow V_1V_2g$ and $qg \rightarrow V_1V_2q$ and, finally, *iii*) two-loop amplitudes for tree-level process $q\bar{q}' \rightarrow V_1V_2$. Once these three ingredients become available, they need to be put together in a self-consistent manner using existing methods for fully differential NNLO computations [43–51]. For the V_1V_2 production, the major unknown is the two-loop amplitude for $q\bar{q}' \rightarrow V_1V_2$; the goal of this paper is to provide it.²

The remainder of the paper is organized as follows. In section 2 we explain the general setup of the calculation. Since two-loop scalar master integrals required for this computation have been recently computed for equal electroweak boson masses in refs. [55, 56] and for unequal masses in refs. [57, 58], we mainly focus on the procedure that allows us to express the various contributions to scattering amplitudes in terms of these integrals. In section 3 we describe checks on our computation, evaluate the scattering amplitudes numerically and discuss numerical stability of the results. We conclude in section 4.

2 The setup of the computation

We are interested in the production of a four-lepton final states in proton collisions $pp \rightarrow (l_1 \bar{l}'_1)(l_2 \bar{l}'_2)$. The four-lepton final states can be produced in three distinct ways:

- through the production of a *pair* of off-shell vector bosons, $pp \to V_1 V_2 \to (l_1 \bar{l}'_1)(l_2 \bar{l}'_2);$
- through the production of a *single* vector boson that decays to *two* vector bosons that, in turn, decay to lepton pairs, $pp \to V_3 \to V_1 V_2 \to (l_1 \bar{l}'_1) (l_2 \bar{l}'_2)$;
- through the production of a single vector boson that decays into a pair of leptons (one of them of-shell), followed by the emission of a vector boson by an of-shell lepton, e.g. pp → V₃ → l₃^{*}l'₂ → (l₃^{*} → V₁l₂)l'₂ → (l₁l'₁)(l₂l'₂).

Each of these three processes depends on different combinations of electroweak couplings; therefore, they are separately invariant under QCD gauge transformations and we can compute QCD corrections to each of them separately. We note in this respect that QCD corrections to processes mediated by single gauge-boson production are simple since they are directly related to QCD corrections to the quark form factor of the vector current $F_V(s)$ whose perturbative expansion through NNLO is well-known [59–61].

However, computation of NNLO QCD corrections to process $pp \to V_1V_2 \to (l_1\bar{l}'_1)(l_2\bar{l}'_2)$, where vector bosons couple only to fermions and not to other vector bosons or primary leptons, is non-trivial. Calculation of this contribution to two-loop QCD amplitudes for four-lepton production processes is the main focus of this paper.

²We note that for the simpler case of processes involving one real photon two-loop amplitudes are known both for the quark [52] and gluon [53] channels. For the case of on-shell WW production, the amplitude was calculated in the high-energy approximation in [54].

We consider quark-antiquark annihilation in to vector bosons

$$q(p_1)\bar{q}'(p_2) \to V_1(p_3)V_2(p_4) \to (V_1(p_3) \to l(p_5)\bar{l}'(p_6)) \ (V_2(p_4) \to l(p_7)\bar{l}'(p_8)), \tag{2.1}$$

and work in the approximation where quarks of the first two generations are massless and quarks of the third generation are consistently neglected. We also set the CKM matrix to an identity matrix. Since we work with massless quarks, helicity is a conserved quantum number; therefore, once the helicity of the incoming quark is specified, the helicity of the incoming anti-quark is completely fixed. We will use this observation when writing the amplitude for quark-antiquark annihilation process in eq. (2.1).

The partonic process in eq. (2.1) can proceed in two different ways since the vector bosons can either couple directly to external fermions $q\bar{q}'$ or to closed loops of virtual fermions, see figure 1. Consequently, we write the scattering amplitude for process in eq. (2.1) as

$$\mathcal{M}(\lambda_{q},\lambda_{5},\lambda_{7}) = i \left(\frac{g_{W}}{\sqrt{2}}\right)^{4} \delta_{i_{1}i_{2}} \mathcal{D}_{3}(p_{3}) \mathcal{D}_{4}(p_{4}) C_{l,V_{2}}^{\lambda_{7}} C_{l,V_{1}}^{\lambda_{5}} \epsilon_{3}^{\mu}(\lambda_{5}) \epsilon_{4}^{\nu}(\lambda_{7}) \\ \times \left[C_{\bar{q}',V_{2}}^{\lambda_{q}} C_{q,V_{1}}^{\lambda_{q}} \mathcal{A}_{\mu\nu}^{(d)}(p_{1}^{\lambda_{q}},p_{3},p_{4},p_{2}^{-\lambda_{q}}) + C_{\bar{q}',V_{1}}^{\lambda_{q}} C_{q,V_{2}}^{\lambda_{q}} \mathcal{A}_{\nu\mu}^{(d)}(p_{1}^{\lambda_{q}},p_{4},p_{3},p_{2}^{-\lambda_{q}}) \right. \\ \left. + C_{V_{1}V_{2}}^{n_{g}} \mathcal{A}_{\mu\nu}^{n_{g}}(p_{1}^{\lambda_{q}},p_{2}^{-\lambda_{q}};p_{3},p_{4}) \right],$$

$$(2.2)$$

where $g_W = e/\sin\theta_W$ is the SU(2) weak coupling, $\mathcal{D}_i = 1/(p_i^2 - m_{V_i}^2 + im_{V_i}\Gamma_{V_i})$ is the V_i -boson propagator, λ_q , λ_5 , λ_7 are helicities of the incoming quark and outgoing leptons, respectively, $C_{\vec{q}',V_2}^{\lambda_q}$, $C_{q,V_1}^{\lambda_q}$ and $C_{l,V_2}^{\lambda_7}C_{l,V_1}^{\lambda_5}$ are helicity-dependent couplings of vector bosons to quarks and leptons, and $\epsilon_{3,4}$ are matrix elements for leptonic decays of V_1 and V_2 that we will specify shortly. The amplitudes $\mathcal{A}^{(d)}$ describe direct coupling of the vector bosons to external fermions and the amplitude \mathcal{A}^{n_g} describes contributions of diagrams where vector bosons couple to loops of virtual fermions, see figure 1. The factor $C_{V_1V_2}^{n_g}$ involves sums over couplings of virtual fermions to gauge bosons.

We note that full amplitude for $q\bar{q}' \to V_1 V_2$ in eq. (2.2) is written as the sum of two $\mathcal{A}^{(d)}$ amplitudes where two vector bosons appear in different order. These amplitudes are, therefore, quantities similar to ordered or primitive amplitudes often used in perturbative QCD computations. For us they are useful because we only need to compute one of them and then obtain the other one from $p_3 \leftrightarrow p_4$ permutation. We stress that amplitudes $\mathcal{A}^{(d)}$ and $\mathcal{A}^{(n_g)}$ in eq. (2.2) are computed assuming that vector bosons couple to a single quark generation and that the Vqq coupling is vector-like with unit coefficient, i.e. γ_{μ} . As we explain below, such amplitudes are sufficient to obtain physical amplitudes for pair production of all electroweak gauge bosons.

We begin by discussing the parametrization of $\mathcal{A}_{\mu\nu}^{(d)}\epsilon_{3}^{\mu}\epsilon_{4}^{\nu}$ amplitude in terms of Lorentzinvariant form factors. A representative diagram that contributes to $\mathcal{A}^{(d)}$ at two loops in perturbative QCD is shown in figure 1. Our goal is to re-write these diagrams in such a way that all Feynman integrals can be dealt with using the integration-by-parts technique [62, 63]. To achieve this goal, we need to express the amplitude in terms of



Figure 1. Representative diagrams that contribute to two-loop amplitude for vector boson production in hadron collisions.

invariant form factors. To this end, we note that the most general form of $\mathcal{A}_{\mu\nu}^{(d)}$ is

$$\mathcal{A}_{\mu\nu}^{(d)}(p_1, p_3, p_4, p_2) = \bar{v}_{p_2} \hat{p}_{\perp} u_{p_1} A_{\mu\nu}^{(1,d)} + \bar{v}_{p_2} \gamma^{\mu} u_{p_1} A_{\nu}^{(2,d)} + \bar{v}_{p_2} \gamma^{\nu} u_{p_1} A_{\mu}^{(3,d)} + \bar{v}_{p_2} \gamma^{[\nu} \hat{p}_{\perp} \gamma^{\mu]} u_{p_1} A_{4}^{(4,d)},$$
(2.3)

where $\gamma^{[\nu}\hat{p}_{\perp}\gamma^{\mu]} = \gamma^{\nu}\hat{p}_{\perp}\gamma^{\mu} - \gamma^{\mu}\hat{p}_{\perp}\gamma^{\nu}$ and the vector p_{\perp} is defined by the Sudakov decomposition of the momenta $p_{3,4}$

$$p_3 = \alpha_3 p_1 + \beta_3 p_2 + p_\perp, \quad p_4 = \alpha_4 p_1 + \beta_4 p_2 - p_\perp.$$
 (2.4)

The transverse momentum p_{\perp} is orthogonal to $p_{1,2}$, $p_{\perp} \cdot p_{1,2} = 0$. The coefficients $\alpha_{3,4}$ and $\beta_{3,4}$ in eq. (2.4) can be written as

$$\alpha_3 = \frac{m_3^2 - u}{s}, \quad \beta_3 = \frac{m_3^2 - t}{s}, \quad \alpha_4 = \frac{m_4^2 - t}{s}, \quad \beta_4 = \frac{m_4^2 - u}{s}, \tag{2.5}$$

where we defined $p_3^2 = m_3^2$, $p_4^2 = m_4^2$ and introduced standard Mandelstam variables $s = (p_1 + p_2)^2$, $t = (p_1 - p_3)^2$, $u = (p_1 - p_4)^2$. The functions $A^{(i,d)}$, $i = 1, \ldots, 4$, introduced in eq. (2.3), depend on momenta and Lorentz indices. To make this dependence explicit, we decompose them into invariant form factors T_i , $i = 1, \ldots, 17$,

$$\begin{aligned} A^{(1,d)}_{\mu\nu} &= T_1 g_{\mu\nu} + T_2 p_{1\mu} p_{1\nu} + T_3 p_{1\mu} p_{2\nu} + T_4 p_{1\mu} p_{\perp\nu} + T_5 p_{2\mu} p_{1\nu} + T_6 p_{2\mu} p_{2\nu} + T_7 p_{2\mu} p_{\perp\nu} \\ &+ T_8 p_{\perp\mu} p_{1\nu} + T_9 p_{\perp\mu} p_{2\nu} + T_{10} p_{\perp\mu} p_{\perp\nu}, \end{aligned} \tag{2.6} \\ A^{(2,d)}_{\mu} &= T_{11} p_{1\mu} + T_{12} p_{2\mu} + T_{13} p_{\perp\mu}, \quad A^{(3,d)}_{\nu} = T_{14} p_{1\nu} + T_{15} p_{2\nu} + T_{16} p_{\perp\nu}, \quad A^{(4,d)} = T_{17}. \end{aligned}$$

We note that not all invariant form factors that appear in eq. (2.6) give independent contributions to physical amplitudes. This happens because we did not use the transversality condition for lepton currents $p_3 \cdot \epsilon_3 = p_4 \cdot \epsilon_4 = 0$. As we will see shortly, when physical amplitudes are computed, the number of relevant form factors will be reduced thanks to the transversality condition.

To calculate the physical amplitude in eq. (2.2), we need to contract $\mathcal{A}_{\mu\nu}^{(d)}$ with polarization vectors of external vector bosons. As we already mentioned, they are given by matrix elements of the vector current between relevant leptonic states. Using the spinor-helicity formalism, we write

$$\epsilon_3^{\mu}(5_L) = \langle 5|\gamma^{\mu}|6], \qquad \epsilon_3^{\mu}(5_R) = [5|\gamma^{\mu}|6\rangle = \langle 6|\gamma^{\mu}|5], \\ \epsilon_4^{\nu}(7_L) = \langle 7|\gamma^{\mu}|8], \qquad \epsilon_4^{\nu}(7_R) = [7|\gamma^{\mu}|8\rangle = \langle 8|\gamma^{\mu}|7].$$
(2.7)

Since, as we already pointed out, the helicity of the incoming quark fully determines the allowed helicity of the incoming antiquark, we need eight helicity amplitudes to fully describe production of two vector bosons. They are $\mathcal{M}(q_L, 5_L, 7_L)$, $\mathcal{M}(q_L, 5_R, 7_L)$, $\mathcal{M}(q_L, 5_L, 7_R)$, $\mathcal{M}(q_L, 5_R, 7_R)$, $\mathcal{M}(q_R, 5_L, 7_L)$, $\mathcal{M}(q_R, 5_R, 7_L)$, $\mathcal{M}(q_R, 5_L, 7_R)$, $\mathcal{M}(q_R, 5_R, 7_R)$. We note that, according to eq. (2.7), a change in lepton helicities can be obtained by interchanging momenta $p_5 \leftrightarrow p_6$ and $p_7 \leftrightarrow p_8$, where necessary. Therefore, we obtain all required helicity amplitudes from $\mathcal{M}(q_L, 5_L, 7_L)$ and $\mathcal{M}(q_R, 5_L, 7_L)$ by simple permutations of lepton momenta.

For left- and right-handed incoming quarks we find

$$\begin{split} A^{(d)}_{LLL}(3,4) &= \mathcal{A}^{(d)}_{\mu\nu}(p_1^L,p_3,p_4,p_2^R)\epsilon^{\mu}_3(5_L)\epsilon^{\nu}_4(7_L) = -F_1\langle 57\rangle[86]\langle 2\hat{3}1] + F_2\langle 15\rangle\langle 17\rangle[16][18]\langle 2\hat{3}1] \\ &+ F_3\langle 15\rangle\langle 27\rangle[16][28]\langle 2\hat{3}1] + F_5\langle 17\rangle\langle 25\rangle[18][26]\langle 2\hat{3}1] + F_6\langle 25\rangle\langle 27\rangle[26][28]\langle 2\hat{3}1] \\ &+ F_{11}\langle 25\rangle\langle 17\rangle[16][18] + F_{12}\langle 25\rangle\langle 27\rangle[16][28] + F_{14}\langle 15\rangle\langle 27\rangle[16][18] + F_{15}\langle 25\rangle\langle 27\rangle[26][18], \\ A^{(d)}_{RLL}(3,4) &= \mathcal{A}^{(d)}_{\mu\nu}(p_1^R,p_3,p_4,p_2^L)\epsilon^{\mu}_3(5_L)\epsilon^{\nu}_4(7_L) = -F_1\langle 57\rangle[86][2\hat{3}1\rangle + F_2\langle 15\rangle\langle 17\rangle[16][18][2\hat{3}1\rangle \\ &+ F_3\langle 15\rangle\langle 27\rangle[16][28][2\hat{3}1\rangle + F_5\langle 17\rangle\langle 25\rangle[18][26][2\hat{3}1\rangle + F_6\langle 25\rangle\langle 27\rangle[26][28][2\hat{3}1\rangle \\ &+ F_{11}[26]\langle 15\rangle\langle 17\rangle[18] + F_{12}[28]\langle 15\rangle[26]\langle 27\rangle + F_{14}[28]\langle 15\rangle[16]\langle 17\rangle + F_{15}[28]\langle 25\rangle[26]\langle 17\rangle, \end{split}$$

where the following *nine* combinations of form factors enter

$$F_{1} = -2T_{1},$$

$$F_{2} = T_{2} - \alpha_{3}\alpha_{4}T_{10} - \alpha_{3}T_{8} + \alpha_{4}T_{4},$$

$$F_{3} = T_{3} - \frac{4T_{17}}{s} - \alpha_{3}\beta_{4}T_{10} - \alpha_{3}T_{9} + \beta_{4}T_{4},$$

$$F_{5} = T_{5} + \frac{4T_{17}}{s} - \beta_{3}\alpha_{4}T_{10} - \beta_{3}T_{8} + \alpha_{4}T_{7},$$

$$F_{6} = T_{6} - \beta_{3}\beta_{4}T_{10} - \beta_{3}T_{9} + \beta_{4}T_{7},$$

$$F_{11} = 2T_{11} + 2\alpha_{4}T_{13},$$

$$F_{12} = 2T_{12} + 2\beta_{4}T_{13},$$

$$F_{14} = 2T_{14} - 2\alpha_{3}T_{16},$$

$$F_{15} = 2T_{15} - 2\beta_{3}T_{16}.$$
(2.9)

We note that form factors only depend on s, t, u and m_3^2, m_4^2 ; therefore, they are invariant under $p_5 \leftrightarrow p_6$ and $p_7 \leftrightarrow p_8$ permutations. For this reason, it is straightforward to obtain all the relevant amplitudes from eq. (2.8). To find $T_{1,\dots,17}$ and $F_{1,2,\dots,15}$, we need to construct projection operators. To accomplish this, we write a generic amplitude as a matrix element of a string of Dirac matrices

$$\mathcal{A}_{\mu\nu} = \bar{v}_{p_2} \hat{\Gamma}_{\mu\nu} u_{p_1}. \tag{2.10}$$

Multiplying it with $\bar{u}_{p_1} \hat{O} v_{p_2}$ and summing over spinor helicities, we find

$$\sum \mathcal{A}_{\mu\nu} \bar{u}_{p_1} \hat{O} v_{p_2} = \operatorname{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{O} \right].$$
(2.11)

Choosing different operators \mathcal{O} , we can project on individual T form factors or their combinations. Below we list all the projection operators and the results that we get when these

operators are convoluted with the amplitude

$$\begin{split} &G_1 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^3} \times p_1^\mu p_2^\nu, \qquad G_1 = T_6, \\ &G_2 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^3} \times p_1^\mu p_2^\nu, \qquad G_2 = T_2, \\ &G_3 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_1^\mu p_2^\nu, \qquad G_3 = T_1 + 2T_{17} + (p_1 \cdot p_2)T_5, \\ &G_4 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_1^\mu p_1^\nu, \qquad G_5 = T_7 p_2^2 + T_{15}, \\ &G_5 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_1^\mu p_1^\nu, \qquad G_6 = T_9 p_2^2 + T_{12}, \\ &G_7 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_2^\mu p_1^\nu, \qquad G_8 = T_4 p_2^2 + T_{14}, \\ &G_8 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_2^\mu p_2^\nu, \qquad G_8 = T_4 p_2^2 + T_{14}, \\ &G_9 = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_2^\mu p_2^\nu, \qquad G_{10} = d T_1 + (T_3 + T_5) p_1 \cdot p_2 + T_{10} p_2^2 + T_{16} + T_{13}, \\ &G_{10} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)^2} \times p_2^\mu, \qquad G_{12} = T_9 p_2^2 + (d - 2)T_{11}, \\ &G_{12} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4(p_1 \cdot p_2)^2} \times p_2^\mu, \qquad G_{12} = T_9 p_2^2 + (d - 2)T_{12}, \\ &G_{13} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4(p_1 \cdot p_2)^2} \times p_1^\mu, \qquad G_{12} = T_9 p_2^2 + (d - 2)T_{14}, \\ &G_{14} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4(p_1 \cdot p_2)^2} \times p_1^\mu, \qquad G_{13} = T_7 p_2^2 + (d - 2)T_{14}, \\ &G_{14} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4(p_1 \cdot p_2)^2} \times p_1^\mu, \qquad G_{15} = T_1 + T_{10} p_1^2 + (d - 2)T_{13} + T_{16}, \quad (2.12) \\ &G_{15} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)} \times p_1^\mu, \qquad G_{15} = T_1 + T_{10} p_1^2 + T_{13} + (d - 2)T_{16}, \\ &G_{15} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)} \times p_1^\mu, \qquad G_{16} = T_1 + T_{10} p_1^2 + T_{13} + (d - 2)T_{16}, \\ &G_{16} = -\frac{\text{Tr} \left[\hat{p}_2 \Gamma_{\mu\nu} \hat{p}_1 \hat{p}_1 \right]}{4p_1^2 (p_1 \cdot p_2)} \times p_1^\mu, \qquad G_{16} = T_1 + T_{10} p_1^2 + T_{13} + (d - 2)T_{16}, \\ &G_{16} = -\frac{\text{Tr$$

Each of the $G_{1...17}$ projections can be calculated from Feynman diagrams by taking traces and integrating over loop momenta. These quantities depend on the Mandelstam invariants of the underlying $2 \rightarrow 2$ process $q\bar{q}' \rightarrow V_1V_2$ but not on the polarization vectors of electroweak bosons or on the momenta of final-state leptons. Thus, they can be expressed in terms of Feynman integrals of the type introduced in refs. [57, 58] and then reduced to master integrals using integration-by-parts identities.

We can use projections $G_{1,\dots,17}$ to derive the T form factors; the result reads

$$\begin{split} T_1 &= \frac{G_{10} - G_9 - G_4 - G_3}{d - 3}, \qquad T_2 = G_2, \\ T_3 &= -\frac{2(d(6 - d)G_4 + (3 - d)G_3 + (d - 4)G_{10} - 7G_4 + (4 - d)G_9 + G_{17})}{s_{12}(d - 3)(d - 4)}, \\ T_4 &= -\frac{G_{14} + (2 - d)G_8}{p_{\perp}^2(d - 3)}, \\ T_5 &= -\frac{2((3 - d)G_4 - G_{17} - (d^2 - 6d + 7)G_3 + (d - 4)G_{10} + (4 - d)G_9)}{s_{12}(d - 3)(d - 4)}, \\ T_6 &= G_1, \qquad T_7 = -\frac{G_{13} + (2 - d)G_5}{p_{\perp}^2(d - 3)}, \qquad T_8 = -\frac{G_{11} + (2 - d)G_7}{p_{\perp}^2(d - 3)}, \\ T_9 &= -\frac{G_{12} + (2 - d)G_6}{p_{\perp}^2(d - 3)}, \qquad T_{10} = -\frac{G_{15} - G_4 - G_3 - dG_9 + G_{16} + G_{10}}{p_{\perp}^2(d - 3)}, \\ T_{11} &= \frac{G_{11} - G_7}{d - 3}, \qquad T_{12} = \frac{G_{12} - G_6}{d - 3}, \qquad T_{13} = \frac{G_{15} - G_9}{d - 3}, \qquad T_{14} = \frac{G_{14} - G_8}{d - 3}, \\ T_{15} &= \frac{G_{13} - G_5}{d - 3}, \qquad T_{16} = \frac{G_{16} - G_9}{d - 3}, \qquad T_{17} = -\frac{G_4 - G_3 + G_{17}}{2(d - 3)(d - 4)}. \end{split}$$

$$(2.13)$$

The F form factors that are used for the evaluation of the amplitude can now be easily computed using eq. (2.9). We do not present these results here since they are not very illuminating.

One point worth emphasizing though is that in eq. (2.13) there are expressions (see e.g. equation for T_{17}) that appear to have spurious singularities in the limit $d \to 4$. It would have been unfortunate if these singularities survive in the final formulas for form factors F since, in such a situation, computation of all pieces needed for the evaluation of G's, including master integrals, to higher orders in ϵ is required. It is therefore pleasing to observe that this does not happen and once results for the F form factors are written in terms of $G_{1,\ldots,17}$ projectors, all spurious (d-4) singularities disappear.

For future reference, we give results for leading order form factors that appear in the physical amplitudes eq. (2.8)

$$F_{1} = \frac{2}{t}, \quad F_{2} = 0, \quad F_{3} = -\frac{2}{st}, \quad F_{5} = \frac{2}{st}, \quad F_{6} = 0, \quad F_{11} = \frac{2(m_{4}^{2} - t)}{st},$$

$$F_{12} = \frac{2(s + m_{3}^{2} - t)}{st}, \quad F_{14} = -\frac{2(s + m_{4}^{2} - t)}{st}, \quad F_{15} = -\frac{2(m_{3}^{2} - t)}{st}.$$
(2.14)

We stress that these results are exact in a sense that no $d \rightarrow 4$ limit was taken to obtain them; in other words, all the *d*-dependence cancels out completely once physical form factors are computed.

Before proceeding to the discussion of various couplings of vector bosons to fermions introduced in eq. (2.2), we want to explain why $\mathcal{A}^{(d)}$ and $\mathcal{A}^{(n_g)}$ amplitudes can be computed assuming that interactions of vector bosons with quarks are mediated by vector currents. The main reason is that we only consider here contributions of massless quarks and that, in such a case, the helicity is conserved. Therefore, we can always write couplings of vector bosons to fermions as linear combinations of left- and right-handed couplings and then move helicity projection operators to external lines. In this way the helicity-dependent couplings are generated and amplitudes that remain can be viewed as originating from pure vector current interactions of gauge bosons and quarks.

Of course, this discussion applies only to diagrams where vector bosons couple directly to external quark lines; in the notation of eq. (2.2) such diagrams contribute to $\mathcal{A}^{(d)}$ amplitudes. However, we will now argue that the same parametrization eq. (2.3) can be used to compute amplitudes $\mathcal{A}^{(n_g)}$, which receive contributions from diagrams where vector bosons couple to closed fermion loops. In fact, this would have been obvious provided that electroweak boson coupling to fermions is vector-like since in this case tracing over spin degrees of freedom in the internal quark loop gives us diagrams that are not very different from the ones that we already considered. Potential problems could be expected with axial couplings and, in particular, with terms where one axial and three vector couplings appear in qqV_1V_2 Green's function. However, all such terms cancel because of C-parity conservation for massless fermions for any final state with two vector bosons [35, 36].³ Contributions of terms with two vector and two axial couplings are not anomalous and must be equal to those with four vector couplings. Hence, we conclude that it is sufficient to consider vector-current couplings of gauge bosons to internal fermion loops, to account for all non-vanishing contributions to the amplitude. The parametrization of the amplitude $\mathcal{A}^{(n_g)}$ is then taken from eq. (2.3) and its expression through invariant form factors is taken from eq. (2.8).

To complete our construction of the scattering amplitude for $q\bar{q}' \rightarrow V_1 V_2$ in eq. (2.2), it remains to specify various helicity-dependent couplings that we introduced there. Below we present these couplings for various pairs of electroweak gauge bosons that can be produced in proton collisions.

2.1 $\gamma^*\gamma^*$ production

Photons are produced in the annihilation of a quark and an antiquark of the same flavor q = q'. Photon interactions with both quarks and leptons are pure vector-like and, therefore, helicity-independent. We find

$$C_{q,\gamma}^{L,R} = -\sqrt{2}Q_q \sin\theta_W, \quad C_{\bar{q},\gamma}^{L,R} = -\sqrt{2}Q_q \sin\theta_W, \quad C_{l,\gamma}^{L,R} = -\sqrt{2}Q_l \sin\theta_W, \quad (2.15)$$

where Q_l and Q_q are electromagnetic quark and lepton charges in units of the positron charge. Finally, $C_{\gamma\gamma}^{n_g}$ is given by the sum of up and down quark charges of the first two generations

$$C_{\gamma\gamma}^{n_g} = 2\sin^2\theta_W \sum_{i_q=1}^2 \left(Q_u^2 + Q_d^2\right) = \frac{20\sin^2\theta_W}{9}.$$
 (2.16)

³More precisely, for C-parity argument to be applicable, all fermions in the loop should have equal masses [35, 36].

2.2 ZZ production

Pairs of Z bosons are produced in annihilation of quarks and anti-quarks of the same flavor. Couplings of Z bosons to quarks and leptons depend on helicity and weak isospin of the corresponding particle. We find

$$C_{q,Z}^{L,R} = C_{\bar{q},Z}^{L,R} = \frac{1}{\sqrt{2}\cos\theta_W} \left(V_q \pm A_q \right), \quad C_{l,Z}^{L,R} = \frac{1}{\sqrt{2}\cos\theta_W} \left(V_l \pm A_l \right), \tag{2.17}$$

where

$$V_{u} = \frac{1}{2} - \frac{4}{3}\sin^{2}\theta_{W}, \qquad A_{u} = \frac{1}{2}, \qquad V_{d} = -\frac{1}{2} + \frac{2}{3}\sin^{2}\theta_{W}, \qquad A_{d} = -\frac{1}{2},$$

$$V_{e} = -\frac{1}{2} + 2\sin^{2}\theta_{W}, \qquad A_{e} = -\frac{1}{2}, \qquad V_{\nu} = \frac{1}{2} \quad A_{\nu} = \frac{1}{2}.$$
(2.18)

The coefficient
$$C_{ZZ}^{n_g}$$
 is given by the sum of up and down quark vector and axial charges of

the first two generations. We find $C_{ZZ}^{n_g} = \frac{1}{\cos^2 \theta_W} \left(V_u^2 + V_d^2 + A_u^2 + A_d^2 \right).$ (2.19)

2.3 $Z\gamma^*$ production

Production of Z and γ^* occurs in annihilation of quarks and anti-quarks of the same flavor. The couplings to external fermions are given in eqs. (2.15), (2.17). The coefficient $C_{\gamma Z}^{(n_g)}$ reads

$$C_{Z\gamma}^{n_g} = -\frac{2\sin\theta_W}{\cos\theta_W} \left(V_u Q_u + V_d Q_d \right).$$
(2.20)

2.4 $W^+\gamma^*$ and for $W^-\gamma^*$ production

The $W^+\gamma^*$ final state is produced in $u\bar{d}$ annihilation, $u\bar{d} \to W^+\gamma^*$. Production of $W^-\gamma^*$ occurs in $d\bar{u} \to W^-\gamma^*$ process. Since W bosons interact with left-handed fermions, we have

$$C_{u,W^{+}}^{\lambda} = \delta_{\lambda,L}, \quad C_{\bar{d},\gamma}^{\lambda} = -\sqrt{2}Q_{d}\sin\theta_{W}, \quad C_{u,\gamma}^{\lambda} = -\sqrt{2}Q_{u}\sin\theta_{W}, \quad C_{\bar{d},W^{+}}^{\lambda} = \delta_{\lambda,L}, \\ C_{d,W^{-}}^{\lambda} = \delta_{\lambda,L}, \quad C_{\bar{u},\gamma}^{\lambda} = -\sqrt{2}Q_{u}\sin\theta_{W}, \quad C_{d,\gamma}^{\lambda} = -\sqrt{2}Q_{d}\sin\theta_{W}, \quad C_{\bar{u},W^{-}}^{\lambda} = \delta_{\lambda,L}.$$
(2.21)
$$C_{l,W^{+}}^{\lambda} = C_{l,W^{-}}^{\lambda} = \delta_{\lambda,L}, \quad C_{l,\gamma}^{\lambda} = -\sqrt{2}Q_{l}\sin\theta_{W}.$$

Coefficients $C_{W^+\gamma}^{n_g}$ and $C_{W^-\gamma}^{n_g}$ vanish identically due to electric charge conservation.

2.5 W^+Z and for W^-Z production

Production processes for $W^{\pm}Z$ are identical to $W^{\pm}\gamma^*$. The relevant couplings are

$$C_{u,W^{+}}^{\lambda} = \delta_{\lambda,L}, \quad C_{\bar{d},W^{+}}^{\lambda} = \delta_{\lambda,L}, \quad , C_{d,W^{-}}^{\lambda} = \delta_{\lambda,L}, \quad C_{\bar{u},W^{-}}^{\lambda} = \delta_{\lambda,L}, \quad C_{l,W^{+}}^{\lambda} = C_{l,W^{-}}^{\lambda} = \delta_{\lambda,L}, \quad C_{q,Z}^{L,R} = C_{\bar{q},Z}^{L,R} = \frac{1}{\sqrt{2}\cos\theta_{W}} \left(V_{q} \pm A_{q} \right), \quad C_{l,Z}^{L,R} = \frac{1}{\sqrt{2}\cos\theta_{W}} \left(V_{l} \pm A_{l} \right), \quad (2.22)$$

Coefficients $C_{W^{\pm}Z}^{n_g}$ vanish identically due to electric charge conservation.

2.6 W^+W^- production

Finally, we display the necessary couplings for the pair production of two W bosons. This process can occur thanks to $u\bar{u}$ and $d\bar{d}$ annihilation. First, consider the process $u(p_1)\bar{u}(p_2) \to W^+(p_3)W^-(p_4)$. In this case,

$$C_{u,W^+}^{\lambda} = \delta_{\lambda L}, \quad C_{\bar{u},W^-}^{\lambda} = \delta_{\lambda,L}, \quad C_{u,W^-}^{\lambda} = 0, \quad C_{\bar{u},W^+}^{\lambda} = 0.$$
 (2.23)

In case of $d(p_1)\overline{d}(p_2) \to W^+(p_3)W^-(p_4)$, the coupling constants read

$$C_{d,W^+}^{\lambda} = 0, \quad C_{\bar{d},W^-}^{\lambda} = 0, \quad C_{d,W^-}^{\lambda} = \delta_{\lambda L}, \quad , \quad C_{\bar{d},W^+}^{\lambda} = \delta_{\lambda,L}.$$
 (2.24)

Couplings to leptons are given in eq. (2.22). The coefficient $C_{W^+W^-}^{(n_g)}$ receives contributions from two generations⁴ and reads

$$C_{W^+W^-}^{(n_g)} = 1. (2.25)$$

3 Calculation of the form factors and the amplitudes

Once projection operators are established, one can compute the form factors and construct the scattering amplitudes for arbitrary di-boson final state. To this end, we note that all integrals that appear in the calculation of form factors can be associated with one of the six topologies, introduced in refs. [57, 58]. Integrals that belong to each of these topologies are closed under the integration-by-parts identities [62, 63]. We use QGRAF [65] to generate Feynman diagrams and FORM [66] for algebraic manipulations and computation of projections $G_{1,...,17}$. We use FIRE [67–69] to reduce all the integrals that appear in this calculation to master integrals. The master integrals were computed by us in refs. [57, 58]. In principle, once all the ingredients are in place, computation of the amplitude becomes straightforward. In practice, however, it requires some effort to put all the pieces together primarily because algebraic expressions that appear e.g. in the course of the reduction to master integrals are quite large in size. The master integrals are expressed in terms of Goncharov polylogarithms up to weight four; we use GiNaC [70] implementation [71] to compute them.

The results for the amplitudes contain infra-red and ultraviolet divergences. The ultraviolet divergences are removed by the renormalization of the strong coupling constant. Since tree-level scattering amplitudes are independent of α_s , we only need its renormalization through one loop. It reads

$$\alpha_s^{(0)} S_\epsilon = \alpha_s \mu^{2\epsilon} \left(1 - \frac{\beta_0}{\epsilon} \left(\frac{\alpha_s}{2\pi} \right) + \mathcal{O}(\alpha_s^2) \right), \tag{3.1}$$

where $\alpha_s^{(0)}$ is the bare and $\alpha_s = \alpha_s(\mu)$ is the renormalized coupling constant. In addition, $\beta_0 = (11C_A - 4T_R n_f)/6 = 11/2 - n_f/3$ is the QCD beta-function, n_f is the number of massless flavors, $C_A = 3$, $T_R = 1/2$ and $S_{\epsilon} = (4\pi)^{\epsilon} e^{-\epsilon \gamma_E}$, with γ_E being the Euler constant.

⁴As we already mentioned several times, we do not consider the third quark generation in this paper.

According to Catani [72], infra-red singularities of UV-renormalized amplitudes at next-to-next-to-leading order are fully determined by leading and next-to-leading order amplitudes. This feature can be used as an important check of the correctness of the computation. To introduce Catani's result, we write a UV-renormalized amplitude as

$$\mathcal{M} = \mathcal{M}_0 + \left(\frac{\alpha_s(\mu)}{2\pi}\right) \mathcal{M}_1 + \left(\frac{\alpha_s(\mu)}{2\pi}\right)^2 \mathcal{M}_2 + \dots$$
(3.2)

Leading order amplitude \mathcal{M}_0 in this formula is finite and, in fact, *d*-independent. The next-to-leading order amplitude contains infra-red divergences that, however, can be written in a factorized form

$$\mathcal{M}_1 = I_1(\epsilon)\mathcal{M}_0 + \mathcal{M}_{\text{fin}}.$$
(3.3)

For the process of vector boson pair production, the final state is neutral. Therefore,

$$I_1(\epsilon) = -\frac{e^{-\epsilon(L_s - \gamma_E)}}{\Gamma(1 - \epsilon)} C_F\left(\frac{1}{\epsilon^2} + \frac{3}{2\epsilon}\right), \qquad (3.4)$$

where $L_S = \ln(-s/\mu^2 - i0) = \ln |s/\mu^2| - i\pi$ and $C_F = 4/3$. In variance with \mathcal{M}_0 , \mathcal{M}_1 and $\mathcal{M}_{1,\text{fin}}$ depend on the dimensional regularization parameter ϵ . This feature is important for proper comparison of Catani's formula with results of explicit computation.

The two-loop amplitude \mathcal{M}_2 can be written in a similar way

$$\mathcal{M}_2 = \hat{I}_2(\epsilon)\mathcal{M}_0 + \hat{I}_1(\epsilon)\mathcal{M}_1 + \mathcal{M}_{2,\text{fin}}, \qquad (3.5)$$

where

$$\hat{I}_2 = -\frac{1}{2}I_1^2(\epsilon) - \frac{\beta_0}{\epsilon}I_1(\epsilon) + \frac{e^{-\epsilon\gamma_E}\Gamma(1-2\epsilon)}{\Gamma(1-\epsilon)} \left(\frac{\beta_0}{\epsilon} + k_q\right)I_1(2\epsilon) + \frac{H_q}{2\epsilon}.$$
(3.6)

The two constants that enter this formula read

$$k_q = \left(\frac{67}{18} - \frac{\pi^2}{6}\right) C_A - \frac{10}{9} T_R n_f, \tag{3.7}$$

$$H_q = C_F^2 \left(-\frac{3}{8} + \frac{\pi^2}{2} - 6\zeta_3 \right) + C_F n_f T_R \left(-\frac{25}{54} + \frac{\pi^2}{12} \right) + C_F C_A \left(\frac{245}{216} - \frac{23\pi^2}{48} + \frac{13\zeta_3}{2} \right).$$

We note that Catani formula can also be used for individual form factors that we introduced earlier to describe physical amplitudes. To this end, we only need to replace tree and loop amplitudes in the above formulas with the corresponding form factors. We have used the above results for the infra-red poles of scattering amplitudes to check the correctness of our computation of the amplitudes $\mathcal{A}^{(d)}$. We also note that the amplitude $\mathcal{A}^{(n_g)}$ appears for the first time at NNLO; therefore, according to Catani's formula it cannot have infra-red $1/\epsilon$ singularities. This is an important check of the correctness of the computation of the amplitude $\mathcal{A}^{(n_g)}$.

We turn to the discussion of numerical results for the scattering amplitudes $\mathcal{A}^{(d)}$, $\mathcal{A}^{(n_g)}$. We define those amplitudes by contracting them with polarization vectors of electroweak bosons

$$\mathcal{A}^{(d,n_g)}(\lambda_q,\lambda_5,\lambda_7) = \mathcal{A}^{(d,n_g)}(p_1^{\lambda_q}, p_3^{\lambda_5}, p_4^{\lambda_7}, p_2^{-\lambda_q}) = \mathcal{A}^{(d,n_g)}_{\mu\nu}(p_1^{\lambda_q}, p_3, p_4, p_2^{-\lambda_q})\epsilon_3^{\mu}(\lambda_5)\epsilon_4^{\nu}(\lambda_7).$$
(3.8)

Helicity	ϵ^{-2}	ϵ^{-1}	ϵ^0	ϵ	ϵ^2
$\tilde{A}_{LLL}^{(d,1)}$	$-\frac{4}{3}$	-2 - i4.1887902	3.2003253 - i 6.2069828	5.4520124 - i 2.9495550	$5.3607865 \!-\! i 3.1814370$
$\tilde{A}_{RLL}^{(d,1)}$	$-\frac{4}{3}$	-2 - i4.1887902	$4.0059079 \!-\! i 3.0593301$	$0.5752861 \!+\! i 5.8440713$	$-9.6769949\!+\!i0.9775875$

Table 1. Ratios of selected one-loop helicity amplitudes and tree amplitudes, see eq. (3.9). Momenta of external particles are given in the main text of the paper. Tree-amplitudes are $\mathcal{A}_{LLL}^{(d,0)} = 4600.82746 - i \, 17933.17244$ and $\mathcal{A}_{RLL}^{(d,0)} = 732.100366 - i \, 1148.55597$.

and write their perturbative expansion as

$$\mathcal{A}^{(d)} = \mathcal{A}^{(d,0)} \left[1 + a_0 \ s^{-\epsilon} \tilde{A}^{(d,1)} + a_0^2 \ s^{-2\epsilon} \tilde{A}^{(d,2)} \right], \qquad \mathcal{A}^{(n_g)} = \mathcal{A}^{(d,0)} a_0^2 \ s^{-2\epsilon} \tilde{A}^{(n_g,2)}, \tag{3.9}$$

where $a_0 = \alpha_s^{(0)}(4\pi)^{\epsilon}\Gamma(1+\epsilon)/(2\pi)$. We note that in eq. (3.9) we choose to expand in bare, rather than renormalized, QCD coupling. Also, we made it explicit in eq. (3.9) that the amplitude $\mathcal{A}^{(n_g)}$ appears at two loops for the first time.

To motivate our choice of kinematics for numerical results for the amplitudes that we present below, we consider $q\bar{q}' \to W^+W^-$ production as the background to Higgs boson signal in $pp \to H \to W^+W^-$. Therefore, we choose the center-of-mass energy \sqrt{s} to be the mass of the Higgs boson $\sqrt{s} = m_H = 125$ GeV. The invariant mass of the vector boson V_1 is set to $p_3^2 = m_W^2$, with $m_W = 80.419$ GeV. The invariant mass of the second vector boson V_2 is set to 25 GeV. We take the vector boson scattering angle in the center-of-mass collision frame to be $\pi/3$ radians. We also take decay angles of the lepton l_5 in the rest frame of the boson V_1 to be $\theta_5 = \pi/4$ and $\varphi_5 = \pi/2$ and decay angles of the lepton l_7 in the rest frame of the boson V_2 to be $\theta_7 = \pi/6$ and $\varphi_7 = \pi$. The four-momenta of initial and final state particles are given by

$$p_{1} = (62.5, 0, 0, 62.5), \quad p_{2} = (62.5, 0, 0, -62.5),$$

$$p_{5} = (48.2561024468725, 13.8697156788798, -28.4324101181205, 36.4400941989053),$$

$$p_{6} = (37.6127597971275, 12.2010429705974, 28.4324101181205, -21.3881346746519),$$

$$p_{7} = (19.5655688780000, -19.2853793247386, 0, 3.29933778517879),$$

$$p_{8} = (19.5655688780000, -6.78537932473856, 0, -18.3512973094322). \quad (3.10)$$

To obtain numerical results for the amplitude, we take the number of massless fermion species n_f to be five. Results for selected helicity amplitudes $A^{(d)}$ and $A^{(n_g)}$ are shown in tables 1, 2, 3. They can be compared to predictions based on Catani's formula.⁵ For both one- and two-loop amplitudes, divergent terms agree perfectly. For the one-loop amplitudes, we can also compare the $\mathcal{O}(\epsilon^0)$ terms against the known results [73]; we find perfect agreement. The amplitude $A^{(n_g)}$ that describes contributions to $q\bar{q}' \to V_1 V_2$ where vector bosons couple to closed loops of fermions is finite as expected since those amplitudes have no tree- and one-loop contributions.

We will now elaborate on the numerical stability of our results. This is an important issue since amplitudes for vector boson pair productions may exhibit numerical instabilities in the limit of forward or backward scattering. In fact, numerical stability depends

⁵Catani's formula needs to be re-written to provide an expansion in the unrenormalized QCD coupling.

Helicity	ϵ^{-4}	ϵ^{-3}	ϵ^{-2}	ϵ^{-1}	ϵ^0
$\tilde{A}_{LLL}^{(d,2)}$	$\frac{8}{9}$	1.388889 + i 5.58505	-16.02478 + i 8.625043	$-29.43232\!-\!i28.086442$	23.53917 - i67.620386
$\tilde{A}_{RLL}^{(d,2)}$	$\frac{8}{9}$	1.388889 + i 5.58505	-17.09889 + i 4.42818	$-8.268265\!-\!i37.414997$	73.483267 + i23.301609

Table 2. Ratios of selected two-loop helicity amplitudes and tree amplitudes, see eq. (3.9). Momenta of external particles are given in the main text of the paper. The number of massless fermion species n_f is taken to be five. Tree-amplitudes are $\mathcal{A}_{LLL}^{(d,0)} = 4600.82746 - i \, 17933.17244$ and $\mathcal{A}_{RLL}^{(d,0)} = 732.100366 - i \, 1148.55597.$

Helicity	$1/\epsilon^4$	$1/\epsilon^3$	ϵ^2	$1/\epsilon$	ϵ^0
$\tilde{A}_{LLL}^{(n_g,2)}$	0	$\mathcal{O}(10^{-14})$	$\mathcal{O}(10^{-10})$	$\mathcal{O}(10^{-8})$	$-0.6207871 + i \ 0.01394607$
$\tilde{A}_{RLL}^{(n_g,2)}$	0	$\mathcal{O}(10^{-13})$	$\mathcal{O}(10^{-8})$	$O(10^{-7})$	$2.6570701 + i \ 4.89645656$

Table 3. Ratios of finite two-loop $\mathcal{A}^{(n_g)}$ helicity amplitudes to tree amplitudes. Momenta of external particles are given in the main text of the paper. Tree-amplitudes are $\mathcal{A}_{LLL}^{(d,0)} = 4600.82746 - i 17933.17244$ and $\mathcal{A}_{RLL}^{(d,0)} = 732.100366 - i 1148.55597$.



Figure 2. Absolute value squared of the ratio of $1/\epsilon$ poles of the *LLL* scattering amplitude computed from Feynman diagrams and using Catani formula, $R_{1/\epsilon} = ||\tilde{A}_{LLL}^{(d,2)}|^2/|\tilde{A}_{C,LLL}^{(d,2)}|^2 - 1|$. The center-of-mass energy, gauge boson masses and lepton scattering angles are given in the text.

on the vector boson transverse momentum since $1/p_{\perp}^2$ singularities appear when Feynman integrals are reduced to master integrals. We have evidence from previous studies about values of transverse momenta where such instabilities arise. In case of one-loop $gg \rightarrow VV$ amplitudes, numerical instabilities start to appear for transverse momenta of an order of a few GeV [74, 75] and sophisticated treatment is required to remove them completely [75]. To explore numerical (in)stability of our results, we study amplitudes $\tilde{A}_{LLL}^{(d,n_g,2)}$ in dependence of the vector boson scattering angle. All other kinematic variables are taken to be identical to what we described above.

In figure 2 we show ratios of $1/\epsilon$ singularities in the squared amplitude $|\tilde{A}_{LLL}^{(d,2)}|^2$ computed directly and using Catani's formula. Deviations of this ratio from one signal numerical instabilities. We observe that $|\tilde{A}_{LLL}^{(d,2)}|^2/|\tilde{A}_{C,LLL}^{(d,2)}|^2 - 1$ is of order $\mathcal{O}(10^{-6} - 10^{-10})$ in the bulk of the phase-space. Significant instabilities are observed for backward scattering (178 degrees), where the transverse momentum is close to 1 GeV. However, the situation improves considerably already for 176 degree scattering where the transverse momentum



Figure 3. Absolute values squared of $\tilde{A}_{LLL}^{(d,2)}$ (left pane) and $\tilde{A}_{LLL}^{(n_g,2)}$ (right pane) as a function of the vector boson scattering angle. The center-of-mass energy, gauge boson masses and lepton scattering angles are given in the text.

is 2 GeV. The forward scattering limit appears to be more stable; even at two degrees, the $1/\epsilon$ contribution is computed properly to within a percent.

In the left pane of figure 3 we show the absolute value squared of the ratio of the finite part of the $\mathcal{A}_{LLL}^{(d,2)}$ amplitude and the leading order amplitude $\mathcal{A}_{LLL}^{(d,0)}$. To understand numerical accuracy of these results, we compared the output obtained with the double-precision version of the Fortran code with the Mathematica implementation. The advantage of the latter is that it provides a possibility to compute amplitudes with arbitrary numerical precision thereby ameliorating the problem of numerical instability. For backward scattering, we find that up to 174 degrees, the finite part is computed to within a per mille. For forward direction, the situation is similar but, perhaps, slightly better.

Finally, in the right pane of figure 3 we show absolute value squared of the ratio of the finite part of the left amplitude $A_{LLL}^{(n_g,2)}$ and the leading order LLL amplitude $\mathcal{A}_{LLL}^{(d,0)}$. Numerical instabilities are apparent for backward scattering. In fact, at 170 degrees ($p_{\perp} \sim$ 5 GeV), the agreement between double-precision Fortran code and the Mathematica code is about ten percent. In the forward direction, the situation is much better — the doubleprecision Fortran results agree with the results obtained using Mathematica implementation to better than a fraction of a percent for scattering angles as small as six degrees.

To summarize, while the two-loop amplitudes that we compute in this paper do exhibit numerical instabilities at small values of vector boson transverse momenta, we believe the stability is acceptable for phenomenological applications. Moreover, there are several ways to improve the situation. For example, it is possible to extend the Fortran code to provide results with quadruple precision. Note that computation of master integrals with arbitrary precision is feasible since Goncharov polylogarithms implementation in GiNaC does provide this functionality [71]. Moreover, it should also be possible to construct expansion of analytic expressions for scattering amplitudes that we obtained in this paper around singular limits, for example for forward or backward scattering and threshold production. If such expansions become available, computation of helicity amplitudes in singular limits will be significantly simplified. We leave these improvements for future work.

4 Conclusions

In this paper we described computation of two-loop scattering amplitudes for the annihilation of a quark and an antiquark into four leptons, that occurs through the production of two electroweak gauge bosons. The invariant masses of gauge bosons are kept arbitrary. We have given explicit formulas for projection operators that allow one to compute contributions of individual Feynman diagrams to invariant form factors. We use these form factors to construct helicity amplitudes for vector boson pair production processes including all off-shell effects and leptonic decays of vector bosons.

Results for two-loop scattering amplitudes obtained in this paper remove the last obstacle for computing the NNLO QCD corrections to the production of pairs of vector bosons with identical and different invariant masses. The two-loop virtual corrections that we compute in this paper will have to be combined with one-loop amplitudes for $pp \rightarrow V_1V_2j$ and with tree amplitudes for $pp \rightarrow V_1V_2jj$. While doing this consistently is non-trivial, the relevant technology is well-understood by now [43–51]. We hope, therefore, that results for NNLO fiducial volume cross sections for pair production of electroweak bosons can be expected in the near future.

Acknowledgments

J.M.H. and A.V.S. are grateful to Institute for Theoretical Particle Physics (TTP) at Karlsruhe Institute of Technology for the hospitality extended to them during the work on this paper. Research of K.M and F.C. was partially supported by US NSF under grant PHY-1214000. K.M. was also partially supported by Karlsruhe Institute of Technology through its distinguished researcher fellowship program. J.M.H. is supported in part by the DOE grant DE-SC0009988 and by the Marvin L. Goldberger fund. F.C. and A.S. are supported in part by DFG through SFB/TR 9. V.S. is supported in part by the Alexander von Humboldt Foundation (Humboldt Forschungspreis).

Open Access. This article is distributed under the terms of the Creative Commons Attribution License (CC-BY 4.0), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] ATLAS collaboration, Measurement of W^+W^- production in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector and limits on anomalous WWZ and WW γ couplings, Phys. Rev. D 87 (2013) 112001 [arXiv:1210.2979] [INSPIRE].
- [2] CMS collaboration, Measurement of WW production rate, CMS-PAS-SMP-12-005 (2012).
- [3] CMS collaboration, Measurement of WW production rate, CMS-PAS-SMP-12-013 (2012).
- [4] D. Curtin, P. Meade and P.-J. Tien, Natural SUSY in Plain Sight, arXiv:1406.0848 [INSPIRE].
- [5] D. Curtin, P. Jaiswal and P. Meade, *Charginos Hiding In Plain Sight*, *Phys. Rev.* D 87 (2013) 031701 [arXiv:1206.6888] [INSPIRE].

- [6] P. Meade, H. Ramani and M. Zeng, Transverse momentum resummation effects in W⁺W⁻ measurements, arXiv:1407.4481 [INSPIRE].
- [7] CMS collaboration, Measurement of the pp to ZZ production cross section and constraints on anomalous triple gauge couplings in four-lepton final states at $\sqrt{s} = 8 \text{ TeV}$, arXiv:1406.0113 [INSPIRE].
- [8] CMS collaboration, Measurement of the $W\gamma$ and $Z\gamma$ inclusive cross sections in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and limits on anomalous triple gauge boson couplings, *Phys. Rev.* **D** 89 (2014) 092005.
- [9] ATLAS collaboration, Measurement of $W\gamma$ and $Z\gamma$ production cross sections in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ and limits on anomalous triple gauge couplings with the ATLAS detector, Phys. Lett. **B** 717 (2012) 49 [arXiv:1205.2531] [INSPIRE].
- [10] P. Jaiswal and T. Okui, Explanation of the WW excess at the LHC by jet-veto resummation, Phys. Rev. D 90 (2014) 073009 [arXiv:1407.4537] [INSPIRE].
- [11] J. Ohnemus and J.F. Owens, An Order α⁻s calculation of hadronic ZZ production, Phys. Rev. D 43 (1991) 3626 [INSPIRE].
- [12] J. Ohnemus, An Order α^{-s} calculation of hadronic $W^{\pm}Z$ production, Phys. Rev. D 44 (1991) 3477 [INSPIRE].
- [13] J. Ohnemus, An Order α^{-s} calculation of hadronic $W^{-}W^{+}$ production, Phys. Rev. D 44 (1991) 1403 [INSPIRE].
- [14] B. Mele, P. Nason and G. Ridolfi, QCD radiative corrections to Z boson pair production in hadronic collisions, Nucl. Phys. B 357 (1991) 409 [INSPIRE].
- [15] S. Frixione, P. Nason and G. Ridolfi, Strong corrections to W Z production at hadron colliders, Nucl. Phys. B 383 (1992) 3 [INSPIRE].
- [16] S. Frixione, A Next-to-leading order calculation of the cross-section for the production of W⁺W⁻ pairs in hadronic collisions, Nucl. Phys. B 410 (1993) 280 [INSPIRE].
- [17] U. Baur, T. Han and J. Ohnemus, QCD corrections and nonstandard three vector boson couplings in W⁺W⁻ production at hadron colliders, Phys. Rev. D 53 (1996) 1098
 [hep-ph/9507336] [INSPIRE].
- [18] L.J. Dixon, Z. Kunszt and A. Signer, Vector boson pair production in hadronic collisions at order α_s: Lepton correlations and anomalous couplings, Phys. Rev. D 60 (1999) 114037 [hep-ph/9907305] [INSPIRE].
- [19] J.M. Campbell and R.K. Ellis, An Update on vector boson pair production at hadron colliders, Phys. Rev. D 60 (1999) 113006 [hep-ph/9905386] [INSPIRE].
- [20] J.M. Campbell, R.K. Ellis and C. Williams, Vector boson pair production at the LHC, JHEP 07 (2011) 018 [arXiv:1105.0020] [INSPIRE].
- [21] S. Frixione and B.R. Webber, Matching NLO QCD computations and parton shower simulations, JHEP 06 (2002) 029 [hep-ph/0204244] [INSPIRE].
- [22] P. Nason and G. Ridolfi, A Positive-weight next-to-leading-order Monte Carlo for Z pair hadroproduction, JHEP 08 (2006) 077 [hep-ph/0606275] [INSPIRE].
- [23] K. Hamilton, A positive-weight next-to-leading order simulation of weak boson pair production, JHEP 01 (2011) 009 [arXiv:1009.5391] [INSPIRE].

- [24] T. Melia, P. Nason, R. Rontsch and G. Zanderighi, W⁺W⁻, WZ and ZZ production in the POWHEG BOX, JHEP 11 (2011) 078 [arXiv:1107.5051] [INSPIRE].
- [25] R. Frederix et al., Four-lepton production at hadron colliders: aMC@NLO predictions with theoretical uncertainties, JHEP 02 (2012) 099 [arXiv:1110.4738] [INSPIRE].
- [26] F. Cascioli et al., Precise Higgs-background predictions: merging NLO QCD and squared quark-loop corrections to four-lepton + 0,1 jet production, JHEP 01 (2014) 046 [arXiv:1309.0500] [INSPIRE].
- [27] F. Campanario, M. Rauch and S. Sapeta, W⁺W⁻ production at high transverse momenta beyond NLO, Nucl. Phys. B 879 (2014) 65 [arXiv:1309.7293] [INSPIRE].
- [28] M. Billóni, S. Dittmaier, B. Jäger and C. Speckner, Next-to-leading order electroweak corrections to $pp \rightarrow W^+W^- \rightarrow 4$ leptons at the LHC in double-pole approximation, JHEP 12 (2013) 043 [arXiv:1310.1564] [INSPIRE].
- [29] A. Bierweiler, T. Kasprzik, J.H. Kühn and S. Uccirati, *Electroweak corrections to W-boson pair production at the LHC, JHEP* 11 (2012) 093 [arXiv:1208.3147] [INSPIRE].
- [30] A. Bierweiler, T. Kasprzik and J.H. Kühn, Vector-boson pair production at the LHC to $\mathcal{O}(\alpha^3)$ accuracy, JHEP **12** (2013) 071 [arXiv:1305.5402] [INSPIRE].
- [31] J. Baglio, L.D. Ninh and M.M. Weber, Massive gauge boson pair production at the LHC: a next-to-leading order story, Phys. Rev. D 88 (2013) 113005 [arXiv:1307.4331] [INSPIRE].
- [32] M. Grazzini, Soft-gluon effects in WW production at hadron colliders, JHEP 01 (2006) 095
 [hep-ph/0510337] [INSPIRE].
- [33] S. Dawson, I.M. Lewis and M. Zeng, Threshold resummed and approximate next-to-next-to-leading order results for W⁺W⁻ pair production at the LHC, Phys. Rev. D 88 (2013) 054028 [arXiv:1307.3249] [INSPIRE].
- [34] Y. Wang, C.S. Li, Z.L. Liu, D.Y. Shao and H.T. Li, Transverse-Momentum Resummation for Gauge Boson Pair Production at the Hadron Collider, Phys. Rev. D 88 (2013) 114017 [arXiv:1307.7520] [INSPIRE].
- [35] E.W.N. Glover and J.J. van der Bij, Vector boson pair production via gluon fusion, Phys. Lett. B 219 (1989) 488 [INSPIRE].
- [36] C. Kao and D.A. Dicus, Production of W^+W^- from gluon fusion, Phys. Rev. **D** 43 (1991) 1555 [INSPIRE].
- [37] T. Binoth, M. Ciccolini, N. Kauer and M. Krämer, Gluon-induced WW background to Higgs boson searches at the LHC, JHEP 03 (2005) 065 [hep-ph/0503094] [INSPIRE].
- [38] T. Binoth, M. Ciccolini, N. Kauer and M. Krämer, Gluon-induced W-boson pair production at the LHC, JHEP 12 (2006) 046 [hep-ph/0611170] [INSPIRE].
- [39] M. Grazzini, S. Kallweit, D. Rathlev and A. Torre, Zγ production at hadron colliders in NNLO QCD, Phys. Lett. B 731 (2014) 204 [arXiv:1309.7000] [INSPIRE].
- [40] F. Cascioli et al., ZZ production at hadron colliders in NNLO QCD, Phys. Lett. B 735 (2014) 311 [arXiv:1405.2219] [INSPIRE].
- [41] C. Anastasiou et al., NNLO QCD corrections to $pp \to \gamma^* \gamma^*$ in the large-N_F limit, arXiv:1408.4546 [INSPIRE].
- [42] T. Gehrmann et al., W⁺W⁻ production at hadron colliders in NNLO QCD, arXiv:1408.5243 [INSPIRE].

- [43] A. Gehrmann-De Ridder, T. Gehrmann and E.W.N. Glover, Antenna subtraction at NNLO, JHEP 09 (2005) 056 [hep-ph/0505111] [INSPIRE].
- [44] S. Catani and M. Grazzini, An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC, Phys. Rev. Lett. 98 (2007) 222002
 [hep-ph/0703012] [INSPIRE].
- [45] S. Catani, L. Cieri, D. de Florian, G. Ferrera and M. Grazzini, Universality of transverse-momentum resummation and hard factors at the NNLO, Nucl. Phys. B 881 (2014) 414 [arXiv:1311.1654] [INSPIRE].
- [46] S. Weinzierl, Subtraction terms at NNLO, JHEP 03 (2003) 062 [hep-ph/0302180] [INSPIRE].
- [47] G. Somogyi, P. Bolzoni and Z. Trócsányi, NNLO jet cross sections by subtraction, Nucl. Phys. Proc. Suppl. 205-206 (2010) 42 [arXiv:1007.4067] [INSPIRE].
- [48] M. Czakon, A novel subtraction scheme for double-real radiation at NNLO, Phys. Lett. B 693 (2010) 259 [arXiv:1005.0274] [INSPIRE].
- [49] M. Czakon, Double-real radiation in hadronic top quark pair production as a proof of a certain concept, Nucl. Phys. B 849 (2011) 250 [arXiv:1101.0642] [INSPIRE].
- [50] M. Czakon and D. Heymes, Four-dimensional formulation of the sector-improved residue subtraction scheme, arXiv:1408.2500 [INSPIRE].
- [51] R. Boughezal, K. Melnikov and F. Petriello, A subtraction scheme for NNLO computations, Phys. Rev. D 85 (2012) 034025 [arXiv:1111.7041] [INSPIRE].
- [52] T. Gehrmann and L. Tancredi, Two-loop QCD helicity amplitudes for $q\bar{q} \to W^{\pm}\gamma$ and $q\bar{q} \to Z^{0}\gamma$, JHEP **02** (2012) 004 [arXiv:1112.1531] [INSPIRE].
- [53] T. Gehrmann, L. Tancredi and E. Weihs, Two-loop QCD helicity amplitudes for $gg \rightarrow Zg$ and $gg \rightarrow Z\gamma$, JHEP 04 (2013) 101 [arXiv:1302.2630] [INSPIRE].
- [54] G. Chachamis, M. Czakon and D. Eiras, W Pair Production at the LHC. I. Two-loop Corrections in the High Energy Limit, JHEP 12 (2008) 003 [arXiv:0802.4028] [INSPIRE].
- [55] T. Gehrmann, A. von Manteuffel, L. Tancredi and E. Weihs, *The two-loop master integrals* for $q\bar{q} \rightarrow VV$, *JHEP* **06** (2014) 032 [arXiv:1404.4853] [INSPIRE].
- [56] T. Gehrmann, L. Tancredi and E. Weihs, Two-loop master integrals for $q\bar{q} \rightarrow VV$: the planar topologies, JHEP 08 (2013) 070 [arXiv:1306.6344] [INSPIRE].
- [57] F. Caola, J.M. Henn, K. Melnikov and V.A. Smirnov, Non-planar master integrals for the production of two off-shell vector bosons in collisions of massless partons, JHEP 09 (2014) 043 [arXiv:1404.5590] [INSPIRE].
- [58] J.M. Henn, K. Melnikov and V.A. Smirnov, Two-loop planar master integrals for the production of off-shell vector bosons in hadron collisions, JHEP 05 (2014) 090
 [arXiv:1402.7078] [INSPIRE].
- [59] R.J. Gonsalves, Dimensionally regularized two loop on-shell quark form-factor, Phys. Rev. D 28 (1983) 1542 [INSPIRE].
- [60] W.L. van Neerven, Dimensional Regularization of Mass and Infrared Singularities in Two Loop On-shell Vertex Functions, Nucl. Phys. B 268 (1986) 453 [INSPIRE].
- [61] G. Kramer and B. Lampe, Integrals for Two Loop Calculations in Massless QCD, J. Math. Phys. 28 (1987) 945 [INSPIRE].

- [62] F.V. Tkachov, A Theorem on Analytical Calculability of Four Loop Renormalization Group Functions, Phys. Lett. B 100 (1981) 65 [INSPIRE].
- [63] K.G. Chetyrkin and F.V. Tkachov, Integration by Parts: The Algorithm to Calculate β -functions in 4 Loops, Nucl. Phys. B 192 (1981) 159 [INSPIRE].
- [64] J.M. Henn, Multiloop integrals in dimensional regularization made simple, Phys. Rev. Lett. **110** (2013) 251601 [arXiv:1304.1806] [INSPIRE].
- [65] P. Nogueira, Automatic Feynman graph generation, J. Comput. Phys. **105** (1993) 279 [INSPIRE].
- [66] J. Kuipers, T. Ueda, J.A.M. Vermaseren and J. Vollinga, FORM version 4.0, Comput. Phys. Commun. 184 (2013) 1453 [arXiv:1203.6543] [INSPIRE].
- [67] A.V. Smirnov, Algorithm FIRE Feynman Integral REduction, JHEP 10 (2008) 107 [arXiv:0807.3243] [INSPIRE].
- [68] A.V. Smirnov and V.A. Smirnov, FIRE4, LiteRed and accompanying tools to solve integration by parts relations, Comput. Phys. Commun. 184 (2013) 2820 [arXiv:1302.5885] [INSPIRE].
- [69] A.V. Smirnov, FIRE5: a C++ implementation of Feynman Integral REduction, arXiv:1408.2372 [INSPIRE].
- [70] C.W. Bauer, A. Frink and R. Kreckel, Introduction to the GiNaC Framework for Symbolic Computation within the C++ Programming Language, J. Symb. Comput. 33 (2002) 1.
- [71] J. Vollinga and S. Weinzierl, Numerical evaluation of multiple polylogarithms, Comput. Phys. Commun. 167 (2005) 177 [hep-ph/0410259] [INSPIRE].
- S. Catani, The Singular behavior of QCD amplitudes at two loop order, Phys. Lett. B 427 (1998) 161 [hep-ph/9802439] [INSPIRE].
- [73] L.J. Dixon, Z. Kunszt and A. Signer, Helicity amplitudes for O(α_s) production of W⁺W⁻, W[±]Z, ZZ, W[±]γ, or Zγ pairs at hadron colliders, Nucl. Phys. B 531 (1998) 3
 [hep-ph/9803250] [INSPIRE].
- [74] N. Kauer and G. Passarino, Inadequacy of zero-width approximation for a light Higgs boson signal, JHEP 08 (2012) 116 [arXiv:1206.4803] [INSPIRE].
- [75] J.M. Campbell, R.K. Ellis and C. Williams, Bounding the Higgs width at the LHC using full analytic results for $gg \rightarrow e^-e^+\mu^-\mu^+$, JHEP **04** (2014) 060 [arXiv:1311.3589] [INSPIRE].