

**Title:** Sex differences in scent-marking in captive red-ruffed lemurs.

**Short running head:** Sex differences in scent-marking in red-ruffed lemurs.

**Authors:** Ellesse D. Janda<sup>1,2</sup>, Kate L. Perry<sup>2</sup>, Emma Hankinson<sup>3</sup>, David Walker<sup>2</sup> and Stefano Vaglio<sup>2,4</sup>

**Affiliations of authors:**

<sup>1</sup> Department of Social Sciences, Oxford Brookes University, Headington Campus, Headington Road, Oxford, OX3 0PB, United Kingdom

<sup>2</sup> Department of Biology, Chemistry and Forensic Science, University of Wolverhampton, City Campus South, Wulfruna Street, Wolverhampton, WV1 1LY, United Kingdom

<sup>3</sup> Department of Science and Technology, Bournemouth University, Talbot Campus, Fern Barrow, Poole, BH12 5BB, United Kingdom

<sup>4</sup> Department of Anthropology & Behaviour Ecology and Evolution Research (BEER) Centre, Science Site, South Road, Durham, DH1 3LE, United Kingdom

**Corresponding author:** Stefano Vaglio [S.Vaglio@wlv.ac.uk](mailto:S.Vaglio@wlv.ac.uk)

**Word count:** 3,804

## Abstract

Primate chemical communication remains underappreciated, as primates are considered to rely on other sensory modalities. However, various lines of evidence suggest that olfaction plays an important role in primate societies, including the conspicuous scent-marking behavior of many strepsirrhines and callitrichines. Although lemurs typically show scent-marking, little is known about this behavior in red-ruffed lemurs (*Varecia variegata rubra*). We combined behavioral observations and semiochemistry analyses to improve our understanding of scent-marking in two captive troops housed at Dudley and Twycross zoos (UK). We collected olfactory behavioral observations by focusing on two family troops ( $N=7$ ) for 132 h. We investigated the volatile compounds of ano-genital scent-marks using solid-phase microextraction and gas chromatography-mass spectrometry and compared volatile chemical profiles with features of the signaller. Males scent-marked most frequently and predominantly in specific meaningful areas of the enclosure, while within females the occurrence of scent-marking was related to their age. We found behavioral sexual dimorphism, with male predominantly depositing secretions via neck and mandible glands and females via ano-genital glands. We identified a total of 32 volatile components of ano-genital gland secretion, including compounds that have already been found in other mammals as sex pheromones and cues to fitness, in ano-genital scent-marks spontaneously left on filter paper by adult females. Our findings suggest that red-ruffed lemurs might use scent-marking to convey information about sex and female age, with male neck-marking behavior playing defensive territorial functions and ano-genital marking related to socio-sexual communication.

**Key-words:** Communication, Signalling, Olfaction, Gas Chromatography–Mass Spectrometry, *Varecia variegata rubra*

## Introduction

Communication plays a fundamental role within animal societies, especially for species displaying complex social systems. The ultimate goal of communication is to spread information that influences the behaviors of receivers (Seyfarth and Cheney 2003). Animals can use various sensory modalities to transfer their messages to other individuals. In particular, olfactory communication is exhibited by several mammal species (reviewed in Scordato and Drea 2007), such as rodents (*e.g.*, Hurst *et al* 1998; Roberts 2007), but also by reptiles (*e.g.*, Muller-Schwarze 2006) and birds (*e.g.*, Leclaire *et al* 2017). Odor secretions are directly linked to the physiological conditions of senders (Harris *et al* 2018) and can be costly to produce (Scordato and Drea 2007), thus they are expected to deliver a more honest signal compared to other forms of communication (Hasson 1997).

Primates have traditionally been considered to be “microsmatic”, relying more upon other sensory modalities than olfaction (Dulac and Torello 2003). Since vision and acoustics are considered to be the main sensory modalities in most primate species little is known about the chemical signals used by non-human primates (Walker 1998). However, several studies support the hypothesis that chemical communication is crucial also for primates (*e.g.*, Porter and Moore 1971; Geissman and Hultegger 1994; Wedekind *et al* 1995; Wedekind and Furi 1997; Smith *et al* 2001; Jacob *et al* 2002; Hayes *et al* 2004, 2006; Heymann 2006; Knapp *et al* 2006; Laidre 2009; Smith 2006; Scordato *et al* 2007; Setchell *et al* 2010, 2011; Vaglio *et al* 2016). Particularly, it is established that some primates also rely heavily on olfaction in addition to vision and auditory senses; for instance, this is the case of squirrel monkeys (Laska *et al* 2000) and several lemurs (Gould and Overdorff, 2002; Scordato and Drea 2007). Strepsirrhines have retained their olfactory complexity (reviewed in Hayes *et al* 2006) due to morphological constraints that limit the visual signals produced by senders (Scordato and Drea 2007).

Lemur behavioral repertoire comprises both olfactory investigative and scent-marking behaviors. Lemurs show both direct and indirect olfactory investigative behaviors (Drea 2015); direct investigations may include behaviors such as sniffing and/or licking a conspecific's skin (palms, soles, eyelids, or nostrils) or genitals, and self-sniffing genitals, while indirect behaviors consist of sniffing and/or licking scent-marks deposited by the signaller. Scent-marking behavior is shown by several terrestrial vertebrates, including mammal and reptile species (Müller-Schwarze 2006; Roberts 2007). Scent-marks may include species-wide pheromones (*i.e.*, chemical substances released by an animal or insect which can affect a conspecific individual; for further details see Vaglio *et al* 2018) as well as highly individual odors. Scent-marking is a very effective form of communication within habitats that make difficult the detection of visual and auditory signals (Gould and Overdorff 2002), which is the case with forests inhabited by lemurs (Sussman *et al* 2003). In particular, this behavior is reasonably common in lemurs and New World monkeys – among which may play several functions, including the reproductive suppression of subordinate females, advertisement of individual “quality”, preparing males to assist in the delivery and care of newborn infants, and territorial defence (*e.g.*, Gould and Overdorff 2002; Pochron *et al* 2005; Heyman 2006) – while is less commonly reported in Old World monkeys and apes (*e.g.*, Freeman *et al* 2012). Especially, among strepsirrhines, social complexity may have selected for olfactory complexity in lemurs (delBarco-Trillo *et al* 2012).

Mammals have a common pattern of scent-marking: glandular secretions, if not feces or urine, are placed at meaningful places such as along paths and territorial boundaries (Gosling and Roberts 2001). Scent-glands have been observed in various lemur species, including all *Eulemur* species (delBarco-Trillo *et al* 2012), ring-tailed lemurs (Scordato and Drea 2007), red-bellied lemurs (Gould and Overdorff 2002), red-fronted lemurs (Hayes *et al* 2006), Milne-Edward's sifakas (Hayes *et al* 2004), black-and-white and red-ruffed lemurs (Gould

and Overdorff 2002). In particular, red-ruffed lemurs have multiple scent-glands (Gould and Overdorff 2002), composed of neck and mandible glands (male), and anogenital glands (male and female) (Pereira *et al* 1988); indicating that olfactory communication should be significant for this species (Elisa *et al* 2004).

The red-ruffed lemur is a large, frugivorous lemur species (Vasey 2006), which inhabit the residual primary forests of the Masoala Peninsula (Andriaholinirina *et al* 2014). Red-ruffed lemurs have a variable social system; in smaller home ranges their group size is usually between 2-5 individuals, whereas larger home ranges have been known to support between 18-32 individuals (Rigamonti 1993). Although red-ruffed lemur communities are not cohesive units, the home range is communally defended. In addition, only females participate in communal home range defense against females from other groups, which includes agonistic behaviors such as chasing, scent-marking, vocalizing, and even physical contact with members of neighboring communities (Vasey 2005; 2007). Females are dominant to males, winning almost all agonistic encounters with them and rarely showing submissive behavior towards them (Raps and White 1995; Meyer *et al* 1999). Communication is commonly observed as vocalisations, emitting species-specific calls which serve several functions and are transmittable between groups (Macedonia and Taylor 1985); however, also chemical communication is thought to play a crucial role in group dynamics (Elisa *et al* 2004).

The overarching aim of this study is to improve our understanding of the role played by chemical communication, particularly focusing on scent-marking behavior, in red-ruffed lemurs. We predict that red-ruffed lemurs advertise information about their sex, age and rank by using scent-marking. We also anticipate that this study may contribute to further exploring the connection between functional and mechanistic levels of lemur scent-marking (Charpenter *et al* 2010).

119

## 120 **Materials and Methods**

121

### 122 *Subjects and Housing*

123         We studied two captive troops of red-ruffed lemurs ( $n=7$ ) housed at Dudley and  
124 Twycross zoos (UK). The troop housed at Dudley Zoological Gardens consisted of two  
125 related (brothers) adult males (13 years old) and one unrelated adult female (12 years old).  
126 The troop housed at Twycross Zoo consisted of one adult male (11 years old), one adult  
127 female (12 years old) and their offspring (two 1.5 years old females). Red-ruffed lemurs are  
128 considered sexually mature at 2 years old, with first conception approximately one year later  
129 (Vasey 2007). Adult females were contracepted, and all individuals in non-breeding season  
130 (*i.e.* regarding red-ruffed lemurs in captivity in the Northern Hemisphere breeding usually  
131 occurs in December-January with births in April-May; Brockman *et al* 1987).

132         We carried out behavioral observations and odor sampling from September to  
133 November 2016 (Twycross Zoo) and from July to September 2018 (Dudley Zoological  
134 Gardens). In both institutions, the troops lived in an indoor enclosure (heated to 28°C) with  
135 access to an outdoor enclosure ('visitor walktrough' enclosures).

136

### 137 *Ethics Statement*

138         This study followed the guidelines for the care and use of captive animals in the  
139 UK, involving non-invasive methods for obtaining both behavioral data and odor samples  
140 from red-ruffed lemurs. Moreover, the study was conducted in compliance with the  
141 Convention on International Trade in Endangered Species of Wild Fauna and Flora (CITES)

and approved by the Life Sciences Ethics committee at the University of Wolverhampton (UK) and the Ethics committees at Dudley Zoological Gardens and Twycross Zoo (UK).

#### *Behavioral Data Collection and Analysis*

We collected behavioral data by instantaneous scan sampling (Altmann 1974), with behaviors recorded at 30-sec intervals over the duration of an hour in three time periods (two during the morning, and one during the afternoon), two days per week, over three months. Alongside the scan sampling we also used *ad libitum* sampling for recording olfactory behaviors (**Table I**), including both scent-marking (ano-genital marking, neck-marking – that is scent-marking via neck and mandible glands) behavior and locations (comprising of “hatches”, “tree branch”, “indoor enclosure”, “wooden hut”, and “climbing frame”). We recorded a total of 132 hours of observations throughout the study period, including 360 scan samples each sampling day on the entire group.

We investigated the relationships between individuals and scent-marking behavior in relation to sex and age of senders. We also investigated the role played by different types of scent-marking behavior, and locations of scent-marks within the enclosure. We tested all variables through Shapiro-Wilk test for normality. As data were not normally distributed, we performed non-parametric Kruskal-Wallis tests followed by pairwise Mann-Whitney U *post-hoc* tests. All tests were carried out using SPSS v.23, and a significance level of  $P < 0.05$  was applied.

[insert Table I here]

#### *Odor Sampling and Analysis*

We collected odor secretions spontaneously released via scent-marking by red-ruffed lemurs on brand-new filter paper fixed on hatches, climbing equipment, and tree trunks and branches (**Figure I**). Unfortunately, we were not able to collect any odor sample from males, while we collected scent-marks deriving from ano-genital marking by all the females (14 samples, 3-4 replicates per individual). In addition, we placed control filter paper in the environment to control for the contact with wood (where there may be chemical compounds deriving from the wood, but also algae, microorganisms, etc.) and we exposed control filter paper also to the air during sampling in order to detect any chemical compounds which did not derive from the red-ruffed lemurs. We collected odor samples immediately after scent-mark deposition by red-ruffed lemurs in the outdoor enclosure. We placed all samples and controls into brand-new sterile vials (Supelco) and immediately stored them at -20°C. We used 10-ml screw-capped clear glass vials (thread: 18O.D. 22.5-mm x H 46-mm) closed by teflon-faced rubber septa and seals (1.3-mm thick).

We conducted laboratory analyses at the Rosalind Franklin Science Centre, University of Wolverhampton (UK). We investigated the volatile components of odor secretions using established solid-phase microextraction (SPME) and gas chromatography-mass spectrometry (GC-MS) and applying the same methods used in our previous work on mandrill odor signals (Setchell *et al* 2010; Vaglio *et al* 2016).

We introduced a 65- $\mu$ m polydimethylsiloxane/divinylbenzene SPME syringe needle through the vial septum and then we exposed the fibre to the headspace above the sample in the vial for 15 minutes at 40°C. We analysed the adsorbed volatile analytes of all samples by using a 5975C mass spectrometer (Agilent Technologies, Santa Clara, CA, USA) EI, 70 eV, coupled directly to a 7890B gas chromatograph (Agilent Technologies, Santa Clara, CA, USA) equipped with a fused silica HP5-MS UI capillary column (Agilent Technologies,



Santa Clara, CA, USA) 30 m x 0.25 mm crossbonded 5%-phenyl-95%-dimethylpolysiloxane, film thickness 0.25  $\mu$ m. We maintained the injector and transfer line temperatures at 270°C and 280°C, respectively. We made injections in splitless mode (purge valve opened after 1 min) with a constant flow of helium carrier gas of 1 mL min<sup>-1</sup>. We started the oven temperature program at 45°C for 2 min, then raised it by 4°C min<sup>-1</sup> to 170°C, and finally by 20°C min<sup>-1</sup> to 300°C.

We assessed potential contamination due to the lab environment through blank analyses of an empty 10-mL vial (Supelco) following the same procedure as for the samples. In addition, we conditioned the fibre at 260 °C pre- and post- injection, for 5 and 20 mins respectively in order to avoid any possible carry-over effects.

We tentatively identified eluted compounds by comparing the experimental spectra with the spectra provided by the mass-spectral library in ChemStation (Agilent Technologies, Santa Clara, CA, USA) and NIST (National Institute of Standards and Technology) Database, version MSD F.01.01.2317 (Agilent Technologies, Santa Clara, CA, USA). We accepted a putative identification when the minimum matching factor was higher than 80%. If more than one compound was a good match for the same GC peak then we considered the chromatographic retention time and compared it with the retention time reported in the literature for the same chromatographic column type (El-Sayed 2016) in order to minimize the chance of misidentification. We created a data matrix using the peak area relative to each identified compound by using the integrated signal of the deconvoluted total ion current (TIC). We analysed all samples in a short period of time (approximately 24 hours) to minimize interassay variability. We removed all the contaminants (*i.e.*, any compounds that appeared in the ‘environmental controls’ and ‘lab blanks’) from the scent-mark results.

[Insert Figure I here: Filter paper attachments]

## Results

### *Behavioral Observations*

During the study period olfactory behaviors were exhibited predominantly by males (40.00%), followed by adult females (34.48%) and subadult females (25.52%). These behaviors included both scent-marking (ano-genital marking and neck-marking) and investigative behaviors (sniffing and/or licking an area within the enclosure, sniffing and/or licking a conspecific, self-licking of the ano-genital glands). Scent-marks were most commonly deposited by males (51.55%), followed by adult females (26.80%) and subadult females (21.65%) (**Figure II**), although differences were not significant between the sexes ( $U=137.5$ ;  $p=0.688$ ).

[Insert Figure II here: Frequency of marks in relation to sex and age]

We observed males ( $N=3$ ) scent-marking significantly more via neck glands rather than via ano-genital glands ( $U=12.5$ ;  $p<0.001$ ). Females ( $N=4$ ) displayed the opposite behavioral pattern; showing a significantly higher frequency of ano-genital marking rather than neck marking ( $U=41.5$ ;  $p=0.022$ ). We found significant differences in neck-marking behavior between individuals, and particularly between males and females ( $N=7$ ;  $U=12.5$ ;  $p=0.003$ ).

We found a significant difference ( $N=7$ ;  $Z = -5.675$ ,  $p<0.001$ ) in scent-marking behaviors between inside and outside locations, with 81.73% of scent-marks occurring in the

outdoor enclosure. Moreover, scent-marks were most commonly deposited near, or upon, the hatches leading to the indoor and off-show enclosure (18.27%). However, scent-marks were also deposited on tree branches next to the path of the walk-through, all climbing frames in the walk-through, a hut providing shelter within the walk-through, and upon furniture in the indoor enclosure.

We also found a preference of location when considering the type of scent-marking performed; “hatches” were used most commonly overall for neck-marking (81.82%) compared to ano-genital marking (18.18%), whereas “climbing frame” was used more for ano-genital marking (71.43%) than neck-marking (28.57%).

We found significant differences in areas used for neck-marking, most commonly upon “hatches” ( $X^2=23.152$ ;  $p<0.001$ ), and upon “tree branches” ( $X^2=9.456$ ;  $p=0.009$ ). Deposition of neck scent-marks upon “hatches” was significantly different between males and females ( $U=04.5$ ;  $p<0.001$ ). Neck-marking on “tree branches” showed a difference between males and females ( $U=30.0$ ;  $p=0.029$ ), and between adult and subadult females ( $U=30.0$ ;  $p=0.029$ ).

We found significant differences in areas used for ano-genital marking, with most frequent occurrences upon “hatches” ( $X^2=11.748$ ;  $p=0.003$ ) and “climbing frame” ( $X^2=13.119$ ;  $p<0.001$ ). Deposition of ano-genital marks upon “hatches” was significantly different between adult and subadult females ( $U=84.0$ ;  $p=0.037$ ). Ano-genital marks upon “climbing frames” were also significantly different between adult and subadult females ( $U=35.0$ ;  $p<0.001$ ).

*Odor secretions*

We identified a total of 32 individual compounds from the analysis of 14 filter paper samples of female ano-genital secretions. These compounds included a range of naturally occurring odorous volatile compounds such as hydrocarbons, terpenes, terpene alcohols and ketones. Tentative identifications are listed in **table II**, while typical chromatograms (1 from the blank control and 1 from a female lemur ano-genital scent-mark) are shown in **figure III**. There was variation in the number and abundance of the compounds observed from sample to sample across different individuals. However, six compounds (benzaldehyde, 2-ethyl-1-hexanol, p-cresol, cis-p-mentha-2,8-dien-1-ol, 2-pinen-4-one, pentadecane) were present in all samples. We were not able to analyse the ratios of individual components in order to compare the volatile profiles with features of the signaller (for instance, adult vs subadult females) due to the small amount of filter paper samples.

*[Insert Table II here. Secretion compounds]*

*[Insert Figure III here. Chromatographs]*

## **Discussion**

Primates rely on olfactory communication in several contexts, including foraging, territorial defense, individual and family recognition, mate choice and mother-offspring bonding (Zeller 1987). Although very little is known about Old World primates, research has been accumulating on chemical communication in strepsirrhines and New World monkeys; particularly, semiochemical data are accessible for few non-human primate species, including various strepsirrhines [galago (Crewe et al 1979); lemurs (delBarco-Trillo *et al* 2011, 2012; Hayes *et al* 2004, 2006; Palagi and Dapporto 2006; Scordato *et al* 2007), owl monkeys (Macdonald *et al* 2008), marmosets and tamarins (Epplé *et al* 1993; Smith *et al* 2001),

macaques (Curtis *et al* 1971) and mandrills (Setchell *et al* 2010, 2011; Vaglio *et al* 2016)]. In this study we focused on scent-marking behavior, via both behavioral and chemical approaches, in two troops of zoo-managed red-ruffed lemurs.

Red-ruffed lemurs, as the well-studied ring-tailed lemurs, are characterised by a female-dominated society. In ring-tailed lemurs males scent-mark more than females (Pochron *et al* 2005) and female age is positively correlated to scent-marking frequency (Kappeler 1990; Gould and Overdorff 2002; Pochron *et al* 2005). Similarly, in our study, males scent-marked significantly more than any female and adult females showed the highest frequency of scent-marking within females. Therefore, our findings support the prediction that scent-marking would differ between individuals based on their sex, as found in other lemur species such as sifakas (Pochron *et al* 2005), and age, as found in ring-tailed lemurs (Kappeler 1990; Gould and Overdorff 2002; Pochron *et al* 2005).

Neck-marking was exhibited by all study subjects, but significantly more by males. In contrast, females exhibited ano-genital scent-marking significantly more than males. This supports the hypothesis of sexual dimorphism in red-ruffed lemur scent-marking, as already observed by Vasey (2003). The preferences shown by males for neck-marking “hatches” and ano-genital marking “tree branches”, and by adult females for ano-genital marking “climbing frames”, confirm behavioral sexual dimorphism. These observable preferences also suggest that scent-marking behaviors might play different roles in males and females, as observed in other primates, including ring-tailed lemurs (Scordato and Drea 2007), black-and-white ruffed lemurs (Pereira *et al* 1988) and mandrills (Vaglio *et al* 2016).

Our results also support the hypothesis that scent-marking might have a territorial function in this species (Pereira *et al* 1988). In particular, males scent-marked mostly specific meaningful places, by using neck-marking for hatches (small openings allowing access from

outdoor to indoor enclosure; *i.e.*, potential role of territorial defense) and ano-genital marking for tree branches and climbing equipment (areas of frequent transit by individuals; *i.e.*, potential role of social communication). This also implies that scent-marks released via neck-marking and ano-genital marking might communicate different messages to the receivers by conveying information about distinct features of the senders. Previous studies have indicated information regarding sex to be conveyed in scent-marks from ring-tailed lemurs (Hayes *et al* 2004, Scordato *et al* 2007), but absent in odorants from sifakas (Hayes *et al* 2004, 2006).

Although scent-marking behavior is observable, it is challenging to decipher the message which is chemically communicated. Therefore, the chemical investigation of odor secretions released by scent-marking is critical to understand the message transferred by this behavior. Since we used spontaneously released scent-marks, we were able to investigate odor secretions released by scent-marking and thus corresponding to the exact message sent by red-ruffed lemurs.

A total of 32 compounds were tentatively identified within the ano-genital secretions released by female study subjects (excluding environmental and lab contaminants as well as co-eluted compounds). This low amount of volatile compounds in comparison to other female lemur ano-genital marks (for example, ring-tailed lemurs and sifakas; Hayes *et al* 2004, Scordato *et al* 2007) might be explained by the fact that breeding *versus* non-breeding season (Scordato & Drea 2007) and chemical contraception (Crawford *et al* 2011) can have significant impacts on semiochemical signals in lemurs. For instance, in ring-tailed lemurs (Crawford *et al* 2011) genital odorants of adult contracepted females were proved altered, including decreased richness, modified relative abundances, and minimized individual chemical distinctiveness of their volatile chemical profiles.

332 Volatile hydrocarbons have previously been identified in odorants deriving from ring-  
333 tailed lemurs and sifakas (Hayes *et al.* 2004; Scordato *et al.* 2007) as well as from Old World  
334 monkeys such as mandrills (Setchell *et al.* 2010; Vaglio *et al.* 2016) and olive baboons (Vaglio  
335 *et al.* in preparation). In particular, high-molecular weight volatile hydrocarbons might act as  
336 a fixative which slows the release of more volatile compounds, as suggested for major  
337 urinary proteins in mice (Green 2016; Hurst *et al.* 1998). The compounds benzaldehyde, p-  
338 cresol (also known as p-methylphenol), hexanal and geranylacetone are commonly  
339 encountered in mammal scent markings (*e.g.*, lions, wild dogs, wolves, mice, red foxes)  
340 ([Osada](#) *et al.* 2015; Roberts *et al.* 2010; Soso & Koziel 2016). The compound benzaldehyde  
341 has already been found in gland secretions released by marmosets (Smith *et al.* 2001), and  
342 functions as sex pheromone in other mammals (reviewed in El-Sayed 2016) and also as cue  
343 to genetic quality (reviewed in Wyatt 2014) in other vertebrates (*e.g.*, in the crested auklet, a  
344 seabird with citrus scent based on decanal and octanal produced by both sexes during the  
345 breeding season, concentration correlates with rank in males). Thus, benzaldehyde might  
346 serve as pheromone and play a role in signalling individual quality also in red-ruffed lemurs.  
347 In addition, ethyl-phenol occurs in rat urine as mate attraction signal and also in beaver urine  
348 as part of a multicomponent signal of range occupation (reviewed in Apps *et al.* 2015). The  
349 compound 2-pine-4-one (also known as verbenone) is a bark beetle antiaggregation  
350 pheromone (Lindgren & Miller 2002), which has similarly been found in other insects (*i.e.*,  
351 bees and butterflies) and is also naturally occurring in plants (reviewed in Bakthavatsalam  
352 2016). 2-ethyl-1-hexanol and cis-p-metha-2,8-dien-1-ol both appear to be associated with  
353 fragrances. Finally, other compounds, such as  $\alpha$ -pinene, are known to derive from plants;  
354 therefore, they might be a by-product and potentially vary with the environmental context but  
355 could also contribute to the message communicated by red-ruffed lemurs through scent-  
356 marking (for instance, convey information about group identity).

## Conclusions

In conclusion, the present study supports the hypotheses (Smith *et al* 2015) of sexual dimorphism and of more than one function served by scent-marking in red-ruffed lemurs. We suggest that scent marking could serve a function in intergroup spacing and intrasexual competition for both sexes, as might be expected in a female-dominant species.

In particular, male neck-marking might have a defensive territorial function while ano-genital marking might play a role in socio-sexual communication in this lemur species. Furthermore, our findings suggest that odor secretions released via ano-genital-marking might convey information about the age of female signallers. Additionally, the similarity of red-ruffed lemur's volatile chemical profiles to those found in other vertebrates would support our previous suggestion (Setchell *et al* 2010) that non-human primates are not as microsmatic as traditionally considered.

Since this study is based on seven animals living in two captive family troops it can only be considered a preliminary work for the red-ruffed lemur species. Future research work should focus on a larger sample size, record behaviors consistently throughout the day, and investigate the odor secretions released by adult non-contracepted females and also by male scent-marks. In addition, it would be crucial to study the perception by the recipient, for instance looking for evidence of behavioral or physiological responses facilitated by scent-marks via bioassay tests (Wyatt 2014). Also, more detailed analysis of the ratios of individual components could form the basis of further studies. Finally, although we focused on the volatile profile of red-ruffed lemur odor, we also recognize the significance of non-volatile components of odor secretions, as high-molecular weight compounds may extend the persistence of volatile signals in scent-marks (Alborne 1984; Belcher *et al* 1990; Hurst and



Beynon 2004).

## **Funding**

This research work was supported by the Faculty of Science and Engineering, University of Wolverhampton (sampling & laboratory analysis), and the Nuffield Foundation (Nuffield Research Placement for KLP).

## **Acknowledgements**

We are grateful to Dudley Zoological Gardens (especially David Beeston, Chris Leeson, Pat Stevens, and Lemur Wood keepers) and Twycross Zoo (especially Mat Liptovszky, Manuela Townsend, and Zak Showell) for their support to the project and assistance with sample collection. We also thank Keith Holding for his assistance with GC-MS chemical analyses at the Rosalind Franklin Science Centre, Wolverhampton, and Clare Everson and Farzana Aslam (Coventry University) for facilitating the Nuffield Research Placement of KLP. Furthermore, we would like to thank Tim Baldwin for his invaluable suggestions. On top of that, we also thank two anonymous reviewers for constructive comments.

## **Conflicts of interest**

The authors declare that they have no conflict of interest.

## **References**

403 Andriaholinirina, N., Baden, A., Blanco, M., Chikhi, L., Cooke, A., Davies, N., ...  
 404 Zaramody, A. (2014). *Varecia rubra*. *The IUCN Red List of Threatened Species* IUCN.  
 405 Available at: <e.T22920A16121712: <http://www.iucnredlist.org/details/22920/0>>.  
  
 406 Alborne, E. S. (1984). *Mammalian semiochemistry*. New York: John Wiley & Sons.  
 407 Altmann, J. (1974). Observational study of behaviour: Sampling methods. *Behaviour*, 49,  
 408 227-267. DOI: 10.1163/156853974X00534  
  
 409 Apps, P. J., Weldon, P. J., & Kramer, M. (2015) Chemical signals in terrestrial vertebrates:  
 410 search for design features. *Natural Product Reports*, 32, 1131-1153. DOI:  
 411 10.1039/C5NP00029G  
  
 412 Bakthavatsalam, N. (2016) Semiochemicals. In: Omkar, K. (editor). *Ecofriendly Pest*  
 413 *Management for Food Security*. New York: Academic Press.  
  
 414 Belcher, A., Epple, G., Greenfield, K. L., Richards, L. E., Kuderling, I., & Smith, A. B.  
 415 (1990). Proteins-biologically relevant components of the scent marks of a primate (*Saguinus*  
 416 *fuscicollis*). *Chemical Senses*, 15, 431-446. DOI: 10.1093/chemse/15.4.431  
  
 417 Brockman, D. K., Willis, M. S., & Karesh, W. B. (1987) Management and husbandry of  
 418 ruffed lemurs, *Varecia variegata*, at the San Diego Zoo. I. Captive population, San Diego  
 419 Zoo housing and diet. *Zoo Biology*, 6, 341-347. DOI: 10.1002/zoo.1430060408  
  
 420 Crawford, J. C., Boulet, M., & Drea, C. (2011) Smelling wrong: hormonal contraception in  
 421 lemurs alters critical female odour cues. *Proceedings of Royal Society of London B*, 278, 122-  
 422 130. DOI: 10.1098/rspb.2010.1203

423 Crewe, R. M., Burger, B. V., Le Roux, M., & Katsir, Z. (1979). Chemical constituents of the  
 424 chest gland secretion of the thick-tailed galago (*Galago crassicaudatus*). *Journal of Chemical*  
 425 *Ecology*, 5, 861-868. DOI: 10.1007/BF00986569

426 Curtis, R. F., Ballantine, J. A., Keverne, E. B., Bonsall, R. W., & Michael R. P. (1971).  
 427 Identification of primate sexual pheromones and properties of synthetic attractants. *Nature*,  
 428 232, 396-398. DOI: 10.1038/232396a0

429 Dapporto, L. (2008). The asymmetric scent: Ringtailed lemurs (*Lemur catta*) have distinct  
 430 chemical signatures in left and right brachial glands. *Die Naturwissenschaften*, 95, 987-991.  
 431 DOI: 10.1007/s00114-008-0407-7

432 delBarco-Trillo, J., Burkert, B. A., Goodwin, T. E., & Drea, C. M. (2011). Night and day: the  
 433 comparative study of strepsirrhine primates reveals socioecological and phylogenetic patterns  
 434 in olfactory signals. *Journal of Evolutionary Biology*, 24, 82-98. DOI: 10.1111/j.1420-  
 435 9101.2010.02145.x

436 delBarco-Trillo, J., Sacha, C., Dubay, G., & Drea, C. M. (2012). *Eulemur*, me lemur: The  
 437 evolution of scent-signal complexity in a primate clade. *Philosophical Transactions of the*  
 438 *Royal Society B*, 367, 1909-1922. DOI: 10.1098/rstb.2011.0225

439 Drea, C. M. (2015). D'scent of man: A comparative survey of primate chemosignaling in  
 440 relation to sex. *Hormones and Behavior*, 68, 117-133. DOI: 10.1016/j.yhbeh.2014.08.001

441 Dulac, C., & Torello, A. T. (2003). Molecular detection of pheromone signals in mammals:  
 442 From genes to behaviour. *Nature Reviews Neuroscience*, 4, 551-562. DOI: 10.1038/nrn1140

443 Elisa, U., Bracchi, P., & Federico, B. (2004). Captive Bred Lemur Behaviour and  
 444 Endocrinology. *Annali della Facolta` di Medicina Veterinaria di Parma*, 24, 193-202.

445 El-Sayed, A. (2016). *The pherobase: Database of pheromones and semiochemicals*.  
 446 Available at: <[www.pherobase.com](http://www.pherobase.com)>.

447 Epple, G., Belcher, A. M., Kuderling, I., Zeller, U., Scolnick, L., Greenfield, K. L., & Smith  
 448 A. B. I. (1993). Making sense out of scents: species differences in scent glands, scent-  
 449 marking behaviour, and scent-mark composition in the Callitrichidae. In: Rylands, A. B.  
 450 (editors). *Marmosets and Tamarins: Systematics, Behaviour, and Ecology*. Oxford: Oxford  
 451 University Press.

452 Freeman, N. J., Pasternak, G. M., Rubi, T. L., Barrett, L., & Henzi, S. P. (2012). Evidence for  
 453 scent marking in vervet monkeys? *Primates*, 53, 311-315. DOI: 10.1007/s10329-012-0304-8

454 Geissman, T., & Hultegger, A. M. (1994). Olfactory communication in gibbons? In: Roeder  
 455 J. J., Thierry, B., Anderson, J. R., & Herrenschildt, N. (editors). *Current primatology: Social*  
 456 *development, learning and behaviour*. Strasbourg: Université Louis Pasteur Press.

457 Gosling, L. M., & Roberts, S. C. (2001). Scent-marking by male mammals: cheat-proof  
 458 signals to competitors and mates. *Advances in the Study of Behavior*, 30, 169-217. DOI:  
 459 10.1016/S0065-3454(01)80007-3

460 Gould, L., & Overdorff, D. (2002). Adult male scent-marking in *Lemur catta* and *Eulemur*  
 461 *fulvus rufus*. *International Journal of Primatology*, 23, 575-596. DOI:  
 462 10.1023/A:1014921701106

463 Greene, L. K., Grogan, K. E., Smyth, K. N., Adams, C. A., Klager, S. A., & Drea, C. M.  
 464 (2016) Mix it and fix it: functions of composite olfactory signals in ring-tailed lemurs. *Royal*  
 465 *Society Open Science*, 3(4):160076. DOI: 10.1098/rsos.160076

466 Hayes, R., Morelli, T., & Wright, P. (2004). Anogenital gland secretions of *Lemur catta* and  
 467 *Propithecus verreauxi coquereli*: A preliminary chemical examination. *American Journal of*  
 468 *Primatology*, 63, 49-62. DOI: 10.1002/ajp.20038

469 Hayes, R., Morelli, T., & Wright, P. (2006). Volatile components of lemur scent secretions  
 470 vary throughout the year. *American Journal of Primatology*, 68, 1202-1207. DOI:  
 471 10.1002/ajp.20319

472 Harris, R. L. Boulet, M., Grogan, K. E., & Drea C. (2018) Costs of injury for scent signalling  
 473 in a strepsirrhine primate. *Scientific Reports*, 8: 9882. DOI: 10.1038/s41598-018-27322-3

474 Hasson, O. (1997). Towards a general theory of biological signalling. *Journal of Theoretical*  
 475 *Biology*, 185, 139-156. DOI: 10.1006/jtbi.1996.0258

476 Heymann, E. W. (2006). Scent marking strategies of New World primates. *American Journal*  
 477 *of Primatology*, 68, 650-661. DOI: 10.1002/ajp.20258

478 Hurst, J. L., & Beynon, R. J. (2004). Scent wars: the chemo-biology of competitive signalling  
 479 in mice. *Bioessays*. 26, 1288-1298. DOI: 10.1002/bies.20147

480 Hurst, J. L., Robertson, D., Tolladay, U., & Beynon, J. (1998). Proteins in urine scent marks  
 481 of male house mice extend the longevity of olfactory signals. *Animal Behavior*, 55, 1289-  
 482 1297. DOI: 10.1006/anbe.1997.0650

483 Jacob, S., McClintock, M. K., Zelano, B., & Ober, C. (2002). Paternally inherited HLA  
 484 alleles are associated with women's choice of male odor. *Nature Genetics*, 30, 175-179. DOI:  
 485 10.1038/ng830

486 Kappeler, P. (1990). Female dominance in *Lemur catta*: More than just feeding priority?  
 487 *Folia Primatologica*, 55, 92-95. DOI: 10.1159/000156504

488 Knapp, L. A., Robson, J., & Waterhouse, J. S. (2006). Olfactory signals and the MHC: A  
 489 review and a case study in *Lemur catta*. *American Journal of Primatology*, 68, 568-584. DOI:  
 490 10.1002/ajp.20253

491 Laidre, M. E. (2009). Informative breath: olfactory cues sought during social foraging among  
 492 Old World monkeys (*Mandrillus sphinx*, *M. leucophaeus*, and *Papio anubis*). *Journal of*  
 493 *Comparative Psychology*, 123, 34-44. DOI: 10.1037/a0013129

494 Laska, M., Seibt, A., & Weber, A. (2000). 'Microsmatic' primates revisited: Olfactory  
 495 sensitivity in the squirrel monkey. *Chemical Senses*, 25, 47-53. DOI: 10.1093/chemse/25.1.47

496 Leclaire, S., Strandh, M., Mardon, J., Westerdahl, H., & Bonadonna, F. (2017). Odour-based  
 497 discrimination of similarity at the major histocompatibility complex in birds. *Proceedings of*  
 498 *the Royal Society of London B*, 284, 1846. DOI: 10.1098/rspb.2016.2466

499 Lindgren, B. S., & Miller, D. R. (2002). Effect of verbenone on five species of bark beetles  
 500 (Coleoptera: Scolytidae) in lodgepole pine forests. *Environmental Entomology*, 31(5): 759-  
 501 765. DOI: 10.1603/0046-225X-31.5.759

502 Macdonald, E. A., Fernandez-Duque, E., Evans, S., & Hagey, L. R. (2008). Sex, age, and  
 503 family differences in the chemical composition of owl monkey (*Aotus nancymae*) subcaudal  
 504 scent secretions. *American Journal of Primatology*, 70, 12-18. DOI: 10.1002/ajp.20450

505 Meyer, C., Gallo, T., & Schultz, S.T. (1999). Female dominance in captive red ruffed lemurs,  
 506 *Varecia variegata rubra* (Primates, lemuridae). *Folia primatologica*, 70(6), 358-361. DOI:  
 507 10.1159/000021718

508 Müller-Schwarze, D. (2006). *Chemical Ecology of Vertebrates*. Cambridge: Cambridge  
509 University Press.

510 Osada, K., Miyazono, S., & Kashiwayanagi, M. (2015). The scent of wolves: Pyrazine  
511 analogs induce avoidance and vigilance behaviors in prey. *Frontiers in Neuroscience*, 9: 363.  
512 DOI: 10.3389/fnins.2015.00363

513 Palagi, E., & Dapporto, L. (2006). Beyond odor discrimination: Demonstrating individual  
514 recognition by scent in *Lemur catta*. *Chemical Senses*, 31, 437-443. DOI:  
515 10.1093/chemse/bjj048

516 Pereira, M., Seeligson, M., & Macedonia, J. (1988). The behavioral repertoire of the black-  
517 and-white ruffed lemur, *Varecia variegata variegata* (Primates: Lemnridae). *Folia*  
518 *Primatologica*, 51, 1-32. DOI: 10.1159/000156353

519 Pochron, S., Morelli, T., Scirbona, J., & Wright, P. (2005). Sex differences in scent-marking  
520 in *Propithecus edwardsi* of Ranomafana National Park, Madagascar. *American Journal of*  
521 *Primatology*, 66, 97-110. DOI: 10.1002/ajp.20130

522 Porter, R. H., & Moore, J. D. (1971). Human kin recognition by olfactory cues. *Physiology &*  
523 *Behavior*, 27, 493-495. DOI: 10.1016/0031-9384(81)90337-1

524 Raps, S., & White, F.J. (1995) Female social dominance in semi-free-ranging ruffed lemurs  
525 (*Varecia variegata*). *Folia Primatologica*, 65(3), 163-168. DOI: 10.1159/000156883

526 Rigamonti, M. (1993). Home range and diet in red ruffed lemurs (*Varecia Variegata Rubra*)  
527 on the Masoala Peninsula, Madagascar. In Kappeler, P., & Ganzhorn, J. (eds). *Lemur Social*  
528 *Systems and Their Ecological Basis*.. New York: Springer.

529 Roberts S. C. (2007). Scent-marking. In: Wolff JO, Sherman PW (eds). *Rodent Societies: An*  
530 *Ecological and Evolutionary Perspective*. Chicago: Chicago University Press.

531 Scordato, E., & Drea, C. (2007). Scents and sensibility: Information content of olfactory  
532 signals in the ring-tailed lemur, *Lemur catta*. *Animal Behaviour*, 73, 301-314. DOI:  
533 10.1016/j.anbehav.2006.08.006

534 Roberts, S. A., Simpson, D. M., Armstrong, S. D., Davidson, A. J., Robertson, D. H.,  
535 McLean, L., Beynon, R. J., & Hurst, J. L. (2010). Darcin: a male pheromone that stimulates  
536 female memory and sexual attraction to an individual male's odour. *BMC Biology*, 8: 75.  
537 DOI: 10.1186/1741-7007-8-75

538 Scordato, E., Dubay, G., & Drea, C. (2007). Chemical composition of scent marks in the  
539 ring-tailed lemur (*Lemur catta*): Glandular differences, seasonal variation, and individual  
540 signatures. *Chemical Senses*, 32, 493-504. DOI: 10.1093/chemse/bjm018

541 Setchell, J., Vaglio, S., Moggi-Cecchi, J., Boscaro, F., Calamai, L., & Knapp, L. (2010).  
542 Chemical composition of scent-gland secretions in an Old World monkey (*Mandrillus*  
543 *sphinx*): Influence of sex, male status, and individual identity. *Chemical Senses*, 35, 205-220.  
544 DOI: 10.1093/chemse/bjp105

545 Setchell, J., Vaglio, S., Abbot, K., Moggi-Cecchi, J., Boscaro, F., Pieraccini, G., & Knapp, L.  
546 (2011). Odour signals major histocompatibility complex genotype in an Old World  
547 monkey. *Proceedings of Royal Society of London B*, 278, 274-280. DOI:  
548 10.1098/rspb.2010.0571



549 Seyfarth, R., & Cheney, D. (2003). Signallers and receivers in animal  
550 communication. *Annual Review of Psychology*, 54, 145-173. DOI:  
551 10.1146/annurev.psych.54.101601.145121

552 Soso, S. B., & Koziel, J. A. (2016). Characterizing the scent and chemical composition of  
553 *Panthera leo* marking fluid using solid-phase microextraction and multidimensional gas  
554 chromatography–mass spectrometry-olfactometry. *Scientific Reports*, 7: 5137. DOI:  
555 10.1038/s41598-017-04973-2

556 Smith, T. (2006). Individual olfactory signatures in common marmosets (*Callithrix jacchus*).  
557 *American Journal of Primatology*, 68, 585-604. DOI: 10.1002/ajp.20254

558 Smith, T., Muchlinski, M., Bhatnagar, K., Durham, E., Bonar, C., & Burrows A. (2015). The  
559 vomeronasal organ of *Lemur catta*. *American Journal of Primatology*, 77, 229-238. DOI:  
560 10.1002/ajp.22326

561 Smith, T., Tomlinson, A., Mlotkiewicz, J., & Abbott, D. (2001). Female marmoset monkeys  
562 (*Callithrix jacchus*) can be identified from the chemical composition of their scent marks.  
563 *Chemical Senses*, 26, 449-458. DOI: 10.1093/chemse/26.5.449

564 Sussman, R., Green, G., Porton, I., Andrianasolondraibe, O., & Ratsirarson, J. (2003). A  
565 survey of the habitat of *Lemur catta* in Southwestern and Southern Madagascar. *Primate*  
566 *Conservation*, 19, 32-57.

567 Vaglio, S., Minicozzi, P., Romoli, R., Boscaro, F., Pieraccini, G., Moneti, G., & Moggi-  
568 Cecchi, J. (2016). Sternal gland scent-marking signals sex, age, rank, and group identity in  
569 captive mandrills. *Chemical Senses*, 41, 177-186. DOI: 10.1093/chemse/bjv077

570 Vaglio, S., Bartels-Hardege, H., & Hardege, J. (2018). Pheromone. In: Vonk, J., &  
 571 Shackelford, T. (eds). *Encyclopedia of Animal Cognition and Behavior*. Springer, Cham.  
 572 DOI: 10.1007/978-3-319-47829-6

573 Vasey, N. (2005) New developments in the behavioral ecology and conservation of ruffed  
 574 lemurs (*Varecia*). *American Journal of Primatology*, 66(1), 1-6. DOI: 10.1002/ajp.20124

575 Vasey, N. (2003). *Varecia*, ruffed lemurs. In: Goodman, S., & Benstead, J. (eds). *The natural*  
 576 *history of Madagascar*. Chicago: University Chicago Press.

577 Vasey, N. (2007) The breeding system of wild red ruffed lemurs (*Varecia rubra*): a  
 578 preliminary report. *Primates*, 48(1), 41-54. DOI: 10.1007/s10329-006-0010-5

579 Walker, S. F. (1998). Animal Communication. In: Mey, J. L. (ed). *Concise Encyclopaedia of*  
 580 *Pragmatics*. Amsterdam: Elsevier.

581 Wedekind, C., & Furi, S. (1997). Body odour preferences in men and women: Do they aim  
 582 for specific MHC combinations or simply heterozygosity? *Proceedings of Royal Society of*  
 583 *London B*, 264, 1471-1479. DOI: 10.1098/rspb.1997.0204

584 Wedekind, C., Seebeck, T., Bettens, F., & Paepke, A. J. (1995). MHC-dependent mate  
 585 preferences in humans. *Proceedings of Royal Society of London B*, 260, 245-249. DOI:  
 586 10.1098/rspb.1995.0087

587 Wyatt, T. (2014). Proteins and peptides as pheromone signals and chemical  
 588 signatures. *Animal Behaviour*, 1-8. DOI: 10.1016/j.anbehav.2014.07.025

589 Zeller, A. C. (1987). Communication by sight and smell. In: Smuts, B. B., Cheney, D. L.,  
590 Seyfarth, R. M., Wrangham, R. W., & Struhsaker, T. T. (editors). *Primates societies*.  
591 Chicago: Chicago University Press.

**Table I. Ethogram (based on Scordato and Drea, 2007, and Vaglio *et al* 2016, modified).**

Behavior	Description
Scent-Marking; Neck / Mandible	Individual rubs neck region against substrate or upon an item within the enclosure
Scent-Marking; Ano-genital	Individual rubs genital region against substrate or upon an item within the enclosure
Sniffing / Licking; Environment	Individual deliberately places nostrils or tongue within 3cm from substrate or an item within the enclosure and sniffs/licks
Sniffing / Licking; Conspecific	Individual deliberately places nostrils or tongue within 3cm from a conspecific and sniffs/licks
Self-Licking	Individual uses tongue to lick an area near a scent gland on their own body

**Table 2 – Volatile compounds present in filter paper samples from female lemur ano-genital secretions identified tentatively using ChemStation and NIST mass spectral databases (v. MSD F.0101.2317). Compounds in bold font were found in all samples.**

Retention Time (mins)	Tentative Compound ID	Molecular Weight
3.906	Hexanal	100
6.057	5-methyl-3-hexanone	114
7.413	Alpha-pinene	136
8.077	1-isopropyl-4-methylenebicyclo[3.1.0]hex-2-one	134
8.268	<b>Benzaldehyde</b>	106
8.623	3,7,7-trimethyl-1,3,5-cycloheptatriene	134
9.096	Phenol	94
9.269	6-methoxy-5-hepten-2-one	126
10.720	<b>2-ethyl-1-hexanol</b>	130
12.362	<b>p-Cresol</b>	108
12.553	cis-Verbenol	152
13.385	<b>cis-p-Mentha-2,8-dien-1-ol</b>	152
14.104	1,7,7-Trimethylbicyclo[2.2.1]hepta-2-one	152
14.536	L-Pinocarveol	152
14.791	trans-Verbenol	152
15.605	p-Ethyl-phenol	122
15.928	Terpinen-4-ol	154
16.415	Alpha-Terpineol	154
16.615	Myrtenol	152
17.047	<b>2-Pinen-4-one</b>	150
18.252	Carvone	150
19.217	p-Mentha-1,8-dien-3-one	150
23.283	4,7,7-Trimethylbicyclo[4.1.0]hept-3-ene-2-one	150
23.443	Tetradecane	198
25.094	Geranylacetone	194
25.899	Isomethylionone	206
26.513	<b>Pentadecane</b>	212
30.871	2,6,10-Trimethylpentadecane	254
32.208	Heptadecane	240


32.372	2,6,10-Trimethylhexadecane	268
34.446	n-Tetracosane	338
34.591	2,6,10,14-Tetramethylhexadecane	282

---

**Fig. I. Sterile filter paper attached to popular scent-marking locations using plastic cable ties. (a) hatches, (b) climbing frame equipment, (c) tree branches.**



**Fig. II. Percentage of occurrence for total scent-marks associated to classes of authors. Males scent-marked the most (51.55%), followed by adult females (26.80%) and subadult females (21.65%), however differences were not statistically significant.**





**Fig III. Example chromatograms from (a) one adult female ano-genital odour secretions, showing contaminants and meaningful biological compounds; and (b) the control sample, showing contaminants. Red arrows indicate the six meaningful biological compounds which were found in all samples: (a) benzaldehyde; (b) 2-ethyl-1-hexanol; (c) p-cresol; (d) cis-p-mentha-2,8-dien-1-ol; (e) 2-pinen-4-one; (f) pentadecane.**

