

Effects of environmental variation during seed production on seed dormancy and germination

Steven Penfield^{1*}, and Dana R. MacGregor^{1,2}

1. Department of Crop Genetics, John Innes Centre, Norwich Research Park, Norwich, NR4 7UH, UK. 2. Current Address: School of Biological and Biomedical Sciences, Durham University, South Road, Durham, DH1 3LE, UK.

*Corresponding author Email: steven.penfield@jic.ac.uk

Abstract

The environment during seed production has major impacts on the behaviour of progeny seeds. It can be shown that for annual plants temperature perception over the whole life history of the mother can affect the germination rate of progeny, and instances have been documented where these affects cross whole generations. Here we discuss the current state of knowledge of signal transduction pathways controlling environmental responses during seed production, focusing both on events which take place in the mother plant and those which occur directly as a result of environmental responses in the developing zygote. We show that seed production environment effects are complex, involving overlapping gene networks active independently in fruit, seed coat and zygotic tissues that can be deconstructed using careful physiology alongside molecular and genetic experiments.

Key Words: seed dormancy, seed development, seed germination, maternal, environment, temperature, flowering time, FLC, seed coat, tannin

Highlight: Environmental signals during seed production strongly affect seed dormancy at harvest and are exploited by plants to generate variation in dormancy levels among progeny. This review discusses the state of the art knowledge of the underlying biological mechanisms.

Introduction

It has been known for more than a century that the mother plant has a significant influence over seed traits, including seed size, dormancy and germination. In many species factors such as age of the mother plant, position of the seed in the fruit, inflorescence or canopy can affect seed properties, often accompanied by a dimorphism either of the seeds themselves or the fruits in which they arise (Roach and Wulff, 1987). In addition to the major effects of the genetics and developmental characters of the mother plant, the environment during seed

production also has a major influence on seed germinability (Fenner, 1991), and this will be the major focus of this review. These environmental effects can be strong and temperature is the most dominant signal across species, with as little as 1 °C differences shown to have important consequences for seed dormancy in sensitive ranges (Springthorpe and Penfield, 2015). This extreme sensitivity stretches the capabilities of modern controlled environment technology in generating reproducible datasets experiment after experiment. This effect of temperature on seeds may be designed in part to transduce environmental noise into variation in dormancy among progeny (Springthorpe and Penfield, 2015) and therefore can be a confounding factor during commercial seed production, and severely restricts the global regions in which high quality seed can be produced for major seed markets.

Researchers typically refer to these environmental effects as the influence of the ‘maternal environment’ on seed properties, which has in the past been used to refer generally to the environmental conditions during seed production (Roach and Wulff, 1987). However, in the era of mechanistic biology it is now necessary to distinguish carefully between effects mediated by the mother plant and thus strictly maternal, from those which are simply direct effects of environmental variation on the developing zygote itself. In practise the two can be differentiated through careful physiological and mechanistic studies coupled with clear genetic analysis demonstrating whether the maternal genotype, or that of the zygote, is key to response to an environmental signal. So here I will define ‘Maternal Environment (ME)’ strictly as *environmental effects involving direct perception by the tissues of the mother plant*, and mediated by either the genome or epigenome of the mother. Typically this involves modification of fruit and/or seed coat tissues, the production of signals from the mother to the maturing zygote, or inheritance of maternal epigenetic states through meiosis into the embryo or endosperm. I will distinguish this from effects of the ‘Zygotic Environment (ZE)’ which I will define as *environmental effects involving direct perception of environmental signals by tissues of the zygote* during seed development and maturation, and genetically under the influence of the genome or epigenome of the embryo and/or endosperm. Carefully distinguishing between ME and ZE effects will be necessary to breed for improved seed characters, especially in crops where F₁ hybrid seeds dominate the market.

Evidence for maternal influences on progeny seed germination

It has long been known that variation in the temperature during seed set strongly affects seed dormancy, with lower temperatures almost always resulting in lower germination. However, what is less clear is whether this really is a maternal process, or whether paternal alleles in the zygote also contribute. One important set of experiments that reveals the presence of signal transduction pathways in the mother plant is the use of environmental manipulations

before first anthesis to manipulate progeny seed germination. For instance, temperature during the vegetative phase can be shown to affect progeny seed dormancy in tobacco, oats and *Arabidopsis* (Thomas and Raper, 1975; Sawhney et al., 1985; Chen et al., 2014), the wide conservation of this affect across diverse angiosperm families showing that this is a very general phenomenon. Because there are no gametes in the vegetative phase these experiments show clearly that it is not just the ZE that affects seed dormancy, but that the temperature experience of one or more of the parent plants can be remembered across time and affect progeny seed properties. In addition to temperature variation, it has been shown that altitude and time of year can affect seed dormancy, probably through differences in ME and ZE during seed production (Fenner, 1991).

The role of the seed coat in the environmental regulation of dormancy

In seeds with physical dormancy such as many legumes, the seed coat acts to prevent water uptake by the zygotic tissues (Bolingue et al. 2010). Seed dormancy in legumes depends on pigments such as tannins, and light-coloured seeds are less dormant. In legumes therefore, seed size and seed dormancy are in general maternally inherited (Davies, 1975). In other species, most notably barley, the seed coat has been shown to be essential for creating a low oxygen environment in the seed, and removing the seed coat permits growth of the embryo at low oxygen concentrations (Lenoir et al., 1986). The key evidence that the seed coat is critical to the imposition of physiological coat imposed dormancy in *Arabidopsis* comes from the very low dormancy of *transparent testa* (*tt*) mutants (Debeaujon et al., 2000). Even when *tt* mutant seed is set at low temperatures, some alleles retain close to 100% germination (MacGregor et al., 2015), far surpassing the consequences of deleting *DELAY OF GERMINATION1* (*DOG1*; Kendall et al., 2011), which can enter dormancy under the same conditions. However, when set at 16°C there is no strict correlation between seed coat colour and dormancy, with some yellow seeded mutants showing much weaker phenotypes than others (MacGregor et al., 2015), and this raises the possibility that the tannins themselves are not the only seed coat factors affecting dormancy. Further evidence for this comes from *Spergularia diandra*, where yellow seeded varieties are more dormant than dark seeded varieties (Gutterman, 1994), although in this case the dark-seed forms appear to have some defects in integument development. In addition to colour mutants, other genetic lesions that affect the ovule integument or seed coat development also affect seed dormancy (Leon-Kloosterziel et al., 1994; MacGregor et al., 2015).

The second candidate for the imposer of coat-imposed dormancy is of course the endosperm. In a key paper Bethke et al (2007) showed that mechanically removing the seed coat from

dormant *Arabidopsis* seeds did not break dormancy, whereas breaching the endosperm did. This appears instead to favour the endosperm as critical to seed dormancy imposition. The important role of the endosperm has further support, including genetic evidence that reactive oxygen production in the endosperm promotes dormancy (Penfield et al., 2006), and more comprehensively that the endosperm synthesises and secretes ABA in dormant *Arabidopsis* seeds (Lee et al., 2010; Kang et al., 2015). It is possible that the seed coat is required not during imbibition, but during seed maturation to impose physiological dormancy, for instance by secretion of growth regulators, or by provision of materials to the cell wall of the endosperm which itself has been implicated directly in dormancy imposition (De Giorgi et al., 2015).

The seed coat is a highly plastic plant organ that responds to environmental signals. Depending on the species this response can lead to a change in the development or metabolism in the seed coat producing a change in dormancy observable upon imbibition under favourable conditions. The genus *Chenopodium* exhibits physiological seed dormancy that depends on seed coat thickness, which in turn is affected by environmental signals experienced by the mother plant during seed maturation. In *Chenopodium polyspermum* and *Chenopodium album* seed coat thickness depends on day length with shorter days leading to thinner seed coats and seeds with higher levels of germination at harvest (Karssen, 1970; Pourrat and Jacques, 1975). A similar effect on seed coat development in *Chenopodium bonus-henricus*, but this time altitude rather than photoperiod was the driver of seed coat development, with higher altitudes leading to thicker seed coats and lower germination (Dorne, 1981), and with high altitude seeds accumulating more polyphenols. These ME effects are often associated with effects on seed coat colour which have been observed in several species including legumes where the seed coat also imposes physical dormancy (Gutterman and Evenari, 1972). Other environmental signals that affect seed coat development include salinity and nutrient levels (Wang et al., 2012). Strikingly, Wang et al (2012) showed that the ratio of brown to black seeds produced by *Suaeda aralocaspicavia* plants depended not only on the maternal environment but also the colour of the seed coat of the starting seed. This shows that maternal effects can persist across whole generations in some cases. With a keen eye differences in coat colour between seed lots matured at different temperatures can be observed in the model species *Arabidopsis*, and measured with a multispectral imager (Figure 1). These are accompanied by changes in seed coat tannin levels, but no obvious effects on seed coat thickness (MacGregor et al., 2015). Measuring the precise impact of these changes in tannin levels on seed dormancy is a challenge because although it is clear the tannin is required for dormancy, separating the contribution of the impacts on tannin levels from impacts on other processes, such as effects on hormone levels and *DOG1* expression, is not straightforward. However, it is clear that *Arabidopsis* is a good model system for investigating

environmental control of seed coat development, although the importance of the process is likely to vary from species to species. In *Arabidopsis* as in other species changes in seed coat thickness or tannin content caused by temperature effects are associated with altered seed coat permeability to water and dyes (Gutterman, 1978; MacGregor et al., 2015),

The fact that a memory of past temperature can be used by plants to control seed coat properties, and that in some species the process is affected by photoperiod, are important clues that seed coat metabolism is influenced by genes involved in flowering time control. *FLOWERING LOCUS C (FLC)*, circadian clock genes and *FLOWERING LOCUS T (FT)* control flowering in response to vernalisation and day length in *Arabidopsis*, and all have been shown to affect seed dormancy (Chiang et al., 2009; Penfield and Hall, 2009; Chen et al., 2014). *FT* expression in the phloem of siliques is more than 100-fold higher than in leaves (Adrian et al., 2010; Chen et al., 2014) and responds to ME signals that affect seed dormancy (Chen et al., 2014). In contrast, the *Arabidopsis FT* promoter is not expressed directly in seed tissues. Phloem-expressed FT-GFP fusions accumulate in the chalazal pole of the seed coat, indicating that the FT protein may be translocated to the seed to control dormancy, although transcript analyses show that FT also acts locally in silique tissues. The effects of *ft* mutation on the silique transcriptome can be phenocopied by exposing the vegetative tissues of the mother plant to lower temperatures (Chen et al., 2014). Together with genetic evidence that FT is necessary and sufficient for ME-regulated changes in seed coats and seed dormancy, this data show the central role of fruit-expressed FT in dormancy regulation by the ME. In *Capsella bursa-pastoris* flowering time was also linked to seed coat properties, with early flowering lines showing lower dormancy and having less seed coat mucilage (Toorop et al., 2012), and in *Ononis sicula* (Fabaceae) lengthening the photoperiod results in yellower seeds which in this case have more dormancy (Gutterman and Evenari, 1972). In *Arabidopsis*, flowering time genes can be directly linked to seed coat metabolism, with *ft-1* mutants showing increased tannin accumulation and increased expression of phenylpropanoid pathway enzymes necessary for tannin biosynthesis (Chen et al., 2014). These changes are associated with an effect on the transcript levels of major regulators of tannin biosynthesis in ovule integuments including *TRANSPARENT TESTA2* and the MADS box transcription factors *TRANSPARENT TESTA16* and *SHATTERPROOF2* (Figure 2).

In *Arabidopsis* studies disagree on whether photoperiod has an effect on dormancy, but our experience is similar to that of Donohue et al., (2005) in that standard photoperiod treatments used in plant physiology applied to the mother plant do not affect seed dormancy, and also do not affect *FT* expression (Adrian et al., 2010; Chen et al., 2014). This observation effectively rules out a role for FT synthesised in leaves in dormancy control. Yet the key photoperiod sensor protein *CONSTANS* is highly expressed in *Arabidopsis* siliques, so it is unclear why

FT expression in *Arabidopsis* fruits is unable to respond to photoperiod changes. A similar uncoupling of photoperiod responses was previously observed in cucumber where plants were daylength insensitive for flowering, but dormancy was promoted by long day signals applied to fruits (Gutterman, 1978).

Flowering time gene manipulation affects hormone metabolism in fruit and seeds

In addition to changes in the seed coat, flowering time genes also affect hormone metabolism in seeds and in this way could influence dormancy in some species. This could be particularly important in soft-fruited seeds such as tomato and cucumber where it has previously been shown that photoperiod affects accumulation of uncharacterised germination inhibitors in the pulp surrounding the seed (Gutterman, 1978a). The fact that photoperiod treatments can be applied to detached fruits and affect seed dormancy is good evidence that FT synthesised in fruits rather than leaves is important in dormancy control. ABA is a strong candidate for this fruit-derived germination inhibitor because in tobacco maternal ABA is important for seed dormancy regulation (Frey et al., 2004). This fruit-derived ABA can also be affected by nutrient status of the mother. Natural variation at *FLC* also influences ABA levels in seeds (Chiang et al., 2009), but this seemed to be directly in zygotic tissues because the effect was preserved in mature seeds. In addition, loss of FT increases the accumulation of both jasmonate and gibberellin (GA; Chen et al., 2014). Notably, *ft-1* mutant seeds accumulate higher levels of GA despite being more dormant than wild type, demonstrating that FT has the potential to increase dormancy, as well as decrease dormancy, acting through different mechanisms. It is also relevant that control of seed dormancy by oxylipins has been proposed to occur via an FT orthologue MOTHER OF FT AND TFL1 (*MFT*; Dave et al., 2016). This is an important observation because it is likely that different overlapping signalling cascades occur in different seed compartments using overlapping gene networks leading to different outcomes controlling dormancy. In embryos this is likely affects levels of hormones and control of hormone signal transduction. For instance, *FLC* has not only been linked to regulation of ABA metabolism through *CYTOCHROME P450 707A* (*CYP707A*) genes (Chiang et al., 2009; Deng et al., 2011), but also to indirect regulation of *GA3OX1* via both *TEMPRANILLO1* and *APETALA1* (Mateos et al., 2015). It is very likely that understanding the environmental control of seed germination will require understanding in detail links between flowering time pathways which carry environmental signals and hormone metabolism and signalling. Interestingly, maternal control of ABA levels has also been linked to the regulation of of seed coat pigmentation (Frey et al., 2004; Gu et al., 2011) and *Arabidopsis della* mutants have reduced seed coat

permeability (Chen et al., 2014), showing that hormone production and signalling also play a role in the development of the seed coat during seed maturation.

Seed dormancy and germination control by the Zygotic Environment

The clearest direct effect of the ZE on seed dormancy is the manipulation of light quality during seed maturation. Changing the red/far-red ratio during seed production changes the dormancy of seeds produced on the mother plant (Hayes and Klein, 1974; Cresswell and Grime, 1981). Importantly, giving far-red during seed maturation induced a red light requirement for germination, and this depended on the green colour of the tissues surrounding the zygote (Cresswell and Grime, 1981). This suggests that the perceptive tissue is the zygote within the green fruit tissues. It is hypothesised that in the embryo or endosperm the Pr:Pfr ratio of phytochrome states at maturity can be affected by light treatments given during seed maturation, and that this can subsequently affect light requirements upon imbibition. However, phytochromes do strongly affect *FT* expression in leaves (Halliday et al., 2003), so it is also possible that phytochrome has an overlapping dormancy-imposing role during seed development. In *Arabidopsis* there are five phytochromes (phyA-E), and loss of phyA, phyB and phyE causes a strong increased dormancy phenotype or a poor germination response to far-red light (Hennig et al., 2002). However, if seeds are matured at cooler temperatures then phyD becomes important for the response to dormancy-breaking temperature treatments (Donohue et al., 2008). This increase in importance for phyD may be explained by the fact that low temperatures during seed maturation strongly reduce *phyB* and *phyE* gene expression in seeds (Kendall et al., 2011). This underlines the close integration of light and temperature signalling pathways in *Arabidopsis* (Halliday et al., 2003; Penfield et al., 2005; Koini et al., 2009). As well as light quality, light intensity during seed maturation has been shown to affect seed dormancy in *Arabidopsis* (He et al., 2016).

Measuring the composition of developing and mature seeds, especially in species in which maternal tissues undergo apoptosis during the final stages of desiccation, is an excellent way to understand effects of ZE on seeds (Kendall et al., 2011; He et al., 2014; Righetti et al. 2015; He et al., 2016). Temperature during seed maturation has a strong effect on *DOG1* transcript levels in mature seeds, and also effects *CYP707A2* gene expression, GA and ABA levels (Kendall et al., 2011; Chiang et al., 2011). Because genetics shows that *DOG1* and ABA affect seed dormancy from the zygotic genotype (Koorneef et al., 1982; Alonso-Blanco et al., 2003) it is possible that these are direct impacts from temperature perception in the zygote itself. Measurement of the effects of light intensity, temperature and nitrate on metabolites in mature seeds show that these signals directly affect the metabolome of the zygote (He et al., 2016), suggesting they act in zygotic tissues. By combining an analysis of several *Arabidopsis* lines

with different levels of dormancy He et al (2016) showed that these effects on metabolites are dependent on the genotype, and that genotypes with similar dormancy levels had similar metabolomes. This suggests that metabolic composition is a consequence of dormancy state or the action of gene networks controlling dormancy state, rather than a causative factor of differences in dormancy levels. Thus we cannot rule out that some of the environmental signals controlling seed composition, dormancy and longevity in response to nutrient levels or light intensity also emanate from the mother plant.

Aside from *DOG1*, the best candidate for a central regulator of temperature response to the ZE is *MOTHER OF FT AND TFL1 (MFT)* (Nakamura et al., 2011). In wheat *MFT* is expressed in the embryonic tissues and is strongly up-regulated by low temperatures during seed maturation (Nakamura et al., 2011). Loss of *MFT* leads to lower dormancy indicating that *MFT* is a germination inhibitor, and variation at *MFT* underlies variation in dormancy among East Asian cultivars of wheat (Chono et al., 2015). Taken together the data show that *MFT* is a low temperature-induced inhibitor of seed germination. In *Arabidopsis* the role of *MFT* as a germination inhibitor is conserved and *MFT* is a direct target of the SPATULA bHLH transcription factor, which in turn is necessary for normal temperature responses in *Arabidopsis* seeds (Penfield et al., 2005; Vaistij et al., 2012). In *Arabidopsis* *MFT* is only weakly temperature-regulated at the transcript level compared to wheat (Kendall et al., 2011). Interestingly, *SPT* is also a regulator of fruit development (Heisler et al., 2001), further underlining the evolutionary links between environmental signalling pathways in the fruit, integuments and zygote. *MFT* expression is strongly up-regulated by the germination inhibiting hormone 12-oxo-phytodienoic acid (OPDA; Dave et al., 2016) suggesting the involvement of oxylipin hormones in temperature responses in seeds, as is the case during cold acclimation (Hu et al., 2013). *MFT* acts to control seed dormancy in a complex gene network including *ABI5*, DELLAs and hormone metabolism (Dave et al., 2016; Figure 2) that still requires more detailed analysis, and this analysis probably needs to extend beyond *Arabidopsis* where temperature effects on *MFT* expression are apparently attenuated. *MFT* is highly conserved among major crops suggesting it is a good target for dormancy manipulation in crop species.

Conclusions

It is interesting that the major effects of the ZE on seed dormancy involve temperature effects on *DOG1*, *MFT* and phytochrome because these appear to mimic those seen in buried seeds during seed dormancy cycling, the induction and loss of secondary dormancy (Footitt et al., 2011). So it is possible that these ZE effects result from a partial activation of secondary dormancy-regulating processes during seed maturation in the zygote. These processes work alongside truly maternal pathways in determining the final dormancy status of the mature

seed. These maternal processes clearly involve modification of fruit and seed coat tissues, but there are also maternally-derived mobile signals such as production of hormones and FT protein that have the potential to interact directly with zygotic tissues to control seed properties. Finally it is clear that some epigenetic states are inherited from mother to zygote, including imprinted genes and genes such as *FLC* which is strongly expressed and temperature sensitive in all tissues of the seed (Sheldon et al., 2008). Because there are high levels of cross talk between seed tissues it is very likely that maternal and zygotic environmental signal transduction pathways interact at several levels that remain to be discovered.

References

- Adrian J, Farrona S, Reimer JJ, Albani MC, Coupland G, Turck F.** 2010. cis-Regulatory elements and chromatin state coordinately control temporal and spatial expression of FLOWERING LOCUS T in Arabidopsis. *The Plant Cell* **22**, 1425-1440.
- Alonso-Blanco C, Bentsink L, Hanhart CJ, Blankestijn-de Vries H, Koornneef M.** 2003. Analysis of natural allelic variation at seed dormancy loci of Arabidopsis thaliana. *Genetics* **164**, 711-729.
- Bethke PC, Libourel IG, Aoyama N, Chung YY, Still DW, Jones RL.** 2007. The Arabidopsis aleurone layer responds to nitric oxide, gibberellin, and abscisic acid and is sufficient and necessary for seed dormancy. *Plant Physiology* **143**, 1173-1188.
- Bolingue W, Vu BL, Leprince O, Buitink J.** 2010. Characterization of dormancy behaviour in seeds of the model legume *Medicago truncatula*. *Seed Science Research* **20**, 97-107.
- Chen M, MacGregor DR, Dave A, Florance H, Moore K, Paszkiewicz K, Smirnov N, Graham IA, Penfield S.** 2014. Maternal temperature history activates Flowering Locus T in fruits to control progeny dormancy according to time of year. *PNAS* **111**, 18787-18792.
- Chiang GC, Barua D, Kramer EM, Amasino RM, Donohue K.** 2009. Major flowering time gene, flowering locus C, regulates seed germination in Arabidopsis thaliana. *Proceedings of the National Academy of Sciences of the United States of America* **106**, 11661-11666.
- Chiang GC, Bartsch M, Barua D, Nakabayashi K, Debieu M, Kronholm I, Koornneef M, Soppe WJ, Donohue K, De Meaux J.** 2011. DOG1 expression is predicted by the seed-maturation environment and contributes to geographical variation in germination in Arabidopsis thaliana. *Molecular Ecology* **20**, 3336-3349.
- Chono M, Matsunaka H, Seki M, Fujita M, Kiribuchi-Otobe C, Oda S, Kojima H, Nakamura S.** 2015. Molecular and genealogical analysis of grain dormancy in Japanese wheat varieties, with specific focus on MOTHER OF FT AND TFL1 on chromosome 3A. *Breeding Science* **65**, 103-109.
- Cresswell EG, Grime JP.** 1981. Induction of a light requirement during seed development and its ecological consequences. *Nature* **291**, 583-585.
- Dave A, Vaistij FE, Gilday AD, Penfield SD, Graham IA.** 2016. Regulation of Arabidopsis thaliana seed dormancy and germination by 12-oxo-phytodienoic acid. *Journal of Experimental Botany* **67**, 2277-2284.

- Davies DR.** 1975. Studies of seed development in *Pisum sativum*. I. Seed size in reciprocal crosses. *Planta* **124**, 297–302
- De Giorgi J, Piskurewicz U, Loubery S, Utz-Pugin A, Bailly C, Mène-Saffrané L, Lopez-Molina L.** 2015. An Endosperm-Associated Cuticle Is Required for Arabidopsis Seed Viability, Dormancy and Early Control of Germination. *PLoS Genetics* **11**, e1005708.
- Debeaujon I, Léon-Kloosterziel KM, Koornneef M.** 2000. Influence of the testa on seed dormancy, germination, and longevity in Arabidopsis. *Plant Physiology* **122**, 403-414.
- Deng W, Ying H, Helliwell CA, Taylor JM, Peacock WJ, Dennis ES.** 2011. FLOWERING LOCUS C (FLC) regulates development pathways throughout the life cycle of Arabidopsis. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 6680-6685.
- Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, Polisetty CR, Schmitt J.** 2005. Environmental and genetic influences on the germination of Arabidopsis thaliana in the field. *Evolution* **59**, 740-757.
- Donohue K, Heschel MS, Butler CM, Barua D, Sharrock RA, Whitelam GC, Chiang GC.** 2008. Diversification of phytochrome contributions to germination as a function of seed-maturation environment. *New Phytologist* **177**, 367-379.
- Dorne, C.J.** 1981. Variation in seed germination inhibition of *Chenopodium bonus-henricus* in relation to altitude of plant growth. *Canadian Journal of Botany* **59**, 1893–1901.
- Fenner M.** 1991. The effects of the parent environment on seed germinability. *Seed Science Research* **1**, 75-84.
- Footitt S, Douterelo-Soler I, Clay H, Finch-Savage WE.** 2011. Dormancy cycling in Arabidopsis seeds is controlled by seasonally distinct hormone-signaling pathways. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 20236-20241.
- Frey A, Godin B, Bonnet M, Sotta B, Marion-Poll A.** 2004. Maternal synthesis of abscisic acid controls seed development and yield in *Nicotiana plumbaginifolia*. *Planta*. **218**, 958-964.
- Gu XY, Foley ME, Horvath DP, Anderson JV, Feng J, Zhang L, Mowry CR, Ye H, Suttle JC, Kadowaki K, Chen Z.** 2011. Association between seed dormancy and pericarp color is controlled by a pleiotropic gene that regulates abscisic acid and flavonoid synthesis in weedy red rice. *Genetics* **189**, 1515-1524.
- Gutterman Y.** 1974. The influence of the photoperiodic regime and red/far-red light treatments of *Portulaca oleracea* L. plants on the germinability of their seeds. *Oecologia* **17**, 27–38.
- Gutterman Y.** 1978 Germinability of seeds as a function of the maternal environments. *Acta Horticulture* **83**, 49–55.
- Gutterman, Y.** 1978. Seed coat permeability as a function of photoperiodical treatments of the mother plants during seed maturation in the desert annual plant *Trigonella arabica*, del. *Journal of Arid Environments* **1**,141-144.
- Gutterman Y.** 1978. Influence of environmental conditions and hormonal treatment of the mother plants during seed maturation on the germination of their seeds. In: Malik, C.P. (ed.) *In Advances. Plant Reproductive Physiology*. Kalyani Publishers, New Delhi, pp. 288–294

- Gutterman Y, Evenari M.** 1972. The influence of day length on seed coat colour, an index of water permeability of the desert annual *Ononis sicula* Guss. *Journal of Ecology* **60**, 713–719.
- Halliday KJ, Salter MG, Thingnaes E, Whitelam GC.** 2003. Phytochrome control of flowering is temperature sensitive and correlates with expression of the floral integrator FT. *Plant Journal* **33**, 875-885.
- Hayes RG, Klein WH.** 1974. Spectral quality influence of light during seed development of *Arabidopsis thaliana* plants regulating seed germination. *Plant and Cell Physiology* **15**, 643-653.
- He H, de Souza Vidigal D, Snoek LB, Schnabel S, Nijveen H, Hilhorst H, Bentsink L.** 2014. Interaction between parental environment and genotype affects plant and seed performance in *Arabidopsis*. *Journal of Experimental Botany* **65**, 6603-6615.
- He H, Willems LA, Batushansky A, Fait A, Hanson J, Nijveen H, Hilhorst HW, Bentsink L.** 2016. Effects of Parental Temperature and Nitrate on Seed Performance are Reflected by Partly Overlapping Genetic and Metabolic Pathways. *Plant and Cell Physiology* **57**, 473-487.
- Heisler MG, Atkinson A, Bylstra YH, Walsh R, Smyth DR.** 2001. SPATULA, a gene that controls development of carpel margin tissues in *Arabidopsis*, encodes a bHLH protein. *Development* **128**, 1089-1098.
- Hennig L, Stoddart WM, Dieterle M, Whitelam GC, Schäfer E.** 2002. Phytochrome E controls light-induced germination of *Arabidopsis*. *Plant Physiology* **128**, 194-200.
- Hu Y, Jiang L, Wang F, Yu D.** 2013. Jasmonate regulates the inducer of cbf expression-C-repeat binding factor/DRE binding factor1 cascade and freezing tolerance in *Arabidopsis*. *The Plant Cell* **25**, 2907-2924.
- Kang J, Yim S, Choi H, Kim A, Lee KP, Lopez-Molina L, Martinoia E, Lee Y.** 2015. Abscisic acid transporters cooperate to control seed germination. *Nature Communications*. **6**, 8113.
- Karssen CM.** 1970. The light promoted germination of the seeds of *Chenopodium album* L. III. Effect of the photoperiod during growth and development of the plants on the dormancy of the produced seeds. *Acta Botanica Neerlandica* **19**, 81–94.
- Kendall SL, Hellwege A, Marriot P, Whalley C, Graham IA, Penfield S.** 2011. Induction of dormancy in *Arabidopsis* summer annuals requires parallel regulation of DOG1 and hormone metabolism by low temperature and CBF transcription factors. *The Plant Cell* **23**, 2568-2580.
- Koini MA, Alvey L, Allen T, Tilley CA, Harberd NP, Whitelam GC, Franklin KA.** 2009. High temperature-mediated adaptations in plant architecture require the bHLH transcription factor PIF4. *Current Biology* **19**, 408-413.
- Koornneef M, Jorna ML, Brinkhorst-van der Swan DL, Karssen CM.** 1982. The isolation of abscisic acid (ABA) deficient mutants by selection of induced revertants in non-germinating gibberellin sensitive lines of *Arabidopsis thaliana* (L.) heynh. *Theoretical and Applied Genetics* **61**, 385-393.
- Lee KP, Piskurewicz U, Turecková V, Strnad M, Lopez-Molina L.** 2010. A seed coat bedding assay shows that RGL2-dependent release of abscisic acid by the endosperm controls embryo growth in *Arabidopsis* dormant seeds. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 19108-19113.

- Lenoir C, Corbineau F, Côme D.** 1986. Barley (*Hordeum vulgare*) seed dormancy as related to glumella characteristics. *Physiologica Plantarum* **68**: 301-307.
- Leon-Kloosterziel KM, Keijzer CJ, Koornneef M.** 1994. A Seed Shape Mutant of *Arabidopsis* That Is Affected in Integument Development. *The Plant Cell* **6**, 385-392.
- MacGregor DR, Kendall SL, Florance H, Fedi F, Moore K, Paszkiewicz K, Smirnov N, Penfield S.** 2015. Seed production temperature regulation of primary dormancy occurs through control of seed coat phenylpropanoid metabolism. *New Phytologist* **205**, 642-652.
- Mateos JL, Madrigal P, Tsuda K, Rawat V, Richter R, Romera-Branchat M, Fornara F, Schneeberger K, Krajewski P, Coupland G.** 2015. Combinatorial activities of SHORT VEGETATIVE PHASE and FLOWERING LOCUS C define distinct modes of flowering regulation in *Arabidopsis*. *Genome Biology* **16**, 31.
- Nakamura S, Abe F, Kawahigashi H, Nakazono K, Tagiri A, Matsumoto T, Utsugi S, Ogawa T, Handa H, Ishida H, Mori M, Kawaura K, Ogihara Y, Miura H.** 2011. A wheat homolog of MOTHER OF FT AND TFL1 acts in the regulation of germination. *The Plant Cell* **23**, 3215-3229.
- Penfield S, Hall A.** 2009. A role for multiple circadian clock genes in the response to signals that break seed dormancy in *Arabidopsis*. *The Plant Cell* **21**, 1722-3172.
- Penfield S, Josse EM, Kannangara R, Gilday AD, Halliday KJ, Graham IA.** 2005. Cold and light control seed germination through the bHLH transcription factor SPATULA. *Current Biology* **15**, 1998-2006.
- Penfield S, Li Y, Gilday AD, Graham S, Graham IA.** 2006. *Arabidopsis* ABA INSENSITIVE 4 regulates lipid mobilization in the embryo and reveals repression of seed germination by the endosperm. *The Plant Cell* **18**, 1887-1899.
- Pourrat Y, Jacques R.** 1975. The influence of photoperiodic conditions received by the mother plant on morphological and physiological characteristics of *Chenopodium polyspermum* L. seeds. *Plant Science Letters* **4**, 273–279.
- Righetti K, Vu JL, Pelletier S, Vu BL, Glaab E, Lalanne D, Pasha A, Patel RV, Provart NJ, Verdier J, Leprince O, Buitink J.** 2015. Inference of Longevity-Related Genes from a Robust Coexpression Network of Seed Maturation Identifies Regulators Linking Seed Storability to Biotic Defense-Related Pathways. *The Plant Cell* **27**, 2692-2708.
- Roach DA, Wulff RD.** 1987. Maternal effects in plants. *Annual Review of Ecology and Systematics* **18**, 209-235.
- Sawhney R, Quick WA, Hsiao AI.** 1985. The effect of temperature during parental vegetative growth on seed germination of wild oats *Avena fatua* L. *Annals of Botany* **55**, 25-28.
- Springthorpe V, Penfield S.** 2015. Flowering time and seed dormancy control use external coincidence to generate life history strategy. *Elife* **4**, 05557.
- Thomas TH, Raper CD.** 1975. Seed germinability as affected by the environmental temperature of the mother plant. *Tobacco Science* **19**, 98-100.
- Toorop PE, Cuerva RC, Begg GS, Locardi B, Squire GR, Iannetta PP.** 2012. Co-adaptation of seed dormancy and flowering time in the arable weed *Capsella bursa-pastoris* (shepherd's purse). *Annals of Botany* **109**, 481-489.

Vaistij FE, Gan Y, Penfield S, Gilday AD, Dave A, He Z, Josse EM, Choi G, Halliday KJ, Graham IA. 2012. Differential control of seed primary dormancy in *Arabidopsis* ecotypes by the transcription factor SPATULA. *Proceedings of the National Academy of Sciences of the United States of America* **110**, 10866-10871.

Wang L, Baskin JM, Baskin CC, Cornelissen JHC, Dong M, Huang Z. 2012. Seed dimorphism, nutrients and salinity differentially affect seed traits of the desert halophyte *Suaeda aralocaspicaviva* multiple maternal effects. *BMC Plant Biology* **12**, 170.

Figures




Figure 1. Replicate seed batches of *Arabidopsis thaliana* Col-0 accession produced at either 16°C or 22°C, to show seed coat colour differences observable between the two treatments. Shown is the multispectral quantification of light reflectance from *Arabidopsis* Col-0 and the tannin-deficient *banyuls-4* (*ban4*) mutant. Differences in reflectance seen in wild type seeds set at different temperatures are no longer observed in *ban-4*.




Figure 2. Summary of characterised processes mediating control of seed dormancy and germination by environmental signals during seed production. Diagram shows cross section of an Arabidopsis fruit revealing the internal structure including seeds. On the left processes which control responses to the ZE, the seed on the right processes controlling seed germination from the ME. Supporting data is to be found in Chiang et al., 2009; Penfield and Hall, 2009; Chen et al., 2014; Kendall et al., 2011; Chiang et al., 2011; Deng et al., 2011; Mateos et al., 2015; Kaufmann et al., 2009; Vaistij et al 2012; Nakamura et al., 2011; Dave et al 2016; He et al 2016). Temperature-regulated genes are shown in blue.