

Phase Separation on Bicontinuous Cubic Membranes: Symmetry Breaking, Re-entrant and Domain Facetting

Fabien Paillusson^{*,†}, Matthew R. Pennington^{*} and Halim Kusumaatmaja^{*}

^{*}*Department of Physics, Durham University, South Road DH1 3LE*

[†]*School of Mathematics and Physics, University of Lincoln, Brayford Pool, LN6 7TS*

(Dated: June 28, 2016)

We study the phase separation of binary lipid mixtures that form bicontinuous cubic phases. The competition between non-uniform Gaussian membrane curvature and line tension leads to a very rich phase diagram, where we observe symmetry breaking of the membrane morphologies and re-entrant phenomena due to the formation of bridges between segregated domains. Upon increasing the line tension contribution, we also find facetting of lipid domains that we explain using a simple argument based on the symmetry of the underlying surface and topology.

Introduction - Lipid self-assembly can adopt an astonishing range of shapes and morphologies, from single bilayer structures to stacks and convoluted periodic structures [1]. Nature has, of course, exploited this polymorphism. A large number of organelles feature lipid-based structures, including synaptic vesicles, endoplasmic reticulum, and Golgi apparatus. At the same time, lipids are indispensable for detergency and foodstuffs industries [2], and membrane-based structures are increasingly exploited in biotechnological and biomedical applications, e.g. as efficient nanoporous scaffolds for tissue engineering [3] or for gene silencing with siRNA [4].

In this letter we will focus on one particular type of mesophases that lipid mixtures in water can adopt, the so-called bicontinuous cubic phases (BCP) [5–7], whereby the lipids form a triply periodic lipid bilayer that separates two percolating and non-intersecting water channels [8–10]. These phases have attracted attention due to their high surface area, continuity of the bilayer surface, and pore network. The amphiphilic nature of the lipids also allows other molecules to be embedded in them; for example, they have high propensity to enable membrane protein crystallization. Although the details remain unclear, it is thought that a combination of curvature induced phase separation on the cubic surface, a local destabilization of the cubic phase to a lamellar phase and a two dimensional reservoir of proteins provided by the cubic phase are responsible for the observed yield [11]. Here our interests are in the aforementioned curvature induced phase separation.

Both in the biological and synthetic systems, these lipid mesophases usually contain more than one lipid species. To the best of our knowledge, the distribution of different lipids across such a cubic surface, especially the possible demixing transitions under the influence of *non-uniform* curvature of the membrane structures, is still not well-understood. Most studies on lipid phase separation focus on much simpler membrane geometries, such as lipid vesicles and supported membranes [12–14]. From a biological perspective, lateral lipid organizations into domains and membrane curvatures are ubiquitous features, and are known to play an important role for




Figure 1. *Visualising the P-surface.* Left panel: normalised Gaussian curvature field $\frac{G(x)}{G_{min}}$ (note that $G(x) \leq 0$) on a single patch with the zero-curvature point p at its centre. Right panel: curvature induced formation of A-lipid domains (in green) in k patches among the 8 available denoted by $\binom{8}{k}$ (for $f_A = 0.07$). For the sake of illustration we show them for $k = 1$ (top) and $k = 8$ (bottom).

the membrane functionalities [15, 16]. From a materials perspective, understanding the distribution of species of interest on a BCP may be the first steps towards a systematic and rational functionalization of BCPs, where active species can be localized into targeted domains. Finally, our work provides a comprehensive phase diagram, with predictions of distinguishing features which we hope will stimulate experimental verifications.

This letter is organized as follows. We first show that if the two species do not interact but induce different bending rigidities, then a single type of curvature induced phase separation occurs at all non zero area fractions. Upon considering interactions between the species, we observe a multiplicity of new modalities for the phase separation, including the formation of bridges between previously disconnected lipid domains. Moreover, we observe *facetting* of domains for which we provide a simple explanation relying on symmetry and topology.

Segregation in absence of line tension - In this paper we consider a binary lipid mixture or, alternatively, a mixture of lipids and proteins that has formed a minimal surface S (with zero mean curvature everywhere), and

ask what are the thermodynamically favoured repartitions of the species and how they depend on the bending rigidities and inter-species interactions. Here our focus is on the triply periodic surfaces which are known to be formed by lipid mixtures as well as mixtures of lipids and proteins in water [5–7]. We use standard notations P, D, and G for the *primitive*, *Diamond*, and *Gyroid* surfaces respectively. Since they are periodic, we characterize their properties per unit cell. To model a binary mixture on a curved surface, we use a straightforward extension of the Helfrich hamiltonian [17] to the case of a binary mixture on a minimal surface that reads (cf. Supplemental Materials (SM) for more details):

$$H_{el}^S(f_A) = \delta\kappa \int_S d\mu_S(x) \sigma_A(x) G(x), \quad (1)$$

where x denotes a point on S , $d\mu_S(x)$ is the area measure on S at x , $G(x) < 0$ is the gaussian curvature at point x , f_A is the imposed area fraction of species A (with $f_B = 1 - f_A$), the field $\sigma_A(x) \in [0, 1]$ is the *mean* occupation number of species A at x , $\delta\kappa = \kappa_g^A - \kappa_g^B$ and $\kappa_g^{A,B}$ are the gaussian bending rigidities associated to the species A and B respectively. Our convention here is $\kappa_g^A < \kappa_g^B$, such that $\delta\kappa < 0$. It is also worth remarking that, since typically $\kappa_g < 0$, B domains are softer than A domains. Such a model may represent a coexistence between L_o (liquid ordered; A -rich) and L_d (liquid disordered; B -rich) domains, or alternatively, between lipid-rich and protein-rich domains.

To model the distribution σ_A in a more tractable way, we first stress that any minimal surface with genus g embedded in a flat torus \mathbb{T}^3 must contain $4(g - 1)$ zero Gaussian curvature points [18]. For the P, D and G surfaces, $g = 3$ and they have 8 zeros. Each zero is located at the centre of a hexagonal area we term as a *patch* (See Fig. 1 here and Fig. 1 in the SM). The unit cell of either of the P, D or G surfaces can thus be partitioned into 8 equivalent patches $\{\Sigma_i\}_{i=1..8}$ such that the unit cell surface $S = \cup_{i=1}^8 \Sigma_i$. We characterise the repartition of the lipids on S by both the area fraction f_A^i of lipid A on each patch Σ_i and the occupation number function $\sigma_A^i(x)$ in it. For each patch Σ_i , given σ_A^i , the entropy then reads $\mathbb{S}_i = -k_B \int_{\Sigma_i} d\mu_S(x) [\sigma_A^i \ln \sigma_A^i + (1 - \sigma_A^i) \ln(1 - \sigma_A^i)]$. Minimising the overall free energy $\mathcal{F} = H_{el}^S(f_A) - T \sum_{i=1}^8 \mathbb{S}_i$ with respect to the occupation number for a given set of area fractions $\{f_A^i\}_{i=1..8}$ leads to the typical Fermi-Dirac distribution $\sigma_A^{i*}(x) = [1 + e^{-\beta\lambda_A^i + \beta\delta\kappa G(x)}]^{-1}$ where λ_A^i is a Lagrange multiplier that imposes the value of f_A^i .

At low temperatures, the Fermi-Dirac distribution will reach a value close to unity for all points x of Σ_i with an energy lower than λ_A^i . The lowest energy point p_i in a patch Σ_i is the symmetry point of the patch which has exactly zero gaussian curvature (cf. Fig. 1). Thus, at low T , the lipids A will *fill* the neighbourhood of p_i until they reach a critical *Fermi* curve \mathcal{C}_F , where $\{x \in \mathcal{C}_F | \delta\kappa G(x) =$

$\lambda_A^i\}$, beyond which there is no more lipids of type A (cf. Fig. 1; a disconnected area occupied by lipid A is termed as a domain). Close to p_i , one may use polar coordinates (ρ_i, θ_i) and, as a crude approximation, the space is assumed euclidean and circularly symmetric near p_i . This allows us to Taylor expand the function G about p_i up to the second order so that the curvature energy reads $H_{el}^{\Sigma_i}(f_A^i) \sim \delta\kappa \int_0^{R_i} 2\pi\rho_i d\rho_i [G''(p_i)\rho_i^2/2] \sim C(f_A^i)^2$, where R_i is the mean radial distance of the Fermi curve from the point p_i such that $f_A^i \approx \pi R_i^2/\mu(\Sigma_i)$. Here C is a constant and $\mu(\Sigma_i)$ is the area of the hexagonal patch Σ_i . Remarkably, in spite of the very crude approximations we have used, the predicted behaviour of the curvature energy $H_{el}^{\Sigma_i}(f_A^i) \propto (f_A^i)^\alpha$, with $\alpha = 2$, is close to what we observed in simulations for the P-surface where the exponent is found to be $\alpha = 1.83$ (cf. SM).

Next, upon minimising the total free energy $\mathcal{F} = C \sum_{i=1}^8 (f_A^i)^2$ with respect to the area fractions f_A^i at fixed total area fraction $f_A = (\sum_{i=1}^8 f_A^i)/8$, it is easy to see that the ground state in repartition among the patches is always $f_A^i = f_A$ for all values of f_A , corresponding to the $\binom{8}{8}$ configuration in Fig. 1.

Effect of the line tension - We have seen that, with only curvature, the A lipids are evenly distributed among the 8 available patches and formed dense domains in the neighbourhood of zero curvature points at low temperature. This begs the question of how this picture changes if the $A - B$ interactions are not negligible *i.e.* if there are line tension effects arising with domain formation, which is a more realistic physical scenario. To answer this question, we now carry out computer simulations of the binary phase separation on the P-surface (the qualitative picture is the same for the D- and G-surfaces, as justified in the SM). There are several known approaches to model bicontinuous cubic membranes, from coarse-grained Molecular Dynamics simulations [19] to continuum field theoretical approaches [20–22]. Here we use Metropolis Monte Carlo simulations [23] to resolve the thermodynamics of the system.

In our approach, we explicitly discretize a piece of the P-surface contained in a cubic cell. This can be efficiently done with the help of the Weierstrass-Enneper (W-E) representation of minimal surfaces [18, 24, 25]. Upon discretization, the binary mixture can then be modelled as an Ising-like problem (see the SM for technical details). A spin variable s is associated to each site and takes either value 0 (for species B) or 1 (for species A). The curvature hamiltonian of Eq. (1) thus maps exactly onto a system of magnetic spins on a network $\mathcal{N}(S)$ with a node-dependent external magnetic field and reads:

$$H_{el}^S(f_A) = \sum_{i \in \mathcal{N}(S)} \delta A_i \delta\kappa G_i s_i, \quad (2)$$

where δA_i is the area of the tile i on the surface. In this language, at any finite f_A , species A (spin variable $s = 1$)




Figure 2. *Phase diagrams* of the number of patches k occupied by lipid A domains among the 8 available as a function of the area fraction f_A and the ratio $J\ell/|\delta\kappa|$. ℓ is the lattice spacing of the cubic unit cell. For clarity, we show the phase diagrams (a) excluding (filled symbols) and (b) including (open symbols) the possibilities of bridge formations between lipid A domains. In (c) are shown configurations with increasing energy from top to bottom for $f_A = 0.75$ and $J\ell/|\delta\kappa| = 0.5$. The boxed configuration with 7 connected domains corresponds to the most stable phase.

will occupy sites with the lowest value of $\delta A_i \delta \kappa G_i$ to minimize the total energy, as we have analyzed with a different vocabulary in the previous section. To model the $A - B$ interspecies interactions, we choose a short-range nearest neighbours interaction which directly translates into the line tension of the lipid domains:

$$H_{A-B}^S(f_A) \equiv J \sum_{i \in \mathcal{N}(S)} \sum_{j \in \mathcal{N}(i)} (s_i + s_j - 2s_i s_j) \delta L_{ij}, \quad (3)$$

where J sets the magnitude of the exchange interactions, δL_{ij} is the length of the edge shared by cells i and j , and $(s_i + s_j - 2s_i s_j) = 1$ when $s_i \neq s_j$ and 0 otherwise.

Symmetry breaking - As it is evident, the hamiltonian in Eq. (3) is equivalent to an Ising model of ferromagnetism and therefore should lead to the same phenomenology: above a critical temperature $T^*(J)$, the system is paramagnetic and the two lipid species are mixed;

while below T^* , the system becomes ferromagnetic and a *symmetry breaking* favouring "lumping" of spins in spatial regions (segregation) occurs. There is, however, one crucial difference between the standard Ising model and our model. For the former, line tension effects always dominate demixing: domains of A lipids coalesce to minimize the overall interfacial energy. In our model, this coalescence mechanism competes with the curvature-induced mechanism described in the previous section.

The first effect of line tension is to re-shuffle the (energy) ranking of configurations $\binom{8}{k}$ with k patches occupied by the A species by shifting *down* the low k configurations (because they have a lower interfacial cost) and *up* the high k ones (because they have a high interfacial cost). A first account of the competition between curvature and line tension consists in assuming that the total energy of a configuration $\binom{8}{k}$ at a given packing fraction f_A would read $k[H_{el}^S(8f_A/k) + H_{A-B}^S(8f_A/k)]$ *i.e.* as the sum of the free energy of individual patches of equal size. This *summation* approximation is valid when isolated domains are formed at the centre of the hexagonal patches, and one finds that increasing f_A at fixed $J\ell/|\delta\kappa|$ always favours, eventually, higher k values, in agreement with Monte Carlo simulation results shown in the phase diagram in Fig. 2(a).

The caveat is that this summation approach is only valid when the A species domains are disconnected. Above certain f_A values, the lowest energy configurations are in fact those in which domains of lipid A span across multiple patches (see Fig. 2). These *bridges* between patches essentially make the domains interact negatively and in a non-pairwise fashion. The location of these bridges coincides with the lowest curvature energy regions at the patch boundary (c.f. Fig. 1). Taking these configurations into account, the phase diagram in Fig. 2(b) shows that the simple picture of Fig. 2(a) only holds for small $J\ell/|\delta\kappa|$ and f_A . In fact, one observes *re-entrant* behaviours whereby a configuration $\binom{8}{k}$ previously unfavored in the disconnected regime, becomes re-favored thermodynamically. We note that when bridges are formed in the $\binom{8}{8}$ configuration (open crosses in Fig. 2(b)), the segregated and continuous phases are effectively *inverted* (lipid B domains are surrounded by A).

Domain facetting - Another distinguishing feature that appears with line tension is the *facetting* of the domains formed by the A lipids. This effect is shown in Fig. 3(b) where the domain almost draws a hexagon compared to Fig. 3(a) where the shape is more rounded, thus the term "facetting". To explain this, we recall that in general if the underlying manifold has an n -fold rotational symmetry, we expect the bounding curve that minimises the perimeter length of a domain with fixed area to be a regular n -gon whose sides are geodesics of the underlying manifold. Moreover, on an anisotropic curved surface, not all orientations of a regular n -gon are equivalent as they lead in principle to different total perimeter lengths.




Figure 3. *Domain faceting*. Shape of a lipid A domain in the neighbourhood of the zero-curvature point p of a single hexagonal patch. Note that only a half of the patch is represented. (a) in absence of line tension and (b) with high line tension ($J\ell/|\delta\kappa| > 1$). (c) Geodesic curvature as a function of the curvilinear coordinate l in absence of line tension (dashed) and with high line tension (solid).

Thus, we interpret the bounding curve in Fig. 3 (b) with 6-fold symmetry to be the curve that minimises both shape and orientation at the same time.

To test the above rationale, we estimate the geodesic curvature along the bounding curves of the two representative examples shown in Fig. 3(a) and (b) (cf. e.g. Ref. [26]). In these figures, the curvilinear coordinate $l \in [0, 1]$ is the normalised arc length of each curve which enables the comparison of the geodesic curvature for curves with different total lengths in Fig. 3(c). In absence of line tension, the geodesic curvature g_c is approximately constant around the boundary. With a large line tension, g_c reaches very high values for l close to zero, but is much smaller than that without line tension as l approaches 1. This is consistent with the above explanation although it shows that the faceting is not perfect. It nevertheless sheds light on what happens as we approach the ideal faceting case: the geodesic vanishes almost everywhere except close to $l \simeq 0$ where it diverges. This divergence is representative of the wedge formed by the intersection of two geodesics of the 6-gon and whose angle γ can be estimated to be (cf. SM) $\gamma = 2\pi/3 - (6|\delta\kappa|)^{-1}H_{el}^{\Sigma_i}(f_A^i)$ for an area $f_A^i\mu(\Sigma_i)$.

Discussions - Let us start by estimating where typical lipid mixtures are located in the phase diagram of Fig. 2. For a mixture of DOPC, sphingomyelin, and cholesterol forming coexisting L_o and L_d domains, $J \simeq 1.2$ pN and $|\delta\kappa| \simeq 3 \times 10^{-19}$ J [27, 28]. Most synthetic BCPs, however, are formed using the lipid Monoolein, which is known to have a low bending rigidity with $|\kappa_G| \sim \kappa_m < 10k_B T$ [29]. Here κ_m is the (mean curvature) bending rigidity. Using these values and taking the typical lattice spacing of a BCP $\ell \simeq 10 - 100$ nm [30–32], this leads to $J\ell/|\delta\kappa|$ in the range of $O(0.1)$ - $O(1)$ considered in this paper.

When the line tension effects can be neglected, the

natural curvature of the surface alone is enough to a) induce segregation in all surfaces and b) the segregation is such that domains form in the same proportions on all available patches on the surface. We then confirmed this theoretical prediction by numerical calculations on the P-surface and looked at the effects of non-zero line tension. The latter gives rise to two important features. (i) Below the demixing critical temperature, it favours the formation of bigger domains in a fewer number of patches available on the surface that we characterize with a corresponding phase diagram. We also observe re-entrances in this patch-occupation space due to the formation of bridges between domains on neighbouring patches. Some of these morphologies should lead to distinguishing features (e.g. different x-ray scattering signatures due to the change in symmetry), and we hope this work will stimulate experimental works to verify our predictions. (ii) In the large line tension limit, we observed a faceting of the domains for which we provided a simple explanation and that we can relate to the curvature energy of a domain on a patch.

Predicting patterning on cubic membranes is the vital first step towards their systematic and rational functionalization. On one hand, the ability to localize molecular species by design into targeted domains can be beneficial for controlled release in drug delivery or of chemical substances [33, 34], and for templating self-assembly [35] or phase separation [36, 37] in the surrounding fluids. On the other hand, suppressing phase separation between lipid species or between lipids and proteins can be desirable in applications such as protein crystallization [11], where segregation at an incorrect stage can strongly hamper the efficiency of the applications.

There are also a number of avenues for future work. Firstly, here we have assumed that the BCP remains a minimal surface. A closer inspection based on the theory developed in [38] for domain-induced budding shows that the conclusions presented here can be qualitatively affected when $\kappa_m/|\delta\kappa| < 0.4$ (cf. SM). However, estimates of this ratio for a wide range of lipid bilayers and monolayers in the literature show that it is only rarely below 1 [39]. This suggests that the minimal surface assumption is very reasonable for realistic parameter values. Further work is however still needed to fully assess how membrane deformation, including budding instability, affects the phase diagram of multicomponent BCPs. Secondly, the present work tacitly assumes that the membrane domains are formed by lipids of the same species in the two leaflets (*registration* phase). Indeed, recent work on flat bilayers suggests that registered domains is the thermodynamically favoured phase for a wide range of lipid mixtures [40]. It would be interesting to relax this assumption to probe how curvature affects registration/anti-registration and how, in turn, registration/anti-registration may affect the bilayer morphology. Thirdly, the system considered here provides an excellent setup to study how non-

uniform curvature may affect the nature of the demixing phase transition.

Acknowledgements - We thank J. M. Seddon, P. D. Olmsted, J. J. Williamson, N. J. Brooks, D. Frenkel, F. Schmid and C. Semprebon for useful discussions, and the Biophysical Sciences Institute in Durham for a summer studentship for MRP. This work is funded by EPSRC (EP/J017566/1).

-
- [1] J. M. Seddon and R. H. Templer, in *Structure and dynamics of membranes: From cells to vesicles*, Handbook of Biological Physics, Vol. 1, edited by R. Lipowsky and E. Sackmann (Elsevier Science Publishers B.V., 1995) Chap. 3.
- [2] L. Sagalowicz and M. E. Leser, *Curr. Opin. Colloid Interface Sci.* **15**, 61 (2010).
- [3] S. C. Kapfer, S. T. Hyde, K. Mecke, C. H. Arns, and G. E. Schröder-Turk, *Biomaterials* **32**, 6875 (2011).
- [4] C. Leal, N. F. Boussein, K. K. Ewert, and C. R. Safinya, *J. Am. Chem. Soc.* **132**, 16841 (2010).
- [5] R. H. Templer, J. M. Seddon, N. A. Warrender, A. Syrykh, Z. Huang, R. Winter, and J. Erbes, *J. Phys. Chem. B* **102**, 7251 (1998).
- [6] B. Angelov, A. Angelova, M. Ollivon, C. Bourgaux, and A. Campitelli, *J. Am. Chem. Soc.* **125**, 7188 (2003).
- [7] A. I. I. Tyler, H. M. G. Barriga, E. S. Parsons, N. L. C. McCarthy, O. Ces, R. V. Law, J. M. Seddon, and N. J. Brooks, *Soft Matter* **11**, 3279 (2015).
- [8] A. D. Benedicto and D. F. O'Brien, *Macromolecules* **30**, 3395 (1997).
- [9] S. T. Hyde and G. E. Schröder-Turk, *Interface Focus* **2**, 529 (2012).
- [10] A. H. Schoen, *Interface Focus* **2**, 638 (2012).
- [11] M. Caffrey, *Acta Crystallogr. F Struct. Biol. Commun.* **71**, 3 (2015).
- [12] S. L. Veatch and S. L. Keller, *Biophys. J.* **85**, 3074 (2003).
- [13] R. Parthasarathy, C.-H. Yu, and J. T. Groves, *Langmuir* **22**, 5095 (2006).
- [14] T. Baumgart, A. T. Hammond, P. Sengupta, S. T. Hess, D. A. Holowka, B. A. Baird, and W. W. Webb, *Proc. Natl. Acad. Sci. USA* **104**, 3165 (2007).
- [15] D. A. Brown and E. London, *Annu. Rev. Cell Dev. Biol.* **14**, 111 (1998).
- [16] H. T. McMahon and J. L. Gallop, *Nature* **438**, 590 (2005).
- [17] R. Lipowsky, *Nature* **349**, 475 (1991).
- [18] W. H. Meeks, *Indiana U. Math. J.* **39**, 877 (1990).
- [19] M. Fuhrmans, V. Knecht, and S. J. Marrink, *J. Am. Chem. Soc.* **131**, 9166 (2009).
- [20] U. S. Schwarz and G. Gompper, *Phys. Rev. E* **59**, 5528 (1999).
- [21] W. B. Lee, R. Mezzenga, and G. H. Fredrickson, *Phys. Rev. Lett.* **99**, 187801 (2007).
- [22] S.-D. Yang, H. G. Lee, and J. Kim, *Computer. Physics. Comm.* **181**, 1037 (2010).
- [23] D. Frenkel and B. Smit, *Understanding Molecular Simulation: From Algorithms to Applications* (Academic Press, 2001).
- [24] U. Dierkes, S. Hildebrandt, A. Küser, and O. Vohlrab, *Minimal Surfaces I: Boundary Value Problems* (Springer, 2013).
- [25] P. Gandy and J. Klinowski, *Chem. Phys. Lett.* **322**, 579 (2000).
- [26] R. D. Kamien, *Rev. Mod. Phys.* **74**, 953 (2002).
- [27] S. Semrau, T. Idema, L. Holtzer, T. Schmidt, and C. Storm, *Phys. Rev. Lett.* **100**, 088101 (2008).
- [28] T. Baumgart, S. T. Hess, and W. W. Webb, *Nature* **425**, 821 (2003).
- [29] G. C. Shearman, O. Ces, R. H. Templer, and J. M. Seddon, *J. Phys.: Condens. Matter* **18**, S1105 (2006).
- [30] H. M. G. Barriga, A. I. I. Tyler, N. L. C. McCarthy, E. S. Parsons, O. Ces, R. V. Law, J. M. Seddon, and N. J. Brooks, *Soft Matter* **11**, 600 (2015).
- [31] Z. A. Almshergqi, S. D. Kohlwein, and Y. Deng, *J. Cell Biol.* **173**, 839 (2006).
- [32] Y. Deng, M. Marko, K. F. Buttler, A. Leith, M. Mieczkowski, and C. A. Mannella, *J. Struct. Biol.* **127**, 231 (1999).
- [33] S. B. Rizwan, B. J. Boyd, T. Rades, and S. Hook, *Expert Opin. Drug Deliv.* **7**, 1133 (2010).
- [34] N. Garti, D. Libster, and A. Aserin, *Food Funct.* **3**, 700 (2012).
- [35] L.-L. Pontani, M. F. Haase, I. Raczowska, and J. Brujic, *Soft Matter* **9**, 7150 (2013).
- [36] M. Böltau, S. Walheim, J. Mlynek, G. Krausch, and U. Steiner, *Nature* **391**, 877 (1998).
- [37] L. Xue, J. Zhang, and Y. Han, *Prog. Polym. Sci.* **37**, 564 (2012).
- [38] Reinhard Lipowsky, *J. Phys. II France* **2**, 1825 (1992).
- [39] M. Hu, J. J. Briguglio, and M. Dserno, *Biophys. J.* **102**, 1403 (2012).
- [40] J. J. Williamson and P. D. Olmsted, *Biophys. J.* **108**, 1963 (2015).