

1 *Title:*

2

3 Changing patterns of daily precipitation totals at the Coweeta Hydrologic Laboratory,
4 North Carolina, USA

5

6

7 *Authors and Affiliations:*

8

9 T. P. Burt^{1*}, C. Ford Miniati^{2,2}, S. H. Laseter^{2,3} and W. T. Swank²

10

11 ¹Department of Geography, Durham University, Durham, DH1 3LE, UK

12 ²USDA Forest Service, Southern Research Station, Coweeta Hydrologic Lab, Otto,
13 NC 28763 USA

14 ³USDA Forest Service, Southern Research Station, Center for Integrated Forest
15 Science, Franklin, NC 28734 USA (current address)

16

17 *Author for correspondence: t.p.burt@durham.ac.uk

18

19

20 **Abstract**

21

22 A pattern of increasing frequency and intensity of heavy rainfall over land has been
23 documented for several temperate regions and is associated with climate change. This
24 study examines the changing patterns of daily precipitation at the Coweeta
25 Hydrologic Laboratory, North Carolina, USA, since 1937 for four rain gauges across
26 a range of elevations. We analyse seasonal total rainfall, number of rain days and the
27 frequency of heavy rainfall. We compare these with several teleconnections, including
28 the Bermuda High Index, the West Bermuda High Index, the North Atlantic
29 Oscillation and the El Niño-Southern Oscillation. Our data show a tendency for
30 increased variability, including major periods of drought, with fewer rain days
31 recently, especially in summer. Only autumn tended to have increases in rainfall
32 frequency and magnitude; this is the season when orographic enhancement is at its
33 strongest. The major driver of precipitation at Coweeta is the strength of the Bermuda
34 High. The strength of the North Atlantic Oscillation is important in summer. The
35 results are relevant to the Southeast USA in general, given that the region comes
36 under the influence of similar air masses during the year. The findings are applicable
37 to the wider Appalachian Mountains and to other mountainous regions where there is
38 significant orographic enhancement.

39

40 Key words: rainfall, Coweeta, orographic enhancement, Bermuda High

41 Introduction

42

43 Globally, precipitation extremes are increasing, a result of increasing air temperatures
44 at low elevations leading to increased evaporation and higher water vapour content in
45 the atmosphere (Meehl, et al., 2007; Allan and Soden, 2008; O’Gorman and
46 Schneider, 2009; Hartmann *et al.*, 2013). Because of the dependence of the saturation
47 water vapor pressure on temperature, as predicted by the Clausius-Clapeyron
48 equation), a 1°C temperature increase leads to an approximate 7% increase in the
49 moisture-holding capacity of the atmosphere. Since about 1950 the number of heavy
50 precipitation events over land has increased, and in North America the frequency or
51 intensity of heavy precipitation has increased, depending on season and region
52 (Hartmann *et al.*, 2013). Since 1910, precipitation across the USA has increased by
53 about 10%, again due to increases in the frequency and intensity of very large
54 precipitation events, with the proportion of total annual precipitation derived from
55 heavy rainfall increasing (Karl and Knight, 1998). This pattern of increasing
56 frequency and intensity of heavy rainfall holds true for the southern Appalachian
57 region within the USA as well (Karl and Knight, 1998). Changes in the proportion of
58 precipitation derived from heavy rainfall (variation) without a proportional increase in
59 the total annual rainfall (mean) must result in increases in the number of dry days.
60 Indeed, for the Southeast (SE) USA a significant increase in dry days during summers
61 is occurring along with increases in heavy rainfall events (Wang *et al.*, 2010). Such
62 variability in dry days has been attributed to changes in the location and intensity of
63 the western ridge of the North Atlantic (“Bermuda”) subtropical high (Li *et al.*, 2011)
64 and in the greater variability in the North Atlantic Oscillation (NAO) since about
65 1950 (Hurrell *et al.*, 2003).

66 The NAO is an important mode of climatic variability in the northern
67 hemisphere including the eastern United States (Hurrell *et al.*, 2003; Folland *et al.*,
68 2009). The strength and position of the NAO varies seasonally, generally expanding
69 over much of the North Atlantic during winter and contracting and shifting poleward
70 during summer. Summer streamflow east of the Appalachians from New England to
71 the Gulf of Mexico is positively correlated with the strength of the NAO (Coleman
72 and Budikova, 2013). A tendency for drier summers in the southern Appalachians
73 (Laseter *et al.*, 2012) might therefore be associated with a long-term decline in
74 summer NAO since the middle of the 20th Century. NAO is in part affected by the
75 behaviour of the Bermuda High, as the NAO depends both on the Bermuda High and
76 the Icelandic Low. Diem (2013) notes that the Bermuda High Index (BHI) had a
77 significant positive trend from the 1970s to 2009, and the western ridge of the
78 Bermuda High moved significantly south-eastward from approximately the mid-
79 1970s to 2009.

80 Complex terrain can also influence precipitation totals, through orographic
81 enhancement and interactions with cloud and fog layers. Integrated analysis of model
82 simulations and observations have demonstrated how low-level feeder-seeder
83 interactions play an important role in orographic enhancement in the southern
84 Appalachians (Hill *et al.*, 1981; Wilson and Barros, 2014). Rainfall at higher
85 elevations is characterised by higher rainfall intensity. This is particularly important in
86 the cooler seasons when convective activity is less vigorous (Prat and Barros, 2010).
87 Stratiform rainfall systems can also interact with local low-level clouds and fog, the
88 latter being persistent at high altitudes in this region (Wilson and Barros, 2014). This
89 can help account for systematic differences in rainfall totals between valley locations
90 and ridge tops. On the other hand, the passage of warm-season intense storm systems

91 including tropical depressions and cool-season synoptic-scale storms may produce
92 rainfall distributions much less well correlated with topography. Localised
93 thunderstorm activity would have the same effect, possibly even favouring (warmer)
94 valleys over (cooler) ridge tops if the valleys are where the storm cells are generated.
95 Interesting diurnal variations emerge seasonally with more afternoon rainfall in
96 summer, as might be expected.

97 Orographic controls on precipitation can also interact with larger-scale climate
98 modes. For example, Coleman and Budikova (2013) suggest that in summer, when
99 the NAO is positive, orographic uplift and local convection occur along the east side
100 of the Appalachians. For the British Isles, Burt and Howden (2013) showed that
101 variations in the strength of the NAO, in this case influenced mainly by the strength of
102 the Icelandic Low, cause large differences in seasonal precipitation totals compared
103 to NAO-neutral conditions, an effect which is amplified with altitude - what Burt and
104 Howden (2013) term “double orographic enhancement”. Similar effects have been
105 noted for the mountains of the Pacific Northwest US and Sri Lanka, both linked to
106 variations in the El Niño-Southern Oscillation (ENSO: Burt and Howden, 2013). In
107 all such cases, it is the movement of storm systems across the upland area which is
108 important in generating the orographic effect. Hurrell and Van Loon (1997)
109 emphasise that studies of long-term trends of temperature or precipitation in local
110 climate records, especially those from high-elevation sites, should also examine
111 parallel changes in atmospheric circulation in order to fully synthesize the available
112 information about regional or global climate change.

113 Our purpose here is to analyse the long-term, *daily* precipitation records at the
114 Coweeta Hydrologic Laboratory, one of the wettest locations in SE USA. Coweeta is
115 among the oldest, continuously operating environmental study sites in North America
116 (Swank and Crossley, 1988; Laseter *et al.*, 2012). Coweeta has observed increased
117 precipitation variability in recent decades with summer months becoming drier over
118 time and autumn months becoming wetter (Laseter *et al.*, 2012). Given its location in
119 the SE USA, Coweeta is particularly influenced by atmospheric conditions prevailing
120 across the Atlantic Ocean. Our analysis includes examining variation with altitude,
121 and possible drivers such as the Bermuda High and the NAO, including the possible
122 amplification of precipitation totals at high altitude. Given Coweeta’s location, it is
123 anticipated that the Bermuda High will be a major driver of precipitation, with a more
124 direct linkage compared to the NAO, However, we also include other climatic indices
125 to investigate whether other teleconnections might exist, in particular links to
126 conditions in the Pacific Ocean. The findings may be relevant to other mountainous
127 regions which are subject to significant orographic enhancement driven by strong
128 atmospheric circulation across a topographic barrier.

129 **Site description and methods**

130

131 *Study Site*

132 The USDA Forest Service, Coweeta Hydrologic Laboratory (2185 ha) lies in the
133 Nantahala Mountains of western North Carolina, USA (35° 03' N, 85° 25' W: Figure
134 1). Established in 1934, it has been a testing ground for theoretical and applied forest
135 hydrological research (Swank and Crossley, 1988). Climate in the Coweeta Basin is
136 classified as perhumid, mesothermal with water surplus in all seasons (Swift *et al.*,
137 1988). Precipitation comprises frequent, low-intensity rain in all seasons with little
138 snow. Occasional intense storms are associated with severe thunderstorms or inland
139 influences from hurricanes (Shepherd *et al.*, 2007). For all rain-gauge sites at
140 Coweeta, March is the wettest month (193 mm) and October is the driest (112 mm).

141

142 *Daily precipitation measurements*

143 Long-term climate and precipitation have been monitored at Coweeta through a
144 network of gauging stations at different elevations and aspects since 1934 (Swift *et*
145 *al.*, 1988). Parameters include temperature, rainfall, wind speed and direction,
146 humidity, solar radiation, air quality, and evaporation. A network of 12 standard rain
147 gauges (SRG, 8-inch Standard Rain Gauge, NWS) and nine recording rain gauges
148 (RRG, Belfort Universal Recording Rain Gauge, Belfort Instrument Co., Baltimore,
149 MD, USA) are currently located throughout the basin; four recording rain gauges are
150 included in this study (Table1). Total precipitation is measured by the SRG; volume
151 and intensity are measured by the RRG. The longest running climate station is CS01
152 at the valley floor, with a co-located recording rain gauge (RRG06). Here,
153 measurements started in 1937. CS01 is also a cooperative station with the National
154 Weather Service. Measurements at the other three (higher-altitude) gauges started as
155 follows: RG13 (1942), RG96 (1943) and RG31 (1958). Complete daily records exist
156 for all four gauges from the start date to present. There is some doubt about the
157 problem of under-catch, in two regards. Hibbert (1966, quoted in Swift *et al.*
158 1998) noted that precipitation totals increase fairly regularly with altitude except near
159 ridge-tops where catch was significantly lower than immediately downslope:
160 reductions of about 30% in a 30 m change of elevation extending about 100 m either
161 side of a ridge. It is possible therefore that totals at RRG31 are less than they should
162 be. There is also some doubt about totals at RRG40 compared to RRG96 because of
163 differential exposure in relation either to topography or shelter from nearby trees (see
164 below). Of more significance is the problem of under-catch for the very lowest
165 observed totals. Histograms of daily precipitation totals should show a smooth,
166 negative exponential distribution but all four gauges show a lower total for catches of
167 0.254 mm (0.01 inch) compared to the next category 0.508 mm (0.02 inch). This is
168 not likely to be a problem unique to Coweeta: for example at Oxford, England, there
169 is the same under-catch for both hourly (tipping bucket) and daily (manual
170 observation) data. Under-catch of the smallest amounts is not at all significant for
171 long-period totals, given that such small amounts are involved, but may well be for
172 counts (Burnette and Stahle, 2013). For that reason, the frequency of wet days (at
173 least 1 mm total) is used here in subsequent analyses instead of rain days.

174

175 *Precipitation Analysis*

176 Analysis of heavy rainfall, as indicated by daily totals, follows Osborn *et al.* (2000)
177 and Maraun *et al.* (2008). We ranked all daily rainfall data, accumulating the totals,
178 and identified the group of highest daily totals that together contributed 10% of the

179 total precipitation (henceforth T10). Following Osborn *et al.* (2000), the cumulative
 180 rainfall totals are only calculated for rain days, defined as a day with total of 0.25 mm
 181 or more. For a long daily rainfall series, only a very small fraction of total rainfall is
 182 thereby excluded. T10 values for the four gauges are listed in Table 1; they range
 183 from 74.9 mm at the lowest gauge to 90.9 mm at the highest. Other precipitation
 184 thresholds examined include daily totals of 25 and 50 mm and the upper quartile (Q3)
 185 for all rain day totals at RRG06: 17.8 mm. Measures of dispersion include the lower
 186 quartile (Q1) and the upper quartile (Q3) and the 10th and 90th percentiles. Seasons are
 187 defined as: winter (Dec–Feb: DJF), spring (Mar–May: MAM), summer (Jun–Aug:
 188 JJA) and autumn (Sep–Nov: SON). Analyses for all four gauges started on 1st
 189 January for the first available complete year of record (Table 1) until 31st December
 190 2012. We used 20-year moving windows (Burt *et al.*, 2008) to fit percentiles and
 191 quartiles to the data series ($n = 57$ for the moving-window correlations). We also use
 192 running monthly totals to identify major periods of below-average rainfall, as trends
 193 in accumulated deficit can be misleading (Linacre, 1992). Use of 24-month totals
 194 highlights more protracted droughts compared to short-term deficits. For analyses of
 195 orographic enhancement, because all rain gauges do not have the same length of
 196 record, we truncate the records for a common period (1959–2013) to allow
 197 comparison.

198

199 *Regional Atmospheric and Sea Surface Temperature Data*

200 NAO data were obtained from the Climatic Research Unit (UEA CRU, 2015). The
 201 NAO is defined as the normalized pressure difference between a station on the Azores
 202 (Ponta Delgada) and one on Iceland (Reykjavik). Seasonal values of the BHI and the
 203 West Bermuda High Index (WBHI) were obtained from Diem (2013). The BHI is an
 204 index of the difference between normalized sea-level pressure over Bermuda and New
 205 Orleans, Louisiana (Stahle and Cleaveland, 1992); seasonal sea-level pressure values
 206 are for grid cells corresponding to 32.5°N, 64°W and 30°N, 90°W. The centre of the
 207 Bermuda High is typically located approximately 2500 km east of Bermuda during
 208 summer; therefore, the BHI shows the standardized pressure gradient across the
 209 western side of the Bermuda High. The WBHI is based on the BHI; it uses 850-hPa
 210 heights and is centred over the Southeast United States. Seasonal 850-hPa
 211 geopotential heights are for grid cells corresponding to 30°N, 75°W and 30°N, 92°W
 212 (Diem, 2013). Other proxy measures of atmospheric circulation used here include the
 213 NINO12, NINO3 and NINO34 indices, which are measures of sea-surface
 214 temperature (SST) anomalies in the Pacific Ocean (Kaplan *et al.*, 1998) and the
 215 Pacific North America (PNA) index. The NAO index tends to be easier to interpret
 216 physically, but is based on only two arbitrary stations. PNA data (starts in 1950; $n =$
 217 63 here), were downloaded (NOAA National Weather Service, 2015), as were
 218 extended SST anomaly data for equatorial regions of the Pacific based on Kaplan *et*
 219 *al.* (1998) (IRI LDEO, 2015). The regions included are the composite regions
 220 NINO12 (0–10°S, 80–90°W) and NINO34 (120–170°W, 5°S–5°N) plus the core
 221 region NINO3 (150°W–90°W, 5°S–5°N). We derive relationships between seasonal
 222 totals of precipitation statistics and averages of climatic indices using ordinary least
 223 squares regression. We also undertook a composite analysis (Bulic and Kucharski,
 224 2012), identifying mean summer rainfall for all years that were more than ± 1 standard
 225 deviation from mean summer NAO. We conducted a *t*-test on these composite data.

226 **Results**

227

228 *Temporal and Spatial Patterns*

229 While precipitation in the Coweeta Basin was not characterized by pronounced
230 seasonality, seasonal differences did exist (Table II). Although summer had the most
231 rain days, winter was the wettest season; while autumn was the driest. Summer had
232 the lowest mean rainfall per rain day and the lowest number of days with totals above
233 25 mm. The highest daily total (196.9 mm) was recorded on 4th October 1964.

234 There were almost no significant trends in rainfall regime in winter, spring and
235 summer, but autumn has become wetter with heavier rainfall more common (Table
236 III). Laseter *et al.* (2012) noted a long-term increase in September rainfall totals,
237 which is clearly a part of this autumn trend. RRG96 had significant positive trends in
238 autumn for total rainfall plus the number of days above 25 mm and above the T10 mm
239 threshold. RRG40 had positive trends only in autumn for the T10 mm threshold. It
240 had negative trends for numbers of rain days per year and the number of days above
241 25 mm in winter; it also had significant negative trends for rain days in spring,
242 summer and for the year as a whole. At the highest-elevation gauge, RRG31, there
243 were significant trends for the number of days above 25 mm in winter (negative) and
244 the number of days above the T10 in summer (positive). Note that, for all four gauges,
245 other than for T10, all the significant trends were negative except those in autumn
246 which were uniformly positive. Results for the T10 index are of limited value given
247 large numbers of zeroes in the record.

248

249 *Variability*

250 Laseter *et al.* (2012) noted a tendency for annual precipitation totals to become more
251 variable over time, with wetter wet years and drier dry years. At RRG06 (Figure 2)
252 there were downward trends ($p < 0.001$) for the 10th percentile ($r = 0.89$) and Q1 ($r =$
253 0.77), and an upward trend for Q3 ($r = 0.66$; Figure 2). Thus, variability over time
254 increased, with increases in the interquartile range (IQR) over time ($r = 0.83$, $p <$
255 0.001). Up to 1974, only one year had a rainfall total more than one standard
256 deviation below the mean; since 1975 there were eight such years. Totals above one
257 standard deviation were similar, seven up to 1974, but only six since. Totals above
258 two standard deviations increased from two to four in the latter half of the study
259 period. Downward trends for the Q1 in winter, spring and summer ($r > 0.69$ and $p <<$
260 0.0001 ; and for Q3 in winter ($r = 0.73$, $p << 0.0001$) and summer ($r > 0.32$, $p < 0.05$)
261 were also observed. In contrast, autumnal Q1 and Q3 trended upward for total rainfall
262 ($r > 0.77$ and $p << 0.0001$ for both seasons).

263 The IQR for numbers of wet days narrowed in winter ($r = -0.3$) and autumn (r
264 $= -0.54$) but widened very significantly in summer ($r = 0.82$). Q1 for numbers of wet
265 days fell significantly in all seasons except autumn and for the year as a whole ($r = -$
266 0.55), but increased significantly in autumn ($r = 0.42$). Q3 for numbers of wet days
267 decreased significantly in winter ($r = -0.54$) but increased significantly in summer ($r =$
268 0.35). The changing frequency of numbers of wet days very much accords with
269 changes in totals, of course, especially in relation to wetter autumns and more variable
270 summers. Negative trends in numbers of rain days in summer and for the year as a
271 whole are not seen for numbers of wet days (Figure 3b). The results for numbers of
272 rain days deserves further attention and might relate to warmer summers, a reduction
273 in the occurrence of light showers, or some other cause. Comparison with other
274 gauges across widely different climatic regimes could also be worthwhile.

275

276 *Intensity*

277 There was no trend for mean rainfall per day except in autumn ($r = 0.27, p <$
 278 0.05), which also was the only season with a significant change (increase) in rainfall
 279 total over time (Table III). Variation in mean rainfall per wet day (as indicated by IQR)
 280 also increased over time in both autumn ($r = 0.27, p < 0.05$) and for the year as a
 281 whole ($r = 0.46, p < 0.001$), whereas variation in summer decreased significantly ($r =$
 282 $-0.68, p < 0.0001$). There was no significant change in winter or spring.

283 Although the number of daily falls above 25 mm each year did not change
 284 significantly over the study period, variation increased over time (IQR, $r = 0.37, p <$
 285 0.01). Autumn was the only season with a significant (positive) trend in numbers of
 286 daily totals above 25 mm (Table IV). All four measures of dispersion used here (Q1,
 287 Q3 and the 10th and 90th percentiles) and the IQR ($r = 0.596, p < 0.001$) increased in
 288 autumn over time. In summer, the Q3 and the 90th percentile increased whilst the 10th
 289 percentile decreased; accordingly the IQR increased ($r = 0.48, p < 0.001$). In winter all
 290 measures of dispersion decreased over time, but the IQR increased ($r = 0.32, p <$
 291 0.05). Spring IQR decreased for numbers of days above 25 mm ($r = -0.43, p < 0.01$).
 292 The results for the upper quartile value 17.8 mm were very similar to those for the 25
 293 mm threshold. Correlations for totals above 50 mm and for the T10 index were
 294 hampered by large numbers of zeroes, but numbers of T10 events increased in autumn
 295 ($r = 0.28, p < 0.05$). Maximum daily rainfall in autumn increased over the 76-year
 296 period of analysis ($r = 0.34, p < 0.01$), and the variability of heavy rainfall increased
 297 in all seasons except spring (Table V). Autumn was the only season where Q1 went
 298 up, but even so, there was a significant increase in the IQR, as a result of a strongly
 299 upward trend for Q3. In winter both Q1 and Q3 fell but nevertheless the IQR
 300 increased. The overall pattern for the year as a whole was an increase in IQR
 301 therefore, with the strongest effect seen in autumn.

302

303 *Drought*

304 Notwithstanding some evidence of more intense precipitation recently, greater
 305 variability (as shown by significant increases in IQR) implies a trend towards more
 306 frequent periods of prolonged drought; and indeed, other than one event in the early
 307 1940s, severe droughts were a feature of the Coweeta climate since the mid-1980s
 308 (Figure 3a). The major periods of rainfall deficit were more clearly related to the
 309 number of heavy falls of rain than to the number of wet days. Whilst, as noted above,
 310 there was no significant long-term trends in the number of daily falls over 25 mm,
 311 variability clearly increased, matching the pattern for rainfall total (Figure 3c).

312

313 *Orographic enhancement*

314 Mean rainfall increased with altitude (Table IV). The gradient in precipitation
 315 between the lowest and highest gauges (RRG06 and RRG31) was 85 mm per 100
 316 metres. The intermediate gauges RRG96 and RRG13 showed differences in the
 317 gradient (99 mm/100m and 60 mm/100m, respectively), likely due to differences in
 318 gauge exposure (Swift et al., 1988) or shelter by surrounding trees. Daily rainfall data
 319 for ridge top and valley bottom locations was relatively similar in all seasons except
 320 autumn, where the gradient is larger (Table VI). The highest daily rainfall for any
 321 Coweeta rain gauge was recorded at the high-elevation gauge RRG31, 232.4 mm on
 322 28th May 1976. There was very little difference in the frequency of rain days at the
 323 ridge top and valley bottom gauges, but given higher autumn totals and lower number
 324 of rain days, the gradient in rain per rain day was significantly higher in autumn
 325 compared to other seasons. The frequency of heavy falls of rain also increased with

326 altitude, although there was little seasonal variation. Whilst totals were often similar
 327 for both RRG31 and RRG06 and there were some days where totals were higher at
 328 the latter low-altitude gauge, the largest differences in total were always in favour of
 329 the high-altitude gauge (Table VII).

330

331 *Atmospheric drivers*

332 Given Coweeta's location in the Southeast USA, strong linkage between the Bermuda
 333 High and Coweeta precipitation is to be expected. We employed two indices of
 334 atmospheric circulation in relation to the Bermuda High: BHI and WBHI (Diem,
 335 2013). Results reported are for the period 1959-2013 ($n=55$). Both the BHI and the
 336 WBHI were positively related to rainfall totals, number of wet days, and the number
 337 of daily rainfalls above 25 mm. Total rainfall at RRG06 was positively related to BHI
 338 in three seasons ($r > 0.39$ and $p < 0.001$: winter, summer and autumn); and to WBHI
 339 in all four seasons ($r > 0.43$ and $p < 0.00003$). The same pattern was seen for number
 340 of wet days: BHI had significant correlations in three seasons ($r > 0.41$ and $p < 0.002$
 341 for winter, summer and autumn), and WBHI had significant correlations in all four
 342 seasons ($r > 0.58$, $p < 0.00001$ for all four seasons). In contrast to Diem (2013), who
 343 found a decline in the number of rain days since the 1970s in the Atlanta region, the
 344 seasonal correlations for both BH indices from 1959–2013 at Coweeta are positive.
 345 This may either relate to different rainfall-generation processes being dominant in the
 346 mountains compared to the piedmont, or more likely reflect the longer window of
 347 analysis used here (Howden *et al.*, 2011). Daily rainfalls totalling more than 25mm
 348 were positively related to BHI in winter and summer ($r > 0.3$, $p < 0.00009$), and to
 349 WBHI in all four seasons ($r > 0.36$ and $p < 0.007$).

350 NAO was a strong atmospheric driver of summer precipitation at Coweeta.
 351 Total summer precipitation was positively correlated with the NAO ($r = 0.37$, $p =$
 352 0.005). Using detrended NAO data made no difference in the correlation with summer
 353 rainfall ($r = 0.38$). Summers with very high NAO (> 0.72 , one standard deviation
 354 (SD) above the mean) and those with very low NAO (< -0.87 , -1 SD) produced
 355 significantly different rainfall totals ($p = 0.002$). Summer NAO declined over time.
 356 The summers of 2009–2012 all had a very low NAO, but conditions reversed in
 357 summer 2013, with a seasonal total of 708 mm, and 9 daily totals over 25 mm. Not
 358 only was NAO correlated with rainfall indices, but the strength of the relationship was
 359 amplified with altitude (Table VIII). With one exception, the regression coefficients
 360 were larger at RRG31; however, only four of the six correlations were stronger at the
 361 high-altitude site. A very similar pattern was seen using BHI: only one regression
 362 coefficient was lower at the high-altitude gauge (number of wet days in winter) and
 363 only two results had lower correlation coefficients at the high-altitude gauge (winter
 364 totals and number of heavy falls in summer, Table IX).

365 Winter rainfall was significantly correlated with the average NAO value for
 366 the previous calendar year ($r = 0.319$, $p = 0.005$) and the previous summer ($r = 0.23$, p
 367 $= 0.04$). The same lagged relationships pertained for daily totals over 25 mm. Much
 368 the same pattern of correlations was found at all gauges and the PNA index: total
 369 winter rainfall was negatively correlated to the PNA index ($r = -0.29$, $p = 0.023$, $n =$
 370 62), as was spring total rainfall and previous winter's PNA ($r = -0.32$). There was also
 371 a positive relationship between for winter PNA and number of rain days ($r = 0.348$, p
 372 $= 0.006$), but none for heavy rainfall. The Pacific Ocean indices NINO12 and
 373 NINO34 produced very few significant correlations with Coweeta rainfall statistics
 374 (RRG06) although the number of wet days in autumn was weakly related to both
 375 NINO12 ($r = 0.22$, $p = 0.054$, $n = 76$) and NINO34 ($r = 0.21$, $p = 0.066$).

376

377 **Discussion**

378 Summer was the only season positively correlated with the NAO in the same season
379 (*i.e.*, zero lag); this was seen at all gauges for total and number of days above 25 mm.
380 This matches the results for streamflow found by Coleman and Budikova (2013).
381 During a positive NAO phase, the pressure gradient between the subtropical and polar
382 regions of the North Atlantic intensifies, the Bermuda High extends westward and
383 south-easterly flow advects the moist air masses inland over the Southern
384 Appalachians. Coleman and Budikova (2013) suggest that this may generate localised
385 convection and orographic uplift along the east side of the Appalachian ridge.
386 Tropical storms deflected around the Bermuda High provide additional, often very
387 heavy, falls of rain. Negative NAO conditions see the Bermuda High considerably
388 weakened, associated with drier northerly air flow (Coleman and Budikova, 2013;
389 Figure 7). The trend since the 1930s has been for summer NAO to decrease but, as
390 noted above, summer rainfall variables at Coweeta also correlate with the detrended
391 NAO index. This gives confidence in a causal relationship, not simply an association
392 related to both variables being collinear with an independent variable, *e.g.* rising
393 summer temperatures. Since inter-annual variations in summer NAO correlate with
394 Coweeta summer rainfall, there is a straightforward explanation in terms of the
395 strength of the Bermuda High and the associated pressure gradient across the polar
396 front.

397 Indeed, rainfall models reveal the Bermuda High to be an important predictor
398 of the precipitation regime at Coweeta (cf. Diem, 2013). At all four gauges, the WBHI
399 was strongly correlated with totals, numbers of wet days and numbers of heavy falls
400 in all four seasons. Of 48 possible correlations (4 gauges, 4 seasons, 3 rainfall
401 variables), 48 were statistically significant. BHI was less powerful as a predictor (34
402 significant correlations) because it never correlated with any of the spring rainfall
403 variables. Diem (2013) showed that increased rainfall variability in the Southeast
404 USA related to increased variability in the WBHI. Extreme negative values of the
405 WBHI occur when the Bermuda High extends over much of the Southeast, and *vice*
406 *versa*. Further work is needed to understand why BHI does not predict spring rainfall
407 at Coweeta.

408 Autumn showed a general pattern of increased precipitation over time, unlike
409 the other seasons where, if there were any trend at all, it tended to be downwards.
410 Autumn totals and numbers of falls above 25 mm showed positive trends at RRG06
411 and RRG40; all four gauges showed upward trends for mean rain per rain day in
412 autumn. These positive trends in autumn were accompanied by increased variability:
413 given 4 gauges, 3 indices of variability (Q1, Q3, IQR) and 4 variables (total, wet days,
414 falls above 25 mm, T10), autumn showed positive trends for 34 out of a possible 48
415 correlations. Excluding numbers of wet days, which tended not to be collinear with
416 other variables, there were significant positive trends for 31 out of 36 possibilities.
417 Even so, autumn rainfall was poorly correlated with atmospheric drivers: only number
418 of wet days correlated weakly with NINO12 and NINO34. Notwithstanding a general
419 context of increasing variability, autumn rainfall totals tended to increase, with heavy
420 falls of rain more frequent. During the El Niño portion of ENSO, increased
421 precipitation falls along the Gulf coast and Southeast USA due to a stronger than
422 normal and more southerly polar jet stream. Both summer and autumn NAO declined
423 over time (this could relate to changes in either or both the Bermuda High and the
424 Icelandic Low), which may have enabled a greater influence of Pacific Ocean
425 conditions across the Caribbean and the Southeast USA. This merits further attention

426 and, as well as looking for more relevant indices of atmospheric circulation for this
427 region and season, it might be fruitful to analyse objectively-derived weather types
428 (Kalnay *et al.*, 1996; Compo *et al.*, 2011) to see why the autumnal rainfall regime has
429 changed at Coweeta (cf. Diem, 2013, Table III, which is for summer only). This is
430 relevant since autumn rainfall shows the greatest rate of orographic enhancement of
431 any season (see also below); the long-term increase in autumn rainfall may well have
432 implications for downstream flood generation given the importance of autumnal
433 rainfall for recharge of soil moisture deficits. The results at Coweeta do not accord
434 with those reported in Shepherd *et al.* (2007) because at Coweeta winter and spring
435 have the highest number of days with heavy falls of rain (daily rainfall totals over 25
436 mm). Even so, the upward trend in autumn for the total of daily falls over 25 mm may
437 well relate to the increase in intense hurricane activity discussed by Shepherd *et al.*
438 (2007). Whilst autumn rainfall was more variable over time (Table V), there was
439 nevertheless a significant upward trend in the number of extreme rainfall events (as
440 measured by the number of daily totals above 25 mm).

441 High winter rainfall totals reflect enhanced westerly airflow and the influence
442 of warm moist air from the Gulf of Mexico. A positive PNA is indicative of the
443 strength and location of the East Asian jet stream and, subsequently, the weather it
444 delivers to North America. As the magnitude of the positive PNA increases, the large-
445 scale weather pattern is increasingly meridional which implies a greater transport of
446 heat, moisture, and momentum between low and high latitudes over North America.
447 The strongest, most extensive correlations between the PNA index and precipitation
448 have been observed in winter and early spring (Leathers *et al.*, 1991). Winter was
449 notably the only season with a significant (negative) correlation between rainfall total
450 and the PNA index at RRG06, although there was also a significant correlation
451 between spring rainfall total and PNA in the preceding winter. None of the gauges
452 had a correlation with numbers of daily totals above 25 mm. Both Leathers *et al.*
453 (1991, Figure 3) and Yin (1994 Figure 7a) confirm that, given its location, Coweeta
454 should have a negative correlation with the PNA index. A positive PNA index sees
455 the polar jet further south and east than usual, resulting in a decrease in precipitation
456 amounts for a large part of central USA including the southern Appalachians.
457 Cyclones are less frequent in this region given changes in track; in addition, polar and
458 arctic continental air masses dominate the region, further decreasing precipitation.
459 With negative PNA index values, the polar front jet is pushed far to the north of its
460 mean position over eastern USA. This allows tropical maritime air masses from the
461 Gulf of Mexico to cross the region more often, increasing precipitation.

462 The gradient in annual precipitation between the lowest and highest gauges
463 was considerably steeper than any gradients measured in the UK (of which the
464 maximum is 45.8 mm/100m in Snowdonia (Unwin, 1969)). For a group of 19 gauges
465 in the Pacific NW USA, the average gradient was 66.2 mm/100 m (Burt and Howden,
466 2013). The result at Coweeta reflects its relatively southern location, its exposure to
467 warm, moist tropical air masses and the steep topography and high elevation locally.
468 Results suggest that autumn has the most favourable conditions conducive to
469 orographic enhancement, but the effect is strong in all seasons. The orographic effect
470 comprises seeder clouds in the form of passing depression systems and feeder clouds
471 in the form of low cloud and fog over the hills; Wilson and Barros (2014) detail the
472 mechanisms involved in this process. In general, the strength of the relationships
473 between NAO, BHI and rainfall are amplified with altitude (Tables VII-IX).
474 However, the results are not fully convincing: not all of the significant regression
475 coefficients or correlation coefficients are larger at RRG31, the high-altitude gauge.

476 This weakens the argument for a “double orographic enhancement” effect in that,
477 although the high-elevation site experiences a stronger absolute response to NAO
478 variability (roughly proportional to mean rainfall), there is not necessarily a stronger
479 correlation at altitude too. This implies that the underlying mechanisms are only to
480 some extent more effective at higher altitudes (cf. Burt and Howden, 2013). This
481 would merit further investigation using hourly recording gauge data and
482 accompanying microphysics measurements (Wilson and Barros, 2014). It is novel that
483 previous results obtained using the NAO index (Burt and Howden, 2013) have been
484 replicated here using BHI.

485 Observations of double orographic enhancement (Burt and Howden, 2013) at
486 Coweeta underline the importance of changes in atmospheric circulation for rainfall
487 generation. Total autumn rainfall increased in line with a weakening of the Bermuda
488 High, allowing more tropical maritime air masses to pass over the Coweeta region
489 from the south-west. The tendency in other seasons was for frequency and magnitude
490 to decline, in response to long-term trends in atmospheric circulation as measured by
491 the proxies NAO and PNA. The most marked orographic enhancement can be
492 expected when the atmospheric flow is at its strongest but the increase of PNA in
493 winter and the decrease of NAO in summer have had the opposite effect. Further
494 research on orographic enhancement at Coweeta using hourly rainfall data would be
495 beneficial.

496

497 **Conclusions**

498 Long-term trends in daily rainfall at the Coweeta Hydrologic Laboratory reflected
499 hemispheric changes in the global circulation. There has been a tendency for greater
500 variability in the WBHI (Diem, 2013), resulting in wetter wet years and drier dry
501 years. The Bermuda High was, not surprisingly, the main influence on precipitation
502 regime at Coweeta, with atmospheric circulation in the North Atlantic more generally
503 (as indicated by the NAO index) influential in summer. There is, of course, evidence
504 of the influence of air masses moving from the Gulf of Mexico (as indicated by the
505 PNA index) but direct associations with the Pacific Ocean were not generally seen.
506 Further work using synoptic weather types as well as circulation indices like WBHI
507 would seem worthwhile (cf. Diem, 2013; Burt *et al.*, 2015). The findings are
508 applicable to the wider southern Appalachian Mountains and to other mountainous
509 regions where there is significant orographic enhancement. The use of the BHI as well
510 as the NAO indices to demonstrate double orographic enhancement is novel, helping
511 to confirm the general applicability of the hypothesis (Burt and Howden, 2013).
512 Further work is needed to explore the downward trend in number of rain days in
513 summer and for the year as a whole, not seen in the wet days’ data. This may well be
514 an artefact of the data rather than a real change in the occurrence of light rainfall. If it
515 reflects the impact of higher temperatures on rain gauge catch, it would be good to see
516 whether this observation is replicated across widely different climatic regimes.

517

518 **Acknowledgements**

519 We thank all scientists, technicians and students who have contributed to the long-
520 term climate and hydrology data at Coweeta. Funding for this research was provided
521 by the USDA Forest Service, Southern Research Station; and the National Science
522 Foundation (NSF), Long-Term Ecological Research (LTER) program (award #DEB-
523 0823293). Any opinions, findings, conclusions or recommendations expressed in the
524 material are those of the authors and do not necessarily reflect the views of the USDA
525 or the NSF. We thank Drs. Doug Miller and Marshall Shepherd for their thorough

526 reviews of a draft manuscript, and Dr. Jeremy Diem for providing the BHI and WBHI
 527 data. The figures were drawn by Chris Orton, Design and Imaging Unit, Department
 528 of Geography, Durham University, UK. We thank two anonymous referees for their
 529 helpful critique of an earlier draft of this paper.

530

531 **References**

- 532 Allan RP, Soden BJ. 2008. Atmospheric warming and the amplification of
 533 precipitation extremes. *Science* **321**(5895):1481-1484. DOI
 534 10.1126/science.1160787
- 535 Barnston AG, Livezey RE. 1987. Classification, Seasonality and Persistence of Low-
 536 Frequency Atmospheric Circulation Patterns. *Monthly Weather Review*
 537 **115**(6):1083-1126. DOI 10.1175/1520-0493(1987)115<1083:Csapol>2.0.Co;2
- 538 Bulic IH, Kucharski F. 2012. Delayed ENSO impact on spring precipitation over
 539 North/Atlantic European region. *Climate Dynamics* **38**(11-12):2593-2612.
 540 DOI 10.1007/s00382-011-1151-9
- 541 Burnette DJ, Stahle DW. 2014. Historical perspective on the dust bowl drought in the
 542 central United States. *Climatic Change* **116**:479-494.
- 543 Burt T, Howden N, Worrall F. 2014. On the importance of very long-term water
 544 quality records. *Wiley Interdisciplinary Reviews: Water* **1**(1):41-48.
- 545 Burt TP, Howden NJK. 2013. North Atlantic Oscillation amplifies orographic
 546 precipitation and river flow in upland Britain. *Water Resources Research*
 547 **49**(6):3504-3515. DOI 10.1002/wrcr.20297
- 548 Burt TP, Howden NJK, Worrall F, Whelan MJ. 2008. Importance of long-term
 549 monitoring for detecting environmental change: lessons from a lowland river
 550 in south east England. *Biogeosciences* **5**(6):1529-1535.
- 551 Burt TP, Jones PD, Howden NJK. 2015. An analysis of rainfall across the British Isles
 552 in the 1870s. *International Journal of Climatology* **35**(10):2934-2947.
 553 10.1002/joc.4184
- 554 Coleman JSM, Budikova D. 2013. Eastern U.S. summer streamflow during extreme
 555 phases of the North Atlantic oscillation. *Journal of Geophysical Research-
 556 Atmospheres* **118**(10):4181-4193. DOI 10.1002/jgrd.50326
- 557 Compo GP, Whitaker JS, Sardeshmukh PD, Matsui N, Allan RJ, Yin X, Gleason BE,
 558 Vose RS, Rutledge G, Bessemoulin P, Bronnimann S, Brunet M, Crouthamel
 559 RI, Grant AN, Groisman PY, Jones PD, Kruk MC, Kruger AC, Marshall GJ,
 560 Maugeri M, Mok HY, Nordli O, Ross TF, Trigo RM, Wang XL, Woodruff
 561 SD, Worley SJ. 2011. The Twentieth Century Reanalysis Project. *Quarterly
 562 Journal of the Royal Meteorological Society* **137**(654):1-28. 10.1002/qj.776
- 563 Diem JE. 2013. Influences of the Bermuda High and atmospheric moistening on
 564 changes in summer rainfall in the Atlanta, Georgia region, USA. *International
 565 Journal of Climatology* **33**(1):160-172. 10.1002/joc.3421
- 566 Folland CK, Knight J, Linderholm HW, Fereday D, Ineson S, Hurrell JW. 2009. The
 567 Summer North Atlantic Oscillation: Past, Present, and Future. *Journal of
 568 Climate* **22**(5):1082-1103. DOI 10.1175/2008JCLI2459.1
- 569 Hartmann D, Klein Tank A, Rustucci M, Alexander L, Bronnimann S, Charabi Y,
 570 Dentener F, Dlugokencky E, Easterling D, Kaplan A. 2013. Observations:
 571 Atmosphere and Surface in: *Climate Change 2013: The Physical Science
 572 Basis. Contribution of working group I to the fifth assessment report of the
 573 intergovernmental panel on climate change. Cambridge University Press,
 574 Cambridge, United Kingdom and New York, NY, USA.*

- 575 Hill F, Browning K, Bader M. 1981. Radar and raingauge observations of orographic
576 rain over south Wales. *Quarterly Journal of the Royal Meteorological Society*
577 **107**(453):643-670.
- 578 Howden NJK, Burt TP, Worrall F, Whelan MJ. 2011. Monitoring fluvial water
579 chemistry for trend detection: hydrological variability masks trends in datasets
580 covering fewer than 12 years. *Journal of Environmental Monitoring*
581 **13**(3):514-521. 10.1039/c0em00722f
- 582 Hurrell JW, Kushnir Y, Ottersen G, Visbeck M. 2003. An overview of the North
583 Atlantic oscillation: Wiley Online Library.
- 584 Hurrell JW, VanLoon H. 1997. Decadal variations in climate associated with the
585 north Atlantic oscillation. *Climatic Change* **36**(3-4):301-326. DOI
586 10.1023/A:1005314315270
- 587 IRI LDEO. 2015. Indices nino EXTENDED [NINO12 , NINO3 , NINO34 , NINO4]
588 Table.
589 <http://iridl.ldeo.columbia.edu/SOURCES/.Indices/.nino/.EXTENDED/table/:table/.html>. accessed on: 1/ 2015
590
- 591 Kalnay E, Kanamitsu M, Kistler R, Collins W, Deaven D, Gandin L, Iredell M, Saha
592 S, White G, Woollen J, Zhu Y, Chelliah M, Ebisuzaki W, Higgins W,
593 Janowiak J, Mo KC, Ropelewski C, Wang J, Leetmaa A, Reynolds R, Jenne
594 R, Joseph D. 1996. The NCEP/NCAR 40-year reanalysis project. *Bulletin of*
595 *the American Meteorological Society* **77**(3):437-471. Doi 10.1175/1520-
596 0477(1996)077<0437:Tnyrp>2.0.Co;2
- 597 Kaplan A, Cane MA, Kushnir Y, Clement AC, Blumenthal MB, Rajagopalan B.
598 1998. Analyses of global sea surface temperature 1856-1991. *Journal of*
599 *Geophysical Research-Oceans* **103**(C9):18567-18589. DOI
600 10.1029/97jc01736
- 601 Karl TR, Knight RW. 1998. Secular trends of precipitation amount, frequency, and
602 intensity in the United States. *Bulletin of the American Meteorological Society*
603 **79**(2):231-241. DOI 10.1175/1520-0477(1998)079<0231:Stopaf>2.0.Co;2
- 604 Kingston DG, Lawler DM, McGregor GR. 2006. Linkages between atmospheric
605 circulation, climate and streamflow in the northern North Atlantic: research
606 prospects. *Progress in Physical Geography* **30**(2):143-174. DOI
607 10.1191/0309133306pp471ra
- 608 Laseter SH, Ford CR, Vose JM, Swift LW. 2012. Long-term temperature and
609 precipitation trends at the Coweeta Hydrologic Laboratory, Otto, North
610 Carolina, USA. *Hydrology Research* **43**(6):890-901. Doi
611 10.2166/Nh.2012.067
- 612 Leathers DJ, Yarnal B, Palecki MA. 1991. The Pacific North-American
613 Teleconnection Pattern and United-States Climate .1. Regional Temperature
614 and Precipitation Associations. *Journal of Climate* **4**(5):517-528. DOI
615 10.1175/1520-0442(1991)004<0517:Tpatpa>2.0.Co;2
- 616 Li WH, Li LF, Fu R, Deng Y, Wang H. 2011. Changes to the North Atlantic
617 Subtropical High and Its Role in the Intensification of Summer Rainfall
618 Variability in the Southeastern United States. *Journal of Climate* **24**(5):1499-
619 1506. DOI 10.1175/2010JCLI3829.1
- 620 Linacre E. 1992. Climate data and resources: a reference and guide: Psychology
621 Press.
- 622 Maraun D, Osborn TJ, Gillett NP. 2008. United Kingdom daily precipitation
623 intensity: improved early data, error estimates and an update from 2000 to

- 624 2006. *International Journal of Climatology* **28**(6):833-842. DOI
625 10.1002/joc.1672
- 626 Meehl GA, Stocker TF, Collins WD, Friedlingstein P, Gaye AT, Gregory JM, Kitoh
627 A, Knutti R, Murphy JM, Noda A, Raper SCB, Watterson IG, Weaver AJ,
628 Zhao Z-C. 2007. Global climate projections. In *Climate Change 2007: The*
629 *Physical Science Basis*, Contribution of Working Group I to the Fourth
630 Assessment Report of the Intergovernmental Panel on Climate Change.
631 Solomon S, Qin D, Manning M, Chen C, Marquis M, Averyt KB, Tignor M,
632 Miller HL (eds), Cambridge University Press: Cambridge, New York.
- 633 NOAA National Weather Service. 2015. Northern Hemisphere Teleconnection
634 Patterns. <http://www.cpc.ncep.noaa.gov/data/teledoc/telecontents.shtml>.
635 accessed on: 1/ 2015
- 636 O'Gorman PA, Schneider T. 2009. The physical basis for increases in precipitation
637 extremes in simulations of 21st-century climate change. *Proceedings of the*
638 *National Academy of Sciences of the United States of America* **106**(35):14773-
639 14777. DOI 10.1073/pnas.0907610106
- 640 Osborn TJ, Hulme M, Jones PD, Basnett TA. 2000. Observed trends in the daily
641 intensity of United Kingdom precipitation. *International Journal of*
642 *Climatology* **20**(4):347-364. DOI 10.1002/(Sici)1097-
643 0088(20000330)20:4<347::Aid-Joc475>3.0.Co;2-C
- 644 Prat OP, Barros AP. 2010. Ground observations to characterize the spatial gradients
645 and vertical structure of orographic precipitation - Experiments in the inner
646 region of the Great Smoky Mountains. *Journal of Hydrology* **391**(1-2):143-
647 158. DOI 10.1016/j.jhydrol.2010.07.013
- 648 Shepherd JM, Grundstein A, Mote TL. 2007. Quantifying the contribution of tropical
649 cyclones to extreme rainfall along the coastal southeastern United States.
650 *Geophysical Research Letters* **34**(23). DOI Artn L23810
651 10.1029/2007gl031694
- 652 Stahle DW, Cleaveland MK. 1992. Reconstruction and analysis of spring rainfall over
653 the southeastern US for the past 1000 years. *Bulletin of the American*
654 *Meteorological Society* **73**(12):1947-1961.
- 655 Swank WT, Crossley DA, editors. 1988. *Ecological Studies, Vol. 66: Forest*
656 *Hydrology and Ecology at Coweeta*. Springer-Verlag: New York.
- 657 Swift LW, Cunningham GB, Douglass JE. 1988. Climate and hydrology. In: Swank
658 WT, Crossley DA, editors. *Ecological Studies, Vol. 66: Forest Hydrology and*
659 *Ecology at Coweeta*. Springer-Verlag: New York.
- 660 UEA CRU. 2015. North Atlantic Oscillation (NAO).
661 <https://crudata.uea.ac.uk/cru/data/nao/>. accessed on: 1/ 2015
- 662 Unwin D. 1969. The areal extension of rainfall records: an alternative model. *Journal*
663 *of Hydrology* **7**(4):404-414.
- 664 Wang H, Fu R, Kumar A, Li WH. 2010. Intensification of Summer Rainfall
665 Variability in the Southeastern United States during Recent Decades. *Journal*
666 *of Hydrometeorology* **11**(4):1007-1018. DOI 10.1175/2010JHM1229.1
- 667 Wilson AM, Barros AP. 2014. An Investigation of Warm Rainfall Microphysics in
668 the Southern Appalachians: Orographic Enhancement via Low-Level Seeder-
669 Feeder Interactions. *Journal of the Atmospheric Sciences* **71**(5):1783-1805.
670 DOI 10.1175/Jas-D-13-0228.1
- 671 Yin ZY. 1994. Moisture Condition in the South-Eastern USA and Teleconnection
672 Patterns. *International Journal of Climatology* **14**(9):947-967. DOI
673 10.1002/joc.3370140902

674
675
676

677 **Table Heading and Figure Legends**

678 Table I. Location, elevation and first record of recording gauges included in this study
679 (Partly based on (Laseter *et al.*, 2012)).

680 Table II. Mean rainfall total, mean number of days above given threshold values,
681 maximum daily rainfall and mean rain per rain day by season at RRG06, 1937–2013.
682 Also shown is mean air temperature (°C). Note that some rows may not be exact totals
683 due to rounding errors.

684 Table III. Correlations for variables which changed significantly during the study
685 period at the RRG06 rain gauge. Correlations significant at $p < 0.05$ are shown in
686 plain font, $p < 0.01$ in bold, and $p < 0.001$ in bold and underlined ($n = 76$). Blank cells
687 indicate no significant correlation.

688 Table IV. Mean annual data for the four recording rain gauges, 1959–2013.

689 Table V. Trends in the lower (Q1) and upper (Q3) quartiles and the inter-quartile
690 range (IQR) at RRG06 for the number of days with totals over 25mm, by season and
691 year, 1937–2013. Correlations significant at $p < 0.05$ are shown in plain font, $p < 0.01$
692 in bold, and $p < 0.001$ in bold and underlined ($n = 57$). Blank cells indicate no
693 significant correlation. Moving window length is 20 years.

694 Table VI. Comparison between daily rainfall statistics for the ridge top (RRG31) and
695 valley floor (RRG06) for the period 1959–2012 inclusive.

696 Table VII. Frequency distribution of differences in daily rainfall totals between
697 RRG31 and RRG06 for the period Nov 1958 through Dec 2013. Only rain days
698 (≥ 0.25 mm) are included ($n = 6780$).

699 Table VIII. Double orographic enhancement (Burt and Howden, 2013) at Coweeta:
700 regression and correlation coefficients for relationships between the NAO and various
701 rainfall statistics. Results for RRG06 and RRG31 only, 1959–2013. Blank cells
702 indicate no significant correlation.

Table IX. Regression and correlation coefficients for relationships between the BHI
and various rainfall statistics at RRG06 and RRG31. * indicates no significant
correlation.

703 Figure 1. The location of the Coweeta Hydrologic Laboratory.

704 Figure 2. Increasing variability of annual rainfall totals at Coweeta (RRG06) over
705 time. Trends for 10th percentile, IQ1 and IQ3 are statistically significant. Trends for
706 annual total and 90th percentile (not shown) are not statistically significant.

707 Figure 3. 24-month running totals at RRG06 for (a) rainfall total, (b) wet days, and (c)
708 days with more than 25 mm.

Table I. Location, elevation and first record of recording gauges included in this study (Partly based on (Laseter *et al.*, 2012)).

Recording gauge (RRG)	Paired standard gauge (SRG)	Elevation (m)	Start of record	Aspect	T10 (mm)
06	19	685	1/1/1937	Valley bottom	74.9
96	96	894	1/1/1944	North-facing	79.5
40	13	961	1/1/1943	South-facing	78.5
31	31	1,366	1/1/1959	High-elevation gap	90.9

Table II. Mean rainfall total, mean number of days above given threshold values, maximum daily rainfall and mean rain per rain day by season at RRG06, 1937–2013. Also shown is mean air temperature (°C). Note that some rows may not be exact totals due to rounding errors.

	Winter	Spring	Summer	Autumn	Annual
Total (mm)	517	474	411	396	1799
Rain days	34	35	39	27	135
Days>17.8 mm (Q3)	10	9	7	7	34
Days>25 mm	7	6	4	5	22
Days>50 mm	2	2	1	2	6
Days>T10 per decade	6	4	3	6	19
Daily maximum (mm)	158.0	192.5	153.4	196.9	196.9
Rain/ rain day (mm/day)	15.0	13.6	10.5	14.6	13.3
Mean air temperature	4.2	12.7	21.3	13.3	12.9

Table III. Correlations for variables which changed significantly during the study period at the RRG06 rain gauge. Correlations significant at $p < 0.05$ are shown in plain font, $p < 0.01$ in bold, and $p < 0.001$ in bold and underlined ($n = 76$). Blank cells indicate no significant correlation. Note that significant correlations for the other three gauges are noted in the text but now shown in a table.

RRG06	Winter	Spring	Summer	Autumn	Annual
Total				0.258	
Wet days					
Days>17.8					
Days>25				0.267	
Days>50					
Days>T10				0.284	
Daily maximum (mm)				0.339	
Rain/ rain day (mm/day)				0.335	0.241
Mean air temperature (°C)		<u>0.427</u>	<u>0.447</u>	0.341	<u>0.459</u>

Table IV. Mean annual data for the four recording rain gauges, 1959–2013. Elevations for the stations increase from left to right (see Table I for specific station elevations).

	RRG06	RRG96	RRG40	RRG31
Total (mm)	1814	2010	1981	2394
Rain days	134	130	129	140
Days>25 mm	22	26	25	31
Days>50 mm	6	8	8	11
Days>T10	2	2	2	2
Daily maximum (mm)	196.9	205.7	198.9	232.4
Rain/ rain day (mm/day)	13.6	15.5	15.4	17.1

Table V. Trends in the lower (Q1) and upper (Q3) quartiles and the inter-quartile range (IQR) at RRG06 for the number of days with totals over 25mm, by season and year, 1937–2013. Correlations significant at $p < 0.05$ are shown in plain font, $p < 0.01$ in bold, and $p < 0.001$ in bold and underlined ($n = 57$). Blank cells indicate no significant correlation. Moving window length is 20 years.

	Q1	Q3	IQR
Winter	<u>-0.619</u>	<u>-0.697</u>	0.321
Spring	-0.32	<u>0.62</u>	<u>-0.435</u>
Summer		0.293	<u>0.484</u>
Autumn	<u>0.803</u>	<u>0.572</u>	<u>0.596</u>
Year	<u>-0.401</u>		<u>0.371</u>

Table VI. Comparison between daily rainfall statistics for the ridge top (RRG31) and valley floor (RRG06) for the period 1959–2012 inclusive.

	Winter	Spring	Summer	Autumn	Annual
Rainfall total ridge	647	621	540	581	2394
Rainfall total valley	506	480	402	422	1814
Gradient (mm/100m)	21	21	20	23	85
Rain days ridge	35	36	40	29	140
Rain days valley	33	35	37	28	134
Gradient (days/100m)	0.3	0.2	0.4	0.1	0.9
Rain per rain day ridge	18.2	17.3	13.4	20.2	17.1
Rain per rain day valley	15.2	13.9	10.4	15.3	13.6
Gradient					
(mm/day/100m)	0.4	0.5	0.4	0.7	0.5
Days \geq 25 mm ridge	9	8	6	8	31
Days \geq 25 mm valley	7	6	4	5	22
Gradient (days/100m)	0.3	0.4	0.3	0.4	1.3

Table VII. Frequency distribution of differences in daily rainfall totals between RRG31 and RRG06 for the period Nov 1958 through Dec 2013. Only rain days (≥ 0.25 mm) are included ($n = 6780$). The bins indicate the difference (mm) between the high-altitude gauge RRG31 and the valley-bottom gauge RRG06. Positive values show higher rain at altitude and *vice versa*.

bins	n	%
> 5	75	1.1%
4	72	1.1%
3	158	2.3%
2	432	6.4%
1	1102	16.3%
0	3597	53.1%
-1	1098	16.2%
-2	147	2.2%
< -2	99	1.5%

Table VIII. Double orographic enhancement (Burt and Howden, 2013) at Coweeta: regression and correlation coefficients for relationships between the NAO and various rainfall statistics. Results for RRG06 and RRG31 only, 1959–2013. Blank cells indicate no significant correlation.

	Total rainfall (mm)	Number of rain days	Daily totals ≥ 25 mm
(a) Regression coefficients			
RRG06 summer	61.0	2.08	1.02
RRG31 summer	76.3	3.04	1.42
RRG06 year	206.2	7.35	3.05
RRG31 year	258.8		4.81
(b) Correlation coefficients			
RRG06 summer	0.370	0.275	0.362
RRG31 summer	0.343	0.378	0.384
RRG06 year	0.365	0.289	0.350
RRG31 year	0.358		0.411

Table IX. Regression and correlation coefficients for relationships between the BHI and various rainfall statistics at RRG06 and RRG31.

	Total rainfall (mm)	Number of rain days	Daily totals ≥ 25 mm
(a) Regression coefficients			
RRG06 summer	82.1	2.47	1.34
RRG31 summer	122.1	2.87	1.84
RRG06 winter	51.1	2.48	0.75
RRG31 winter	62.8	0.19	0.92
(b) Correlation coefficients			
RRG06 summer	0.533	0.388	0.503
RRG31 summer	0.587	0.382	0.53
RRG06 winter	0.39	0.415	0.301
RRG31 winter	0.381	0.509	0.324




Figure 1. The location of the Coweeta Hydrologic Laboratory.




Figure 2. Increasing variability of annual rainfall totals at Coweeta (RRG06) over time. Trends for 10th percentile, IQ1 and IQ3 are statistically significant. Trends for annual total and 90th percentile (not shown) are not statistically significant.




Figure 3. 24-month running totals at RRG06 for (a) rainfall total, (b) rain days and (c) days with more than 25 mm.