

1 **Calreticulin is required for development of the cumulus oocyte**
2 **complex and female fertility**

3 Keizo Tokuhira¹, Yuhkoh Satouh¹, Kaori Nozawa^{1,2}, Ayako Isotani^{1,3,4}, Yoshitaka Fujihara¹,
4 Yumiko Hirashima⁵, Hiroyuki Matsumura^{1,4}, Kazuhiro Takumi^{1,4}, Takashi Miyano⁵, Masaru
5 Okabe¹, Adam M. Benham^{1,6}, and Masahito Ikawa^{1,2,3,4*}

6
7 ¹ Research Institute for Microbial Diseases, Osaka University, Suita, Osaka, Japan

8 ² Graduate School of medicine, Osaka University, Suita, Osaka, Japan

9 ³ Immunology Frontier Research Center, Osaka University, Suita, Osaka, Japan

10 ⁴ Graduate School of Pharmaceutical Sciences, Osaka University, Suita, Osaka, Japan

11 ⁵ Graduate School of Agricultural Science, Kobe University, Kobe, Hyogo, Japan

12 ⁶ School of Biological and Biomedical Sciences, Durham University, Durham, United Kingdom

13
14 * Corresponding author

15 Masahito Ikawa,

16 E-mail: ikawa@biken.osaka-u.ac.jp (MI)

17

18

1 **Abstract**

2 Calnexin (CANX) and calreticulin (CALR) chaperones mediate nascent glycoprotein folding in
3 the endoplasmic reticulum. Here we report that these chaperones have distinct roles in male and
4 female fertility. *Canx* null mice are growth retarded but fertile. *Calr* null mice die during
5 embryonic development, rendering indeterminate any effect on reproduction. Therefore, we
6 conditionally ablated *Calr* in male and female germ cells using *Stra8* (mKO) and *Zp3* (fKO)
7 promoter-driven Cre recombinase, respectively. *Calr* mKO male mice were fertile, but fKO
8 female mice were sterile despite normal mating behavior. Strikingly, we found that *Calr* fKO
9 female mice had impaired folliculogenesis and decreased ovulatory rates due to defective
10 proliferation of cuboidal granulosa cells. Oocyte-derived, TGF-beta family proteins play a major
11 role in follicular development and molecular analysis revealed that the normal processing of GDF9
12 and BMP15 was defective in *Calr* fKO oocytes. These findings highlight the importance of
13 CALR in female reproduction and demonstrate that compromised CALR function leads to ovarian
14 insufficiency and female infertility.

15

16

1 Introduction

2 Proper folding in the endoplasmic reticulum (ER) is a prerequisite for correct localization and
3 function of most secreted and transmembrane proteins ¹. Failures in protein folding and quality
4 control compromise cellular functions and cause disease including amyloidosis, cystic fibrosis,
5 and diabetes ^{2,3}. Failure in ER quality control also results in male infertility ^{4,6}. In the ER, soluble
6 calreticulin (CALR) and membrane bound calnexin (CANX) were originally discovered as
7 homologous calcium binding proteins and later shown to be lectin like chaperones that chiefly
8 mediate nascent glycoprotein folding ⁷⁻⁹. Despite their extensive homology, CALR and CANX
9 have contrasting functions. For example, proteins in *Calr*^{-/-} cells have accelerated folding with an
10 accumulation of misfolded proteins whereas folding is significantly impaired in *Canx*^{-/-} cells ¹⁰.
11 Differences in CALR and CANX function are further reflected in the distinct phenotypes of
12 knock-out mice. *Calr*^{-/-} mice are embryonic lethal due to defective heart development whereas
13 *Canx*^{-/-} mice are viable but growth retarded with neurological deficits ¹¹⁻¹³. However, the client
14 specificity of calnexin/calreticulin *in vivo* has not been fully established.

15
16 We have previously demonstrated that calmegin (CLGN) and calsperin (CALR3) are testis
17 specific homologues of CANX and CALR, respectively, and are expressed in the ER of
18 spermatogenic cells ^{4,5,14}. CLGN mediates the heterodimerization of ADAM1A/ADAM2 that is
19 required for the maturation of ADAM3, a sperm membrane protein. In contrast to this indirect
20 activity, CALR3 binds directly to ADAM3 and regulates its maturation. Although a pseudogene in
21 humans ¹⁵, mouse ADAM3 is essential for sperm migration from the uterus into the oviduct and
22 *Adam3* null male mice are sterile ^{16,17}. Both *Clgn* and *Calr3* null mice lack ADAM3 on their sperm
23 surface and null males are infertile. However, other membrane and secretory proteins are normally
24 present in these mutant spermatozoa, indicating the importance of CLGN/CALR3 rather than
25 CANX/CALR, for the maturation of ADAM3 related proteins ^{4,5}. The restricted client specificity
26 of CLGN and CALR3 is further highlighted by the fact that the N-glycosylated sperm membrane
27 fusion protein IZUMO1 is functionally presented on *Clgn/Calr3* mutant spermatozoa ¹⁸.

28
29 CLGN and CALR3 are not expressed in the ovary and thus female germ cells rely on the

1 CANX/CALR system for quality control of nascent proteins in the ER. During oogenesis, oocytes
2 secrete many factors that regulate the growth and differentiation of granulosa cells, including
3 GDF9 and BMP15. These two proteins are structurally complex with extensive disulfide-bonds
4 and N-linked glycans. Mutations that alter their ability to fold have been implicated as causes of
5 premature ovarian failure¹⁹⁻²². *Gdf9*^{-/-} and *Bmp15*^{-/-} females are sterile or subfertile, respectively
6^{23,24} and a recent paper has reported the physiological importance of the GDF9:BMP15 heterodimer
7²⁵.

8
9 In the present study, we have established mouse lines in which either *Canx* or *Calr* has been
10 ablated to investigate their roles in male and female germ cell development and function. Although
11 growth retardation was observed, both male and female *Canx* knockout mice were fertile. For
12 CALR, we generated male and female germ cell specific knockout mice to circumvent the
13 embryonic lethality. Whereas CALR was dispensable for spermatogenesis and sperm fertilizing
14 ability, it was required in the oocyte for the maturation of the TGF-beta family proteins, GDF9 and
15 BMP15, as well as the subsequent development of the cumulus oocyte complex (COC). Our
16 results highlight the importance of CALR for female reproduction and suggest that compromised
17 CALR function can lead to ovarian insufficiency and female sterility.

18

1 Results

2 *Generation of conditional knockout (KO) mice for Canx and Calr genes*

3 Mice with genetically disrupted *Canx* and *Calr* genes are growth retarded or die during embryonic
4 development, respectively¹¹⁻¹³. Thus, we generated male (mcKO) and female (fcko) germ
5 cell-specific conditional knockout mice to investigate the role of these genes in fertility. We first
6 generated mice carrying floxed alleles (fl/+) using gene targeting vectors that floxed exons 3 and 4
7 for *Canx* and exons 4-7 for *Calr* (**Supplementary Fig. S1A and Supplementary Fig. S2A,**
8 **related to Fig. 1**). The correct gene targeting event in embryonic stem cells and subsequent
9 germ-line transmission was confirmed by PCR analysis (**Supplementary Fig. S1B and**
10 **Supplementary Fig. S2B**). To remove the floxed exons in male and female germ cells, we used
11 transgenic mouse lines expressing Cre recombinase under the *Stra8* and *Zp3* promoters,
12 respectively^{26,27}. Homozygous knockout (-/-) mice were generated by crossing heterozygous
13 mutant (+/-) mice.

15 *Fertility in Canx knockout mice*

16 When we crossed *Canx*^{+/-} females with *Canx*^{+/-} males, *Canx*^{-/-} mice were born in expected
17 Mendelian ratios (+/+:+/-:-/=14:27:16, n=8 litters from 3 breeding pairs). However, 50% of the
18 homozygous null pups died within 48 hours and very few survived to three months, as reported
19 previously¹². Similar postnatal lethality was not observed in *Canx* gene trapped mice despite the
20 complete absence of CANX protein¹³. Paradoxically, truncated CANX protein was present in one
21 of the two lines in the earlier report with the more severe phenotype. Using antibodies against the
22 N- and C- terminal residues, CANX protein was not detected in our -/- mice (**Supplementary Fig.**
23 **S1D**). Therefore the variance in postnatal viability reported earlier may reflect the combined effect
24 of CANX deficiency and other factors, including genetic background and animal husbandry.

26 To examine *Canx*^{-/-} mouse fertility, adult (8 week-old) *Canx*^{-/-} females and males were mated with
27 wild-type (+/+) males and females, respectively. Although the testis size was smaller in *Canx*^{-/-}
28 male mice, the ratio of testis weight to body weight was comparable to that of wild-type ($0.37 \pm$
29 0.03% and $0.39 \pm 0.06\%$, respectively) and normal spermatogenesis was observed

1 **(Supplementary Fig. S3, related to Fig. 1).** *Canx*^{-/-} males were fertile and average litter sizes
2 were 7.1 ± 2.2 and 7.5 ± 0.8 (avg. \pm s.d. pups, n=6 litters from 3 males) in normal and null mice
3 respectively.

4
5 *Canx*^{-/-} females were also fertile, but litter sizes were smaller (3.3 ± 1.5 pups, n=4 litters from 4
6 females) possibly due to their smaller body size. We produced oocyte specific *Canx* fckO mice by
7 introducing the *ZP3-cre* transgene (*Canx*^{fl/-}; *Zp3-cre*). Mice were genotyped by PCR analysis and
8 the lack of CANX protein was confirmed by immunoblot (**Supplementary Fig. S1C, related to**
9 **Fig. 1**). *Canx* fckO female mice had normal fertility when mated with wild type males (8.4 ± 2.6
10 pups, n=9 litters from 3 females) (**Fig. 1A, B**). All 76 pups carried the knockout allele, which
11 confirmed successful excision of the floxed exons by the *Zp3* promoter driven Cre recombinase.
12 Therefore, we conclude that CANX is not required for either male or female reproduction.

14 ***Female infertility in Calr conditional knockout mice***

15 It was reported that the *Stra8* promoter driven Cre recombinase was expressed at the postnatal day
16 3 in early-stage spermatogonia and the recombination efficiency was >95%²⁶. In the present study,
17 with our *Stra8-cre* transgenic line, *Calr* disruption was confirmed in most (84.2%, 219/260) of the
18 testicular germ cells as determined by immunostaining (**Supplementary Fig. S4A, related to Fig.**
19 **1**). When mated with normal female mice, *Calr* mcKO males had comparable fertility (10.1 ± 1.5
20 pups, n=26 litters from 6 males) with normal male mice (10.6 ± 1.3 pups, n=8 litters from 3 males).
21 Whereas some pups inherited the floxed allele, the majority (82.1%, 215/262) of the offspring
22 inherited the knockout allele, which was consistent with the aforementioned observations in
23 testicular germ cells.

24
25 We next used *ZP3-cre* transgenic lines to disrupt the *Calr* gene during oogenesis (**Supplementary**
26 **Fig. S2C and Supplementary Fig. S4B, related to Fig. 1**). Whereas control female mice
27 (*Calr*^{fl/+}; *Zp3-cre*) had normal fertility, *Calr* fckO female mice had a profound decrease in
28 fecundity (**Fig. 1A, B**). Only 2 litters with 1 pup each were obtained from 16 copulations with 4
29 *Calr* fckO females, whereas 10 litters from 10 copulations were obtained with 4 control females.

1 The average litter sizes were 0.1 ± 0.3 and 8.4 ± 2.6 pups, respectively. When we superovulated
2 *Calr* fcKO females with gonadotropins, successful copulation was observed. Thus, the female
3 infertility was not caused by disrupted mating behavior. However, ovulation was severely
4 impaired in the *Calr* fcKO female mice (**Table 1**).

5
6 To elucidate the cause of infertility in the *Calr* fcKO females, we superovulated 3 (**Fig. 1C-F**) or
7 12 (**Supplementary Fig. S5C-H**) week-old female mice and prepared ovarian sections 2 hours
8 before the anticipated time of ovulation. In *Calr* fcKO mice, the ovary was smaller, possibly due to
9 defective folliculogenesis (**Table 2**). A few preovulatory follicles appeared, but most of the
10 follicles at the surface of the ovary were immature. Follicular development was arrested at the
11 early antral stage, the number of cumulus cells that surrounded an oocyte after germinal vesicle
12 breakdown in cross sections was reduced (33.1 ± 13.6 in fcKO and 141.6 ± 30.5 in control) and
13 cumulus expansion was impaired (**Fig. 1C-F and Supplementary Fig. S5A-H**). Corpora lutea
14 were rarely observed in *Calr* fcKO ovaries. Because antral follicles were present, we assayed *in*
15 *vitro* maturation and *in vitro* fertilization. Comparable numbers of oocytes were collected from
16 control and *Calr* fcKO mice 46-48 hours after PMSG injection. Both control and mutant oocytes
17 underwent germinal vesicle breakdown (GVBD) and matured to the metaphase II (MII) stage *in*
18 *vitro*. The oocytes collected from the *Calr* fcKO had slightly larger diameters than those collected
19 from control mice ($77.6 \pm 0.3 \mu\text{m}$ in fcKO and $73.5 \pm 0.4 \mu\text{m}$ in control, $P < 0.01$), as reported in
20 *Gdf9* KO mice²³. Although the efficiency was lower than in controls, these MII eggs could be
21 fertilized and developed to term after transfer into pseudopregnant females (**Table 3**).

22 23 ***CALR mediated quality control of GDF9 and BMP15***

24 *Calr* fcKO had impaired ovulation with defects in cumulus expansion. GDF9 and BMP15 are two
25 growth factors that are secreted from oocytes and stimulate cumulus cell expansion^{23,28}. To
26 examine the effect of CALR disruption on the production of these proteins, primary mouse
27 embryonic fibroblast (MEF) cells were prepared from *Calr*^{+/+} and *Calr*^{-/-} mice and transfected
28 with mouse *Gdf9* or *Bmp15* expression vectors. Using co-immunoprecipitation, we confirmed that
29 CALR associates with GDF9 and BMP15 in wild-type cells (**Fig. 2A**). In *Calr*^{+/+} MEFs, GDF9

1 and BMP15 were secreted into the extracellular fluid for 4-5 days, whereas in *Calr*^{-/-} MEFs,
2 secretion stopped within 1-2 days (**Fig. 2B**). When carefully observed, GDF9 proproteins
3 appeared as a doublet on SDS-PAGE²⁹, but the upper band was not present in *Calr*^{-/-} cell
4 supernatants (**Fig. 2C**). Recovery of the doublet was observed after co-transfection of a *Calr*
5 expression vector. Similar results were observed for BMP15. To examine whether GDF9 is
6 expressed and/or secreted in *Calr* fcKO oocytes, we collected 300 oocytes and examined their
7 lysates by immunoblot (**Fig. 2D**). Whereas comparable amounts of GDF9 proproteins (57kDa)
8 were observed in control and fcKO oocytes, mature type GDF9 (17kDa) was detected in control,
9 but not in CALR deficient oocytes. There were comparable amounts of other fertilization related
10 N-glycoproteins, including ZP3 and CD9.

11

12 ***Complementation of defective GOC (granulosa-oocyte complex) development by recombinant*** 13 ***GDF9 and BMP15***

14 External addition of recombinant BMP15 or GDF9 improves cumulus cell proliferation,
15 differentiation, and steroidogenesis in *Bmp15/Gdf9* null mice^{30,31}. In the present study, we
16 examined whether GDF9 and BMP15 supplements could restore the defective GOC development
17 in *Calr* fcKO mice using an *in vitro* follicle culture system. Without supplementation, granulosa
18 cells proliferated and the size of the follicles gradually increased when CALR was present. In *Calr*
19 fcKO GOC, the follicle size did not increase compared to controls (**Fig. 3**). When the GOC growth
20 medium was supplemented with either recombinant GDF9 or BMP15, both factors significantly
21 enhanced *Calr* fcKO GOC growth. Conditioned medium from cells expressing both GDF9 and
22 BMP15 did not show a synergistic effect.

23

1 Discussion

2 Although CANX is not essential for mouse development *in vivo*^{12,13}, it has been difficult to
3 investigate its role in the reproductive system since few *Canx* KO mice survived to adulthood and
4 they were smaller than their littermates. In the present study, we showed that *Canx* KO males were
5 able to copulate and successfully impregnate females. *Canx* KO females showed reduced litter size,
6 but oocyte specific disrupted cKO females were fully fertile. Therefore we conclude that CANX is
7 dispensable for both male and female reproductive systems.

8
9 We next examined the role of CALR, the soluble homologue of CANX, in the reproductive system.
10 *Calr* KO mice are embryonic lethal but ectopic expression of calcineurin in the heart enabled *Calr*
11 KO mice to survive until adulthood³². However, the surviving mice exhibited growth retardation,
12 hampering any study of their reproduction. Here, we generated male germ cell specific cKO mice
13 and showed that CALR is dispensable for spermatogenesis and sperm fertilizing ability. Although
14 CANX and CALR have contrasting functions in other tissues^{12,32}, their major roles may be
15 redundant and complementary to each other, at least in developing male germ cells. In contrast, the
16 homologues of CANX and CALR, CLGN and CALR3, respectively, have different substrate
17 specificity and both, albeit by different mechanisms, are required for fertilization by controlling
18 ADAM3 presentation on the sperm surface^{4,5}. Thus, the present study reinforces the uniqueness
19 and importance of the CLGN/CALR3 ER chaperone system in the male reproductive system.

20
21 In the female reproductive system, we have shown that CALR has a novel and indispensable role
22 in COC development and female fertility. Since GDF9 and BMP15 are secreted proteins, soluble
23 CALR might be more accessible to these molecules than membrane-tethered CANX. However it
24 is also reported that soluble CANX and CALR have different substrate specificity³³ and an
25 in-depth analysis of the molecular interactions of these chaperones with these two growth factors
26 requires future investigation. Our data indicated CALR plays an important role in regulating the
27 folding of GDF9 and BMP15 in the ER. The folding defects could cause various effects such as,
28 protein instability, abnormal cleavage or secretion defects, on post-translational modifications of
29 GDF9 and BMP15 at later stages of the secretory pathway.

1
2 In the present study we could not detect mature GDF9 and BMP15 proteins in MEF cells. The
3 defect in processing machinery may cause different behaviors in unfolded GDF9 and BMP15 in
4 MEF and oocytes. It has been reported that the mature recombinant mouse BMP15 is not
5 processed in HEK 293T cells and CHO cells ³⁴. Further, the post-translational processing of
6 BMP15 is precisely regulated during meiosis and the mature BMP15 can be detected 9 hours
7 after the hCG injection in mice ²⁸. The studies imply that oocytes may have a unique
8 post-translational processing mechanism for GDF9 and BMP15.

9
10 The follicular development in the *Gdf9* KO is arrested at the primary stage and the *Bmp15* KO
11 showed normal folliculogenesis despite decreased ovulation and rates of fertilization ^{23,24}. However,
12 follicular development in *Calr* fckO mice arrested at the early antral stage. In *Calr* fckO mice,
13 GDF9 proprotein is expressed in oocytes, but not cleaved. In rat, macaque and human, GDF9
14 proprotein or BMP15 is detected in the follicular fluid ^{35,36}. In addition, mutation in the protease
15 cleavage site of GDF9 causes female infertility in ewes ³⁷. The hypoplastic ovaries of homozygous
16 mutated ewe lambs contain large numbers of primordial follicles and developing follicles up to the
17 early antral stage. These data suggest that the proprotein of GDF9 accounts for the different
18 phenotypes in *Gdf9* KO and *Calr* fckO mice.

19
20 The mechanism of COC development has been well investigated and many key factors have been
21 identified in both oocyte and surrounding granulosa cells ^{38,39}. Among these factors, GDF9 and
22 BMP15 are secreted from developing oocytes and stimulate granulosa cell proliferation and
23 expansion ^{24,31,40}. GDF9 and BMP15 have a unique disulfide bond arrangement among TGF-beta
24 family proteins ⁴¹⁻⁴³, and showed different mobility in SDS electrophoresis under reducing or
25 non-reducing conditions ^{31,44}, indicating the importance of this disulfide bond formation during
26 their proper folding, targeting, and function. Here we showed that CALR interacts with GDF9 as
27 well as BMP15 and the lack of CALR compromised their secretion. This is reminiscent of the
28 misfolding and disappearance of ADAM3 in *Calr3* knockout male mice. In testis, CALR3 recruits
29 PDILT (protein disulfide isomerase like in testis) and assists in the quality control of ADAM3 ⁶.

1 From studies of other nascent glycoproteins folding in somatic cells ⁴⁵⁻⁴⁷, CALR may cooperate
2 with PDIA3 (ERP57) to regulate the quality control of GDF9/BMP15.

3
4 GDF9 and BMP15 play important roles in energy metabolism and/or cholesterol biosynthesis in
5 granulosa cells ^{25,48,49}. Because TGF-beta/Smad3 signaling regulates glucose and energy
6 homeostasis, and *Smad3* KO mice exhibit improved glucose tolerance and enhanced
7 glucose-stimulated insulin secretion from pancreatic islet beta cells ^{50,51}, it will be interesting to
8 investigate the role of CALR in the secretion of TGF-beta family proteins in other tissues. That
9 might also explain why *Calr* KO mice, transgenically rescued from embryonic lethality, still die
10 mainly due to metabolic failure ³². Of interest, it was also reported that TGF-beta stimulates cells
11 and induces extracellular matrix secretion that depends on CALR mediated Ca²⁺ signaling ⁵². Thus
12 the ability of CALR to regulate calcium availability might also be important downstream of the
13 TGF-beta signaling pathway. In the present study, we could not fully recover GOC development
14 with recombinant GDF9/BMP15 (**Fig. 3**), implying that there are other CALR dependent factors
15 required for complete oocyte development.

16
17 In conclusion, we document that CALR plays an indispensable role in the COC development by
18 controlling the maturation of the TGF-beta proteins GDF9 and BMP15. Because GDF9 and
19 BMP15 are known to be involved in premature ovarian failure and polycystic ovarian syndrome
20 ^{22,53-55}, targeting CALR and the ER quality control system in follicular development should be
21 considered as a novel avenue for female infertility treatment.

22

1 **Methods**

2 **Animal experimentation**

3 All animal experiments were carried out in accordance with the protocols approved by the Animal
4 Care and Use Committee of the Research Institute for Microbial Diseases, Osaka University. The
5 *Stra8*-cre mouse line was established in our laboratory by injecting a transgene that was
6 constructed by replacing the CAG promoter in pCX-cre vector with the *Stra8* promoter⁵⁶. The
7 *Zp3-cre* mouse line was obtained from the Jackson laboratory²⁷.

8 **Generation of KO and cKO mice**

9 Targeting vectors were constructed by placing *Canx* genomic fragments (1.7 kbp, 2.0 kbp and 4.0
10 kbp) and *Calr* genomic fragments (2.0 kbp, 1.4 kbp and 3.5 kbp) into pNT1.1⁵⁷. D3 embryonic
11 stem cells were electroporated with the NotI-linearized targeting vector. After G418/GANC
12 selection, drug resistant clones (3/96 for *Canx* and 5/672 for *Calr*) with homologous recombinants
13 (tm1a) were identified by PCR analysis. Three targeted clones were injected into C57BL/6
14 blastocysts, resulting in the birth of coat-color chimeric male mice. The neomycin-resistant
15 cassette was removed by crossing with transgenic mice expressing FLPe under the CAG promoter
16 and this allowed the generation of mice with the conditional floxed (fl or tm1b) allele (*Canx*^{fl/+} or
17 *Calr*^{fl/+}). These mice were crossed with transgenic mice expressing Cre under the CAG promoter
18 to generate *Canx*^{+/-} or *Calr*^{+/-} mice (- or tm1c). *Canx*^{-/-} or *Calr*^{-/-} mice were generated by mating
19 with each heterozygous mutated mouse that did not contain CAG-FLPe and CAG-Cre. To
20 generate cKO mice, *Canx*^{fl/+} or *Calr*^{fl/+} mice were crossed with transgenic mice expressing Cre
21 under the control of *Stra8* or *Zp3* promoters. For *Canx* fcKO, *Canx*^{fl/+}; *Zp3-cre* mice were crossed
22 with *Canx*^{fl/-} to produce mice with the following genotype: *Canx*^{fl/+}; *Zp3-cre* (for control) and
23 *Canx*^{fl/-}; *Zp3-cre* (for fcKO). *Calr* cKO were generated through the same breeding strategy as
24 *Canx* cKO using *Stra8-cre* or *Zp3-cre* transgenic mice. Pr-FlpeF; 5'-
25 ccacctaaggctcctggttcgtcagtttg -3' and pr-FlpeR; 5- atacaagtggatgatcctacccttgccg -3' primers
26 were used for the *Flpe* transgene in addition to the primers indicated in Figure S1 and Figure S2.

27 **Histology of the ovary**

1 Female mice were injected with pregnant mare serum gonadotropin (PMSG) and human chorionic
2 gonadotropin (hCG) at 48 hour intervals. Ovaries were collected at 10 hours after hCG injection
3 and fixed with 4% paraformaldehyde/PBS overnight. Then the ovaries were embedded into glycol
4 methacrylate (Technobit 8100: Heraeus Kulzer, Germany) after treatment with a graded ethanol
5 series. Plastic thin sections (5 μm) were stained with hematoxylin and eosin. For Periodic acid
6 Schiff (PAS) staining, sections were rehydrated, and treated with 2% metaperiodic acid for 15
7 minutes, followed by treatment with Schiff's reagent (Wako, Japan) for 20 minutes. The sections
8 were stained with hematoxylin prior to imaging.

9 10 **Immunohistochemistry**

11 Ovaries were collected from adult mice and fixed in 4% (wt/vol) paraformaldehyde/PBS overnight
12 at 4 $^{\circ}\text{C}$, cryopreserved in graded 10–30% (wt/vol) sucrose, and embedded in TissueTek OCT
13 compound (Sakura Finetechnical). Frozen sections (5 μm) were mounted on APS
14 (aminosilane)-coated glass slides. After washing with PBS, slides were blocked with 10%
15 (vol/vol) newborn calf serum (NBCS)/PBS for 1 hour and incubated with anti CALR antibody in
16 10% (vol/vol) NBCS/PBS at 4 $^{\circ}\text{C}$ overnight. After washing with PBS, the slides were incubated
17 with secondary antibodies with Alexa Fluor 488 (Invitrogen) in 10% (vol/vol) NBCS/PBS for 2
18 hours. After washing with PBS, the slides were imaged with an Olympus IX-70 fluorescence
19 microscope.

20 21 ***In vitro* maturation and fertilization**

22 For *in vitro* maturation⁵⁸, immature GV (germinal vesicle) oocytes were collected from ovaries 46
23 hours after injection with PMSG. Antral follicles were punctured with 26G needles in FHM⁵⁹ with
24 100 μM dibutyryl-cyclic AMP (Sigma). After dissociating the cumulus cells by pipetting, oocytes
25 were washed 3 times and cultured in Minimum Essential Medium Alpha (GIBCO) with 3mg BSA
26 (A3311, Sigma). After 14 hours, partial zona dissection of MII oocytes was performed using a
27 piezo-micromanipulator with a glass capillary needle (diameter of 5–10 μm)⁶⁰ and incubated in
28 TYH medium⁶¹ with $2 \times 10^5/\text{ml}$ B6D2F1 capacitated sperm. Fertilization was determined by
29 formation of two-cell embryos, which were then transferred into the oviduct of pseudopregnant

1 mice.

2

3 **Immunoblot and immunoprecipitation**

4 Oocytes were collected from ovaries 46 hours after PMSG injection. Cumulus or granulosa cells
5 were dissociated using glass pipettes and collected oocytes were boiled in SDS-PAGE sample
6 buffer. Proteins were separated by SDS-PAGE and run under reducing conditions.
7 Immunoprecipitation was performed as described ⁵. Antibodies used included rabbit antisera
8 against CAXN and CALR ⁵, rabbit antibody against the N-terminus of CAXN (sc-11397, Santa
9 Cruz Biotechnology), penta-His mouse monoclonal antibody (Qiagen), horseradish peroxidase
10 conjugated goat anti-rabbit IgG and goat anti-mouse IgG antibodies (Jackson Immuno Research
11 Laboratories); and a GDF9 monoclonal antibody ⁶², kindly provided by Dr. Martin M. Matzuk.

12

13 **Preparation of *Calr* deficient mouse embryonic fibroblasts (MEF)**

14 MEFs were isolated from 13.5-14.5 day-old embryos. The head and all internal organs were
15 removed from the embryo. The remaining tissue was minced with scissors and incubated in 0.25%
16 trypsin containing penicillin (100 U/ml, Nacalai) and streptomycin (100 µg/ml, Nacalai) for 10-20
17 min. The cells were pipetted and plated onto a 10 cm tissue culture dish in culture medium
18 [Dulbecco's Modified Eagle Medium (DMEM, GIBCO) with 10% FCS, penicillin (100 U/ml) and
19 streptomycin (100 µg/ml)]. The next day, the medium was replaced and cells were cultured at
20 37 °C until confluent, when the cells were harvested and frozen.

21

22 **Production of recombinant GDF9 and BMP15**

23 The HIV-1-based self-inactivating-type lentiviral vector plasmid pLV-EGFP was constructed by
24 replacing the EGFP cDNA with mouse GDF9 or BMP15 cDNA, respectively, and isolated for
25 infection of HEK293T cells ⁶³.

26

27 ***In vitro* follicle culture**

28 Ovaries were collected from 12 day-old mice from which follicles were mechanically isolated
29 prior to *in vitro* culture ⁶⁴. Each follicle was placed in a 30µl droplet of DMEM (GIBCO) with

1 10% FCS, penicillin (100 U/ml) and streptomycin (100 µg/ml). The droplets were placed in 60mm
2 uncoated culture dishes (Iwaki) and covered with paraffin oil (Nacalai). After overnight culture,
3 half of the culture medium was replaced by HEK293T cell-conditioned medium containing
4 recombinant GDF9 or BMP15. Culture medium from non-transfected HEK293T cells was used as
5 a negative control. Half of the medium was replaced every other day. Follicle volumes were
6 calculated as described ⁶⁵.

8 **Purification of His tagged GDF9 or BMP15 secreted into MEF cell media**

9 Mouse *Gdf9* or *Bmp15* cDNA tagged with Flag/Hisx6 at the C-terminus was inserted into a
10 pNCAG vector consisting of the CAG-promoter and rabbit β globin poly-adenylation signal,
11 respectively ⁵⁶. *Calr*^{+/+} or *Calr*^{-/-} MEF cells were cultured in 6-well plates and transfected with
12 pNCAG-Gdf9 Flag/His or pNCAG-Bmp15 Flag/His using lipofectamine LTX (Invitrogen). After
13 18 hours, cell debris was removed by two washes in PBS (GIBCO) and the medium was replaced
14 [DMEM with 3%FCS, penicillin (100 U/ml) and streptomycin (100 µg/ml)]. The culture medium
15 was collected every 24 hours for 4 days. Purification of His tagged recombinant protein in the
16 culture medium was performed using µMACS anti-His Isolation Kit (Miltenyi Biotec). Two
17 milliliters of culture medium were used for purification and the proteins were eluted into 300 µl.

19 **Statistical analysis**

20 The values were the means ± standard deviation or standard error of the mean from at least three
21 independent experiments. Statistical analyses were performed using Student's t-test. Differences
22 were considered significant at P<0.05.

23

References

- 1 Braakman, I. & Bulleid, N. J. Protein folding and modification in the mammalian endoplasmic reticulum. *Annu Rev Biochem* 80, 71-99 (2011).
- 2 Cohen, F. E. & Kelly, J. W. Therapeutic approaches to protein-misfolding diseases. *Nature* 426, 905-909 (2003).
- 3 Gregersen, N., Bross, P., Vang, S. & Christensen, J. H. Protein misfolding and human disease. *Annu Rev Genomics Hum Genet* 7, 103-124 (2006).
- 4 Ikawa, M. *et al.* The putative chaperone calmegin is required for sperm fertility. *Nature* 387, 607-611 (1997).
- 5 Ikawa, M. *et al.* Calsperin is a testis-specific chaperone required for sperm fertility. *J Biol Chem* 286, 5639-5646 (2011).
- 6 Tokuhira, K., Ikawa, M., Benham, A. M. & Okabe, M. Protein disulfide isomerase homolog PDILT is required for quality control of sperm membrane protein ADAM3 and male fertility [corrected]. *Proc Natl Acad Sci U S A* 109, 3850-3855 (2012).
- 7 Helenius, A. & Aebi, M. Roles of N-linked glycans in the endoplasmic reticulum. *Annu Rev Biochem* 73, 1019-1049 (2004).
- 8 Molinari, M. N-glycan structure dictates extension of protein folding or onset of disposal. *Nat Chem Biol* 3, 313-320 (2007).
- 9 Aebi, M., Bernasconi, R., Clerc, S. & Molinari, M. N-glycan structures: recognition and processing in the ER. *Trends Biochem Sci* 35, 74-82 (2010).
- 10 Molinari, M. *et al.* Contrasting functions of calreticulin and calnexin in glycoprotein folding and ER quality control. *Mol Cell* 13, 125-135 (2004).
- 11 Coppelino, M. G. *et al.* Calreticulin is essential for integrin-mediated calcium signalling and cell adhesion. *Nature* 386, 843-847 (1997).
- 12 Denzel, A. *et al.* Early postnatal death and motor disorders in mice congenitally deficient in calnexin expression. *Mol Cell Biol* 22, 7398-7404 (2002).
- 13 Kraus, A. *et al.* Calnexin deficiency leads to dysmyelination. *J Biol Chem* 285, 18928-18938 (2010).
- 14 Watanabe, D. *et al.* Molecular cloning of a novel Ca(2+)-binding protein (calmegin) specifically expressed during male meiotic germ cell development. *J Biol Chem* 269, 7744-7749 (1994).
- 15 Frayne, J. & Hall, L. The gene for the human tMDC I sperm surface protein is non-functional: implications for its proposed role in mammalian sperm-egg recognition. *The Biochemical journal* 334 (Pt 1), 171-176 (1998).
- 16 Shamsadin, R. *et al.* Male mice deficient for germ-cell cyritestin are infertile. *Biology of reproduction* 61, 1445-1451 (1999).
- 17 Yamaguchi, R. *et al.* Disruption of ADAM3 impairs the migration of sperm into oviduct in mouse. *Biology of reproduction* 81, 142-146 (2009).
- 18 Inoue, N., Ikawa, M. & Okabe, M. Putative sperm fusion protein IZUMO and the role of N-glycosylation. *Biochem Biophys Res Commun* 377, 910-914 (2008).
- 19 Zhao, H. *et al.* Analyses of GDF9 mutation in 100 Chinese women with premature ovarian failure. *Fertility and sterility* 88, 1474-1476 (2007).

- 1 20 Dixit, H. *et al.* Mutational screening of the coding region of growth differentiation factor
2 9 gene in Indian women with ovarian failure. *Menopause* 12, 749-754 (2005).
- 3 21 Di Pasquale, E., Beck-Peccoz, P. & Persani, L. Hypergonadotropic ovarian failure
4 associated with an inherited mutation of human bone morphogenetic protein-15 (BMP15)
5 gene. *Am J Hum Genet* 75, 106-111 (2004).
- 6 22 Otsuka, F., McTavish, K. J. & Shimasaki, S. Integral role of GDF-9 and BMP-15 in
7 ovarian function. *Mol Reprod Dev* 78, 9-21 (2011).
- 8 23 Dong, J. *et al.* Growth differentiation factor-9 is required during early ovarian
9 folliculogenesis. *Nature* 383, 531-535 (1996).
- 10 24 Yan, C. *et al.* Synergistic roles of bone morphogenetic protein 15 and growth
11 differentiation factor 9 in ovarian function. *Mol Endocrinol* 15, 854-866 (2001).
- 12 25 Peng, J. *et al.* Growth differentiation factor 9:bone morphogenetic protein 15
13 heterodimers are potent regulators of ovarian functions. *Proc Natl Acad Sci U S A* 110,
14 E776-785 (2013).
- 15 26 Sadate-Ngatchou, P. I., Payne, C. J., Dearth, A. T. & Braun, R. E. Cre recombinase
16 activity specific to postnatal, premeiotic male germ cells in transgenic mice. *Genesis* 46,
17 738-742 (2008).
- 18 27 de Vries, W. N. *et al.* Expression of Cre recombinase in mouse oocytes: a means to study
19 maternal effect genes. *Genesis* 26, 110-112 (2000).
- 20 28 Yoshino, O., McMahon, H. E., Sharma, S. & Shimasaki, S. A unique preovulatory
21 expression pattern plays a key role in the physiological functions of BMP-15 in the
22 mouse. *Proc Natl Acad Sci U S A* 103, 10678-10683 (2006).
- 23 29 Gilchrist, R. B. *et al.* Immunoneutralization of growth differentiation factor 9 reveals it
24 partially accounts for mouse oocyte mitogenic activity. *Biology of reproduction* 71,
25 732-739 (2004).
- 26 30 Sugiura, K. *et al.* Oocyte-derived BMP15 and FGFs cooperate to promote glycolysis in
27 cumulus cells. *Development* 134, 2593-2603 (2007).
- 28 31 Hayashi, M. *et al.* Recombinant growth differentiation factor-9 (GDF-9) enhances growth
29 and differentiation of cultured early ovarian follicles. *Endocrinology* 140, 1236-1244
30 (1999).
- 31 32 Guo, L. *et al.* Cardiac-specific expression of calcineurin reverses embryonic lethality in
32 calreticulin-deficient mouse. *J Biol Chem* 277, 50776-50779 (2002).
- 33 33 Wada, I., Imai, S., Kai, M., Sakane, F. & Kanoh, H. Chaperone function of calreticulin
34 when expressed in the endoplasmic reticulum as the membrane-anchored and soluble
35 forms. *J Biol Chem* 270, 20298-20304 (1995).
- 36 34 Hashimoto, O., Moore, R. K. & Shimasaki, S. Posttranslational processing of mouse and
37 human BMP-15: potential implication in the determination of ovulation quota. *Proc Natl*
38 *Acad Sci U S A* 102, 5426-5431 (2005).
- 39 35 Gode, F. *et al.* Influence of follicular fluid GDF9 and BMP15 on embryo quality.
40 *Fertility and sterility* 95, 2274-2278 (2011).
- 41 36 Duffy, D. M. Growth differentiation factor-9 is expressed by the primate follicle
42 throughout the periovulatory interval. *Biology of reproduction* 69, 725-732 (2003).

1 37 Souza, C. J., McNeilly, A. S., Benavides, M. V., Melo, E. O. & Moraes, J. C. Mutation in the protease cleavage site of GDF9 increases ovulation rate and litter size in heterozygous
2 the protease cleavage site of GDF9 increases ovulation rate and litter size in heterozygous
3 ewes and causes infertility in homozygous ewes. *Animal genetics* 45, 732-739(2014).

4 38 Juengel, J. L. & McNatty, K. P. The role of proteins of the transforming growth
5 factor-beta superfamily in the intraovarian regulation of follicular development. *Hum*
6 *Reprod Update* 11, 143-160 (2005).

7 39 Scaramuzzi, R. J. *et al.* Regulation of folliculogenesis and the determination of ovulation
8 rate in ruminants. *Reprod Fertil Dev* 23, 444-467 (2011).

9 40 Otsuka, F. *et al.* Bone morphogenetic protein-15. Identification of target cells and
10 biological functions. *J Biol Chem* 275, 39523-39528 (2000).

11 41 McPherron, A. C. & Lee, S. J. GDF-3 and GDF-9: two new members of the transforming
12 growth factor-beta superfamily containing a novel pattern of cysteines. *J Biol Chem* 268,
13 3444-3449 (1993).

14 42 Laitinen, M. *et al.* A novel growth differentiation factor-9 (GDF-9) related factor is
15 co-expressed with GDF-9 in mouse oocytes during folliculogenesis. *Mech Dev* 78,
16 135-140 (1998).

17 43 Dube, J. L. *et al.* The bone morphogenetic protein 15 gene is X-linked and expressed in
18 oocytes. *Mol Endocrinol* 12, 1809-1817 (1998).

19 44 Li, Q., Rajanahally, S., Edson, M. A. & Matzuk, M. M. Stable expression and
20 characterization of N-terminal tagged recombinant human bone morphogenetic protein
21 15. *Mol Hum Reprod* 15, 779-788 (2009).

22 45 Oliver, J. D., van der Wal, F. J., Bulleid, N. J. & High, S. Interaction of the
23 thiol-dependent reductase ERp57 with nascent glycoproteins. *Science* 275, 86-88 (1997).

24 46 Hughes, E. A. & Cresswell, P. The thiol oxidoreductase ERp57 is a component of the
25 MHC class I peptide-loading complex. *Curr Biol* 8, 709-712 (1998).

26 47 Morrice, N. A. & Powis, S. J. A role for the thiol-dependent reductase ERp57 in the
27 assembly of MHC class I molecules. *Curr Biol* 8, 713-716 (1998).

28 48 Sugiura, K., Pendola, F. L. & Eppig, J. J. Oocyte control of metabolic cooperativity
29 between oocytes and companion granulosa cells: energy metabolism. *Dev Biol* 279,
30 20-30 (2005).

31 49 Su, Y. Q. *et al.* Oocyte regulation of metabolic cooperativity between mouse cumulus
32 cells and oocytes: BMP15 and GDF9 control cholesterol biosynthesis in cumulus cells.
33 *Development* 135, 111-121 (2008).

34 50 Lin, H. M. *et al.* Transforming growth factor-beta/Smad3 signaling regulates insulin gene
35 transcription and pancreatic islet beta-cell function. *J Biol Chem* 284, 12246-12257
36 (2009).

37 51 Yadav, H. *et al.* Protection from obesity and diabetes by blockade of TGF-beta/Smad3
38 signaling. *Cell Metab* 14, 67-79 (2011).

39 52 Zimmerman, K. A., Graham, L. V., Pallero, M. A. & Murphy-Ullrich, J. E. Calreticulin
40 regulates transforming growth factor-beta-stimulated extracellular matrix production. *J*
41 *Biol Chem* 288, 14584-14598 (2013).

42 53 Simpson, J. L. Genetic and phenotypic heterogeneity in ovarian failure: overview of
43 selected candidate genes. *Ann N Y Acad Sci* 1135, 146-154 (2008).

1 54 Rossetti, R. *et al.* BMP15 mutations associated with primary ovarian insufficiency cause
2 a defective production of bioactive protein. *Hum Mutat* 30, 804-810 (2009).

3 55 Tiotiu, D. *et al.* Variants of the BMP15 gene in a cohort of patients with premature
4 ovarian failure. *Hum Reprod* 25, 1581-1587 (2010).

5 56 Matsumura, H., Hasuwa, H., Inoue, N., Ikawa, M. & Okabe, M. Lineage-specific cell
6 disruption in living mice by Cre-mediated expression of diphtheria toxin A chain.
7 *Biochem Biophys Res Commun* 321, 275-279 (2004).

8 57 Fujihara, Y. *et al.* Sperm equatorial segment protein 1, SPESP1, is required for fully
9 fertile sperm in mouse. *Journal of cell science* 123, 1531-1536 (2010).

10 58 Miki, H., Ogonuki, N., Inoue, K., Baba, T. & Ogura, A. Improvement of cumulus-free
11 oocyte maturation in vitro and its application to microinsemination with primary
12 spermatocytes in mice. *J Reprod Dev* 52, 239-248 (2006).

13 59 Lawitts, J. A. & Biggers, J. D. Culture of preimplantation embryos. *Methods in*
14 *enzymology* 225, 153-164 (1993).

15 60 Yamagata, K. *et al.* Sperm from the calmeglin-deficient mouse have normal abilities for
16 binding and fusion to the egg plasma membrane. *Dev Biol* 250, 348-357 (2002).

17 61 Toyoda, Y., Yokoyama, M. & Hoshi, T. Studies on the fertilization of mouse eggs in
18 vitro I: In vitro fertilization of eggs by fresh epididymal sperm. *Jpn J Anim Reprod* 16,
19 147-151 (1971).

20 62 Elvin, J. A., Clark, A. T., Wang, P., Wolfman, N. M. & Matzuk, M. M. Paracrine actions
21 of growth differentiation factor-9 in the mammalian ovary. *Mol Endocrinol* 13,
22 1035-1048 (1999).

23 63 Okada, Y. *et al.* Complementation of placental defects and embryonic lethality by
24 trophoblast-specific lentiviral gene transfer. *Nat Biotechnol* 25, 233-237 (2007).

25 64 Cortvrindt, R., Smitz, J. & Van Steirteghem, A. C. In-vitro maturation, fertilization and
26 embryo development of immature oocytes from early preantral follicles from prepuberal
27 mice in a simplified culture system. *Hum Reprod* 11, 2656-2666 (1996).

28 65 Janik, P., Briand, P. & Hartmann, N. R. The effect of estrone-progesterone treatment on
29 cell proliferation kinetics of hormone-dependent GR mouse mammary tumors. *Cancer*
30 *Res* 35, 3698-3704 (1975).

31

1 **Acknowledgements**

2 We thank Yumiko Maruyama and Akiko Kawai for generating mutant mice and Dr. Jurrien Dean
3 and Samantha A.M. Young (B. Sc. Hons) for review of the manuscript. This work was partly
4 supported by the Ministry of Education, Culture, Sports, Science, and Technology of Japan.

5

6

1 **Author Contributions**

2 K. Tokuhira, A.M.B., and M.I. designed research; K. Tokuhira, Y.S., K.N., A.I., Y.F., Y.H., H.M.,
3 K. Takumi, T.M., and M.I. performed research; K. Tokuhira, M.O., A.M.B., and M.I. analyzed
4 data; and K. Tokuhira, A.M.B., and M.I. wrote the paper.

5

1 **Additional Information**

2

3 **Competing financial interests**

4 The authors declare no competing financial interests.

1 **Figure legends**

2 **Figure 1. Abnormal ovarian follicular development in *Calr* fcKO mice.**

3 (A) Pregnancy rate (pregnancy/vaginal plug formation) obtained by mating control (*Calr*^{fl/+};
4 *Zp3-cre*), *Calr* fcKO (*Calr*^{fl/-}; *Zp3-cre*) and *Canx* fcKO (*Canx*^{fl/-}; *Zp3-cre*) females with B6D2F1
5 wild-type male mice. The total number of plugs observed is indicated in parentheses. (B) Average
6 litter sizes obtained by mating control, *Calr* fcKO and *Canx* fcKO females with B6D2F1 wild-type
7 male mice. (C-F) Histological analysis of ovarian sections. Preovulatory follicles from *Calr*
8 fcKO mice demonstrate immature follicles just under the surface of the ovary. The cumulus mass
9 surrounding oocytes is composed of fewer cells than control. Scale bar=300 μ m (C and E) and 50
10 μ m (D and F). Error bars represent standard deviation (B).

12 **Figure 2. CALR is essential for extracellular secretion of GDF9 and BMP15**

13 (A) *Calr*^{+/+} or *Calr*^{-/-} MEFs expressing Flag/His tagged GDF9 or BMP15 were
14 immunoprecipitated (IP) with anti-CALR antibody from the cell lysates (50 μ g) and the
15 immunoprecipitates were probed with anti-His antibody on immunoblots. (B) Secreted GDF9 and
16 BMP15 from *Calr*^{+/+} or *Calr*^{-/-} MEFs were examined every 24 hours after transfection. *Calr*^{-/-}
17 MEFs stopped secreting these proteins after 1-2 days. (C) Secreted GDF9 and BMP15 were both
18 observed as a single band in *Calr*^{-/-} MEFs. The upper band was recovered by co-transfection with
19 a plasmid expressing *Calr*. (D) The 57kDa GDF9 proprotein was present, but not the 17.5 kDa
20 mature GDF9 in CALR deficient oocytes. Other glycoproteins, CD9 and ZP3, were normally
21 expressed. 300 oocytes (GDF9) or 20 oocytes (ZP3 and CD9) were loaded per lane.

23 **Figure 3. Recombinant GDF9 and BMP15 partially restored follicular development of *Calr* 24 cKO**

25 (A) Control and *Calr* fcKO follicles were collected from 12 day-old mice and cultured *in vitro* for
26 9 days. In *Calr* fcKO mice, the addition of GDF9 or BMP15 partially promoted granulosa cell
27 proliferation and increased the follicular volume of *Calr* fcKO. (B) The average follicular
28 volumes from three independent experiments were determined. Error bars represent standard error
29 of the mean. Columns with the same letter are not significantly different ($P > 0.05$). (n=43, 39, 43,

1 38 and 12 for Control (*Calr*^{fl/+}; *Zp3-cre*), *Calr* fckO, *Calr* fckO+GDF9, *Calr* fckO+BMP15 and
2 *Calr* cKO + GDF9/BMP15, respectively).

3

1 Tables

2 **Table 1. Fertility of *Calr* fckO female mice after PMSG/hCG treatment**

Genotype	N ^a	Females ^b	Total eggs ^c	Normal eggs ^d	Fertilization rate ^e (%)
<i>Calr</i> ^{fl/+} ; <i>Zp3-cre</i>	4	5	28.2±4.1	22.8±6.8	70.2±19.7
<i>Calr</i> ^{fl/-} ; <i>Zp3-cre</i>	4	6	0.8±1.0	0.5±0.8	0

3 ^a number of replicates. ^b number of mice used. ^c number of ovulated eggs/animal. ^d number of MII
4 eggs/animal. ^e number of 2 cell embryos/number of inseminated eggs. Data are presented as
5 mean ± SD.

7 **Table 2. Ovarian characteristics after PMSG/hCG hormonal treatment**

Genotype	N ^a	Ovarian weight (mg)	Antral follicles ^b	Antral follicles with expanded cumulus ^c
<i>Calr</i> ^{fl/+} ; <i>Zp3-cre</i>	6	7.88 ±1.26	27.8 ±5.7	26.2 ±4.8
<i>Calr</i> ^{fl/-} ; <i>Zp3-cre</i>	6	4.18 ±1.36**	44.7 ±11.5**	1.2 ±1.5***

8 ^a number of mice used. ^b number of antral follicles/section. ^c number of antral follicles with
9 expanded cumulus/section. ***P* < 0.01, ****P* < 0.001 Data are presented as mean ± SD.

11 **Table 3. Development of *in vitro* matured and fertilized *Calr* fckO oocytes**

Genotype	N ^a	Females ^b	MII/GV ^c (%±SD)	2cell/MII ^d (%±SD)	Pups /transferred ^e
<i>Calr</i> ^{fl/+} ; <i>Zp3-cre</i>	7	11	320/585 (54.7±21.0)	139/248 (55.6 ±12.2)	ND
<i>Calr</i> ^{fl/-} ; <i>Zp3-cre</i>	8	16	575/961 (63.7±12.9)	147/471 (31.4±14.6)	8/64 [#] (12.5%)

12 ^a number of replicates. ^b number of mice used. ^c number of *in vitro* matured MII eggs/number of
13 GV oocytes. ^d number of 2 cell embryos/number of inseminated eggs. ^e number of pups/number
14 of transferred embryos. [#]Eight pups were obtained from two foster mothers.

15 Data are presented as mean ± SD.





