PROBING THE z > 6 UNIVERSE WITH THE FIRST HUBBLE FRONTIER FIELDS CLUSTER A2744*

HAKIM ATEK¹, JOHAN RICHARD², JEAN-PAUL KNEIB^{1,3}, BENJAMIN CLEMENT⁴, EIICHI EGAMI⁴, HARALD EBELING⁵,

MATHILDE JAUZAC⁶, ERIC JULLO³, NICOLAS LAPORTE^{7,8}, MARCEAU LIMOUSIN³, AND PRIYAMVADA NATARAJAN⁹

¹ Laboratoire d'Astrophysique, Ecole Polytechnique Fédérale de Lausanne, Observatoire de Sauverny, CH-1290 Versoix, Switzerland

² CRAL, Observatoire de Lyon, Université Lyon 1, 9 Avenue Ch. André, 69561 Saint Genis Laval Cedex, France

³ Aix Marseille Université, CNRS, LAM (Laboratoire d'Astrophysique de Marseille) UMR 7326, 13388, Marseille, France

⁴ Steward Observatory, University of Arizona, 933 North Cherry Avenue, Tucson, AZ, 85721, USA

⁵ Institute for Astronomy, University of Hawaii, 2680 Woodlawn Drive, Honolulu, Hawaii 96822, USA

⁶ Astrophysics and Cosmology Research Unit, School of Mathematical Sciences, University of KwaZulu-Natal, Durban, 4041 South Africa

⁷ Instituto de Astrofisica de Canarias (IAC), E-38200 La Laguna, Tenerife, Spain

⁸ Departamento de Astrofisica, Universidad de La Laguna (ULL), E-38205 La Laguna, Tenerife, Spain

Department of Astronomy, Yale University, 260 Whitney Avenue, New Haven, CT 06511, USA

Received 2013 November 29; accepted 2014 March 13; published 2014 April 16

ABSTRACT

The Hubble Frontier Fields program combines the capabilities of the *Hubble Space Telescope (HST)* with the gravitational lensing of massive galaxy clusters to probe the distant universe to an unprecedented depth. Here, we present the results of the first combined *HST* and *Spitzer* observations of the cluster A-2744. We combine the full near-infrared data with ancillary optical images to search for gravitationally lensed high-redshift ($z \ge 6$) galaxies. We report the detection of 15 I_{814} dropout candidates at $z \sim 6-7$ and one Y_{105} dropout at $z \sim 8$ in a total survey area of 1.43 arcmin² in the source plane. The predictions of our lens model also allow us to identify five multiply imaged systems lying at redshifts between $z \sim 6$ and $z \sim 8$. Thanks to constraints from the mass distribution in the cluster, we were able to estimate the effective survey volume corrected for completeness and magnification effects. This was in turn used to estimate the rest-frame ultraviolet luminosity function (LF) at $z \sim 6-8$. Our LF results are generally in agreement with the most recent blank field estimates, confirming the feasibility of surveys through lensing clusters. Although based on a shallower observations than what will be achieved in the final data set including the full Advanced Camera for Survey observations, the LF presented here goes down to $M_{\rm UV} \sim -18.5$, corresponding to $0.2L^{\star}$ at $z \sim 7$ with one identified object at $M_{\rm UV} \sim -15$ thanks to the highly magnified survey areas. This early study forecasts the power of using massive galaxy clusters as cosmic telescopes and its complementarity to blank fields.

Key words: galaxies: high-redshift – gravitational lensing: strong – surveys

Online-only material: color figures

1. INTRODUCTION

The discovery of the first galaxies to assemble in the universe has long been one of the most exciting challenges of observational cosmology. The identification of these high-redshift sources in deep blank fields relies mostly on the photometric detection of the intergalactic medium (IGM) absorption blueward Ly α , or the Lyman break technique (Steidel et al. 1996; Giavalisco et al. 2004). Great progress has been made in characterizing the early galaxy population at $z \sim 6-7$ through the determination of their UV colors, stellar masses, or ages (McLure et al. 2011; Bouwens et al. 2012; Richard et al. 2011) owing to the unprecedented capabilities of the Wide Field Camera 3 (WFC3) on board *Hubble Space Telescope (HST)*. Far from the several thousands of galaxies confirmed up to a redshift of 6.5, we are starting to spectroscopically confirm a few galaxies at z > 7 (Vanzella et al. 2011; Schenker et al. 2012; Ono et al. 2012; Finkelstein et al. 2013). The decrease of the prevalence of Ly α emitters at z > 7 might be a direct indication of the increase in the opacity of the IGM, which greatly attenuates the Ly α line through resonant scattering (e.g., Stark et al. 2010). Recent studies also suggest that the Ly α luminosity function (LF) decreases at z > 7, indicating an increase in the neutral gas fraction in the IGM (Kashikawa et al. 2006; Ouchi et al. 2010). However, such conclusions are still prone to large uncertainties because of the small sample size, as near-infrared (NIR) observation of faint sources at z > 7 is extremely challenging. Of course, one key driver for these studies of early galaxies is to determine the sources responsible for re-ionization of the high-redshift universe.

A complementary approach is to exploit the gravitational lensing offered by massive galaxy clusters (Kneib & Natarajan 2011), which magnifies the brightness of intrinsically faint sources. This has been successfully used to detect galaxies over a wide redshift range, taking advantage of the flux magnification (e.g., Kneib et al. 2004; Bouwens et al. 2009; Richard et al. 2011; Alavi et al. 2014) and the higher spatial resolution for detailed, small-scale studies of high-redshift galaxies (e.g., Brammer et al. 2012). In this framework, the CLASH¹⁰ program (Postman et al. 2012) yielded important results in a variety of fields using

^{*} Based on observations made with the NASA/ESA *Hubble Space Telescope* (*HST*), which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with programs 13495 and 11689. Based in part on observations made with the *Spitzer Space Telescope*, which is operated by the Jet Propulsion Laboratory, California Institute of Technology under a contract with NASA. This work utilizes gravitational lensing models produced by PIs Ebeling, Merten, and Zitrin, and Sharon funded as part of the *HST* Frontier Fields program conducted by STScI. STScI is operated by the Association of Universities for Research in Astronomy, Inc. under NASA contract NAS 5-26555. The lens models were obtained from the Mikulski Archive for Space Telescopes (MAST).

¹⁰ http://www-int.stsci.edu/~postman/CLASH/Home.html

THE ASTROPHYSICAL JOURNAL, 786:60 (9pp), 2014 May 1

 Table 1

 Summary of HST Observations

Instrument/Filter	No. Orbits ^a	5σ Depth ^b	Obs Date
WFC3/F160W	24	28.4	2013 Oct/Nov
WFC3/F140W	10	28.5	2013 Oct/Nov
WFC3/F125W	12	28.5	2013 Oct/Nov
WFC3/F105W	24	28.6	2013 Oct/Nov
ACS/F814W	5	27.4	2009 Oct
ACS/F606W	5	27.5	2009 Oct
ACS/F435W	6	27.6	2009 Oct
IRAC/3.6	90.9	0.139	2013 Sep
IRAC/4.5	90.9	0.225	2013 Sep

Notes.

^a Exposure time in ks for IRAC.

^b 3σ depth in μ Jy for IRAC data.

25 lensing clusters, resulting in significant progress in cluster mass modeling (e.g., Zitrin et al. 2012; Bradley et al. 2013; Coe et al. 2013; Vanzella et al. 2014). Combining the exquisite capabilities of *HST* with the power of "gravitational telescopes", the new *HST* Frontier Fields¹¹ (HFF) initiative aims to peer deeper into the distant universe by initially devoting a total of 560 orbits to observe four clusters down to a magnitude limit of ~29. Over Cycles 21 and 22, the program will obtain deep optical and NIR imaging in seven filters using the Advanced Camera for Survey (ACS) and WFC3 for both the clusters and the parallel blank fields.

Here we present the first results of the NIR observations of A2744, where we search for high-redshift candidates at $z \sim 6-8$ behind the galaxy cluster. In Section 2, we present the observational data set and reduction steps leading to the construction of the source catalog. The lensing model is described in Section 3. The procedure used to select high-redshift candidates is detailed in Section 4 together with the sources of contamination. In Section 5, we present the luminosity distribution at $z \sim 6-8$ and compare to previous blank field results. We use a standard Λ CDM cosmology with $H_0 = 71$ km s⁻¹ Mpc⁻¹, $\Omega_{\Lambda} = 0.73$, and $\Omega_m = 0.27$. Magnitudes are in the AB system.

2. OBSERVATIONS AND DATA REDUCTION

In this paper, we combine existing imaging and spectroscopic data obtained in previous *HST* campaigns targeting the A2477 cluster with new *HST* IR observations that were taken as part of the Frontier Fields program. The resulting pseudo-color image is shown in Figure 3.

2.1. Previous HST Data

Optical observations were obtained with the ACS on board *HST* in Cycle 17 (GO 11689, PI: Dupke) and used in the strong lensing analysis presented in Merten et al. (2011). Images were taken with the WFC using three broadband filters F435W, F606W, and F814W. A summary of the data set is presented in Table 1. For the basic reduction steps, we used the CALACS package v2012.2 that includes charge transfer efficiency corrections, which were not available in the publicly released reductions. Then, all exposures in each filter were median combined using the Astrodrizzle task in the new STSDAS/Drizzlepac package.¹² First, the different exposures were corrected for small

12 http://drizzlepac.stsci.edu

misalignments using Tweakreg before rejection of cosmic rays and correction for geometrical distortions. The final images have a pixel scale of 0''.05 pixel⁻¹ and reach a 5σ depth (for point sources in 0'.4 aperture) of 27.6, 27.5, and 27.4 in F435W, F606W, and F814W, respectively. Depths were determined by measuring the median standard deviation from 100 apertures of 0''.4 diameter positioned randomly on the sky.

2.2. Hubble Frontier Fields Data

The NIR observations of the first cluster in the HFF program (GO/DD 13495), using the WFC3, started on 2013 October 25. This includes imaging in four filters, F105W, F125W, F140W, and F160W, that achieves a total exposure time of 24, 12, 10, and 24 orbits, respectively. Here we use the full NIR observations summarized in Table 1. This is the first epoch observations of A2744 that will be completed with ACS observations during the second epoch scheduled for 2014 June. Basic reductions were once again performed using HSTCAL and the most recent calibration files. Individual frames were coadded using Astrodrizzle after registration to the ACS reference image using the Tweakreg shift file. After an iterative process, we achieve an alignment accuracy of 0.1 pixel between the WFC3 and ACS images. The final images have a 0.13 pixel size and the 5σ depth reached in the NIR filters is 28.6 (F105W), 28.5 (F125W), 28.5 (F140W), and 28.4 (F160W), respectively.

2.3. Spitzer Data

Spitzer imaging of A2744 using the Infrared Array Camera (IRAC) was obtained on 2013 September as part of the Frontier Field Spitzer program (PI: T. Soifer). The data set we used here represents 50% of the planned observations for the cluster. The total exposure time is 25 hr on sources in each of the 2 IRAC channels of 3.6 and 4.5 μ m. We used corrected Basic Calibrated Data (cBCD) images, which are provided by the Spitzer Science Center and automatically corrected by pipeline for various artifacts (such as muxbleed, muxstripe, and pulldown). The cBCD frames and associated mask and uncertainty images were processed, drizzled, and combined into final mosaics using the standard SSC reduction software MOPEX. The mosaic has a 2σ point source sensitivity (measured from the noise in a clean area) of 0.093 and 0.148 μ Jy in the 3.6 and 4.5 μ m channels, respectively.

2.4. Photometric Catalogs

We used a deep (F125W+F140W) image for source detection with SExtractor software (Bertin & Arnouts 1996). First, all images were point-spread function (PSF)-matched to the F160W image with a PSF model derived using Tiny Tim (Krist et al. 2011) and to a pixel scale of 0.'13. Then, we ran Sextractor in dual mode with the deep IR image as detection frame and the image in each filter to perform the photometry within the same aperture. The isophotal (AUTO) magnitude is adopted for the color (total flux) and a local background calculation is adopted for the photometry. The final drizzling was performed with the inverse variance map to account for all the sources of uncertainty in the image. Then this weight map was transformed to an rms map including the correlated noise correction (Casertano et al. 2000) to derive flux uncertainties during source extraction.

3. CLUSTER MASS MODEL

The strong lensing (SL) mass model will be presented in detail in a forthcoming publication (J. Richard et al. 2014, in



Figure 1. Survey area as a function of magnification in A2744. The different curves represent the cumulative area at redshift z = 7 probed for a given minimum magnification for different models. We use the mass models available on MAST (http://archive.stsci.edu/prepds/frontier/lensmodels/) that use a parametric approach (see Section 3 for details).

(A color version of this figure is available in the online journal.)

preparation). Here, we outline the main points of the methodology. We follow our previous lensing work (e.g., Richard et al. 2010; Limousin et al. 2012), adopting a parametric mass model combining both large-scale (cluster or group size) mass clumps and galaxy-scale mass clumps centered on each cluster galaxy (selected using a color-magnitude diagram), using scaling laws based on their magnitude in order to assign a mass to each galaxy (Limousin et al. 2007). The number of large-scale mass clumps is driven by the goal of reproducing as accurately as possible the location of the multiply imaged systems used as constraints.

Starting with an earlier model by Merten et al. (2011), we looked for multiply imaged systems within the ACS field of view. We finally identified 52 images coming from 17 multiply imaged background sources. Five of them (corresponding to two background sources) are located close to sub-clumps (group scale) located to the north and northwest limits of the ACS coverage (outside the WFC3 field of view). We targeted the core of this cluster with FORS2 on the ESO Very Large Telescope (VLT) and measured spectroscopic redshifts for two systems, allowing to reliably calibrate the mass model. The optimization is performed in the image plane, using the LENSTOOL software (Jullo et al. 2007). We found that a five component mass model is able to reproduce the observational constraints with an rms in the image plane equal to 0?. Three large-scale mass clumps are located in the core of the cluster and are associated with the brightest galaxies. The other two large-scale mass clumps are associated with the two north and northwest sub-clumps. On top of these large-scale mass clumps, perturbations associated with cluster members are considered. From this mass model, we generate the critical lines and the multiple-image region and compute the magnification map. Figure 1 shows the survey area as a function magnification using four lens models.

4. CANDIDATE SELECTION

For the selection of high-redshift candidates, we adopted the commonly used dropout criteria based on the Lyman break technique (Steidel et al. 1996; Giavalisco et al. 2004). For I_{814} dropouts we adopted the following criteria (see, for example, Oesch et al. 2010):

$$(I_{814} - Y_{105}) > 0.8$$

$$(I_{814} - Y_{105}) > 0.6 + 2(Y_{105} - J_{125})$$
 (1)

$$(Y_{105} - J_{125}) < 0.8.$$



Figure 2. Color–color selection of $z \sim 6-7$ (top panel) and $z \sim 8$ (bottom panel) candidates. The dropout candidates are represented by green circles with 1σ uncertainties and the selection window by the shaded region. We also show the expected color tracks as a function of redshift for starburst galaxies (solid lines) and elliptical galaxies (dotted lines). The color code indicates an increase of attenuation from $A_V = 0$ (in blue) to $A_V = 2$ (in red). We used standard galaxy templates from Coleman et al. (1980) and Kinney et al. (1996). The magenta points represent the colors of cool stars from the Chabrier et al. (2000) catalog.

(A color version of this figure is available in the online journal.)

In order to more efficiently reject low-*z* interlopers, the second criterion is more stringent than what has been used previously to identify similar sources (see Figure 2). We also require that the candidates be detected in the Y_{105} and J_{125} bands with a minimum of 5σ significance while they remain undetected in the optical B_{435} and V_{606} bands at less than 2σ . Objects that are not detected in the I_{814} filter are assigned a 3σ lower limit for their continuum break. To identify the Y_{105} dropouts, we applied the following color selection:

$$(Y_{105} - J_{125}) > 0.5$$

$$(Y_{105} - J_{125}) > 0.4 + 1.6(J_{125} - H_{140})$$
(2)

$$(J_{125} - H_{140}) < 0.5.$$

Similarly, we require a 5σ detection in the bands redward of the break and no detection in all the bands blueward of the break. The selection procedure is illustrated in Figure 2, where the top and bottom panels show the $z \sim 6-7$ and $z \sim 8$



Figure 3. Location of the drop-out images is superposed on the strong lensing model for A2744 (J. Richard et al., in preparation). The pseudo-color image is a combination of the F606W, F814W, and a deep (four-IR-band stack) WFC3/IR image. Overlaid in red are the critical lines for the background sources at z = 7. The positions of the candidates at z = 6-7 and $z \sim 8$ are marked with cyan and magenta circles, respectively. The green circles show the positions of the multiple-image systems identified with the predictions from the mass model, while the yellow line is the predicted region for multiple images.

(A color version of this figure is available in the online journal.)

selections, respectively. Objects satisfying our dropout selection (shaded box) are represented by green circles with 1σ uncertainties. We also plot the color tracks of different galaxy types as a function of redshift and for three values of attenuation. We used the standard libraries of Coleman et al. (1980) to calculate the tracks of elliptical galaxies (dotted lines) and Kinney et al.

(1996) for star-forming galaxies (solid lines). The blue, orange, and red colors represent an attenuation of $A_V = 0$, 1, and 2, respectively. All objects are visually inspected to remove spurious detections or contaminated photometry from close objects. Our final sample consists of 15 candidates at $z \sim 7$ and 1 candidate at $z \sim 8$. Figure 3 shows a color image of A2744 with the

positions of the candidates. Object 3284 is very uncertain due to its point-like morphology. Therefore, we decided to not include this candidate in the $z \sim 7$ LF determination of Section 5. The Y_{105} dropout shows a flux excess in the 4.5 μ m IRAC band while it is undetected in the 3.5 μ m channel. This is possibly due to the contribution of the redshifted $[O III] + H\beta$ emission lines. A detailed analysis of the properties of this candidate and a discussion of its high-redshift solution are presented in Laporte et al. (2014). Gravitational lensing also provides higher angular resolution allowing the study of spatially resolved star formation and the internal structure of distant galaxies (Frye et al. 2012; Brammer et al. 2012). The HFF observations will extend this study to the most magnified high-redshift galaxies. Candidate 561, for instance, shows an arc-like extended morphology. The source reconstruction of such objects will enable us to analyze the clumps and morphology of the building blocks of present day galaxies and compare them to other results at intermediate redshifts (Zitrin et al. 2011; Bradley et al. 2012). Because the optical ACS data do not match the depth of the new IR observations, we restricted the search to relatively bright candidates. Specifically, in the case of objects undetected in the I_{814} filter, we measured the 2 σ limit in the same aperture as in the IR. Then, we retained only those candidates that satisfy the selection criteria 1, replacing the I_{814} magnitude by its limit. For both I and Y dropouts, to ensure non-detections in the optical bands blueward of the break, the detection limit in the combined optical image was about one magnitude deeper than the object magnitude in the detection band. Consequently, the faintest object in our catalogue was about $J_{125} \sim 27.5$ mag. The identification of fainter objects will be possible with upcoming deep ACS observations that will be taken as part of the same HFF program.

4.1. Contamination

Given the stringent selection criteria we used here, we expect a very low contamination rate from spurious sources or lowredshift interlopers. As we have seen before, the IR observations are much deeper than the faintest candidate in our sample. This ensures that all candidates were detected in several filters with a very high significance, which basically excludes spurious detections.

Since the infrared data were taken much later than the ACS data, transient sources, such as supernovae that have exploded recently, can be detected only in the infrared, satisfying the continuum break criterion. We first compared the four epochs of IR observations spanning a period of 5 weeks and verified the consistency between the photometry in all the bands. Furthermore, such sources would have appeared as point sources. The inspection of the high-resolution images of the *HST* shows that the candidates are all spatially resolved.

Possible sources of contamination also include low-redshift galaxies that enter the selection space. This can occur when strong nebular emission lines contribute to the total flux in one filter, mimicking a continuum break (Atek et al. 2011). However, the use of multiple and contiguous filters greatly mitigates the contamination from such sources, since the emission line should be isolated and the flux enhancement restricted to only one filter (or two in the case of a combination of lines such as H α and [O III]). Low-*z* interlopers with unusually high reddening or an extremely old stellar population, although rare, could also be selected mistakenly as high-*z* dropouts (Hayes et al. 2012). The colors of most of our candidates suggest a rather blue or flat continuum incompatible with the existence of such a red population.

Finally, cool dwarf stars can have colors similar to highredshift galaxies. As we can see in Figure 2, the brown dwarf color track is more of a concern for $z \sim 8$ galaxy selection since it comes close to our candidate. However, we stress again that stars can be visually identified as point-like sources from their light profiles or the stellarity parameter of SExtractor (i.e., stellarity >0.6).

4.2. Photometric Redshifts

Additionally, we computed photometric redshifts for our candidates by fitting the photometric data with spectral energy distribution templates using the Hyperz software (Bolzonella et al. 2000). We used standard stellar population libraries of Bruzual & Charlot (2003) and a Salpeter (1955) initial mass function. We find that the probability distributions of the photometric redshifts all indicate a high-*z* solution in agreement with the color–color selection. The photometric and color information of the candidates are presented in Table 2. We also include in Table 2 the best-fit redshift given by the probability distribution function for each galaxy, together with the magnification factors. Figures 4 and 5 show the image cutouts of the *z* ~ 6–7 and 8 candidates in the ACS, WFC3, and IRAC filters.

4.3. Multiple Images

Strong lensing creates multiple images of the same background galaxy, whose locations can be predicted by our lens model (J. Richard et al., in preparation). For each object selected by our color-color selection, we visually search the field for additional counter-images according to their photometric redshift, color, and position. We also looked for multiply imaged systems that did not satisfy our selection criteria, mainly because of their low signal-to-noise ratio or contamination from cluster members. In Table 2, we keep only the brightest image among the multiples of the same object. The full catalog of multiple images is shown in Table 3. We identified two double-image, two tripleimage, and one quadruple-image system between $6 \leq z \leq 8$, which are also presented in Figure 3 with their respective identifying indices. The counter-images 1.2, 2.1, 5.2, and 5.4 were too severely affected by bright nearby objects and/or had too low of a signal-to-noise ratio to be included in our initial dropout selection. Their photometric redshift was, however, consistent with their dropout counterpart. In order to further assess the robustness of these multiply imaged candidates, we included each of them separately in the mass model as new constraints. We found that all candidates fit into the mass model without significant deviation, their individual rms being of the order of the total rms. Since parametric strong lensing mass modeling is very sensitive to redshift misidentification, we consider this test as a further confirmation of the robustness of these systems.

Some of these counter-images will likely be confirmed with the forthcoming ACS observations of A2744 which will provide deeper optical images allowing us to test their multiplicity, and thus corroborating the high-redshift nature of these candidates (e.g., Ellis et al. 2001; Kneib et al. 2004) as well as the robustness of our lens modeling procedure.

5. THE UV LUMINOSITY FUNCTION AT z = 7-8

We now turn to the estimate of the rest-frame UV LF at redshift $z \sim 6-7$ and 8 using the two galaxy samples assembled above. We first determine the absolute magnitude of each source in the J_{125} band at a mean redshift of $z \sim 6-7$ and in the H_{140}



Figure 4. Postage stamps of the $z \sim 6-7$ candidates in the ACS F435W, F606W, F814W; WFC3 F105W, F125W, F140W, F160W; and IRAC 3.6 μ m and 4.5 μ m bands. The size of each cutout is about 5" and the white circle denotes the source position. Sources show a strong $I_{814} - Y_{105}$ and remain undetected at a 2σ level in the B_{435} and V_{606} bands.

(A color version of this figure is available in the online journal.)



Figure 5. Same as Figure 4 for a $z \sim 8$ candidate. The source has a strong $Y_{105} - J_{125}$ break with no detection at a 2σ level in the B_{435} , V_{606} , and I_{814} bands. It is detected in all WFC3 filters redward of the break with a minimum of 5σ significance. (A color version of this figure is available in the online journal.)

Table 2Photometric and Color Measurements for the $z \sim 6-7$ Dropouts

Target	R.A. (J2000)	Decl. (J2000)	$I_{814} - Y_{105}$	$Y_{105} - J_{125}$	$J_{125}{}^{a}$	Magnification ^b	Photo-z
93	3.593807	-30.415442	>2.08	-0.03 ± 0.02	26.39 ± 0.01	3.42 ± 0.19	6.8
171	3.570648	-30.414662	1.89 ± 0.31	0.08 ± 0.02	26.49 ± 0.03	1.57 ± 0.03	6.3
410	3.585800	-30.411740	1.04 ± 0.15	-0.03 ± 0.02	27.06 ± 0.02	3.69 ± 0.18	5.8
561	3.603225	-30.410330	>1.28	-0.10 ± 0.03	27.04 ± 0.03	3.75 ± 0.19	7.5
632	3.593541	-30.409719	1.46 ± 0.16	-0.06 ± 0.02	26.57 ± 0.02	6.28 ± 0.56	5.9
841	3.606378	-30.407279	>1.60	-0.01 ± 0.03	26.73 ± 0.02	2.27 ± 0.07	6.4
1127	3.580442	-30.405039	>1.38	0.09 ± 0.03	26.73 ± 0.02	4.71 ± 0.36	6.4
1192	3.576128	-30.404494	1.23 ± 0.10	0.06 ± 0.01	26.46 ± 0.01	2.92 ± 0.14	6.1
1265	3.570062	-30.403720	>1.51	0.15 ± 0.02	27.01 ± 0.03	1.95 ± 0.05	7.0
1266	3.601099	-30.403956	>1.36	0.19 ± 0.03	26.94 ± 0.02	3.51 ± 0.22	6.5
1291	3.584396	-30.403395	1.12 ± 0.17	-0.05 ± 0.03	27.63 ± 0.03	72.75 ± 9.75	6.0
1521	3.600541	-30.401804	1.12 ± 0.22	-0.03 ± 0.04	27.50 ± 0.05	3.04 ± 0.16	5.9
2770	3.606230	-30.386646	1.88 ± 0.15	0.16 ± 0.02	26.34 ± 0.03	1.52 ± 0.03	5.5
3284	3.576655	-30.391364	2.59 ± 0.07	0.22 ± 0.00	24.05 ± 0.00	5.82 ± 0.55	6.5
3458	3.603540	-30.393106	0.91 ± 0.10	0.03 ± 0.02	26.68 ± 0.02	1.77 ± 0.05	6.0
Target	R.A. (J2000)	Decl. (J2000)	$Y_{105} - J_{125}$	$J_{125} - H_{140}$	H_{140}		
2070	3.604522	-30.380463	1.07 ± 0.03	0.30 ± 0.02	26.22 ± 0.01	1.47 ± 0.02	8.35

Notes.

^a Total magnitude.

^b This is the flux amplification factor.

Table 3Multiple-image Systems at $z \sim 6-8$

Image	Target	R.A. (J2000)	Decl. (J2000)	$I_{814} - Y_{105}$	$Y_{105} - J_{125}$	J ₁₂₅ ^a
1.1	1192	3.576128	-30.404494	1.23 ± 0.10	0.06 ± 0.01	26.46 ± 0.02
1.2	3822	3.591436	-30.396687	1.66 ± 0.22	-0.07 ± 0.03	27.34 ± 0.03
2.1	1291	3.584396	-30.403395	1.12 ± 0.17	-0.05 ± 0.03	27.83 ± 0.09
2.2	1291b	3.584741	-30.403147			>27.8
3.1	632	3.593541	-30.409719	1.46 ± 0.16	-0.06 ± 0.02	26.57 ± 0.02
3.2	1521	3.600541	-30.401804	1.12 ± 0.22	-0.03 ± 0.04	27.64 ± 0.03
4.1	410	3.585800	-30.411740	1.04 ± 0.15	-0.03 ± 0.02	27.15 ± 0.03
4.2	1180	3.600068	-30.404382	1.47 ± 0.26	0.23 ± 0.04	27.21 ± 0.03
4.3	673	3.596525	-30.409031	1.69 ± 0.43	0.33 ± 0.07	27.32 ± 0.04
5.1	1127	3.580442	-30.405039	1.89 ± 0.36	0.09 ± 0.03	26.75 ± 0.02
5.2	1956	3.585321	-30.397960	>0.36	0.09 ± 0.03	27.18 ± 0.02
5.3	3761	3.597850	-30.395970	>0.23	0.36 ± 0.04	26.74 ± 0.04
5.4	1327	3.587268	-30.403279	>0.18	$0.12\pm0.0.04$	27.48 ± 0.05

Note.^a Total magnitude.

filter at $z \sim 8$. The observed values need to be corrected for the gravitational lensing magnification.

5.1. Effective Volume

While amplifying the intrinsic flux of a given source, strong lensing also distorts and stretches the source plane volume where it lies. The resultant drawback is that a higher-magnification region will necessarily probe a smaller comoving volume. Therefore, we need to account for these two effects in our LF estimates. The LF is given by

$$\phi(M)dM = \frac{N_i}{V_{\text{eff}}(M_i)},\tag{3}$$

where N_i is the number of galaxies in the *i*th bin and V_{eff} is the associated effective survey volume.

Our effective volume is also determined by the shape of the redshift selection function and the incompleteness. We generated starburst templates from the Kinney et al. (1996) library, which were shifted to the desired redshifts 5 < z < 10,

then we applied attenuation of $A_V = 0-1$, and calculated synthetic fluxes using the HST filter throughputs. Then we created artificial galaxies using the ARTDATA package in IRAF, exploring the parameter space of observed magnitude, color, shape, and position in the image. We generated 10,000 galaxies randomly added to the actual optical and IR images, using 100 objects at a time. We applied a distribution of absolute magnitudes between -23 and -14 and applied the colors derived from the spectral templates. We used a log-normal distribution of half-light radii derived from the observed sizes of spectroscopically confirmed $z \sim 4$ LBGs of Vanzella et al. (2009). We applied a factor of $(1 + z)^{-1}$ to account for redshift-evolution in size (see Grazian et al. 2011). The sizes are computed in the source plane and the galaxy positions chosen randomly in the image plane. Then we used Lenstool to simulate the size and flux of the magnified galaxies in the image plane using our A2744 mass model. In the completeness calculation, we keep only the brightest image of each system.

We applied our source extraction and selection criteria 1 and 2 to the final images and compared to the input object catalogs.



Figure 6. Redshift selection function showing the relative efficiency in selecting I_{814} (blue curve) and Y_{105} (red) galaxies. This is determined using the simulations of color–color selection and completeness described in Section 5.1. (A color version of this figure is available in the online journal.)

Given our stringent selection criteria, the incompleteness function is dominated by the contamination by bright sources in the crowded field. We incorporate this incompleteness correction in the selection function that goes into the effective volume calculation. Figure 6 shows the resulting redshift selection function at $z \sim 6-7$ and 8.

The effective survey volume for each magnitude bin is calculated according to the following equation:

$$V_{\rm eff} = \int_0^\infty \int_{\Omega > \Omega_{\rm min}} \frac{dV_{\rm com}}{dz} f(z, m, \mu) d\Omega(\mu, z) dz, \qquad (4)$$

where Ω_{\min} is the source plane area with a minimum magnification μ_{\min} required to detect a galaxy with an apparent magnitude *m*. $f(z, m, \mu)$ is the completeness function including the redshift selection function, and $d\Omega(\mu)$ is the area element in the source plane as a function of the magnification factor.

Several mass models for the HFF clusters, including ours, are accessible through the STScI website. In order to estimate the uncertainties in the magnification maps and the differences between the models, we derived the LF using each model. Among the other five available models, only three offer the Kappa and Gamma maps that we integrate to derive the effective volume. These are the models of Sharon et al., Zitrin NFW (Navarro–Frenk–White), and Zitrin LTM (Light–Traces–Mass). In addition to the effective surface that shows variations from one model to another, we also computed the amplification factor for our candidates using the four models.

5.2. UV LF Results

The results are shown in Figure 7 where our luminosity distribution is compared with previous results at $z \sim 6-8$. At redshifts z = 6-7, we use two bins in magnitude (-18.5, -19.5) plus one object around $M_{\rm abs} \sim -15$, whereas we only have one candidate at $z \sim 8$. We stress here that object 1291 is close to the z = 7 critical line where the uncertainties become very high. Therefore, the amplification, and hence the intrinsic magnitude of this candidate, must be taken with caution.

The top panel of Figure 7 illustrates how the uncertainties in the different lensing models affect the LF determination. The four models yield a very similar LF, although at high amplification (see Table 3) the differences become significant (at



Figure 7. Rest-frame UV luminosity function at $z \sim 6-8$ based on our I_{814} and Y105 dropout samples. Top: UV LF determination using four different lensing models repeated by different symbols (see the legend). The blue symbols represent the LF determination at $z \sim 6-7$ compared to previous results in the blank fields at $z \sim 6$ (green line; Bouwens et al. 2006) and $z \sim 7$ (blue line; Schenker et al. 2012). The red symbols are the estimate at $z \sim 8$ compared to the HUDF12 determination at the same redshift (red line; Schenker et al. 2012). Error bars correspond to 1σ uncertainties. Bottom: UV LF averaged over the four models. The blue circles are the LF at $z \sim 6-7$ compared to the HUDF12 results at $z \sim 7$ (blue open squares) and $z \sim 8$ (red open squares).

(A color version of this figure is available in the online journal.)

 $M_{\rm abs} \sim -15$). The uncertainties on the UV LF determination also include the cosmic variance and Poisson errors. We estimated the 1σ fractional uncertainties of the galaxy number counts using the (Trenti & Stiavelli 2008) cosmic variance tool with the selection function and survey volume as inputs. Because of the small number statistics, the fractional Poisson errors $(1/\sqrt{N})$ are comparable to the uncertainties due to cosmic variance, which account for up to ~30%.

We show in the bottom panel of Figure 7 the averaged LF constructed from all the models. Overall, our derived LF is in good agreement with blank field results from the Hubble Ultra Deep Field (HUDF; Bouwens et al. 2006) and HUDF12 (Schenker et al. 2012). Most notably, we report here the faintest galaxy discovered at these redshifts, even though the uncertainties associated with its proximity to the critical line (where amplification is virtually infinite) are very large. The nature of this candidate will be better constrained with the advent of deep ACS data as part of the second epoch of the HFF observations. In the meantime, our results show that strong lensing will enable us to probe the distant universe at very faint magnitudes, comparable to intermediate redshift results, shedding light on the potential sources of cosmic reionization.

6. CONCLUSION

The first HFF result we report here is important in regards to unraveling the properties and distribution of high-redshift galaxies, since it confirms previous survey results derived from completely independent fields and therefore unequivocally demonstrates the feasibility of surveys using gravitational lensing fields. The agreement with previous blank field determinations of the LF also points to the robustness of our mass models, and hence our modeling procedure. Despite the current limited depth of the observations (soon to be rectified once the HFF/ACS observations are completed), our LF robustly reaches an intrinsic magnitude of $M_{abs} \sim -18.5$ at $z \sim 6-7$, which corresponds to about $0.2L_{z=7}^{*}$, and extends down to $M_{abs} \sim -15$ with a highly amplified object (a factor of 30–70). Albeit with large uncertainties, this result bolsters the advantage gained from gravitational magnification.

With the completed observations of the proposed six lensing clusters, the HFF program will probe the high-redshift universe to unprecedented depths, about two magnitudes deeper than typical blank field surveys, with the aid and enhancement provided by these "cosmic telescopes". We have shown here the feasibility and effectiveness of such studies and the robustness of our cluster mass models, which will gain even more accuracy as new candidates and ever more multiply imaged systems are discovered, identified, and confirmed. The LF of these highestredshift galaxies is a key signature and determinant of the sources responsible for re-ionization of the universe. Unmasking these sources with the help of the additional magnifying power offered by cluster-lenses looks not only promising but also feasible as presented here with the first HFF results.

H.A. and J.P.K. are supported by the European Research Council (ERC) advanced grant "Light on the Dark" (LIDA). P.N. acknowledges support from NSF theory grant AST-1044455 and a theory grant from the Space Telescope Science Institute HST-AR1214401.A. M.L. acknowledges support from CNRS. J.R. acknowledges support from the ERC starting grant CALENDS.

REFERENCES

Alavi, A., Siana, B., Richard, J., et al. 2014, ApJ, 780, 143 Atek, H., Siana, B., Scarlata, C., et al. 2011, ApJ, 743, 121

- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393
- Bolzonella, M., Miralles, J.-M., & Pelló, R. 2000, A&A, 363, 476
- Bouwens, R. J., Illingworth, G. D., Blakeslee, J. P., & Franx, M. 2006, ApJ, 653, 53
- Bouwens, R. J., Illingworth, G. D., Bradley, L. D., et al. 2009, ApJ, 690, 1764
- Bouwens, R. J., Illingworth, G. D., Oesch, P., et al. 2012, ApJ, 754, 83
- Bradley, L. D., Bouwens, R. J., Zitrin, A., et al. 2012, ApJ, 747, 3
- Bradley, L. D., Zitrin, A., Coe, D., et al. 2013, arXiv e-prints

Brammer, G. B., Sánchez-Janssen, R., Labbé, I., et al. 2012, ApJL, 758, L17

- Bruzual, G., & Charlot, S. 2003, MNRAS, 344, 1000
- Casertano, S., de Mello, D., Dickinson, M., et al. 2000, AJ, 120, 2747
- Chabrier, G., Baraffe, I., Allard, F., & Hauschildt, P. 2000, ApJ, 542, 464
- Coe, D., Zitrin, A., Carrasco, M., et al. 2013, ApJ, 762, 32
- Coleman, G. D., Wu, C.-C., & Weedman, D. W. 1980, ApJS, 43, 393
- Ellis, R., Santos, M. R., Kneib, J.-P., & Kuijken, K. 2001, ApJL, 560, L119
- Finkelstein, S. L., Papovich, C., Dickinson, M., et al. 2013, Natur, 502, 524
- Frye, B. L., Hurley, M., Bowen, D. V., et al. 2012, ApJ, 754, 17
- Giavalisco, M., Dickinson, M., Ferguson, H. C., et al. 2004, ApJL, 600, L103
- Grazian, A., Castellano, M., Koekemoer, A. M., et al. 2011, A&A, 532, A33
- Hayes, M., Laporte, N., Pelló, R., Schaerer, D., & Le Borgne, J.-F. 2012, MNRAS, 425, L19
- Jullo, E., Kneib, J.-P., Limousin, M., et al. 2007, NJPh, 9, 447
- Kashikawa, N., Shimasaku, K., Malkan, M. A., et al. 2006, ApJ, 648, 7
- Kinney, A. L., Calzetti, D., Bohlin, R. C., et al. 1996, ApJ, 467, 38
- Kneib, J.-P., Ellis, R. S., Santos, M. R., & Richard, J. 2004, ApJ, 607, 697
- Kneib, J.-P., & Natarajan, P. 2011, A&ARv, 19, 47
- Krist, J. E., Hook, R. N., & Stoehr, F. 2011, Proc. SPIE, 8127, 81270J
- Laporte, N., Streblyanska, A., Clement, B., et al. 2014, A&A, 562L, 8L
- Limousin, M., Ebeling, H., Richard, J., et al. 2012, A&A, 544, A71
- Limousin, M., Richard, J., Jullo, E., et al. 2007, ApJ, 668, 643
- McLure, R. J., Dunlop, J. S., de Ravel, L., et al. 2011, MNRAS, 418, 2074
- Merten, J., Coe, D., Dupke, R., et al. 2011, MNRAS, 417, 333
- Oesch, P. A., Bouwens, R. J., Illingworth, G. D., et al. 2010, ApJL, 709, L16
- Ono, Y., Ouchi, M., Mobasher, B., et al. 2012, ApJ, 744, 83
- Ouchi, M., Shimasaku, K., Furusawa, H., et al. 2010, ApJ, 723, 869
- Postman, M., Coe, D., Benítez, N., et al. 2012, ApJS, 199, 25
- Richard, J., Kneib, J.-P., Ebeling, H., et al. 2011, MNRAS, 414, L31
- Richard, J., Smith, G. P., Kneib, J.-P., et al. 2010, MNRAS, 404, 325 Salpeter, E. E. 1955, ApJ, 121, 161
- Schenker, M. A., Stark, D. P., Ellis, R. S., et al. 2012, ApJ, 744, 179
- Stark, D. P., Ellis, R. S., Chiu, K., Ouchi, M., & Bunker, A. 2010, MNRAS, 408, 1628
- Steidel, C. C., Giavalisco, M., Pettini, M., Dickinson, M., & Adelberger, K. L. 1996, ApJL, 462, L17
- Trenti, M., & Stiavelli, M. 2008, ApJ, 676, 767
- Vanzella, E., Fontana, A., Zitrin, A., et al. 2014, ApJL, 783, L12
- Vanzella, E., Giavalisco, M., Dickinson, M., et al. 2009, ApJ, 695, 1163
- Vanzella, E., Pentericci, L., Fontana, A., et al. 2011, ApJL, 730, L35
- Zitrin, A., Broadhurst, T., Coe, D., et al. 2011, MNRAS, 413, 1753
- Zitrin, A., Moustakas, J., Bradley, L., et al. 2012, ApJL, 747, L9