

Title: Acute effects of active gaming on ad-libitum energy intake and appetite sensations of 8-to-11 year-old boys.

Authors: Susan Allsop^{1*}, Caroline J Dodd-Reynolds², Benjamin P Green¹, Dorothee Debuse¹, Penny L S Rumbold¹.

Institutional affiliations:

¹Faculty of Health and Life Sciences, Department of Sport, Exercise and Rehabilitation Northumbria University, Northumberland Building, Newcastle upon Tyne, NE1 8ST, UK.

²School of Applied Social Sciences, Durham University, Durham, DH1 3HN, UK.

*Corresponding author: S Allsop, tel: +44(0)1912437018, fax: +44(0)191 227 3190, email s.allsop@northumbria.ac.uk.

Running title: Active gaming effects on intake and appetite.

Key words: Children: Active gaming: Energy intake: Physical activity: Appetite.

28 **Abstract**

29 The present study examined the acute effects of active gaming on energy intake (EI) and
30 appetite responses in 8-11 y boys in a **school-based setting**. Using a randomized crossover design,
31 21 boys completed four individual 90 min gaming bouts, each separated by 1 week. The gaming
32 bouts were 1) seated gaming no food or drink; 2) active gaming no food or drink; 3) seated gaming,
33 food and drink offered ad-libitum and 4) active gaming, food and drink offered ad-libitum. In the
34 two gaming bouts during which foods and drinks were offered, EI was measured. Appetite
35 sensations; hunger, prospective food consumption and fullness were recorded using visual analogue
36 scales (VAS) during all gaming bouts at 30 min intervals and at two 15 min intervals post-gaming.
37 In the two bouts with food and drink, no significant differences were found in acute EI (MJ)
38 ($p=0.238$). Significant differences were detected in appetite sensations for hunger, prospective food
39 consumption and fullness between the four gaming bouts at various time-points. The **relative energy**
40 **intake** calculated for the two gaming bouts with food and drink (active gaming 1.42 ± 0.28 MJ;
41 seated gaming 2.12 ± 0.25 MJ) was not statistically different. Acute EI in response to active gaming
42 was no different to seated gaming and appetite sensations were influenced by whether food was
43 made available during the 90 min gaming bouts.

Introduction

By the age of 10-11 y, one in three English children are now classified as being either overweight or obese ⁽¹⁾. In England, only 21% of boys and 16% of girls aged between 5 and 15 y achieve physical activity (PA) guidelines, and time spent being sedentary appears to be increasing ⁽²⁾. It is widely believed that this lack of PA has become a major contributor to children's positive energy balance ⁽³⁾.

Seated media activities, including television viewing, computer use and playing computer games are thought to reduce the time children spend undertaking sports and other physical activities ⁽⁴⁻⁸⁾. For children aged ≤ 11 years, associations have been found between sedentary activities such as television viewing and computer use and the spontaneous intake of unhealthy foods and drinks ⁽⁹⁻¹¹⁾. The more recently introduced active video games however require physical movement from the player. Linked to a television set via a console, active games need the player to physically interact with on-screen images through a tracking device within a camera (The Eye Toy, Sony Computer Entertainment[®]), a handheld controller (Nintendo Wii[™], Nintendo[®]) or a web-cam device (Xbox 360 Kinect, Microsoft). As such, these video games might present an appealing way to increase children's PA and thus offset any spontaneous energy intake (EI) ⁽¹²⁾. Recent laboratory-controlled investigations have indeed established that active video game play can increase children's energy expenditure (EE) threefold, in comparison to sedentary pursuits (watching television or playing seated video games) ⁽¹³⁻¹⁶⁾. Some active game play has been shown to elicit an intensity of 5 METS (moderate PA) with games such as "EyeToy Knockout" (PlayStation 2, Sony, Tokyo, Japan) ^(17, 18). Moreover, 15 min of Nintendo Wii Fit jogging, has been found to elicit an average of 5.35 METS in obese children ⁽¹⁹⁾. Such findings suggest that active video games have the potential to contribute significantly to levels of EE and help children meet the recommended moderate to vigorous physical activity (MVPA) levels.

To the authors' knowledge Mellecker and colleagues (2010) were the first to explore acute EI during active gaming in children ⁽²⁰⁾. During two, 1 hour laboratory gaming sessions (seated and activity enhanced), snacks were made available ad-libitum to 9-13 y children. No significant differences in snack consumption were found between the seated and activity enhanced video gaming conditions. During both trials Mellecker and Colleagues (2010) found that the mean EI was on average, 66% above resting levels ⁽²⁰⁾. This suggests that the additional PA elicited by the active gaming bout may not actually offset the EI in this group. No measures of appetite were explored however and EE was not estimated, failing to provide insight into any potential mechanisms for these changes in EI.

In relation to seated gaming, there have also been no differences reported in appetite when compared with resting conditions, in both male adolescents (aged 15–17 y)⁽²¹⁾ and boys (aged 9–14 y)⁽²²⁾. In the adolescent group, EE and ad libitum EI were significantly higher than after resting⁽²¹⁾, whilst the food intake of the younger boys (9–14 y) was lower after 30 minutes of seated video game-play versus the identical period of resting⁽²²⁾. Nonetheless, when EE was subtracted from EI, both groups were found to be in positive relative energy intake^(21, 22). The findings of paediatric gaming studies thus far however, suggest that active game play might be a healthier substitute for seated media activities.

To date, the effects of active gaming on appetite and EI have not been explored in 8–11 y boys. Furthermore, all of the previous studies have been strictly laboratory-controlled and the gaming protocols employed did not resemble the active game play of young children. We have recently published data describing the active gaming practices of 7–11 y children from Newcastle-upon-Tyne (North East England, UK)⁽²³⁾ to enable active gaming interventions to be designed that are representative of young people's habitual gaming practices.

The primary aims of the present study were therefore to explore acute EI and appetite sensations during active gaming and seated gaming, in 8–11 y boys. Secondary aims were to measure PA, estimate both EE and relative energy intake and establish time to eating onset.

Materials and methods

Design

A randomised, cross-over design was used to compare acute EI and appetite sensations of 8–11-year-old boys during four gaming bouts each separated by 1 week, utilising methods identified in a previous study⁽²³⁾. The four gaming bouts were: 1) 90 min seated gaming, no food or drink offered; 2) 90 min active gaming, no food or drink offered; 3) 90 min seated gaming with food and drink offered ad-libitum; 4) 90 min active gaming with food and drink offered ad-libitum. The boys were stratified according to school year into two groups of two so that a total of four boys were tested on each occasion. They were randomly assigned to a different gaming bout every week either with or without food. This meant that food and drinks were available to all four boys at the same time so that appetite sensations were not influenced by the sight of food and another boy eating. By the end of four weeks they had completed each of the four trials.

Ethical approval for the study was granted by the University of Northumbria, Faculty of Health and Life Sciences Ethics Committee. Written informed consent was obtained from both the parent (or main carer) and from the child, prior to data collection.

110 **Participants**

111 To recruit 8-11 y males, consent was obtained from head teachers of two primary schools
112 located within the city of Newcastle upon Tyne (North East England, UK of matching achievement
113 levels⁽²⁴⁾. The researchers distributed recruitment packs to all eligible boys who expressed an
114 interest in participating and they were asked to take this home. The pack contained a letter
115 addressed to their parent/main carer with a full explanation of the study and consent forms for them
116 and their child to sign and return to school. Signed consent was received for 22 boys and 21 of these
117 (mean age 9.8 ± 0.9 y) took part in and completed the study. The boy who did not complete the study
118 was excluded due to the parent being unable to standardize his EI. Boys were excluded from
119 participating when injury or illness prevented them from being able to play active video games or if
120 they had intolerances or allergies to the foods provided in the study.

121 **Preliminary measures**

122 Prior to the first gaming bout, the researchers visited the school to meet the children and
123 their parent or main carer. The purpose of the visit was to familiarise the boys (and where
124 applicable their parent or main carer) with the gaming consoles, the games, the gaming session
125 format, the self-reported weighed food diaries and visual analogue scales (VAS) used to measure
126 appetite sensations. A demonstration of the right hip placement of accelerometers (Actigraph LLC[®]
127 GT3X+) to enable measurement of PA during the gaming bouts was also provided. The boys were
128 asked to complete a food preference questionnaire during the familiarisation session to ensure the
129 foods and drinks offered during the study were not disliked by them. The boys were also
130 familiarised with the appetite and mood VAS and completed the Dutch Eating Behaviour
131 Questionnaire for children (DEBQ-C), as a measure of dietary restraint⁽²⁵⁾.

132 Anthropometric measurements were taken, with stature and seated height measured to the
133 nearest 0.01 m using a Harpenden Portable Stadiometer (Holtain Limited, Pembs, UK). Body
134 weight was measured to the nearest 0.1 kg using portable SECA scales (SECA United Kingdom).
135 Waist circumference was measured to the nearest 0.01 m with a non-elastic flexible tape at each
136 boy's natural waist whilst standing as a indicator of central adiposity⁽²⁶⁾.

137 **Protocol**

138 Each boy was provided with a self-report, weighed food diary and a set of food weighing
139 scales (Salter[®], Kent, UK) to use prior to all intervention days. With the help of their parent or
140 main carer they were asked to weigh and record all foods and drinks they consumed from 17:00 the

141 evening preceding, until after they had consumed breakfast on the morning of each intervention
142 day. A photocopy of this food diary was provided to each parent and they were asked to replicate
143 their child's food and drink intake prior to each gaming bout on three further intervention days. The
144 boys were also asked to abstain from all physical education at school on the day of the study and
145 PA from 17:00 the evening preceding.

146 The boys attended school as normal at 08:55. If any of the boys usually consumed a snack
147 during their morning break (10:40), they were provided with this in each intervention week by the
148 research team. The snack was dependant on the personal food intake of the boy and was the same
149 each week. At lunchtime (12:00) in the first week, each boy consumed a packed lunch prepared by
150 the research team which comprised their preferred food and drink items. The food and drink items
151 consumed were weighed and recorded by the research team so that an identical lunch could be
152 provided in each of the three subsequent visits.

153 Gaming sessions took place, at the end of the school day (15:15) on school premises as an
154 after-school club and commenced at 15:30 until 17:00. The gaming sessions were implemented for
155 90 min, as we found this to be the average time 7-11 y children spent playing active gaming
156 consoles⁽²³⁾. The boys were tested in sub-groups of four, with each of the four intervention arms
157 taking place on the same school day of each week, for four consecutive weeks.

158 *Gaming interventions*

159 The design of the individual gaming bouts was based on data we have published which
160 described the active gaming practices of 7-11 y children from Newcastle upon Tyne⁽²³⁾. Thus the
161 gaming console used for the active gaming bouts was Nintendo Wii™ and the game used was
162 Nintendo Wii™ Sports tennis. During each gaming bout, two boys played Nintendo Wii™ Sports
163 tennis and two the seated game. The seated game utilised was 'Mario and Sonic at the London 2012 Olympic
164 Games', played on Nintendo 3DS. The pair who were assigned to the Nintendo Wii™ played together
165 against the computer, whilst the two boys who played the Nintendo 3DS played individually against
166 the computer. In doing this, peer influence related to winning or losing⁽²⁷⁾ was avoided along with
167 any subsequent effects on EI.

168 The food and drink items provided during the gaming sessions were also based on the
169 previous findings⁽²³⁾ and were 130g apples (raw, slices and cored), 50g crisps [potato chips
170 (Walkers®, ready salted)], 250 mL semi-skimmed milk, 350 mL Robinson's apple and blackcurrant
171 squash (no added sugar). All food items were pre-weighed by the researchers to the nearest gram
172 using electronic portable scales (Salter®, Kent, UK) and all drinks were measured to the nearest

173 millilitre. The crisps and apple were placed in clear plastic bags and the milk and squash were
174 placed in coloured drinks bottles so that volumes were not identifiable. They were all numerically
175 coded by the researchers and placed at a station designated to each individual boy. All of the foods
176 and drinks were offered ad-libitum. The researchers noted each bag or bottle taken by the boys and
177 then weighed or measured anything left over so that amounts consumed could be calculated and
178 recorded. If further food and drink items were required during the gaming bouts, additional portions
179 were served.

180 To estimate EI from the food and drink items served, individual food labels, an online
181 resource (www.asda.com) and MicroDiet (Downlee Systems [©], Derbyshire, UK) were employed.
182 For each boy, exercise EE was subtracted from the amount of energy consumed during each gaming
183 bout to calculate relative energy intake. When the gaming bouts commenced, the time of the first
184 eating episode for each boy was recorded.

185 ***Appetite***

186 Hunger, fullness and prospective food consumption were assessed using paper-based VAS.
187 ‘How full do you feel now?’ anchored by very full (0) and not full at all (100), and prospective food
188 consumption ‘‘How much would you like to eat now?’’ anchored by a lot (0) and nothing at all
189 (100). The boys were requested to place a vertical mark along the 100 mm horizontal lines at set
190 times, prior to, during and following gaming cessation on all intervention days. Scales were
191 collected at baseline (0 min: 15:30), 30 min (16:00), 60 min (16:30), at the end (90 min: 17:00), 15
192 min post (17:15) and 30 min post gaming (17:30).

194 ***Physical activity assessment***

195 During every gaming bout, the PA levels of each boy were measured by accelerometry
196 using an Actigraph[©] LLC, GT3X+ worn on the right hip⁽²⁸⁾. The majority of accelerometer research
197 favours placement on the right hip as there is evidence to support this as being the optimum site^{(29,}
198 ³⁰⁾. Furthermore, when 11-17 y children played Nintendo WiiTM Sports tennis, right hip
199 accelerometer placement was found to have a closer relationship with EE than when positioned on
200 the right or left wrist⁽³¹⁾. The boys wore the accelerometer between 15:30 and 17:00 on each
201 gaming intervention day, with PA counts recorded at 10 s epochs. At the end of every gaming
202 session, the data was downloaded utilising Actilife 6 data analysis software and interpreted using
203 recommended child-appropriate activity cut-off values⁽³²⁾. Activity counts were integrated into 60 s
204 epochs utilising the child appropriate cut-offs of Evenson et al.,(2008) so that they could then be
205 converted into mean metabolic equivalents of time (METs) using the algorithm of⁽³³⁾ within the

206 Actilife version 6 software (ActiGraph Ltd, Pensacola, FL, USA). The following MET thresholds
207 recommended for use with children were used to categorise data based on PA intensity: sedentary <
208 1.5 METs; light 1.5 to < 4 METs; moderate 4 to < 6 METs; vigorous > 6 METS⁽³⁴⁾.

209 *Energy expenditure*

210 For each boy, Henry's body mass, stature and sex-specific equations were used to calculate
211 basal metabolic rate (BMR)⁽³⁵⁾. Energy expenditure was then calculated as recommended by Ridley,
212 Ainsworth and Olds (2008), as follows; METS x BMR (MJ·min⁻¹·d) x 90 min gaming = MJ⁽³⁶⁾. This
213 particular method of EE estimation accounts for age, sex, body mass (kg) and stature (m), unlike
214 other prediction equations which utilise only one or two of these physiological characteristics<sup>(34, 37-
215 39)</sup>.

217 **Statistical analysis**

218 Means and SEM are presented for all data. VAS ratings for subjective appetite sensations (hunger,
219 prospective food consumption and fullness) were calculated as time-averaged area under the curve
220 (AUC) for the gaming (15:30 – 17:00) and post gaming period (17:00 – 17:30) (120 min). PASW
221 Statistics (version 18.0, SPSS Inc., Chicago, Illinois) was used for all statistical analyses. One-way
222 repeated measures ANOVA were used to detect differences between mean PA, EE and baseline
223 appetite values. Two-way repeated measures ANOVA (trial × time) were used to detect differences
224 in appetite sensations and following a significant interaction effect, simple main effects analyses
225 were utilised. This approach enabled comparison between the four gaming bouts across all time
226 points. A Bonferroni correction was made when significant differences were identified. Energy
227 intake, % macronutrient intake [carbohydrate (CHO, fat and protein (PRO)], relative energy intake
228 and time to eating onset were analysed using dependent t-tests. For significant differences found in
229 the t-test analyses, Cohen's *d* effect size was calculated and interpreted against the effect size
230 categories of ≤0.20 = small effect, ~ 0.50 = moderate effect, and ≥0.80 = large effect⁽⁴⁰⁾. Statistical
231 significance was set at $p < 0.05$ for all analyses.

233 **Results**

234 **Population characteristics**

235 Preliminary measurements established mean (SD) stature 1.39 (SD 0.06) m; body mass 35.5
236 (SD 7.6) kg; waist circumference 64.3 (SD 8.4) cm and BMI 18.4 (SD 3.65) kg/m² of the boys.

237 According to UK age and gender-specific BMI centiles ⁽⁴¹⁾, the majority of the boys were classified
238 as having a healthy body mass (71.4%), 14.3% were classified as overweight and 14.3% as obese.
239 The mean maturity offset was -1.0 (SD 1.0) y from peak height velocity, indicating that the boys
240 were of similar maturation status. In addition, all boys were identified as being unrestrained eaters
241 according to the Dutch Eating Behaviour Questionnaire⁽²⁵⁾, with a mean (SD) dietary restraint score
242 of 1.7 (SD 0.5) categorized as being average for boys of this age (1.53 SD 1.95) ⁽²⁵⁾.

243 Appetite

244 All values for appetite are displayed in Table 1. There were no detectable differences in
245 mean baseline appetite sensations (hunger, prospective food consumption or fullness) between any
246 of the four gaming conditions as illustrated in Table 1. The time-averaged area under the curve
247 appetite values however, revealed significant differences between gaming conditions. Participants
248 felt more hungry during seated gaming without food compared with when they were both seated
249 ($p=0.006$) and active gaming with food ($p=0.009$). More specifically, they felt more hunger at 30
250 min, 60 min and 90 min during the above gaming conditions, as indicated in Figure 1. In relation to
251 prospective food consumption, the participants felt they wanted to eat more during seated gaming
252 without food, compared with during seated gaming with food ($p=0.002$) and active gaming with
253 food ($p=0.008$). They felt they wanted to eat more during the above gaming conditions at 60 min
254 and 90 min ($p=0.042$), as shown in Figure 2. They felt less full during seated gaming without food
255 compared with when they were seated gaming with food ($p=0.003$). They also felt less full when
256 active gaming without food compared with when they were both seated ($p=0.002$) and active
257 gaming with food ($p=0.014$). The boys felt less full during the above gaming bouts at 60 min, as
258 illustrated in Figure 3.

260 Physical activity METS and Energy Expenditure (MJ)

261 All values for PA (METS) and EE are displayed in Table 2. Active gaming elicited only
262 light PA and seated gaming was sedentary. The PA METS during active gaming with food were
263 significantly greater than seated gaming with food ($p<0.001$, effect size 0.6). Likewise the PA
264 METS during active gaming without food were significantly greater than seated gaming without
265 food ($p<0.001$, moderate effect size 0.7). As expected, no differences were found between active
266 gaming with or without food ($p=1.000$) and between seated gaming with or without food ($p=0.389$).

268 Energy intake (MJ), relative energy intake [EI – EE (MJ)] and time to eating onset (min)

All values for EI, relative energy intake and time to eating onset, are displayed in Table 2.

No significant differences were found in total EI (MJ) between the seated and active gaming bouts in which the foods and drinks were offered ($p=0.238$). Mean relative energy intake (MJ) was significantly greater after seated gaming in comparison to active gaming ($p=0.031$, effect size 0.3). The average time to eating onset (min) was significantly longer during active gaming with food, in comparison to seated gaming with food ($p=0.017$, effect size 1.0).

Discussion

The present study found no differences in the acute ad-libitum EI of 8-11 y boys during 90 min of active gaming when compared with seated gaming. Despite the lack of difference in EI, it took considerably longer for the first eating episode during active gaming to occur (17.10 min versus seated gaming 6.90 min). Sensations of hunger, fullness and prospective food consumption appeared to be influenced by whether food was made available during the gaming bouts. This was illustrated by the boys feeling more hungry, less full and wanted to eat more during the gaming conditions without food. Furthermore there were no differences in appetite between the seated and active gaming bouts with food.

To the best of our knowledge this is the first study to have investigated acute ad-libitum EI during active gaming using a genuine active game, in 8-11 y boys. The lack of difference in EI and macronutrient intake between active and seated gaming found in the present study is consistent with findings in adults when playing Nintendo Wii™ and Xbox 360⁽⁴²⁾. In the previously cited study, EI during active gaming when compared with seated gaming was also not significantly different. Over the one hour gaming periods, the adults consumed an average of 3.13 ± 2.26 MJ when seated versus 2.32 ± 2.08 MJ when active.

The only comparison that can be made with paediatric active gaming studies might be with the study by Mellecker and colleagues (2010), during which 9-13 y old children played a seated video game or an enhanced activity gaming device. In their investigation, the total energy consumed by the children during the seated gaming bout was equivalent to $1.57 \text{ MJ} \cdot \text{h}^{-1}$ and during active gaming was $1.60 \text{ MJ} \cdot \text{h}^{-1}$. When calculated per hour, the values of Mellecker and colleagues (2010) were similar to those found in the present study (1 h of seated gaming 1.92 MJ; 1h of active gaming 1.53 MJ). Although in the present study, EI during seated gaming was higher and during active gaming was lower than found by Mellecker and colleagues⁽²⁰⁾. The differences in EI findings between the previously cited study⁽²⁰⁾ and those determined presently, might be due to the former study being conducted in the unfamiliar setting of the laboratory and with the active gaming being a

302 seated gaming device played whilst walking on a treadmill. As such the active gaming format might
303 not have been as stimulating or challenging for the children as an actual active video game. Our
304 intention with the use of Nintendo Wii Sports™ tennis and the primary school settings was to
305 provide an intervention in an environment the participants were more accustomed to than the
306 laboratory.

307 Total EI during 90 min of seated gaming was calculated to be 2.88 MJ, and for active
308 gaming 2.30 MJ. When considering the daily estimated average requirement (EAR) for energy for
309 UK males aged 9 y (the average age of the present study population) is 7.70 MJ⁽⁴³⁾, the EI due to
310 seated and active gaming equated to 34% and 27% of daily EAR, respectively. Relative energy
311 intake which was estimated for seated and active gaming by subtracting the value for estimated EE
312 from EI which was 2.42±0.33 MJ and 1.64±0.27 MJ, respectively. As such EI during active gaming
313 with food was not offset by the greater estimated energy expenditure. Instead, a positive relative
314 energy intake was produced by both games which could contribute to a state of positive energy
315 balance. The consumption of food or drinks by children during active gaming therefore should not
316 be encouraged. Furthermore, the estimated EE from active gaming was equivalent to only light PA
317 and so in contrast to the findings of O'Donovan, Roche and Hussey (2014), would not contribute to
318 children's MVPA⁽¹⁹⁾.

319 Due to this present study being an acute investigation, the EI of the boys was not monitored
320 after the gaming bouts had ended (17:00) so it is not known whether compensation for the gaming
321 EI occurred later. The only paediatric study⁽⁴⁴⁾ thus far to have investigated compensation due to
322 active gaming EE, also did not establish any difference in EI in a post gaming meal, when
323 compared with 1 h of resting and seated gaming. At the end of the active gaming trial however, the
324 participants in the previously cited study, were established as being in negative energy balance,
325 which was then compensated for 24 h later by an increase in EI⁽⁴⁴⁾. However, the previous study
326 offered the food *ad-libitum* following the active gaming in a post-trial test meal and not during the
327 conditions, as in the present study and this might explain the difference in findings. In the current
328 study, it is possible that the boys compensated for the extra EI during both gaming trials at a later
329 time, either by a down-regulation in EI or an increase in EE. If no compensation did occur however
330 such substantial levels of energy surplus could contribute to a state of positive energy balance,
331 which could be clinically meaningful with regards to weight status. Particularly when a reduction of
332 only 0.46 to 0.69 MJ per day might be all that is required to reduce the energy gap to bring about a
333 decrease in body weight in children⁽⁴⁵⁾.

334 In 15-19 y adolescents, EI was monitored following seated gaming for the remainder of the
335 day but no compensation was found to have occurred for the extra food consumed ⁽²¹⁾. Although we
336 acknowledge that in relation to an increase in EE, it is possible for children to compensate and
337 increase EI 72 hours later to try and restore energy balance⁽⁴⁶⁾. As such, future research that
338 examines compensation post active gaming is warranted.

339 In relation to appetite, the boys felt significantly more hungry, less full and wanted to eat
340 more during the gaming conditions without food, in comparison to the gaming conditions with
341 food. However appetite sensations were no different between active and seated gaming with food.
342 Thus far, only two other paediatric studies have investigated the acute effects of active gaming on
343 subjective appetite ^(44, 47) and both of these reported similar findings to the present study. The
344 appetite sensations of healthy male adolescents were no different during 1 h of resting, seated
345 gaming and active gaming ⁽⁴⁴⁾ or in obese adolescent males during 1 h of resting, seated gaming,
346 active gaming and cycling ⁽⁴⁷⁾. As such, appetite and EI do not appear to be coupled due to active or
347 seated gaming as they are with exercise ⁽⁴⁸⁾.

348 These findings are also similar to those of seated gaming when compared with resting
349 conditions ^(21, 22). Both of the cited studies also observed no significant differences in appetite
350 sensations, whilst EI obtained from an ad-libitum post-gaming meal differed in the two
351 populations^(21, 22). In 11-13 y boys, EI was found to be significantly lowered by 0.25 MJ after 30
352 min of seated gaming⁽²²⁾, whilst 15 to 19 y males consumed similar amounts following seated
353 gaming and resting⁽²¹⁾. It should be noted however that in the more recent study of 11-13 y boys, a
354 glucose pre-load was administered at the start of the session which might have suppressed
355 subsequent food intake⁽²²⁾.

356 The lack of difference in appetite sensations and EI between seated and active gaming with
357 food observed in the present study population and during seated gaming in 15-19 y adolescents,
358 could be that both seated and active gaming might lead the boys to over-consume without an
359 increase in appetite sensations, as previously reported with television watching⁽⁴⁹⁾. As such, both
360 active and seated gaming might have the same distractive effect as television which appears to
361 cause fullness sensations to be ignored, resulting therefore in an over-compensation in EI for
362 gaming EE. Whether this over-compensation is due to the mental-stress-induced reward system⁽⁵⁰⁾
363 or an impairment in satiety signalling, has not yet been established ^(21, 51). Future active gaming
364 research with children might therefore consider the objective measurement of appetite hormones
365 alongside VAS due to the latter being a subjective measure.

366 To the best of our knowledge this is the first paediatric study to investigate the influences of
367 active gaming on acute appetite sensations and EI. The strengths of the study were, that it utilised
368 an intervention designed from actual survey findings that had established the active gaming
369 practices of 7-11 y children. In addition, the gaming bouts were implemented in a school-based
370 setting, as an after-school club thus creating a more relaxed and familiar environment for
371 participants, in contrast to that of a laboratory. Such a free-living and holistic design has evolved
372 from our latest paediatric exercise and appetite research.

373 A limitation of the study was that only Nintendo Wii™ Sports tennis was utilised during the
374 two, 90 min active gaming sessions. We felt however that it was important for the boys to play the
375 same game to enable accurate comparison of the individual gaming bouts. Furthermore, the
376 Nintendo Wii™ Sports tennis game utilises both upper and lower limbs during play, which should
377 allow for greater body movement and thus higher activity counts and EE ^(18, 52). The authors also
378 recognise that the prediction of EE from METS obtained by accelerometry is not without error,
379 particularly in children ^(36, 38). The accelerometers were placed on the right hip however, and there
380 is evidence to support this as being the optimum location ^(28, 29). Furthermore, in 11-17 y males, the
381 PA METS recorded from hip placement during 15 min play of Nintendo Wii™ Sport tennis, were
382 shown to have the closest relationship to EE, than other body sites tested, which were the right and
383 left wrists ⁽⁵²⁾. We considered it important however to implement the active gaming sessions in a
384 manner that was most true-to-life, outside of the laboratory with two boys playing against one
385 another. For this it was necessary to measure PA by accelerometry and thus estimate EE.
386 Subsequent estimations of relative energy intake therefore will also not be without error. Therefore,
387 we believe this study to have been exploratory in nature and we encourage researchers to use the
388 paediatric responses to active gaming presented in this manuscript to help power future studies.

389 To conclude, the availability of food had a significant effect on appetite sensations during
390 the gaming bouts. However there were no differences in the acute EI or appetite sensations of 8-11
391 y boys between 90 min active video gaming bout with food, when compared with seated gaming
392 bout with food. Acute EI due to eating and drinking during active gaming was calculated as 37%
393 (2.88 ± 0.32 MJ) and during seated gaming was 30% (2.30 ± 0.28 MJ) of daily EAR (based on 9 y old
394 boys). The relative energy intake estimated to have been produced from active gaming and acute
395 ad-libitum EI was 1.64 ± 0.27 MJ and for seated gaming was 2.42 ± 0.33 MJ, which might contribute
396 to positive energy balance. Appetite and EI responses to active and seated gaming require further
397 exploration, in order to establish whether the observed acute over-compensation in EI is offset by
398 subsequent EI and EE, after a gaming session has ended.

399 **Acknowledgements**

400 We thank all of the schools, the children and their parents who participated in this study. The
401 authors are grateful to Ashleigh Keenan, Ryan Cosgrove, and Mairead Fowler for assistance with
402 data collection. The project received no external funding. SA, PLS, DD and CJDR designed the
403 study, SA and BPG led and carried out the data collection and all authors contributed to data
404 analysis, interpretation and writing of the manuscript. The authors declare no conflicts of interest.

405

406

Figure 1






Figure 2



409

Figure 3



410

411 **Figure captions**

412 **Figure 1**

413 Mean_(SEM) hunger sensations for all participants (n = 21). *At 30 min, participants felt less hungry
414 during ○seated gaming with food compared with ●seated gaming without food ($p=0.010$). *At 60
415 min, they felt less hungry during Δactive gaming with food compared with ●seated gaming without
416 food ($p=0.025$). †At 60 min they felt significantly more hungry during ▲active gaming without
417 food in comparison to both ○seated ($p=0.049$) and Δactive gaming with food ($p=0.013$). *At 90
418 min, participants felt significantly less hungry during Δactive gaming with food compared with both
419 ●seated ($p=0.032$) and ▲active gaming without food ($p=0.029$).

420 **Figure 2**

421 Mean_(SEM) prospective food consumption for all participants (n = 21). *At 60 min participants
422 wanted to eat significantly more when ●seated gaming without food compared with Δactive gaming
423 with food ($p=0.030$). *At 90 min, participants wanted to eat significantly more when ●seated
424 gaming without food compared with ○seated gaming with food ($p=0.042$).

425

426 **Figure 3**

427 Mean_(SEM) fullness sensations for all participants (n = 21). *At 60 min, the participants felt
428 significantly less full when ●seated gaming without food than when ○seated gaming with food
429 ($p=0.011$) † Δand active gaming with food ($p=0.011$). They also felt more full when ‡ Δactive
430 gaming with food compared with ▲active gaming without food ($p=0.023$).

431

432 **Tables**

433 **Table 1**

434 **Table 1.** Mean and standard error values for all baseline and time-averaged area under the curve
435 (AUC) appetite sensations for all gaming bouts.

		Hunger (mm)		Prospective food consumption (mm)		Fullness (mm)	
		Mean	SEM	Mean	SEM	Mean	SEM
Baseline	Seated no food	43	8	46	8	55	9
	Active no food	52	7	55	7	60	7
	Seated with food	39	7	41	7	71	6
	Active with food	37	7	38	7	58	8
Time-averaged (AUC)	Seated no food	32	5	32	5	66	5
	Active no food	35	6	38	6	70	6
	Seated with food	47	4	55	6	50	4
	Active with food	50	4	50	5	52	5

436

Table 2

Table 2. Mean (SEM) PA METS, energy expenditure (EE) (MJ), energy intake (EI) (MJ), **relative energy intake** (MJ), time to eating onset (min) for all participants for each gaming bout.

	Seated no food		Active no food		Seated with food		Active with food	
	Mean	SEM	Mean	SEM	Mean	SEM	Mean	SEM
PA levels (METs)	1.38	0.07	2.14*	0.10	1.49	0.74	2.08†	0.90
EE (MJ)	0.43	0.02	0.69‡	0.03	0.51	0.03	0.66§	0.03
EI (MJ)					2.88	0.26	2.30	0.28
Relative energy intake (MJ)					2.42	0.25	1.64	0.28
Time of eating onset (min)					6.90	1.52	17.10¶	4.00

*PA (METs) were greater during active gaming without food than seated gaming without food ($p<0.001$).

†PA (METs) were greater during active gaming with food than seated gaming with food ($p<0.001$).

‡EE was significantly greater during active gaming without food than seated gaming without food ($p<0.001$).

§EE was significantly greater during active gaming with food than seated gaming with food ($p<0.001$).

||Relative energy intake was significantly lower when active gaming with food than when seated gaming with food ($p=0.031$).

¶Time of eating onset (min) was significantly longer during active gaming with food ($p=0.017$).

447 **References**

- 448 1. Boodhna G. Children's body mass index, overweight and obesity. London: The Health and
449 Social Care Information Centre 2013.
- 450 2. Scholes S, Mindell J. Health Survey for England: Volume 1: Chapter 3: Physical activity in
451 children. Health and Social Care Information Centre, 2012.
- 452 3. Almond L, Biddle S, Bull F, Cavill N, Ferguson R, Foster C, et al. Start Active, Stay Active.
453 A report on physical activity for health from the four home countries'. 2011.
- 454 4. Cummings HM, Vandewater EA. Relation of adolescent videogame play to time spent in
455 other activities Archives of Pediatric Adolescent Medicine. 2007;161(7):684-9.
- 456 5. Hands BP, Chivers PT, Parker HE, Beilin L, Kendall G, Larkin D. The associations between
457 physical activity, screen time and weight from 6 to 14 yrs: The Raine Study. Journal of Science and
458 Medicine in Sport. 2011;14(5):397-403.
- 459 6. Huang WY, Wong SH, Salmon J. Correlates of physical activity and screen-based behaviors
460 in Chinese children. Journal of Science and Medicine in Sport. 2013;16(6):509-14.
- 461 7. Vandewater EA, Shim M-s, Caplovitz AG. Linking obesity and activity level with children's
462 television and video game use. Journal of Adolescence. 2004;27(1):71-85.
- 463 8. Marshall SJ, Gorely T, Biddle SJH. A descriptive epidemiology of screen-based media use
464 in youth: A review and critique. Journal of Adolescence. 2006;29(3):333-49.
- 465 9. Pearson N, Biddle SJH. Sedentary Behavior and Dietary Intake in Children, Adolescents,
466 and Adults. A Systematic Review. American Journal of Preventive Medicine. 2011;41(2):178-88.
- 467 10. Moag-Stahlberg A, Miles A, Marcello M. What kids say they do and what parents think kids
468 are doing: The ADAF/Knowledge Networks 2003 Family Nutrition and Physical Activity Study.
469 Journal of the American Dietetic Association. 2003;103(11):1541-6.
- 470 11. Thivel D, Tremblay MS, Chaput J, P. Modern Sedentary Behaviors Favor Energy
471 Consumption in Children and Adolescents. Current obesity reports. 2012;2(1):50-7.
- 472 12. Mathieu M-E, Kakinami L. Active video games could be the solution to the increased
473 energy intake reported with sedentary video games. The American Journal of Clinical Nutrition.
474 2011;94(4):1150-1.
- 475 13. Graf DL, Pratt LV, Hester CN, Short KR. Playing Active Video Games Increases Energy
476 Expenditure in Children. Pediatrics. 2009;124(2):534-41.
- 477 14. Lanningham-Foster L, Foster RC, McCrady SK, Jensen TB, Mitre N, Levine JA. Activity
478 promoting games and increased energy expenditure. Journal of Pediatrics. 2009;154(6):819-23.

- 479 15. Straker L, Abbott R. Effect of Screen-Based Media on Energy Expenditure and Heart Rate in
480 9-to-12-Year-Old Children. *Pediatric Exercise Science*. 2007;19(4):459-71.
- 481 16. Lanningham-Foster L, Jensen TB, Foster RC, Redmond AB, Walker BA, Heinz D, et al.
482 Energy Expenditure of Sedentary Screen Time Compared With Active Screen Time for Children.
483 *Pediatrics*. 2006;118(6):E1831-E5.
- 484 17. White K, Schofield G, Kilding AE. Energy expended by boys playing active video games.
485 *Journal of Science and Medicine in Sport*. 2011;14(2):130-4.
- 486 18. Maddison R, Mhurchu CN, Jull A, Yannan J, Prapavessis H, Rodgers A. Energy Expended
487 Playing Video Console Games: An Opportunity to Increase Children's Physical Activity? *Pediatric*
488 *Exercise Science*. 2007;19(3):334-43.
- 489 19. O'Donovan C, Roche EF, Hussey J. The energy cost of playing active video games in
490 children with obesity and children of a healthy weight. *Pediatric Obesity*. 2014;9(4):310-7.
- 491 20. Mellecker RR, Lanningham-Foster L, Levine JA, McManus AM. Energy intake during
492 activity enhanced video game play. *Appetite*. 2010;55(2):343-7.
- 493 21. Chaput J, P, Visby T, Nyby S, Klingenberg L, Gregersen NT, Tremblay A, et al. Video
494 game playing increases food intake in adolescents: a randomized crossover study. *Am J Clin Nutr*.
495 2011;93(6):1196-203.
- 496 22. Branton A, Akhavan T, Gladanac B, Pollard D, Welch J, Rossiter M, et al. Pre-meal video
497 game playing and a glucose preload suppress food intake in normal weight boys. *Appetite*.
498 2014;83:256-62.
- 499 23. Allsop S, Rumbold PLS, Debus D, Dodd-Reynolds CJ. Real life active gaming practices of
500 7-11-year-old children. *Games for Health Journal: Research, Development and Clinical*
501 *Applications*. 2013;2(6):347-53.
- 502 24. Ofsted Inspection Reports [Internet]. Crown Copyright. 2014 [cited 23rd December 2014].
503 Available from: [http://reports.ofsted.gov.uk/inspection-reports/find-inspection-](http://reports.ofsted.gov.uk/inspection-reports/find-inspection-report/results/1/any/any/any/any/chillingham%20road/any/any/any/any/0/0)
504 [report/results/1/any/any/any/any/chillingham%20road/any/any/any/any/0/0](http://reports.ofsted.gov.uk/inspection-reports/find-inspection-report/results/1/any/any/any/any/chillingham%20road/any/any/any/any/0/0).
- 505 25. van Strien T, Oosterveld P. The Children's DEBQ for Assessment of Restrained, Emotional,
506 and External Eating in 7- to 12-Year-Old Children. *International Journal Eating Disorders*.
507 2008;41(1):72-81.
- 508 26. McCarthy HD, Jarrett KV, Crawley H, F. The development of waist circumference
509 percentiles in British children aged 5.0 - 16.9 y. *European Journal of Clinical Nutrition*.
510 2001;55(10):902-7.

- 511 27. Smith AL. Peer relationships in physical activity contexts: a road less traveled in youth sport
512 and exercise psychology research. *Psychology of Sport and Exercise*. 2003;4(1):25-39.
- 513 28. Rowlands A. Accelerometer Assessment of Physical Activity in Children: An Update.
514 *Pediatric Exercise Science*. 2007;19(3):252-66.
- 515 29. Reilly JJ, Penpraze V, Hislop J, Davies G, Grant S, Paton JY. Objective measurement of
516 physical activity and sedentary behaviour: review with new data. *Archives of Disease in Childhood*.
517 2008;93(7):614-9.
- 518 30. Ward DS, Evenson KR, Vaughn A, Rodgers AB, Troiano RP. Accelerometer Use in
519 Physical Activity: Best Practices and Research Recommendations. *American College of Sports*
520 *Medicine*. 2005;37(11):S582-S8.
- 521 31. Graves LE, Ridgers ND, Stratton G. The contribution of upper limb and total body
522 movement to adolescents' energy expenditure whilst playing Nintendo Wii. *European Journal of*
523 *Applied Physiology*. 2008;104(4):617-23.
- 524 32. Evenson KR, Catellier D, J, Gill K, Ondrak K, S, McMurray R, G. Calibration of two
525 objective measures of physical activity for children. *Journal of Sports Sciences*. 2008;24(14):1557-
526 65.
- 527 33. Sasaki JE, John D, Freedson PS. Validation and comparison of ActiGraph activity monitors.
528 *Journal of Science and Medicine in Sport*. 2011;14(5):411-6.
- 529 34. Trost SG, Loprinzi PD, Moore R, Pfeiffer KA. Comparison of Accelerometer Cut Points for
530 Predicting Activity Intensity in Youth. *Methodological Advances*. 2010;11(4307):1360-8.
- 531 35. Henry C. Basal metabolic rate studies in humans: measurement and development of new
532 equations. *Public Health Nutrition*. 2005;8(7a):1133-52.
- 533 36. Ridley K, Ainsworth BE, Olds TS. Development of a Compendium of Energy Expenditures
534 for Youth. *International Journal of Behavioral Nutrition and Physical Activity*. 2008;5(1):45.
- 535 37. Corder K, Brage S, Mattocks C, Ness A, Riddoch C, Wareham NJ, et al. Comparison of
536 Two Methods to Assess PAEE during Six Activities in Children. *Medicine & Science in Sports &*
537 *Exercise*. 2007;39(12):2180-8.
- 538 38. Nilsson A, Brage S, Riddoch C, Anderssen SA, Sardinha LB, Wedderkopp N, et al.
539 Comparison of equations for predicting energy expenditure from accelerometer counts in children.
540 *Scandinavian Journal of Medicine & Science in Sports*. 2008;18(5):643-50.
- 541 39. Trost SG, Way R, Okely AD. Predictive Validity of Three ActiGraph Energy Expenditure
542 Equations for Children. *Medicine & Science in Sports & Exercise*. 2006;38(2):380-7.
- 543 40. Cohen J. A power primer. *Psychological Bulletin*. 1992;112(1):155-9.

- 544 41. Cole TJ, Freeman JV, Preece MA. Body mass index reference curves for the UK, 1990.
545 Arch Dis Child. 1995;73(1):25-9.
- 546 42. Lyons EJ, Tate DF, Ward DS, Wang X. Energy intake and expenditure during sedentary
547 screen time and motion-controlled video gaming. Am J Clin Nutr. 2012;96(2):234-9.
- 548 43. Jackson A. Dietary Recommendations for Energy. Scientific Advisory Committee on
549 Nutrition 2010. London: Scientific Advisory Committee on Nutrition, 2011.
- 550 44. Gribbon A, McNeil J, Jay O, Tremblay MS, Chaput J-P. Active video games and energy
551 balance in male adolescents: a randomized crossover trial. The American Journal of Clinical
552 Nutrition. 2015.
- 553 45. Wang YC, Gortmaker SL, Sobol AM, Kuntz KM. Estimating the energy gap among US
554 children: A counterfactual approach. Pediatrics. 2006;118(6):e1721-e33.
- 555 46. Rumbold PLS, St Clair Gibson A, Allsop S, Stevenson E, Dodd-Reynolds CJ. Energy intake
556 and appetite following netball exercise over 5 days in trained 13-15 year old girls. Appetite.
557 2011;56(3):621-8.
- 558 47. Chaput J, P, Schwartz C, Boirie Y, Duclos M, Tremblay A, Thivel D. Energy intake
559 adaptations to acute isoenergetic active video games and exercise are similar in obese adolescents.
560 European Journal of Clinical Nutrition. 2015;1(5).
- 561 48. Thivel D, Chaput J-P. Are Post-Exercise Appetite Sensations and Energy Intake Coupled in
562 Children and Adolescents? Sports Med. 2014;44(6):735-41.
- 563 49. Temple JL, Giacomelli AM, Kent KM, Roemmich JN, Epstein LH. Television watching
564 increases motivated responding for food and energy intake in children. The American Journal of
565 Clinical Nutrition. 2007;85(2):355-61.
- 566 50. Dallman MF. Stress-induced obesity and the emotional nervous system. Trends in
567 Endocrinology & Metabolism. 2010;21(3):159-65.
- 568 51. Marsh S, Ni Mhurchu C, Maddison R. The non-advertising effects of screen-based
569 sedentary activities on acute eating behaviours in children, adolescents, and young adults. A
570 systematic review. Appetite. 2013;71(0):259-73.
- 571 52. Graves L, Stratton G, Ridgers ND, Cable NT. Energy expenditure in adolescents playing
572 new generation computer games. British Journal of Sports Medicine. 2008;42(7):592-4.

573