

1 **TITLE**

2 Phylogeny, phylogenetic inference and cranial evolution in pitheciids and *Aotus*

3

4 **SHORT TITLE**

5 Phylogeny and cranial evolution in pitheciids and *Aotus*

6

7 **AUTHORS**

8 Alexander Bjarnason*, Department of Anthropology, University College London, 14 Taviton
9 Street, London, WC1H 0BW

10

11 Christophe Soligo, Department of Anthropology, University College London, 14 Taviton
12 Street, London, WC1H 0BW

13

14 Sarah Elton, Department of Anthropology, Durham University, Dawson Building, South
15 Road, Durham, DH1 3LE

16

17 *corresponding author, ucsaobj@ucl.ac.uk

18 **ABSTRACT**

19 Pitheciids, one of the major radiations of New World monkeys endemic to South and Central
20 America, are distributed in the Amazon and Orinoco basins, and include *Callicebus*, *Cacajao*,
21 *Chiropotes* and *Pithecia*. Molecular phylogenetics strongly support pitheciid monophyly,
22 while morphological analyses infer a range of phylogenies including a sister relationship
23 between *Aotus* and *Callicebus*. We collected geometric morphometric cranial data from
24 pitheciids and *Aotus*, and used cranial data for distance-based phylogenetic analysis and tests
25 of phylogenetic signal. Phylogenetic analyses of pitheciids were repeated with *Lagothrix*,
26 *Callimico* and *Saimiri* outgroups for Procrustes shape with and without *Aotus* based on the
27 whole cranium and six anatomical regions. All phylogenetic signal tests were significant, and
28 tree lengths were shortest and had the least morphological change over the phylogeny for
29 Procrustes residuals from the cranial base and palate. The majority of phylogenetic analyses
30 of Procrustes shape for pitheciids without *Aotus* supported the molecular phylogeny, and with
31 *Aotus* included the majority inferred an *Aotus-Callicebus* clade, although three analyses with
32 *Callimico* as outgroup supported the molecular phylogeny. The morphological similarity of
33 *Aotus* and *Callicebus* is likely a mix of plesiomorphy, allometry and homoplasy, and future
34 phylogenetic inference of living and extinct platyrrhine taxa should consider the impact of
35 these factors alongside outgroup selection and cranial region.

36

37 **Key words: allometry; homoplasy; geometric morphometrics; platyrrhines**

38

39 **INTRODUCTION**

40 The pitheciids (family Pitheciidae; parvorder Platyrrhini) are one of the three major adaptive
41 radiations of primates endemic to South and Central America, and recent molecular analyses
42 estimate the pitheciid clade split from the atelids and cebids around 25 million years ago
43 (MYA) [Perelman et al., 2011; Wilkinson et al., 2011; Jameson Kiesling et al., 2015]. The
44 extant pitheciids are split into two subfamilies: Callicebinae for the smaller-bodied,
45 frugivorous titi monkeys (*Callicebus*), and Pitheciinae (the pitheciins), the larger-bodied,
46 specialized seed predators that includes sakis (*Pithecia*), bearded sakis (*Chiropotes*), and
47 uacaris (*Cacajao*).

48

49 Pitheciids are distributed in the Amazon and Orinoco basins, inhabit a range of habitats, are
50 arboreal and have a mixed locomotor repertoire [Kinzey, 1997; Norconk 2011]. The smallest
51 pitheciids belong to the genus *Callicebus*, with body masses of around 1kg, and the largest
52 pitheciid is the moderately sexually dimorphic *Cacajao*, with mean male body masses around
53 3.1 – 3.5 kg, depending on species, and females are about 20% smaller [Ford & Davis, 1992;
54 Smith & Jungers, 1997]. *Callicebus* and *Pithecia* have a relatively small brain size compared
55 to *Cacajao* and *Chiropotes*, which are both highly encephalized [Isler et al., 2008; Hartwig et
56 al., 2011]. The *Callicebus* diet is primarily frugivorous with some seed consumption, whereas
57 *Cacajao*, *Chiropotes* and *Pithecia* are predominantly seed predators [Norconk et al., 2009].
58 Seed predation involves sclerocarpic foraging and morphological adaptations to access hard,
59 thick fruits from which seeds are extracted, chewed and swallowed [Kinzey & Norconk,
60 1990, 1993; Kinzey, 1997].

61

62 Monophyly of *Cacajao*, *Chiropotes* and *Pithecia* have been acknowledged in all major
63 primate taxonomic classifications [Kinzey, 1992; Rosenberger et al., 1996]. Morphology-
64 based phylogenetic analyses of platyrrhines have also supported a pitheciin clade with
65 *Cacajao-Chiropotes* sister to *Pithecia* [Rosenberger, 1984; Ford, 1986; Kay, 1990, Horovitz,
66 1999]. However, the systematics of the family are not entirely straightforward. In particular,
67 the relationship with the nocturnal *Aotus* is controversial and there have been debates over the
68 position of *Callicebus*. An *Aotus-Callicebus* clade distantly related to the pitheciins has been
69 suggested [Ford, 1986], and *Aotus-Callicebus* has been placed as sister to the pitheciins
70 [Rosenberger, 1984]. Alternatively, *Callicebus* has been inferred as the basal-most
71 platyrrhine [Kay, 1990], or sister only to pitheciins [Horovitz, 1999].

72

73 Morphology and molecules appear to tell different stories with respect to *Callicebus* and
74 *Aotus*. Platyrrhine molecular phylogenetic data strongly support a pitheciid clade with
75 *Callicebus* basal-most and a sister relationship between *Pithecia* and *Cacajao-Chiropotes*,
76 and *Aotus* more closely related to *Cebus-Saimiri* and callitrichines than it is to *Callicebus* or
77 the pitheciids [Fig. 1: Wildman et al., 2009; Jameson Kiesling et al., 2015; Schneider &
78 Sampaio, 2015]. Despite the molecular data, *Aotus* and *Callicebus* have similar body masses
79 of around 1kg, are both primary frugivores with tall thin incisors and high
80 temporomandibular joints, are socially monogamous, have small group sizes, and low sexual
81 dimorphism [Kinzey, 1997; Rosenberger & Tejedor, 2013]. The two taxa are sympatric in
82 parts of Peru, and resource competition could be avoided through the evolution of nocturnal
83 behaviour in *Aotus* and reliance on alternative secondary dietary resources [Norconk et al.,
84 2009]. The morphological and behavioural similarities of *Aotus* and *Callicebus* have led
85 some researchers to consider them closely-related sister taxa [Rosenberger, 1981, 1984, 1992,
86 2002; Kinzey, 1992; Rosenberger et al., 2009; Rosenberger & Tejedor, 2013]. Nonetheless,

87 the two groups have some major biological differences, primarily because the nocturnal and
88 cathemeral activity of *Aotus* is unique among platyrrhines, resulting in its distinctive very
89 large orbits [Kinzey, 1997], and *Aotus* has a wider distribution across Central and South
90 America than pitheciids [Kinzey, 1997].

91

92 While both morphological and molecular data provide important information about
93 evolutionary biology, molecular phylogenetics have become ubiquitous as they tend to be
94 more robust and reliable approximations of evolutionary relationships [Scotland et al., 2003].
95 Morphological datasets generally contain hundreds of characters or anatomical landmarks,
96 whereas next-generation DNA and genome sequencing creates datasets with tens to hundreds
97 of thousands of characters per species for use in phylogenetic inference [Yang & Rannala,
98 2012]. These large molecular datasets use sophisticated statistics and models of evolution,
99 and combined with increased number of independent traits used, provide a clear advantage
100 over morphology-based analyses [Whelan et al., 2001]. However, molecular phylogenies can
101 vary due to differences between gene trees and species trees, the source of DNA (e.g. nuclear
102 or mitochondrial genomes) and use of coding or non-coding regions, variation in rates of
103 evolution, homoplasy, incomplete lineage sorting, and introgression amongst other factors
104 [Degnan & Rosenberg, 2009, Davalos et al., 2012]. They will not invariably recover the
105 ‘correct’ relationship, and as Perez & Rosenberger [2014] point out, major disparities are still
106 evident in relationships recovered for platyrrhines. Although there are discrepancies in the
107 position of *Aotus* in relation to callitrichines and *Cebus-Saimiri*, on balance it is likely the
108 molecular phylogenetic separation of *Aotus* and *Callicebus* is accurate.

109

110 This separation of *Aotus* from the pitheciids in turn suggests the proposed morphological
111 affinity of *Aotus* and *Callicebus* reflects either homology and retention of ancestral
112 platyrrhine plesiomorphic traits or homoplasy and convergence between the two taxa, but not
113 evidence of recent common ancestry. As molecular studies indicate the two groups last
114 shared a common ancestor approximately 25 million years ago [Perelman et al., 2011;
115 Wilkinson et al., 2011; Jameson Kiesling et al., 2015], it raises important research questions
116 applicable to platyrrhines and the palaeontological study of primates more generally. What
117 factors influenced *Aotus* and *Callicebus* convergence or lack of divergence from the common
118 ancestral form? If *Aotus* had gone extinct 1 million years ago and was only known from the
119 fossil record, given its social, ecological and biological similarities with *Callicebus*, would
120 the two groups be erroneously classified as closely related sister taxa? Given that recoverable
121 DNA is absent from most fossil taxa, resolving the “tree of life” of both extant and extinct
122 taxa will require sound and reliable phylogenetic inference using morphology [Wiens, 2004].

123

124 The development of geometric morphometric methods has provided new opportunities for
125 quantification and statistical analysis of morphology [Adams et al., 2004] which can be
126 applied to analyse morphological and phylogenetic relationships. Previous morphological
127 analyses that recovered a close sister relationship between *Aotus* and *Callicebus* were based
128 on character-state and cladistic techniques despite high levels of homoplasy across the
129 platyrrhine clade and most characters showing parallel evolution [Lockwood, 1999; Kay et
130 al., 2008]. In contrast, several large-scale studies of primates demonstrated geometric
131 morphometric data, with its ability to capture small yet significant shape variation, may find
132 greater congruence between molecular and morphological phylogenies [Lockwood et al.,
133 2004; Cardini & Elton, 2008b]. A major benefit of geometric morphometric methods is the
134 ability to separate size from shape, which can be used to investigate allometry, the study of

135 size and its consequences, particularly the relationship between body size and traits including
136 morphology, diet, behaviour, and ecology [Gould, 1966; Cheverud, 1982; Fleagle, 1995;
137 Mitteroecker et al., 2013]. Interspecific allometry –size-related differences between adults of
138 different species [Martin, 1990; Fleagle, 1995] – is important for pitheciid evolution, as the
139 largest taxon *Cacajao* is approximately three times larger than the smallest taxa *Callicebus*;
140 the similarities in body mass between the latter and *Aotus* could explain their morphological
141 and behavioural similarities.

142

143 Additionally, a combined geometric morphometric and modular approach to phylogenetic
144 inference using cranial variation can highlight which regions are congruent, and incongruent,
145 with molecular phylogenetic results. Modularity involves interaction and co-variation
146 between traits/variables in a shared region that are partially independent, with modules
147 partially distinct from each other in structure and function [Klingenberg, 2008]. If modules of
148 the cranium reflect alternative functional, developmental and evolutionary roles, the pattern
149 of similarity and utility of modules for accurate phylogenetic inference should vary [Wood &
150 Lieberman, 2001; Harvati & Weaver, 2006]. It is unlikely a single cranial anatomical region
151 will accurately infer phylogenetic relationships for all primate clades [von Cramon-Taubadel,
152 2014], creating the need to investigate each group individually. By examining whether
153 molecular clades are consistently inferred in some regions of the cranium compared to others,
154 the most informative regions may be targeted for phylogenetic reconstructions in fossil taxa,
155 provided appropriate specimens are available for study.

156

157 An important concept for understanding the relationship between molecular and
158 morphological evolution is the phylogenetic signal, where closely related taxa will be

159 phenotypically more similar to each other than either is to more distantly related taxa,
160 whereas a weak phylogenetic signal occurs when taxa are more similar to distant relatives or
161 similarity is distributed randomly across the phylogeny [Blomberg et al., 2003, Klingenberg
162 & Gidaszewski, 2010, Kamilar & Cooper, 2013]. The phylogenetic signal can also be
163 considered a statistical measure of the non-independence of trait similarity shared by taxa due
164 their phylogenetic relationships [Revell et al., 2008]. A strong phylogenetic signal is
165 predicted under a Brownian motion model of evolution, while the strength of phylogenetic
166 signal is phenotype and phylogeny dependent and can be lowered by adaptation,
167 measurement error of traits, and error in phylogenetic topology and branch lengths
168 [Blomberg & Garland, 2002, Kamilar & Cooper, 2013]. The phylogenetic signal of primates
169 across a range of phenotypic traits has provided insight into their evolution [Kamilar &
170 Cooper, 2013], and comparative study and quantification of which areas of morphology have
171 stronger or weaker phylogenetic signals can suggest which areas will be informative for
172 phylogenetic inference and help inform our understanding cranial evolution in groups of
173 interest.

174

175 In this paper, we examine the evolutionary relationships and phylogenetic signal of pitheciids
176 and *Aotus* based on geometric morphometric data from the cranium. We test two primary
177 hypotheses – [1] there is a phylogenetic signal in the pitheciid cranium, and a particular
178 cranial region and outgroup will find greater congruence between morphological and
179 molecular phylogenies; [2] that phylogenetic analysis of geometric morphometric data will
180 differentiate between *Aotus* and *Callicebus* and find little support for an *Aotus-Callicebus*
181 clade.

182 **METHODS**

183 This research complied with the American Society of Primatologists Principles for the Ethical
184 Treatment of Primates, protocols of the appropriate Institutional Animal Care Committee,
185 and legal requirements of each country housing collections.

186

187 Morphometric data, consisting of sixty-three 3D anatomical landmarks quantifying
188 morphological variation in the cranium (Table I) were collected from museum collections for
189 *Callicebus cupreus*, *Callicebus hoffmannsi*, *Callicebus moloch*, *Callicebus torquatus*,
190 *Cacajao calvus*, *Cacajao melanocephalus*, *Chiropotes satanas*, *Pithecia pithecia*, *Pithecia*
191 *monachus*, *Aotus azarae*, *Aotus lemurinus*, *Aotus vociferans*, *Aotus trivirgatus*, and outgroup
192 taxa *Callimico goeldii*, *Lagothrix lagotricha* and *Saimiri sciureus* (Table II). Museum
193 specimens were originally wild caught except for *Callimico goeldii* specimens that were all
194 captive. Despite the large number of pitheciid species recognized in recent taxonomic
195 classifications, adequate sample sizes are difficult to obtain from museum collections. The
196 3D anatomical landmarks were analysed with geometric morphometric methods (GMM) that
197 measure and preserve the geometry of structures being studied by removing non-biological
198 variation in scale, orientation and position of landmarks [Rohlf & Slice, 1990; Adams et al.,
199 2004]. The GMM methods used Generalised Procrustes Analysis (GPA), which has the
200 highest accuracy of available superimposition methods in estimating mean shape, lowest
201 error estimates, and greatest power to test for differences in mean shape between taxa
202 [Gower, 1975; Goodall, 1991; Rohlf, 2000a,b, 2003]. Procrustes shape coordinates describing
203 shape are distinct from the measure of size, centroid size, the square root of summed squared
204 distances between landmarks and their centroid [Mitteroecker et al., 2013] are produced
205 following GPA.

206

207 Geometric morphometric analysis was carried out in MorphoJ v1.06 (University of
208 Manchester, Manchester, UK; http://www.flywings.org.uk/morphoj_page.htm). Centroid
209 size, the square root of the sum of squared distances of landmarks from the centroid, is the
210 measure of size provided by GMM [Zelditch et al., 2004]. MorphoJ allows geometric
211 morphometric data to be mapped onto a phylogeny, in this case based on molecular
212 phylogenetic relationships of pitheciids with and without *Aotus*, using squared-change
213 parsimony to examine and quantify the phylogenetic signal. The phylogenetic signal will be
214 strongest when closely related taxa are phenotypically more similar to each other and occupy
215 similar morphometric space compared to more distantly related taxa [Klingenberg &
216 Gidaszewski, 2010]. This approach quantifies tree length based on the total sum of squared
217 change along all landmark coordinates and branches of the phylogeny, providing a single
218 measure of morphological change over the phylogeny provided, and morphometric data with
219 a stronger phylogenetic signal will have less shape change across the branches of the
220 phylogenetic tree and shorter tree lengths, whereas morphometric data with a lower
221 phylogenetic signal will exhibit greater morphological change along branches of the
222 phylogeny and have longer tree lengths [Klingenberg & Gidaszewski, 2010]. The
223 measurement of the phylogenetic signal uses permutations to test the null hypothesis of no
224 phylogenetic signal by resampling taxa, recalculating tree length, and providing a *P* value for
225 the proportion of resampled datasets with a shorter or equal tree length compared to the
226 original dataset [Klingenberg & Gidaszewski, 2010]. If the null hypothesis of no
227 phylogenetic signal is true, the permutation test that randomly swaps the morphometric
228 values at the tip of the phylogeny should not alter tree length and morphological change
229 compared to the original data, while the tree length would increase if the permutation acted
230 on morphometric data with a phylogenetic signal. Different phylogenetic signal results are

231 best considered comparatively where the same phylogeny and alternative shape data, or
232 alternative phylogenies and the same shape data, are used.

233

234 The phylogenetic signal in both shape (based on Procrustes coordinates) and size (based on
235 log centroid size) were analysed with and without *Aotus* included, and no outgroup, requiring
236 separate input phylogenies to quantify the phylogenetic signal based on the molecular
237 analyses of all platyrrhines. These phylogenies, based on relationships supported by multiple
238 molecular phylogenetic studies had *Aotus* sister to pitheciids, within which *Callicebus* is
239 basal-most and *Pithecia* is sister to *Cacajao-Chiropotes*, and for analyses of just pitheciids
240 the same phylogenetic relationships with *Aotus* removed [Perelman et al., 2011; Jameson
241 Kiesling et al., 2015; Schneider & Sampaio, 2015]. As neither Perelman and colleagues
242 [2011] nor Jameson Kiesling and colleagues [2015] used the neighbor-joining method for
243 phylogenetic inference, for consistency we accessed their publically available molecular
244 datasets and ran neighbor-joining in PAUP 4 (Sinauer Associates, Sunderland,
245 Massachusetts, USA; <http://paup.sc.fsu.edu/>), which supported the previously described
246 pitheciid relationships and placement of *Aotus* within cebids. Considering the species-level
247 relationships within *Callicebus* and *Aotus* are not fully resolved, the relationships within each
248 genus were treated as unresolved polytomies.

249

250 Euclidean morphological distances were used for phylogenetic construction using neighbor-
251 joining in the Neighbor module of Phylip 3.6 (University of Washington, Seattle,
252 Washington, USA; <http://evolution.genetics.washington.edu/phylip.html>). Neighbor-joining
253 constructs a phylogeny with a stepwise additive method based on a divisive cluster algorithm
254 that minimizes overall branch length, is statistically consistent, inferring the correct

255 evolutionary tree when distances accurately reflect phylogeny, assumes distances between
256 two taxa are equal to the distance between each respective group and a shared node, and roots
257 the tree using an outgroup taxa [Saitou & Nei, 1987; Kuhner & Felsenstein, 1994; Yang,
258 2006].

259

260 Selection of outgroup taxa can impact phylogenetic inference of morphology [e.g. Bjarnason
261 et al., 2011, 2015], and although a plesiomorphic fossil platyrrhine taxa would make an ideal
262 outgroup, in the absence of an adequately large sample size of specimens, using geometric
263 morphometric data for fossil taxa is difficult due to increased error rates in estimating mean
264 shape with low sample sizes [Cardini & Elton, 2008b], and distortion to fossil specimens can
265 require considerable virtual reconstruction [e.g. Zollikofer et al., 2005, Spoor et al., 2015]. As
266 two of the five major extant platyrrhine clades, pitheciids and *Aotus*, are ingroup taxa, one
267 outgroup was sampled from each of the three remaining clades, with phylogenetic inference
268 repeated using an atelid, callitrichine and cebine outgroup. The atelid *Lagothrix lagotricha*
269 was selected as it is likely the closest to the ancestral atelid phenotype and least derived
270 extant group in that clade [Rosenberger & Strier, 1989, Bjarnason et al., 2015], and *Callimico*
271 *goeldii* has lost multiple typically callitrichine traits in morphology and reproduction and
272 likely acquired secondarily derived traits similar to the ancestral platyrrhine [Martin, 1992,
273 Pastorini et al., 1998, Scott, 2015]. As allometry and the size of outgroups, and its impact on
274 phylogenetic inference, is of interest [Bjarnason et al., 2011], we selected outgroups that were
275 considerably larger (*Lagothrix lagotricha*) and smaller (*Callimico goeldii*) than ingroup taxa,
276 in addition to a third outgroup (*Saimiri sciureus*) that is derived in morphology but shares
277 ancestral platyrrhine body size with *Aotus* and *Callicebus* [Ford & Davis, 1992].

278

279 Statistical support for clades was quantified using a jack-knife method where phylogenetic
280 analysis and Procrustes superimposition was repeated with each landmark removed, with
281 percentage clade support the number of times a clade was present in each phylogenetic
282 analysis, and results were collated using the Consensus module in Phylip [Felsenstein, 2005].
283 Majority consensus trees were drawn using TreeView (University of Glasgow, Glasgow, UK;
284 <https://www.ctu.edu.vn/~dvxe/Bioinformatic/Software/Rod%20Page/treeview.html>) and
285 TreeGraph 2 (University of Münster, Münster, Germany;
286 <http://treegraph.bioinfweb.info/Download>). As with the tests of a phylogenetic signal, the
287 neighbour-joining phylogenetic analysis was repeated to include pitheciids only, and with
288 pitheciids and *Aotus* as ingroup taxa.

289

290 Tests for phylogenetic signal and neighbour-joining phylogenetic analysis were all repeated
291 with morphometric data from the whole cranium, and hypothesized modules within the
292 cranium. Cranial modules of the orofacial and neurocranium are recognized with further
293 subdivision into the face, palate/oral, nasal, zygomatic, cranial base and cranial vault
294 [Cheverud, 1982; Hallgrímsson et al., 2004], in addition to larger modules for the
295 chondrocranium of the cranial base and dermatocranium of the face and cranial vault based
296 on mode of ossification [Hallgrímsson et al., 2004; Cardini & Elton, 2008a]. Cardini & Elton
297 [2008a] have shown sampling error becomes high in modules with low numbers of
298 landmarks, and we are unable to analyse orbit and zygomatic modules in our cranial dataset
299 due to the low number of landmarks. Modules of the cranial vault and palate region had too
300 few landmarks to be analysed as individual modules, but were combined with the face and
301 cranial base in a series of landmark combinations. Overall, seven regions were analysed: the
302 cranium (landmarks 1-63), face (landmarks 1-15), face and palate (landmarks 1-15, 30-38),
303 face and cranial vault (landmarks 1-26, including landmarks 17-19 from the zygomatic arch),

- 304 cranial base (landmarks 40-63), cranial base and vault (landmarks 16, 20-26, 40-63), and
- 305 cranial base and palate (landmarks 30-63, including landmark 39 that falls between regions).

306 **RESULTS**

307 The measures of phylogenetic signal for Procrustes coordinates and log centroid size, without
308 and with *Aotus*, are presented in Table III based on tree length and a permutation test of
309 significance. The permutation test of significance takes morphometric values at the tip of a
310 phylogeny and randomly swaps them, which will have no effect on tree length if there is no
311 phylogenetic signal, but will be significantly different to the tree length from the original data
312 if a phylogenetic signal is present- our results show a phylogenetic signal is present for all
313 iterations, rejecting the null hypothesis there is no phylogenetic signal in cranial data. Tree
314 length quantifies the combined morphological change across all branches of a phylogeny,
315 with lower tree lengths signifying less morphological change and a stronger phylogenetic
316 signal, and larger tree lengths involving greater morphological change and a weaker
317 phylogenetic signal. For each cranial region in pitheciid analyses without *Aotus*, log centroid
318 size tree lengths were longer than for Procrustes coordinates with the exception of the cranial
319 base and palate. For pitheciid analyses including *Aotus*, tree lengths were longer than for
320 analyses without *Aotus* as expected considering the increased taxa sampling, and for each
321 cranial region the tree lengths from Procrustes coordinates were longer than for log centroid
322 size except for the cranial base and palate, and face and palate. For shape coordinates, for
323 pitheciids both with and without *Aotus*, the region with the strongest phylogenetic signal,
324 shortest tree lengths and least morphological change across the phylogeny was the cranial
325 base and palate, followed by the cranium, cranial base and vault, cranial base, face and
326 cranial vault, face, and the weakest phylogenetic signal was in the face and palate.

327

328 The results of neighbour-joining phylogenetic analysis are provided at the genus level as
329 majority consensus trees (Figs. 2-3) and jack-knife clade support (Tables IV-V) for pitheciids
330 with and without *Aotus* included as ingroup taxa. Phylogenetic analysis of pitheciids-only

331 (Fig. 2 and Table IV) supported the molecular phylogeny with *Cacajao-Chiropotes* sister to
332 *Pithecia* and *Callicebus* basal-most in eleven of twenty-one analyses, supported a dichotomy
333 between *Callicebus-Pithecia* and *Cacajao-Chiropotes* in nine analyses, and *Callicebus* sister
334 to *Cacajao-Chiropotes* and *Pithecia* basal-most in one analysis.

335

336 Phylogenetic analyses of pitheciids with *Aotus* (Fig. 3 and Table V) supported an *Aotus-*
337 *Callicebus* clade in sixteen of twenty-one analyses. Eleven analyses placed *Cacajao-*
338 *Chiropotes* basal-most and *Pithecia* sister to *Aotus-Callicebus*, and three analyses inferred
339 *Aotus-Callicebus* basal-most and *Pithecia* sister to *Cacajao-Chiropotes*. A further three
340 analyses inferred *Cacajao-Chiropotes* sister to *Pithecia* in a clade with *Aotus*, and *Callicebus*
341 basal-most, and one analysis inferred a dichotomy between *Aotus-Callicebus* and *Cacajao-*
342 *Chiropotes* with *Pithecia* basal-most. Pitheciid monophyly and the molecular phylogeny with
343 *Cacajao-Chiropotes* sister to *Pithecia*, *Callicebus* within the pitheciids and *Aotus* basal-most
344 was inferred for three analyses with *Callimico* as outgroup.

345 **DISCUSSION**

346 Phylogenetic analysis of pitheciid cranial variation confirms the first hypothesis of the
347 presence of a phylogenetic signal, with a complex mix of congruence between molecular and
348 morphological phylogenies depending on ingroup taxa, outgroup selection and cranial region.
349 However, considering the majority of phylogenies constructed including pitheciids and *Aotus*
350 inferred an *Aotus-Callicebus* clade, we reject the second hypothesis that phylogenetic
351 analysis of geometric morphometric data would differentiate between the two taxa in the
352 majority of analyses, and support earlier findings of a morphological affinity between
353 *Callicebus* and *Aotus* [e.g. Rosenberger, 1984, 2002; Kinzey, 1992; Rosenberger et al., 2009;
354 Rosenberger & Tejedor, 2013].

355

356 Rosenberger & Tejedor [2013] view the similarity of *Aotus* and *Callicebus* as phylogenetic,
357 and propose that long-branch attraction in molecular phylogenetics has mis-placed *Aotus*
358 outside of the pitheciids. However, there are a number of other evolutionary scenarios that
359 could explain similarities between *Aotus* and *Callicebus*: (a) *Aotus* and *Callicebus* have
360 maintained plesiomorphic primitive ancestral traits in size, morphology and behaviour, for
361 over 25 million years; (b) *Aotus* and *Callicebus* have undergone major homoplasy, whereby
362 similarity shared by taxa is not due to common ancestry [Lockwood & Fleagle, 1999], and
363 converged upon the same size, morphology and behaviour via convergence in similar
364 ecological and social environments; or (c) a complex mix of the two, with a combination of
365 ancestral and convergent traits.

366

367 Interpretation of the early platyrrhine fossil record is important for considering the extent of
368 plesiomorphy and homoplasy found in *Aotus* and pitheciids, although the topic is contentious.
369 The long lineage hypothesis considers extant platyrrhines a more ancient radiation and

370 positions early fossil taxa such as *Tremacebus* and *Soriacebus* within clades alongside extant
371 groups [e.g. Rosenberger et al., 2009, Rosenberger, 2010], whereas the layered hypothesis
372 views extant clades and fossil taxa descended from the crown group common ancestor as a
373 more recent radiation and places several of the earliest platyrrhine fossil taxa outside the
374 crown group as stem platyrrhines [e.g. Kay, 1990, 2015, Kay et al., 2008]. Both hypotheses
375 require extensive homoplasy [Rosenberger 2002, Kay & Fleagle 2010], but differ in an
376 important interpretation of living and fossil groups fundamental to understanding the
377 similarity of *Aotus* and *Callicebus*. The long lineage hypothesis views seed predation in
378 *Soriacebus* as providing an ecophylogenetic link to pitheciids and traits in orbit morphology
379 in *Tremacebus* and *Aotus* are due to shared ancestry [Rosenberger, 2010], indicating traits
380 connecting *Aotus* and *Callicebus* are similarly derived and phylogenetic. In contrast, the
381 layered hypothesis views *Tremacebus* and *Soriacebus* as stem platyrrhines rather than close
382 relatives of *Aotus* and pitheciids [Kay et al., 2008, Kay, 2015], with many similarities
383 between stem and crown groups primitive traits, indicating *Aotus* and *Callicebus* shared traits
384 are ancestral for platyrrhines.

385

386 With debate still ongoing over the long lineage and layered hypotheses, we propose the
387 molecular phylogenetic separation of *Aotus* and *Callicebus* is accurate and that a mix of
388 plesiomorphy, allometry and homoplasy combines to drive morphological and behavioural
389 similarity rather than recent common ancestry. While *Aotus* and *Callicebus* may retain the
390 plesiomorphic platyrrhine body size [Ford & Davis, 1992] alongside several other ancestral
391 traits, the callitrichine-like body size of the earliest platyrrhine fossil *Perupithecus* [Bond et
392 al., 2015] suggests a smaller ancestral body size and convergent size evolution in *Aotus* and
393 *Callicebus*, although that interpretation depends on whether *Perupithecus* belongs to a crown
394 or stem group and is representative of the platyrrhine common ancestor. Whether shared body

395 size is ancestral or derived in *Aotus* and *Callicebus*, it seems probable they will share other
396 plesiomorphic traits, yet homoplasy remains a pervasive evolutionary reality [Kay & Fleagle,
397 2010]. Platyrrhine morphological characters are known to have high levels of homoplasy
398 [Lockwood, 1999], nearly all phylogenetically informative traits from the platyrrhine
399 cladistic analysis of Kay and colleagues [2008] showed some parallel evolution, and due to
400 the high levels of homoplasy morphological characters can be used in support of most
401 phylogenetic relationships [Kay, 2015]. As homoplasy is widespread in the platyrrhine clade,
402 allometry is a particularly powerful intrinsic factor in morphological homoplasy [Lockwood
403 & Fleagle, 1999; Kay & Fleagle, 2010], and post-cranial traits shared by *Aotus* and
404 *Callicebus* have been linked to parallel evolution [Lockwood, 1999], it is likely some of the
405 traits shared by *Aotus* and *Callicebus* are due to homoplasy.

406

407 The body size similarity and allometric link between *Aotus* and *Callicebus* contributes to
408 shared morphological similarity, but a key factor in morphology-based phylogenetic
409 inference is also the allometric relationship between outgroup and ingroup taxa. This issue
410 has been previously highlighted in hominoids, where allometric scaling and cranial shape
411 linked to brain size in *Hylobates* and *Homo* complicate accurate phylogenetic inference
412 [Creel, 1986; Bjarnason et al., 2011]. The phylogenetic analyses of pitheciids including *Aotus*
413 with *Saimiri* as outgroup inferred an *Aotus-Callicebus* clade in all seven analyses, and *Aotus*,
414 *Callicebus* and *Saimiri* share a similar body size. Using the much larger-bodied *Lagothrix*
415 outgroup supported *Aotus-Callicebus* in six of seven analyses, whereas the smaller-bodied
416 *Callimico* outgroup inferred *Aotus-Callicebus* in two analyses, and the molecular phylogeny
417 in three. This does not mean using a smaller-bodied outgroup will reduce the influence of
418 allometry on all morphology-based phylogenetic analyses as it will be dependent up the
419 allometric relationships within the ingroup, as in Old World monkeys [e.g. Gilbert & Rossie,

420 2007; Gilbert et al., 2009] and between ingroup and outgroup taxa, and the issue remains
421 pertinent for accuracy of phylogenetic inference and study of primate groups.

422

423 The relative lack of support for a monophyletic pitheciid clade when *Aotus* is included in
424 analyses contrasts with the eleven analyses that support the molecular phylogenetic
425 relationships when only pitheciid cranial data is analysed. This reflects the evolution of
426 multiple traits including morphological adaptations, diet, and relative brain size, which
427 broadly follow a morphocline, with *Callicebus* expressing a relatively ancestral or primitive
428 phenotype, *Pithecia* an intermediate or partially derived condition, and *Cacajao* and
429 *Chiropotes* sharing a derived phenotype [Kinzey, 1992]. For example, in cranial morphology
430 the differentiation in phylogenetic analysis between *Callicebus* and the pitheciins *Cacajao*,
431 *Chiropotes* and *Pithecia* reflects the latter as specialized sclerocarpic foragers with incisor
432 and canine adaptations and enlarged temporalis and masseter muscles able to generate high-
433 forces to open hard-tusked fruits [Kinzey & Norconk, 1990, 1993; Kinzey, 1992, 1997].
434 Allometry also helps maintain a phylogenetic signal with inference of the smallest lineage
435 *Callicebus* basal-most and a sister relationship between the two largest genera, *Chiropotes*
436 and *Cacajao*. The choice of outgroup is clearly also important, as six of seven phylogenetic
437 analyses with *Callimico* inferred the pitheciid molecular phylogeny, whereas six of seven
438 analyses using *Saimiri* as outgroup inferred a dichotomy including a *Pithecia-Callicebus*
439 clade not supported by molecular phylogenetics.

440

441 From our data, all cranial regions had a phylogenetic signal, but there were clear differences
442 in tree lengths for different regions. The region with the strongest phylogenetic signal, the
443 cranial base and palate, had a tree length one third of the tree length for the region with the
444 weakest phylogenetic signal, the face and palate, meaning there has been greater

445 morphological change over the phylogeny in the face and palate. The maintenance of a
446 stronger phylogenetic signal in cranial base morphology has been hypothesized as due to
447 strong genetic control and a role in multiple functional systems compared to the more plastic
448 face that is shaped by environmental factors [e.g. Olson, 1981; Lieberman et al., 1996;
449 Lieberman, 1997]. However, Revell and colleagues [2008] cautions against linking strong
450 and weak phylogenetic signals with concepts of conserved or plastic traits, as an array of
451 evolutionary processes and rates of evolution can create a similar phylogenetic signal, and
452 very similar processes can lead to varied phylogenetic signals.

453

454 While the region of the cranial base and palate has the strongest phylogenetic signal of the
455 regions investigated here in pitheciids and *Aotus*, the phylogenetic signal in phenotypic traits
456 will likely vary dependent on the taxonomic and phylogenetic level [Kamilar & Cooper,
457 2013], and no single cranial region will maintain the strongest phylogenetic signal across all
458 primates [von Cramon-Taubadel, 2014]. It is worth considering an additional issue; how a
459 region can have a strong phylogenetic signal, yet phylogenetic inference based on data from
460 that region often fails to support evolutionary relationships strongly supported by molecular
461 data. For our three regions with the strongest phylogenetic signal, the cranial base and palate,
462 cranium, and cranial base and vault, phylogenetic inference that included pitheciids and *Aotus*
463 inferred non-molecular clades in each analysis using *Lagothrix* and *Saimiri* outgroups, but
464 inferred the molecular phylogeny in all three analyses with *Callimico* as outgroup. This
465 suggests the presence of a strong phylogenetic signal is not, of itself, enough to find
466 congruence between molecular and morphological phylogenies, but as has been shown in
467 other primate groups [e.g. Bjarnason et al., 2011, 2015] methodological decisions such as
468 outgroup selection and rooting are integral to using a strong phylogenetic signal for accurate
469 phylogenetic inference.

470

471 To return to one of our original questions, if *Aotus* was known only from the fossil record and
472 included in a phylogenetic analysis with pitheciids, it would probably be erroneously
473 classified as sister to *Callicebus* – our study, in common with several others demonstrates the
474 morphological similarity between the two taxa despite their deep divergence. This
475 morphological connection is likely to be a mix of the retention of ancestral platyrrhine traits
476 and convergence, both with a link to allometry and similar dietary niches, body mass and
477 cranial form in *Aotus* and *Callicebus*. By considering the effects of allometry, outgroup
478 selection and modularity on phylogenetic analysis alongside the benefits of including fossil
479 taxa, combined datasets, molecular scaffolds and character weighting, it should be possible to
480 have greater confidence in assessing phylogenetic relationships and derived similarity in the
481 platyrrhine fossil record than appears initially from the *Aotus-Callicebus* example.

482 **ACKNOWLEDGMENTS**

483 This project was originally conceived and planned with Charlie Lockwood. We thank the
484 following institutions for access to their collections: Natural History Museum, London; Field
485 Museum of Natural History, Chicago; Museum für Naturkunde, Berlin; Naturhistorisches
486 Museum, Vienna; Smithsonian National Museum of Natural History, Washington, DC;
487 Naturhistoriska Riksmuseet, Stockholm; and the Anthropological Institute & Museum of the
488 University of Zurich. We thank Louise Tomsett, Roberto Portela Miguez, Paula Jenkins, Bill
489 Stanley, Bettina Wimmer, Frieder Mayer, Barbara Herzig, Olavi Gronwall, Tea Jashashvili,
490 and Marcia Ponce de Leon for access to collections, and Andrea Cardini, Brian Villmoare,
491 Alfie Rosenberger, Chris Klingenberg, Jim Rohlf, and Marilyn Norconk for help and advice.
492 We would like to thank the Executive Editor, Paul A. Garber, the Review Editor, Donald C.
493 Dunbar, and both anonymous reviewers for their insightful and helpful comments, and the
494 time they have taken to review our work. A. Bjarnason received financial support for the
495 research undertaken from the Department of Anthropology (University College London), the
496 Graduate Research Fund (UCL Graduate School), the Central Research Fund (University of
497 London), and SYNTHESYS.

498 **REFERENCES**

- 499 Adams DC, Rohlf FJ, Slice DE. 2004. Geometric morphometrics: Ten years of progress
500 following the 'revolution'. *Italian Journal of Zoology* 71:5-16.
- 501 Bjarnason A, Chamberlain AT, Lockwood CA. 2011. A methodological investigation of
502 hominoid craniodental morphology and phylogenetics. *Journal of Human Evolution* 60:47-
503 57.
- 504 Bjarnason A, Soligo C, Elton S. 2015. Phylogeny, ecology, and morphological evolution in
505 the atelid cranium. *International Journal of Primatology* 36: 513-529.
- 506 Blomberg SP, Garland Jr T. 2002. Tempo and mode in evolution: phylogenetic inertia,
507 adaptation and comparative methods. *Journal of Evolutionary Biology* 15: 899-910.
- 508 Blomberg SP, Garland Jr T, Ives AR. 2003. Testing for phylogenetic signal in comparative
509 data: behavioural traits are more labile. *Evolution* 57:717-745.
- 510 Bond M, Tejedor MF, Campbell KE Jr, Chornogubsky L, Novo N, Goin F. 2015 Eocene
511 primates of South America and the African origins of New World monkeys. *Nature* 520: 538-
512 541.
- 513 Cardini A, Elton S. 2008a. Does the skull carry a phylogenetic signal? *Evolution and*
514 *modularity in the guenons*. *Biological Journal of the Linnean Society* 93:813-834.
- 515 Cardini A, Elton S. 2008b. Variation in guenon skulls I: species divergence, ecological and
516 genetic differences. *Journal of Human Evolution* 54: 615-637.
- 517 Cheverud JM. 1982. Phenotypic, genetic, and environmental morphological integration in the
518 cranium. *Evolution* 36:499-516.
- 519 Creel, N. 1986. Size and phylogeny in hominoid primates. *Systematic Zoology* 18:81-99.

- 520 Davalos LM, Cirranello AL, Geisler JH, Simmons NB. 2012. Understanding phylogenetic
521 incongruence: lessons from phyllostomid bats. *Biological Reviews* 87:991-1024.
- 522 Degnan, JH, Rosenberg NA. 2009. Gene tree discordance, phylogenetic inference and the
523 multispecies coalescent. *Trends in Ecology & Evolution* 24:332-340.
- 524 Fleagle JG. 1995. Size and adaptation in primates. In: Jungers WL, editor. *Size and Scaling in*
525 *Primate Biology*. New York: Plenum Press. p 1-19.
- 526 Ford SM. 1986. Systematics of the New World monkeys. In: Swindler DR, Erwin J, editors.
527 *Comparative Primate Biology Volume 1: Systematics, Evolution, and Anatomy*. New York:
528 Alan R. Liss, Inc. p 73-135.
- 529 Ford SM, Davis LC. 1992. Systematics and body size: Implications for feeding adaptations in
530 New World monkeys. *American Journal of Physical Anthropology* 88:415-468.
- 531 Gilbert CC, Rossie JB. 2007. Congruence of molecular and morphology using a narrow
532 allometric approach. *Proceedings of the National Academy of Sciences* 104:11910-11914.
- 533 Gilbert CC, Frost SR, Strait DS. 2009. Allometry, sexual dimorphism, and phylogeny: a
534 cladistic analysis of extant African papionins using craniodental data. *Journal of Human*
535 *Evolution* 57:298-320.
- 536 Goodall C. 1991. Procrustes methods in the statistical analysis of shape. *Journal of the Royal*
537 *Statistical Society Series B (Methodological)* 53:285-339.
- 538 Gould SJ. 1966. Allometry and size in ontogeny and phylogeny. *Biological Reviews* 41:587-
539 640.
- 540 Gower JC. 1975. Generalized Procrustes analysis. *Psychometrika* 40:33-51.

- 541 Hallgrímsson B, Willmore K, Dorval C, Cooper DM. 2004. Craniofacial variability and
542 modularity in macaques and mice. *Journal of Experimental Zoology. Part B Molecular and*
543 *Developmental Evolution* 302:207-225.
- 544 Hartwig WC, Rosenberger AL, Norconk M, Owi MY. 2011. Relative brain size, gut size, and
545 evolution in New World monkeys. *The Anatomical Record* 294:2207-2221.
- 546 Harvati K, Weaver T. 2006. Human cranial anatomy and the differential preservation of
547 population history and climate signatures. *The Anatomical Record* 288:1225–1233.
- 548 Horowitz I. 1999. A phylogenetic study of living and fossil platyrrhines. *American Museum*
549 *Novitates* 3269:1-40.
- 550 Isler K, Kirk EC, Miller JMA, Albrecht GA, Gelvin BR, Martin RD. 2008. Endocranial
551 volume of primate species: Scaling analyses using a comprehensive and reliable dataset.
552 *Journal of Human Evolution* 55:967-978.
- 553 Jameson Kiesling NM, Yi SV, Xu K, Gianluca Sperone F, Wildman DE. 2015. The tempo
554 and mode of New World monkey evolution and biogeography in the context of
555 phylogenomic analysis. *Molecular Phylogenetics and Evolution* 82:386-399.
- 556 Kamilar JM, Cooper N. 2013. Phylogenetic signal in primate behaviour, ecology and life
557 history. *Philosophical Transactions of the Royal Society B* 368:20120341.
- 558 Kay RF. 1990. The phyletic relationships of extant and fossil Pitheciinae [Platyrrhini,
559 Anthropoidea]. *Journal of Human Evolution* 19:175-208.
- 560 Kay RF. Biogeography in deep time- What do phylogenetics, geology, and paleoclimate tell
561 us about early platyrrhine evolution? *Molecular Phylogenetics and Evolution* 82: 358-374.
- 562 Kay RF, Fleagle JG. 2010. Stem taxa, homoplasy, long lineages, and the phylogenetic
563 position of *Dolichocebus*. *Journal of Human Evolution* 59:218-222.

- 564 Kay RF, Fleagle JG, Mitchell TRT, Colbert M, Brown T, Powers DW. 2008. The anatomy of
565 *Dolichocebus gaimanensis*, a stem platyrrhine monkey from Argentina. *Journal of Human*
566 *Evolution* 54:323-382.
- 567 Kinzey WG. 1992. Dietary and dental adaptations in the Pitheciinae. *American Journal of*
568 *Physical Anthropology* 88:499-514.
- 569 Kinzey WG. 1997. *New World Primates: Ecology, Evolution, and Behavior*. New York:
570 Aldine de Gruyter. 437 p.
- 571 Kinzey WG, Norconk MA. 1990. Hardness as a basis of fruit choice in two sympatric
572 primates. *American Journal of Physical Anthropology* 81:5-15.
- 573 Kinzey WG, Norconk MA. 1993. Physical and chemical properties of fruit and seeds eaten
574 by *Pithecia* and *Chiropotes* in Surinam and Venezuela. *International Journal of Primatology*
575 14:207-227.
- 576 Klingenberg CP. 2008. Novelty and "homology-free" morphometrics. *Evolutionary Biology*
577 35:186-190
- 578 Klingenberg CP, Gidaszewski NA. 2010. Testing and quantifying phylogenetic signals and
579 homoplasy in morphometric data. *Systematic Biology* 59:245-261.
- 580 Kuhner MK, Felsenstein J. 1994. A simulation comparison of phylogeny algorithms under
581 equal and unequal evolutionary rates. *Molecular Biology and Evolution* 11:459-468.
- 582 Lieberman DE. 1997. Making behavioural and phylogenetic inferences from hominid fossils:
583 Considering the developmental influence of mechanical forces. *Annual Review of*
584 *Anthropology* 26:185–210.

- 585 Lieberman DE, Wood BA, Pilbeam, DR. 1996. Homoplasy and early *Homo*: An analysis of
586 the evolutionary relationships of *H. habilis sensu stricto* and *H. rudolfensis*. Journal of
587 Human Evolution 30:97–120.
- 588 Lockwood CA. 1999. Homoplasy and adaptation in the atelid postcranium. American Journal
589 of Physical Anthropology 108: 459-482.
- 590 Lockwood CA, Fleagle JG. 1999. The recognition and evaluation of homoplasy in primate
591 and human evolution. American Journal of Physical Anthropology 110:189-232.
- 592 Lockwood CA, Kimbel WH, Lynch JM. 2004. Morphometrics and hominoid phylogeny:
593 Support for a chimpanzee-human clade and differentiation among great ape subspecies.
594 Proceedings of the National Academy of Sciences 101:4356-4360.
- 595 Martin RD. 1990. Primate Origins and Evolution: A Phylogenetic Reconstruction. London:
596 Chapman Hall. 828 p.
- 597 Mitteroecker P, Gunz P, Windhager S, Schaefer K. 2013. A brief review of shape, form, and
598 allometry in geometric morphometrics, with applications to human facial morphology.
599 Hystrix, the Italian Journal of Mammalogy 24:59-66.
- 600 Norconk MA. 2011. Sakis, uakaris, and titi monkeys. In: Campbell CJ, Fuentes A,
601 MacKinnon KC, Bearder SK, Stumpf RM, editors. Primates in Perspective. Oxford: Oxford
602 University Press. p122-139.
- 603 Norconk MA, Wright BW, Conklin-Brittain NL, Vinyard CJ. 2009. Mechanical and
604 nutritional properties of food as factors in platyrrhine dietary adaptations. In: Garber PA,
605 Estrada A, Bicca-Marques JC, Heymann EW, Strier KB, editors. South American Primates
606 Comparative Perspectives in the Study of Behaviour, Ecology, and Conservation. New York:
607 Springer. p279-319.

- 608 Olson, T. R. 1981. Basicrania and evolution of the pliocene hominids. In: Stringer, CB,
609 editor. Aspects of Human Evolution. London: Taylor and Francis. p99-128.
- 610 Pastorini J, Forstner MR, Martin RD, Melnick DJ. A reexamination of the phylogenetic
611 position of *Callimico* (Primates) incorporating new mitochondrial DNA sequence data. 1998.
612 Journal of Molecular Evolution 47:32-41.
- 613 Perelman P, Johnson WE, Roos C, Seuanez HN, Horvath JE, Moreira MA, Kessing B,
614 Pontius J, Roelke M, Rumppler Y, Schneider MP, Silva A, O'Brien SJ, Pecon-Slattery J. 2011.
615 A molecular phylogeny of living primates. PLOS Genetics 7:e1001342.
- 616 Perez SI, Rosenberger AL 2014. The status of platyrrhine phylogeny: A meta-analysis and
617 quantitative appraisal of topological hypotheses. Journal of Human Evolution 76: 177-187.
- 618 Revell LJ, Harmon LJ, Collar DC. 2008. Phylogenetic signal, evolutionary process, and rate.
619 Systematic Biology 57:591-601.
- 620 Rohlf FJ. 2000a. On the use of shapes spaces to compare morphometric methods. Hystrix
621 11:1-17.
- 622 Rohlf FJ. 2000b. Statistical power comparisons among alternative morphometric methods.
623 American Journal of Physical Anthropology 111:463-478.
- 624 Rohlf FJ. 2003. Bias and error in estimates of mean shape in geometric morphometrics.
625 Journal of Human Evolution 44:665-683.
- 626 Rohlf FJ, Slice D. 1990. Extensions of the Procrustes method for the optimal superimposition
627 of landmarks. Systematic Zoology 39:40-59.
- 628 Rosenberger AL. 1981. Systematics: the higher taxa. In: Coimbra-Filho AF, Mittermeier,
629 R.A., editor. Ecology and Behaviour of Neotropical Primates. Rio de Janeiro: Academia
630 Brasilia de Ciencias. p9-27.

631 Rosenberger AL. 1984. Fossil New World monkeys dispute the molecular clock. *Journal of*
632 *Human Evolution* 13:737-742.

633 Rosenberger AL. 1992. Evolution of feeding niches in New World monkeys. *American*
634 *Journal of Physical Anthropology* 88:525-562.

635 Rosenberger AL. 2002. Platyrrhine paleontology and systematics: The paradigm shifts. In:
636 Hartwig WC, editor. *The Primate Fossil Record*. Cambridge: Cambridge University Press.
637 p151-160.

638 Rosenberger AL. 2010. Platyrrhines, PAUP, parallelism, and the Long Lineage Hypothesis:
639 A reply to Kay et al. (2008). *Journal of Human Evolution* 59:214-217.

640 Rosenberger AL, Strier KB. 1989. Adaptive radiation of the ateline primates. *Journal of*
641 *Human Evolution* 18:717-750.

642 Rosenberger AL, Tejedor MF. 2013. The misbegotten: long lineages, long branches and the
643 interrelationships of *Aotus*, *Callicebus* and the saki-uakaris. In: Barnett AL, Veiga L, Ferrari
644 S, Norconk M, editors. *Evolutionary Biology and Conservation of Titis, Sakis and Uacaris*.
645 Cambridge: Cambridge University Press. p 13-22.

646 Rosenberger AL, Norconk MA, Garber PA. 1996. New perspectives on the Pitheciines. In:
647 Norconk MA, Rosenberger AL, and Garber PA, editors. *Adaptive Radiations of Neotropical*
648 *Primates*. New York: Plenum Press. p329-333.

649 Rosenberger AL, Tejedor MF, Cooke S, Halenar L, Pekkar S. 2009. Platyrrhine
650 ecophylogenetics, past and present. In: Garber PA, Estrada A, Bicca-Marques JC, Heymann
651 EW, Strier KB, editors. *South American Primates Comparative Perspectives in the Study of*
652 *Behaviour, Ecology and Conservation*. New York: Springer. p69-113.

- 653 Saitou N, Nei M. 1987. The neighbor-joining method: a new method for reconstructing
654 phylogenetic trees. *Molecular Biology and Evolution* 4:406-425.
- 655 Schneider H, Sampaio I. 2015. The systematics and evolution of New World primates - A
656 review. *Molecular Phylogenetics and Evolution* 82:348-357.
- 657 Scotland RW, Olmstead RG, Bennet JR. 2003. Phylogeny reconstruction: The role of
658 morphology. *Systematic Biology* 52:539-548.
- 659 Scott JE. 2015. Lost and found: the third molars of *Callimico goeldii* and the evolution of the
660 callitrichine postcanine dentition. *Journal of Human Evolution* 83:65-73.
- 661 Smith RJ, Jungers WL. 1997. Body mass in comparative primatology. *Journal of Human*
662 *Evolution* 32:523-559.
- 663 Spoor F, Gunz P, Neubauer S, Stelzer S, Scott N, Kwekason A, Dean MC. 2015.
664 Reconstructed *Homo habilis* type OH 7 suggests deep-rooted species diversity in early
665 *Homo*. *Nature* 519:83-86.
- 666 von Cramon-Taubadel N. 2014. The microevolution of modern human cranial variation:
667 Implications for human and primate evolution. *Annals of Human Biology* 41:323-335.
- 668 Whelan S, Lio P, Goldman N. 2001. Molecular phylogenetics: state-of-the-art methods for
669 looking into the past. *Trends in Genetics* 17:262-272.
- 670 Wiens JJ. 2004. The role of morphological data in phylogeny reconstruction. *Systematic*
671 *Biology* 53:653-661.
- 672 Wildman DE, Jameson NM, Opazo JC, Yi SV. 2009. A fully resolved genus level phylogeny
673 of Neotropical primates (Platyrrhini). *Molecular Phylogenetics and Evolution* 53:694-702.

- 674 Wilkinson RD, Steiper ME, Soligo C, Martin RD, Yang Z, Tavaré S. 2011. Dating primate
675 divergences through an integrated analysis of palaeontological and molecular data.
676 *Systematic Biology* 60:16-31.
- 677 Wood BA, Lieberman DE. 2001. Craniodental variation in *Paranthropus boisei*: A
678 developmental and functional perspective. *American Journal of Physical Anthropology*
679 116:13-25.
- 680 Yang Z. 2006. *Computational Molecular Evolution*. Oxford: Oxford University Press. 376 p.
- 681 Yang Z, Rannala B. 2012. Molecular phylogenetics: principles and practice. *Nature Reviews*
682 *Genetics* 13:303-314.
- 683 Zelditch ML, Swiderski DL, Sheets HD, Fink WL. 2004. *Geometric Morphometrics For*
684 *Biologists: A Primer*. London: Elsevier Academic Press. 416 p.
- 685 Zollikofer CP, de León MS, Lieberman DE, Guy F, Pilbeam D, Likies A, Mackaye HT,
686 Vignaud P, Brunet M. 2005. Virtual cranial reconstruction of *Sahelanthropus tchadensis*.
687 *Nature* 434:755-759.
- 688

689 **Figure Legends**

690 Figure 1 Platyrrhine genus-level molecular phylogenetic relationships

691

692 Figure 2 Consensus genus-level phylogenetic relationships inferred from pitheciid analyses

693 without *Aotus*. (a) Face, and the face and cranial vault with *Lagothrix* as outgroup, the cranial694 base and palate for both *Callimico* and *Saimiri* outgroups, and the cranium, face, face and695 cranial vault, cranial base, cranial base and vault for *Saimiri* as outgroup. (b) Molecular

696 phylogeny for the face and palate with all three outgroups, from the cranium, and cranial base

697 for both *Lagothrix* and *Callimico* outgroups, for the cranial base and palate with *Lagothrix* as698 outgroup, and the face, face and cranial vault, and cranial base and vault for *Callimico* as699 outgroup. (c) Cranial base and vault data with *Lagothrix* outgroup.


700

701 Figure 3 Consensus genus-level phylogenetic relationships inferred from Procrustes shape for

702 pitheciid and *Aotus* analyses. (a) Face and cranial vault with *Callimico* outgroup, and cranial703 base and palate, and face and palate for *Saimiri* outgroup. (b) Cranial base for all three704 outgroups, cranium, face, and face and cranial vault for *Lagothrix* and *Saimiri* outgroups,705 face and palate for *Lagothrix* outgroup, and cranial base and vault for *Saimiri* outgroup. (c)706 Face, and face and palate for *Callimico*, and cranial base and palate for *Lagothrix* outgroup.707 (d) Cranial base and vault with *Lagothrix* outgroup. (e) Cranium, cranial base and palate, and708 cranial base and vault for *Callimico* outgroup, and congruent with the molecular phylogeny.

709


710 Figure 1 Platyrrhine genus-level molecular phylogenetic relationships



711

712


713 Figure 2 Consensus genus-level phylogenetic relationships inferred from pitheciid analyses
 714 without *Aotus*. (a) Face, and the face and cranial vault with *Lagothrix* as outgroup, the cranial
 715 base and palate for both *Callimico* and *Saimiri* outgroups, and the cranium, face, face and
 716 cranial vault, cranial base, cranial base and vault for *Saimiri* as outgroup. (b) Molecular
 717 phylogeny for the face and palate with all three outgroups, from the cranium, and cranial base
 718 for both *Lagothrix* and *Callimico* outgroups, for the cranial base and palate with *Lagothrix* as
 719 outgroup, and the face, face and cranial vault, and cranial base and vault for *Callimico* as
 720 outgroup. (c) Cranial base and vault data with *Lagothrix* outgroup.



721

722

723 Figure 3 Consensus genus-level phylogenetic relationships inferred from Procrustes shape for
 724 pitheciid and *Aotus* analyses. (a) Face and cranial vault with *Callimico* outgroup, and cranial
 725 base and palate, and face and palate for *Saimiri* outgroup. (b) Cranial base for all three
 726 outgroups, cranium, face, and face and cranial vault for *Lagothrix* and *Saimiri* outgroups,
 727 face and palate for *Lagothrix* outgroup, and cranial base and vault for *Saimiri* outgroup. (c)
 728 Face, and face and palate for *Callimico*, and cranial base and palate for *Lagothrix* outgroup.
 729 (d) Cranial base and vault with *Lagothrix* outgroup. (e) Cranium, cranial base and palate, and
 730 cranial base and vault for *Callimico* outgroup, and congruent with the molecular phylogeny.



731

732

733 **Table I list of cranial anatomical landmarks**

734

1. Piriform aperture nasospinale
2. Piriform aperture point of greatest width
3. Piriform aperture meeting of nasal and maxilla
4. Piriform aperture rhinion, most anterior midline
5. Nasion suture meeting of fronto nasals
6. Glabella midline point on frontal between supraorbital ridges
7. Supraorbital superior
8. Frontomolare orbitale
9. Frontomolare temporal
10. Zygo-max superior
11. Zygo-max inferior
12. Zygomatic foramen inferior
13. Infraorbital foramen inferior
14. Lacrimal duct fossa bottom
15. Optic foramen most medial
16. Upper posterior maxilla
17. Maximum point of curvature on upper zygomatic
18. Zygo-temp superior
19. Zygo-temp inferior
20. Meeting point of sphenoid and zygomatic
21. Meeting point of sphenoid, parietal and zygomatic process of temporal
22. Midpoint between glabella and bregma

23. Bregma
24. Midpoint between bregma and lambda
25. Lambda
26. Asterion
27. Auditory meatus anterior
28. Auditory meatus posterior
29. Auditory meatus inferior
30. Incisor I1 septum
31. Canine septum
32. Premolar P2 septum
33. Molar M1 septum
34. Midpoint of septum at end of dentition
35. Incisive foramen posterior
36. Meeting point of maxilla and palatine
37. Palatine foramen posterior/lateral
38. Max curvature of posterior edge of palatine
39. Nasal spine midpoint where wings split
40. Midpoint between basisphenoid and basioccipital
41. Petrous apex meeting point of petrous, basiosphenoid and basioccipital
42. Foramen larelli
43. Meeting point of petrous, sphenoid and zygomatic process of temporal
44. Petrous greatest central projection
45. Stylomastoid foramen
46. Jugular foramen distal

47. Jugular foramen medial
48. Carotid foramen anterior
49. Midpoint between basion and basisphen-basioccipital
50. Basion anterior
51. Occipital condyle anterior apex
52. Occipital condyle posterior midpoint
53. Hypoglossal canal
54. Opisthion posterior
55. Midway between opisthion and inion
56. Inion
57. Greatest curvature on posterior zygomatic process of temporal
58. Temporal meeting point between sphenoid and zygomatic process of
59. Tip of post glenoid process
60. Deepest point within mandibular fossa
61. Articular eminence medial
62. Articular eminence midpoint
63. Articular eminence lateral

735

736

737 **Table II Pitheciid and outgroup taxa sample sizes for phylogenetic analyses**

Taxa	Sample size		
	Female	Male	Pooled
<i>Aotus azarae</i>	10	6	16
<i>Aotus lemurinus</i>	10	10	26
<i>Aotus vociferans</i>	10	10	20
<i>Aotus trivirgatus</i>	11	13	24
<i>Callicebus cupreus</i>	9	10	19
<i>Callicebus hoffmannsi</i>	10	9	19
<i>Callicebus moloch</i>	15	13	28
<i>Callicebus torquatus</i>	9	12	21
<i>Cacajao calvus</i>	10	13	23
<i>Cacajao melanocephalus</i>	17	13	30
<i>Chiropotes satanas</i>	9	14	23
<i>Pithecia pithecia</i>	10	12	22
<i>Pithecia monachus</i>	13	14	27
Outgroups			
<i>Callimico goeldii</i>	11	11	22
<i>Lagothrix lagotricha</i>	10	10	20
<i>Saimiri sciureus</i>	33	15	48

738

739

740 **Table III Test of phylogenetic signal as measured by tree length (total amount of shape change across all phylogenetic branches) and**
 741 **statistical significance (comparing tree length for original data against permutation with random swapping of values) for Procrustes**
 742 **coordinates and log centroid size of pitheciids without and with *Aotus***
 743

	Pitheciid without <i>Aotus</i>				Pitheciid with <i>Aotus</i>			
	Procrustes coordinates		Log centroid size		Procrustes coordinates		Log centroid size	
	Tree length	<i>P</i>	Tree length	<i>P</i>	Tree length	<i>P</i>	Tree length	<i>P</i>
Cranial base	0.0130	<0.0001	0.0337	<0.001	0.0190	<0.0001	0.0413	<0.0001
Cranial base & palate	0.0079	<0.001	0.0367	<0.001	0.0101	<0.0001	0.0450	<0.0001
Cranial base & vault	0.0107	<0.0001	0.0412	<0.001	0.0156	<0.0001	0.0489	<0.0001
Cranium	0.0102	<0.0001	0.0351	<0.001	0.0153	<0.0001	0.0408	<0.0001
Face	0.0244	<0.001	0.0225	<0.001	0.0343	<0.0001	0.0229	<0.0001
Face & cranial vault	0.0137	<0.0001	0.0339	<0.001	0.0204	<0.0001	0.0364	<0.0001
Face & palate	0.0253	<0.0001	0.0316	<0.001	0.0363	<0.0001	0.0322	<0.0001

744

745

746 **Table IV Jack-knife clade support for phylogenetic analysis of Procrustes shape of pitheciids.**

747

Cranial region	Cranium			Face			Face & palate			Face & cranial vault			Cranial base			Cranial base & vault			Cranial base & palate		
	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>
Outgroup																					
Molecular clades																					
<i>Cacajao</i>	100	100	100	100	86.6	100	100	95.8	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Callicebus</i>	100	100	100	93.3	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Pithecia</i>	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Cacajao-Chiropotes</i>	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	87.5	100	100	100
<i>Cacajao-Chiropotes</i> <i>-Pithecia</i>	100	92	<20	100	<20	<20	100	79.2	100	100	<20	<20	<20	100	<20	100	<20	<20	38.2	100	<20
Non-molecular clades																					
<i>Pithecia-Callicebus</i>	<20	<20	100	<20	86.6	100	<20	20.8	<20	<20	80.8	100	100	<20	100	<20	<20	100	61.8	<20	100
<i>Cacajao-Chiropotes</i> <i>-Callicebus</i>	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	19.2	<20	<20	<20	<20	<20	87.5	<20	<20	<20	<20

748

749

750 **Table V Jack-knife clade support for phylogenetic analysis of Procrustes shape of pitheciids and *Aotus*.**

Cranial region	Cranium			Face			Face & cranial vault			Face & palate			Cranial base			Cranial base & vault			Cranial base & palate		
	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>	<i>Callimico</i>	<i>Lagothrix</i>	<i>Saimiri</i>
Molecular clades																					
<i>Aotus</i>	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Cacajao</i>	100	100	100	100	93	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Callicebus</i>	100	100	100	100	100	100	100	96	100	100	100	100	100	100	100	100	100	100	100	82	100
<i>Pithecia</i>	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Cacajao-Chiropotes</i>	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100	100
<i>Cacajao-Chiropotes-Pithecia</i>	100	<20	<20	<20	<20	<20	100	46	100	100	<20	<20	<20	<20	<20	100	<20	<20	100	94	100
<i>Cacajao-Chiropotes-Pithecia-Callicebus</i>	100	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	<20	100	<20	<20	100	<20	<20
Non-molecular clades																					
<i>Aotus-Callicebus</i>	<20	100	100	<20	100	100	42	100	100	100	100	100	96	100	100	<20	100	100	<20	<20	100
<i>Aotus-Callicebus-Pithecia</i>	<20	97	100	<20	93	100	<20	54	<20	<20	100	100	88	100	100	<20	<20	100	<20	<20	<20
<i>Aotus-Cacajao-Chiropotes-Pithecia</i>	<20	<20	<20	100	<20	<20	54	<20	<20	<20	<20	<20	<20	<20	<20	<20	97	<20	<20	88	<20

751

752