## Distinguishing mesoscopic quantum superpositions from statistical mixtures in periodically shaken double wells

## **Christoph Weiss**

Institut für Physik, Carl von Ossietzky Universität, D-26111 Oldenburg, Germany Department of Physics, Durham University, Durham DH1 3LE, United Kingdom

Abstract. For Bose-Einstein condensates in double wells, N-particle Rabi-like oscillations often seem to be damped. Far from being a decoherence effect, the apparent damping can indicate the emergence of quantum superpositions in the many-particle quantum dynamics. However, in an experiment it would be difficult to distinguish the apparent damping from decoherence effects. The present paper suggests using controlled periodic shaking to quasi-instantaneously switch the sign of an effective Hamiltonian, thus implementing an "echo" technique which distinguishes quantum superpositions from statistical mixtures. The scheme for the effective time-reversal is tested by numerically solving the time-dependent N-particle Schrödinger equation.

PACS numbers: 03.75.Lm,67.85.Hj,03.75.-b

Small Bose-Einstein condensates (BECs) of some 1000 [1] or even 100 atoms [2] have been a topic of experimental research for several years. Recently, the investigation of many-particle wave-functions of BECs in phase space became experimentally feasible [3]. This experimental technique will lead to further investigations of beyond-mean-field (Gross-Pitaevskii) behaviour for small BECs.

For a BEC initially loaded into one of the wells of a double-well potential, the manyparticle oscillations often seem to be damped compared to the mean-field behaviour. Figure 1 shows such an apparent damping, which in fact is a collapse which will eventually be followed by at least partial revivals [cf. Refs. [4, 5], Fig 1 (c)], for N = 100particles. This apparent damping coincides with an increase of the fluctuations of the number of particles in each well [Fig. 1 (b)].

In order to numerically calculate the many-particle dynamics, the Hamiltonian in the two-mode approximation [6] is used,

$$\hat{H}_0 = -J\left(\hat{c}_1^{\dagger}\hat{c}_2 + \hat{c}_2^{\dagger}\hat{c}_1\right) + \frac{U}{2}\sum_{j=1}^2 \hat{n}_j \left(\hat{n}_j - 1\right) , \qquad (1)$$

where  $\hat{c}_{j}^{(\dagger)}$  are the boson creation and annihilation operators on site j,  $\hat{n}_{j} = \hat{c}_{j}^{\dagger}\hat{c}_{j}$  are the number operators, J is the hopping matrix element and U the on-site interaction energy.

The experimentally measurable [7] population imbalance is useful to quantify the oscillations depicted in Fig. 1:

$$\frac{\langle z \rangle(\tau)}{2} \equiv \frac{\langle n_2 \rangle(\tau) - \langle n_1 \rangle(\tau)}{2N} , \qquad (2)$$

where  $\tau$  is the dimensionless time:

$$\tau \equiv \frac{tJ}{\hbar} \,. \tag{3}$$

The variance of the population imbalance can be quantified by using the experimentally measurable [7] quantity

$$F_{z} \equiv \frac{\langle (\hat{n}_{1} - \hat{n}_{2})^{2} \rangle - \langle \hat{n}_{1} - \hat{n}_{2} \rangle^{2}}{N} , \qquad (4)$$

with  $0 \leq F_z \leq N$ . If all atoms are in the same single-particle state, this can be expressed by using the atomic coherent states [8],

$$|\theta, \phi\rangle_N = \sum_{n=0}^N \binom{N}{n}^{1/2} \cos^n(\theta/2) \sin^{N-n}(\theta/2) \\ \times e^{i(N-n)\phi} |n, N-n\rangle .$$
 (5)

For the product wave-functions defined by Eq. (5),  $F_z$  can be calculated using properties of the classical binomial distribution, with probabilities

$$P_{n_1} = \binom{N}{n_1} p^{n_1} (1-p)^{N-n_1}, \quad p = \cos^2(\theta/2).$$



Figure 1. (Colour online) a) Population imbalance  $\langle z \rangle/2$  [Eq. (2)] as a function of dimensionless time  $\tau$  (3) for small BEC in a double-well potential. Initially, N = 100 atoms are in well 1, the quantum dynamics is given by the Hamiltonian (1) (NU/J = 0.4). b) Variance of the population imbalance (4). c) Same curve as in panel (a) but for longer times. The time-evolution appears to be damped up to  $\tau \approx 200$  and is then followed by a partial revival for times above  $\tau \approx 300$ . Experimentally, it will be difficult to distinguish the apparent damping and the increased fluctuations (which are both triggered by a collapse and revival phenomenon) from true damping introduced, e.g., by decoherence.

The result reads:

$$F_z = 4 \left[ \cos^2(\theta/2) - \cos^4(\theta/2) \right]$$
  
=  $\sin^2(\theta)$ , (6)

and hence  $0 \leq F_z \leq 1$ . Thus, while  $F_z < 1$  would not be sufficient to distinguish product states, as in Eq. (5), from "spin-squeezed states" [7], for which  $F_z < 1$  is also true, a pure state with  $F_z > 1$  has to be in a quantum superposition. For numeric solutions of the Schrödinger equation corresponding to Hamiltonians like the one given in Eq. (1), one always knows that the system is in a pure state and thus that any state with  $F_z > 1$  is a quantum superposition. In an experiment, the situation would be more complicated. The focus of the present paper lies on providing a way to distinguish quantum superpositions with  $F_z > 1$  from statistical mixtures with  $F_z > 1$  via an 'echo' technique.

For pure states, Eq. (4) coincides with a quantum Fisher information [9]. Like the spin-squeezed states investigated in Ref. [7] (and references therein), quantum superpositions with large fluctuations are also relevant to improve interferometric measurements beyond single-particle limits. A prominent example of a quantum superposition relevant for interferometry are the NOON-states [10]

$$|\psi_{\text{NOON}}\rangle = \frac{1}{\sqrt{2}} \left(|N,0\rangle + |0,N\rangle\right) , \qquad (7)$$

i.e., quantum superpositions of all particles either being in well one or in well two;  $|n_1, n_2\rangle$  refers to the Fock state with  $n_1$  particles in well 1 and  $n_2$  particles in well 2. Suggestions how such states can be obtained for ultra-cold atoms can be found in Refs. [5, 11, 12, 13, 14, 15, 16, 17] and references therein. For pure states,  $F_z > 1$ indicates that this quantum superposition is relevant for interferometry [9]. However, it remains to be shown that the increased fluctuations are really due to pure states rather than statistical mixtures.

It might sound tempting to use the revivals investigated in Refs. [4, 5] to identify pure quantum states. However, while such revivals can be observed, e.g., for two-particle systems [18], the situation for a BEC in a double well is more complicated. In principle, very good revivals of the initial wave-function should occur as long as the system is described by the Hamiltonian (1). While partial revivals [cf. Fig 1 (c)] can easily be observed, (nearly) perfect revivals might occur for times well beyond experimental timescales – in particular if the experiment is performed under realistic conditions subject to decoherence effects ‡. It is thus not obvious how such an apparent damping might be distinguished experimentally from decoherence effects which would lead to statistical mixtures with (now truly) damped oscillation similar to Fig. 1. The focus of this paper thus lies on an experimentally realisable "echo" technique to distinguish statistical mixtures from quantum superpositions by using periodic shaking.

Periodic shaking [19] is currently being established experimentally to control tunnelling of BECs [20, 21, 22, 23, 24, 25]. For the model (1), periodic shaking can be included via

$$\hat{H} = \hat{H}_0 + \frac{K}{2}\cos(\omega t)(\hat{n}_2 - \hat{n}_1) , \qquad (8)$$

where K is the strength of shaking and  $\omega$  its (angular) frequency. For large shaking frequencies § and not-too-large interactions, the time-dependent Hamiltonian (8) can

<sup>‡</sup> For computer simulations, numerical errors might produce an effective decoherence which would again prevent nearly perfect revivals from occurring at very long time-scales.

<sup>§</sup> While the validity of this approximation also depends on the values chosen for the interaction, driving frequencies as low as  $\hbar\omega \approx 6J$  can sometimes be considered large. Choosing higher frequencies will



Figure 2. (Colour online) a) Sketch of a double well which is shaken periodically to control tunnelling for ultra-cold atoms. b) For high shaking frequencies, the tunnelling rate is modified by the  $\mathcal{J}_0$ -Bessel function [see Eq. (10)]. The two squares indicate a pair of shaking amplitudes for which the Bessel function has equal modulus and opposite sign  $(x_1 \simeq 1.69 \text{ and } x_2 \simeq 3.83 \text{ with } |J_0(x_{1,2})| \simeq 0.403 - \text{it will be shown in Fig. 5 (d)}$  that it is not essential to determine the values of  $x_1$  and  $x_2$  with very high accuracy).

be replaced by a time-independent effective Hamiltonian:

$$\hat{H}_{\text{eff}} = -J_{\text{eff}} \left( \hat{c}_1^{\dagger} \hat{c}_2 + \hat{c}_2^{\dagger} \hat{c}_1 \right) + \frac{U}{2} \sum_{j=1}^2 \hat{n}_j \left( \hat{n}_j - 1 \right)$$
(9)

with

$$J_{\text{eff}} = J \mathcal{J}_0 \left( K_0 \right); \quad K_0 \equiv \frac{K}{\hbar \omega}$$
(10)

where  $\mathcal{J}_0$  is the Bessel-function depicted in Fig. 2 (b). Such effective Hamiltonians have been successfully tested experimentally in optical lattices, see, e.g., Refs. [20, 26]; negative  $J_{\text{eff}}$  have been experimentally investigated in Refs. [22, 25]. While numerically it is much easier to investigate the two-site Hamiltonian (9) or its time-dependent counterpart (8), experimentally it might be preferable to realise such models in optical lattices for parameters in which the one-band approximation is good [20, 21]. There are, however, also examples [27, 28] for which two or more Bessel function are needed to understand the tunnelling dynamics.

In the present situation, the effective description (9) offers the possibility to quasiinstantaneously switch the sign of both the kinetic energy (via shaking, cf. Fig. 2) and the interaction (via a Feshbach-resonance [29]). Contrary to special cases where the wave-function [30, 31] can be changed to obtain time-reversal, for periodically driven systems the *Hamiltonian* can be changed by quasi-instantaneously changing both the tunnelling term [by switching the shaking amplitude, e.g., between values shown in Fig. 2 (b)] and the sign of the interaction via a Feshbach-resonance [29];

$$\hat{H}_{\text{ideal}} \equiv \begin{cases} +H_{\text{eff}}(\tau=0) &: \tau < \tau_0 \\ -\hat{H}_{\text{eff}}(\tau=0) &: \tau \ge \tau_0 \end{cases}$$
(11)

improve the approximation. However, as this will, in general, also increase the driving amplitude, for too high frequencies the two-mode approximation (1) no longer is valid.



Figure 3. (Colour online) Two-dimensional projection of the modulus-squared of the scalar product of the wave-function with the product states (5) as a function of both  $\phi$  and  $z = \cos(\theta)$ . a) Parameters as in Fig. 1 except for  $K_0 \simeq 1.6917$  and  $\hbar\omega = 32J$ . The wave-function after the apparent damping is displayed for  $\tau \simeq 42.41$ . This quantum superposition can be characterised by  $F_z \simeq 38.8$  and it could thus be used to improve interferometric measurements. b) If all 100 atoms occupy the same single-particle state (here:  $z = 0, \phi = \pi/2$ ), the wave-function is much narrower [see Eq. (6)]. Furthermore, no product state would be interesting for quantum enhanced interferometry.

The corresponding unitary time-evolution is given by

$$U(0,\tau) = \begin{cases} \exp\left(-\frac{\mathrm{i}\tau\hat{H}_{\mathrm{eff}}(\tau=0)}{\hbar J}\right) & : \quad \tau < \tau_0 \\ \exp\left(\frac{\mathrm{i}(\tau-2\tau_0)\hat{H}_{\mathrm{eff}}(\tau=0)}{\hbar J}\right) & : \quad \tau \ge \tau_0 \end{cases},$$
(12)

with perfect return to the initial state at  $\tau = 2\tau_0$ . However, the turning point  $\tau_0$  has to be chosen with care: only by taking  $\tau_0$  close to the maximum of the shaking can unwanted excitations be excluded (cf. Refs. [32, 33, 34]). Recent related investigations of the influence of the initial phase of the driving [replacing  $\cos(\omega t)$  in the Hamiltonian (8) by  $\cos(\omega t + \phi)$ ] can be found in Refs. [35, 36, 37].

In the following, the time-reversal is demonstrated by numerically solving the full, time-dependent Hamiltonian (8) corresponding to the ideal time-reversal Hamiltonian (11) using the Shampine-Gordon routine [38]. Contrary to time-reversal schemes on the level of the Gross-Pitaevskii equation [39, 40], here time-reversal is used to distinguish interesting quantum superpositions from statistical mixtures. Before implementing the time-reversal, Fig. 3 shows the wave-function for N = 100 particles which were initially in one well. After several oscillations, the wave-function no longer is in a product state. Both the population imbalance and the phase can be measured experimentally [7]; in Fig. 3 the squared modulus of the scalar product with the atomic Distinguishing mesoscopic quantum superpositions from mixtures

coherent states (5) is plotted. The angle  $\theta$  corresponds to a population imbalance of

$$\frac{\langle z \rangle}{2} = \frac{\cos(\theta)}{2} \tag{13}$$

Ideally, it should be possible to show that the wave-function of Fig. 3 (a) indeed is a quantum superposition by using the time-reversal of Eq. (11) and then investigating

$$\langle z_{\rm end} \rangle \equiv \langle z \rangle (2\tau_0) \,.$$
 (14)

Firstly, there is only one many-particle wave-function which fulfils

$$\langle z_{\rm end} \rangle = 1. \tag{15}$$

Secondly, the unitary evolution of solutions of the Schrödinger equation guarantees that for two different solutions  $|\psi_1(2\tau_0)\rangle = U(\tau_0, 2\tau_0)|\psi_1(\tau_0)\rangle$  and  $|\psi_2(2\tau_0)\rangle = U(\tau_0, 2\tau_0)|\psi_2(\tau_0)\rangle$ , the scalar product would be the same at  $\tau = \tau_0$  and at  $\tau = 2\tau_0$ (as  $U^{\dagger}U = 1$ ).

However, the Hamiltonian (11) is a high-frequency approximation and it has thus to be shown that this works for realistic driving frequencies (cf. Fig. 4). Furthermore, although there is only one wave-function at  $\tau = \tau_0$  which exactly leads to the value  $\langle z \rangle_{\text{end}} = 1$  at  $\tau = 2\tau_0$ , other (less interesting) wave-functions might lead to values close to  $\langle z \rangle_{\text{end}} = 1$  (cf. Fig. 5).

Figure 4 shows that the time-reversal dynamics is indeed feasible. On time-scales for which there is not even a partial revival [cf. Fig 1 (c)] of the initial state characterised by  $\langle z \rangle = 1$ , the proposed time-reversal dynamics leads to final values above  $\langle z_{end} \rangle/2 = 0.45$ (Fig. 5 shows that this is enough to show that the wave-function at  $\tau = \tau_0$  was indeed a quantum superposition). In order to show that the scheme does not rely on the switching to be truly instantaneous at  $t = t_0$  [where  $t_0$  is linked to  $\tau_0$  via Eq. (3)], the amplitude in Fig 4 (d) was switched according to

$$K_0(t) = K_0^{(1)} + \left(K_0^{(2)} - K_0^{(1)}\right) \frac{1 + \tanh\left(\frac{\omega(t-t_0)}{\gamma}\right)}{2};$$
(16)

the switching between the two interaction values was chosen analogously (for instantaneous switching, the switching time can also slightly deviate from the ideal switching time, it just has to be close to the shaking maximum).

Figure 4 shows that the time-reversal dynamics is feasible. If one implements the time-reversal dynamics without decoherence, one will get a reasonably good return to the initial state. However, does this imply the statement that if at  $\tau = 2\tau$  the system has approximately returned to the initial state, it automatically has been in a quantum superposition at  $\tau = \tau_0$ ? In order to answer that question, Figure 5 investigates the dynamics of states which are in a product state (5) at  $\tau = \tau_0$ :

$$|\psi(\tau_0)\rangle = |\theta_0, \phi_0\rangle_N, \qquad (17)$$

and are subject to the same shaking as Fig. 4 (b). Figure 5 (a) shows that

$$\langle z_{\max} \rangle \equiv \max \left\{ \langle z \rangle(\tau) \right\} |_{1.99\tau_0 \le \tau \le 2\tau_0}$$
(18)



Figure 4. (Colour online) Time-reversal of the quantum dynamics of a small Bose-Einstein condensate in a periodically shaken double well [Eq. (8)]. a) Population imbalance (2) as a function of time  $\tau = tJ/\hbar$  for the same parameters as in Fig. 3 (a). b) Red/dark solid line: all other parameters as in panel (a) except for  $\tau > \tau_0 = 13.5\pi \simeq 42.41$ :  $K_0 = 3.8317 J$  and U = -0.4J/N; the revival of the initial state is visible near  $\tau \approx 85$ . c) Population imbalance for the same situation as in panel (a) but for much longer time-scales. d) If the switching takes place continuously rather than instantaneously [Eq. (16)], the revival of the initial state can still be observed [same parameters as for panel (b)] ( $\gamma = 0$  corresponds to instantaneous switching; in the limit  $\hbar\omega/J \to \infty$  the maximum of this curve would be at  $\gamma = 0$ ).

[shown as a two-dimensional projection as a function of both  $\cos(\theta_0)$  and  $\phi_0$ ] lies well below the values achieved in time-reversal [Fig. 4 (b)]. Furthermore, it does not change dramatically on short time-scales, as can be seen in Fig. 5 (b) which uses a  $\langle z_{\min} \rangle$  which is analogously defined to Eq. (18) to calculate:

$$\Delta z \equiv \frac{\langle z_{\rm max} \rangle - \langle z_{\rm min} \rangle}{2}, \qquad (19)$$

which indicates how accurately  $\langle z \rangle_{\text{max}}$  can be determined. In addition to not approaching  $\langle z \rangle = 1$ , many product states lead to very large fluctuations [Fig. 5 (c)];

0

1

 $\phi_0 / \pi$ 

2



**Figure 5.** (Colour online) **a)** For the Hamiltonian which leads to the curve in Fig. 4 (b), at  $\tau = \tau_0$  product states [Eqs. (5) and (17);  $z_0 = \cos(\theta_0)$ ] are implemented. Displayed is the two-dimensional projection of  $\langle z_{\max} \rangle/2$  as a function of both the initial phase and the initial population imbalance. This lies well below the values obtained in Fig. 4 (b), thus indicating that any wave-function which leads to values of  $\langle z \rangle_{\text{end}} \tau = 2\tau_0$  comparable to Fig. 4 (b) was indeed a quantum superposition at  $\tau = \tau_0$ . **b)** Two-dimensional projection of  $\Delta z$  [Eq. (19)], which indicates how accurately  $\langle z \rangle_{\text{max}}$  can be determined, shows that  $\langle z \rangle$  [as for Fig. 4 (b)] does not change dramatically on short time-scales. Thus, choosing the time of observation is not that crucial. **c)**  $F_z(\tau_0)$  (4) as a two-dimensional projection. Many product states lead to considerably larger fluctuations than obtained for the curve in Fig. 4 (b) (which goes below  $F_z = 1$  near  $\tau = 2\tau_0$ ). **d)** If the time-reversal scheme of Fig. 4 (b) for the red/dark curve is repeated with non-ideal driving amplitudes,  $\langle z_{\text{end}} \rangle/2$  (shown as a two-dimensional projection as a function of both driving amplitudes, normalised by their ideal values - cf. Fig. 2) still lies well above the values shown in panel (a).

0.996

1

 $K_0^{(1)}/ideal$ 

1.004

these fluctuations are particularly large if one compares them with the tiny values of  $F_z(2\tau_0) \simeq 0.4$  for the curve in Fig. 4 (b) ||. Carefully investigating how the product states (5) with large contributions to Fig. 3 (a) behave offers an additional route to distinguish quantum superpositions as in Fig. 3 (a) from statistical mixtures. Figure 5 (d) shows that the time-reversal scheme is feasible even if the driving amplitudes only approximately meet the ideal values [Fig. 5 (d)]. Many of the features displayed Fig. 5 (a)-(c) could be understood by comparing them to the mean-field behaviour including the fact that the fluctuations are large for some parameters and low for others (cf. Ref. [41]). However, the main focus within this paper lies on showing that the return to the initial state displayed in Fig. 5 (a) is much lower than what can be obtained in Fig. 5 (b). In addition, the fluctuations displayed in Fig. 5 (c) are much larger than would be obtained by the time-reversal dynamics displayed in Fig. 5 (b).

To conclude, time-reversal via quasi-instantaneously changing the sign of the effective Hamiltonian is experimentally feasible for ultra-cold atoms in a periodically shaken double well. The change of the sign of the Hamiltonian is achieved by changing both the driving amplitude and the sign of the interaction; a particularly useful initial state is the state with all particles in one well. The numeric investigations show that the revival of the initial state can be used to distinguish damping introduced via decoherence from the apparent damping related to a collapse phenomenon. Even if the revival of the initial state is not perfect, the scheme clearly distinguishes product states from quantum superpositions with potential interferometric applications. While the present paper focuses on on an experimentally relevant example for which the time-reversal dynamics can distinguish intermediate quantum superpositions from statistical mixtures, a perfect return to the initial state would prove that for all initial states. For general cases [which might include general product states (5)], more precise measurements [3] of both the initial and the final state including both population imbalance and fluctuations as well as measuring the phase distribution will be necessary.

## Acknowledgements

I would like to thank S.A. Gardiner and M. Holthaus for their support and T.P. Billam, B. Gertjerenken, E. Haller, C. Hoffmann, J. Hoppenau, A. Ridinger, T. Sternke and S. Trotzky for discussions.

## References

- M. Albiez, R. Gati, J. Folling, S. Hunsmann, M. Cristiani, and M. K. Oberthaler. *Phys. Rev. Lett.*, 95:010402, 2005.
- [2] C.-S. Chuu, F. Schreck, T. P. Meyrath, J. L. Hanssen, G. N. Price, and M. G. Raizen. Phys. Rev. Lett., 95:260403, 2005.
- [3] T. Zibold, E. Nicklas, C. Gross, and M. K. Oberthaler. Phys. Rev. Lett., 105:204101, 2010.
- [4] M. Holthaus and S. Stenholm. Eur. Phys. J. B, 20:451, 2001.
- || In the limit  $\hbar \omega/J \to \infty$ , one would even have  $F_z(2\tau_0) = 0$ .

- [5] K. Ziegler. J. Phys. B, 44:145302, 2011.
- [6] G. J. Milburn, J. Corney, E. M. Wright, and D. F. Walls. Phys. Rev. A, 55:4318, 1997.
- [7] J. Esteve, C. Gross, A. Weller, S. Giovanazzi, and M. K. Oberthaler. Nature (London), 455:1216, 2008.
- [8] L. Mandel and E. Wolf. Cambridge University Press, Cambridge, 1995.
- [9] L. Pezzé and A. Smerzi. Phys. Rev. Lett., 102:100401, 2009.
- [10] V. Giovannetti, S. Lloyd, and L. Maccone. Science, 306:1330, 2004.
- [11] A. Micheli, D. Jaksch, J. I. Cirac, and P. Zoller. Phys. Rev. A, 67:013607, 2003.
- [12] K. W. Mahmud, H. Perry, and W. P. Reinhardt. Phys. Rev. A, 71:023615, 2005.
- [13] A. I. Streltsov, O. E. Alon, and L. S. Cederbaum. J. Phys. B, 42:091004, 2009.
- [14] D. Dagnino, N. Barberan, M. Lewenstein, and J. Dalibard. Nature Phys., 5:431, 2009.
- [15] B. Gertjerenken, S. Arlinghaus, N. Teichmann, and C. Weiss. Phys. Rev. A, 82:023620, 2010.
- [16] M. A. García-March, D. R. Dounas-Frazer, and L. D. Carr. Phys. Rev. A, 83:043612, 2011.
- [17] G. Mazzarella, L. Salasnich, A. Parola, and F. Toigo. Phys. Rev. A, 83:053607, 2011.
- [18] S. Folling, S. Trotzky, P. Cheinet, M. Feld, R. Saers, A. Widera, T. Muller, and I. Bloch. Nature, 448:1029, 2007.
- [19] M. Grifoni and P. Hänggi. Phys. Rep., 304:229, 1998.
- [20] C. Sias, H. Lignier, Y. P. Singh, A. Zenesini, D. Ciampini, O. Morsch, and E. Arimondo. Phys. Rev. Lett., 100:040404, 2008.
- [21] E. Haller, R. Hart, M. J. Mark, J. G. Danzl, L. Reichsöllner, and H.-C. Nägerl. Phys. Rev. Lett., 104:200403, 2010.
- [22] J. Struck, C. Ölschläger, R. Le Targat, P. Soltan-Panahi, A. Eckardt, M. Lewenstein, P. Windpassinger, and K. Sengstock. *Science*, 2011.
- [23] Y.-A. Chen, S. Nascimbène, M. Aidelsburger, M. Atala, S. Trotzky, and I. Bloch. Phys. Rev. Lett., 107:210405, 2011.
- [24] R. Ma, M. E. Tai, P. M. Preiss, W. S. Bakr, J. Simon, and M. Greiner. Phys. Rev. Lett., 107:095301, 2011.
- [25] D. Ciampini, O. Morsch, and E. Arimondo. J. Phys.: Conf. Ser., 306:012031, 2011.
- [26] C. E. Creffield, F. Sols, D. Ciampini, O. Morsch, and E. Arimondo. Phys. Rev. A, 82:035601, 2010.
- [27] N. Teichmann, M. Esmann, and C. Weiss. Phys. Rev. A, 79:063620, 2009.
- [28] M. Esmann, J. D. Pritchard, and C. Weiss. Laser Phys. Lett. doi: 10.1002/lapl.201110109, ArXiv:1109.2735, 2011.
- [29] D. M. Bauer, M. Lettner, C. Vo, G. Rempe, and S. Durr. Nature Phys., 5:339, 2009.
- [30] G. Morigi, E. Solano, B.-G. Englert, and H. Walther. Phys. Rev. A, 65:040102(R), 2002.
- [31] T. Meunier, S. Gleyzes, P. Maioli, A. Auffeves, G. Nogues, M. Brune, J. M. Raimond, and S. Haroche. *Phys. Rev. Lett.*, 94:010401, 2005.
- [32] A. Ridinger and N. Davidson. Phys. Rev. A, 76:013421, 2007.
- [33] A. Ridinger and C. Weiss. Phys. Rev. A, 79:013414, 2009.
- [34] P. W. Cleary, T. W. Hijmans, and J. T. M. Walraven. Phys. Rev. A, 82:063635, 2010.
- [35] C. E. Creffield and F. Sols. *Phys. Rev. A*, 84:023630, 2011.
- [36] K. Kudo and T. S. Monteiro. Phys. Rev. A, 83:053627, 2011.
- [37] S. Arlinghaus and M. Holthaus. Phys. Rev. B, 84:054301, 2011.
- [38] L. F. Shampine and M. K. Gordon. Freeman, San Francisco, 1975.
- [39] J. Martin, B. Georgeot, and D. L. Shepelyansky. Phys. Rev. Lett., 101:074102, 2008.
- [40] N. Tsukada, H. Yoshida, and T. Suzuki. Phys. Rev. A, 78:015601, 2008.
- [41] C. Weiss and N. Teichmann. Phys. Rev. Lett., 100:140408, 2008.