

Human observers impact habituated samango monkeys' perceived landscape of fear

Katarzyna Nowak^{1,2,3*}, Aliza le Roux², Shane A. Richards⁴, Ciska Scheijen⁵, Russell A. Hill^{1,3}

1. Evolutionary Anthropology Research Group, Department of Anthropology, Durham University, South Road, Durham, DH1 3LE, UK; katarzyna.nowak@durham.ac.uk

2. Department of Zoology and Entomology, University of the Free State, Qwaqwa, Phuthaditjhaba 9866, South Africa

3. Primate & Predator Project, Lajuma Research Centre, P.O. Box 522, Makhado 0920, South Africa

4. Department of Biological and Biomedical Sciences, Durham University, South Road, Durham, DH1 3LE, UK

5. Department of Animal Management, Van Hall Larenstein, 8901 BV Leeuwarden, The Netherlands

*Corresponding author

1 **Human observers impact habituated samango monkeys' perceived landscape of fear**

2 **Short title:** Observing the landscape of fear

3 **Abstract:** Humans and human infrastructure are known to alter the relationship between
4 predators and prey, typically by directly or indirectly shielding one of the species from the other.
5 In addition to these overt changes to animals' behavior, observers may have more subtle impacts
6 on animals' foraging decisions. However, the anthropogenic alteration of risk-taking behavior
7 has rarely been acknowledged or quantified, particularly in behavioral ecological studies reliant
8 on habituated animals. We tested the magnitude of the "human shield effect" experimentally on
9 two groups of samango monkeys, *Cercopithecus mitis erythrarcus*, at a site with high natural
10 predator density and no human hunting pressure. In general, giving up densities (GUDs) – the
11 density of food remaining in a patch when a forager leaves – were greatest at ground level (0.1m)
12 relative to three tree canopy levels (2.5m, 5m and 7.5m), highlighting a strong vertical axis of
13 fear. When human followers were present, however, GUDs were reduced at all four heights;
14 furthermore, for one group, the vertical axis disappeared in the presence of observers. Our results
15 suggest that human observers lower monkeys' perceived risk of terrestrial predators and thereby,
16 affect their foraging decisions at or near ground level. These results have significant implications
17 for future studies of responses to predation risk based on habituation and observational methods.

18 **Lay Summary:** Wild animals make trade-offs between food and safety. Using an experimental
19 approach, we show that habituated arboreal monkeys at a predator-rich site feel safer in the
20 presence of human observers. Specifically, they exploit more food at and near ground level,
21 altering their typical anti-predator responses. Research on risk-sensitive behavior thus needs to
22 account for observers' influence on mammals' landscape of fear, and employ indirect and
23 experimental methods in combination with, or instead of direct observations.

24 INTRODUCTION

25 Many studies of behavioral ecology rely on the habituation process for the collection of
26 detailed observational data on focal species. After repeated and non-threatening contact with
27 humans, we often assume that animals' behavior becomes relatively independent of our presence
28 (Crofoot et al. 2010). However, human observers are known to influence naturalistic interactions,
29 for example between predators and prey by changing prey animals' behavior including routine
30 anti-predator activities, or by affecting the timing and frequency of predation attempts (Isbell and
31 Young 1993; Stanford 1998; Berger 2007; Meshesha 2013). These effects are a manifestation of
32 predators and prey tending to respond differentially to human activity (Ngoprasert, Lynam, and
33 Gale 2007; Rogala et al. 2011). For example, in Gombe National Park, Tanzania, red colobus
34 monkeys (*Procolobus badius tephrosceles*) fled human observers following habituated
35 chimpanzees (*Pan troglodytes schweinfurthii*), which in turn used this "flushing" of colobus by
36 humans to capture the panicked monkeys (Stanford 1998). Isbell and Young (1993) observed a
37 higher rate of predation on vervet monkeys (*Chlorocebus aethiops*) by leopards (*Panthera*
38 *pardus*) in Amboseli National Park, Kenya when observers were away from the field site,
39 suggesting that observers shield habituated monkeys from ground predators. The potential
40 impact of "non-invasive" observers on small-scale anti-predator responses of prey animals (such
41 as alarm calling, or abandoning food to seek shelter) is seldom explicitly considered in
42 behavioral ecological research, even though the "human shield effect" (Berger 2007) can bring
43 about broad-scale changes in the distribution and movement patterns of prey and predator
44 species.

45 Quantitatively assessing subtle changes in animals' behavior due to the presence of
46 human observers can be challenging, particularly in matching conditions with and without

47 observers present. The amount of food that a forager leaves behind in an experimental food
48 patch, the “giving up density” (GUD), provides one technique for measuring animals’ perceived
49 risk without the need for human observers (Brown 1988; Tadesse and Kotler 2012). Animals are
50 predicted to feed more intensively in areas where they feel safe, and relatively lower GUDs
51 indicate preference for a specific area or food patch (Brown 1988; Verdolin 2006). In addition to
52 reflecting trade-offs between resource use and predation risk, GUDs reflect “missed opportunity
53 costs,” activities foregone while foraging in a patch (Brown 1988). A forager should cease
54 feeding in a depletable food patch when the value of its harvest rate (H) no longer exceeds the
55 sum of its energetic cost of foraging (C), predation risk (P) and missed opportunity cost (MOC):
56 $H = C + P + MOC$ (Brown 1988).

57 GUDs have been used to measure foraging effort under predation risk in a variety of taxa
58 such as rodents (Brown 1988; Baker and Brown 2010), marsupials (Stokes et al. 2004) and
59 domestic goats (*Capra hircus*) (Shrader et al. 2008). The method has a range of applications,
60 including the assessment of avian microhabitat preferences (Oyugi and Brown 2003), the impact
61 of tourism on Nubian ibex (*Capra nubiana*) (Tadesse and Kotler, 2012), and how arboreal
62 primate species perceive predation risk in both the horizontal and vertical dimensions (Emerson,
63 Brown, and Linden 2011). In samango monkeys (*Cercopithecus mitis erythrarcus*), GUDs
64 decrease with height, highlighting a strong vertical axis of fear in this species (Emerson, Brown,
65 and Linden 2011) as opposed to the horizontal axis (sightlines and vegetation cover) more
66 typical of semi-terrestrial primates such as vervet monkeys (*Chlorocebus aethiops*) (Makin et al.
67 2012). Importantly, the GUD method allows researchers to quantify prey animals’ risk-taking
68 behavior without the need for direct observation.

69 We used GUDs and vertical arrangements of feeding stations (similar to Emerson,
70 Brown, and Linden 2011) to compare the depletion of artificial foraging patches by two groups
71 of samango monkeys in the presence and absence of human observers. We expected that human
72 observers should modulate monkeys' typical responses to predation risk. The baseline response
73 was assessed in our first prediction: an increase in GUDs at lower and upper stations in response
74 to higher perceived predation risks from terrestrial and aerial predators respectively (Emerson,
75 Brown, and Linden 2011). Anthropogenic changes to this baseline anti-predator response were
76 predicted as follows: monkeys would exhibit lower GUDs on days when observers are present.
77 Since humans are terrestrial, the "shielding" effect would be more pronounced in feeding stations
78 closer to the ground, as reflected in differential changes in GUDs along the vertical axis. Finally,
79 monkeys were expected to improve in patch exploitation over the course of the experiment, with
80 each additional visit day, a reflection of increased experience with the specific experiment
81 (Emerson and Brown 2013).

82 **METHODS**

83 **The study species**

84 The samango monkey (*Cercopithecus mitis erythrarcus*) is an arboreal guenon confined to
85 forests in southern Africa, the southernmost range extent of the polytypic *Cercopithecus mitis*
86 (Lawes 1990). Samango monkeys are medium sized monkeys (adult females ~4.4kg, adult males
87 ~7.6kg) that form single-male, multi-female groups), with group sizes ranging from 4-65
88 (Lawes, Cords, and Lehn 2011). The diet of samango monkeys is primarily frugivorous, but also
89 incorporates leaves, insects, and flowers (Lawes, Henzi, and Perrin 1990; Lawes 1991).

90

91 **The study area**

92 The study was conducted at the Lajuma Research Centre, in the Soutpansberg Mountains, South
93 Africa (23°02'23''S, 29°26'05''E) (Fig. 1). Local climate is temperate/mesothermal, with cool
94 dry winters from April-September and warm to hot wet summers from October-March (Willems,
95 Barton, and Hill 2009). The study area has natural fragments of tall forest (10-20m height)
96 occurring amongst areas of natural short forest (5-10m height). Confirmed predators of monkeys
97 at this site include leopard (*Panthera pardus*) (Chase Grey, Kent, and Hill 2013), as well as
98 caracal (*Caracal caracal*), crowned eagle (*Stephanoaetus coronatus*), African black eagle
99 (*Aquila verreauxii*) and African rock python (*Python sebae*) (Willems and Hill 2009; Ian
100 Gaigher, pers. comm. 2013; Coleman and Hill 2014). Terrestrial carnivores are ambush hunters,
101 leading to stronger risk responses close to ground level, while aerial predators increase primates'
102 predation risk close to the canopy (Seyfarth, Cheney, and Marler 1980; Emerson et al. 2011).
103 Sympatric diurnal primates are chacma baboon (*Papio ursinus*) and vervet monkey (*Chlorocebus*
104 *aethiops*). There is no hunting of primates at this site, and since the monkeys forage naturally,
105 without access to crops, there is no human-wildlife conflict or anthropogenic persecution of the
106 monkeys. Leopards face persecution from landowners below the mountains for perceived
107 livestock depredation, and are also subject to legal and illegal hunting in parts of the
108 Soutpansberg (Chase Grey 2011; Chase Grey, Kent, and Hill 2013) such that the leopard
109 population is likely to be wary of humans in this environment.

110

111 **Field experiments**

112 We studied two habituated samango groups during the winter (dry) season from May to August
113 2013. “Barn” group (40 individuals) has been regularly followed since September 2009, while
114 “House” group (60 individuals) has been under systematic observation since July 2011. This

115 persistent contact with study groups has resulted in their habituation to people, i.e. a reduction in
116 fear of human observers (Williamson and Feistner, 2011), presumably allowing the study
117 subjects to exhibit their natural behavioral patterns despite the presence of observers. “Follows”
118 were observation sessions in which single observers followed groups from dawn until dusk to
119 conduct standard behavioral data collection independent of our particular experiment. Groups
120 were not followed every day, but an average of 3-4 days per week. We exploited this “natural”
121 variation in observer presence, using follow days as a predictor variable throughout our study.

122 We generated 16 random points (8 per group) within the groups’ exclusive winter ranges
123 (outside the groups’ range overlap area), mapped using data from the previous year, and selected
124 emergent trees within 20m of those points based on tree characteristics (height 11-12m, with
125 diameter at breast height >25cm) and habitat type (semi-deciduous forest bordering tall
126 evergreen mist-belt forest) (Fig. 1). Group arrays were placed roughly equidistant from a known
127 crowned eagle perch, although the Barn group array was relatively closer to the active crowned
128 eagle nest-site (Fig. 1). Eagle calls were heard almost daily from both groups’ ranges. Ground
129 predators (leopard and caracal) were captured on camera traps visiting the covered GUD basins
130 during the night in groups’ ranges.

131 We modified methodology from Emerson, Brown, and Linden (2011), who previously
132 validated the GUD protocol for studying samango monkey ecology at our site. We established
133 artificial foraging patches at each tree, consisting of plastic basins covered with a rope mesh to
134 reduce access and so slow foraging rates, suspended at 0.1m, 2.5m, 5.0m, and 7.5m (four basins
135 per tree, one basin at each height). The 0.1m basin level predominantly tested terrestrial
136 predation risk and the 7.5m basin level, just below the canopy, tested risk from eagles. Basins
137 were filled with 4 litres of sawdust, “baited” with 25 shelled peanuts before 07:00, and sieved

138 after 16:00 on each of 20 experimental days (4 consecutive days per week for 5 weeks). We
139 recorded the number of peanuts left (the GUD) and the amount of sawdust spilled (in litres) from
140 all four basins at each tree (n=64 basins, 32 per group over 20 days), topping off any spilled
141 sawdust so that each basin contained 4L at the start of the next experimental day. We also noted
142 if researchers were following the monkeys that day for observational data collection. On follow
143 days, human observers were present at or in the vicinity of the feeding stations, but did not
144 actively interfere with monkeys' foraging behavior. Barn group visited GUD stations on 20/20
145 days, of which 8 were follow days, and House group visited trees on 17/20 days, of which 11
146 were follow days. On non-follow days and in the absence of observers in the GUD areas, camera
147 trap data confirmed that no other animals visited GUD stations, as was found in Emerson and
148 Brown's study at this same site (2013).

149

150 **Data analysis**

151 Examination of the numbers of peanuts remaining in basins revealed significant variation
152 throughout the study period (Fig. S1). This created a statistical analysis challenge, as the
153 experimental design created three scales at which random processes were likely to influence
154 GUD variation. Firstly, the trees onto which we placed our basins may have varied in their
155 perceived risk due to subtle differences in, for example, canopy cover and understory foliage
156 density (Emerson and Brown 2013). Secondly, basin utilization may have varied between days
157 due to variation in environmental factors that influence risk (e.g. cloud cover), which could
158 affect all monkeys equally on a given day. Thirdly, the utilisation of basins on a tree on a given
159 day may have varied stochastically due to local chance events that may impact individual
160 monkeys differentially (e.g. a monkey becoming startled). It is important that these biologically

161 relevant sources of stochastic variation are accounted for in the statistical analysis because
 162 ignoring them could lead to inflated type I errors (Richards 2008). As we could not find a
 163 statistical package capable of explicitly describing the above-mentioned correlated variance
 164 structure (including GLMMs), we instead developed a likelihood function linking the
 165 experimental design with our hypotheses on the effect of environmental factors and observation
 166 on risk-taking behavior (see equation 2). Specifically, we developed a model that allowed us to
 167 look for evidence that GUDs were influenced by basin height (Z), the presence of human
 168 followers that day (F), and the number of days the tree had been previously visited during the
 169 study (V). For a basin placed at height z_k on tree i , the logit-transformed probability that each
 170 peanut remained in the basin on sampling day j was described by:

$$\text{logit } \bar{p}_{ijk} = (\beta_0 + \beta_F x_{ij}) + (\beta_Z + \beta_{ZF} x_{ij}) z_k + (\beta_V + \beta_{VF} x_{ij}) v_{ij}, \quad (1)$$

171 where v_{ij} is the number of days the tree had previously been visited, x_{ij} indicates if the group was
 172 followed that day (0 = no, 1 = yes), and the β are model parameters describing effect sizes. The
 173 three above-mentioned sources of variation were included by considering the following
 174 likelihood function:

$$L(\theta|\text{data}) = \prod_{j=1}^{20} \int_{u=-\infty}^{\infty} f_n(u|0,1) \prod_{i=1}^8 \prod_{k=1}^4 f_{bb}(y_{ijk}|25, T[\bar{p}_{ijk}, u\sigma + w_i], \phi) du, \quad (2)$$

175 where θ is the set of parameters that define the statistical model, $f_n(u|0,1)$ is the density function
 176 for the standard normal, $f_{bb}(y|N,p,\phi)$ is the beta-binomial distribution with variance parameter ϕ
 177 (Richards 2008), and $T[\bar{p}, \tau] = e^\lambda / (1 + e^\lambda)$ where $\lambda = \ln(\bar{p}/(1 - \bar{p})) + \tau$. Potential

178 differences in tree affinities (i.e. the first source of stochastic variation) were included by
179 associating each tree i with a parameter w_i (a positive value indicates that, on average, more
180 peanuts were left in the tree's basins throughout the study). The transformation T ensures that the
181 probability a peanut remains in a basin when placed in a specific tree on any day is bounded by
182 $(0, 1)$. The standard normal distribution accounts for the second source of variation (a positive u
183 indicates that GUDs were higher than average on that day), with the degree of daily variation
184 quantified by σ . The third source of GUD variation mentioned is accounted for by incorporating
185 the beta-binomial distribution. We found that the likelihood function described by equation (2)
186 was able to describe the structured variation in peanut numbers observed (Fig. S1).

187 Data were effectively excluded for non-visit days by setting f_{bb} to 1, i.e., on days when a
188 tree was not visited, as these conditions provide no information on GUDs. Further, we also set f_{bb}
189 to 1 when more than 1L of sawdust was spilled from a basin (2.9% of basins). These non-visits
190 and excessive spillage occurred at a very low rate, and preliminary analyses indicated that these
191 events were random and thus unlikely to cause significant parameter bias. The Barn and House
192 groups' feeding behavior was inferred using GUDs from 590 and 463 basin samples,
193 respectively (32 basins x 20 experimental days yielded 640 maximum possible samples). Models
194 describing patterns of foraging by the Barn and House groups were coded according to equations
195 (1) and (2) and fit using maximum likelihood. For both groups, we used likelihood ratio tests
196 (LRTs) to look for statistical evidence that the parameters describing the effects of factors: F, Z
197 and V, differed from zero. Specifically, the importance of each of these three factors was
198 evaluated by comparing the full model with the model that removed the factor of interest. A
199 human shield effect is inferred if factor F is deemed statistically significant and GUDs are
200 reduced in the presence of a follower. In this case, the statistical significance of factors Z and V

201 reveals how followers modulate the monkeys' vertical axis of fear and their propensity to change
202 their level of fear over time.

203 **RESULTS**

204 Both groups of samango monkeys responded in the predicted way to terrestrial, but not aerial,
205 predation risk. The presence of followers modulated one group's baseline responses to potential
206 risk from ground predators. The likelihood ratio tests, when applied to the Barn group data,
207 indicated that GUDs varied with height and the presence of a follower but not with the number
208 of days that the tree had been visited by the group during the study (Table I). GUDs were
209 reduced with increasing height on days when the group was not followed, but GUDs remained
210 low at all heights when the group was followed (Fig. 2A,C). Like Barn group, GUDs of House
211 group were also influenced by basin height and observer follows; however, GUDs were also
212 influenced by tree experience (Table I). For House group, GUDs tended to decrease with height,
213 were lower when the group was followed, and became lower the more the tree was visited (Fig.
214 2B, D). Importantly, the height effect on GUDs would not have been detected for Barn group if
215 the analysis only included data taken from days the group were followed (Table I; Fig. 2A). In
216 other words, while in the House group the effect of basin height on GUDs remained detectable in
217 the presence of observers, in the Barn group the height effect disappeared when observers were
218 present suggesting that a vertical axis of fear would not have been detected if the group was
219 always followed. In general, Barn group had lower GUDs than House group (Fig. 2). Table S1
220 provides the maximum likelihood parameter estimates for each of the models presented in Table
221 I.

222 **DISCUSSION**

223 Our results suggest that human observers can alter the vertical axis of fear in habituated, arboreal
224 monkeys. As predicted, we found a strong vertical axis of fear, consistent with Emerson, Brown,
225 and Linden (2011), with monkeys depleting the least food at patches near ground level. More
226 significantly, however, monkeys' variation in perceived risk along this vertical gradient was
227 affected by the presence of human observers (Fig. 1). This effect was most apparent at lower
228 levels in the tree, suggesting that behavioral researchers were perceived as shields against
229 terrestrial predators in particular. We speculate that this observed pattern may be due to humans
230 passively deterring predators from the immediate area, rather than playing the role of active
231 sentinels. This is consistent with findings from ungulates, where mountain nyala (*Tragelaphus*
232 *buxtoni*) avoid hyenas (*Crocuta crocuta*) by over-nighting next to human settlements (Meshesha
233 2013), and moose (*Alces alces*) avoiding calf depredation by brown bears (*Ursus arctos*) by
234 birthing near to human infrastructure (Berger 2007). In our case, monkeys did not actively seek
235 out humans to avoid predators, although monkeys' tolerance of human followers could in part be
236 due to anti-predator and other perceived benefits (e.g. displacement of other groups not
237 habituated to humans).

238 Contrary to predictions from the "Group Size Effect", which posits that larger groups
239 perceive themselves as less vulnerable to ground predators and are expected to deplete more food
240 in risky areas (Quenette 1990; Miller 2002; Makin et al. 2012), overall GUDs were lower (more
241 food was depleted) by the smaller Barn group. This discrepancy may be explained by this
242 group's greater prior experience with both observers and GUD experiments (Emerson, Brown,
243 and Linden 2011; Emerson and Brown 2013). In contrast, the less experienced House group
244 showed an expected response to increasing experience with GUDs reducing with repeat visits,
245 although GUDs decreased more rapidly in the presence of observers. These results suggest that

246 even a slight difference in experience with human observers can affect subjects' behavior and
247 that GUD experiments themselves – and animals' learning of foraging tasks – may influence
248 future experiments (Dukas 2008). Even though future experiments may yield lower GUDs due to
249 past experience, it seems that GUDs, nonetheless, are able to measure differences between
250 treatments, such as height.

251 Our results add to the growing literature suggesting that observer presence never becomes
252 truly “neutral” to study animals (Jack et al. 2008; Crofoot et al. 2010; McDougall 2012) and
253 highlight the importance of considering the effects that habituation has on animal behavior,
254 particularly where study populations may be at risk from other human activities, for example
255 poaching, poisoning or conflict with agriculturalists (Williamson and Feistner 2011).

256 Importantly, we would not have detected a vertical axis of fear for Barn group if they were
257 always followed (Fig. 2A). If observer presence can significantly alter GUDs in study animals
258 (Fig. 2), and modify the effects of learning/experience, this clearly has important implications for
259 studies of vigilance and predator-prey interactions. Researchers thus need to be cautious in
260 interpreting the effect sizes of predation parameters where the extent of human presence may
261 vary (e.g. Isbell and Young 1993), and use indirect and experimental methods in combination
262 with or instead of direct observation. For example, while behavioral research on the same
263 population of samango monkeys found the landscape of fear from eagles to be the most
264 significant determinant of range use (Coleman and Hill 2014), the study reported no response to
265 the threat of predation from leopards. Further work would be required to determine whether this
266 represents a true reflection of the landscape of fear from terrestrial predators or whether human
267 shield effects influence ranging patterns in the presence of observers. Similarly, habituation has
268 been shown to interact with the effects of risk allocation in blackbirds (*Turdus merula*) in

269 determining flight initiation distances in the face of ‘usual’ and ‘novel’ predators (Rodriguez-
270 Prieto et al. 2009). Nevertheless, these insights about fear, behavior and predator-prey
271 relationships can assist conservation managers in understanding how anthropogenic effects
272 influence species distribution, habitat selection and risk-sensitive behavior (Berger 2007).

273 **ACKNOWLEDGMENTS**

274 We thank Ian Gaigher for permission and support in working at Lajuma. For field assistance, we
275 thank Tyler Carver, Noeks Cilliers, Christel Dalhuizen, Ellie Darbey, Leanne Fitzgerald, Marion
276 Gohier, Trevor Jones, Amy Kuczynski, Gregoire Moutardier, Kasim Rafiq, Rachel Stokes, Kyle
277 Stuart, Oldrich van Schalkwyk, and Chloë Wright. The project was approved by the Limpopo
278 Province Department of Economic Development and Tourism, with ethical approval from
279 Durham University's Life Sciences Ethical Review Process Committee and from the University
280 of the Free State's Inter-Faculty Animal Ethics Committee. This work was supported by
281 postdoctoral grants to KN from the Claude Leon Foundation and from the Durham University
282 COFUND research fellowship program with additional fieldwork support from Earthwatch.

283

284 **REFERENCES**

285 Abu Baker MA, Brown JS. 2010. Islands of fear: effects of wooded patches on habitat suitability
286 of the striped mouse in a South African grassland. *Funct. Ecol.* 24: 1313-1322.

287 Berger J. 2007. Fear, human shields and the redistribution of prey and predators in protected
288 areas. *Biol. Lett.* 3: 620-623.

289 Brown JS. 1988. Patch use as an indicator of habitat preference, predation risk, and competition.
290 *Behav. Ecol. Sociobiol.* 22: 37-47.

291 Chase Grey J. 2011. Leopard population dynamics, trophy hunting and conservation in the
292 Soutpansberg Mountains, South Africa. PhD thesis, Durham University, UK.

293 Chase Grey J, Kent VT, Hill RA. 2013. Evidence of a high density population of harvested
294 leopards in a montane environment. *PLOS ONE* 8: e82832.

295 Coleman BT, Hill RA. 2014. Living in a landscape of fear: the impact of predation, resource
296 availability and habitat structure on primate range use. *Anim. Behav.* 88: 165-173.

297 Crofoot MC, Lambert TD, Kays R, Wikelski MC. 2010. Does watching a monkey change
298 its behaviour? Quantifying observer effects in habituated wild primates using automated
299 radiotelemetry. *Anim. Behav.* 80: 475-480.

300 Dukas R. 2008. Life history of learning: Performance curves of honeybees in the wild. *Ethology*
301 114: 1195-1200.

302 Emerson SE, Brown JS, Linden JD. 2011. Identifying Sykes' monkeys', *Cercopithecus*
303 *albugularis erythrarchus*, axes of fear through patch use. *Anim. Behav.* 81: 455-462.

304 Emerson SE, Brown JS. 2013. Identifying preferred habitat of samango monkeys
305 (*Cercopithecus (nictitans) mitis erythrarchus*) through patch use. *Behav. Process.* 100:
306 214-221.

307 Isbell LA, Young TP. 1993. Human presence reduces predation in a free-ranging vervet monkey
308 population in Kenya. *Anim. Behav.* 45: 1233-1235.

309 Jack KM, Lenz BB, Healan E, Rudman S, Schoof VA, Fedigan L. 2008. The effects of
310 observer presence on the behavior of *Cebus capucinus* in Costa Rica. *Am. J. Primatol.*
311 70: 490-494.

312 Lawes MJ. 1990. The distribution of the samango monkey (*Cercopithecus mitis erythrarcus*
313 Peters, 1852 and *Cercopithecus mitis labiatus* I. Geoffroy, 1843) and forest history in
314 Southern Africa. *J. Biogeogr.* 17: 669-680.

315 Lawes MJ, Henzi SP, Perrin MR. 1990. Diet and feeding behavior of samango
316 monkeys (*Cercopithecus mitis labiatus*) in Ngoye Forest, South-Africa. *Folia Primatol.*
317 54: 57-69.

318 Lawes MJ. 1991. Diet of samango monkeys (*Cercopithecus mitis erythrarchus*) in the Cape
319 Vidal Dune Forest, South-Africa. *J. Zool.* 224: 149-173.

320 Lawes MJ, Cords M, Lehn C. 2011. *Cercopithecus mitis* profile. In J. Kingdon & J. Kalina (eds.)
321 *Primates of Africa volume II, The mammals of Africa*. Berkley, CA: University of
322 California Press.

323 Makin DF, Payne HFP, Kerley GIH, Shrader AM. 2012. Foraging in a 3-D world: how does
324 predation risk affect space use of vervet monkeys? *J. Mammal.* 93: 422-428.

325 McDougall P. 2012. Is passive observation of habituated animals truly passive? *J. Ethol.* 30: 219-
326 223.

327 Meshesha AA. 2013. *Landscape genetics and behavioural ecology of mountain nyala*
328 (*Tragelaphus buxtoni*) in the Southern Highlands of Ethiopia, PhD thesis, University of
329 Oslo.

330 Miller LE. 2002. The role of group size in predator sensitive foraging decisions for wedge-
331 capped capuchin monkeys (*Cebus olivaceus*). In: Miller LE, editor. *Eat or be eaten:*
332 *predator-sensitive foraging among primates*. Cambridge, UK: Cambridge University
333 Press. p. 95-104.

334 Ngoprasert D, Lynam AJ, Gale GA. 2007. Human disturbance affects habitat use and behaviour
335 of Asiatic leopard *Panthera pardus* in Kaeng Krachan National Park, Thailand. *Oryx* 41:
336 343-351.

337 Oyugi JO, Brown JS. 2003. Giving-up densities and habitat preferences of European starlings
338 and American robins. *Condor* 105: 130-135.

339 Quenette PY. 1990. Functions of vigilance behavior in mammals: a review. *Acta Oecol.* 11: 801-
340 818.

341 Richards SA. 2008. Dealing with overdispersed count data in applied ecology. *J. Appl. Ecol.* 45:
342 218-227.

343 Rodriguez-Prieto I, Fernández-Juricic E, Martín J, Regis Y. 2009. Antipredator behavior in
344 blackbirds: habituation complements risk allocation. *Behav. Ecol.* 20: 371-377.

345 Rogala JK, Hebblewhite M, Whittington J, White CA, Coleshill J, Musiani M. 2011.
346 Human activity differentially redistributes large mammals in the Canadian Rockies
347 national parks. *Ecol. Soc.* 16: 16-39.

348 Seyfarth RM, Cheney DL, Marler P. 1980. Monkey responses to three different alarm
349 calls: evidence of predator classification and semantic communication. *Science* 210: 801-
350 803.

351 Shrader AM, Brown JS, Kerley GIH, Kotler BP. 2008. Do free-ranging goats show “landscapes
352 of fear”? Patch use in response to habitat features and predator cues. *J. Arid Environ.* 72:
353 1811-1819.

354 Stanford C. 1998. *Chimpanzee and red colobus: the ecology of predator and prey*. Harvard:
355 Harvard University Press.

356 Stokes VL, Pech RP, Banks PB, Arthur AD. 2004. Foraging behaviour and habitat use
357 by *Antechinus flavipes* and *Sminthopsis murina* (Marsupialia: Dasyuridae) in response to
358 predation risk in eucalypt woodland. *Biol. Conserv.* 117: 331-342.

359 Tadesse SA, Kotler BP. 2012. Impact of tourism on Nubian ibex (*Capra nubiana*) revealed
360 through assessment of behavioral indicators. *Behav. Ecol.* 1-6.


361 Verdolin JL. 2006. Meta-analysis of foraging and predation risk trade-offs in terrestrial systems.
362 *Behav. Ecol. Sociobiol.* 60: 457-464.

363 Willems EP, Hill RA. 2009. Predator-specific landscapes of fear and resource distribution:
364 effects on spatial range use. *Ecology* 90: 546-555.

365 Willems EP, Barton RA, Hill RA. 2009. Remotely sensed productivity, home range selection and
366 local range use by an omnivorous primate. *Behav. Ecol.* 20: 985-992.

367 Williamson EA, Feistner ATC. 2011. Habituating primates: Processes, techniques, variables and
368 ethics. In Setchell JM and Curtis DJ, editors. *Field and laboratory methods in*
369 *primatology: a practical guide*. Cambridge: Cambridge University Press. p. 33-50.

370 **Fig. 1** Study area showing locations of 16 experimental trees. Minimum convex polygons
 371 represent winter ranges of two samango monkey study groups based on 773 waypoints for Barn
 372 group and 695 waypoints for House group from the previous winter. The vegetation map is based
 373 on unpublished data. The active crowned eagle nest and common perch site are shown.




374

375

376

377 **Fig. 2** Observed GUDs for both groups split according to whether or not data were collected on
 378 days when the group was followed or not. Panels A and B show basin height effects. Panels C
 379 and D show how GUDs change with tree experience for the lowest basins that were hung at a
 380 height of 0.1m. Error bars depict 95% confidence intervals, assuming data are drawn from a
 381 beta-binomial distribution, and were calculated using the profile likelihood approach.



382

383

384 **Table I:** Summary of the likelihood ratio tests (LRTs) used to look for evidence that GUDs were
385 affected by the presence of a follower (F), the height of the food basin (Z), and the number of
386 days the group had previously visited the experimental tree (V). Results are presented for both
387 groups. The first set of tests considers all data (i.e. days followed and not followed), whereas the
388 second set only examines data collected on days when a group was followed. Factor removed
389 indicates which factor the LRT is testing for statistical significance.

Models compared	Factor removed	G	df	P-value
Barn Group				
<i>Consider days when the group was followed and days when it was not followed</i>				
M(F+Z+V), M(Z+V)	F	14.86	3	0.002
M(F+Z+V), M(F+V)	Z	29.01	2	< 0.001
M(F+Z+V), M(F+Z)	V	0.04	2	0.980
<i>Only consider days when the group was followed</i>				
M(Z+V), M(V)	Z	0.12	1	0.733
M(Z+V), M(Z)	V	0.07	1	0.796
House Group				
<i>Consider days when the group was followed and days when it was not followed</i>				
M(F+Z+V), M(Z+V)	F	10.94	3	0.012
M(F+Z+V), M(F+V)	Z	18.25	2	< 0.001
M(F+Z+V), M(F+Z)	V	9.19	2	0.010
<i>Only consider days when the group was followed</i>				
M(Z+V), M(V)	Z	8.49	1	0.004
M(Z+V), M(Z)	V	0.77	1	0.380

390 **Supplementary Material**

391 **Table S1:** Maximum likelihood parameter estimates for models fit to both sets of group data and
 392 used for the likelihood ratio tests. Model descriptions that include a Z, F, or V, indicate that GUD is
 393 affected by basin height, whether the group is followed that day, and the number of times the tree
 394 had been previously visited, respectively. K is the number of model parameters estimated and LL_{\max}
 395 is the maximum log-likelihood. The seven independent model parameters that describe variation
 396 among trees, w_1 - w_7 , are not presented here. Note that $w_8 = -\sum_{i=1}^7 w_i$. Fits are presented for both
 397 groups. The first set of parameter estimates are for all the data (i.e. days followed and not followed),
 398 whereas the second set only examines data collected on days when a group was followed.

Maximum-likelihood model parameter estimates

Model	β_0	β_F	β_Z	β_{ZF}	β_V	β_{VF}	ϕ	σ	K	LL_{\max}
-------	-----------	-----------	-----------	--------------	-----------	--------------	--------	----------	-----	-------------

Barn Group

Consider days when the group was followed and days when it was not followed

M(Z+V)	-1.483	0	-0.079	0	-0.011	0	0.160	0.583	12	-1352.1
M(F+V)	-1.519	-0.720	0	0	-0.007	0.004	0.171	0.455	13	-1359.2
M(F+Z)	-1.185	-1.058	-0.110	0.094	0	0	0.156	0.473	13	-1344.7
M(F+Z+V)	-1.163	-1.021	-0.112	0.096	-0.001	0.000	0.157	0.473	15	-1344.7

Only consider days when the group was followed

M(V)	-2.235	NA	0	NA	-0.007	NA	0.151	0.203	11	-481.0
M(Z)	-2.261	NA	-0.010	NA	0	NA	0.150	0.208	11	-481.0
M(Z+V)	-2.210	NA	-0.010	NA	-0.005	NA	0.151	0.203	12	-480.9

House Group

Consider days when the group was followed and days when it was not followed


M(Z+V)	-0.002	0	-0.103	0	-0.098	0	0.371	0.496	12	-1253.0
M(F+V)	-0.002	-1.169	0	0	-0.108	0.087	0.401	0.297	13	-1256.9
M(F+Z)	-0.387	-0.654	-0.120	0.044	0	0	0.371	0.447	13	-1252.1
M(F+Z+V)	-0.002	-0.852	-0.097	0.017	-0.083	0.058	0.372	0.330	15	-1247.5

Only consider days when the group was followed

M(V)	-1.224	NA	0	NA	-0.016	NA	0.384	0.065	11	-822.5
M(Z)	-1.055	NA	-0.076	NA	0	NA	0.366	0.079	11	-818.6
M(Z+V)	-0.953	NA	-0.075	NA	-0.016	NA	0.365	0.070	12	-818.2


400

401 (A) Barn group when not followed



402


403 (B) Barn group when followed



404


405

406 (C) House group when not followed



407

408 (D) House group when followed



409

410 **Figure S1:** Peanuts remaining in basins for two groups of samango monkeys across 20 sampling
 411 days. Note that sampling days span a 5-week period. Data are segregated according to whether or not
 412 the group was followed on the sampling day. Each panel depicts peanuts remaining at a specific tree
 413 for four basin heights. These data show very high variation in peanut numbers across sampling days
 414 and even among basins on a given day, supporting our assumption that the data are beta-binomial
 415 distributed. The data also suggest some degree of between-tree variation, which supports our model
 416 incorporating tree as a random factor.