

Obtaining Planarity by Contracting Few Edges^{*}

Petr A. Golovach¹, Pim van 't Hof¹, and Daniël Paulusma²

¹ Department of Informatics, University of Bergen, Norway
{petr.golovach,pim.vanthof}@ii.uib.no

² School of Engineering and Computing Sciences, Durham University, UK
daniel.paulusma@durham.ac.uk

Abstract. The PLANAR CONTRACTION problem is to test whether a given graph can be made planar by using at most k edge contractions. This problem is known to be NP-complete. We show that it is fixed-parameter tractable when parameterized by k .

Keywords. Planar graphs, edge contractions, FPT algorithms

1 Introduction

Numerous problems in algorithmic graph theory, with a large variety of applications in different fields, can be formulated as graph modification problems. A graph modification problem takes as input an n -vertex graph G and an integer k , and the question is whether G can be modified into a graph that belongs to a prescribed graph class, using at most k operations of a certain specified type. Some of the most common graph operations that are used in this setting are vertex deletions, edge deletions and edge additions, leading to famous problems such as FEEDBACK VERTEX SET, ODD CYCLE TRANSVERSAL, MINIMUM FILL-IN and CLUSTER EDITING, to name but a few. More recently, the study of graph modification problems allowing only *edge contractions* has been initiated, yielding several results that we will survey below. The contraction of an edge removes both end-vertices of an edge and replaces them by a new vertex, which is made adjacent to precisely those vertices that were adjacent to at least one of the two end-vertices. Choosing edge contraction as the only permitted operation leads to the following decision problem, for each graph class \mathcal{H} .

\mathcal{H} -CONTRACTION

Instance: A graph G and an integer k .

Question: Does there exist a graph $H \in \mathcal{H}$ such that G can be contracted to H , using at most k edge contractions?

Heggernes et al. [12] presented a $2^{k+o(k)} + n^{O(1)}$ time algorithm for \mathcal{H} -CONTRACTION when \mathcal{H} is the class of paths. Moreover, they showed that in this case the problem has a linear vertex kernel. When \mathcal{H} is the class of trees, they showed

^{*} This paper is supported by EPSRC (EP/G043434/1) and Royal Society (JP100692), and by the Research Council of Norway (197548/F20). The results of this paper will be presented at MFCS 2012 [7].

that the problem can be solved in $4.98^k n^{O(1)}$ time, and that a polynomial kernel does not exist unless $\text{NP} \subseteq \text{coNP/poly}$. When the input graph is a chordal graph with n vertices and m edges, then \mathcal{H} -CONTRACTION can be solved in $O(n + m)$ time when \mathcal{H} is the class of trees and in $O(nm)$ time when \mathcal{H} is the class of paths [10]. Heggenes et al. [11] proved that \mathcal{H} -CONTRACTION is fixed-parameter tractable when \mathcal{H} is the class of bipartite graphs and k is the parameter. This also follows from a more general result from a recent paper by Marx, O’Sullivan, and Razgon [18] on generalized bipartization. Golovach et al. [8] considered the class of graphs of minimum degree at least d for some integer d . They showed that in this case \mathcal{H} -CONTRACTION is fixed-parameter tractable when both d and k are parameters, $\text{W}[1]$ -hard when k is the parameter, and para-NP-complete when d is the parameter.

The combination of planar graphs and edge contractions has been studied before in a closely related setting. Kamiński, Paulusma and Thilikos [13] showed that for every fixed graph H , there exists a polynomial-time algorithm for deciding whether a given planar graph can be contracted to H . Very recently, this result was improved by Kamiński and Thilikos [14]. They showed that, given a graph H and a planar graph G , the problem of deciding whether G can be contracted to H is fixed-parameter tractable when parameterized by $|V(H)|$.

Our Contribution. We study \mathcal{H} -CONTRACTION when \mathcal{H} is the class of planar graphs, and refer to the problem as PLANAR CONTRACTION. This problem is known to be NP-complete due to a more general result on \mathcal{H} -CONTRACTION by Asano and Hirata [2]. We show that the PLANAR CONTRACTION problem is fixed-parameter tractable when parameterized by k . This result complements the following results on two other graph modification problems related to planar graphs. The problem of deciding whether a given graph can be made planar by using at most k vertex deletions was proved to be fixed-parameter tractable independently by Marx and Schlotter [17], who presented a quadratic-time algorithm for every fixed k , and by Kawarabayashi [15], whose algorithm runs in linear time for every k . Kawarabayashi and Reed [16] showed that deciding whether a graph can be made planar by using at most k edge deletions can also be done in linear time for every fixed k .

Our algorithm for PLANAR CONTRACTION starts by finding a set S of at most k vertices whose deletion transforms G into a planar graph. Such a set can be found by using either the above-mentioned linear-time algorithm by Kawarabayashi [15] or the quadratic-time algorithm by Marx and Schlotter [17]. The next step of our algorithm is based on the irrelevant vertex technique developed in the graph minors project of Robertson and Seymour [20,22]. We show that if the input graph G has large treewidth, we can find an *edge* whose contraction yields an equivalent, but smaller instance. After repeatedly contracting such irrelevant edges, we invoke Courcelle’s Theorem [4] to solve the remaining instance in linear time.

We finish this section by making two remarks that show that we cannot apply the techniques that were used to prove fixed-parameter tractability of the vertex deletion and edge deletion variants of PLANAR CONTRACTION. First, a crucial

observation in the paper of Kawarabayashi and Reed [16] is that any graph that can be made planar by at most k edge deletions must have bounded genus. This property is heavily exploited in the case where the treewidth of the input graph is large. The following example shows that we cannot use this technique in our setting. Take a complete biclique $K_{3,r}$ with partition classes A and B , where $|A| = 3$ and $|B| = r$ for some integer $r \geq 3$. Now make the vertices in A pairwise adjacent and call the resulting graph G_r . Then G_r can be made modified into a planar graph by contracting one of the edges in A . However, the genus of G_r is at least the genus of $K_{3,r}$, which is equal to $\frac{r-2}{2}$ [3].

Second, the problem of deciding whether a graph can be made planar by at most k vertex deletions for some fixed integer k , i.e., k is not part of the input, is called the k -APEX problem. As observed by Kawarabayashi [15] and Marx and Schlotter [17], the class of so-called k -apex graphs (graphs that can be made planar by at most k vertex deletions) is closed under taking minors. This means that the k -APEX problem can be solved in cubic time for any fixed integer k due to deep results by Robertson and Seymour [21]. However, we cannot apply Robertson and Seymour's result on PLANAR CONTRACTION, because the class of graphs that can be made planar by at most k edge contractions is not closed under taking minors, as the following example shows. Take the complete graph K_5 on 5 vertices. For each edge $e = uv$, add a path P_e from u to v consisting of p new vertices for some integer $p \geq k$. Call the resulting graph G_p^* . Then G_p^* can be made planar by contracting an arbitrary edge of the original K_5 . However, if we remove all edges of this K_5 , we obtain a minor of G_p^* that is a subdivision of the graph K_5 . In order to make this minor planar, we must contract all edges of a path P_e , so we need at least $p + 1 > k$ edge contractions.

2 Preliminaries

Throughout the paper we consider undirected finite graphs that have no loops and no multiple edges. Whenever we consider a graph problem, we use n to denote the number of vertices of the input graph. We refer to the text book of Diestel [5] for undefined graph terminology and to the monographs of Downey and Fellows [6] for more information on parameterized complexity.

Let $G = (V, E)$ be a graph and let S be a subset of V . We write $G[S]$ to denote the subgraph of G induced by S , i.e., the subgraph of G with vertex set S and edge set $\{uv \mid u, v \in S \text{ with } uv \in E\}$. We write $G - S = G[V \setminus S]$, and for any subgraph H of G , we write $G - H$ to denote $G - V(H)$. We say that two disjoint subsets $U \subseteq V$ and $W \subseteq V$ are *adjacent* if there exist two vertices $u \in U$ and $w \in W$ such that $uw \in E$. Let H be a graph that is not necessarily vertex-disjoint from G . Then $G \cup H$ denotes the graph with vertex set $V(G) \cup V(H)$ and edge set $E(G) \cup E(H)$, and $G \cap H$ denotes the graph with vertex set $V(G) \cap V(H)$ and edge set $E(G) \cap E(H)$.

The *contraction* of edge uv in G removes u and v from G , and replaces them by a new vertex made adjacent to precisely those vertices that were adjacent to u or v in G . A graph H is a *contraction* of G if H can be obtained from G by a

sequence of edge contractions. Alternatively, we can define a contraction of G as follows. An H -witness structure \mathcal{W} is a partition of $V(G)$ into $|V(H)|$ nonempty sets $W(x)$, one for each $x \in V(H)$, called H -witness sets, such that each $W(x)$ induces a connected subgraph of G , and for all $x, y \in V(H)$ with $x \neq y$, the sets $W(x)$ and $W(y)$ are adjacent in G if and only if x and y are adjacent in H . Clearly, H is a contraction of G if and only if G has an H -witness structure; H can be obtained by contracting each witness set into a single vertex. A *witness edge* is an edge of G whose end-vertices belong to two different witness sets.

Let \mathcal{W} be an H -witness structure of G . For our purposes, we sometimes have to contract edges in G such that the resulting graph does *not* contain H as contraction. In order to do this it is necessary to *destroy* \mathcal{W} , i.e., to contract at least one witness edge in \mathcal{W} . After all, every edge that is not a witness edge has both its end-vertices in the same witness set of \mathcal{W} , which means that contracting such an edge yields an H -witness structure of a contraction of G . Hence, contracting all such non-witness edges transforms G into H itself. Note that if we destroy \mathcal{W} by contracting a witness edge e in \mathcal{W} , the obtained graph still has H as a contraction if e was a non-witness edge in some other H -witness structure of G . Hence, in order to obtain a graph that does not have H as a contraction, it is necessary and sufficient to destroy *all* H -witness structures of G .

A *planar graph* G is a graph that can be embedded in the plane, i.e., that can be drawn in the plane so that its edges intersect only at their end-vertices. A graph that is actually drawn in such a way is called a *plane graph*, or an *embedding* of the corresponding planar graph. A plane graph G partitions the rest of the plane into a number of connected regions, called the *faces* of G . Each plane graph has exactly one unbounded face, called the *outer face*; all other faces are called *inner faces*. Let C be a cycle in a plane graph G . Then, by the Jordan curve theorem (cf. [5]), C divides the plane in exactly two regions: the *outer* region of C , containing the outer face of G , and the *inner* region of C . We say that a vertex u of G lies *inside* C if u is in the inner region of C . Similarly, u lies *outside* C if u is in the outer region of C . The *interior* of C with respect to G , denoted $\text{interior}_G(C)$, is the set of all vertices of G that lie inside C . We also call these vertices *interior* vertices of C . We say that C *separates* the vertices that lie inside C from the vertices that lie outside C . A sequence of mutually vertex-disjoint cycles C_1, \dots, C_q in a plane graph is called *nested* if there exist disks $\Delta_1, \dots, \Delta_q$ such that C_i is the boundary of Δ_i for $i = 1, \dots, q$, and $\Delta_{i+1} \subset \Delta_i$ for $i = 1, \dots, q - 1$. We also refer to such a sequence of nested cycles as *layers*. We say that a vertex u lies *between* two nested cycles C_i and C_j with $i < j$ if u lies in the inner region of C_i and in the outer region of C_j .

A graph G contains a graph H as a *minor* if G can be modified to H by a sequence of edge contractions, edge deletions and vertex deletions. Note that a graph G contains a graph H as a minor if and only if G contains a subgraph that contains H as a contraction. The *subdivision* of an edge $e = uv$ in a graph G removes e from G and replaces it by a new vertex w that is made adjacent to u and v . A *subdivision* of a graph G is a graph obtained from G after performing a sequence of edge subdivisions. In Figure 1, three examples of an *elementary wall*

are given. The unique cycle that forms the boundary of the outer face is called the *perimeter* of the wall. A *wall* W of height h is a subdivision of an elementary wall of height h and is well-known to have a unique planar embedding. We also call the facial cycle of W corresponding to the perimeter of the original elementary wall the *perimeter* of W , and we denote this cycle by $P(W)$.




Fig. 1. Elementary walls of height 2, 3, and 4 with perimeters of length 14, 22, and 30, respectively.

The $r \times r$ *grid* has all pairs (i, j) for $i, j = 0, 1, \dots, r - 1$ as the vertex set, and two vertices (i, j) and (i', j') are joined by an edge if and only if $|i - i'| + |j - j'| = 1$. The *side length* of an $r \times r$ grid is r . A well-known result of Robertson and Seymour [19] states that, for every integer r , any planar graph with no $r \times r$ grid minor has treewidth at most $6r - 5$. Although it is not known whether a largest grid minor of a planar graph can be computed in polynomial time, there exist several constant-factor approximation algorithms. In our algorithm we will use one by Gu and Tamaki [9]. For any graph G , let $\text{gm}(G)$ be the largest integer r such that G has an $r \times r$ grid as a minor. Gu and Tamaki [9] showed that for every constant $\epsilon > 0$, there exists a constant $c_\epsilon > 3$ such that an $r \times r$ grid minor in a planar graph G can be constructed in time $O(n^{1+\epsilon})$, where $r \geq \text{gm}(G)/c_\epsilon$. Because we can obtain a wall of height $\lfloor r/2 \rfloor$ as a subgraph from an $r \times r$ grid minor by deleting edges and vertices, their result implies the following theorem.

Theorem 1 ([9]). *Let G be a planar graph, and let h^* be the height of a largest wall that appears as a subgraph in G . For every constant $\epsilon > 0$, there exists a constant $c_\epsilon > 3$ such that a wall in G with height at least h^*/c_ϵ can be constructed in time $O(n^{1+\epsilon})$.*

In parameterized complexity theory, we consider the problem input as a pair (I, k) , where I is the main part and k the parameter. A problem is *fixed-parameter tractable* if an instance (I, k) can be solved in time $f(k)|I|^c$, where f denotes a computable function that only depends on k , and where c is a constant independent of k .

3 Fixed-Parameter Tractability of PLANAR CONTRACTION

For our algorithm we need the aforementioned result of Kawarabayashi [15].

Theorem 2 ([15]). *For every fixed integer k , it is possible to find in $O(n)$ time a set S of at most k vertices in an n -vertex graph G such that $G - S$ is planar, or conclude that such a set S does not exist.*

We also need the three following lemmas.

Lemma 1. *If a graph $G = (V, E)$ can be contracted to a planar graph by using at most k edge contractions, then there exists a set $S \subseteq V$ with $|S| \leq k$ such that $G - S$ is planar.³*

Proof. Let $G = (V, E)$ be a graph on n vertices that can be contracted to a planar graph H by using $k' \leq k$ edge contractions. Because each edge contraction reduces the number of vertices by exactly one, H has $\ell = n - k'$ vertices x_1, \dots, x_ℓ . Let \mathcal{W} be an H -witness structure of G . We write $|W(x_i)| = w_i$ for $i = 1, \dots, \ell$. Note that $w_1 + \dots + w_\ell = n$.

For each witness set $W(x_i)$, we arbitrarily remove a set S_i of $w_i - 1$ vertices from $W(x_i)$. We let $S = S_1 \cup \dots \cup S_\ell$. Because each witness set $W(x_i)$ has size 1 after removing the vertices of S_i , the graph $G - S$ is a spanning subgraph of H . Because H is planar and the class of planar graphs is closed under edge deletion, this means that $G - S$ is planar. Moreover, $|S| = w_1 - 1 + \dots + w_\ell - 1 = n - \ell = k' \leq k$. This completes the proof of Lemma 1. \square

When k is fixed, we write k -PLANAR CONTRACTION instead of PLANAR CONTRACTION. A seminal result of Courcelle [4] states that on any class of graphs of bounded treewidth, every problem expressible in monadic second-order logic can be solved in time linear in the number of vertices of the graph.

Lemma 2. *For every fixed integer k , the k -PLANAR CONTRACTION problem can be expressed in monadic second-order logic.*

Proof. By Kuratowski's Theorem (cf. [5]), a graph is planar if and only if it does not contain K_5 or $K_{3,3}$ as a minor, i.e., if it cannot be modified into K_5 or $K_{3,3}$ by a sequence of edge contractions, edge deletions and vertex deletions. Recall that a graph G contains a graph H as a minor if and only if G contains a subgraph that contains H as a contraction, i.e., that has an H -witness structure.

Let G be a graph and let $k \geq 0$ be an integer. We claim that G can be contracted to a planar graph by at most k edge contractions if and only if G contains a set of edges $\{e_1, \dots, e_{k'}\}$ for some $k' \leq k$ such that for all K_5 -witness structures and all $K_{3,3}$ -witness structures of all subgraphs of G , at least one of the edges in $\{e_1, \dots, e_{k'}\}$ is a witness edge.

First suppose that G can be contracted to a planar graph H by at most k edge contractions. Let $E' \subseteq E(G)$ be a set of at most k edges such that contracting all edges in E' transforms G into H . Suppose, for contradiction, that there is a subgraph F of G that has a K_5 -witness structure or a $K_{3,3}$ -witness structure \mathcal{W} for which none of the edges in E' is a witness edge. Then the graph F' that is obtained from F by contracting all edges in $E' \cap E(F)$ also has a K_5 -witness structure or a $K_{3,3}$ -witness structure. Since F' is a subgraph of H , this means

³ As an aside, we point out that the reverse of this statement is not true. For instance, take a K_5 and subdivide each of its edges $p \geq 3$ times. The resulting graph can be made planar by one vertex deletion, but at least $p - 1$ edge contractions are required.

that H contains K_5 or $K_{3,3}$ as a minor, contradicting the assumption that H is planar.

For the reverse direction, suppose that G contains a set of edges $\{e_1, \dots, e_{k'}\}$ for some $k' \leq k$ such that for all K_5 -witness structures and all $K_{3,3}$ -witness structures of all subgraphs of G , at least one of the edges in $\{e_1, \dots, e_{k'}\}$ is a witness edge. Let H be the graph obtained from G by contracting each of the edges $e_1, \dots, e_{k'}$, and let \mathcal{W} be an H -witness structure of G . We claim that H is planar. Suppose, for contradiction, that H is not planar. Then H contains a subgraph F' that has a K_5 -witness structure or a $K_{3,3}$ -witness structure \mathcal{W}' . We consider the subgraph F of G induced by the union of the H -witness sets $W(x)$ of \mathcal{W} with $x \in V(F')$. For each witness set $W'(a)$ of \mathcal{W}' we define the set $W^*(a)$ to be the union of the H -witness sets $W(x)$ of \mathcal{W} with $x \in W'(a)$. Then the sets $W^*(a)$ form a K_5 -witness structure or a $K_{3,3}$ -witness structure of F in which none of the edges $e_1, \dots, e_{k'}$ is a witness edge, contradicting our assumption on the set $\{e_1, \dots, e_{k'}\}$.

We observe that for any fixed graph H , the property “having H as a subgraph” can be expressed in monadic second-order logic. Also “having an H -witness structure” can be expressed in monadic second-order logic, as we can express the properties “being connected” and “being adjacent” in monadic second-order logic. This completes the proof of Lemma 2. \square

It is possible that the following lemma or a close variant of it is already known in the literature. However, because we could not find a reference, we give its proof here.

Lemma 3. *Let B be a planar graph that has an embedding with two nested cycles C_1 and C_2 , such that C_1 is the boundary of its outer face and C_2 is the boundary of an inner face, and such that there are at least two vertex-disjoint paths that join vertices of C_1 and C_2 . Let I be a graph with $B \cap I = C_2$ such that $R = B \cup I$ is planar. Then R has an embedding such that C_1 is the boundary of the outer face.*

Proof. Let \mathcal{B} be a plane graph that is an embedding of B such that C_1 is the boundary of the outer face, and C_2 is the boundary of an inner face. Let \mathcal{R} be an embedding of R such that C_2 lies inside C_1 and, subject to this condition, the set of vertices and edges of \mathcal{R} that lie in the outer region of C_1 has minimum size. To simplify our arguments, we assume that each edge of \mathcal{R} that lies outside C_1 has at least one endvertex not on C_1 , i.e., that also lies outside C_1 ; otherwise we can subdivide each edge outside C_1 that has both its end-vertices in C_1 without loss of generality.

Let X be the set of vertices of \mathcal{R} that are in the outer region of C_1 . If $X = \emptyset$, then we have the desired embedding. Suppose that $X \neq \emptyset$. Let H be a connected component of $\mathcal{R}[X]$. Let u_1, \dots, u_k be the vertices of C_1 that are adjacent to H , and assume that the vertices u_1, \dots, u_k appear on C_1 in this order (see also Figure 2). Note that $k \geq 2$, as otherwise H could be embedded inside C_1 , contradicting the minimality of X .

We claim that H contains no vertex of I . In order to obtain a contradiction, suppose that $I \cap H$ contains a vertex s . Because H is connected and every u_i has a neighbor in H by definition, the subgraph of \mathcal{R} induced by $V(H) \cup \{u_1, \dots, u_k\}$ is connected. Hence it contains a path from s to a vertex in $B - C_2$, namely u_1 (or any other vertex u_j). This path contains no vertex of C_2 . Consequently, it must contain an edge between a vertex of $I - C_2$ and a vertex of $B - C_2$. This is not possible, because $B \cap I = C_2$.

Now let Y be the set of vertices and edges of \mathcal{R} that lie in the inner region of C_2 . Let \mathcal{B}' be the plane graph obtained from \mathcal{B} after removing all vertices of $X \setminus V(H)$ together with their incident edges and all elements of Y . If we remove all vertices of H from \mathcal{B}' , we obtain a plane graph \mathcal{B}'' . Because H contains no vertex of I , all its vertices belong to $B - (C_1 \cup C_2)$. Then, by the definition of \mathcal{B} , H must be embedded in the inner region of C_1 in \mathcal{B} , and hence also in \mathcal{B}' . Moreover, because there are at least two vertex-disjoint paths that join C_1 and C_2 , there exists a face in \mathcal{B}'' whose boundary contains a cycle $C \notin \{C_1, C_2\}$ such that all vertices and edges of H lie in the inner region of C in \mathcal{B}' . By the definition of H , the only vertices of B that are adjacent to vertices of H are the vertices u_1, \dots, u_k . As a result, cycle C contains the vertices u_1, \dots, u_k , and they appear in the same order on C as they do on C_1 . Moreover, C does not contain any vertex of H , as C is a cycle in \mathcal{B}'' , and \mathcal{B}'' does not contain any vertex of H by definition. Because C also does not contain any vertex from $(X \setminus V(H)) \cup Y$, we find that all vertices of C not on C_1 or C_2 lie between C_1 and C_2 in \mathcal{R} (see Figure 2 for an illustration).




Fig. 2. An example of a plane graph \mathcal{R} , in which H is the only connected component of $R[X]$. The cycle C is drawn as the bold (blue) cycle. The shaded area indicates the empty region D , in which the connected component H can be embedded.

Let f denote the inner region of C in \mathcal{R} . We say that a path in \mathcal{R} that joins two vertices of C and that lies in f is a *chord* of C . A chord P *separates* u_i and u_j if u_i and u_j are not on the boundary of the same inner face of $C \cup P$ in \mathcal{R} .

Claim 1. The graph \mathcal{R} has no chords separating u_i and u_j for all $1 \leq i < j \leq k$.

To obtain a contradiction, let x and y be the end-vertices of a chord P in \mathcal{R} that separates u_i and u_j for some $1 \leq i < j \leq k$. By definition, $x, y \in V(C)$. Because

P separates u_i and u_j , at least one of the vertices x, y does not belong to C_2 . We assume without loss of generality that x is not on C_2 . Hence, x is a vertex of $B - I$.

We claim that all vertices and edges of P belong to B . As a matter of fact, we prove something stronger by showing that all vertices of $P - \{y\}$ belong to $B - I$, and that y either belongs to $B - I$ or to C_2 . In order to see this, suppose that not all vertices of P belong to $B - I$. Let z be the first vertex on P that belongs to I when we start traversing P from x . Because $B \cap I = C_2$, we find that z is on C_2 . Recall that in \mathcal{R} all vertices inside C lie between C_1 and C_2 as shown in Figure 2. Hence $z = y$, which implies that y belongs to C_2 .

Because all vertices and edges of P belong to B , we find that P is embedded inside C_1 in \mathcal{B} by definition. However, then the connected subgraph H cannot be embedded in the inner region of C_1 in \mathcal{B} . This contradiction completes the proof of Claim 1.

For $k = 2$, Claim 1 immediately implies that the embedding \mathcal{R} of R can be modified in such a way that H is placed in f , as we can find an empty open disk D inside C , such that no vertex or edge of \mathcal{R} lies in D and the boundary of D contains u_1, \dots, u_k (see Figure 2 for an illustration). This contradicts the minimality of X . If $k \geq 4$, then there cannot exist a path in f that joins two vertices u_i, u_j such that $2 \leq |i - j| \leq k - 2$ as a result of Claim 1. Hence we can, like in the case $k = 2$, find an empty open disk D inside f , and obtain a new embedding of R by placing H inside D .

For the case $k = 3$, we need the following claim in addition; we can prove this claim by the same arguments as we used in the proof of Claim 1.

Claim 2. There is no vertex v in \mathcal{R} that is in f such that v is joined with u_1, u_2, u_3 by disjoint paths in f .

Claims 1 and 2 together imply that we can find a desired empty region D inside C in the case $k = 3$, just as in the cases $k = 2$ and $k \geq 4$. This completes the proof of Lemma 3. \square

We are now ready to present our main theorem, which shows that PLANAR CONTRACTION is fixed-parameter tractable when parameterized by k . At some places in our proof of Theorem 3 we allow constant factors (independent of k) to be less than optimal in order to make the arguments easier to follow.

Theorem 3. *For every fixed integer k and every constant $\epsilon > 0$, the k -PLANAR CONTRACTION problem can be solved in $O(n^{2+\epsilon})$ time.*

Proof. Let G be a graph on n vertices, and let k be some fixed integer. If G has connected components L_1, \dots, L_q for some $q \geq 2$, then we solve for every possible tuple (k_1, \dots, k_q) with $\sum_{i=1}^q k_i = k$ the instances $(L_1, k_1), \dots, (L_q, k_q)$. Hence, we may assume without loss of generality that G is connected. We apply Theorem 2 to decide in $O(n)$ time whether G contains a subset S of at most k vertices such that $G - S$ is planar. If not, then we return **no** due to Lemma 1. Hence, from now on we assume that we have found such a set S . We write $H = G - S$.

Choose $\epsilon > 0$. We apply Theorem 1 on the graph H to find in $O(n^{1+\epsilon})$ time a subgraph W of H that is a wall with height $h \geq h^*/c_\epsilon$, where h^* denotes the height of a largest wall in H and $c_\epsilon > 3$ is some constant.

Suppose that $h \leq \lceil \sqrt{2k+1} \rceil (12k+10)$. Then $h^* \leq c_\epsilon h \leq c_\epsilon \lceil \sqrt{2k+1} \rceil (12k+10)$, i.e., the height of a largest wall in H is bounded by a constant. Consequently, the treewidth of H is bounded by a constant [19]. Since deleting a vertex from a graph decreases the treewidth by at most 1, the treewidth of G is at most $|S| \leq k$ larger than the treewidth of H . Because k is fixed, this means that the treewidth of G is bounded by a constant as well. Then Lemma 2 tells us that we may apply Courcelle's Theorem [4] to check in $O(n)$ time if G can be modified into a planar graph by using at most k edge contractions.




Fig. 3. On the left, a schematic depiction of the wall W with height $h > \lceil \sqrt{2k+1} \rceil (12k+10)$ and the way the subwalls W_1, \dots, W_{2k+1} , each with height $12k+8$, are packed within W . On the right, a more detailed picture of a subwall W_i of height 8 in case $k = 0$. The bold blue edges indicate the perimeter of the smaller subwall W'_i of height 6.

Now suppose that $h > \lceil \sqrt{2k+1} \rceil (12k+10)$. We consider some fixed planar embedding of H . For convenience, whenever we mention the graph H below, we always refer to this fixed embedding. The wall W is contained in some connected component \tilde{H} of H , and we assume without loss of generality that all other connected components of H lie outside $P(W)$. Inside $P(W)$, we choose $2k+1$ mutually vertex-disjoint subwalls W_1, \dots, W_{2k+1} of height $12k+8$ that are packed inside W in $\lceil \sqrt{2k+1} \rceil$ rows of $\lceil \sqrt{2k+1} \rceil$ subwalls, such that vertices of distinct subwalls are not adjacent; see Figure 3. Inside each W_i , we choose a subwall W'_i of height $12k+6$ such that the perimeters of W_i and W'_i are vertex-disjoint; see Figure 3 for a depiction of W_i and W'_i in case $k = 0$. By definition, the inner region of $P(W_i)$ is the region that contains the vertices of W'_i , and the inner region of $P(W'_i)$ is the region that contains no vertex of $P(W_i)$. Note that the interiors of $P(W_i)$ and $P(W'_i)$ are defined with respect to (the fixed planar embedding of) the graph H . Hence, these interiors may contain vertices of H that do not belong to W , as W is a subgraph of H .

We now consider the graph G . Recall that $H = G - S$, and that G is not necessarily planar. Hence, whenever we speak about the interior of some cycle below, we always refer to the interior of that cycle with respect to the fixed planar embedding of H . For $i = 1, \dots, 2k + 1$, let $S_i \subseteq S$ be the subset of vertices of S that are adjacent to an interior vertex of $P(W'_i)$. Observe that the sets S_i are not necessarily disjoint, since a vertex of S might be adjacent to interior vertices of $P(W'_i)$ for several values of i . Also note that no vertex of S belongs to W , since W is a wall in the graph $H = G - S$. We can construct the sets S_i in $O(n)$ time, because the number of edges of G is $O(n)$. The latter can be seen as follows. The number of edges in G is equal to the sum of the number of edges of H , the number of edges between H and S , and the number of edges of $G[S]$. Because H is planar, the number of edges of H is at most $5|V(H)| \leq 5n$. Hence, the number of edges of G is at most $5n + kn + \frac{1}{2}k(k-1) = O(n)$ for fixed k .

We say that a set S_i is of *type 1* if S_i is non-empty and if every vertex $y \in S_i$ also belongs to some set S_j for $j \neq i$, i.e., every vertex $y \in S_i$ is adjacent to some vertex z that lies inside $P(W'_j)$ for some $j \neq i$; see Figure 4 for an illustration. Otherwise we say that S_i is of *type 2*. We can check in $O(n)$ time how many sets S_i are of type 1. We claim the following.

Claim 1. If there are at least $k + 1$ sets S_i of type 1, then (G, k) is a no-instance.

We prove Claim 1 as follows. Suppose that there exist $\ell \geq k + 1$ sets S_i of type 1, say these sets are S_1, \dots, S_ℓ . Then for each $i = 1, \dots, \ell$ we can define a K_5 -witness structure \mathcal{X}_i of a subgraph of G as follows. We divide the perimeter of W_i into three connected non-empty parts in the way illustrated in Figure 4. The vertices of each part will form a separate witness set of \mathcal{X}_i ; let us call these




Fig. 4. Two subwalls W_i and W_j , where the bold blue edges indicate the perimeters $P(W'_i)$ and $P(W'_j)$ of the smaller subwalls W'_i and W'_j . Vertex y is in S_i , since it is adjacent to an interior vertex x of $P(W'_i)$. If, for every $y \in S_i$, there is an edge between y and an interior vertex z of $P(W'_j)$ for some $j \neq i$, then S_i is of type 1. The three shaded areas indicate how the perimeter of W_i is divided into three non-empty parts, each forming a separate witness set of a K_5 -witness structure \mathcal{X}_i of a subgraph of G .

witness sets X_i^1, X_i^2, X_i^3 . Let H'_i be the subgraph of H induced by the vertices that lie inside $P(W_i)$ in H . The fourth set X_i^4 of the witness structure \mathcal{X}_i consists of all the vertices of the connected component of H'_i that contains W'_i . Let G'_i

be the graph obtained from G by deleting all the vertices of $P(W_i)$ and all the vertices that lie inside $P(W_i)$ in H , i.e., $G'_i = G - P(W_i) - \text{interior}_H(P(W_i))$. Let D be the connected component of G'_i that contains the perimeter $P(W)$ of the large wall W . It is clear that D contains all vertices that are on or inside $P(W_j)$ for every $j \neq i$. Hence, due to the assumption that S_i is of type 1, all vertices of S_i also belong to D . The fifth set X_i^5 is defined to be the vertex set of D . Let us argue why these five sets form a K_5 -witness structure of a subgraph of G .

It is clear that each of the sets X_i^1, X_i^2, X_i^3 is connected, and that they are pairwise adjacent. The set X_i^4 is connected by definition. The choice of the subwall W'_i within W_i ensures that X_i^4 is adjacent to each of the sets X_i^1, X_i^2, X_i^3 . Let us consider the set X_i^5 . By definition, X_i^5 is connected. Since X_i^5 contains the perimeter $P(W)$ of the large wall W and W_i lies inside $P(W)$, set X_i^5 is adjacent to X_i^1, X_i^2 and X_i^3 . Since S_i is of type 1 and hence non-empty, there is a vertex $y \in S_i$ that is adjacent to a vertex x that lies inside $P(W'_i)$ by the definition of S_i . We already argued that X_i^5 contains all vertices of S_i , so $y \in X_i^5$. Recall that \tilde{H} is the unique connected component of H that contains the wall W , and that all connected components of H other than \tilde{H} were assumed to lie outside $P(W)$ in H . Because x lies inside $P(W'_i)$, this means that x is in the connected component of H'_i that contains W'_i , implying that $x \in X_i^4$. Consequently, the edge between x and y ensures the adjacency between X_i^4 and X_i^5 .

We now consider the ℓ different K_5 -witness structures \mathcal{X}_i of subgraphs of G defined in the way described above, one for each $i \in \{1, \dots, \ell\}$. Let us see how such a K_5 -witness structure \mathcal{X}_i can be destroyed by using edge contractions only. Denote by E_i the set of edges of G incident with the vertices of $X_i^1 \cup \dots \cup X_i^4$ for $i = 1, \dots, \ell$. We can only destroy a witness structure \mathcal{X}_i by edge contractions if we contract the edges of at least one path that has its endvertices in different witness sets of \mathcal{X}_i and its inner vertices (in case these exist) not belonging to any witness set of \mathcal{X}_i . Clearly, such a path always contains an edge of E_i . Hence, in order to destroy \mathcal{X}_i , we have to contract at least one edge of E_i . Because the sets E_1, \dots, E_ℓ are pairwise disjoint by the construction of the witness structures \mathcal{X}_i , we must use at least $\ell \geq k + 1$ edge contractions in order to make G planar. Hence, G is a no-instance. This proves Claim 1.

Due to Claim 1, we are done if there are at least $k + 1$ sets S_i of type 1. Note that every step in our algorithm so far took $O(n^{1+\epsilon})$ time, as desired. Suppose that we found at most k sets S_i of type 1. Because the total number of sets S_i is $2k + 1$, this means that there are at least $k + 1$ sets S_i of type 2. Let S_i be a set of type 2. If S_i is non-empty, then S_i contains a vertex x that is not adjacent to an interior vertex of $P(W'_j)$ for any $j \neq i$, as otherwise S_i would be of type 1. Consequently, S_i is the only set of type 2 that contains x . Since $|S| \leq k$ and there are at least $k + 1$ sets of type 2, at least one of them must be empty. Without loss of generality, we assume from now on that $S_1 = \emptyset$.

We will now exploit the property that $S_1 = \emptyset$, i.e., that none of the vertices in the interior of $P(W'_1)$ is adjacent to any vertex of S . We define a *triple layer* as the perimeter of a wall with the perimeters of its two largest proper subwalls

inside, such that the three perimeters are mutually vertex-disjoint, and the *middle* perimeter is adjacent to the *outer* and *inner* perimeter; see Figure 5 for an illustration. We define a sequence of nested triple layers in the same way as we defined a sequence of layers in Section 2. Because W'_1 has height $12k + 6$, there exist two adjacent vertices u and v inside $P(W'_1)$, such that u and v are separated from the vertices outside $P(W'_1)$ by $2k + 1$ nested triple layers L_1, \dots, L_{2k+1} , i.e., u and v lie inside the inner perimeter of triple layer L_{2k+1} (see also Figure 5).




Fig. 5. The wall W'_1 of height $12k + 6 = 18$ in case $k = 1$. The black bold edge uv is separated from the vertices that lie outside W'_1 by $2k + 1 = 3$ nested triple layers. The three shaded areas indicate the sets Y_1 , Y_2 , and Y_3 .

Let G' be the graph obtained from G after contracting uv . The following claim shows that uv is an “irrelevant” edge, i.e., that uv may be contracted without loss of generality.

Claim 2. (G, k) is a yes-instance if and only if (G', k) is a yes-instance.

We prove Claim 2 as follows. First suppose that (G, k) is a yes-instance. This means that G can be modified into a planar graph F by at most k edge contractions. Let $E' \subseteq E(G)$ be a set of at most k edges whose contraction modifies G into F . Observe that we can contract the edges in E' in any order to obtain F from G . If $uv \in E'$, then we can first contract uv to obtain the graph G' , and then contract the other edges in E' to modify G' into the planar graph F . If $uv \notin E'$, then we first contract the edges in E' to modify G into F , and then contract the edge uv . This leads to a graph F' . Since planar graphs are closed under edge contractions, F' is planar. Moreover, F' can also be obtained from G' by contracting the edges in E' . We conclude that (G', k) is a yes-instance.

Now suppose that (G', k) is a yes-instance. This means that G' can be modified into a planar graph F' by at most k edge contractions. Let $E' \subseteq E(G')$ be a set of at most k edges whose contraction modifies G' into F' . Let F be the graph obtained from G by contracting all the edges of E' . We will show that F is planar as well.

Recall that $S_1 = \emptyset$, and that we defined $2k + 1$ triple layers L_1, \dots, L_{2k+1} inside $P(W'_1)$. Let $Q_i, Q'_i,$ and Q''_i denote the three perimeters in H that form the triple layer L_i for $i = 1, \dots, 2k + 1$, where Q_i is the outer perimeter, Q'_i the middle perimeter, and Q''_i the inner perimeter. Let Y_i be the set of all vertices of H that are in $Q_i \cup Q'_i \cup Q''_i$ or that lie in the intersection of the inner region of Q_i and the outer region of Q''_i , i.e., Y_i is the set of vertices that lie on or “in between” the perimeters Q_i and Q''_i in H ; see Figure 5 for an illustration. Because we applied at most k edge contractions in G' , there exists a set Y_i , for some $1 \leq i \leq 2k + 1$, such that none of its vertices is incident with an edge in E' . This means that L_i is a triple layer in F' as well. We consider a planar embedding of F' , in which Q''_i is in the inner region of Q'_i , and Q'_i is in the inner region of Q_i ; for convenience, we will denote this planar embedding by F' as well.

We will now explain how to apply Lemma 3. We define C_1 and C_2 to be the perimeters Q'_i and Q''_i , respectively. We define B as the subgraph of F' induced by the vertices that either are in $Q'_i \cup Q''_i$ or lie between Q'_i and Q''_i in F' . Here, we assume that B is connected, as we can always place connected components of B that do not contain vertices from $Q'_i \cup Q''_i$ outside Q'_i . Because Q'_i and Q''_i are perimeters of subwalls in H , and Q''_i is contained inside Q'_i , there exist at least two vertex-disjoint paths P_1, P_2 in H joining Q'_i and Q''_i using vertices of Y_i only. Because none of the vertices in Y_i is incident with an edge in E' , the two paths P_1, P_2 are also vertex-disjoint in F' , and consequently in B .

We now construct the graph I . Because F' is a contraction of G' , and G' is a contraction of G , we find that F' is a contraction of G . Let \mathcal{W} be an F' -witness structure of G corresponding to contracting exactly the edges of $E' \cup \{uv\}$ in G . Then we define I to be the subgraph of G induced by the union of the vertices of all the witness sets $W(x)$ with x on or inside Q''_i in F' . Just as we may assume that B is connected, we may also assume that the subgraph of F' induced by the vertices that lie on or inside Q''_i is connected. Because witness sets are connected by definition, we then find that I is connected.

Because the edge uv is contracted when G is transformed into F' , u and v belong to the same witness set of \mathcal{W} . Let x^* be the vertex of F' , such that u and v are in the witness set $W(x^*)$. Recall that all the vertices of Q''_i and the vertices u and v belong to the wall W'_1 . Since u and v lie inside Q''_i in H and walls have a unique plane embedding, x^* lies inside Q''_i in F' . Hence, u and v are vertices of I . Also recall that none of the vertices of Y_i , and none of the vertices of Q'_i in particular, is incident with an edge of E' . Hence, the vertices of Q''_i correspond to witness sets of \mathcal{W} that are singletons, i.e., that have cardinality 1. This means that we can identify each vertex of Q''_i in F' with the unique vertex of G in the corresponding witness set. Hence, we obtain that $B \cap I = Q''_i = C_2$.

We now prove that $R = B \cup I$ is planar. For doing this, we first prove that B contains no vertex x with $W(x) \cap S \neq \emptyset$, and that I contains no vertex from S . To see that B contains no vertex x with $W(x) \cap S \neq \emptyset$, assume that x is a vertex of B and s is a vertex of S with $s \in W(x)$. Recall that no vertex from Q'_i is incident with an edge in E' . Hence, we can identify each vertex in Q'_i in F' with

the unique vertex of the corresponding witness set, just as we did earlier with the vertices of Q_i'' . Because $s \in W(x)$, this means that x is not in Q_i' . Because B is connected, we find that F' contains a path from x to a vertex y in Q_i' that contains no vertex from Q_i . Note that since y is in Q_i' , y is a vertex in G as well. Because $W(x)$ induces a connected subgraph of G by definition, this path can be transformed into a path in G from s to y that does not contain a vertex from Q_i . This is not possible, because $S_1 = \emptyset$ implies that every path in G from s to y must go through Q_i .

We now show that I contains no vertex from S . In order to obtain a contradiction, assume that I contains a vertex $s \in S$. Because I is connected, this means that G contains a path from s to a vertex in Q_i'' that contains no vertex from Q_i (and also no vertex from Q_i'). This is not possible, because $S_1 = \emptyset$.

Let R' be the subgraph of G induced by the vertices in the sets $W(x)$ with $x \in V(B)$ and the vertices of I . Since we proved that R' contains no vertices from S , R' is a subgraph of H . Consequently, R' is planar because H is planar. As a result, R is planar because R can be obtained from R' by contracting all edges in every set $W(x)$ with $x \in V(B)$, and planar graphs are closed under edge contractions. As we have shown that $R = B \cup I$ is planar, we are now ready to apply Lemma 3. This lemma tells us that R has an embedding \mathcal{R} , such that $Q_i' = C_1$ is the boundary of the outer face. Now consider the embedding that we obtain from the (plane) graph F' by removing all vertices that lie inside Q_i' . We combine this embedding with \mathcal{R} to obtain a plane embedding of a graph F^* . We can obtain F from F^* by contracting all edges in E' that are incident to a vertex in I ; recall that u and v are both in I and that uv is not an edge of E' . Because planar graphs are closed under edge contractions, this means that F is planar. This completes the proof of Claim 2.

We can find the irrelevant edge uv mentioned just above Claim 2 in $O(n)$ time. Since all other steps took $O(n^{1+\epsilon})$ time, we used $O(n^{1+\epsilon})$ time so far. After finding the edge uv , we contract it and continue with the smaller graph G' . Because removing S will make G' planar as well, we can keep S instead of applying Theorem 2 again. Hence, we apply Theorem 2 only once. Because G has n vertices, and every iteration reduces the number of vertices by exactly one, the total running time of our algorithm is $O(n^{2+\epsilon})$. This completes the proof. \square

4 Conclusions

We proved that PLANAR CONTRACTION is fixed-parameter tractable when parameterized by k . Very recently, Abello et al. [1] independently showed that the closely related problem that is to test whether a given graph can be made planar by contracting the edges of at most k mutually vertex-disjoint subgraphs, each of which of size at most ℓ , can be solved in quadratic time for any fixed k and $\ell \geq 2$. Their algorithm can easily be modified to show that k -PLANAR CONTRACTION can be solved in quadratic time for any fixed k (just as we can modify our algorithm to solve their problem).

A natural direction for future work is to consider the class \mathcal{H} that consists of all H -minor free graphs for some graph H and to determine the parameterized complexity of \mathcal{H} -CONTRACTION for such graph classes. Our proof techniques rely on the fact that we must contract to a planar graph, and as such they cannot be used directly for this variant. Hence, we pose this problem as an open problem.

References

1. J. Abello, P. Klavík, J. Kratochvíl and T. Vyskočil. Matching and ℓ -subgraph contractibility to planar graphs. Manuscript (2012) arXiv:1204.6070.
2. T. Asano and T. Hirata. Edge-contraction problems. *Journal of Computer and System Sciences* **26** (1983) 197–208.
3. A. Bouchet. Orientable and nonorientable genus of the complete bipartite graph. *Journal of Combinatorial Theory, Series B* **24** (1978) 24–33.
4. B. Courcelle. The monadic second-order logic of graphs. I. Recognizable sets of finite graphs. *Information and Computation* **85** (1990) 12–75.
5. R. Diestel. *Graph Theory*. Springer-Verlag, Electronic Edition, 2005.
6. R.G. Downey and M.R. Fellows. *Parameterized Complexity*. Springer, 1999.
7. P.A. Golovach, P. van ’t Hof and D. Paulusma. Obtaining planarity by contracting few edges. In: Proc. MFCS 2012, LNCS, to appear.
8. P.A. Golovach, M. Kamiński, D. Paulusma and D.M. Thilikos. Increasing the minimum degree of a graph by contractions. In: Proc. IPEC 2011, LNCS 7112, 67–79.
9. Q.-P. Gu and H. Tamaki. Constant-factor approximations of branch-decomposition and largest grid minor of planar graphs in $O(n^{1+\varepsilon})$ time. *Theoretical Computer Science* **412** (2011) 4100–4109.
10. P. Heggernes, P. van ’t Hof, B. Lévêque, and C. Paul. Contracting chordal graphs and bipartite graphs to paths and trees. In: Proc. LAGOS 2011, ENDM 37, 87–92.
11. P. Heggernes, P. van ’t Hof, D. Lokshtanov, and C. Paul. Obtaining a bipartite graph by contracting few edges, In: Proc. FSTTCS 2011, 217–228.
12. P. Heggernes, P. van ’t Hof, B. Lévêque, D. Lokshtanov, and C. Paul. Contracting graphs to paths and trees, In: Proc. IPEC 2011, LNCS 7112, 55–66.
13. M. Kamiński, D. Paulusma and D.M. Thilikos. Contractions of planar graphs in polynomial time. In: Proc. ESA 2010, LNCS 6346, 122–133.
14. M. Kamiński and D.M. Thilikos. Contraction checking in graphs on surfaces. In: Proc. STACS 2012, 182–193.
15. K. Kawarabayashi. Planarity allowing few error vertices in linear time. In: Proc. FOCS 2009, 639–648.
16. K. Kawarabayashi and B.A. Reed. Computing crossing number in linear time. In: Proc. STOC 2007, 382–390.
17. D. Marx and I. Schlotter. Obtaining a planar graph by vertex deletion. *Algorithmica* **62** (2012) 807–822.
18. D. Marx, B. O’Sullivan and I. Razgon. Finding small separators in linear time via treewidth reduction. Manuscript (2012) arXiv:1110.4765.
19. N. Robertson and P.D. Seymour. Quickly excluding a planar graph. *Journal of Combinatorial Theory, Series B* **62** (1994) 323–348.
20. N. Robertson and P.D. Seymour. Graph minors XIII: The disjoint paths problem. *Journal of Combinatorial Theory, Series B* **63** (1995) 65–110.
21. N. Robertson and P.D. Seymour. Graph minors XX: Wagner’s conjecture. *Journal of Combinatorial Theory, Series B* **92** (2004) 325–357.
22. N. Robertson and P.D. Seymour. Graph minors XXII: Irrelevant vertices in linkage problems. *Journal of Combinatorial Theory, Series B* **102** (2012) 530–563.