

Editorial Manager(tm) for Vegetation History and Archaeobotany  
Manuscript Draft

Manuscript Number:

Title: Neolithic human impact on the landscapes of North-East Hungary inferred from pollen and settlement records

Article Type: Original Article

Corresponding Author: Enikő Magyari, Ph.D.

Corresponding Author's Institution:

First Author: Enikő Magyari, Ph.D.

Order of Authors: Enikő Magyari, Ph.D.; John Chapman, Ph.D.; Andrew S. Fairbairn, Ph.D.; Mark Francis; Margarita de Guzman, Ph.D.

**Abstract:** In this article, we discuss the Neolithic and Early Copper Age (ECA) part of two pollen records from the Middle Tisza Floodplain in association with the local archaeological settlement record. We address the hypothesis of Willis and Bennett (2004) that there was little human impact by farmers on the environment of SE Europe until the Bronze Age. Contrary to this hypothesis, our results show that small-scale agriculture and woodland clearance is already attestable in the earliest Neolithic in Eastern Hungary, there are signs of expanding scale of mixed farming in the Middle Neolithic and very strong evidence for extensive landscape alterations with enhanced pasturing and mixed farming in the Late Neolithic (LN) and ECA. The main vegetation exploitation techniques in the alluvial plain of Sarló-hát were selective tree felling (mainly oak), coppicing (mainly hazel and elm), small-scale crop farming and woodland clearance to establish grazing pastures. Comparison with other well-dated pollen diagrams from Eastern Hungary suggested that, in the Early and Middle Neolithic (6000-5000 cal B.C.), hazel and elm coppicing were ubiquitous, while pastoral activities and associated woodland clearance distinguished the LN (5000-4500 cal B.C.). Comparison with regional and Northern Hemisphere paleoclimate proxy records suggested possible alignments between local environmental and Northern Hemisphere climatic fluctuations triggered by solar activity changes but, in several cases, climate change was attenuated by the buffering capacity of the floodplain (e.g. during the Szatmár II phase) or prevailed through transmissions modulated by precipitation and temperature changes in the watershed of the Tisza river (e.g. during the ECA). The interrelation between climatic and economic changes especially during the LN and ECA suggested a shift to moister summer conditions in the alluvium, which may contributed to decisions towards settlement dispersion and increased reliance on animal husbandry in the NE Hungarian Plain.

1 Enikő K. Magyari<sup>1</sup>, John Chapman<sup>2</sup>, Andrew S. Fairbairn<sup>3</sup>, Mark Francis<sup>4</sup>, Margarita  
2 de Guzman<sup>5</sup>  
3  
4

## 5 **Neolithic human impact on the landscapes of North-East** 6 **Hungary inferred from pollen and settlement records** 7 8 9

10  
11 <sup>1</sup>Hungarian Academy of Sciences - Hungarian Natural History Museum Research Group for  
12 Paleontology, 1476 Budapest, P. O. Box 222, Hungary

13 <sup>2</sup>Department of Archaeology, Durham University, South Road, Durham, DH1 3LE, United Kingdom

14 <sup>3</sup>School of Social Science, The University of Queensland, St Lucia, QLD 4072, Australia

15 <sup>4</sup> 10 Vara Drive, Belfast, Northern Ireland BT13 3BY

16 <sup>5</sup> Circle CRM Group Inc., 304, 716 - 3rd Avenue NW, Calgary, Alberta T2N 0J1, Canada  
17  
18

19 **Correspondent author:** Enikő K. Magyari, [magyari@bot.nhmus.hu](mailto:magyari@bot.nhmus.hu), tel.: +36705625614  
20  
21

### 22 **Abstract**

23 In this article, we discuss the Neolithic and Early Copper Age (ECA) part of two  
24 pollen records from the Middle Tisza Floodplain in association with the local  
25 archaeological settlement record. We address the hypothesis of Willis and Bennett  
26 (2004) that there was little human impact by farmers on the environment of SE  
27 Europe until the Bronze Age. Contrary to this hypothesis, our results show that small-  
28 scale agriculture and woodland clearance is already attestable in the earliest Neolithic  
29 in Eastern Hungary, there are signs of expanding scale of mixed farming in the  
30 Middle Neolithic and very strong evidence for extensive landscape alterations with  
31 enhanced pasturing and mixed farming in the Late Neolithic (LN) and ECA. The main  
32 vegetation exploitation techniques in the alluvial plain of Sarló-hát were selective tree  
33 felling (mainly oak), coppicing (mainly hazel and elm), small-scale crop farming and  
34 woodland clearance to establish grazing pastures. Comparison with other well-dated  
35 pollen diagrams from Eastern Hungary suggested that, in the Early and Middle  
36 Neolithic (6000-5000 cal B.C.), hazel and elm coppicing were ubiquitous, while  
37 pastoral activities and associated woodland clearance distinguished the LN (5000-  
38 4500 cal B.C.). Comparison with regional and Northern Hemisphere paleoclimate  
39 proxy records suggested possible alignments between local environmental and  
40 Northern Hemisphere climatic fluctuations triggered by solar activity changes but, in  
41 several cases, climate change was attenuated by the buffering capacity of the  
42 floodplain (e.g. during the Szatmár II phase) or prevailed through transmissions  
43 modulated by precipitation and temperature changes in the watershed of the Tisza  
44 river (e.g. during the ECA). The interrelation between climatic and economic changes  
45 especially during the LN and ECA suggested a shift to moister summer conditions in  
46 the alluvium, which may contributed to decisions towards settlement dispersion and  
47 increased reliance on animal husbandry in the NE Hungarian Plain.  
48  
49  
50  
51  
52  
53

54 **Keywords:** human impact, pollen analysis, archaeology, Neolithic, Great Hungarian  
55 Plain  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Introduction

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

Ever since the interpretation of plant taxa from pollen records in terms of anthropogenic effects, there has been a debate about the extent of the impact of prehistoric cultivation and/or animal keeping on the contemporary environment (Iversen 1941; Lüning and Kalis 1992). In SE Europe, an area which was decisive in the early spread of agriculture, Willis and Bennett (1994) for example argued that there was little human impact by farmers on the environment until the Bronze Age. The implication is that, despite the large number of Neolithic and Copper Age settlements in this region (Marinova 2007; Marinova et al. this issue), their subsistence practices were relatively small-scale and they had little overall impact on their environments. It is a matter of regret that, with a few recent exceptions (e.g., Ecsegfalva: Whittle *et al.* 2007, Willis 2007), all of the pollen sites investigated so far in Hungary lie some distance from key early farming sites with well-dated on-site evidence for mixed farming (e.g. Willis et al. 1998, Sümegi 1999; Magyari et al. 2001; Gardner 2002; Magyari et al. 2008). This is also true of our paper, since the archaeological sites in question have not been excavated yet. Nonetheless, intensive fieldwalking has revealed well-dated surface artifact scatters, suggesting localised occupation near the pollen coring sites. The first aim of this paper is to address Willis and Bennett's hypothesis through the integration of palynological and archaeological data in a poorly known semi-arid river basin of North-East Hungary.

The identification of human impact in long-term pollen sequences leads to a second aim in this paper – the diagnosis of the relative importance of anthropogenic and climatic factors in accounting for major phases of cultural change. In recent decades, archaeologists have been extremely sceptical about invoking environmental factors to account for cultural and social change (Renfrew 1984). A recent example from Eastern Hungary is Parkinson's (2006) attempt to explain the shift from Late Neolithic settlement nucleation on tells and large flat sites to Early Copper Age dispersed farmsteads in terms of Johnson's (1984) 'scalar stress' hypothesis. In part, this is due to the difficulties in differentiating the impact of anthropogenic from climatic variables in the interpretation of pollen sequences (Behre 1981; Berglund 2003) and there is also the residue of the reaction against environmental determinism characteristic of some processual archaeologists (Jones et al. 1999) and environmental scientists (Kertész and Sümegi 1999). But the main reason for mistrusting a simple equation between climatic change and cultural and social change is the multi-level intervening variables that must be taken into account in such an assessment. The most obvious intervening variables are the settlement structure and the strategies for obtaining food and drink and specific preferences for plant and animal species; the buffering potential of successful participation in exchange networks; the overall cultural values which promote a specific life-style; and, last but not least, the options which people may or may not have chosen from the wide range of available options when faced by potential climatic change. Attention to some of these variables can create a framework for studying detailed regional and local palaeo-environmental data; a full consideration of all the associated cultural and symbolic variables goes beyond the boundaries of this paper. Here, two pollen cores from the Sarló-hát meander (Figure 1) provide a fine-grained record that can be interrogated to help in assessing the significance of climatic and cultural factors over the period of the Neolithic and Early Copper Age in North East Hungary (6000 – 4000 cal B.C.).

1 The occupation sequence in this micro-region is set against the framework of the  
2 regional settlement pattern, as documented in the Upper Tisza Project (Chapman et al.  
3 2003, 2010a) and Hungarian investigations along the line of the M-3 Motorway  
4 (Hajdú and Nagy 1999; Raczky et al. 2007, 2011). The pollen cores are then used to  
5 determine local human impacts and wider regional climatic factors and to compare  
6 these results with the archaeological sequence. The paper concludes with an attempt  
7 to paint a broader picture of Neolithic and Early Copper Age subsistence through a  
8 comparison of the Sarló-hát diagrams with other coeval Hungarian pollen diagrams.  
9

## 10 11 **Materials and Methods**

### 12 *Study site*

13  
14  
15  
16  
17 The oxbow lake of Sarló-hát is situated 2 km northwest of the village of  
18 Tiszagyulaháza in the northern part of the Great Hungarian Plain (Figure 1).  
19 Geomorphological and palaeoenvironmental investigation of the area started under the  
20 aegis of an Anglo-Hungarian interdisciplinary project, the Upper Tisza Project  
21 (Chapman et al. 2003).  
22  
23

24 The *Sarló-hát meander* (Figure 1c) lies in an area that has been the focus of settlement  
25 since the Early Middle Neolithic, owing to its wide range of productive habitats  
26 suitable for farming, animal husbandry, hunting and gathering. From the meander,  
27 two undisturbed sediment cores were taken, one from the vicinity of a site dominated  
28 by Middle Neolithic discard (SH-WOOD, ca. 20 m from the shore), the other near a  
29 site with signs of intensive Bronze Age discard and less intensive Middle Neolithic  
30 discard (SH-II, ca. 90 m from the shore) (Figures 1 and 2).  
31  
32  
33

34 The study area is dominated by meadow soils (chernozems) on the Pleistocene  
35 elevated levees, and raw unstable soils on the floodplains (Chapman et al. 2003,  
36 2010a). Parts of the Pleistocene levee localities are covered by saline soils (Marosi  
37 and Somogyi 1990), whereas, in the northern part of the area, brown-earth developed  
38 in very few places.  
39  
40

41 From the standpoint of climate, the northern part of the study area is relatively humid  
42 (annual precipitation: 600 mm); however, towards the south (Újtikos, Polgár: Figure  
43 1) rainfall decreases abruptly, and in the Hortobágy hardly attains 500 mm (Szász and  
44 Tókei 1997). The mean annual temperature ranges 9.7-9.9 °C. The difference between  
45 the mean temperature of the coldest and warmest months is also the greatest here in  
46 the Alföld, 22-24 °C. The above parameters would suggest a vegetation cover  
47 intermediate between steppe and warm temperate woodland (Walter 1974); on the  
48 basis of climatic parameters, Zólyomi (1946) described the potential vegetation of the  
49 area as warm continental forest steppe.  
50  
51  
52

53 According to the results of geomorphological surveys, the Polgár landscape looked  
54 quite different in the Neolithic from today. Neotectonic subsidence and climate  
55 change over the last 30,000 years brought frequent and dramatic riverbed changes in  
56 this area (Borsy 1989; Tímár et al. 2005; Sümegi et al. 2005). The recently elevated,  
57 loess-mantled surfaces, where most Neolithic tells and settlements are found (Figure  
58 2) are located in the southern part of the study area (e.g. Polgár Island on Figure 1c).  
59  
60  
61  
62  
63  
64  
65

1 These are remnants of an ancient alluvial plain formed by the activity of NE-SW  
2 running rivers that drained the North Hungarian Mid Mountains and charged into the  
3 paleo-Tisza river that ran further south during the last glacial period (Sümegei et al.  
4 2005). These large channels (e.g. Hódos-ér and Kengyel-ér on Figure 1c) and their  
5 alluvium accumulated loess, on which fertile chernozem soils developed in the early  
6 Holocene (Chapman et al. 2010a). Following a major Neotectonic subsidence,  
7 sometimes between 20-30,000 years ago (Tímár et al. 2005), the Tisza river was  
8 relocated into this area, and its meandering channels incised into the former alluvial  
9 surface developing a floodplain that now lies at 90-91.5 m, i.e. 2.5-4 m below the Late  
10 Pleistocene lag surface (Gillings 1997). In this landscape four lateglacial and  
11 Holocene channel generations were identified (Magyari et al. 2010). The Holocene  
12 rivers laid down silty deposits on which stray Neolithic finds, but also some  
13 settlements, were found (Figure 2). Overall, in the Neolithic period, the Pleistocene  
14 lag-surfaces were already in an elevated position, well above the level of the seasonal  
15 floods that on the other hand inundated lower lying areas below 91.5-92 m (Gillings  
16 1998). The Sarló-hát meander is situated in the northern part of the Holocene  
17 alluvium, about 10 km away from the Polgár Island, but much nearer to a large,  
18 elevated Pleistocene sand and loess mantled hill near Tiszadob (1.5 - 2 km), where  
19 Middle and Late Neolithic discards are abundant, but so far tells were not found  
20 (Figure 2).  
21  
22  
23  
24

25 Using landscape morphological considerations, Sümegei et al. (2005) argued, that prior  
26 to the Neolithic occupation, the Pleistocene alluvial surfaces were covered by  
27 temperate wooded steppe and natural saline steppe communities in the ancient  
28 backswamp areas, with temperate gallery and floodplain forests probably on the  
29 Holocene alluvium. Since Neolithic occupation was mainly on the elevated surfaces,  
30 it is likely that Neolithic farmers/foragers/shepherds chose the naturally semi-open  
31 part of the landscape for settlement, where they cleared woodland stands. Little is  
32 known, however, about their impact on the alluvial forests. Our previously published  
33 paper from Sarló-hát suggested that this part of the Hungarian Plain supported oak  
34 wooded steppes throughout the Early and Middle Holocene and cultural steppe during  
35 the last 3,000 years (Chapman et al. 2009; Magyari et al. 2010). In the present paper,  
36 we examine how Neolithic land-use modified the species composition and spatial  
37 extent of the alluvial forests, and what sort of social practices (crop farming,  
38 gathering, coppicing, tree felling, burning) were established in the floodplain.  
39  
40  
41  
42  
43

#### 44 *Fieldwork, laboratory and numerical analyses*

45  
46 Overlapping core sequences were collected from two localities in the palaeochannel  
47 using hand-held Russian and Livingstone piston corers (Figures 1c and 2).  
48  
49

50 The lithostratigraphy was described using the system of Troels-Smith (1955) as  
51 modified by Aaby and Digerfeldt (1986) and the physical characteristics of the  
52 sediment were determined by calculating loss of weight on ignition (Heiri et al. 2001).  
53 Subsamples of 1 cm<sup>3</sup> taken at 4-8 cm intervals were processed for pollen and  
54 microscopic charcoal analysis following method A of Berglund and Ralska-  
55 Jasiewiczowa (1986), with *Lycopodium* spore tablets used to determine concentrations  
56 (Stockmarr 1971). Pollen identification was performed using Moore et al. (1992),  
57 Reille (1992, 1995, 1998), and Punt *et al.* (1984) for a detailed identification of  
58 Umbelliferae pollen types. For this plant family a cumulative curve is displayed on  
59  
60  
61  
62  
63  
64  
65

1 the pollen diagrams that include the following pollen types: *Oenanthe*, *Cicuta*,  
2 *Angelica*, *Apium*-type and *Pastinaca*-type. A minimum of 500 terrestrial pollen grains  
3 per slide was counted concurrently with charcoal particles using the point count  
4 estimation method of Clark (1982).

5  
6 AMS  $^{14}\text{C}$  dating was performed by the Poznan Radiocarbon Laboratory on eleven  
7 terrestrial plant macrofossils and one mussel shell from core SH-WOOD.

8  
9 Numerical analyses were performed using the routines within Psimpoll 4.10 (Bennett  
10 1997). Pollen data was zoned by optimal splitting of information content; the age-  
11 depth relationship was modelled using Bayesian modelling (for details see Magyari et  
12 al. 2010); and rarefaction analysis were performed on pollen percentage data.

### 13 14 15 16 *Estimation of anthropogenic disturbance on the natural vegetation*

17  
18 Human disturbance on the vegetation is often associated with increasing floristic  
19 diversity, with woodland clearance creating secondary habitats with rich herbaceous  
20 vegetation (Birks and Birks 1980). One way to make quantitative estimates of human  
21 impact on the natural vegetation is to apply diversity measures to fossil pollen spectra  
22 (Birks and Line 1992; Seppä 1998). We use rarefaction analysis for this purpose. All  
23 identified terrestrial pollen taxa were included and the basic pollen sum for  
24 calculations was set to  $E(T_{500})$ .

25  
26  
27  
28 Pollen percentages of herbaceous indicator taxa were also invoked to characterise the  
29 land-use of the non-forested area. The categories considered are crop field, fallow and  
30 footpath ruderal communities, wet and dry meadows (Table 1). Herbaceous pollen  
31 types were assigned to these categories of land use according to previous studies as  
32 follows: (i) the anthropogenic indicator-species approach of Behre (1981); (ii) the  
33 ecological indications in the Hungarian flora according to Ujvárosi (1957) and Fekete  
34 et al. (1997). Percentages of the indicator species were pooled in each land-use  
35 category. Table 1 also provides information on the occurrence of these non-arboreal  
36 pollen types in natural vegetation types of the Middle Tisza Plain. As is evident from  
37 this table, several pollen-types considered as anthropogenic indicators are elements of  
38 the natural floodplain vegetation. The most problematic taxon is Chenopodiaceae.  
39 Species of this family grow preferably in floodplain ruderal mudflat communities and  
40 on saline lake bottoms (Table 1, Fekete et al. 1997). The former habitat appears in the  
41 seasonally flooded alluvium, where floodwater pools persist until mid-summer  
42 (Fekete et al. 1997). In the shallow nutrient rich water cyanobacteria and green algae  
43 produce large biomass (*Zygnema* and *Pediastrum* species) and after desiccation, the  
44 nutrient rich substrate becomes covered by fast growing annual ruderals (mainly  
45 *Chenopodium* and *Polygonum* species). Since we suspect that in the lakeshore SH-  
46 WOOD pollen record and possibly also in the lake-central SH-II pollen record  
47 Chenopodiaceae pollen was mainly derived from these natural communities during  
48 the Neolithic (see further discussion below), we excluded this taxon from the  
49 anthropogenic indicator list (Table 1).

### 50 51 52 53 54 55 56 *Spatial representation of the pollen and microcharcoal records*

57  
58 Modelling and empirical studies (Sugita, 1993, 2007ab; Soepboer et al., 2007)  
59 indicate that for a lake such as Sarló-hát (elongated oxbow lake with 200–240 m  
60  
61  
62  
63  
64  
65

1 diameter today and ca. 960 m diameter mire with a central island in the past, see  
2 Figure 1b), the correlation between pollen abundances and vegetation composition is  
3 not improved by considering vegetation more than 400–600 m from the lake. The  
4 regionally ‘uniform’ background pollen component, representing vegetation between  
5 600 m and tens of kilometres from the lake, accounts for ca. 45% of the total pollen  
6 (Soepboer et al. 2007). The Sarló-hát pollen data thus provide an integrated  
7 palaeovegetation record for the landscape around the lake and the surrounding region,  
8 with pollen from extra-local (20 m - 2 km) and regional sources (2 km - 200 km)  
9 predominating. Assuming a prevailing northerly wind direction in the past, as at  
10 present, the regional and extra-regional components principally reflect the vegetation  
11 of the Taktaköz and southern North Hungarian Mid Mountains (Figure 2b). Pollen  
12 from local sources, up to tens of metres from the sampling point, will be better  
13 represented in the SH-WOOD lake margin core than in the central SH-II core.  
14  
15

16  
17 Charcoal contained in lake sediments is an important proxy for fire and has been  
18 extensively used as an indicator of past burning regimes (Tinner et al. 2005;  
19 Colomabroli et al. 2008). Microscopic charcoal records (particles between 20-180  
20  $\mu\text{m}$  in our case) can be representative of both local and regional signal of burning  
21 (Whitlock and Larsen 2001), and similarly to the empirical relationships with pollen,  
22 the diameter of the lake basin can influence the proportion of charcoal received from a  
23 regional or local source. Our microscopic charcoal record comes from the lake-  
24 marginal SH-WOOD core. This location is expected to receive a greater amount of  
25 input from the environment immediately around the basin, because the lake surface  
26 area is relatively small in this position relative to the edge area, which acts against  
27 large-scale aerial transport (Whitlock and Larsen 2001). Therefore, the microscopic  
28 charcoal record from SH-WOOD is expected to hold a strong local and extra-local fire  
29 history component beside the regional signal.  
30  
31  
32

### 33 34 *Archaeological fieldwalking*

35  
36 The fieldwalking in the Tiszagyulaháza area was carried out by staff and experienced  
37 students, using the tried and tested techniques of the Upper Tisza Project in the Polgár  
38 Block (Chapman et al. 2003, 2010a). This involved walking in parallel lines at 10m  
39 spacing across the main axes of ploughed fields. A total of 0.6 km<sup>2</sup> (Figure 1) was  
40 covered in this way in 12 person-days, indicating a mean of 20 person-days per sq.  
41 km., which shows moderately high fieldwalking intensity for such conditions. This  
42 spacing of fieldwalkers meant that not only were no ‘site scatters’ missed but that  
43 there was a high probability of the recording of single finds (or ‘off-site’ discard) near  
44 the sites.  
45  
46  
47

## 48 **Results**

### 49 50 *The Sarló-hát archaeological survey*

51  
52 Earlier systematic fieldwalking by the Upper Tisza Project led into the mapping of  
53 Neolithic sites and stray finds in the Polgár area. These data are presented on Figure 2  
54 and show that the earliest Neolithic discard belongs to the Early Middle Neolithic  
55 Szatmár II group dating between 7450-7250 cal B.P. (5500-5300 cal B.C.). Site and  
56 discard densities increase in the Middle Neolithic AVK (Alföld Linear Pottery)  
57 cultural phase (7250-6950 cal B.P., 5300-5000 cal B.C.), which is followed by  
58  
59  
60  
61  
62  
63  
64  
65

1 smaller number of large Late Neolithic sites belonging to the Tisza-Herpály-  
2 Csőszhalom (THCS) cultures (6950-6450 cal B.P., 5000-4500 cal B.C.) (Chapman  
3 1994; Raczky and Anders 2008). Despite the relatively low number of sites,  
4 population density was the highest in the Late Neolithic, as the nucleated horizontal  
5 settlement at Csőszhalom had in itself 2100-2600 inhabitants (Raczky et al. 2002). In  
6 the Early Copper Age (not shown on Figure 2), a small number of scattered sites  
7 (probably farmsteads) were found in the Polgár area suggesting settlement dispersion  
8 and decreasing human population (Chapman 1994).  
9

10 Our intensive, systematic fieldwalking around the Sarló-hát meander led to the  
11 discovery of five new sites on the lakeshore (Figure 3):  
12  
13

14 *Tiszagyulaháza 9* is a 0.4-ha multi-period site on the North slope of a low hill.  
15 Widespread and dense Middle Neolithic discard was found, with one 40 x 10m  
16 concentration, partially overlapping with daub and lithic concentrations; low-density  
17 Copper Age discard; undated lithics, daub and ground stone discard. No magnetic  
18 anomalies were located and no cultural material was found in the single soil core.

19 *Tiszadob 28* is a 0.4-ha multi-period site on the South slope of a low hill, 100m North  
20 of the SH-II pollen core. Widespread and dense Middle Neolithic (Tiszadob group)  
21 discard was found with one 40 x 40m concentration that matches one large and two  
22 small magnetic anomalies as well as three soil cores with signs of a cultural layer and  
23 the highest soil phosphate values; strong overlap with lithic and daub concentrations  
24 and some overlap with the ground stone discard; small low-density Copper Age  
25 discard, with Early Copper Age finds in one grid square.

26 *Tiszadob 29* had one Middle Neolithic sherd and 10 lithics in a scatter measuring 40 x  
27 40m. There was no evidence of a cultural layer in the single soil core and no magnetic  
28 anomalies were detected.

29 *Tiszadob 30* had a generally low-density discard. It is a 0.1-ha multi-period site on the  
30 South slope of a low hill with widespread Middle Neolithic discard but lacking sherd  
31 concentrations; small, high-density Copper Age discard, relating to the discovery of a  
32 perforated polished stone axe fragment and a lithic peak in the same grid square as the  
33 soil core, which showed evidence of daub fragments at depth; ground stone discard  
34 more widespread than on most other sites. No magnetic anomalies were recorded.

35 *Tiszadob 31* is a 0.5-ha multi-period site on the South slope of a low hill, 50m North  
36 of the SH-WOOD pollen core. It had small Early Middle Neolithic (Sztalmár II)  
37 discard in one grid square where the soil core shows signs of a cultural layer and  
38 possibly a pit; widespread but discontinuous Middle Neolithic (Tiszadob group)  
39 discard, with two sherd concentrations measuring 30 x 10m and 30 x 20m  
40 respectively, both concentrations match magnetic anomalies and soil cores with traces  
41 of a cultural layer and overlap with lithic concentrations; discontinuous, low-density  
42 Copper Age discard, including some Late Copper Age material, with a peak in one  
43 grid square; traces of a cultural layer in one soil core.

44 These findings may be interpreted as indications of diachronic changes in the intensity  
45 of occupation around the Sarló-hát meander. After a small and/or short-lived Early  
46 Middle Neolithic (Sztalmár II) discard, there was varied Middle Neolithic (Tiszadob  
47 group) discard on four of the five sites, with traces of cultural layers in the soil coring,  
48 magnetic anomalies, daub and lithic concentrations in two sites. Notable is the total  
49 absence of Late Neolithic ceramic discard, while two small low-density scatters were  
50 dated respectively to the Early and the Late Copper Age, with a cultural layer in the  
51 latter.  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



## *Sediment lithostratigraphy, radiocarbon dating and age-depth modelling*

Figure 4 displays the sediment stratigraphies, loss-on-ignition results from SH-II and SH-WOOD, uncalibrated  $^{14}\text{C}$  dates and the Bayesian age–depth model for the SH-WOOD core. This core was dated by 12 AMS  $^{14}\text{C}$  dates. Because no material suitable for  $^{14}\text{C}$  age determinations was found in core SH-II, the age–depth model for this core was estimated using age estimates from the SH-WOOD core for pollen stratigraphical features apparent in both diagrams and used to define LPAZ boundaries. Further details on the chronology are discussed in Magyari et al. (2010).

Four principal lithostratigraphic units were recognized in both cores (Figure 4). Fluvial sands forming the lower parts of both sequences were overlain by silty clay and fine silt layers, and the Neolithic/ Early Copper Age (ECA) sediment layers were dominated by claysilt in both cores with no sign of mottling suggesting that these layers were not affected by periodic ground water fluctuation. Sand lenses were also missing, indicating stable lake conditions and no direct flooding of the core locations during the Neolithic and ECA. Organic content was generally low in the lake-central SH-II core; typical values were around 7-10%, with a single peak of 14% at 6400 cal B.P. Notable is that, in SH-WOOD, increased organic content (12-13%) was detected between ca. 8350 and 7000 cal B.P. (279–234 cm) coinciding with the pre-Neolithic, Early and Middle Neolithic periods, followed by a return to lower values (8-9%) during the Late Neolithic and ECA. Similar changes were not seen in the SH-II core.

### *The pollen records*

The relative frequency distributions of selected terrestrial pollen types from the lakeshore SH-WOOD and lake-central SH-II profiles are plotted on Figures 5 and 6, while wetland and aquatic pollen types from SH-WOOD are shown in Figure 7. In order to aid the interpretation of the vegetation changes, Table 2 summarizes the pollen results by providing averages of the most characteristic arboreal and herbaceous pollen types for each Neolithic and Early Copper Age cultural period, as well as microcharcoal concentrations and palynological richness. Figure 8 shows cumulative pollen curves for the various land use categories according to Table 1. The following interpretation of the Neolithic and Early Copper Age vegetation changes are based on these results. Nine pollen assemblage zones have been defined in both pollen diagrams (see Magyari et al. 2010). The Neolithic period is part of the longest assemblage zone SH-5 (between 8400-5400 cal B.P.), which is characterised by frequent, small-scale pollen frequency changes, but is otherwise the longest Holocene period with stable oak- (*Quercus*), hazel- (*Corylus*) and elm- (*Ulmus*) dominated arboreal pollen assemblages and 30-40% herbaceous pollen (NAP) whose grasses (Poaceae), wormwoods (*Artemisia*) and chenopods (Chenopodiaceae) are the most prominent. The two pollen diagrams indicate that, in the early Holocene, the vegetation of the higher floodplain around the Sarló-hát meander consisted of alternating patches of mixed hazel-oak-elm (*Corylus-Quercus-Ulmus*) woodland and continental steppe vegetation (Figures 5 and 6; Magyari et al. 2010). Along the channels, gallery-forests rich in willow (*Salix*), ash (*Fraxinus excelsior/angustifolia*) and oak (*Quercus*) were characteristic. A major change in this vegetation took place around 8400 cal B.P., i.e. about 1000 years earlier than the first Neolithic settlers appeared in the region.

1 *Pre-Neolithic vegetation changes (8400 – 7950 cal B.P.; 6450 – 6000 cal B.C.)*

2  
3  
4 Changes in the pollen assemblages are prominent in both the terrestrial and aquatic  
5 records at 8400 cal B.P. (6450 cal B.C.). Most conspicuous is a shift from hazel  
6 (*Corylus*) to oak (*Quercus*) dominance in both pollen records, and a decrease in ash  
7 (*Fraxinus excelsior/angustifolia*) in the lakeshore SH-WOOD record (Figure 5).  
8 These imply a change from hazel-, oak- and ash-dominated forests to oak- and elm-  
9 (*Ulmus*) dominated forests in the floodplain. Relatively high arboreal pollen  
10 frequencies (70-75% AP) in both records furthermore suggest that the woodland cover  
11 probably reached its Early-Mid Holocene maximum, with oak-wooded steppes on the  
12 elevated Pleistocene levees ('Polgár Island' and 'Tiszadob Island' *sensu* Sümegi et al.  
13 2005) and oak-hazel-elm-ash forests in the seasonally flooded alluvium (Figures 1 and  
14 4). The combination of woodland compositional change and high microcharcoal  
15 concentrations between 8500-8200 cal B.P. (6550 – 6250 cal B.C.) suggests that  
16 woodland fires probably facilitated the spread of oak, as well as the small-scale  
17 temporary spread of beech (*Fagus*) (2.5% at 8100 cal B.P.) (6150 cal B.C.).  
18  
19  
20  
21

22 The herbaceous pollen flora showed different changes in the two pollen records. An  
23 increase in chenopods (Chenopodiaceae) and sorrel (*Rumex acetosa/acetosella*)  
24 characterised the lakeshore SH-WOOD, while cereal pollen was detected in nearly all  
25 samples after 8500 cal B.P. (6550 cal B.C.) in both records. Unfortunately, none of  
26 the aforementioned herbs can be treated as definitive indicators of human impact in  
27 this region. As discussed in Magyari et al. (2010), chenopods and sorrels can both live  
28 on the seasonally exposed bottoms of the palaeochannels in saline and ephemeral  
29 plant communities, but they are also characteristic plants of paths, disturbed land  
30 surfaces, crop fields and pastures (Behre 1981; Fekete et al. 1997). Wild cereals can  
31 also be natural constituents of oak wooded steppes (e.g. *Hordeum hystrix* and *H.*  
32 *secalinum*). Wild rye (*Secale sylvestre*) is also a typical natural constituent of open  
33 calcareous sand meadows in the Danube-Tisza Interfluve, but it does not appear in the  
34 less calcareous sand areas of the NE Hungarian Plain (Fekete et al. 1997). Overall, the  
35 herbaceous pollen flora can be indicative of both natural processes, i.e. enhanced  
36 seasonal fluctuation in the water-level and hence seasonally exposed lakeshore with  
37 ephemeral and saline vegetation, and anthropogenic landscape disturbance as foragers  
38 opened up clearings for more effective hunting and gathering. Taken together with  
39 changes in the wetland pollen record in this period, such as the decreasing abundance  
40 of open-water indicator green algae (*Pediastrum boryanum* and *P. simplex*), a rapid  
41 increase of lakeshore-dwelling galingale (*Cyperus*), floating fern (*Salvinia natans*)  
42 and redshank (*Polygonum persicaria*) and the increasing sediment organic content, we  
43 interpreted these changes as reflecting an enhanced seasonal lake-level fluctuation  
44 with high spring and early summer water table followed by midsummer lake  
45 desiccation and proliferation of ruderal mudflat communities and pioneer floodplain  
46 vegetation with *Cyperus fuscus* on the lakeshore (Magyari et al. 2010) rather than  
47 human disturbance of the terrestrial vegetation, although the latter cannot be ruled out.  
48 Palynological richness was stable in both records in this period (20-23) suggesting  
49 that the woodland compositional change did not affect the herb flora and was  
50 associated with minimal disturbance. In SH-WOOD a single peak in palynological  
51 richness is seen in the uppermost sample, at 7950 cal B.P. (27), this coincides with a  
52 decrease in AP:NAP ratios and may suggest woodland disturbance in the vicinity of  
53 this core.  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 *Early Neolithic vegetation changes (Körös Culture with no settlement evidence: 7950*  
2 *– 7450 cal B.P.; 6000 – 5500 cal B.C.)*  
3

4  
5 Although local settlement is not indicated by archaeological fieldwalking, the fall in  
6 arboreal pollen percentages and AP:NAP ratios between 7950 and 7450 cal B.P.  
7 (6000 – 5500 cal B.C.) in SH-WOOD suggests small-scale forest clearance in the  
8 vicinity of the meander. Hazel and oak were probably both felled, with the saw-  
9 toothed pollen curves implying periodic clearances already in this period. It is notable  
10 that microcharcoal concentrations decreased in this phase, suggesting that the  
11 woodland was not burnt but, rather, that trees were selectively removed. In the case of  
12 hazel, coppicing may also be assumed. Since similar changes are not detectable in the  
13 lake-central SH-II pollen record, we assume that forest disturbance was confined to  
14 the small hill directly North of the meander in the vicinity of SH-WOOD (Figures 1-  
15 3). The increasing abundance of chenopods (Chenopodiaceae) in this period in SH-  
16 WOOD, together with sedge (Cyperaceae) and galingale (*Cyperus*) pollen  
17 percentages, suggests low summer water levels and further lakeward expansion of the  
18 wetland macrophytes. The continuous nature of the cereal pollen curve in this period  
19 in SH-WOOD, with wheat (*Triticum*-type) pollen grains at the end of this phase,  
20 (Figure 8) may suggest small-scale crop farming after 7600 cal B.P. (5650 cal B.C.).  
21 The absence of cereal pollen types in SH-II supports the local source of the cereal  
22 pollen in SH-WOOD. Palynological richness values showed no change in SH-II  
23 relative to the pre-Neolithic period, suggesting no major disturbance in the vicinity of  
24 this core, in line with the pollen record. In SH-WOOD however, peak terrestrial  
25 pollen diversity values at 7900 cal B.P. were followed by a drop (from 27 to 18), and  
26 increased values followed afterwards. This may imply vegetation reorganization after  
27 a single disturbance event at 7900 cal B.P. Overall, evidence for vegetation  
28 disturbance and crop farming is strong in the SH-WOOD pollen record in contrast to  
29 the lack of sites dating to the Early Neolithic.  
30  
31  
32  
33  
34  
35

36 *Middle Neolithic vegetation changes (Szatmár II and Early AVK: 7450 – 7250 cal*  
37 *B.P.; 5500 -5300 cal B.C.)*  
38  
39

40 The burning of woodland is suggested by the continuous decrease in AP:NAP ratios  
41 and total arboreal pollen abundance in the first part of this short occupation phase in  
42 both records, that is associated with a microcharcoal concentration peak at 7280 cal  
43 B.P. (5330 cal B.C.) in SH-WOOD. The decrease of mainly hazel in both records is  
44 conspicuous and may suggest its selective felling or coppicing. Towards the end of  
45 this phase, a short term woodland recovery can be inferred from the total arboreal  
46 pollen, oak and ash pollen percentage increases in the vicinity of SH-WOOD.  
47 The appearance of cereal (*Triticum*-type and *Secale*) and crop-field weed pollen types  
48 especially in SH-II but also in SH-WOOD suggest small-scale crop farming in the  
49 vicinity of the meander (Figure 7).  
50  
51  
52

53 There is a further notable change in the wetland pollen and green algae records, where  
54 a small increase in *Pediastrum boryanum* and *P. simplex* indicates temporary  
55 increases in the lake-level (Figure 7). On the other hand, the Szatmár II cultural period  
56 also coincides with the spread of Redshank (*Polygonum persicaria*-type, likely *P.*  
57 *periscaria* or *P. hydrolapathum*) in SH-WOOD that is indicative of wet meadows and  
58 seasonally exposed mudflat communities on the lakeshore. Redshank is a natural  
59  
60  
61  
62  
63  
64  
65

1 element of the alluvial landscapes, but also a weed occurring in nitrogen-rich  
2 disturbed habitats and even in wet crop fields (Fekete et al. 1997). Its continuous  
3 pollen curve in SH-WOOD starting in the Szatmár II period is likely to reflect the  
4 natural succession of the landscape, but an anthropogenic spread cannot be ruled out.  
5 In line with the decreasing AP frequencies in SH-II, palynological diversity shows a  
6 prominent peak in this record (26) suggesting that episodic woodland clearance  
7 probably opened up the forest canopy allowing for the spread of herbs. It is notable  
8 that the pollen-inferred human impact on the vegetation is stronger in the vicinity of  
9 SH-II, even though Early Middle Neolithic (Szatmár II) discards were recovered in  
10 the vicinity of SH-WOOD (Dob 31, Figure 3).

11  
12  
13 *Middle Neolithic vegetation changes (Late AVK with strong archaeological evidence*  
14 *for occupation: 7250 – 6950 cal B.P.; 5300 – 5000 cal B.C.)*  
15

16  
17 The most remarkable change that can be associated with human impact is the increase  
18 of wormwood (*Artemisia*) pollen abundances in SH-WOOD, indicating the expansion  
19 of alluvial pasture. Wormwood pollen abundances do not increase in SH-II,  
20 confirming that it spread locally in the vicinity of SH-WOOD, where two late AVK  
21 sites were detected by fieldwalking (Dob 29 and 30 on Figure 3). Since most  
22 wormwood species living in the north-east Hungary today are unpalatable to cattle  
23 and sheep, and are also favoured by soil nitrogen enrichment (especially *A. vulgaris*,  
24 Ujvárosi 1957, Molnár 1996; Fekete et al. 1997), grazing increases their abundance in  
25 pastureland, and this is the probable explanation of their pollen increase in this phase.  
26 After a short recovery, the woodland cover decreased again during the late AVK  
27 occupation phase around SH-WOOD, but increasing AP% and AP:NAP ratios in SH-  
28 II suggest a woodland recovery in this part of the meander. High ash (*Fraxinus*  
29 *excelsior/angustifolia*) pollen frequencies in SH-WOOD probably indicate a  
30 secondary woodland succession following the clearance episodes. Somewhat  
31 surprisingly, palynological diversity decreased in this cultural phase in both pollen  
32 records (Figures 5 and 6), despite the evidence for woodland clearance and increasing  
33 meadow cover in the alluvium.  
34  
35  
36  
37  
38

39 *Late Neolithic vegetation changes (Tisza-Herpály-Csőszhalom group with major*  
40 *occupation in the Polgár Block but no archaeological evidence for occupation in the*  
41 *Sarló-hát area: 6950 – 6450 cal B.P.; 5000 – 4500 cal B.C.)*  
42

43  
44 Rye (*Secale*) pollen is present in SH-WOOD all-along in this cultural period that  
45 together with the occurrence of crop field indicator weeds likely indicate crop fields in  
46 the vicinity of the meander. Its co-occurrence with wheat (*Triticum*-type) and other  
47 cereal pollen suggests that rye pollen more likely derived from crop-fields than the  
48 natural steppes. Similarly to the AVK phase, the high values of wormwood  
49 (*Artemisia*) pollen in SH-WOOD point to the ongoing use of the floodplain for  
50 pasture. Moreover, the combination of decreasing AP:NAP ratios in SH-WOOD with  
51 a microcharcoal concentration peak at 6890 cal BP (4940 cal B.C.) suggests that  
52 selective felling and woodland burning were both frequent occurrences in the vicinity  
53 of the meander near SH-WOOD. Furthermore, several features of the SH-WOOD  
54 pollen record imply that the alluvium became wetter during the summer months. The  
55 gradual increase in Umbelliferae pollen abundances (with its characteristic  
56 components: *Oenanthe*, *Cicuta*-type, *Angelica* and *Pastinaca*-type) and the fall in  
57 organic content, sedge and galinagale pollen abundance all imply less severe seasonal  
58  
59  
60  
61  
62  
63  
64  
65

1 fluctuation in the water table and slightly decreasing productivity in the lake. These  
2 Umbelliferae forbs prefer wet meadows with a seasonally more stable water level than  
3 seasonal ruderals (Fekete et al. 1997; Magyari et al. 2010). It is notable that the SH-II  
4 pollen record does not indicate strong human impact on the vegetation in this cultural  
5 phase, in contrast with the SH-WOOD record. The most prominent feature is the  
6 increase of hazel (*Corylus*) pollen against oak (*Quercus*) that may indicate selective  
7 felling of oak trees. Palynological richness values are stable in SH-II, while they show  
8 a strong increase in SH-WOOD (Figure 5), supporting stronger vegetation disturbance  
9 in the vicinity of this core, although Late Neolithic discards were not found in the  
10 vicinity of Sarló-hát.

11  
12  
13 *Early Copper Age vegetation changes (Tiszapolgár group with limited archaeological*  
14 *evidence for occupation: 6450 – 5950 cal B.P.; 4500 – 4000 cal B.C.)*

15  
16  
17 The most remarkable change at the onset of this occupation phase is the rapid increase  
18 in pollen types indicating wet meadows, mainly in SH-WOOD. The probably rapid  
19 expansion of Umbelliferae forbs in the lakeshore suggests a further increase in  
20 summer water-table stability. The rise in hazel pollen percentages as compared to oak  
21 suggests either the cessation of Neolithic practices of woodland management (hazel  
22 coppicing, leaf foddering) or the selective felling of oak. Decreasing AP abundances  
23 are accompanied by a microcharcoal concentration peak at 6150 cal B.P. (4200 cal  
24 B.C.) in SH-WOOD, indicative of episodic woodland clearances by burning. That  
25 clearings were probably made in the high-floodplain zone can be inferred from the  
26 post-fire expansion of ash (*Fraxinus excelsior/angustifolia*) in SH-WOOD, probably  
27 indicating a secondary woodland succession following the disturbance event. The  
28 absence of Rye (*Secale*) in the cereal pollen and only sporadic wheat (*Triticum*-type)  
29 pollen occurrence in SH-II with arable weeds in both pollen records means that the  
30 evidence for crop fields in the alluvium in the Early Copper Age is ambiguous.  
31 Overall, our results suggest continuing human impact in the floodplain, with strong  
32 evidence for episodic woodland clearances and a probably climatic-induced change in  
33 the wetland vegetation of the floodplain.  
34  
35  
36  
37  
38

## 39 Discussion

40  
41 *Contrasting the archaeological evidence for occupation with pollen-based inferences*  
42 *on human impact on the alluvial vegetation*

43  
44  
45 *The early – middle 6<sup>th</sup> millennium cal B.C.: the first farmers in North East Hungary?*

46  
47 The archaeological record from the Polgár area provided no evidence so far for  
48 occupation in the Earliest Neolithic (Körös / Criş groups) (Chapman 1994, Chapman  
49 et al. 2010a, Raczky et al. 1994, 2008, 2011). The pollen record, on the other hand,  
50 hinted at episodic woodland clearances and selective felling or coppicing of hazel in  
51 this period (7950-7450 cal B.P.; 6000 – 5500 cal B.C.) and *Triticum*-type pollen was  
52 also recorded suggesting small-scale crop farming (Figure 8). So the two records are  
53 seemingly in discrepancy with each other for the Earliest Neolithic. Although  
54 hypothetical, this paradox can be explained in several ways. One would be the  
55 burying of small Körös sites by alluvial deposits or their reduced artefact discard  
56 related to short-lived or seasonal occupations. Körös sites are usually found in  
57 elevated relic channel banks or natural levees (Nandris 1970, Kosse 1979, Sherratt  
58  
59  
60  
61  
62  
63  
64  
65

1 1982/3; Gillings 2007). However the recent discovery of several small sites in the  
2 north GHP on less elevated terrain (Domboróczki 2005; Domboróczki and Raczky  
3 2010) may suggest the possibility of hitherto undetected sites buried under alluvium  
4 or hillwash, even in survey blocks where there has been intensive systematic  
5 fieldwalking (e.g., Blocks 1 and 2 of the Upper Tisza Project: Chapman et al. 2010a,  
6 2010b).

7 The second possibility is that some Szatmár II sites may date earlier than suggested by  
8 Hertelendi et al. (1995). In this respect the radiocarbon dates of recently discovered  
9 northernmost Körös sites (Tizzaszőlős-Domaháza-puszta and Ibrány-Nagyerdő) are  
10 important. They date around 7550-7450 cal B.P. (5600 – 5500 cal B.C.)  
11 (Domboróczki 2005; Domboróczki and Raczky 2010), indicating a chronological as  
12 well as a spatial overlap with Szatmár II populations. This overlap mirrors the  
13 overlaps between the Körös and the Szatmár II groups in the south Alföld, suggesting  
14 continuity in material culture development across the Early – Middle Neolithic  
15 transition.  
16  
17  
18

19 Finally, it is also possible that we see the impact of Mesolithic foragers between 7950-  
20 7450 cal B.P. (6000 – 5500 cal B.C.). In this respect, contrasting opinions have been  
21 formulated by Domboróczki (2005) and Sümegei and Kertész (1998). While the first  
22 author argues that the Mesolithic population was too small to exert any influence on  
23 the southern Körös sites and probably also on the natural vegetation, the second two  
24 authors suggested that a relatively large Mesolithic population lived north of the river  
25 Körös, whose environmental impact is detectable in several East Hungarian pollen  
26 records (Sümegei 2005). According to the interpretation of Sümegei (2005) and Juhász  
27 (2007) (cf. Zvelebil 1994), Mesolithic hunter-gatherers regularly burnt the early  
28 Holocene mixed forests down to create hunting trails and clearings, as well as  
29 promoting the spread of hazel and other edible species. A similar interpretation was  
30 provided by Kuneš et al. (2008) in the Czech Republic on the basis of a strong  
31 correlation between local hazel pollen peaks and Mesolithic intensity of occupation.  
32 However, if we look at the Sarló-hát pollen records (Figures 5 and 6), forest fires in  
33 fact decreased in the period under consideration, and hazel pollen declined  
34 periodically in this period, suggesting that the classical Mesolithic land-use strategy  
35 cannot be invoked to interpret anthropogenic vegetation changes. Overall, it is less  
36 likely that we see the impact of Mesolithic groups here in comparison with undetected  
37 Early Neolithic groups acting as triggers of the detected vegetation changes. However,  
38 it remains an open question whether the pollen-inferred vegetation disturbances in the  
39 Sarló-hát pollen records are attributable to early, undetected Körös groups or late  
40 hunter-gatherer communities who practised incipient cultivation, as in other areas of  
41 Europe (cf. Poska et al. 2004; Tinner et al. 2007).

42 In the subsequent Early Middle Neolithic Szatmár II phase, fieldwalking disclosed  
43 several stray finds in the vicinity of the Sarló-hát meander, pointing to low-intensity  
44 occupation on the lakeshore. This agrees well with the pollen-inferred evidence for  
45 increasing woodland clearance at 7400 cal B.P. (5450 cal B.C.) that probably  
46 involved 'slash-and-burn' clearance as reflected by a single microcharcoal peak in SH-  
47 WOOD. We found cereal pollen and crop field indicator weeds in SH-II, which lies  
48 further from the registered Szatmár II sites than SH-WOOD (Tiszadob 31; Figure 3).  
49 Thus, the pollen and archaeological records support each other and refer to the  
50 likelihood of small-scale crop-farming on the small hill 'Renyő-hát' that lies North-  
51 East of the study site (Figure 1). This is an important result for the Szatmár II group,  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 whose settlement traces are often slight since a high residential mobility is implied by  
2 the small number of small, widely-spaced sites (Chapman et al. 2003).

### 3 *The late 6<sup>th</sup> millennium cal B.C.: changes in farming practices*

4  
5  
6 Middle Neolithic AVK groups using Tiszadob-style pottery were the most abundant  
7 in the alluvial zone around Sarló-hát. Discard was found on four out of the five sites,  
8 all suggesting permanent occupation or seasonal activities near the floodplain zone. It  
9 is notable that both pollen records suggest a short oak woodland recovery between the  
10 Szatmár II and AVK occupation phases. This is followed by a clear AVK-associated  
11 oak clearance, periodic hazel coppicing and woodland burning that probably favoured  
12 the temporary spread of ash (*Fraxinus excelsior/angustifolia*) and dewberry (*Rubus*).  
13 Since ash is a typical re-sprouter and pioneer tree in the floodplain woodlands, its  
14 spread indicates woodland recovery after clearance (Fekete et al. 1997). Overall, the  
15 increased evidence for human modification of the floodplain zone for wetland pasture  
16 is well matched by the increased exploitation of the oak woodlands.  
17  
18  
19

20  
21 Middle Neolithic (AVK) settlement represents the peak of dispersion in the Neolithic  
22 (Figure 3), with a larger number of smaller sites than in the Early Neolithic of  
23 southern Hungary (Sherratt 1982/3; Makkay 1982a). The discard traces around the  
24 Sarló-hát lake are typical of the smaller range of AVK settlement clusters found in the  
25 Békés County surveys of the Hungarian Archaeological Topography (Ecsedy et al.  
26 1982, Jankovitch et al. 1989, Chapman 1997). What is interesting is that there is a  
27 stronger pastoral than arable signal in the AVK segment of the pollen cores – perhaps  
28 an indication of an early emphasis on cattle and caprine husbandry in alluvial pastures  
29 but also with cereal cultivation in a mixed farming system. The inference of an AVK  
30 manuring scatter from field survey evidence in Multi-Community Zone 12, near the  
31 village of Timár, in the Upper Tisza Project Block II, (Chapman and  
32 Laszlovszky 1993; Chapman et al. 2010b) supports the idea of the integration of  
33 farming with a pastoralist (animal manure) input. The high residential mobility of  
34 AVK sites fits well with the notion of a low-intensity farming regime.  
35  
36  
37  
38

### 39 *The Late Neolithic – a boom in cattle husbandry?*

40  
41 The Late Neolithic is a period of maximum settlement nucleation in Eastern Hungary  
42 (Makkay 1982b; Kalicz and Raczky 1989; Parkinson 2006). Parkinson (2006: 139 –  
43 156) characterises the settlement pattern of this period as a small number of clusters  
44 of tightly grouped sites in a river valley, with sites grouped around a central ‘super-  
45 site’ (open site or tell). These clusters were separated by large expanses of seemingly  
46 ‘unoccupied’ territories, often with just as high a land-use potential as the settled  
47 areas. It appears at first sight that Sarló-hát lay in one such unoccupied tract, since  
48 there was discard of not a single sherd of Late Neolithic pottery. Some 10km to the  
49 South lay the tell and horizontal settlement of Polgár-Czósshalom, acting as the local  
50 ritual centre for a Late Neolithic cluster, at the heart of an extensive exchange network  
51 based on limnoquartzite, rocks for polished stone tools, pottery and some obsidian  
52 (Raczky et al. 2002, 2007; Chapman et al. 2010a).  
53  
54  
55  
56

57 Despite the lack of evidence for local sites, the lakeshore SH-WOOD pollen profile  
58 provided evidence for a major change in the scale of floodplain exploitation. The  
59 expansion of floodplain, lakeshore pastures shows more intensive pastoralist  
60  
61  
62  
63  
64  
65

1 clearances than in the AVK phase. Open vegetation probably replaced some of the  
2 lakeshore oak-ash-elm woods and must have served for grazing. The creation of  
3 floodplain pastures probably reflects improved land management in this phase, but it  
4 is also conceivable that cooler and wetter summers in the Late Neolithic mitigated  
5 against cultivation, making more viable the practice of extensive pasturing, which is  
6 reflected in deforestation and grassland and meadow expansion in the Sarló-hát pollen  
7 diagrams. A similar shift towards extensive pasturing was observed and connected to  
8 climate change in several Neolithic pollen records in NW Europe (Berglund 2003).  
9 On the other hand, the archaeological and archaeobotanical records from Polgár  
10 provide evidence for technological improvement, as a greater diversity of crops and  
11 more integrated agro-pastoral strategies were found (Fairbairn 1992, 1993; Raczky et  
12 al. 2011). In addition, crop field indicator weeds and cereal pollen representing some  
13 of the species found in the Late Neolithic plant assemblage at Csőszhalom (Fairbairn  
14 1992) were found in the SH-WOOD profile (see below), probably reflecting the  
15 increasing area covered by crop-fields and open steppe-like disturbed vegetation.  
16  
17  
18

19 The discrepancy between the archaeological and the palaeo-botanical data at Sarló-hát  
20 is greatest in the Late Neolithic, given that this local area lay within one of the so-  
21 called ‘unoccupied’ territories between Late Neolithic settlement clusters. Two  
22 possible explanations may be proposed. The first explanation is that intensive,  
23 systematic survey has not been carried out for an area between Sarló-hát and the  
24 reasonably completely surveyed ‘Tiszadob Island’ (Chapman et al. 2003). It is thus  
25 possible that a small to medium-sized Late Neolithic connected to the mixed farming  
26 around Sarló-hát may yet be discovered South of the ‘Tiszadob Island’. By contrast,  
27 between Sarló-hát and Csőszhalom, but closer to the former, lay two small Late  
28 Neolithic sites – a very small (20 x 20m) high-density cluster (Tiszagyulaháza 002)  
29 and a low-density scatter covering 0.5 ha (Tiszagyulaháza 005). One further low-  
30 intensity Late Neolithic scatter lay North of Sarló-hát on the ‘Tiszadob Island’ –  
31 Tiszadob 001 (0.5 ha) (Chapman et al. 1993). Each of these unexcavated sites are  
32 consistent with small rural sites, perhaps dispersed ‘homesteads’ or even summer  
33 seasonal field huts, rather than major Late Neolithic nucleated sites. It is maybe from  
34 such sites that the extensive land-use attested in the Sarló-hát diagrams was  
35 conducted. Despite the lack of evidence for local Late Neolithic sites, the lakeshore  
36 SH-WOOD pollen profile provided evidence for a major change in the scale of  
37 floodplain exploitation. For the first time, the expansion of floodplain, lakeshore  
38 pastures shows more intensive pastoralist and arable clearances than in the AVK  
39 phase.  
40  
41  
42  
43  
44  
45

46 The second explanation was that there was not even limited, seasonal occupation in  
47 these zones – perhaps because of slightly moister summers causing a slight rise in  
48 Sarló-hát lake levels and longer stay of the seasonal floods in the alluvium. Instead,  
49 the area beyond the intensively cultivated ‘infield’ of the Csőszhalom tell was utilised  
50 first for a limited amount of (apparently manure-free) cultivation and then, around  
51 Sarló-hát, as an extensive zone of pastoral land-use, with cowherds being careful not  
52 to discard Late Neolithic pottery, integrated with far more restricted arable cultivation.  
53 The three small Late Neolithic sites near Sarló-hát could have acted as subsidiary sites  
54 to the Csőszhalom tell in extensive subsistence practices. This proposal has the  
55 advantage of accounting for the pollen data. It is also consonant with the expanded  
56 scale of Late Neolithic agro-pastoralism and particularly the emphasis on cattle  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 husbandry, which was more noticeable on the horizontal ('domestic') site than on the  
2 Csőszhalom ('ritual') tell (Schwartz in Raczky et al. 2002).

3  
4 The extension of our considerations to the macro-territory of the Csőszhalom tell has  
5 implications for the social structure of the tell in the Late Neolithic, viz., the  
6 possibility that the substantial cattle herds pastured in the tell macro-territory (10 km  
7 radius) were owned by significant lineages, households or individuals living in the  
8 horizontal site, if not on the tell itself. It is impossible at this juncture to make a  
9 precise calculation of the size of these herds but an initial estimate has been  
10 attempted, based on the contribution of the pollen in the Csőszhalom macro-territory  
11 to the Sarló-hát diagram (Chapman et al. in prep.). Assuming the grazing intensity of  
12 one head of cattle per 1.2 ha, the use of 20% of a 3 km territory around the tell would  
13 have provided sufficient fodder for over 400 cattle. But if the territory is extended to a  
14 6 km radius, the herd size could have increased to over 1,800 head. The extra-local  
15 pollen for open grassland in the Sarló-hát diagram is consistent with large open areas  
16 in the tell macro-territory capable of supporting an extensive strategy of cattle herding  
17 by the 5<sup>th</sup> millennium BC. These estimates are reminiscent of Sherratt's answer to the  
18 question 'what did plains-dwellers provide in exchange for upland resources in the  
19 Late Neolithic?' – large herds of cattle, as mobile wealth on the hoof (Sherratt 1982).  
20 This scenario would fit well with the amount of exotic material deposited at  
21 Csőszhalom. The Sarló-hát pollen diagram is the first diagram located within the  
22 macro-territory of a Late Neolithic nucleated site and is therefore the first with  
23 evidence for the large-scale animal husbandry long suspected of the period (Bökönyi  
24 1959, 1974).  
25  
26  
27  
28  
29

### 30 *Copper Age dispersion and cattle husbandry*

31  
32 The subsequent Early Copper Age is represented by minor discard on the Sarló-hát  
33 meander shores. These sites probably represent the high residential mobility typical  
34 for much of the Early Copper Age. In this phase, there was a larger number of  
35 settlement clusters, which were separated from each other by smaller tracts of land  
36 (Parkinson 2006). While the ECA settlement clusters covered larger areas, there were  
37 more sites in a cluster and the sites were smaller than in the Late Neolithic.  
38 The pollen record suggested an even moister alluvium than in the Late Neolithic, with  
39 a less pronounced summer water table fluctuation. We inferred episodic, fire-  
40 generated woodland clearances as the main impact on the alluvial vegetation, while  
41 decreasing wormwood (*Artemisia*) pollen frequencies and increasing Umbelliferae  
42 pollen percentages suggested a change in the composition of the floodplain pastures.  
43 There was a reduction in arable markers such as weeds of cultivation, which probably  
44 indicates a reduction in the scale of farming near the Sarló-hát. What appears to  
45 represent continuity from the Late Neolithic is the continuing use of floodplain  
46 pastures, even though their scale of use, too, was slightly reduced. The ECA changes  
47 in settlement structure at each integrative level, and in the direction of less complexity  
48 (Parkinson 2006), are likely to have resulted in changes in the scale of agro-pastoralist  
49 practices, whether in a major Early Copper Age cluster or in a re-occupied, small  
50 cluster, such as at Sarló-hát.  
51  
52  
53  
54  
55  
56

57 The scenario of major Late Neolithic cattle husbandry has implications for the Early  
58 Copper Age at Sarló-hát and more generally in the Hungarian Plain. Depending on the  
59 scale of Late Neolithic cattle husbandry, the move away from tell living and the  
60  
61  
62  
63  
64  
65

1 division of nucleated groups capable of managing large herds of cattle into individual  
2 households meant a re-alignment of dwelling practices with land-use. In contrast to  
3 the Late Neolithic combination of nucleated people and dispersed agro-pastoral land-  
4 use, Early Copper Age groups were dispersed over the landscape in much the same  
5 way as their cattle and fields. It is hard to be sure that there was a major decline in the  
6 scale of cattle-keeping in the later period – or even less social differentiation between  
7 ‘rich’ farms with large cattle herds and ‘poor’ farms with small herds.  
8

9  
10 *Neolithic subsistence and plant exploitation: comparisons between the pollen record*  
11 *and the Late Neolithic archaeo-botanical records of Polgár-Csőszhalom*  
12

13 During the excavations of two Neolithic sites in the Polgár area (Middle Neolithic  
14 Polgár 10: Gyulai 2010; Late Neolithic Csőszhalom tell and horizontal site; Fairbairn  
15 1992, 1993; Raczky et al. 2002; Gyulai 2010) rich archaeo-botanical material was  
16 recovered through flotation. These settlement data provide a detailed picture on the  
17 arable economies of the Middle and Late Neolithic occupation period in this area.  
18 Additional file 1 summarizes these findings, together with the equivalent pollen types  
19 in the Sarló-hát pollen records.  
20  
21

22  
23 On the basis of the diverse weed assemblages that included both dry- and wet ground  
24 taxa (e.g. *Papaver sp.*, *Stipa sp.*, *Solanum nigrum*, *Schoenoplectus lacustris*, *Juncus*  
25 *sp.*) Fairbairn argued that in the Middle Neolithic both dry- and wet-grounds were  
26 used for crop production (Fairbairn 1992). Both Fairbairn (1992) and Gyulai (2010)  
27 explained the dichotomy in the weed flora in terms of a risk management system,  
28 whereby the main crops were grown on the high floodplain to obtain high yields in  
29 ‘dry’ years, while a secondary crop was grown on the elevated lower yielding uplands  
30 (‘Pleistocene lag surfaces’ *sensu* Sümegei 2005) to ensure against total crop loss due to  
31 flooding of the alluvium. Fairbairn later slightly modified his conclusion concerning  
32 the risk-management system operating in the Late Neolithic and argued for the siting  
33 of fields along the Pleistocene riverbanks and elevated levees (Fairbairn 1993).  
34  
35  
36

37  
38 The Sarló-hát pollen records come from the high-yielding, but risky floodplain zone,  
39 and as we have pointed out above, they indicate small-scale crop farming in the  
40 alluvium during the Neolithic (mainly during the Szatmár II and Late Neolithic  
41 phases). We found low percentages of *Triticum* pollen (0.5-1%) accompanied by crop  
42 field weeds. This is in contrast with some alluvial sites in Poland and South-East  
43 Hungary, where near crop-field mires displayed 2-7% cereal pollen in the Neolithic  
44 layers (Wasytkova 1996; Willis 2007), but agrees with other alluvial sites, where crop  
45 fields were inferred to lie at a distance from the investigated meanders (Behre 2007).  
46 Our inference from this is that the majority of crop fields must have been located on  
47 the Pleistocene levees in the Polgár area, and the seasonally flooded alluvium was  
48 only opportunistically exploited for crop farming during the Neolithic. This inference  
49 accords with Fairbairn (1992) and Gyulai (2010), but the pollen data suggest only  
50 small-scale use of the alluvium for crop-farming. The relatively large proportion of  
51 wet-ground herbs in the settlement seed assemblages can probably be interpreted  
52 differently, as suggested by Kreuz et al. (2007) in connection with the Central  
53 European LBK sites. Accordingly, these wet-ground species could have been brought  
54 to the fields with the dung of cattle which grazed the floodplains as well as the  
55 harvested fields. Bogaard et al. (2008) also argued that, during the Early Neolithic of  
56 Hungary and the AVK, crop-fields were located near the settlements and managed  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 with intensive labour input to maintain fertility, while floodplain soils were avoided.  
2 But Bogaard's intensive, local horticultural model may not be appropriate to the Late  
3 Neolithic, in which extensive agro-pastoral practices are clearly attested. The higher  
4 summer lake-levels attested in the Late Neolithic segment of the Sarló-hát pollen core  
5 would also have increased the risk of spring planting. There is no definitive evidence  
6 yet that high-yield, high-risk strategy of floodplain farming was not attempted in the  
7 Late Neolithic at times of nucleated population; in any case, it is recognised that there  
8 was always the back-up of a low-risk, low-yield strategy on the higher areas. Overall,  
9 it seems that, even in the Late Neolithic, fields were probably located in different  
10 parts of the landscape – on the Pleistocene lag surfaces for security and in the  
11 alluvium for spectacular high-yield successes as well as major failures. Moreover,  
12 evidence for probably other kinds of exploitation of the floodplain vegetation comes  
13 from on-site recovery of typical wetland plants, e.g. dwarf elder (*Sambucus ebulus*),  
14 giant bulrush (*Schoenoplectus lacustris*), reed (*Phragmites australis*) and water  
15 dock (*Rumex hydrolapathum*) (Gyulai, 2010).  
16  
17  
18

19 Comparison of the Csőszhalom plant-macro-fossils with the Sarló-hát Late Neolithic  
20 pollen flora suggests considerable variability in species cultivated, weeds of  
21 cultivation and pastoral indicators (Additional file 1). Despite many common  
22 elements, quite a few crop plants and weeds are missing from the pollen spectra. For  
23 example, *Pisum*, *Lathyrus/Vicia*-type and *Linum* were grown by the Late Neolithic  
24 settlers but they fail to show up in the alluvial pollen record. This may be because  
25 they are insect-pollinated or low pollen producers, or because of different cultivation  
26 practices for different species. Conversely, *Plantago lanceolata* seeds were present in  
27 the weed flora at Csőszhalom, while this pollen type also appeared in the Late  
28 Neolithic segment of the Sarló-hát core, suggesting that the alluvium around the  
29 meander was disturbed.  
30  
31  
32

33  
34 An overall reason for differences in species representativity may have been the  
35 varying taphonomic filters that applied to the preservation of on-settlement plant  
36 macro-fossils and those responsible for creating the pollen assemblage. But what is  
37 interesting is that there are so many identical species at Csőszhalom and Sarló-hát,  
38 suggesting that, in the Late Neolithic at least, there were comparable agro-pastoral  
39 practices in place, linking persons on and off the tell in daily subsistence and ritual  
40 tasks.  
41  
42

### 43 *Neolithic human impact in Eastern Hungary: a regional comparison*

44  
45  
46 The eastern part of Hungary comprises two contrasting landscape types: the Great  
47 Hungarian Plain, that has always been dominated by its watercourses and varying  
48 water availability dependent on the distance from watercourses; and the North  
49 Hungarian Mid-Mountains with its closed deciduous forests since at least the early  
50 Holocene (Figure 1b; Willis et al. 1997). From these two landscapes, five well-dated  
51 pollen records offer possibility for comparison (Kismohos (Willis et al. 1998),  
52 Nagymohos (Magyari et al. 2001), Sirok (Gardner 2002), Bábtava (Magyari et al.  
53 2008) and Kiri-tó (Willis 2007), see Figure 1b), albeit only one of them (Ecsegfalva:  
54 Kiri-tó) comes from within 1 km of an excavated Early Neolithic site (Whittle 2007;  
55 Willis 2007). In the Neolithic period, all of these pollen records are characterized by  
56 hazel (*Corylus*) and oak (*Quercus*) dominance and notably all records show signs of  
57 Neolithic woodland management starting at slightly different dates. Figure 9 displays  
58  
59  
60  
61  
62  
63  
64  
65

1 the early and mid Holocene parts of two NE Hungarian pollen records, Nagymohos  
2 from the Putnok Hills and Báltava from the Bereg Plain, both area considered as a  
3 zone of low population density during the Neolithic (Sümegei 1999; Bánffy 2005),  
4 although the number of known archaeological sites in 50 km radius of both sites is  
5 high during the Middle Neolithic AVK and Bükk occupation phases (Bánffy 2005;  
6 Csengeri 2005). Moreover, recent excavations in the vicinity of Nagymohos  
7 recovered a small Middle Neolithic settlement of the Bükk culture, and painted  
8 pottery sherds probably from this settlement were found on the shores of Nagymohos  
9 (Csengeri 2005). Changes in these pollen records (starting at 7500 cal B.P. in  
10 Nagymohos and 7200 cal B.P. in Báltava) involved periodic declines in hazel  
11 (*Corylus*) and elm (*Ulmus*), coincident increases in oak (*Quercus*) and ash (*Fraxinus*  
12 *excelsior* type) and increasing representation of disturbance indicator herbs (e.g.,  
13 *Plantago lanceolata* at Nagymohos on Figure 9). These proxies were attributed to  
14 direct human activity (Magyari et al. 2001, 2008). Management of the woodland by  
15 coppicing of hazel, collection of leaves and young shoots of hazel and elm for leaf  
16 fodder and forest grazing were assumed. The similar saw-teeth shape of the hazel  
17 pollen frequency curve in the SH-WOOD pollen record (Figure 5) suggests a similar  
18 exploitation of hazel in the Körös and Szatmár II occupation phases (Early and Early  
19 Middle Neolithic) in the Upper Tisza Alluvium. Overall, it seems that woodland  
20 management can be traced in several pollen records in NE Hungary in the AVK  
21 occupation phase, and in case of Sarló-hát even earlier, since the start of the 6<sup>th</sup>  
22 millennium cal B.C. (ca. 8000 cal B.P.). In the Nagymohos pollen diagram, woodland  
23 management comes to a halt at ca. 6900 cal B.P. (4950 cal B.C.) (Figure 9), when  
24 increasing microcharcoal concentrations are accompanied by arboreal pollen  
25 decreases, suggesting forest clearance by burning. Similarly to SH-WOOD (Figure 5),  
26 wormwood (*Artemisia*) pollen frequencies show an episodic increase in the Late  
27 Neolithic, between ca. 6800-6500 cal B.P. (4850 – 4550 cal B.C.), pointing to a  
28 similar landscape management that likely involved more intensive grazing in the  
29 forest openings and woodland edges, but *Artemisia* can also indicate footpaths and  
30 fallows in the hill zone (Behre 1986; Fekete et al. 1997). Its mass occurrence was  
31 found somewhat earlier in SH-WOOD, between 7100-6700 cal B.P. (5150 – 4750 cal  
32 B.C.), but very characteristically it was also associated with forest clearance, and  
33 directly followed the woodland management phase. This suggests that this change in  
34 vegetation exploitation strategy is connectable with a technological and cultural  
35 change between the Middle and Late Neolithic. Delayed timing of both the woodland  
36 management and woodland clearance (to create grazing pastures) phases in the North  
37 Hungarian Mid Mountains is a clear feature of these pollen records, probably  
38 indicating a later spread of the Middle Neolithic AVK culture to the hills (represented  
39 by the Bükk culture) and also a delayed spread of the Late Neolithic groups and their  
40 subsistence strategy to the hill zone. This agrees with the archaeological interpretation  
41 of the Bükk culture as a late phase of the AVK (Kalicz and Makkay 1977). The  
42 settlement of the lower slopes of the Northern Mid Mountains from 4700 cal B.C., if  
43 not earlier (Raczky 1995; Pavúk 2007), by communities using Early Lengyel pottery  
44 means that the *Artemisia* rise in the Nagymohos pollen diagram may have reflected  
45 the grazing practices of these groups.  
46 It is important to note, however, that the Báltava diagram does not show an *Artemisia*  
47 increase in the Late Neolithic; instead, the woodland management phase is followed  
48 by an episodic spruce (*Picea*) and fern (Filicales) rise between 6800-6700 cal B.P.  
49 (4850 – 4750 cal B.C.), interpreted as a short-term cold episode or a by-product of  
50 recurrent fluvial activity (Magyari et al. 2008). Subsequently, human impact is weak  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 in this record during the Late Neolithic, in line with the local archaeological evidence  
2 for scarce occupation.

3  
4 Evidence for Late Neolithic and Copper Age woodland management was also found  
5 in the Sirok pollen profile derived from the Mátra Hills (Figure 1b). Here Gardner  
6 (2002) demonstrated periodically-reduced hazel pollen frequencies (72-288 years  
7 cycles) coincident with hornbeam (*Carpinus betulus*) pollen increases between 6900-  
8 5200 cal B.P. (4950 – 3550 cal B.C.) and interpreted these changes as reflecting  
9 regular coppicing or pollarding of hazel. The relatively late start of the woodland  
10 management agrees with the hinterland position of Sirok, but also provides evidence  
11 that coppicing was an important woodland exploitation method in the hill zone during  
12 the Late Neolithic, Early Copper Age and Middle Copper Age as well.

13  
14 Turning to the south-eastern part of the Hungarian Plain, the pollen record of Kiri-tó  
15 (Figure 1b) has special importance, as it is located adjacent to an Early Neolithic  
16 Körös settlement whose cultural origins lie to the South (Whittle 2007). The pollen  
17 record was dominated by oak and hazel during the Körös occupation phase (8000-  
18 7500 cal B.P.), and Willis (2007) argued that inhabitants of the very small settlement  
19 exerted little impact on the forest composition, not even clearing woodland pockets on  
20 the alluvium or the loess-mantled elevated surfaces because there was enough open,  
21 steppe-covered area suitable for farming. Although evidence for Early Neolithic  
22 occupation is missing in the Sarló-hát area, our pollen records suggested stronger  
23 local human impact on the alluvial forest via selective felling and/or coppicing of  
24 hazel, small-scale clearance, and crop farming after 7600 cal B.P. (5650 cal B.C.).  
25 The general impression is that the impact of the Early Neolithic groups on the  
26 vegetation was larger in the northern part of the plain, probably because of the higher  
27 proportion of woodland here that required clearance.

28  
29 Another important feature of the Kiri-tó pollen record was a significant reduction of  
30 arboreal pollen types and hence woodland cover after 7000 cal B.P. (5050 cal B.C.),  
31 that can be attributed to local AVK groups. As cereal pollen types also increased  
32 coincidentally (attaining 5%) without an increase in wormwood (*Artemisia*) pollen, it is  
33 reasonable to assume that clearance was made to establish new crop fields and not  
34 grazing pastures. This feature of the Kiri-tó pollen record again differs from Sarló-hát,  
35 where the main impact of the AVK groups was the establishment of alluvial grazing.  
36 However, it is also possible that these landscape exploitation differences can be  
37 explained by the different position of the pollen coring sites relative to the main  
38 settlements, being nearer in the case of Kiri-tó. Consequently, crop fields were  
39 implicitly nearer to the pollen core location of the latter.

40  
41  
42  
43  
44  
45  
46  
47 *Disentangling climatic and human factors in the Sarló-hát Neolithic pollen record:*  
48 *short-term climate changes in East-Central Europe and the global climatic record*  
49

50  
51 In this final part of the discussion, we make structured comparisons between the local  
52 proxy data, the Neolithic and Early Copper Age cultural phases of Eastern Hungary  
53 and the regional and global climatic trends, inferred from a range of climatic proxy  
54 records. Figure 10 shows these records. What is apparent for a first sight is the  
55 coincidence of the 8.2 ka cal B.P. cooling event (seen in the NGRIP  $\delta^{18}\text{O}$ , residual  $^{14}\text{C}$   
56 and PP10 stalagmite curves on Figure 10) with the organic content increase in SH-  
57 WOOD. Although the 8.2 event was associated with high lake levels in West-Central  
58 Europe due to the intensification of westerlies and lower summer temperatures  
59  
60

1 (Magny 2007), the East-Central European record of Lake St Ana (Magyari et al.  
2 2009) suggests low water-depths with a minimum at 8.1 ka cal B.P. (6150 cal B.C.),  
3 in comparison with hints from the Sarló-hát record at enhanced seasonal water-level  
4 fluctuation with low summer water tables during this event. We also found increased  
5 fire frequencies. On the whole, our data suggest that the 8.2 ka event was associated  
6 with increased summer drought in the Carpathian Basin, but the underlying climatic  
7 circulation changes are yet unresolved to explain the discrepancy with West-Central  
8 Europe. The spread of the Körös Culture into the Carpathian Basin does not coincide  
9 with this climatic perturbation, but directly follows it (~ 7950 cal B.P.: ~ 6000 cal  
10 B.C.). The organic content record from SH-WOOD suggests that the lake basin  
11 stabilized in a different form after the 8.2 ka event; we inferred that, during the Körös  
12 and Szatmár II occupation phases, high spring floods were associated with high water  
13 temperatures, followed by strong summer drought and desiccation of the lakeshore. It  
14 is notable that the onset of the Szatmár II occupation phase (7450-7250 cal B.P.;  
15 5500-5300 cal B.C.) coincides with a solar activity minimum (see the residual <sup>14</sup>C  
16 curve on Figure 10) and a concurrent high lake-level phase in West-Central Europe  
17 (Magny et al. 2003; Magny 2007). The SH-WOOD record also showed a minor  
18 increase in *Pediastrum* green algae (Figure 10) interpreted by us as a sign of  
19 increasing water-level, but the sustained high organic content and high *Cyperus* pollen  
20 percentages in the lake marginal SH-WOOD record suggested that the major trend of  
21 strong seasonal water level fluctuation persisted in this phase. The coincidence of a  
22 short-lived climatic change and a cultural change is nonetheless evident in this phase,  
23 and bivalve fauna based studies in the southern Tisza plain also support an  
24 environmental change at the onset of the early AVK, with stronger flood cycles  
25 leading into the dominance of fluvial bivalves on archaeological sites (Gulyás 2011).  
26  
27  
28  
29  
30

31 At the onset of the Late Neolithic Tisza-Herpály-Csőszhalom (THCS) occupation  
32 phase, a remarkable coincidence is seen between the cultural change and the organic  
33 content plus galingale (*Cyperus*) pollen decreases in SH-WOOD (Figure 10) that  
34 implies slightly increasing summer water tables in the floodplain. Falling in a high  
35 solar activity phase that coincides with a period of low lake-levels in West-Central  
36 Europe (Figure 10), the Sarló-hát proxy records are apparently in mismatch with other  
37 climatic records. A possible interpretation of this discrepancy can be a change in the  
38 water budget of the floodplain. We may see a threshold situation here, whereby the  
39 gradually decreasing summer insolation and the PP10 stalagmite inferred gradual  
40 increase in convective rainfall over the Neolithic period (McDermott et al. 2011) in  
41 east-Central Europe (in this respect see also the gradually increasing lake level of St  
42 Ana) resulted in lower summer evapo-transpiration and hence increasing mid-late  
43 summer moisture availability in the floodplain zone after the recession of floods  
44 (without a need for stronger spring floods).  
45  
46  
47  
48

49 Finally, the onset of the Copper Age around 6550 cal B.P. (4600 cal B.C.) coincided  
50 with a marked increase in Umbelliferae pollen as a wet-meadow indicator in the  
51 lakeshore SH-WOOD profile and also with prominent changes in many European  
52 proxy records, e.g. with a solar activity minimum period and a high West-Central  
53 European lake levels phase that this time coincides with a further increase in Lake St  
54 Ana water depths (Figure 10). The overall impression is that the Tisza floodplain got  
55 wetter in the summer during the ECA, and this may bear a relation with the observed  
56 cultural and landscape exploitation strategies that involved settlement dispersion  
57  
58  
59  
60  
61  
62  
63  
64  
65

(Parkinson 2006) and increased importance of animal husbandry relative to crop farming (Gyulai 2010).

On the whole, this comparison suggests an apparent accordance between cultural changes and climatic oscillations during the Neolithic and ECA. The Sarló-hát proxy records nevertheless show that the alluvial landscape and its vegetation did not respond to solar activity changes directly, as did the West-Central European lake levels (Magny et al. 2003). Rather, these effects were attenuated by the buffering capacity of the floodplain (e.g. during the Szatmár II phase) or prevailed through transmissions modulated by precipitation and temperature changes in the watershed of the Tisza river (e.g. during the ECA). What is less reliable is the inference that any of these climatic fluctuations actually caused the development of specific archaeological cultures. Most prehistoric groups did not live at the extremes of their subsistence possibilities but had a built-in buffering system to cope with unforeseen changes. The rather weak tendencies towards wetter or warmer conditions in the Great Hungarian Plain may have offered better potential for specific responses in terms of settlement or subsistence practices but we are far from demonstrating those inner socio-cultural links.

## Conclusions

In this study, we used results from a carefully designed inter-disciplinary project seeking to integrate archaeology and palaeobotany. Results from a systematic archaeological survey were paired with pollen analysis from the Sarló-hát oxbow lake in the NE Great Hungarian Plain, where deposits covered the last 11,700 years cal BP (9750 cal BC) and provided well-preserved pollen that is rather exceptional in this semi-arid area. Dated by 12 AMS  $^{14}\text{C}$  dates, this record is also exceptional in that it managed to overcome the problems of dating alluvial deposits of low organic content. We focused on the Neolithic and Early Copper Age vegetation changes and their implications for subsistence practices. Sarló-hát is adjacent to a large area with exceptionally good settlement pattern data from intensive, systematic fieldwalking and excavations due to motorway interventions. Several Neolithic and Copper Age sites/discards appear on the lakeshore in direct vicinity of the pollen core locations.

The results of this study allow us to draw the following main conclusions:

- 1) The Sarló-hát pollen records show without any doubt that the hitherto influential Willis and Bennett (1994) hypothesis that there was little human impact by farmers on the environment of SE Europe during the Neolithic is imprecise. While we documented small-scale agriculture in the earliest phases (coeval with Körös; Early Szatmár) in the Sarló-hát area, there are signs of expanding scale of mixed farming in the Middle Neolithic (AVK) and very good evidence for extensive landscape alterations with enhanced pasturing in the Late Neolithic and Early Copper Age. Thus, the strong contrast that Willis and Bennet (1994) drew between the Neolithic and the Bronze Age is fundamentally flawed. More attention to archaeological settlement forms would have made this finding more apparent. The main mitigating factor in revealing Neolithic human impact on the vegetation must have been the distance of most pollen records used in the Willis and Bennett (1994) study from Neolithic and Early Copper Age settlements.
- 2) We recognised, and sought explanations for, some major discrepancies between the pollen proxies for human impact and the archaeological patterns

1 of artefact discard, as a proxy for settlement. We found good fits between  
2 pollen and surface artefact discard for the Middle Neolithic AVK and Early  
3 Copper Age occupation phases. However, considerable mismatches were  
4 demonstrated for the Early Neolithic Körös and Late Neolithic THCS  
5 occupation phases. In case of the first, we explained the discrepancy by either  
6 the alluvial burial of small and hitherto undetected Körös/early Szatmár II sites  
7 or late Mesolithic impact. In case of the LN, the large-scale pastoral land use  
8 suggested by the SH-WOOD pollen record was explained by seasonal use of  
9 alluvial meadows by people living in dispersed homesteads in or near the  
10 alluvium or people living on large, but hitherto undetected horizontal  
11 settlements on the 'Tiszadob Island'.

- 12  
13 3) Comparison with other well-dated pollen diagrams in Eastern Hungary (both  
14 on-site and off-site) pointed to several similarities in land use, such as hazel  
15 and elm coppicing in the Early and Middle Neolithic and more intensive  
16 evidence for pastoral activities and associated woodland clearance in the Late  
17 Neolithic. In addition, there were signs of regional variability, mainly in the  
18 scale and timing of agricultural practices – later and smaller-scale in the North  
19 Hungarian Mid-Mountains in comparison with the GHP. Some of this may  
20 well be related to the distance of Neolithic and ECA sites from the coring site;  
21 other elements include differences in the local environment.  
22  
23 4) Structured comparison with palaeoclimatic proxy records for the Neolithic and  
24 ECA suggest some general alignments between local environmental  
25 fluctuations documented at Sarló-hát and wider climatic fluctuations related to  
26 solar activity changes. Good matches between local and Northern Hemisphere  
27 climatic oscillations at the transitions to the Early Middle Neolithic (Szatmár  
28 II) and the ECA suggest links between cultural change and climate change.  
29 However, we found also mismatches which can be explained in terms of local  
30 buffering effects. The implication from this is that, rather than posing  
31 environmentally deterministic models of changing social practices, we have  
32 been able to suggest some periods when there were more direct responses to  
33 climatic fluctuations and other periods when various kinds of buffering –  
34 ecological mediations or intervening settlement or subsistence practices – were  
35 in operation.  
36  
37  
38  
39

40 One final implication from this study is that it is essential to have high-precision  
41 dating of both sets of data – archaeological and palaeobotanical. Pollen diagrams  
42 with many <sup>14</sup>C AMS (ideally over 10) dates and local settlement records with  
43 multiple AMS <sup>14</sup>C dates are essential. Without this, there is no possibility of  
44 making structured comparisons between these important data sets and thereby  
45 providing the basis not only for dating changes but also proposing causal relations  
46 between the changes.  
47  
48

## 49 **Acknowledgment**

50 This research was funded by the European Commission through a Marie Curie  
51 Postdoctoral Fellowship held by E.K.M. (MEIF-CT-2003-5005001), the Bólyai  
52 Scholarship(BO/00518/07) and the OTKA Research Funds (F026036, PD73234).  
53 Financial support for the fieldwork was provided by the British Academy. This is  
54 MTA-MTM Paleo Contribution No. 141. We thank the help of László Rupnik and  
55 Zoltán Czajlik in preparation of Figure 1. The first author would also thank to Zsolt  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 Molnár for helpful discussions on the potential vegetation and habitat types of the  
2 study area.  
3

#### 4 **Figure legend**

5  
6 **Fig. 1** Location of the study site. (a) Location of the study area in Europe. (b)  
7 Location of the Sarló-hát study site in NE Hungary. Other sites and regions mentioned  
8 in the text are also shown. Key to sites: 1. Kismohos, Nagymohos (Willis et al., 1997,  
9 1998; Magyari et al., 2001); 2. Sirok (Gardner, 2002); 3. Bábtava (Magyari et al.,  
10 2008); 4. Sarló-hát (this study); 5. Ecsegfalva, Kiri-tó (Willis 2007); (c) Map of the  
11 Polgár Region showing the location of Sarló-hát oxbow lake, core locations (1. SH-II  
12 core; 2. SH-WOOD core) and other local place names mentioned in the text; (d)  
13 Second military survey map of the study region in which the Sarló-hát meander  
14 appears as a large swampy area, core locations are also shown (map made between  
15 1806-1869, source: Arcanum 2006).  
16  
17  
18

19 **Fig. 2** Results of the archaeological survey in the Polgár Block indicating the position  
20 of Neolithic sites, stray discards, tell and horizontal settlements. Redrawn and  
21 modified from Chapman et al. (1994).  
22  
23

24 **Fig. 3** Map of fieldwalked areas in the vicinity of the Sarló-hát meander showing the  
25 location of archaeological sites Dob 28-31, TG9 and sediment cores SH-II and SH-  
26 WOOD.  
27  
28

29 **Fig. 4** Sarló-hát lithology and chronology. (a) Sediment lithology and organic content  
30 for the SH-II and SH-WOOD cores plotted against depth. (b)  $^{14}\text{C}$  dates and Bayesian  
31 age–depth model constructed using Bchron (Parnell et al. 2008) for the SH-WOOD  
32 core. (c) local pollen assemblage zones (LPAZ) of SH-WOOD. ECA: Early Copper  
33 Age.  
34  
35

36 **Fig. 5** Pollen percentage diagram for the SH-WOOD core, Sarló-hát, Tiszagyulaháza.  
37 Percentages of selected terrestrial pollen taxa are plotted against age (cal B.P.), the  
38 depth scale also being indicated. AP:NAP ratios, microcharcoal concentrations,  
39 palynological richness and sediment organic content are also shown. . ECA: Early  
40 Copper Age; THCS: Tisza-Herpály-Csőszhalom Cultures.  
41  
42  
43  
44

45 **Fig. 6** Pollen percentage diagram for the SH-II core, Sarló-hát, Tiszagyulaháza.  
46 Percentages of selected terrestrial pollen taxa are plotted against age (cal B.P.), the  
47 depth scale also being indicated. AP:NAP ratios, palynological richness and sediment  
48 organic content are also shown. ECA: Early Copper Age; THCS: Tisza-Herpály-  
49 Csőszhalom Cultures.  
50  
51

52 **Fig. 7** Percentage diagram of selected algae, aquatic and wetland pollen types from  
53 the SH-WOOD core, Sarló-hát, Tiszagyulaháza. Organic content is also shown. ECA:  
54 Early Copper Age; THCS: Tisza-Herpály-Csőszhalom Cultures.  
55  
56

57 **Fig. 8** Cumulative percentages of pollen types reflecting different human land use  
58 types in the pollen records of SH-WOOD and SH-II, Sarló-hát, Tiszagyulaháza,  
59  
60  
61  
62  
63  
64  
65

1 Hungary. ECA: Early Copper Age; AVK: Alföld Linear Pottery Culture; THCS:  
2 Tisza-Herpály-Csőszhalom Cultures.  
3

4 **Fig. 9** Percentage diagram of selected pollen types for the Early and Mid Holocene  
5 from Nagymohos and Bábtava peat bogs (Magyari et al. 2001, 2008) and the number  
6 of Neolithic settlements in 50 km radius of the sites. For site locations see Figure 1b.  
7

8 **Fig. 10** Comparison between various Northern Hemisphere paleoclimate series and  
9 the SH-WOOD pollen, microcharcoal and organic content records. (1) 50 yr averaged  
10 NGRIP  $\delta^{18}\text{O}$  proxy for temperature (North Greenland Ice Core Project members  
11 2004); (2) residual  $^{14}\text{C}$  data calculated from IntCal04  $\Delta^{14}\text{C}$  (%) with a 2000 yr moving  
12 average subtracted (Reimer et al. 2004); (3) PP10 stalagmite  $\delta^{18}\text{O}$  record from Poleva  
13 Cave, SW Romania (Constantin et al. 2007); (4) Macrofossil based relative lake level  
14 changes in Lake St Ana, Romanian Carpathians; the curve shows a hydrological  
15 gradient from shallow to deep lake represented here by principal component axis 2  
16 (Magyari et al. 2009); (5) High lake level phases in West-Central Europe according to  
17 Magny (2007); (6) organic content, relative frequencies of pollen types and green  
18 algae, ratio of arboreal and non-arboreal pollen (AP:NAP) and microcharcoal  
19 concentrations from SH-WOOD, Sarló-hát, NE Hungary (this study); (7) Summer  
20 (red curve) and winter (black curve) insolation values ( $\text{W}/\text{m}^2$ ) at  $60^\circ\text{N}$  (Berger and  
21 Loutre, 1991). ECA: Early Copper Age; THCS: Tisza-Herpály-Csőszhalom Cultures;  
22 Tiszapolg.: Tiszapolgár Culture.  
23  
24  
25  
26  
27

28 **Additional file 1** Comparative table of plant macrofossil taxa recorded by the  
29 archaeobotanical investigation of Late Neolithic settlements in the Polgár area and  
30 their equivalent pollen types in the Sarló-hát (SH-II, SH-WOOD) pollen diagrams.  
31 The Late Neolithic settlement, Polgár-10 was analysed by Fairbairn (1992, 1993) and  
32 the main core of the taxon list enumerates his data. In addition, taxa not found by  
33 Fairbairn but determined by Gyulai (2010) in course of an independent survey of the  
34 Csőszhalom tell are shown and marked (GY-CSH).  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## References

- 1  
2 Aaby B Digerfeldt G (1986) Sampling techniques for lakes and bogs. In: Berglund BE (ed) Handbook  
3 of Holocene Palaeoecology and Palaeohydrology John Wiley and Sons Chichester UK, pp 181–194  
4 Arcanum (2006) A második magyar katonai felmérés 1806-1869 (The Second Military Survey 1806-  
5 1869) DVD Arcanum Kft Budapest, in Hungarian  
6 Behre KE (1981) The interpretation of anthropogenic indicators in pollen diagrams. *Pollen et Spores*  
7 23:225–45  
8 Behre KE (ed) (1986) *Anthropogenic Indicators in Pollen Diagrams*. Balkema, Rotterdam  
9 Behre KE (2007) Evidence for Mesolithic agriculture in and around central Europe? *Veg Hist*  
10 *Archaeobot* 16:203–219  
11 Bánffy E (2005) The Csaroda area during the Mesolithic the Neolithic and the Copper Age In: Gál E  
12 Juhász I Sümegei P (eds) *Environmental archaeology in North-Eastern Hungary*. VAH 19 Budapest,  
13 pp 207-222  
14 Bennett KD (1997) PSIMPOLL 30 & PSCOMB 103 Manual Program. URL:  
15 <http://www.chronoqubacuk/psimpoll/psimpoll.html>  
16 Berger A, Loutre MF (1991) Insolation values for the climate of the last 10 million of years. *Quatern*  
17 *Sci Rev* 10:297–317  
18 Berglund BE (2003) Human impact and climate changes—synchronous events and a  
19 causal link. *Quatern Int* 105:7–12  
20 Berglund BE, Ralska-Jasiewiczowa M (1986) Pollen analysis and pollen diagrams. In: Berglund BE  
21 (ed) *Handbook of palaeoecology and palaeohydrology*. John Wiley and Sons, New York, pp 455-  
22 479  
23 Birks HJ, Birks HH (1980) *Quaternary palaeoecology*. University Park Press, Baltimore  
24 Birks HJB, Line JM (1992) The use of rarefaction analysis for estimating palynological richness from  
25 Quaternary pollen-analytical data. *Holocene* 2:1-10  
26 Bogaard A, Bending J, Jones G (2008) Crop husbandry and its social significance in the Körös and  
27 LBK cultures In: Bailey D Whittle A & Hofmann D (eds) *Living well together? Settlement and*  
28 *materiality in the Neolithic of South-East and Central Europe*. Oxbow Books, Oxford, pp 131-139  
29 Bökönyi S (1959) Die frühalluviale Wirbeltierfauna Ungarns (vom Neolithikum bis zur La Tène Zeit).  
30 *Acta Archaeol Hung* 11:39-102  
31 Bökönyi S (1974) *History of domestic mammals in Central and Eastern Europe*. Akadémiai Kiadó,  
32 Budapest  
33 Borsy Z, Csongor É, Félegyházi E (1989) A Bodrogtörzs kialakulása és vízhálózatának változásai  
34 (Formation of the Bodrogtörzs landscape and changes of its river network). *Alföldi Tanulmányok*  
35 13:65-81 (in Hungarian)  
36 Chapman J (1994) Social power in the early farming communities of Eastern Hungary - perspectives  
37 from the Upper Tisza region. *A Jósza András Múzeum Évkönyve* 36:79-100  
38 Chapman J (1997) Places as timemarks – the social construction of landscapes in Eastern Hungary. In:  
39 Chapman J, Dolukhanov P (eds) *Landscapes in Flux*. *Colloquenda Pontica* 3, Oxbow Books,  
40 Oxford, pp 137-162  
41 Chapman J, Laszlovszky J (1993) The Upper Tisza Project 1992. *Archaeological Reports for 1992*,  
42 University of Durham, pp 13-19  
43 Chapman J, Shiel R, Passmore D, Magyari E (2003) The Upper Tisza Project: studies in Hungarian  
44 landscape archaeology. [http://adsahdsacuk/catalogue/projArch/uppertisza\\_ba\\_2003/indexcfm](http://adsahdsacuk/catalogue/projArch/uppertisza_ba_2003/indexcfm)  
45 Accessed 5 July 2011  
46 Chapman J, Magyari E, Gaydarska B (2009) Contrasting subsistence strategies in the Early Iron Age?  
47 New results from the Alföld Plain Hungary and the Thracian Plain Bulgaria. *Oxford J Archaeol*  
48 28/2:155 – 187  
49 Chapman J, Gillings M, Shiel R, Gaydarska B, Bond C (2010a) The Upper Tisza Project Book 4,  
50 Lowland settlement in North East Hungary: excavations at the Neolithic settlement site of Polgár-  
51 10. BAR International Series 2089, Archaeopress, Oxford  
52 Chapman J, Gillings M, Shiel R, Gaydarska B, Bond C (2010b) The Upper Tisza Project Book 3,  
53 Settlement patterns in the Bodrogtörzs block. BAR International Series 2087, Archaeopress, Oxford  
54 Clark RL (1982) Point count estimation of charcoal in pollen preparations and thin sections in  
55 sediments. *Pollen et Spores* 24:523-535  
56 Colombaroli D, Vannièrè B, Chapron E, Magny M, Tinner W (2008) Fire-vegetation interactions  
57 during the Mesolithic-Neolithic at Lago dell'Accesa Italy. *Holocene* 18: 679-692  
58  
59  
60  
61  
62  
63  
64  
65

- Constantin S, Bojar AV, Lauritzen SE, Lundberg J (2007) Holocene and Late Pleistocene climate in the sub-Mediterranean continental environment: A speleothem record from Poleva Cave (Southern Carpathians Romania). *Palaeogeogr Palaeoclimatol* 243:322–338
- Csengeri P (2005) The Neolithic and the Copper Age in the Sajó-Bódva Interfluve. In: Gál E, Juhász I, Sümegei P (eds) *Environmental Archaeology in North-eastern Hungary*. *Varia Archaeologica Hungarica* 19: 223–235
- Domboróczki L (2005) A Körös-kultúra északi elterjedési határának problematikája a Tiszaszőlös-Domaháza-Pusztán végzett ásítás eredményének fényében (The problem of the northern boundary of the Körös Culture in light of excavations at Tiszaszőlös-Domaháza-Pusztá). *Archeometriai Műhely* 2005/2: 5-15 (in Hungarian with English summary)
- Domboróczki L, Raczky P (2010) Excavations at Ibrány-Nagyerdő and the northernmost distribution of the Körös culture in Hungary. In: Kozłowski JK, Raczky P (eds) *Neolithization of the Carpathian Basin: Northernmost distribution of the Starčevo/Körös culture*. Polish Academy of Arts & Sciences and Institute of Archaeological Sciences ELTE, Kraków-Budapest, pp 191-218
- Ecsedy I, Kovács L, Maráz B, Torma I (eds) (1982) Békés megye régészeti topográfiája IV/1 A szeghalmi járás (Archaeological topography of Hungary, County Békés). *Magyarország Régészeti Topográfiája* 6, Akadémiai Kiadó, Budapest (in Hungarian)
- Fairbairn AS (1992) Archaeobotanical investigations at Csöszhalom: a Late Neolithic tell site in north-east Hungary. MSc Thesis, Institute of Archaeology, University College London
- Fairbairn AS (1993) Plant husbandry at the prehistoric Hungarian tell sites of Csöszhalom and Kenderföld. Final Report for the British Academy Applied Sciences in Archaeology Fund, Durham University, Durham
- Fekete G, Molnár Zs, Horváth F (eds) (1997) A magyarországi élőhelyek leírása határozója és a Nemzeti Élőhely-osztályozási Rendszer (Description and identification key to Hungarian vegetation types according to habitats). A Nemzeti Biodiverzitás-monitorozó Rendszer Kézikönyvei 2, Természettudományi Múzeum, Budapest (in Hungarian)
- Gardner AR (2002) Neolithic to Copper Age woodland impacts in northeast Hungary? Evidence from pollen and sediment chemistry records. *Holocene* 12(5):541-553
- Gillings M (1997) Spatial organization in the Tisza floodplain: landscape dynamics and GIS. In: Chapman J, Dolukhanov P (eds) *Landscapes in flux: Central and Eastern Europe in Antiquity*. Oxbow Books, Oxford, pp 163-179
- Gillings M (1998) Embracing uncertainty and challenging dualism in the GIS-based study of a paleo-flood plain. *European Journal of Archaeology* 1(1):117-144
- Gillings M (2007) The Ecsefalva landscape: affordance and inhabitation. In: Whittle A (ed) *The Early Neolithic on the Great Hungarian Plain, Investigations of the Körös culture site of Ecsefalva 23, County Békés*. Institute of Archaeology, Budapest, pp 31-46
- Gulyás S (2011) Az édesvízi kagylók szerepe a kárpát-medencei neolitik közösségek gazdálkodásában és az ártéri folyóvízi környezet lokális és regionális adottságainak rekonstrukciójában (The role of freshwater bivalves in the economy of Neolithic communities and in the reconstruction of the local and regional floodplain environment). PhD Thesis, University of Szeged, Hungary (in Hungarian with English summary)
- Gyulai F (2010a) Archaeobotany in Hungary: Seed Fruit Food and Beverage Remains in the Carpathian Basin from the Neolithic to the Late Middle Ages. *Archeolingua*, Budapest
- Hajdú Zs & Nagy E (1999) Rövid jelentés az M3-as autópálya Hajdú-Bihar megyei szakaszán azonosított régészeti lelőhelyeken 1993-1998 között végzett munkálatokról (Short report on archaeological sites detected between 1993-1998 along the M3 motorway line in county Hajdú-Bihar). *A Debreceni Déri Múzeum Évkönyve 1997/98*: 143-154 (in Hungarian)
- Heiri O, Lotter AF, Lemcke G (2001) Loss on ignition as a method for estimating organic and carbonate content in sediments: reproducibility and comparability of results. *J Paleolimnol* 25:101-110
- Hertelendi E, Kalicz N, Raczky P, Horváth F, Veres M, Svngor É, Futó I, Bartosiewicz L (1995) Re-evaluation of the Neolithic in Eastern Hungary based on calibrated radiocarbon dates. *Radiocarbon* 37:239-244
- Iversen J (1941) *Landnam i Danmarks Stenalder: En pollenanalytisk Undersøgelse over det første Landbrugs Indvirkning paa Vegetationsudviklingen* (Dansk tekst 7-59 Engl text 60-65). *Danmarks Geologiske Undersøgelse II række* 66:1-68 (reprinted 1964)
- Jankovich BD, Makkay J, Szőke BM (eds) (1989) Békés megye régészeti topográfiája IV/2: szarvasi járás (Archaeological topography of county Békés IV/2: the Szarvas municipality). *Magyarország Régészeti Topográfiája* 8, Akadémiai Kiadó, Budapest (in Hungarian)

- 1 Johnson GA (1984) Organizational structure and scalar stress. In: Renfrew C, Rowlands MJ, Seagraves  
2 BA (eds) *Theory and explanation in archaeology: the Southampton conference*. Academic Press,  
3 New York, pp 389-421
- 4 Jones G, Bogaard A, Halstead P, Charles M, Smith H (1999) Identifying the intensity of crop  
5 husbandry practices on the basis of weed floras. *Annual of the British School at Athens* 94:167-189.
- 6 Juhász I, Sümegei P, Szántó Zs, Svingor É, Molnár M, Jakab G (2007) The Little Balaton region and the  
7 Balaton uplands. In: Zatykó Cs, Juhász I, Sümegei P (eds) *Environmental Archaeology in*  
8 *Transdanubia*. *Varia Archaeologica Hungarica* 20, Budapest, pp 25-78
- 9 Kalicz N, Makkay J (1977) Die Linienbandkeramik in der Großen Ungarischen Tiefebene. *Studia*  
10 *Archaeologica* 7, Budapest
- 11 Kalicz N, Raczky P (eds) (1987) *The Late Neolithic of the Tisza region*. Szolnok County Museums,  
12 Budapest-Szolnok
- 13 Kertész R, Sümegei P (1999) Teóriák kritika és egy modell: Miért állt meg a Körös-Starčevo kultúra  
14 terjedése a Kárpát-medencében? (Theories, critic and a model: why did the spread of the Körös-  
15 Starčevo Culture stop in the Carpathian Basin?) *Tisicum* 11: 9-23 (in Hungarian)
- 16 Kosse K (1979) Settlement ecology of the Early and Middle Neolithic Körös and Linear Pottery  
17 cultures in Hungary. *BAR International Series I-64*, Oxford
- 18 Kreuz A (2007) Archaeobotanical perspectives on the beginning of agriculture north of the Alps. In:  
19 Colledge S Conolly J (eds) *Archaeobotanical perspectives on the origin and spread of agriculture in*  
20 *southwest Asia and Europe*. UCL Press, London
- 21 Kuneš P, Pokorný P, Šída P (2008) Detection of impact of Early Holocene hunter-gatherers on  
22 vegetation in the Czech Republic using multivariate analysis of pollen data. *Veg Hist Archaeobot*  
23 17: 269-287
- 24 Lüning J, Kalis AJ (1992) The influence of Early Neolithic settlers on the vegetation of the Lower  
25 Rhineland and the determination of cleared areas based on archaeological and palynological  
26 criteria. In: Frenzel B (ed) *Evaluation of land surfaces cleared from forests by prehistoric man in*  
27 *Early Neolithic times and the time of migrating Germanic tribes*. *Paläoklimaforschung* 8, pp 41-46
- 28 Magny M, Begeot C, Guiot J, Peyron, O (2003) Contrasting patterns of hydrological changes in Europe  
29 in response to Holocene climate cooling phases. *Quatern Sci Rev* 22: 1589-1596
- 30 Magny M (2007) Lake level studies - West-Central Europe. In: *Encyclopedia of Quaternary Studies*,  
31 Volume 2, Elsevier, Amsterdam, pp 1389-1399
- 32 Magyari E, Sümegei P, Braun M, Jakab G, Molnár M (2001) Retarded wetland succession:  
33 anthropogenic and climatic signals in a Holocene peat bog profile from the NE Carpathian Basin. *J*  
34 *Ecol* 89:1019-1032
- 35 Magyari EK, Jakab G, Sümegei P, Szőőr Gy (2008) Holocene vegetation dynamics in the Bereg Plain  
36 NE Hungary – the Báb-tava pollen and plant macrofossil record. *Acta Geographica Debrecina* 42:1-  
37 16
- 38 Magyari EK, Buczkó K, Jakab G, Braun M, Pál Z, Karátson D (2009) Palaeolimnology of the last  
39 crater lake in the Eastern Carpathian Mountains - a multiproxy study of Holocene hydrological  
40 changes. *Hydrobiologia* 631: 29-63
- 41 Magyari EK, Chapman JC, Passmore DG, Allen JRM, Huntley JP, Huntley B (2010) Holocene  
42 persistence of wooded steppe in the northern Great Hungarian Plain. *Journal of Biogeogr* 37:915-  
43 935
- 44 Makkay J (1982a) Some comments on the settlement pattern of the Alföld Linear Pottery. In: Pavúk J  
45 (ed) *Siedlungen der Kultur mit Linearkeramik in Europa*. *Archäologisches Institute der*  
46 *Slowakischen Akademie der Wissenschaften, Nitra*, pp 157-166
- 47 Makkay J (1982b) A magyarországi neolitikum kutatásának új eredményei (New results of the  
48 Hungarian Neolithic research). *Akadémiai Kiadó, Budapest* (in Hungarian)
- 49 Marinova E (2007) Archaeobotanical data from the early Neolithic of Bulgaria. In: Connolly J,  
50 Colledge S (eds) *Early Neolithic in Southwest Asia and Europe: Archaeobotanical perspectives in*  
51 *Neolithic Plant Economies*. UCL Press, London, pp 93-109
- 52 Marinova E, Tonkov S, Bozilova E, Vajsov I (in press) Holocene anthropogenic landscapes in the  
53 Balkans: the paleobotanical evidence from southwestern Bulgaria. *Veg Hist Archaeobot*
- 54 Marosi M, Somogyi M (1990) *Magyarország Kistájainak Katasztere I (Cadastre of small regions in*  
55 *Hungary I)*. MTA Földrajztudományi Kutatóintézet, Budapest (in Hungarian)
- 56 McDermott F, Atkinson TC, Fairchild IJ, Baldini LM, Matthey DP (2011) A first evaluation of the  
57 spatial gradients in  $\delta^{18}\text{O}$  recorded by European Holocene speleothems. *Global Planet Change*,  
58 doi:10.1016/j.gloplacha.2011.01.005

- 1 Molnár Zs (1996) Flood plain vegetation at Tiszadob and Kesznyéten (Middle Tisza Valley) II. History  
2 and present state of the hardwood floodplain woodlands (Fraxini-Pannonicae-Ulmetum). *Botanikai*  
3 *Közlemények* 83:51-69 (in Hungarian with English summary)
- 4 Moore PD, Webb JA, Collinson ME (1992) *Pollen analysis*, 2<sup>nd</sup> edition, Blackwell Scientific  
5 Publications, Oxford
- 6 Nandris J (1970) Ground water as a factor in the First Temperate Neolithic of the Körös region.  
7 *Zbornik Narodnog Muzeja* (Beograd) 6:59-71
- 8 North Greenland Ice Core Project members (2004) High-resolution record of Northern Hemisphere  
9 climate extending into the last interglacial period. *Nature* 431:147-151
- 10 Parkinson WA (2006) *The social organization of Early Copper Age tribes on the Great Hungarian*  
11 *Plain*. BAR International Series I-1573, Archaeopress, Oxford
- 12 Parnell AC, Haslett J, Allen JRM, Buck CE, Huntley B (2008) A flexible approach to assessing  
13 synchronicity of past events using Bayesian reconstructions of sedimentation history. *Quatern Sci*  
14 *Rev* 27: 1872-1885
- 15 Pavúk J (2007) Zur Frage der Entstehung und Verbreitung der Lengyel-Kultur. In: Kozłowski JK,  
16 Raczky P (eds) *The Lengyel Polgár and related cultures in the Middle/Late Neolithic in Central*  
17 *Europe*. Polish Academy of Arts & Sciences and Institute of Archaeological Sciences ELTE,  
18 Kraków, pp 11-28
- 19 Poska A, Saarse L, Veski S (2004) Reflections of pre- and early-agrarian human impact in the pollen  
20 diagrams of Estonia. *Palaeogeogr Palaeoclimatol* 209: 37-50
- 21 Punt W, Clarke GCS (eds) (1984) *The Northwest European Pollen Flora IV*. Elsevier, Amsterdam
- 22 Raczky P, Meier-Arendt W, Kurucz K, Hajdú Zs, Szikora Á (1994) Polgár-Csőszhalom, a Late  
23 Neolithic settlement in the Upper Tisza Region and its cultural connections, preliminary report  
24 (Polgár-Csőszhalom. Egy késő neolitikus lelőhely kutatása a Felső-Tisza vidéken és annak  
25 kulturális összefüggései) *A nyíregyházi Jósza András Múzeum Évkönyve* 36: 231-240 (in  
26 Hungarian)
- 27 Raczky P (1995) Late Neolithic settlement patterns in the Tisza region of Hungary. In: Aspes A (ed)  
28 Symposium "Settlement patterns between the Alps and the Black Sea 5<sup>th</sup> – 2<sup>nd</sup> millennium BC.  
29 *Memorie del Museo Civico di Storia Naturale di Verona* II/4, pp 77-86
- 30 Raczky P, Anders A (2008) Late Neolithic spatial differentiation at Polgár-Csőszhalom eastern  
31 Hungary. In: Bailey DW, Whittle A, Hofmann D (eds): *Living Well Together? Settlement and*  
32 *materiality in the Neolithic of south-east and central Europe*. Oxbow Books, Oxford, pp 35-53
- 33 Raczky P, Domboróczki L, Hajdú Zs (2007) The site of Polgár-Csőszhalom and its cultural and  
34 chronological connections with the Lengyel culture. In: Kozłowski JK, Raczky P (eds) *The Lengyel*  
35 *Polgár and related cultures in the Middle/Late Neolithic in Central Europe*. Polish Academy of Arts  
36 & Sciences and Institute of Archaeological Sciences ELTE, Kraków, pp 49-70
- 37 Raczky P, Meier-Arendt W, Anders A, Hajdú Zs, Nagy E, Kurucz K, Domboróczki L, Sebők K,  
38 Sümegi P, Magyari E, Szántó Zs, Gulyás S, Dobó K, Bácskay ET, Biró K, Schwartz C (2002)  
39 Polgár-Csőszhalom (1989–2000): Summary of the Hungarian-German Excavations on a Neolithic  
40 Settlement in Eastern Hungary. In: Aslan R, Blum S, Kastl G, Schweizer F, Thumm D (eds)  
41 *Mauerschau: Festschrift für Manfred Korfmann Band 2*, Remshalden-Grunbach, pp 833-860
- 42 Raczky P, Anders A, Bartosiewicz L (2011) The Enclosure System of Polgár-Csőszhalom and its  
43 Interpretation. In: Hansen S, Müller J (eds) *Sozialarchäologische Perspektiven: Gesellschaftliche*  
44 *Wandel 5000-1500 v. Chr. zwischen Atlantik und Kaukasus*, Archäologie in Eurasien 24. Verlag  
45 Philipp Von Zabern, Darmstadt, pp. 57-79
- 46 Reille M (1992) *Pollen et spore D'Europe et D'Afrique du Nord*. Laboratoire de Botanique Historique  
47 et Palynologie, Marseille, France
- 48 Reille M (1995) *Pollen et spore D'Europe et D'Afrique du Nord*, Supplement 1. Laboratoire de  
49 Botanique Historique et Palynologie, Marseille, France
- 50 Reille M (1998) *Pollen et spore D'Europe et D'Afrique du Nord*, Supplement 2. Laboratoire de  
51 Botanique Historique et Palynologie Marseille, France
- 52 Reimer PJ, Baillie MGL, Bard E et al (2004) IntCal04 terrestrial radiocarbon age calibration 0–26 cal  
53 kyr BP. *Radiocarbon* 46:1029-1058
- 54 Renfrew C (1984) *Approaches to social archaeology*. Edinburgh University Press, Edinburgh
- 55 Seppä H (1998) Postglacial trends in palynological richness in the northern Fennoscandian tree-line  
56 area and their ecological interpretation. *Holocene* 8:43-53
- 57 Sherratt A (1982) Mobile resources: settlement and exchange in early agricultural Europe. In: Renfrew  
58 C, Shennan SJ (eds) *Ranking resources and exchange*. Cambridge University Press, Cambridge, pp  
59 13-26
- 60  
61  
62  
63  
64  
65

- 1 Sherratt A (1982/3) The development of Neolithic and Copper Age settlement in the Great Hungarian  
 2 Plain Parts I and II. *Oxford J Archaeol* 1: 287-316 and 2: 13-41
- 3 Soepboer W, Sugita S, Lotter AF, van Leeuwen JFN, van der Knaap WO (2007) Pollen productivity  
 4 estimates for quantitative reconstruction of vegetation cover on the Swiss Plateau. *Holocene* 17:65-  
 5 77
- 6 Stockmarr J (1971) Tablets with spores used in absolute pollen analysis. *Pollen et Spores* 13: 614-621
- 7 Sugita S (1993) A model of pollen source area for an entire lake surface. *Quaternary Res* 39:239-44
- 8 Sugita S (2007a) Theory of quantitative reconstruction of vegetation I: pollen from large sites  
 9 REVEALS regional vegetation composition. *Holocene* 17:229-241
- 10 Sugita S (2007b) Theory of quantitative reconstruction of vegetation II: all you need is LOVE  
 11 *Holocene* 17:243-57
- 12 Sümegei P (1999) Reconstruction of flora soil and landscape evolution and human impact on the Bereg  
 13 Plain from late-glacial up to the present based on palaeoecological analysis. In: Hamar J, Sárkány-  
 14 Kiss A (eds) *The Upper Tisza Valley*. Tiscia Monograph Series, Szeged, pp 171-203
- 15 Sümegei P, Kertész R (1998) A Kárpát-medence ökoszisztémájának jellemzői – egy ökológiai csapda az  
 16 újkőkorszakban? (Palaeoenvironmental characteristics of the Carpathian Basin – an ecological pitfall in  
 17 the Neolithic?). *Jászok* 44(3-4):144-157
- 18 Sümegei P (2005) Preneolithic development in North-Eastern Hungary. In: In: Gál E, Juhász I, Sümegei  
 19 P (eds) *Environmental Archaeology in North-East Hungary*. *Varia Archaeologica Hungarica*,  
 20 Budapest, pp 15-24.
- 21 Sümegei P, Csökmei B, Persaits G (2005) The evolution of Polgár Island a loess-covered lag surface and  
 22 its influences on the subsistence of settling human cultural groups. In: Hum L, Gulyás S, Sümegei P  
 23 (eds) *Environmental historical studies from the Late Tertiary and Quaternary of Hungary*.  
 24 Department of Geology and Paleontology, University of Szeged, Szeged, pp 141-164
- 25 Szász G, Tökei L (1997) Meteorológia mezőgazdáknek kertészeknek erdészeknek (Meteorology for  
 26 agronomists horticulturists and foresters). *Mezőgazda Kiadó*, Budapest (in Hungarian)
- 27 Tímár G, Sümegei P, Horváth F (2005) Late Quaternary dynamics of the Tisza River: Evidence of  
 28 climatic and tectonic controls. *Tectonophysics* 410(1-4): 97-110
- 29 Tinner W, Conedera M, Ammann B, Lotter AF (2005) Fire ecology north and south of the Alps since  
 30 the last ice age. *Holocene* 15:1214-1226
- 31 Tinner W, Nielsen EH, Lotter AF (2007) Mesolithic agriculture in Switzerland? A critical review of the  
 32 evidence. *Quatern Sci Rev* 26:1416-1431
- 33 Troels-Smith J (1955) Karakterisering af løse jordarter Danmarks. *Geologiske Undersøgelse Series IV*  
 34 3(10):1-73
- 35 Ujvárosi M (1957) Gyomnövények gyomirtás (Weeds and chemical weed control). *Mezőgazdasági*  
 36 *Kiadó*, Budapest
- 37 Walter H (1974) Vegetationsmonographien der einzelnen Großräume, Band VII, Die Vegetation  
 38 Osteuropas Nord- und Zentralasiens. Gustav Fischer Verlag, Stuttgart
- 39 Wasylikowa K (1996) Environmental changes during Neolithic times in the Cracow area. In: Frenzel B  
 40 (ed) *Evaluation of land surfaces cleared from forests by prehistoric man in Early Neolithic times*  
 41 *and the time of migrating Germanic tribes*. *Palaeoclimate Research Volume 8*, Gustav Fisher  
 42 Verlag, Stuttgart, pp 73-91
- 43 Whitlock C, Larsen CPS (2001) Charcoal as a Fire Proxy. In: Smol JP, Birks HJB, Last WM (eds)  
 44 *Tracking Environmental Change Using Lake Sediments, Volume 3: Terrestrial Algal and Siliceous*  
 45 *indicators*. Kluwer Academic Publishers, Dordrecht, pp 75-97
- 46 Whittle A (ed) (2007) *The Early Neolithic on the Great Hungarian Plain Investigations of the Körös*  
 47 *culture site of Ecsefalva 23 County Békés*. Institute of Archaeology, Budapest
- 48 Willis KJ and Bennett KD (1994) The Neolithic transition - Fact or Fiction? Palaeoecological evidence  
 49 from the Balkans. *Holocene* 4:326-330
- 50 Willis KJ, Sümegei P, Braun M, Tóth A (1997) Does soil change cause vegetation change or vice versa?  
 51 A temporal perspective from Hungary. *Ecology* 78(3):740-750
- 52 Willis KJ, Sümegei P, Braun M, Bennett KD, Tóth A (1998) Prehistoric land degradation in Hungary:  
 53 who how and why? *Antiquity* 72:101-113
- 54 Willis KJ (2007) The impact of the Early Neolithic Körös culture on the landscape: evidence from  
 55 palaeoecological investigations of the Kiri-tó. In: Whittle A (ed) *The Early Neolithic on the Great*  
 56 *Hungarian Plain Investigations of the Körös culture site of Ecsefalva 23 County Békés*. Institute of  
 57 Archaeology, Budapest, pp 83-98
- 58 Zólyomi B (1946) Természetes növénytakaró a tiszafüredi öntözőrendszer területén (Natural vegetation  
 59 on the Irrigation system-area of Tiszafüred). *Öntözésügyi Közlemények* 7-8:62-74 (in Hungarian)

Zvelebil M (1994) Plant use in the Mesolithic and its role in the transition to farming. P Prehist Soc  
60:35-74

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65



1 **Table 1** Selection of non-arboreal pollen types as anthropogenic indicators in the  
 2 Neolithic section of cores SH-WOOD and SH-II, Sarló-hát, Tiszagyulaháza, Hungary.  
 3 The category of land use was assigned to the pollen types following Behre (1981),  
 4 Ujvárosi (1957) and Fekete et al. (1997). Grey fill indicates the inclusion of a taxon in the  
 5 pooled pollen frequency sum of each land use category. On the right, natural plant  
 6 communities are listed in which the species of the pollen type occur in the Middle Tisza  
 7 Plain.  
 8  
 9

Pollen type	Land use type and ecology							Natural communities in the Upper Tisza floodplain
	Crop fields (Cereals)*	Crop fields (Weeds)	Fallow land / balk herbs	Footpath and ruderal vegetation	Wet pasture/ hay meadow	Dry pasture/ meadow		
<i>Secale</i>								<i>S. sylvestre</i> : calcareous sand steppes (unlikely in the area)
<i>Triticum</i>								-
<i>Cirsium</i>								marshy meadows, forest steppe, saline communities, alder carr
<i>Polygonum aviculare</i>								mudflat communities
<i>Ambrosia</i> -type								<i>X. strumarium</i> , pioneer mudflat and floodplain marsh communities
<i>Melilotus</i>								-
<i>Potentilla</i>								mudflat communities, tall sedge and tall herb wet meadows, marshy habitats, saline mud, forest steppe, sand steppe
<i>Rumex acetosa / acetosella</i>								sand steppe, dry meadows
<i>Apium</i> -type								wet floodplain meadows and marshy habitats
<i>Bellis</i>								-
<i>Cannabis/Humulus</i>								alder and willow carr, woodland edge
Cruciferae								reed swamp, wet meadows mainly
<i>Euphorbia</i>								oak-ash-elm forest edge, dry oak forest edge, sand steppes
<i>Gentiana</i>								wet floodplain meadows
<i>Pedicularis</i>								tall sedge fens
<i>Peucedanum</i> -type								alder carr, tall sedge fen, sand steppe, oak forest steppe, loess oak forest,
<i>Plantago major-media</i>								seasonally flooded meadows
<i>Ranunculus acris</i>								wet floodplain meadows
<i>Ranunculus</i> sp.								wet floodplain meadows, willow carr, tall sedge fen, saline steppe, dry oak forest, mudflat communities
<i>Rumex</i> sp.								reedswamp, wet alluvial meadows and marshy habitats, floodplain ruderal mudflat vegetation
<i>Scabiosa</i>								dry steppe
<i>Stachys</i>								oak-ash-elm forest, oak forest steppe, willow carr, reed swamp, tall sedge fen
<i>Stellaria</i> -type								includes several genera, wide distribution in floodplain plant communities
Umbelliferae* <sup>1</sup>								marshy oxbow lakeshores with summer-dry habitat (Rorippo-Oenanthetum), willow carr, alder carr, permanently wet-soil floodplain meadows
<i>Artemisia</i>								<i>A. vulgaris</i> * <sup>2</sup> saline wormwood steppe, saline tall herb wet meadow, forest steppe
<i>Veronica</i>								oak forest, alder carr, shallow oxbow lakeshores (Rorippo-Oenanthetum), tall herb fens, steppe meadows
<i>Cerastium</i> -type								saline wormwood steppe, sand steppe, floodplain meadows
<i>Dianthus</i>								sand steppe, forest steppe, saline tall herb wet meadow, meadow steppe
<i>Filipendula vulgaris</i>								tall herb wet meadows, loess oak forest steppe
<i>Lotus</i> -type								wormwood steppe, wet meadows
<i>Verbascum</i>								loess steppe
Chenopodiaceae* <sup>3</sup>								floodplain ruderal mudflat vegetation (seasonally flooded lakes and channels with organic debris accumulation hence high nutrient availability), wet saline meadows, saline lake bottoms, willow carr

\* Poaceae > 40µm were not included because of the suspected presence of wetland Gramineae in this group (e.g. *Glyceria*)

\*<sup>1</sup> includes *Oenanthe*, *Cicuta*, *Angelica* and Umbelliferae undifferentiated

\*<sup>2</sup> *Artemisia vulgaris* is a typical constituent of disturbed tall herb wet floodplain meadows

\*<sup>3</sup> Chenopodiaceae were not included in any land use category because of its suggested abundance in the natural mudflat communities (see the methods section)

**Table 2** Pollen assemblage characteristics of Neolithic and Early Copper Age cultural periods in the SH-WOOD and SH-II pollen records (Figs. 5-8)

Cultural period	Age	Characteristics of cultural period (principal or distinctive pollen, microcharcoal, palynological richness)
Early Neolithic Körös	7950 – 7450 cal B.P. 6000 – 5500 cal B.C.	Strong relative frequency fluctuation of <i>Corylus</i> (SH-WOOD: 12-23%, av. 18%, SH-II: 22.5-33%, av. 28%) and <i>Quercus</i> (SH-WOOD: 21-30%, av. 25%, SH-II: 11.5-21%, av. 17%) in both records; <i>Fagus</i> increase in SH-II (max.1%); <i>Chenopodiaceae</i> increase in SH-WOOD, decrease in SH-II; <i>Anthemis</i> -type increase in SH-WOOD; <i>Triticum</i> -type pollen is present in SH-WOOD; <i>Cyperus</i> and <i>Cyperaceae</i> pollen maximum in SH-WOOD; decreasing microcharcoal concentration in SH-WOOD; AP:NAP ratio decrease in SH-WOOD, increase in SH-II; palynological richness ranges 20-22 in SH-WOOD and 21-23 in SH-II
Middle Neolithic Szatmár II, Early AVK	7450 – 7250 cal B.P. 5500 -5300 cal B.C.	<i>Corylus</i> (SH-WOOD: 16-19%, av. 18%, SH-II: 19%) and <i>Quercus</i> (SH-WOOD: 17-32%, av. 24.8%, SH-II: 24%) relative frequency fluctuation in both records but in different direction; <i>Triticum</i> -type, <i>Secale</i> and crop field weed pollen present in SH-II, while <i>Triticum</i> and crop field weed pollen in SH-WOOD; Apophyta pollen and <i>Artemisia</i> increase in SH-WOOD; AP:NAP ratio minimum in both records; microcharcoal concentration peak in SH-WOOD; palynological richness ranges 22.5-25 in SH-WOOD and 21-25 in SH-II.
Middle Neolithic Late AVK	7250 – 6950 cal B.P. 5300 – 5000 cal B.C.	<i>Fraxinus excelsior/angustifolia</i> type pollen increase in SH-WOOD (max. 5.2%); <i>Carpinus betulus</i> episodic increase in SH-II (3%); <i>Artemisia</i> increase in SH-WOOD (from 4.5 to 10%); wet meadow and dry pasture indicator herb increase in SH-WOOD, but decrease in SH-II after an initial peak; <i>Triticum</i> -type and <i>Secale</i> pollen present in SH-WOOD; foothpath and ruderal herb increase in SH-II; AP:NAP ratio decrease following an episodic increase at 7230 cal B.P. in SH-WOOD, AP:NAP ratio increase in SH-WOOD; palynological richness ranges 22.5-25 in SH-WOOD and 18.5-24 in SH-II; <i>Corylus</i> pollen frequencies in SH-WOOD: 20.3-20.7%, av. 20.5%, in SH-II: 25.2-25.6%, av. 25.4%; <i>Quercus</i> pollen frequencies in SH-WOOD: 16-24%, av. 21%, in SH-II: 15-17%, av. 16%
Late Neolithic Tisza-Herpály-Csőszhalom	6950 – 6450 cal B.P. 5000 – 4500 cal B.C.	Further increase in <i>Artemisia</i> relative frequencies in SH-WOOD (max. 11.4%), but not in SH-II; stable <i>Corylus</i> (SH-WOOD:18.7-23% ; SH-II: 27-30%) and <i>Quercus</i> pollen frequencies (SH-WOOD: 18-21%; SH-II: 16.2-18%); <i>Carpinus betulus</i> gradual increase in SH-WOOD (max. 2%); generally low, but fluctuating AP:NAP ratios (SH-WOOD: 1.4-2.3, SH-II: 2.8-3.1); single microcharcoal concentration peak at 6830 cal B.P. (4880 cal B.C.) in SH-WOOD; wet meadow and dry pasture herbs increase in SH-WOOD until 6730 cal B.P. (4780 cal B.C.), decrease afterwards; <i>Triticum</i> -type and <i>Secale</i> present in SH-WOOD; <i>Cyperus</i> and <i>Cyperaceae</i> pollen frequency decrease in SH-WOOD; palynological richness ranges 24-29.5 in SH-WOOD and 18.5-23 in SH-II
Early Copper Age Tiszapolgár	6450 – 5950 cal B.P. 4500 – 4000 cal B.C.	<i>Quercus</i> relative frequencies increase in SH-II (max. 25.6%), <i>Corylus</i> decrease in SH-II (from 30% to 19%); <i>Umbelliferae</i> increase in SH-WOOD (from 2 to 7%); significant AP:NAP ratio decrease in SH-WOOD (from 2 to 1.4), and small-scale decrease in SH-II; microcharcoal concentration maximum at 6100 cal B.P. (4150 cal B.C.) and subsequent increase in <i>Fraxinus excelsior/angustifolia</i> pollen in SH-WOOD; <i>Triticum</i> -type and <i>Secale</i> pollen present in SH-II; meadow indicators are high in SH-WOOD and gradually increase in SH-II; small increase in <i>Sparganium/Typha angustifolia</i> pollen in SH-WOOD; palynological richness ranges 21-27 in SH-WOOD and 20-25.5 in SH-II

Plant remains from Late Neolithic sites in the Polgár Area	Equivalent pollen types in SH-II & SH-WOOD
<b>Domestic plants</b>	
<i>Triticum monococcum</i>	Triticum-type
<i>Triticum dicoccum</i>	
<i>Triticum turgidum/durum</i>	
<i>Triticum aestivum</i>	
<i>Triticum compactum</i>	
<i>Hordeum vulgare</i>	Cereales > 40 µm
<i>Avena</i> sp.	– ( <i>Avena</i> -type)
<i>Panicum miliaceum</i> (GY-CSH)	Gramineae
<i>Lens</i> sp.	– ( <i>Lathyrus/Vicia</i> )
<i>Pisum sativum</i>	– ( <i>Pisum</i> )
<i>Pisum elatius</i>	
<i>Vicia</i> spp.	– ( <i>Lathyrus/Vicia</i> )
<i>Vicia</i> cf. <i>V. sativa</i> (GY-CSH)	
<i>Linum usitatissimum</i>	– ( <i>Linum</i> )
<b>Wild plants</b>	
<i>Polygonum arenastrum</i>	<i>Polygonum aviculare</i> -type
<i>Hordeum bulbosum</i> (GY-CSH)	Cereales > 40 µm
<i>Polygonum</i> cf. <i>P. persicaria</i>	<i>Polygonum persicaria</i> -type
<i>Fallopia convolvulus</i>	– ( <i>Fallopia convolvulus</i> )
<i>Chenopodium hybridum</i>	Chenopodiaceae
<i>Chenopodium polyspermum</i>	
<i>Chenopodium vulparia</i>	
<i>Chenopodium murale</i>	
<i>Chenopodium album</i>	
<i>Chenopodium/Atriplex</i> spp.	
<i>Papaver</i> spp.	<i>Papaver</i>
<i>Solanum nigrum</i>	<i>Solanum nigrum/Physalis</i>
<i>Galium</i> sp.	Galium
<i>Galium</i> cf. <i>verum</i>	
<i>Galium</i> cf. <i>palustre</i>	
Gramineae indet.	Gramineae
<i>Bromus</i> spp.	
<i>Phleum</i> sp.	
<i>Phalaris</i> sp.	
<i>Stipa</i> sp.	
<i>Setaria</i> sp.	
<i>Schoenoplectus lacustris</i>	
<i>Carex</i> spp.	
<i>Juncus</i> spp.	<i>Juncus</i>
Caryophyllaceae spp.	Caryophyllaceae undiff.
<i>Stellaria media/Cerastium</i> sp.	<i>Cerastium</i> -type/ <i>Stellaria</i> -type
<i>Sagina</i> sp.	<i>Sagina</i>
<i>Lychnis</i> sp.	<i>Lychnis</i> -type
<i>Silene</i> sp. (CSH)	<i>Silene</i>
<i>Fragaria</i> sp.	<i>Fragaria</i>
<i>Prunella vulgaris</i>	<i>Prunella</i>
Liliaceae indet.	– ( <i>Liliaceae</i> undiff.)
<i>Dryopteris filix-mas</i>	<i>Dryopteris</i>
<i>Rumex acetosella</i> (CSH)	<i>Rumex acetosa/acetosella</i>
<i>Rumex</i> sp. (CSH)	<i>Rumex</i> sp.
<i>Rumex hydrolapathum</i> (GY-CSH)	
<i>Potentilla</i> sp. (CSH)	<i>Potentilla</i>
Leguminoseae indet. (CSH)	Leguminoseae undiff.
Compositae indet. (CSH)	Compositae Subfamily Liguliflorae / Compositae Subfamily Tubuliflorae
<b>Trees &amp; Shrubs</b>	
<i>Corylus</i> sp.	<i>Corylus</i>
<i>Prunus</i> cf. <i>P. spinosa</i>	<i>Prunus</i>
<i>Cornus mas</i>	<i>Cornus mas</i>
<i>Sambucus nigra</i>	<i>Sambucus</i>

**Additional file 1** Comparative table of plant macrofossil taxa recorded by the archaeobotanical investigation of Late Neolithic settlements in the Polgár area and their equivalent pollen types in the Sarló-hát (SH-II, SH-WOOD) pollen diagrams. The Late Neolithic settlement, Csőszhalom was analysed by Fairbairn (1992, 1993) and the main core of the taxon list enumerates his data. In addition, taxa not found by Fairbairn but determined by Gyulai (2010) in course of an independent survey of the Csőszhalom tell are shown and marked (GY-CSH).

Figure 1




Figure 2





Figure 3

807

808

809



294

293

0 .5km

Marsh  
Fieldwalked areas  
Sites

Road

Road

DOB31 DOB30 DOB29 DOB28  
SH-WOOD SH-II

TG9

Figure 4




Figure 5

# SH-WOOD






Figure 6




Figure 7




Figure 8





Figure 9

Nagymohos (394 m a.s.l.)



Bábtava (106 m a.s.l.)




Figure 10

