




Contents lists available at ScienceDirect

Advances in Mathematics

www.elsevier.com/locate/aim

Rasmussen's spectral sequences and the $\mathfrak{sl}_N$ -concordance invariants

Lukas Lewark¹*Mathematical Sciences, Durham University, South Road, Durham DH1 3LE, UK*

ARTICLE INFO

Article history:

Received 22 November 2013

Accepted 5 April 2014

Available online 25 April 2014

Communicated by Tomasz S. Mrowka

MSC:

57M25

Keywords:

Knot concordance

Khovanov–Rozansky homologies

Slice genus

Rasmussen invariant

Spectral sequences

Pretzel knots

ABSTRACT

Combining known spectral sequences with a new spectral sequence relating reduced and unreduced $\mathfrak{sl}_N$ -homology yields a relationship between the HOMFLYPT-homology of a knot and its $\mathfrak{sl}_N$ -concordance invariants. As an application, some of the $\mathfrak{sl}_N$ -concordance invariants are shown to be linearly independent.

© 2014 The Author. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

Contents

1.	Introduction	60
2.	The Khovanov–Rozansky homologies	63
2.1.	Unreduced homology	63
2.2.	Reduced homology	64
2.3.	Filtered homology	65
2.4.	HOMFLYPT-homology	66

E-mail address: lukas.lewark@durham.ac.uk.

URL: <http://www.maths.dur.ac.uk/~vxhn54>.

¹ Supported by the EPSRC grant EP/K00591X/1.

<http://dx.doi.org/10.1016/j.aim.2014.04.003>

0001-8708/© 2014 The Author. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/3.0/>).

2.5.	The relationship of HOMFLYPT-homology and reduced $\mathfrak{sl}_N$ -homology	66
2.6.	Calculating HOMFLYPT-homology	67
3.	The reduced–unreduced spectral sequence	69
4.	From HOMFLYPT-homology to the $\mathfrak{sl}_N$ -concordance invariants	70
5.	Slice-torus knot concordance invariants	74
6.	Linear independence of some of the $\mathfrak{sl}_N$ -concordance invariants	78
	Acknowledgments	81
	References	82

1. Introduction

The Khovanov–Rozansky homologies [20,21] are categorifications of the HOMFLYPT-polynomial and its specialisations. There is a wealth of different homology theories. As an astonishing and powerful application, some of them induce concordance invariants which give lower bounds for the smooth slice genus of a knot [47,57,31,33]. This paper has two goals: to explore how the different Khovanov–Rozansky homologies are related; and to show that the said concordance invariants, though very close to each other, are not equal. Apart from trying to deepen the understanding of the Khovanov–Rozansky homologies themselves, there is a geometric motivation: for example, the Khovanov–Rozansky concordance invariants may detect free summands in the group of topologically slice modulo smoothly slice knots [29]; and they could even be used to disprove the smooth Poincaré conjecture [6]. The main result is the following:

Theorem 1. *Let τ be the concordance invariant from knot Floer homology [41,44], and for all $N \geq 2$, s_N the concordance invariant from $\mathfrak{sl}_N$ -homology (see Proposition 2.6).*

- (i) *Neither τ nor s_2 is a linear combination of $\{s_N\}_{N \geq 3}$, and for any fixed $N \geq 3$, $\{\tau, s_2, s_N\}$ are linearly independent. (Proof without computer calculations.)*
- (ii) *s_3 is not a linear combination of $\{s_N\}_{N \geq 4}$, and for any fixed $N \geq 4$, $\{\tau, s_2, s_3, s_N\}$ are linearly independent. (Proof relies on computer calculations.)*

This prompts the following conjecture:

Conjecture. *The concordance invariants $\{\tau\} \cup \{s_i\}_{i \geq 2}$ are linearly independent.*

See [13,15] for similar results in Heegaard–Floer homology.

The Khovanov–Rozansky homologies considered in this paper are: the triply graded homology $\overline{\llbracket \cdot \rrbracket}_\infty$ categorifying the HOMFLYPT-polynomial; for each $N \geq 1$, the doubly graded reduced homology $\overline{\llbracket \cdot \rrbracket}_N$ and unreduced homology $\llbracket \cdot \rrbracket_N$ categorifying the reduced and unreduced $\mathfrak{sl}_N$ -polynomial, respectively; and the deformation of the unreduced homology, the filtered homology $\llbracket \cdot \rrbracket_N^f$. On the uncategorified level, the different polynomials are specialisations or multiples with a fixed factor of one another. On the categorified

level, relations normally take the form of spectral sequences; but the understanding of the interdependence of the different Khovanov–Rozansky homologies is far from complete. The following theorem clarifies the relationship between unreduced and reduced $\mathfrak{sl}_N$ -homology. Let D be a diagram of a link L with a marked component, and $N \geq 1$ an integer. Let $C_N(D)$ be the graded chain complex defined by Khovanov and Rozansky [20] whose homology is $[[L]]_N$.

Theorem 2. *There is a filtration of $C_N(D)$ respected by the differential such that the induced spectral sequence satisfies the following properties (where r denotes the grading associated to the new filtration):*

- (i) *Its differentials respect the q -degree; it converges on the N -th page, and forgetting the r -grading, the limit is isomorphic to $[[L]]_N$.*
- (ii) *Its first page is isomorphic to*

$$\frac{(qr)^N - (qr)^{-N}}{qr - (qr)^{-1}} \cdot \overline{[[L]]}_N.$$

- (iii) *The higher pages are invariants of links with a marked component.*

Let us have a closer look at the Khovanov–Rozansky concordance invariants. They belong to the broader class of slice-torus invariants introduced (without a name) by Livingston [28]:

Definition. A *slice-torus knot invariant* is a homomorphism ν from the smooth knot concordance group to the real numbers that satisfies the following conditions:

- (**slice**) For all knots K , $\nu(K)$ is a lower bound to twice the slice genus: $\nu(K) \leq 2g_4(K)$.
- (**torus**) For positive torus knots, this bound is sharp, i.e.

$$\forall p, q \in \mathbb{Z}^+: \quad (p, q) = 1 \quad \implies \quad \nu(T(p, q)) = 2g_4(T(p, q)) = (p-1)(q-1).$$

Note that we chose a different normalisation than Livingston. More importantly, we consider real-valued instead of integer-valued invariants, in order to include the normalised Khovanov–Rozansky concordance invariants. However, every slice-torus invariant discussed in this text takes values only in $\frac{1}{n}\mathbb{Z}$ for some fixed n .

Slice-torus knot invariants form a closed convex subset of the space of all real concordance homomorphisms. The slice-torus conditions are quite restrictive; all slice-torus invariants have e.g. the same value on quasi-positive knots. In this paper, the sharper slice-Bennequin inequality is generalised to slice-torus invariants, thereby showing that all slice-torus invariants have the same value on homogeneous knots (see Theorem 5 in Section 5).

Up to convex linear combination, 2τ (shown to be distinct from the Rasmussen invariant in [12]) is the only known slice-torus invariant not stemming from the Khovanov–Rozansky homologies. The oldest Khovanov–Rozansky concordance invariant is the Rasmussen invariant s_2 [47], which comes from Khovanov homology over a field of characteristic 0. Later on, generalisations were defined: an invariant s_N coming from $\mathfrak{sl}_N$ -homology for arbitrary $N \geq 2$ [33]; and an invariant $s_2^{\mathbb{F}_p}$, obtained from $\mathfrak{sl}_2$ -homology over a prime field $\mathbb{F}_p$ [2,53] (see also [34]). So far, computer calculations indicate that $s_2 \neq s_2^{\mathbb{F}_2}$ [27,51]; and that $s_3 \neq s_2$ [24,25]. The following theorem, whose proof uses [Theorem 2](#), the Rasmussen spectral sequences [45] and the Lee–Gornik spectral sequences [23,8], is a means to distinguish the s_N .

Theorem 3. *Let K be a knot. For all $N \geq 2$, let*

$$X_N = \{ \alpha + N\beta \mid \overline{[K]}_\infty \text{ has a generator of degree } q^\alpha a^\beta \text{ in homological degree } 0 \} \subset 2\mathbb{Z}.$$

Then for the unnormalised $\mathfrak{sl}_N$ -concordance invariant s'_N , we have $\min X_N \leq s'_N(K) \leq \max X_N$, or equivalently, for the normalised invariant:

$$\frac{\max X_N}{1-N} \leq s_N(K) \leq \frac{\min X_N}{1-N}.$$

[Theorem 3](#) combined with the sharper slice–Bennequin inequality enables us to calculate the $\mathfrak{sl}_N$ -concordance invariants of certain three-stranded pretzel knots:

Theorem 4. *Let ℓ and m be odd integers and $\ell > m \geq 3$. Then*

$$\begin{aligned} \text{(i)} \quad & s_2(P(\ell, -m, 2)) = \ell - m, \quad \text{and} \\ \text{(ii)} \quad & \forall N \geq 3: \quad s_N(P(\ell, -m, 2)) \in \left\{ \ell - m - 2, \ell - m - 2 + \frac{2}{(N-1)} \right\}. \end{aligned}$$

Let $n \geq 4$ be an even integer. Then

$$\begin{aligned} \text{(iii)} \quad & s_2(P(\ell, -m, n)) = \begin{cases} \ell - m & \text{if } m > n \\ \ell - m - 2 & \text{if } m < n, \end{cases} \quad \text{and} \\ \text{(iv)} \quad & \forall N \geq 3: \quad s_N(P(\ell, -m, n)) = \ell - m - 2. \end{aligned}$$

These pretzel knots are the examples used to prove [Theorem 1](#). The remainder of the paper is organised as follows: [Section 2](#) details the needed results on the Khovanov–Rozansky homologies. So while this paper is de facto self-contained, nevertheless some familiarity with [20] is advisable. The two following sections contain the proofs of [Theorem 2](#) and [Theorem 3](#), respectively. In [Section 5](#), known results about slice-torus invariants are collected, and the sharper slice Bennequin inequality ([Theorem 5](#)) is proven. [Section 6](#) finally applies the tools to examples such as the family of pretzel knots and contains the proofs of [Theorem 4](#) and [Theorem 1](#).




Fig. 1. The $(5, -3, 2)$ -pretzel knot K with $s_2(K) = 2$, $s_3(K) = 1$ and $s_N(K) \in \{0, 2/(N-1)\}$ for all $N \geq 4$.

2. The Khovanov–Rozansky homologies

This section gives an overview over the different Khovanov–Rozansky homologies and the known spectral sequences relating them. Notations and conventions are clarified on the way. Comparing Khovanov–Rozansky homologies over different base fields leads to interesting open questions (cf. [27,34,53]). However, we restrict our attention to characteristic 0, and consider all chain complexes to be over the complex numbers.

2.1. Unreduced homology

Let D be a diagram of a link L . For all integers $N \geq 1$, Khovanov and Rozansky [20] define a chain complex $C_N(D)$ of graded vector spaces (technically, it is a cochain complex). Any Reidemeister move gives rise to a quasi-isomorphism of $C_N(D)$, and so the homology $[[L]]_N$ is a link invariant, called the $\mathfrak{sl}_N$ -homology. Note that all links have the same $\mathfrak{sl}_1$ -homology, while $\mathfrak{sl}_2$ -homology is isomorphic to Khovanov homology [18], and $\mathfrak{sl}_3$ -homology to a homology theory defined previously via webs and foams [19,35].

We regard this chain complex and his homology as a doubly graded vector space, with a *homological* (t), and a *quantum* (q) degree. In general, for such a graded space V , let us write V^i for the subspace of homological degree i , and, if V has finite dimension, $\text{xdim } V \in \mathbb{N}[t^{\pm 1}, q^{\pm 1}]$ (where $\mathbb{N} = \{0, 1, 2, \dots\}$) for the graded dimension of V . If f is a homogeneous homomorphism of such spaces and has (t, q) -degree (i, j) , we denote by $\text{xdeg } f$ the monomial $t^i q^j$.

The $\mathfrak{sl}_N$ -homology categorifies the $\mathfrak{sl}_N$ -polynomial P_N [48], i.e. $\text{xdim}([L]_N)(-1, q) = P_N(L)$. The $\mathfrak{sl}_N$ -polynomial is given by its value of $[N]_q = (q^{-N+1} + q^{-N+3} + \dots + q^{N-1})$ on the unknot and the following skein relation:

$$q^N \cdot P_N \left(\begin{array}{c} \text{crossing} \end{array} \right) - q^{-N} \cdot P_N \left(\begin{array}{c} \text{crossing} \end{array} \right) = (q - q^{-1}) \cdot P_N \left(\begin{array}{c} \text{two crossings} \end{array} \right).$$

In fact, the $\mathfrak{sl}_N$ -homology theory is richer than that, being defined for tangles as well: to every tangle diagram D with boundary ∂D (the boundary being a finite sequence

of signs) an object in some category $\mathcal{C}_{\partial D}$ is associated. This is done in such a way that the gluing of tangle diagrams corresponds to the tensor product of the associated objects (i.e. this is a *canopolis*-construction, cf. [2]). The category $\mathcal{C}_{\emptyset}$ is equivalent to the category of graded chain complexes over $\mathbb{C}$. In this way, Reidemeister invariance of the $\mathfrak{sl}_N$ -homology of a link can be proven simply by showing that the objects associated to the two small tangle diagrams which correspond to each of the Reidemeister moves are isomorphic.

2.2. Reduced homology

The reduced version of this homology categorifies the reduced $\mathfrak{sl}_N$ -polynomials $\overline{P}_N = P_N/[N]_q$. Let D be a diagram of the link L with a marked component. Let $A = \mathbb{C}[X]/(X^N)$, an algebra with grading $\deg X^i = 2i$ for $i \in \{0, \dots, N-1\}$. Then $C_N(D)$ has the structure of a free graded A -module. This structure is respected by the differential of $C_N(D)$, and it may depend on the choice of the marked component. Let $\tilde{\mathcal{C}}$ be the graded A -module $A/(X)$ with a shift of $N-1$ in the q -grading. Let $\overline{C}_N(D) = C_N(D) \otimes_A \tilde{\mathcal{C}}$. The following proposition, which is essential for the proof of Theorem 2(iii), is implicit in [20, end of section 7]. Let us give an explicit proof.

Proposition 2.1. *If two base-pointed diagrams D and D' are connected by a Reidemeister move that avoids the base-point, then there is a chain homotopy equivalence respecting the A -module structure between $C_N(D)$ and $C_N(D')$.*

Proof. This proof supposes greater familiarity with the details of the construction [20] than the rest of the paper: in fact, we consider link diagrams D with *marks*. Marks form a finite subset of D that avoids the crossings, such that any interval connecting two crossings contains at least one mark. The complement of the marks is the disjoint union of components, all of which are either a positive or negative crossing, a line or a circle. The chain complex $C_N(D)$ is then defined as the tensor product (over adequate rings) of the elementary chain complexes associated to these simple pieces. Adding or removing marks produces a homotopy equivalent chain complex, and this chain homotopy equivalence respects the A -module structure (this follows from [20, Proposition 22]). So, without loss of generality, assume that $D = D_1 \cup D_3$ and $D' = D_2 \cup D_3$ each split along marks into two tangle diagrams: small ones D_1 and D_2 , in which the Reidemeister move takes place, who have the same complement D_3 . Then, $C_N(D) = C_N(D_1) \otimes C_N(D_3)$ and $C_N(D') = C_N(D_2) \otimes C_N(D_3)$. These tensor products are A -modules because the second factor is. There is a chain homotopy equivalence φ between $C_N(D_1)$ and $C_N(D_2)$. The tensor product of φ with the identity of $C_N(D_3)$ gives a chain homotopy equivalence between $C_N(D)$ and $C_N(D')$ that respects the A -module structure. $\square$

Corollary 2.2. (See [20].) *The homology $\overline{\mathbb{L}}_N$ of the reduced complex is an invariant of links with a marked component.*

Proof. Two base-pointed diagrams D and D' represent the same base-pointed link if and only if they are connected by a finite sequence of Reidemeister moves which avoid the base point. $\square$

2.3. Filtered homology

There is a filtered version of $\mathfrak{sl}_N$ -homology, whose associated graded is the original unreduced $\mathfrak{sl}_N$ -homology. As usual, a filtered complex gives rise to a spectral sequence. Let us briefly clarify the indexing convention: a spectral sequence is a sequence $(E_k)_{k \in \{0,1,2,\dots\}}$ of graded chain complexes, such that for all $k \geq 0$, forgetting the differential on E_{k+1} yields the homology of E_k . If not specified otherwise, the differential d_k on the k -th page has (t, q) -degree $(1, k)$; this is non-standard, but convenient in our context.

Proposition 2.3. (See [23] for $N = 2$, [8] for all $N \geq 2$.) *There is a spectral sequence starting at unreduced $\mathfrak{sl}_N$ -homology and converging to filtered $\mathfrak{sl}_N$ -homology.*

Proposition 2.4. (See [23] for $N = 2$, [57, Theorem 1.2] for all $N \geq 2$.) *The higher pages of the Lee–Gornik spectral sequence are link invariants.*

The following detail has not been explicitly stated:

Proposition 2.5. *The differential on the k -th page of the Lee–Gornik spectral sequence vanishes unless k is a multiple of $2N$.*

Proof. Note that the differentials of Gornik’s filtered chain complex preserve the q -degree mod $2N$ (see [8]). So the chain complex decomposes as a direct sum of N terms (for $i \in \{0, \dots, N-1\}$, the i -th term containing the generators of q -degree equal to $2i$ mod $2N$), and so does the induced spectral sequence. $\square$

As a consequence, it makes sense to forget all pages with vanishing differentials and renumber: from now on, by the “ k -th page” of the Lee–Gornik spectral sequences, we actually refer to the $(2Nk)$ -th page. It is still an open conjecture that this spectral sequence converges on the second page.

Proposition 2.6. (See [47] for $N = 2$, [8, 57, 31, 33] for all $N \geq 3$.) *Let K be a knot. The filtered $\mathfrak{sl}_N$ -homology of K is isomorphic to the unreduced $\mathfrak{sl}_N$ -homology of the unknot, with a q -shift by some even integer which we denote by $s'_N(K)$. Its normalisation $s_N(K) = s'_N(K)/(1-N) \in \frac{2}{N-1}\mathbb{Z}$ is a slice-torus invariant called the $\mathfrak{sl}_N$ -concordance invariant.*

Note that unlike the Rasmussen and twice the τ -invariant, the $\mathfrak{sl}_N$ -concordance invariants of a knot need not be even integers, or integers at all.

2.4. HOMFLYPT-homology

Khovanov and Rozansky [21] introduce a chain complex $C_\infty(D)$ of doubly graded complex vector spaces defined for a braid diagram D . Its homology is a link invariant called the HOMFLYPT-homology, which categorifies the HOMFLYPT-polynomial $P_\infty \in \mathbb{Z}[q^{\pm 1}, a^{\pm 1}]$. The HOMFLYPT-polynomial is determined by its value of 1 on the unknot, and the following skein relation:

$$a \cdot P_\infty \left(\text{crossing} \right) - a^{-1} \cdot P_\infty \left(\text{crossing} \right) = (q - q^{-1}) \cdot P_\infty \left(\text{cup/cap} \right).$$

There are several versions of HOMFLYPT-homology. Rasmussen [45] e.g. works with a reduced and an unreduced version, and an interpolation of the two; but all these versions carry the same information (as is not the case for the reduced and unreduced version of $\mathfrak{sl}_N$ -homology). In this text, we stick to the reduced version, denoted by $\overline{\overline{\cdot}}_\infty$. For a knot this is, unlike the unreduced version, a finite dimensional space. We follow similar grading conventions as Mackaay and Vaz [36], but exchanging t and t^{-1} , i.e.

$$\text{xdim } \overline{\overline{T(3, 2)}}_\infty = a^{-2}q^2 + t^2a^{-2}q^{-2} + t^3a^{-4}.$$

In [45], Rasmussen follows still another grading convention; the monomial $q^i a^j t^k$ in that convention corresponds to the monomial $q^i a^j t^{(k-j)/2}$ in ours.

There is yet another version of HOMFLYPT- and $\mathfrak{sl}_N$ -homology, only defined for two-component links: *totally reduced homology*, denoted by $\overline{\overline{\cdot}}_\infty$ and $\overline{\overline{\cdot}}_N$, respectively. We will not give its definition, because we only ever use the totally reduced homology of the positive Hopf link (as calculated e.g. in [36]):

$$\text{xdim } \overline{\overline{T(2, 2)}}_\infty = (a^{-1}q^2 + ta^{-1} + t^2a^{-1}q^{-2} + t^3a^{-3}) \cdot t^{-1/2}.$$

2.5. The relationship of HOMFLYPT-homology and reduced $\mathfrak{sl}_N$ -homology

The Rasmussen spectral sequences show that in a certain sense, HOMFLYPT-homology is the stabilisation of the $\mathfrak{sl}_N$ -homologies as $N \rightarrow \infty$.

Proposition 2.7. (See [45].) *Let L be a link with a marked component. For every $N \geq 1$, there is a spectral sequence with first page $\overline{\overline{L}}_\infty$. Its limit is, after a regrading, isomorphic to the reduced $\mathfrak{sl}_N$ -homology of L . Explicitly, the regrading of the (t, q, a) -degree is $(i, j, \ell) \mapsto (i, j + N\ell)$. The differential on the k -th page of the spectral sequence has degree $tq^{2Nk}a^{-2k}$. The higher pages are invariants of links with a marked component. If L is a knot, then for sufficiently large N , this sequence converges on the first page.*




Fig. 2. The knots $K_{\pm}$ and the link L_0 .

2.6. Calculating HOMFLYPT-homology

It is in fact easier to calculate the HOMFLYPT-homology of a knot than its $\mathfrak{sl}_N$ -homology for some N . See [36] for an exemplary calculation. Let us present the part of the tool-kit which is necessary for the calculations in this paper. HOMFLYPT-homology is well-behaved under taking the connected sum:

Proposition 2.8. (See [45, Lemma 7.8].) *Let L_1 and L_2 be links, and L_3 any connected sum of L_1 and L_2 . Then $\overline{\mathbb{L}}_3 \cong \overline{\mathbb{L}}_1 \otimes \overline{\mathbb{L}}_2$.*

Definition. (See [46].) Let the δ -grading on $\overline{\mathbb{L}}_{\infty}$ be defined by $\delta(t^i q^j a^k) = 2i + j + 2k$. A knot is *KR-thin* if its HOMFLYPT-homology is supported in a single δ -degree that is equal to minus its signature.

Lemma 2.9. *The HOMFLYPT-homology of a KR-thin knot K is determined by its HOMFLYPT-polynomial $P_{\infty}(K)$ and its signature $\sigma(K)$:*

$$\text{xdim } \overline{\mathbb{L}}_{\infty} = (-t)^{-\sigma(K)/2} \cdot P_{\infty}(qt^{-1/2}, at^{-1}).$$

Proof. This immediately follows from the fact that $\text{xdim } \overline{\mathbb{L}}_{\infty}(-1, q, a) = P_{\infty}(q, a)$. $\square$

Proposition 2.10. (See [46], [45, Corollary 1].) *Two-bridge knots are KR-thin.*

Remark 2.11. Quasi-alternating links are a generalisation of alternating links introduced in [42]. Quasi-alternating links have thin Khovanov and knot Floer homology [39], and in particular the Rasmussen and twice the τ -invariant of quasi-alternating knots equal their signature. This can be proven via an unoriented skein relation. For $N \geq 3$, however, $\mathfrak{sl}_N$ -homology does not satisfy such a relation, and indeed there are even alternating knots which are not KR-thin [45]. Still, the $\mathfrak{sl}_N$ -concordance invariants of alternating knots (but not, in general, of quasi-alternating knots) equal their signature for all N , since this is true of all slice-torus invariants (see Corollary 5.9).

Proposition 2.12 (The skein long exact sequences). (See [45, Lemma 7.6].) *Let K_+ , K_- and L_0 be two knots and one two-component link which look the same everywhere except near one crossing, where they differ as shown in Fig. 2. Then for all $N \geq 2$, there is a*

long exact sequence

$$\cdots \longrightarrow \overline{\overline{K_-}}_N \xrightarrow{(-N, \frac{1}{2})} \overline{\overline{L_0}}_N \xrightarrow{(-N, \frac{1}{2})} \overline{\overline{K_+}}_N \xrightarrow{(2N, -2)} \overline{\overline{K_-}}_N \longrightarrow \cdots.$$

The maps' (t, q) -degree is indicated above the arrows.

This proposition refers to $\mathfrak{sl}_N$ -homology; to make a statement about HOMFLYPT-homology, we need the following technical lemma. Let $\leq$ denote the partial order of polynomials given as follows: $A \leq B$ if and only if there is a polynomial C with non-negative coefficients such that $A + C = B$.

Lemma 2.13. *Let $A, B \in \mathbb{N}[q^{\pm 1}, a^{\pm 1}]$. Suppose that for infinitely many N , $A(q, q^N) \leq B(q, q^N)$. Then $A(q, a) \leq B(q, a)$.*

Proof. Let $i_{\max}$ and $i_{\min}$ be the maximal and minimal exponent of q occurring in A or B . Choose N such that $A(q, q^N) \leq B(q, q^N)$ and $|N| > i_{\max} - i_{\min}$. Then different monomials in $A(q, a)$ and $B(q, a)$ yield different monomials in $A(q, q^N)$ and $B(q, q^N)$. To show this, consider two monomials $c \cdot q^i a^j$ and $c' \cdot q^{i'} a^{j'}$ in $A(q, a)$ (with $c, c' \neq 0$). Then $cq^{i+Nj} = c'q^{i'+Nj'}$ implies $c = c'$ and $i + Nj = i' + Nj' \implies i - i' = N(j' - j) \implies |N| \cdot |j' - j| \leq i_{\max} - i_{\min} \implies j' = j \implies i = i'$. Let c_{ij} and c'_{ij} be the coefficients of the monomial $q^i a^j$ in $A(q, a)$ and $B(q, a)$, respectively. Then c_{ij} and c'_{ij} are also the respective coefficients of the monomial q^{i+Nj} in $A(q, q^N)$ and $B(q, q^N)$, and thus $c_{ij} \leq c'_{ij}$. $\square$

Corollary 2.14. *Suppose $K_{\pm}$ and L_0 are given as in Fig. 2, then*

$$\begin{aligned} \text{xdim } \overline{\overline{K_+}}_{\infty} &\leq t^2 \cdot a^{-2} \cdot \text{xdim } \overline{\overline{K_-}}_{\infty} + t^{1/2} \cdot a^{-1} \cdot \text{xdim } \overline{\overline{L_0}}_{\infty} \quad \text{and} \\ \text{xdim } \overline{\overline{K_-}}_{\infty} &\leq t^{-2} \cdot a^2 \cdot \text{xdim } \overline{\overline{K_+}}_{\infty} + t^{-1/2} \cdot a \cdot \text{xdim } \overline{\overline{L_0}}_{\infty}. \end{aligned}$$

Proof. We will just prove the first equation, the second one follows similarly. The long exact sequence can be broken up into short ones; i.e., for some quotient space A of $\overline{\overline{L_0}}_N$ and subspace B of $\overline{\overline{K_-}}_N$ there is a short exact sequence

$$0 \longrightarrow A \xrightarrow{(-N, \frac{1}{2})} \overline{\overline{K_+}}_N \xrightarrow{(-2N, 2)} B \longrightarrow 0.$$

This is equivalent to $\overline{\overline{K_+}}_N \cong (q^{-N} t^{1/2} \cdot A) \oplus (q^{-2N} t^2 \cdot B)$. In terms of graded dimensions, this implies

$$\begin{aligned} \text{xdim } \overline{\overline{K_+}}_N &= q^N \cdot t^{-1/2} \cdot \text{xdim } A + q^{2N} \cdot t^{-2} \cdot \text{xdim } B \\ \implies \text{xdim } \overline{\overline{K_+}}_N &\leq q^N \cdot t^{-1/2} \cdot \text{xdim } \overline{\overline{L_0}}_N + q^{2N} \cdot t^{-2} \cdot \text{xdim } \overline{\overline{K_-}}_N. \end{aligned}$$

For large enough N , the three polynomials in this inequality stabilise (see Proposition 2.7), i.e.

$$\begin{aligned}
(\mathrm{xdim} \, \overline{\overline{K_+}}_\infty)(q, q^N) &= (\mathrm{xdim} \, \overline{\overline{K_+}}_N)(q), \\
(\mathrm{xdim} \, \overline{\overline{L_0}}_\infty)(q, q^N) &= (\mathrm{xdim} \, \overline{\overline{L_0}}_N)(q), \\
(\mathrm{xdim} \, \overline{\overline{K_-}}_\infty)(q, q^N) &= (\mathrm{xdim} \, \overline{\overline{K_-}}_N)(q).
\end{aligned}$$

So using [Lemma 2.13](#), the statement follows. $\square$

Remark 2.15. It may seem cumbersome to prove [Corollary 2.14](#) by first considering the $\mathfrak{sl}_N$ -homologies, and then HOMFLYPT-homology as their stabilisation. But as Rasmussen remarks, who proves the KR-thinness of two-bridge knots in the same way [\[45\]](#), it is unclear how to work directly on HOMFLYPT-homology.

3. The reduced–unreduced spectral sequence

This section is devoted to the proof of [Theorem 2](#). We need the following technical lemma, whose proof is left to the reader.

Lemma 3.1. *Let (C, ∂) be a filtered chain complex, whose filtration we denote by $\mathcal{F}$. Let there be an additional grading $C = \bigoplus_{i \in \mathbb{Z}} C_i$ that is respected by the differential. The filtration $\mathcal{F}$ induces a filtration on each C_i by $\mathcal{F}_j C_i := C_i \cap \mathcal{F}_j C$. If C , as a filtered vector space, is the sum of the filtered C_i , we say that the filtration is compatible with the additional grading. In this case, the spectral sequence induced by $\mathcal{F}$ respects the additional grading on C .*

Proof of Theorem 2. (i): Recall that $C_N(D)$ is a module over $A = \mathbb{C}[X]/(X^N)$. Let us introduce a filtration $\mathcal{R}$ on A , given by $\mathcal{R}_{2i-N+1}A = (X^i)$. Explicitly, we have

$$A = \mathcal{R}_{-N+1}A \supset \mathcal{R}_{-N+3} \supset \dots \supset \mathcal{R}_{N-1}A \supset \{0\}.$$

This induces a filtration on $C_N(D)$. Let us denote it by $\mathcal{R}$ as well, and call the induced grading the r -grading. Since the differential of $C_N(D)$ commutes with the A -scalar multiplication, it also respects this filtration. So $\mathcal{R}$ induces a spectral sequence $E_\bullet$, which (forgetting the additional degree) converges to $\overline{[L]}_N$. Note moreover that $\mathcal{R}$ and the q -grading are compatible in the sense of [Lemma 3.1](#). Hence $E_\bullet$ respects the q -degree, and its differential on the k -th page has degree $tr^{2k}q^0$.

(ii): Let us analyse the 0-th page of that spectral sequence, i.e. the associated graded chain complex. We have $\overline{C_N(D)} = C_N(D) \otimes_A \tilde{\mathbb{C}}$. This is isomorphic to $C_N(D)/((X) \cdot C_N(D)) = \mathcal{R}_{-N+1}C_N(D)/\mathcal{R}_{-N+3}C_N(D)$, with a q -shift of $N-1$. Note that for $i \in \{0, \dots, N-1\}$, the multiplication by X^i is an isomorphism between $\mathcal{R}_{-N+1}A/\mathcal{R}_{-N+3}A$ and $\mathcal{R}_{2i-N+1}A/\mathcal{R}_{2i-N+3}A$. Since $C_N(D)$ is a free A -module, this is true for $C_N(D)$ as well: the multiplication by X^i is an isomorphism

$$\mathcal{R}_{-N+1}C_N(D)/\mathcal{R}_{-N+3}C_N(D) \rightarrow \mathcal{R}_{2i-N+1}C_N(D)/\mathcal{R}_{2i-N+3}C_N(D)$$

of complex vector spaces. Because the A -scalar multiplication commutes with the differential, this map is an isomorphism of chain complexes. It shifts the q -grading and the r -grading by $2i$. The 0-th page of the spectral sequence is the sum of the $\mathcal{R}_{2i-N+1}C_N(D)/\mathcal{R}_{2i-N+3}C_N(D)$, and thus

$$E_0 \cong \bigoplus_{i \in \{-N+1, -N+3, \dots, N-1\}} (qr)^i \cdot \bar{C}_N(D).$$

Taking homology yields the stated result for the first page.

(iii): To prove the invariance of the higher pages, let us use the following lemma proved e.g. in [40, Theorem 3.5] (also used by Rasmussen [47, Lemma 6.1]):

Lemma 3.2. *Let $f : C \rightarrow C'$ be a map of filtered chain complexes. Let $E_\bullet$ and $E'_\bullet$ be the respective spectral sequences associated to C and C' , and for all $r \geq 0$, let f_r be the induced graded map from E_r to E'_r . If f_R is an isomorphism for some R , then f_r is also an isomorphism for all $\infty \geq r \geq R$.*

So the map of Proposition 2.1 induces an isomorphism between the higher pages of the spectral sequence associated to diagrams related by a Reidemeister move. $\square$

Fig. 3 shows the reduced–unreduced spectral sequence for the pretzel knot $P(5, -3, 2)$.

4. From Homflypt-homology to the $\mathfrak{sl}_N$ -concordance invariants

The following lemma (a direct generalisation of [6, Theorem 5.1]) describes the decategorification of a spectral sequence:

Lemma 4.1. *Let $(E_\bullet, d_\bullet)$ be a spectral sequence of $\mathbb{Z}^n$ -graded finite dimensional vector spaces. Then for all $k \geq 1$ there are polynomials $f_k \in \mathbb{N}[x_1^{\pm 1}, \dots, x_n^{\pm 1}]$, such that for all $\ell \geq 1$ the following decomposition holds:*

$$\text{xdim } E_\ell = \text{xdim } E_{\ell+1} + \sum_{k=1}^{\ell} (1 + \text{xdeg } d_k) \cdot f_k.$$

In particular,

$$\text{xdim } E_1 = \text{xdim } E_\infty + \sum_{k=1}^{\infty} (1 + \text{xdeg } d_k) \cdot f_k.$$

The spectral sequence converges on the ℓ -th page if and only if $\forall k \geq \ell: f_k = 0$.




Fig. 3. On the left, the reduced $\mathfrak{sl}_3$ -homology of the $P(5, -3, 2)$ -pretzel knot. In the middle, the first page of the reduced-unreduced spectral sequence. The r -grading is expressed by colours (and underlining/italics). On this page, “red may kill black and black green”. On the second page (not shown), only “red may kill green”. The total differential on the first and second page has rank 2 and 5, respectively. On the right, the third page of homology, which is the limit of the spectral sequence. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 4. The proof of [Proposition 4.2](#) and [Theorem 3](#) in a nutshell. Double arrows stand for spectral sequences, single arrows for other relations.

Proposition 4.2. *Let K be a knot, and let $N \geq 2$. There are polynomials $\bar{P}'_N \in \mathbb{N}[t^{\pm 1}, q^{\pm 1}, a^{\pm 1}]$, $P'_N \in \mathbb{N}[t^{\pm 1}, q^{\pm 1}, r^{\pm 1}]$ and for all $k \geq 1$ polynomials $f_N^k \in \mathbb{N}[t^{\pm 1}, q^{\pm 1}, a^{\pm 1}]$, $g_N^k \in \mathbb{N}[t^{\pm 1}, q^{\pm 1}, r^{\pm 1}]$, $h_N^k \in \mathbb{N}[q^{\pm 1}, t^{\pm 1}]$, such that for large enough N we have $\forall k: f_N^k = 0$, and such that the following decompositions hold:*

$$\begin{aligned} \text{xdim } \overline{[K]}_\infty &= \bar{P}'_N(t, q, a) + \sum_{k=1}^{\infty} (1 + tq^{-2Nk} a^{2k}) f_N^k(t, q, a), \\ \text{xdim } \overline{[K]}_N &= \bar{P}'_N(t, q, q^N), \\ \text{xdim } \overline{[K]}_N \cdot [N]_{qr} &= P'_N(t, q, r) + \sum_{k=1}^{N-1} (1 + tr^{2k}) g_N^k(t, q, r), \\ \text{xdim } [K]_N &= P'_N(t, q, 1), \\ \text{xdim } [K]_N &= q^{s'_N(K)} \cdot [N]_q + \sum_{k=1}^{\infty} (1 + tq^{2Nk}) h_N^k(q, t). \end{aligned}$$

Proof. Apply [Lemma 4.1](#) to Rasmussen's spectral sequence ([Proposition 2.7](#)), the reduced–unreduced spectral sequence ([Theorem 2](#)) and the Lee–Gornik spectral sequence with renumbered pages ([Proposition 2.3](#)). See also [Fig. 4](#). $\square$

Given $\text{xdim } \overline{[K]}_\infty$, there are only finitely many choices for the auxiliary polynomials in the proposition and for the $s'_N(K)$. So HOMFLYPT-homology induces restrictions on the $\mathfrak{sl}_N$ -concordance invariants (and thus a lower bound on the slice genus). [Proposition 4.2](#) may appear unwieldy; and in fact, we will only use [Theorem 3](#), which skips all intermediary steps between the HOMFLYPT-homology and the $\mathfrak{sl}_N$ -concordance invariants.

Proof of Theorem 3. Let us use the equations of [Proposition 4.2](#), climbing from the bottom up:

$$\begin{aligned}
 q^{s'_N(K)-N+1} &\leq \text{xdim} \llbracket K \rrbracket_N^f \\
 \implies q^{s'_N(K)-N+1} &\leq \text{xdim} \llbracket K \rrbracket_N \\
 \implies q^{s'_N(K)-N+1} r^i &\leq P'_N \quad \text{for some } i \\
 \implies q^{s'_N(K)-N+1} r^i &\leq \text{xdim} \llbracket \overline{K} \rrbracket_N \cdot [N]_{qr} \\
 \implies q^{s'_N(K)-N+1+j} &\leq \text{xdim} \llbracket \overline{K} \rrbracket_N \quad \text{for some } j \in \{1-N, 3-N, \dots, N-1\} \\
 \implies q^\alpha a^\beta &\leq \overline{P}'_N \quad \text{for some } \alpha, \beta \text{ with } \alpha + N\beta \leq s'_N(K) \\
 \implies q^\alpha a^\beta &\leq \text{xdim} \llbracket \overline{K} \rrbracket_\infty.
 \end{aligned}$$

An analogous reasoning yields $q^{s'_N(K)+N-1} \leq \text{xdim} \llbracket K \rrbracket_N \implies q^{\alpha'} a^{\beta'} \leq \llbracket \overline{K} \rrbracket_\infty$ for some α', β' with $s'_N(K) \leq \alpha' + N\beta'$. $\square$

Note that the power of [Proposition 4.2](#) is limited:

Proposition 4.3. *Let K be a knot, and let $N \geq 2$. Suppose there are polynomials $i_N^0, \dots, i_N^{N-1} \in \mathbb{N}[t^{\pm 1}, a^{\pm 1}, q^{\pm 1}]$ and some α, β such that the following decomposition holds:*

$$\text{xdim} \llbracket \overline{K} \rrbracket_\infty = q^\alpha a^\beta + \sum_{k=0}^{N-1} (1 + ta^{-2} q^{2k}) \cdot i_N^k.$$

Then there is also a decomposition as in [Proposition 4.2](#) with $\alpha + N\beta$ at the place of $s'_N(K)$, i.e. the theorem cannot be used to show $s'_N(K) \neq \alpha + N\beta$.

Proof. Let N be fixed. Setting $f_N^k = 0$ for all k , $h_N^k = 0$ for $k \geq 2$ and

$$\begin{aligned}
 h_N^1 &= \sum_{k=0}^{N-1} [N-k]_q \cdot q^k \cdot i_N^k, \\
 g_N^k &= [N-k]_{qr} \cdot (qr)^{-k} \cdot i_N^{N-k}
 \end{aligned}$$

gives the desired decomposition. $\square$

Remark 4.4. If Rasmussen's spectral sequence from $\llbracket \overline{K} \rrbracket_\infty$ to the regraded version of $\llbracket \overline{K} \rrbracket_1$ converges on the second page, it gives a decomposition as in the above proposition, with $i_N^k = 0$ for $k \neq 1$ and $\alpha = -\beta$. So for any knot for which that spectral sequence converges on the second page, [Proposition 4.2](#) alone is not strong enough to distinguish the $\mathfrak{sl}_N$ -concordance invariants.




Fig. 5. A cobordism of Euler characteristic -1 inserting a positive crossing. The cobordism consists of a Reidemeister I move and a saddle move.

5. Slice-torus knot concordance invariants

This section is largely independent from Sections 2–4. It collects and extends what is known about slice-torus invariants. Let us start by listing some invariants that are not slice-torus: the classical knot signature σ or the concordance invariant δ from the Floer homology of double branched covers [38] give slice genus bounds, but they are not sharp for torus knots; the Lipshitz–Sarkar invariant [27] on the other hand is not a concordance homomorphism.

The following results are mainly due to Livingston [28], who built on the work of Rudolph [49]. Although Livingston only considers slice-torus invariants which take even integer values, his results and proofs carry over unchanged to the general case of real-valued invariants. Note that we opted for a different normalisation of the slice-torus invariants. Throughout this section, let ν denote an arbitrary slice-torus invariant. The proof of the following proposition is standard:

Proposition 5.1.

- (i) (See [28, Cor. 2].) For all knots K , the absolute value of ν is a lower bound to twice the slice genus: $|\nu(K)| \leq 2g_4(K)$.
- (ii) If there is a connected smooth cobordism of Euler characteristic χ between two knots K_0 and K_1 , then

$$|\nu(K_0) - \nu(K_1)| \leq -\chi.$$

Lemma 5.2. *There is a cobordism of Euler characteristic -1 inserting or resolving a positive or negative crossing,*

Proof. See Fig. 5. $\square$

Proposition 5.3. (See [28, Cor. 3].) *If K_+ and K_- are knots that have diagrams that are identical but for the sign of one crossing, which is given in the subscript (see Fig. 2), then*

$$0 \leq \nu(K_+) - \nu(K_-) \leq 2.$$

Lemma 5.4. (See [28, Cor. 7].) Let B be a positive braid, i.e. a braid whose word contains only the σ_i , not the σ_i^{-1} . Suppose B has n strands and k crossings. If the closure $\text{tr}(B)$ of B is a knot, then $\nu(\text{tr}(B)) = 2g_4(\text{tr}(B)) = 2g_3(\text{tr}(B)) = 1 + k - n$.

Quasi-positivity has been introduced and studied by Rudolph, see [49].

Definition. A braid B is said to be *quasi-positive* if it is the product of braid-words that are conjugate to one of the σ_i ; i.e. $B = \prod_j w_j \sigma_{i_j} w_j^{-1}$, where w is any braid-word.

Proposition 5.5. (See [49].) Let B be a quasi-positive braid with k_+ positive crossings, k_- negative crossings, writhe $w = k_+ - k_-$ and n strands. If $\text{tr}(B)$ is a knot, then $2g_4(\text{tr}(B)) = 1 + w - n$.

The following proposition has been proven for the Rasmussen invariant by Shumakovitch [52]; for 2τ it is an immediate consequence of the results of Plamenevskaya [43]. The relationship between the τ -invariant, quasi-positivity and fibredness were studied by Hedden [11].

Proposition 5.6. *Slice-torus invariants detect the slice-genus of quasi-positive knots.*

Proof. Let B be a quasi-positive braid. Let B' be the braid obtained by switching every negative crossing of B to a positive one. By Lemma 5.4, $\nu(\text{tr}(B')) = 1 + k_+ + k_- - n$, and by Proposition 5.3, $\nu(\text{tr}(B')) - \nu(\text{tr}(B)) \leq 2k_- \implies \nu(\text{tr}(B)) \geq 1 + w - n$. But $1 + w - n = g_4(\text{tr}(B))$ by the previous proposition. $\square$

Corollary 5.7. *Let D be a positive knot diagram of a knot K , i.e. a diagram with only positive crossings. Then ν detects the slice genus of K .*

Proof. Follows from the previous proposition since positive links are quasi-positive [50]. $\square$

Proposition 5.8. (See [47].) Let D be a positive knot diagram of a knot K , with k crossings, and n Seifert circles. Then $g_4(K) = 1 + k - n$.

One of the strongest restrictions that can be deduced from the slice-torus conditions is an inequality à la Bennequin [3]. The first version was proven by Rudolph [49] for the slice-genus, and by Rasmussen [47], Shumakovitch [52] and Plamenevskaya [43] for the Rasmussen invariant. It was subsequently sharpened by Kawamura [16]; her version was generalised by Wu to the s_N -invariants [56]. Then, it was honed yet more independently by Lobb [32] and Kawamura [17]. Given a diagram D of a knot K , the sharper slice-Bennequin inequality gives an upper and lower bound for $\nu(K)$. Those bounds are easily computable from D , depending only the Seifert graph $\Gamma(D)$:




Fig. 6. A diagram of the knot $11n_{53}$, and what happens to it in the proof of [Theorem 5](#). Diagrams drawn with [knotscape \[14\]](#).

Definition. The *Seifert graph* $\Gamma(D)$ of a link diagram D is a planar bipartite graph whose edges carry a sign (+ or −). It is constructed as follows: the vertices of $\Gamma(D)$ correspond to the circles of the Seifert resolution of D . A fixed crossing of D is adjacent to two different Seifert circles, which correspond to two vertices in $\Gamma(D)$. For any crossing, let $\Gamma(D)$ have an edge between these two vertices. The edge's sign indicates if the crossing is positive or negative. Let $\Gamma^+(D)$ ($\Gamma^-(D)$) be the subgraph of $\Gamma(D)$ that contains only the positive (negative) edges. Let $O^\pm(D)$ be the number of connected components of $\Gamma^\pm(D)$.

Theorem 5 (*The sharper slice-Bennequin inequality*). *Let D be a diagram of a knot K , with writhe w and n Seifert circles. Then*

$$-1 + w - n + 2O^+ \leq \nu(K) \leq 1 + w + n - 2O^-.$$

Proof (following [1]). Let us only prove the lower bound, since the upper bound then follows from $\nu(-K) = -\nu(K)$. For an example of the following constructions, see [Fig. 6](#). Let $G(D)$ be the graph obtained from $\Gamma(D)$ by contracting all positive edges; or more explicitly, the graph that has as vertices the components of $\Gamma^+(D)$, and has for each negative edge in $\Gamma(D)$ an edge between the corresponding vertices of $\Gamma^+(D)$. Then $G(D)$ is a connected graph with O^+ many vertices and k^- many edges. Pick $O^+ - 1$ edges that form a tree T . Gluing together $k^- - O^+ + 1$ many copies of the cobordism of [Lemma 5.2](#)

gives a connected cobordism with Euler characteristic $-k^- + O^+ - 1$ between K and a link with diagram D' , such that $G(D')$ is the tree T . The $O^+ - 1$ many negative crossings of D' are thus nugatory and may be removed by twists; each twist diminishes the number of Seifert circles by one. The ensuing diagram D'' is positive, with k^+ many crossings and $n - O^+ + 1$ many Seifert circles. If the link represented by D'' has c components, connect them by adding $c - 1$ positive crossings using the cobordism of [Lemma 5.2](#). This gives a cobordism to a knot K' with diagram D''' of Euler characteristic $1 - c$. Overall, there is a cobordism between K and K' of Euler characteristic $-c - k^- + O^+$. The diagram D''' is positive with $k^+ + c - 1$ many crossings and $n - O^+ + 1$ many Seifert circles, hence $\nu(K') = k^+ + c - n + O^+ - 1$. Finally, by [Proposition 5.1\(ii\)](#),

$$\begin{aligned} \nu(K') - \nu(K) &\leq c + k^- - O^+ \\ \implies \nu(K) &\geq (k^+ + c - n + O^+ - 1) - c - k^- + O^+ \\ \implies \nu(K) &\geq 1 + w - n + 2O^+. \quad \square \end{aligned}$$

Corollary 5.9. *On an alternating knot, slice-torus invariants take the same value as the signature.*

Proof. Follows from [\[22, Proposition 3.3\]](#). $\square$

If D is a knot diagram with $O^+ + O^- = n + 1$, then the lower bound of [Theorem 5](#) equals the upper bound, and thus ν is determined by the inequalities. Such diagrams are called *homogeneous*, and, consequently, a link is called homogeneous if it has a homogeneous diagram. This notion was introduced by Cromwell [\[5\]](#) and its relationship with the Rasmussen invariant was studied by Abe [\[1\]](#).

For the sake of completeness, let us cite results stated in [\[30\]](#) and [\[54\]](#) about $2\mathbb{Z}$ -valued slice-torus invariants of certain satellite knots. The results and their proofs (which are therefore omitted) carry through mostly unchanged to real-valued slice-torus invariants. For a knot K , let $D_{\pm}(K, t)$ be the t -twisted positive or negative Whitehead double. Notice that $g_4(D_{\pm}(K, t)) \leq 1$, and thus $\nu(D_{\pm}(K, t)) \in [-2, 2]$. Let $TB(K)$ be the Thurston–Bennequin number. Then:

Proposition 5.10. (See [\[30\]](#).)

- (i) $|\nu(D_-(K, t))| + |\nu(D_+(K, t))| \leq 2$. In particular, $\nu(D_+(K, t)) = \pm 2 \implies \nu(D_-(K, t)) = 0$.
- (ii) Let $N : \mathbb{Z} \rightarrow \mathbb{R}$ be given by $t \mapsto \nu(D_+(K, t))$. Then N is non-increasing; for $t \leq TB(K)$, we have $N(t) = 2$, and for $t \geq -TB(-K)$, we have $N(t) = 0$.

For two coprime integers m, n , let $K_{m,n}$ be the (m, n) -cable of K , i.e. the satellite with companion K and pattern the (m, n) -torus knot.

Proposition 5.11. (See [54].) *Let us fix a knot K and some $m > 0$, and define $h : \mathbb{Z} \rightarrow \mathbb{R}$ by $h(n) = \nu(K_{m,n}) - (m-1) \cdot n$. Then h is non-increasing and bounded, and $\sup h - \inf h \leq 2(m-1)$.*

6. Linear independence of some of the $\mathfrak{sl}_N$ -concordance invariants

This section contains the proofs of [Theorem 4](#) and [Theorem 1](#). Pretzel knots are a practical family of candidates to disprove the conjecture that all the $\mathfrak{sl}_N$ -concordance invariants are equal: they show sufficiently complex behaviour, yet their diagrams allow easy calculations, because they invite an inductive approach.

Remark 6.1. In the proof of [Theorem 4](#), we will only use [Theorem 2](#) and not the potentially stronger [Proposition 4.2](#). But even [Proposition 4.2](#) would not be strong enough to completely determine the value of $s_N(P(\ell, -m, 2))$. For example, using Webster's programme [55] or the skein long exact sequence, one finds that

$$\begin{aligned} \overline{\llbracket P(5, -3, 2) \rrbracket}_\infty &= t^{-3}a^2q^4 + t^{-2}q^6 + t^{-1}a^2 + (2q^2 + 1) \\ &\quad + t(a^{-2}q^4 + a^2q^{-4}) + 2t^2q^{-2} + t^3a^{-2} + t^4q^{-6} + t^5a^{-2}q^{-4}. \end{aligned}$$

But this polynomial has several different decompositions as in [Proposition 4.3](#), among them one with $\alpha = \beta = 0$, and one with $\alpha = 2, \beta = 0$.

Let us start by verifying the stated values for the Rasmussen invariant s_2 .

Proof of Theorem 4(i) & (iii). Khovanov homology of three stranded pretzels has been completely computed, see [37]. Since pretzel knots have homological width 3, their Khovanov homology determines their Rasmussen invariant. But the essential case $\ell > m > n \geq 2$ has a quicker proof: in that case, the $(\ell, -m, n)$ -pretzel knot is quasi-alternating (see Champanerkar and Kofman [4] and Greene [10]), and hence its s_2 -invariant (and twice its τ -invariant) equals its signature (see [Remark 2.11](#)):

$$s_2(P(\ell, -m, n)) = 2\tau(P(\ell, -m, n)) = \sigma(P(\ell, -m, n)) = \ell - m.$$

This value of the signature can be easily computed using Göttritz matrices and the formula of Gordon and Litherland [7]. $\square$

Let us continue by calculating the higher $\mathfrak{sl}_N$ -concordance invariants.

Lemma 6.2. *For odd $\ell \geq 3$, we have*

$$\overline{\llbracket T(\ell, 2) \rrbracket}_\infty = a^{1-\ell}q^{\ell-1} \cdot \left(1 + (t^2q^{-4} + t^3a^{-2}q^{-2}) \cdot \frac{t^{\ell-1}q^{2-2\ell} - 1}{t^2q^{-4} - 1} \right).$$

Proof. First, one may inductively calculate the HOMFLYPT-polynomial of $T(\ell, 2)$, using its defining skein relation. Then, since the $(\ell, 2)$ -torus knot is two-bridge, [Proposition 2.10](#) gives the HOMFLYPT-homology. $\square$

Lemma 6.3. *For all $N \geq 2$ and odd $\ell \geq 5$,*

$$s_N(P(\ell, 2 - \ell, 2)) \in \left\{ 0, \frac{2}{N-1} \right\}$$

$$s_N(P(\ell, 2 - \ell, 4)) \in \begin{cases} \{0, 2\} & N = 2, \\ [\frac{4}{N-1} - 2, 0] & N \geq 3. \end{cases}$$

Proof. Let $K_- = P(\ell, 2 - \ell, 2)$. Switching one of the two negative crossings of the last pretzel strand, one obtains the sum of two torus knots: $K_+ = T(\ell, 2) \# T(2 - \ell, 2)$. Resolving that crossing, one obtains the positive Hopf link (to get its standard diagram, apply $(\ell - 2)$ Reidemeister II moves). The homology of those torus knots has been computed in [Lemma 6.2](#), and the homology is well-behaved with respect to the connected sum (see [Proposition 2.8](#)). This gives us $\text{xdim } \overline{[K_+]_\infty}$. The totally reduced homology of the Hopf link is known, too, see [Section 2](#). So using the skein long exact sequence (see [Corollary 2.14](#)), one finds that

$$\text{xdim } \overline{[P(\ell, 2 - \ell, 2)]_\infty}^0 \leq (\ell - 2)q^{-2} + 1.$$

By virtue of [Theorem 3](#), this proves the first statement of the lemma. Notice also that

$$\text{xdim } \overline{[P(\ell, 2 - \ell, 2)]_\infty}^2 \leq (\ell - 4)q^{-6} + a^{-2}.$$

Now let $K_- = P(\ell, 2 - \ell, 4)$, and fix one of the negative crossings of the last pretzel strand. Then $K_+ = P(\ell, 2 - \ell, 2)$, and once again L_0 is the positive Hopf link. So

$$\text{xdim } \overline{[P(\ell, 2 - \ell, 4)]_\infty}^0 \leq (\ell - 4)a^2q^{-6} + 2.$$

Applying [Theorem 3](#) concludes the proof of the second statement. $\square$

Lemma 6.4. *Let ν be any slice-torus invariant, ℓ and m be odd positive integers, and n an even positive integer. Then*

$$\nu(P(\ell, -m, n)) \in [\ell - m - 2, \ell - m].$$

Proof. The standard diagram of the $(\ell, -m, n)$ -pretzel knot, as shown exemplarily in [Fig. 1](#), has writhe $(\ell - m - n)$, $(n + 1)$ many Seifert circles, $O^+ = n$ and $O^- = 1$. So the statement follows from the sharper slice Bennequin inequality ([Theorem 5](#)). $\square$

Let us now assemble the proof:

Proof of Theorem 4(ii) & (iv). Let $\ell > m \geq 3$ be odd and $n \geq 4$ even, and let $N \geq 3$. By Lemma 6.3, we have

$$s_N(P(m+2, -m, 4)) \in \left[\frac{4}{N-1} - 2, 0 \right].$$

It takes $\frac{n-4}{2}$ many crossing switches from positive to negative, and $\frac{\ell-m-2}{2}$ many crossing switches from negative to positive to go from $P(m+2, -m, 4)$ to $P(\ell, -m, n)$. Thus by Proposition 5.3 we have

$$s_N(P(\ell, -m, n)) \in \left[\frac{4}{N-1} + 2 - n, \ell - m - 2 \right].$$

But by Lemma 6.4, $s_N(P(\ell, -m, n)) \in [\ell - m - 2, \ell - m]$. This leaves $s_N(P(\ell, -m, n)) = \ell - m - 2$ as only value in the intersection of the two intervals.

Let us now consider the special case $n = 2$. By the same method one finds that

$$s_N(P(\ell, -m, 2)) \in [\ell - m - 2, \ell - m - 2 + 2/(N-1)].$$

The intersection of this interval with $\frac{2}{N-1}\mathbb{Z}$ leaves $\ell - m - 2$ and $\ell - m - 2 + 2/(N-1)$ as only possible values. $\square$

The linear independence of τ , s_2 and s_3 from the s_N with $N \geq 4$ now follows quickly.

Proof of Theorem 1. (i): To prove the first statement, note that any linear combination of $\{s_N\}_{N \geq 3}$ vanishes for $P(7, -5, 4)$, but s_2 and τ do not. Concerning the second statement, invariance of s_2 and τ is due to [12], so it suffices to produce a knot K with $s_2(K) = \tau(K) = 0$, but $s_N(K) \neq 0$. This is accomplished by the quasi-alternating knot $K = P(5, -3, 2) \# -T(3, 2)$.

(ii): Consider a linear combination u of $\{s_N\}_{N \geq 4}$. For u to be equal to s_3 , the linear combination needs to be convex (i.e. the sum of coefficients is equal to 1). Therefore, $u(P(5, -3, 2)) \in [0, 2/3]$; but $s_3(P(5, -3, 2)) = 1$. Next, the linear independence of $\{\tau, s_2, s_3\}$ has been proven in (i). So it is enough to show the existence of a knot K with $\tau(K) = s_2(K) = s_3(K) = 0$, but $s_N(K) \neq 0$. For this purpose, take

$$K = P(5, -3, 2) \# P(5, -3, 2) \# -P(7, -5, 4) \# -T(3, 2).$$

We have $s_2(K) = \tau(K) = 0$, and using FoamHo [25], $s_3(P(5, -3, 2)) = 1 \implies s_3(K) = 0$. On the other hand, $s_N(P(5, -3, 2)) \leq 2/(N-1) \implies s_N(K) \leq 4/(N-1) - 2 \neq 0$ since $N \geq 4$. $\square$

Let us compute another example, to illustrate that the Rasmussen invariant does not necessarily give the best slice genus bound among the $\mathfrak{sl}_N$ -concordance invariants.




Fig. 7. From left to right: $12n_{340}$, 10_{141} and 8_9 (drawn with Knotscape [14]).

Example 6.5. Let $K = 12n_{340}$, then $s_2(K) = 0$, $s_3(K) = 1$ and for $N \geq 4$: $s_N(K) \in \{2 - 2/(N - 1), 2\}$.

Proof. The value of s_2 and s_3 may be computed using JavaKh [9] and FoamHo [25], respectively; the other values can be read from $\overline{\llbracket K \rrbracket}_\infty$, which we are going to compute using the skein long exact sequence. Notice that the calculation is rather quick, and that we do not need to determine the HOMFLYPT-homology of K completely (this would be possible though, using Rasmussen's spectral sequences Proposition 2.7).

Resolving the crossing indicated in Fig. 7 gives K as K_+ , 10_{141} as K_- and the positive Hopf link as L_0 . Resolving once more the indicated crossing of 10_{141} gives 10_{141} as K_+ , 8_9 as K_- and the positive Hopf link as L_0 . The knot 8_9 is two-bridge, so its reduced HOMFLYPT-homology is determined by its HOMFLYPT-polynomial and its signature. One finds

$$\text{xdim } \overline{\llbracket 8_9 \rrbracket}_\infty^{-4} = q^4 a^2.$$

Applying Corollary 2.14 twice gives

$$\text{xdim } \overline{\llbracket K \rrbracket}_\infty \leq t^4 a^{-4} \text{xdim } \overline{\llbracket 8_9 \rrbracket}_\infty + (t^{1/2} a^{-1} + t^{5/2} a^{-3}) \text{xdim } \overline{\llbracket T(2, 2) \rrbracket}_\infty,$$

and therefore

$$\text{xdim } \overline{\llbracket K \rrbracket}_\infty^0 \leq q^4 a^{-2} + q^2 a^{-2}.$$

By Theorem 3 it follows that

$$\forall N \geq 2: \quad s_N(K) \in \{2 - 2/(N - 1), 2\}. \quad \square$$

Acknowledgments

This paper is extracted from my thesis [26], and I thank C. Blanchet for having been my adviser. Thanks to A. Lobb for comments on a first version of the paper.

References

- [1] T. Abe, The Rasmussen invariant of a homogeneous knot, *Proc. Amer. Math. Soc.* 139 (7) (2011) 2647–2656, <http://dx.doi.org/10.1090/S0002-9939-2010-10687-1>.
- [2] D. Bar-Natan, Khovanov’s homology for tangles and cobordisms, *Geom. Topol.* 9 (2005) 1443–1499, <http://dx.doi.org/10.2140/gt.2005.9.1443>.
- [3] D. Bennequin, Entrelacements et équations de Pfaff, *Astérisque* 107 (8) (1982) 87–161.
- [4] A. Champanerker, I. Kofman, Twisting quasi-alternating links, *Proc. Amer. Math. Soc.* 137 (7) (2009) 2451–2458, <http://dx.doi.org/10.1090/S0002-9939-09-09876-1>.
- [5] P.R. Cromwell, Homogeneous links, *J. Lond. Math. Soc.* (2) 39 (3) (1989) 535–552, <http://dx.doi.org/10.1112/jlms/s2-39.3.535>.
- [6] M.H. Freedman, R. Gompf, S. Morrison, K. Walker, Man and machine thinking about the smooth 4-dimensional Poincaré conjecture, *Quantum Topol.* 1 (2) (2010) 171–208, <http://dx.doi.org/10.4171/QT/5>.
- [7] C.M. Gordon, R.A. Litherland, On the signature of a link, *Invent. Math.* 47 (1) (1978) 53–69, <http://dx.doi.org/10.1007/BF01609479>.
- [8] B. Gornik, Note on Khovanov link cohomology, <http://arxiv.org/abs/math/0402266>, 2004.
- [9] J. Green, S. Morrison, JavaKh, computer program available at <http://katlas.math.toronto.edu/>, 2005.
- [10] J. Greene, Homologically thin, non-quasi-alternating links, *Math. Res. Lett.* 17 (1) (2010) 39–49.
- [11] M. Hedden, Notions of positivity and the Ozsváth–Szabó concordance invariant, *J. Knot Theory Ramifications* 19 (5) (2010) 617–629, <http://dx.doi.org/10.1142/S0218216510008017>.
- [12] M. Hedden, P. Ording, The Ozsváth–Szabó and Rasmussen concordance invariants are not equal, *Amer. J. Math.* 130 (2) (2008) 441–453, <http://dx.doi.org/10.1353/ajm.2008.0017>.
- [13] J. Hom, An infinite rank summand of topologically slice knots, <http://arxiv.org/abs/1310.4476>, 2013.
- [14] J. Hoste, M. Thistlethwaite, Knotscape version 1.01, computer program available at www.math.utk.edu/~morwen/knotscape.html, 1999.
- [15] S. Jabuka, Concordance invariants from higher order covers, *Topology Appl.* 159 (10–11) (2012) 2694–2710, <http://dx.doi.org/10.1016/j.topol.2012.03.014>.
- [16] T. Kawamura, The Rasmussen invariants and the sharper slice-Bennequin inequality on knots, *Topology* 46 (1) (2007) 29–38, <http://dx.doi.org/10.1016/j.top.2006.10.001>.
- [17] T. Kawamura, An estimate of the Rasmussen invariant for links.
- [18] M. Khovanov, A categorification of the Jones polynomial, *Duke Math. J.* 101 (3) (2000) 359–426, <http://dx.doi.org/10.1142/S0218216505003750>.
- [19] M. Khovanov, $sl(3)$ link homology, *Algebr. Geom. Topol.* 4 (1) (2004) 1045–1081, <http://dx.doi.org/10.1142/S0218216505003750>.
- [20] M. Khovanov, L. Rozansky, Matrix factorizations and link homology, *Fund. Math.* 199 (1) (2008) 1–91, <http://dx.doi.org/10.4064/fm199-1-1>.
- [21] M. Khovanov, L. Rozansky, Matrix factorizations and link homology II, *Geom. Topol.* 12 (3) (2008) 1387–1425, <http://dx.doi.org/10.2140/gt.2008.12.1387>.
- [22] E.S. Lee, The support of the Khovanov’s invariants for alternating knots, <http://arxiv.org/abs/math/0201105>, 2002.
- [23] E.S. Lee, An endomorphism of the Khovanov invariant, *Adv. Math.* 197 (2) (2005) 554–586, <http://dx.doi.org/10.1016/j.aim.2004.10.015>.
- [24] L. Lewark, sl_3 -foam homology calculations, *Algebr. Geom. Topol.* 13 (6) (2013) 3661–3686, <http://dx.doi.org/10.2140/agt.2013.13.3661> (electronic).
- [25] L. Lewark, *FoamHo*, computer program available at www.maths.dur.ac.uk/~vxhn54, 2013.
- [26] L. Lewark, Khovanov–Rozansky homologies, knotted weighted webs and the slice genus, PhD thesis, Institut de mathématiques de Jussieu (Université Paris VII), 2013, www.maths.dur.ac.uk/~vxhn54/.
- [27] R. Lipshitz, S. Sarkar, A refinement of Rasmussen’s s -invariant, <http://arxiv.org/abs/1206.3532>, 2012.
- [28] C. Livingston, Computations of the Ozsváth–Szabó knot concordance invariant, *Geom. Topol.* 8 (2004) 735–742, <http://dx.doi.org/10.2140/gt.2004.8.735> (electronic).
- [29] C. Livingston, Slice knots with distinct Ozsváth–Szabó and Rasmussen invariants, *Proc. Amer. Math. Soc.* 136 (1) (2008) 347–349, <http://dx.doi.org/10.1090/S0002-9939-07-09276-3> (electronic).
- [30] C. Livingston, S. Naik, Ozsváth–Szabó and Rasmussen invariants of doubled knots, *Algebr. Geom. Topol.* 6 (2006) 651–657, <http://dx.doi.org/10.2140/agt.2006.6.651> (electronic).

- [31] A. Lobb, A slice genus lower bound from $\mathrm{sl}(n)$ Khovanov–Rozansky homology, *Adv. Math.* 222 (4) (2009) 1220–1276, <http://dx.doi.org/10.1016/j.aim.2009.06.001>.
- [32] A. Lobb, Computable bounds for Rasmussen’s concordance invariant, *Compos. Math.* 147 (2) (2011) 661–668, <http://dx.doi.org/10.1112/S0010437X10005117>.
- [33] A. Lobb, A note on Gornik’s perturbation of Khovanov–Rozansky homology, *Algebr. Geom. Topol.* 12 (1) (2012) 293–305, <http://dx.doi.org/10.2140/agt.2012.12.293>.
- [34] M. Mackaay, P. Turner, P. Vaz, A remark on Rasmussen’s invariant of knots, *J. Knot Theory Ramifications* 16 (3) (2007) 333–344, <http://dx.doi.org/10.1142/S0218216507005312>.
- [35] M. Mackaay, P. Vaz, The foam and the matrix factorization sl_3 link homologies are equivalent, *Algebr. Geom. Topol.* 8 (1) (2008) 309–342, <http://dx.doi.org/10.2140/agt.2008.8.309>.
- [36] M. Mackaay, P. Vaz, The reduced HOMFLY-PT homology for the Conway and the Kinoshita–Terasaka knots, <http://arxiv.org/abs/0812.1957>, 2008.
- [37] A. Manion, The rational Khovanov homology of 3-strand pretzel links, <http://arxiv.org/abs/1110.2239>, 2011.
- [38] C. Manolescu, B. Owens, A concordance invariant from the Floer homology of double branched covers, *Int. Math. Res. Not. IMRN* 2007 (20) (2007), <http://dx.doi.org/10.1093/imrn/rnm077>, Art. ID rnm077, 21.
- [39] C. Manolescu, P. Ozsváth, On the Khovanov and knot Floer homologies of quasi-alternating links, in: *Proceedings of Gökova Geometry–Topology Conference 2007, Gökova Geometry/Topology Conference (GGT), Gökova, 2008*, pp. 60–81.
- [40] J. McCleary, *A User’s Guide to Spectral Sequences*, 2nd edition, *Cambridge Stud. Adv. Math.*, vol. 58, Cambridge University Press, Cambridge, 2001.
- [41] P. Ozsváth, Z. Szabó, Knot Floer homology and the four-ball genus, *Geom. Topol.* 7 (2003) 225–254, <http://dx.doi.org/10.2140/gt.2003.7.615>.
- [42] P. Ozsváth, Z. Szabó, On the Heegaard Floer homology of branched double-covers, *Adv. Math.* 194 (1) (2005) 1–33, <http://dx.doi.org/10.1016/j.aim.2004.05.008>.
- [43] O. Plamenevskaya, Transverse knots and Khovanov homology, *Math. Res. Lett.* 13 (4) (2006) 571–586, <http://dx.doi.org/10.1016/j.top.2006.10.001>.
- [44] J.A. Rasmussen, Floer homology and knot complements, PhD thesis, Harvard University, 2003, <http://arxiv.org/abs/math/0306378>.
- [45] J.A. Rasmussen, Some differentials on Khovanov–Rozansky homology, <http://arxiv.org/abs/math/0607544>, 2006.
- [46] J.A. Rasmussen, Khovanov–Rozansky homology of two-bridge knots and links, *Duke Math. J.* 136 (3) (2007) 551–583, <http://dx.doi.org/10.1215/S0012-7094-07-13635-4>.
- [47] J.A. Rasmussen, Khovanov homology and the slice genus, *Invent. Math.* 182 (2) (2010) 419–447, <http://dx.doi.org/10.1007/s00222-010-0275-6>.
- [48] N.Y. Reshetikhin, V.G. Turaev, Ribbon graphs and their invariants derived from quantum groups, *Comm. Math. Phys.* 127 (1) (1990) 1–26.
- [49] L. Rudolph, Quasipositivity as an obstruction to sliceness, *Bull. Amer. Math. Soc. (N.S.)* 29 (1) (1993) 51–59, <http://dx.doi.org/10.1090/S0273-0979-1993-00397-5>.
- [50] L. Rudolph, Positive links are strongly quasipositive, in: *Proceedings of the Kirbyfest, Berkeley, CA, 1998*, in: *Geom. Topol. Monogr.*, vol. 2, Geom. Topol. Publ., Coventry, 1999, pp. 555–562 (electronic).
- [51] C. Seed, knotkit, computer program available at <https://github.com/cseed/knotkit>, 2013.
- [52] A. Shumakovitch, Rasmussen invariant, slice-Bennequin inequality, and sliceness of knots, *J. Knot Theory Ramifications* 16 (10) (2007) 1403–1412, <http://dx.doi.org/10.1142/S0218216507005889>.
- [53] P.R. Turner, Calculating Bar-Natan’s characteristic two Khovanov homology, *J. Knot Theory Ramifications* 15 (10) (2006) 1335–1356, <http://dx.doi.org/10.1142/S0218216506005111>.
- [54] C.A. Van Cott, Ozsváth–Szabó and Rasmussen invariants of cable knots, *Algebr. Geom. Topol.* 10 (2) (2010) 825–836, <http://dx.doi.org/10.2140/agt.2010.10.825>.
- [55] B. Webster, Kr.m2, computer program available from katlas.math.toronto.edu/wiki/User:Ben/KRhomology, 2005.
- [56] H. Wu, The Khovanov–Rozansky cohomology and Bennequin inequalities, <http://arxiv.org/abs/math/0703210>, 2007.
- [57] H. Wu, On the quantum filtration of the Khovanov–Rozansky cohomology, *Adv. Math.* 221 (1) (2009) 54–139, <http://dx.doi.org/10.1016/j.aim.2008.12.003>.