

Manuscript Number: JAES-D-10-00328R1

Title: Reconciling the Intertropical Convergence Zone, Himalayan/Tibetan tectonics, and the onset of the Asian monsoon system

Article Type: Special Issue Asian Climate & Tectonics

Keywords: Intertropical Convergence Zone; Himalayas; Cenozoic; climate; monsoon

Corresponding Author: Dr. Mark Allen,

Corresponding Author's Institution:

First Author: Mark Allen

Order of Authors: Mark Allen; Howard A Armstrong

Abstract: Numerous climate proxies from across Central, South and East Asia have yielded different ages for the start or intensification of a monsoon climate system. Common estimates include the early part of the early Miocene (~23-20 Ma) and the late Miocene (~11-8 Ma). In the early Miocene the average position of the Intertropical Convergence Zone (ITCZ) was likely to be the >2000 km closer to the Himalaya than at present (based on published data for the palaeolatitude of the Central Pacific ITCZ), such that Himalayan climate was marked by high precipitation, but not necessarily seasonality. Here we propose that increased seasonality in the late Miocene in the Himalaya and neighbouring regions was a response to an increase in the distance between the ITCZ and the Himalaya/Tibet, such that the ITCZ was only brought northwards during the northern hemisphere summer each year. This is essentially the pattern of the modern South Asian monsoon system. These climatic changes coincide with a switch from north-south extensional shear on the northern side of the High Himalaya, thrusting on the Main Central Thrust and rapid metamorphic exhumation, to thrusts further south in the Himalaya (Main Boundary Thrust) and a reduction in High Himalayan exhumation rates. We speculate that the tectonic changes were at least in part a response to a reduction in precipitation over the High Himalaya: the Himalayan thrust belt re-organised to maintain a critical taper appropriate to a drier orogen. The reduction in metamorphic exhumation after the early Miocene would also have led to a reduction in the flux of metamorphic CO₂ to the atmosphere, thereby promoting the global shift to a cooler climate in the mid Miocene.

1 **Reconciling the Intertropical Convergence Zone, Himalayan/Tibetan tectonics,**
2 **and the onset of the Asian monsoon system**

3

4 Mark B. Allen* and Howard A. Armstrong

5

6 Department of Earth Sciences, University of Durham, Durham, DH1 3LE, UK

7 *m.b.allen@durham.ac.uk

8

9 ABSTRACT

10 Numerous climate proxies from across Central, South and East Asia have yielded
11 different ages for the start or intensification of a monsoon climate system. Common
12 estimates include the early part of the early Miocene (~23-20 Ma) and the late
13 Miocene (~11-8 Ma). In the early Miocene the average position of the Intertropical
14 Convergence Zone (ITCZ) was likely to be the >2000 km closer to the Himalaya than
15 at present (based on published data for the palaeolatitude of the Central Pacific ITCZ),
16 such that Himalayan climate was marked by high precipitation, but not necessarily
17 seasonality. Here we propose that increased seasonality in the late Miocene in the
18 Himalaya and neighbouring regions was a response to an increase in the distance
19 between the ITCZ and the Himalaya/Tibet, such that the ITCZ was only brought
20 northwards during the northern hemisphere summer each year. This is essentially the
21 pattern of the modern South Asian monsoon system. These climatic changes coincide
22 with a switch from north-south extensional shear on the northern side of the High
23 Himalaya, thrusting on the Main Central Thrust and rapid metamorphic exhumation,
24 to thrusts further south in the Himalaya (Main Boundary Thrust) and a reduction in
25 High Himalayan exhumation rates. We speculate that the tectonic changes were at

least in part a response to a reduction in precipitation over the High Himalaya: the Himalayan thrust belt re-organised to maintain a critical taper appropriate to a drier orogen. The reduction in metamorphic exhumation after the early Miocene would also have led to a reduction in the flux of metamorphic CO₂ to the atmosphere, thereby promoting the global shift to a cooler climate in the mid Miocene.

Keywords: Intertropical Convergence Zone, Himalaya, Cenozoic, climate, monsoon.

1. Introduction

In a recent paper (Armstrong and Allen, 2011) we argued that the changing palaeolatitudes of the Intertropical Convergence Zone (ITCZ) and the Himalaya had an important control on the Asian climate during the late Cenozoic. The specific argument was that the latitudes of the ITCZ and the Himalaya were >2000 km closer in the early Miocene (~20 Ma) than they are today, such that the range would have received precipitation on a scale equivalent to the modern South Asian summer monsoon (roughly June-September) all through the year. This hypothesis is based on reconstructions of the Pacific ITCZ (Lyle et al., 2002) and the northward drift of the Indian plate (Molnar and Stock, 2009). The consequences of this proximity in the early Miocene would have been high erosion rates, consistent with the exhumation data for the range for this time (White et al., 2002), and possibly even the development of “channel flow” tectonics (Beaumont et al., 2001; Searle, 2010) and the generation of crustal melt leucogranites (Harris, 2007).

In this review we develop this hypothesis to address 1) why different climate proxies have produced different ages for the onset of the Asian monsoon system, and

2) in particular, why many results from both north and south of the Himalaya/Tibet indicate a late Miocene onset or intensification age. Our conclusion is that in the late Miocene the increasing divergence of the ITCZ and the Himalaya enhanced seasonality within the range and to its south, whilst overall precipitation declined.

2. Miocene Himalayan/Tibetan tectonics and palaeoaltitudes

As many authors have produced reviews of Himalayan/Tibetan tectonics (e.g. Harrison et al., 1992; Yin and Harrison, 2000; Hodges, 2000; Yin, 2006; Yin, 2010) we do not provide a detailed account, but summarise Miocene events. The Miocene epoch began long after the commonly-assumed initial collision at ~50 Ma (e.g. Dupont-Nivet et al., 2010), such that it is a long-standing issue how and where plate convergence was accommodated for the first 30 million years of collision (the “Eohimalayan” phase of tectonics; Hodges, 2000). Increasing evidence is emerging for deformation and consequent uplift and exhumation through this time (Najman et al. 2008; Aikman et al. 2008), but it is still scant.

The early Miocene interval was a time of dramatic change, with the initiation of the South Tibetan Detachment System (STDS) and the Main Central Thrust (MCT) within the Himalaya (Burchfiel et al., 1992; Hodges et al., 1998; Searle et al., 2008; Kellett et al. 2009). Rapid exhumation of rock in the intervening wedge in the High Himalaya was associated with the generation of leucogranite via decompression (Harris and Massey, 1994; White et al., 2002). The most rapid exhumation, leucogranite generation and ductile shear all took place in the early Miocene and diminished in the middle-late Miocene (see Catlos et al., 2004), with some evidence for diachroneity from west to east along the range (Harris, 2007). For example, Searle

et al. (1999) noted that in the Zaskar region of the High Himalaya (Fig. 1), exhumation rates dropped from 6-10 mm/yr between 21 and 18.5 Ma to 0.4 mm/yr afterwards. The distinctive early Miocene tectonics of the Himalaya has led to the development of channel flow models for the mid and lower crust (Beaumont et al., 2001), although some authors see the synchronous extensional and thrust-sense shear as the readjustment of a critically-tapered orogenic wedge (Robinson et al., 2006; Kali et al., 2010).

At roughly the time activity on the MCT and STDS declined in the middle Miocene, normal faulting started in southern Tibet, indicating rough east-west extension rather than the top-to-north extensional shear on the STDS (Molnar and Tapponnier, 1978). The age constraints on this younger normal faulting are patchy, such that it is not possible to get a clear picture of it through space and time, or its relations to the STDS. The oldest ages found so far for faulting are 14 Ma (Coleman and Hodges, 1995) and 13.5 Ma (Blisniuk et al., 2001), i.e. middle Miocene. Faulting is active (Molnar and Tapponnier, 1978; Molnar and Lyon-Caen, 1989). North-south dykes in southern Tibet, dated at 18.3 ± 2.7 Ma, have been taken to mean east-west extension at this time (Williams et al., 2001; Fig. 1), but this does not necessarily mean contemporary normal faulting: σ_3 could be oriented east-west, in a constrictional stress field. The north-south Dinggye normal fault cross-cuts the STDS, and began its motion >11 Ma (Leloup et al., 2010). See Maheo et al. (2007) for a contrary view, suggesting that normal faulting in southern Tibet is mainly a Pliocene-Quaternary phenomenon.

100 One explanation for the onset of the east-west extension is partial loss of the
 101 lower continental lithosphere, with resultant surface uplift following isostatic
 102 readjustment (Houseman et al., 1981; England and Houseman, 1989; Molnar et al.,
 103 1993). However, this is difficult to reconcile with palaeoaltitude reconstructions that
 104 place southern and central Tibet at close to their present elevations by 15 Ma or
 105 earlier. Most of these studies are based on carbonate $\delta^{18}\text{O}$ values. The pattern is
 106 emerging that southern and central Tibet has been at or close to present day elevations
 107 through the Neogene at least (Garzione et al., 2000; Rowley et al., 2001; Rowley and
 108 Currie, 2006; DeCelles et al., 2007; Fig. 1), and possibly once higher than at present
 109 (Murphy et al., 2009; Saylor et al., 2009). Another approach was taken by Spicer et al.
 110 (2003), who used fossil leaf analysis to infer that southern Tibet has been close to its
 111 present elevation since 15 Ma. Lithosphere loss is not confirmed by recent estimates
 112 of plate thickness, based on earthquake shear wave velocity gradients (Priestley and
 113 McKenzie, 2006), which show a >200 km thick “core” under Tibet. Therefore an
 114 alternative to lithosphere detachment/delamination models may be necessary. Cook
 115 and Royden (2008) modelled extension in southern Tibet as the result of eastward
 116 crustal flow towards the southeast plateau corner, without requiring significant change
 117 in plateau surface elevation in southern Tibet. Evidence for middle Miocene surface
 118 uplift (~ 15-10 Ma) has been found in southeast Tibet and the Longmenshan (Clark et
 119 al., 2005; Ouimet et al., 2010), based on fission track and (U-Th)/He dates from
 120 fluvial gorges (Fig. 1). Fluvial incision further north along the east side of Tibet
 121 appears to have begun later, at ~6 Ma (Kirby et al., 2002). This is consistent with the
 122 model of Cook and Royden (2008), without being conclusive proof.
 123

Middle and late Miocene Himalayan thrusting focused to the south of the MCT, in particular the Main Boundary Thrust and the Main Frontal Thrust (Schelling, 1992; Meigs et al., 1995; Lave and Avouac, 2000; Huyghe et al., 2001; Herman et al., 2010). Present day surface deformation in the Himalaya is focussed along the front of the range. The structure of the Himalaya is also marked by pronounced syntaxes at its western and eastern ends, named after the mountains of Nanga Parbat and Namche Barwa respectively. Both of these regions show extremely high rates of exhumation during the Pliocene-Pleistocene (Zeitler et al., 1993; Seward et al., 2008), but both show evidence for an origin as far back as 10 Ma (Treloar et al., 2000; Booth et al., 2009).

3. Monsoon climate records

The modern Asian monsoon climate is marked by two main characteristics: seasonally reversing winds and heavy summer rainfall. The South and East Asian monsoons have different timings, characteristics and dynamics (Molnar et al., 2010). The South Asian monsoon affects the region south of the Himalaya and also Indochina and the South China Sea. Peak rainfall is from June-September. The East Asian monsoon affects China and adjacent countries to its east. Winters are not generally as dry as regions affected by the South Asian monsoon, and heavy summer rainfall arrives somewhat earlier. Tibet and the Himalaya may drive both forms of the monsoon, but in different ways (Boos and Kuang, 2010; Molnar et al., 2010). The high topography may act as a heat source, or a barrier to southward air flow, in either mechanism creating the South Asian monsoon by bringing a northward flow of tropical air in the northern hemisphere summer. The East Asian monsoon may result

148 more from the Tibetan plateau blocking the subtropical jet stream, which passes south
149 of Tibet in winter and north in summer.

150

151 The modern Asian summer monsoon involves the northward movement of the
152 Intertropical Convergence Zone over the Indian subcontinent and neighbouring areas
153 of South Asia (Fig. 2), partly because it tracks the solar zenith through the year, but
154 mainly because it is drawn northwards by a zone of low atmospheric pressure over
155 northern India, the Himalaya and Tibet (Ruddiman and Kutzbach, 1989). Regardless
156 of whether the dominant mechanism is heating of the overlying atmosphere or
157 blocking the southward flow of cool air (Boos and Kuang, 2010), the Himalaya and
158 Tibet are primary influences on atmospheric circulation patterns and hence climate.
159 For this reason the surface uplift history of the Himalayan-Tibetan orogen has been
160 suggested to be closely linked to the development of the Asian monsoon (Molnar et
161 al., 1993; Clift et al., 2008), with feedbacks operating in both directions. A higher,
162 wider Tibetan plateau would have intensified the monsoon, while a wetter climate
163 would have increased erosion rates in the Himalaya, in turn influencing relief and
164 exhumation patterns (e.g. Kutzbach et al., 1989). Late Cenozoic surface uplift and
165 erosion of the Himalaya and Tibet have been linked to global climate change, via high
166 chemical weathering rates and organic carbon burial, causing a drawdown of
167 atmospheric CO₂, and hence global cooling (Raymo et al. 1988; Raymo and
168 Ruddiman, 1992). Increasing seawater strontium isotope values in the late Cenozoic
169 are interpreted as reflecting increasing erosion of the Himalayan-Tibetan system (Fig.
170 3a; Richter et al., 1992).

171

No consensus exists for the timing of initial monsoon development or intensification. Proposed ages cluster in the early Miocene (~22-20 Ma; e.g. Guo et al., 2002), middle Miocene (16-11 Ma; e.g. Clift et al., 2004) and late Miocene (11-8 Ma; e.g. Molnar et al., 1993), although middle Miocene ages (Clift et al., 2004) may be viewed as the peak of conditions which originated in the early Miocene (Clift et al., 2008). The locations of sites that have provided data, and inferred ages, are shown in Fig. 1. Table 1 summarises these observations. We do not refer to the numerous records of climate change in the Pliocene-Quaternary, although these have been argued as evidence for rapid tectonic change in the Himalaya and/or Tibet. Refer to Molnar (2005) and Molnar et al. (2010), who succinctly showed the flaws in the idea.

3.1. Early Miocene ages

Many of the studies relevant to the timing of the onset of the Asian monsoon system are derived from data collected at one or several localities, with necessary extrapolation to regional scales. Sun and Wang (2005) presented a different approach, by collating “palaeophytogeographical patterns” across the whole of China. Their results show a shift from an east-west arid belt across China in the Paleogene to a Neogene pattern of aridity restricted to northwest China, similar to the present day. The transition occurred close to the Oligocene-Miocene boundary, but is not precisely datable because of the broad resolution of the study.

Early Miocene ages are also drawn from loess deposits close to the Liupan Shan in northern China (Fig. 1), which record alternations between the loess itself and paleosols as far back as 22 Ma (Guo et al., 2002). The loess is interpreted to result from dust derived from northerly winds during the winter monsoon (although at the

present day much more dust is generated in the Spring: Roe, 2009), while the paleosols represent wetter intervals influenced by the summer monsoon. Therefore the loess record provides good evidence for seasonality, but not necessarily for enhanced humidity. A record of humid early Miocene climate is provided by pollen data from fluviolacustrine sediments from the same region (Jiang and Ding, 2009), dating from 20 to 0.08 Ma. The assemblages are generally dominated by steppe flora throughout the Neogene and Quaternary, but variations between the proportions of *Artemisia*, more halophytic taxa such as *Humulus*, and deciduous broad-leaved trees such as *Juglans*, are interpreted by Jiang and Ding (2009) to mean a relatively humid climate from 20.13 to 14.25 Ma (i.e. early Miocene to early middle Miocene), influenced by the East Asian summer monsoon. This was followed by a significant decline in the summer monsoon and a more arid climate from 14.25 to 11.35 Ma. There has been a relatively arid climate and weak summer monsoon ever since. Further west, but still in the north Tibetan region (Qaidam Basin; Fig. 1), a shift to wetter conditions in the early Miocene was inferred by Wang et al. (1999) on the basis of pollen assemblages and a decrease in the proportion of xerophytes near the Oligocene-Miocene boundary.

The onset of aeolian deposition within the interior of the Junggar Basin began at ~24 Ma, i.e. just before the Oligocene-Miocene boundary (Sun et al., 2010; Fig. 1), suggesting that the modern system of westerly winds in this area began at this time. Dust was supplied to the Junggar Basin from Central Asia, in contrast to the northerly winds that supply the Chinese loess plateau.

Sedimentary $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ data from the northern side of the Tibetan plateau show a Neogene increase in both $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ values (Kent-Corson et al., 2009),

222 interpreted as a climatic shift to more arid conditions and basin isolation. The trends
223 vary between localities, such that some places show no clear pattern through the
224 Noogene and others have abrupt shifts that might indicate climatic change in the
225 middle or late Miocene. The $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$ records of Rieser et al. (2009) in the
226 Qaidam Basin show no clear overall trends.

227
228 Chemical and mineralogical proxies have been used to infer enhanced
229 monsoon climate, on the basis that runoff is a principal control of chemical
230 weathering rates. Wei et al. (2006) and Clift et al. (2008) applied this approach to
231 sediments in the South China Sea (ODP site 1148; Fig. 1), derived from the Pearl
232 River drainage basin. This river system was affected by Asian monsoon climate, but
233 not by high elevation source areas that would also carry a record of enhanced physical
234 erosion related to surface uplift of the Himalaya and Tibet. Clift et al. (2008) showed
235 a trend of gradually increasing chemical weathering (and hence inferred precipitation)
236 from 24 to 10 Ma. Chemical weathering rates declined between 10 and 3.5 Ma, and
237 rose thereafter. Indices from the Indus and Bengal fans (Fig. 1) show similar patterns,
238 with the caveat that the physical erosion record of the Himalaya must be present in
239 both fans.

240
241 The combination of high precipitation and mountainous relief should produce
242 high erosion rates, ultimately recorded as high sedimentation rates in basins at the end
243 of sediment routing systems. Assuming greater Asian monsoon intensity to be
244 expressed by higher precipitation rates, this should feed through to greater
245 accumulation rates in basins with the accommodation space to receive the sediment.
246 Clift and Gaedicke (2002), Clift (2006) and Clift et al. (2008) found high sediment

flux rates for the Indus fan (Fig. 1), which receives detritus from the Himalayan/Tibetan orogen, in the early and middle Miocene, followed by a decrease in the late Miocene. Bengal fan records are less complete, but chemical weathering indices show a relatively wet climate in the sediment source area for ODP Site 718 in the late early and middle Miocene, with a shift to inferred drier climate at 8-7 Ma (Clift et al., 2008). These results were interpreted by Clift et al. (2008) to mean an Asian monsoon climate in operation since the early Miocene. From a similar approach Clift et al. (2004) found peak sedimentation rates offshore of the Mekong drainage system beginning in the early Miocene, somewhat earlier than many basins on the East Asian margin (Fig. 1).

In summary, there are now many indicators of South and East Asian climate change in the early Miocene, consistent with the onset or intensification of climatic conditions akin to the modern range of climates (Table 1; Fig. 3b). A note of caution is needed when interpreting these changes in terms of monsoon intensification: the early Miocene was generally a warm, and presumably humid, time on a global scale (e.g. Zachos et al., 2001), such that increased humidity in South or East Asia may not simply be a regional climatic change.

3.2. Middle Miocene ages

There are studies that claim climatic shifts in the middle Miocene (16-11 Ma), largely derived from high sedimentation rates in East Asian basins (e.g. Clift et al., 2004), but the emphasis is now towards a monsoonal climate at this time being a continuation from early Miocene times (Clift et al., 2008), with high global sediment fluxes being influenced by climatic instability, specifically the shift from the early

middle Miocene Climatic Optimum to colder conditions (Zachos et al., 2001; Clift, 2010; Fig. 3a).

High middle Miocene sedimentation rates in the Indus fan and Arabian Sea were interpreted by Clift and Gaedicke (2002) and Clift et al. (2008) as indicating monsoonal conditions in the sediment source areas of the Himalaya and Tibet, continuing or even intensifying from an early Miocene onset. Similar enhanced sedimentation rates were found by Clift et al. (2004) from basins along the East Asian continental margin, and Burma (Fig. 1). However, other authors have inferred increased aridity in the Himalaya and Tibet from the middle Miocene. For example, Wei et al (2006) and Wan et al (2007) noted decreases in the degree of chemical weathering of sediment delivered to the South China Sea, that they attributed to increased aridity in Asian sources areas at about 16-15 Ma. Further increases in aridity were inferred for about 8 Ma and 3 Ma. Neogene pollen records from across China also indicate a shift towards a drier climate in the mid part of the middle Miocene (14.25 to 11.35 Ma), with a decrease in the strength of the East Asian summer monsoon (Jiang et al., 2008; Jiang and Ding, 2009). A decrease in the East Asian summer monsoon was also deduced from 12 Ma, in the Sikouzi section, northern China, based on grain-size analysis of windblown sediments (Jiang and Ding, 2010). The increase in the 10-70 μm fraction was interpreted to mean a much stronger winter monsoon at this time.

Lacustrine carbonate isotopic records for a long (29 Myr) sedimentary record in the Linxia Basin, on the northeast side of Tibet (Fig. 1) show stability in climate from 29-12 Ma, with carbonate $\delta^{18}\text{O}$ at roughly -10.5‰, with a switch to more arid

conditions at 12 Ma ($\delta^{18}\text{O}$ at -9‰) that essentially continued through to Pliocene times (Dettman et al., 2003). These authors interpreted 12 Ma as the time at which the Tibetan plateau grew high and wide enough to cause a significant rain shadow in this part of Tibet.

Within the Himalayan foreland in northern Pakistan, Zaleha (1997) found paleosols consistent with a warm, humid climate in the middle Miocene. There was also evidence for pronounced seasonality, in the form of alternations between nodular calcite horizons and iron oxide nodules. The former are evidence for a marked dry season, the latter for a wet season. Therefore the combination and alternation suggests monsoonal conditions similar to the present day.

In summary, high sedimentation rates in many South and East Asian basins are possible evidence for rapid erosion and presumably high precipitation rates by the middle Miocene (Clift et al., 2004). It is not clear if these conditions began at this time or were enhanced, and to what degree they relate to global climatic instability (Clift, 2010) rather than tectonic changes in the Himalaya and Tibet. There is some evidence that a monsoon climate was in operation south of the Himalaya during the middle Miocene, to produce both high precipitation rates and seasonality (Zaleha, 1997). However, other studies suggest an increase in aridity in East Asia at this time (Wei et al., 2006; Wan et al., 2007).

3.3. Late Miocene ages

As many proxy data are now available to suggest early and middle Miocene monsoon climates in south and east Asia, as summarised above, it is notable how

many datasets imply climatic change in the late Miocene (11-5 Ma), particularly in the first half of the sub-epoch (Table 1; Fig. 3b). late Miocene ages include localities to the south of the Himalaya, either in deposits of the Himalayan foreland or further south in the Arabian Sea. But there are more widespread localities with changes at this time, including the South China Sea and the loess-paleosol succession of north and northeast China.

Pedogenic carbonate stable isotopes (carbon and oxygen) have proven to be very useful climate proxies within the syn-orogenic deposits associated with the Himalaya and Tibet. Quade et al (1989) documented late Miocene shifts in $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ towards more positive values from northern Pakistan (Fig. 1). The change in $\delta^{13}\text{C}$ was on the order of 10‰ in 2 Myr, and for $\delta^{18}\text{O}$, ~3‰. This change began at 7.5 – 8 Ma for the $\delta^{13}\text{C}$ shift (using the Cande and Kent (1995) magnetic polarity timescale), and slightly earlier for $\delta^{18}\text{O}$. Quade et al. interpreted the $\delta^{13}\text{C}$ results as recording the shift from C_3 to C_4 type vegetation, i.e. forest to grassland, given that plant-respired CO_2 is the main source of carbon in pedogenic carbonate, and C_3 vegetation has more negative $\delta^{13}\text{C}$ than C_4 vegetation. The spread of C_4 grasslands is in turn taken to represent onset of strong monsoon conditions in the region, because it implies seasonal drought that does not favour forest – even though total rainfall would normally be less than that required to support forest in the same area. Harrison et al. (1993) and Quade et al. (1995) found a similar shift in $\delta^{13}\text{C}$ in pedogenic carbonates from southeast Nepal, at 7 Ma (Fig. 1). Quade et al. (1995) also reported a positive shift in $\delta^{18}\text{O}$ of ~4‰ from the Nepalese sections, occurring at ~6 Ma, i.e. about 2 Myr younger than in Pakistan. Precipitation $\delta^{18}\text{O}$ and hence soil carbonate $\delta^{18}\text{O}$ is controlled by several factors, including the amount of precipitation, evaporation and

the elevation of the source areas. It is not clear what controls these positive shifts, but an increase in precipitation would normally mean a decrease in $\delta^{18}\text{O}$ values while an increase in evaporation would cause an increase in $\delta^{18}\text{O}$ values. Therefore the results might be consistent with an increase in aridity and/or seasonality, but not an overall increase in precipitation.

Increases in $\delta^{13}\text{C}$ and decreases in $\delta^{18}\text{O}$ values of soil carbonate nodules from the Himalayan foreland basin in India were interpreted by Sanyal et al. (2004) to indicate i) a switch from C_3 to C_4 type vegetation, and ii) monsoon onset by 6 Ma, with a possible earlier peak at 10 Ma. Note that the decrease in $\delta^{18}\text{O}$ is in the opposite direction to the results of Quade et al. (1989) and Harrison et al (1993), such that both sets of authors use their results to indicate monsoon intensification at about the same time, but for very different reasons. Galy et al. (2010) noted an increase in $\delta^{13}\text{C}$ for organic matter in the Bengal fan at 7.4 Ma, also consistent with a change to C_4 type vegetation in the sediment source areas.

Dettman et al. (2001) used $\delta^{18}\text{O}$ trends in freshwater bivalve shells and mammal teeth to study seasonal variation in Himalayan foreland surface waters in the late Miocene and Pliocene. The rather limited change in values over this period was used to infer a strong Asian monsoon from 10.7 Ma, with the implication that the Tibetan plateau was high and wide enough through this time to create a monsoon system similar to the present day. Wet-season rainfall was significantly more negative (-9.5‰) before 7.5 Ma than afterwards (-6.5‰). This could be evidence for increased aridity from 7.5 Ma.

372 The isotopic shifts from the Himalayan foreland are supported by
 373 palaeontological evidence from the same region. Vertebrate palaeontology studies
 374 were at the forefront of evidence for climatic change near the Himalaya in the late
 375 Cenozoic. Flynn and Jacobs (1982); Barry et al. (1985) and Barry (2002) used
 376 changes in rodents and larger herbivores to infer a shift towards grassland habitats in
 377 northern Pakistan by ~7 Ma. This is supported by the fluvial sedimentology of the
 378 host strata for these fossils, which shows a change to a more seasonal discharge and
 379 increased aridity (Barry et al., 2002). Fossil leaf and pollen show a transition from
 380 forest to grassland in central Nepal at ~8 to 6.5 Ma (Hoorn et al., 2000).

381

382 Sedimentary records from the western Arabian Sea provide evidence for
 383 increased seasonality around 8 Ma, attributed to strengthening monsoon winds and
 384 increased upwelling in this region (Kroon et al., 1991; Prell et al., 1992). The key data
 385 point is the increased component of the planktonic foraminifer *Globigerina bulloides*
 386 in the sediment at this time, jumping from a few percent of the annual mass
 387 accumulation to tens of percent. *G. bulloides*, a subpolar species, flourishes in
 388 monsoon wind induced-upwelling in the coastal waters of the tropics like the NW
 389 Arabian Sea, making it an effective proxy for seasonality in this region, although not
 390 necessarily a gauge of which wind system is the stronger, summer or winter.

391

392 Loess records from northern China have provided support for a climate change
 393 in the late Miocene, and the appearance of such deposits may be taken as evidence for
 394 the establishment of wind systems close to the modern regime, influenced by the East
 395 Asian monsoon. Qiang et al (2001) derived a magnetostratigraphic age of 8.35 ± 0.05
 396 Ma for the base of aeolian red clay beneath the Plio-Pleistocene loess-paleosol

succession at Jiaxian, northern China (Fig. 1). These deposits are heavily modified by pedogenesis, and suggest a warm and humid climate developed at this time. Xu et al. (2009) have recently taken the eastern loess record back to 11 Ma, based on magnetostratigraphy. A pulse of aeolian dust in the North Pacific at ~8 Ma was recorded in sediments from ODP site 885/886 (Rea et al., 1998), interpreted as transported by stronger westerlies from northeast Asia, thereby strengthening the case for increased dust production within northern China at this time.

Saylor et al. (2009) used isotope data from molluscs and plants from the Zhada Basin, southwest Tibet (Fig. 1), to infer a cold, arid climate similar to the modern since ~9 Ma, but with a shift from C₃ to C₄ type vegetation. Palaeoaltimetry estimates suggested a drop in elevation of ~1.5 km in this time. These results indicate no late Miocene climate change, but a change in vegetation patterns that presumably results from other causes. Fan et al. (2007) studied $\delta^{18}\text{O}$ records of Neogene deposits of the Linxia Basin (Fig. 1), aged between 13.1 and 4.3 Ma. Between 13.1 and 8 Ma, strong oscillations between dry and wet conditions were inferred from fluctuations in the $\delta^{18}\text{O}$ records, and interpreted as changes from hydrographically closed and open lake systems. A short-lived arid pulse took place from 9.6 to 8.5 Ma. After ~8 Ma, the climate was more stable, less arid and the lake system was hydrographically open. Cooler and/or drier conditions prevailed from 5.3 Ma.

Zheng et al (2004) found a decrease of the abundance ratio of planktonic foraminifera *Globigerinoides sacculifer*/*G. ruber* and increase of *Neoglobobulimina* at ~8 Ma at ODP site 1146 in the South China Sea, which they interpreted as a lowering of the sea surface temperature and increased productivity, related to an

intensification of East Asian winter monsoon winds and upwelling. Steinke et al. (2010) have recently examined temperature independent variations in seawater $\delta^{18}\text{O}$ (i.e. a proxy for sea surface salinity) for the ODP site 1146 using a combined approach of planktonic foraminiferal Mg/Ca ratios and $\delta^{18}\text{O}$ analysis. They found that local seawater $\delta^{18}\text{O}$ increased (became heavier) by 0.34‰, interpreted as a sharp reduction in the East Asian summer monsoon intensity at ~7.5 Ma. This is consistent with results of Wan et al (2007) and Wei et al (2006) who studied shifts in chemical weathering patterns of detrital sediment derived from southeast Asia (within the South China Sea), and interpreted an intensification of the winter monsoon and a decline in the summer monsoon at ~8 Ma.

Sediment flux records for the late Miocene commonly show a decline for regions sourced from the Himalaya and Tibet, in contrast to the increase expected if the monsoon intensified at this time (Burbank et al., 1993; Clift et al., 2008). Such flux reductions are true of both the Indus and Bengal fans (Burbank et al., 1993; Clift et al., 2008), which are major recipients of Himalayan and Tibetan sediment.

Inferred climatic changes in Asia at 10-8 Ma have to be put in a global context. There is evidence from the Indian, equatorial Pacific and southern Atlantic oceans of a biogenic bloom in the late Miocene – early Pliocene (Farrell et al., 1995; Dickens and Owen, 1999). Gupta et al (2004) reported a major increase in biogenic productivity in the eastern Indian Ocean at 10-8 Ma, based on the establishment of benthic biofacies that indicate a year-round high flux of organic matter from the sea surface to the ocean floor. They suggested that the oceanic high productivity occurred across a larger region than that affected by the Asian monsoons or detritus produced

by monsoonal climate, and that increased upwelling driven by strengthening wind systems was a more likely cause, itself created by high-southern-latitude cooling and increased ice volume.

4. Changing latitudes of the ITCZ and the Himalaya

We reviewed the evidence for the palaeolatitude of the ITCZ in the Pacific in Armstrong and Allen (2011) and so only summarise the data here (Fig. 3c). Different lines of evidence used include the position of the smectite-illite transition (because this marks the boundary between wind-blown clay derived predominantly from Asia and from Central/South America; Lyle et al., 2002), and the composition of manganese crusts (because this changes depending on changing biologic and detrital fluxes) (Kim et al., 2006; Hyeong et al., 2005). Both sets of records indicate that the mean position of the ITCZ has moved southwards in the Neogene. Differences in latitudes between the Central and Western Pacific are consistent with present day variation, with Western Pacific palaeolatitudes consistently lying closer to the equator.

Himalayan palaeolatitudes are hard to know precisely, given that the overall plate convergence of India and Eurasia has been partly taken up in the range (perhaps 1000 km out of 2500 km total convergence), making the exercise more difficult than tracking the relative motion of stable plate interiors. There is no single set of values that represents the whole of the Himalaya. Molnar and Stock (2009) calculated palaeo-positions for two localities on the northwest and northeast side of the Indian plate. Armstrong and Allen (2011) added 500 km to the latitude of these points for each time, to more closely represent the Himalayan palaeolatitudes – with the caveats

about the “concertina” effect of Himalayan thrusting just noted. These positions are plotted on Fig. 3c alongside the Pacific ITCZ values. As Armstrong and Allen (2011) discussed, the early Miocene (~20 Ma) latitudes of the Central Pacific ITCZ and the Himalaya were over 2000 km closer than at present, and may have overlapped, at least in the northwest Himalaya. Here, we note the degree of divergence by 10 Ma. The palaeolatitudes had separated: the Central Pacific ITCZ had migrated ~1500 km south of its early Miocene position (Lyle et al., 2002) and the northern margin of the Indian plate/Himalaya had migrated northwards, probably by 200-400 km (Molnar and Stock, 2009).

5. Discussion and conclusions

The above review demonstrates that there are abundant data to indicate Asian monsoon intensification in both the early and late Miocene. If the question is strictly “when did a monsoon system begin?” the answer appears to be the early Miocene. Data from this timeframe indicate many of the climatic characteristics associated with the modern monsoon: high erosion rates from the Himalaya (attributed to high precipitation); climatic seasonality (related to the shift in wind direction between summer and winter); palaeobotanical evidence for humid climate through eastern China (consistent with the range of the modern East Asian summer monsoon); loess deposition in northern China (equivalent to the modern dust blown in to this region from further north). Given that the early Miocene time period is roughly 30 million years after initial India-Eurasia collision, it follows that a threshold was crossed at this time to trigger the monsoon in both its Indian and East Asian versions. There was plausibly a combination of the lateral and/or vertical growth of the Himalaya and Tibet (Tapponnier et al., 2001; Harris, 2007), the convergence of the ITCZ and the

497 Himalaya (Armstrong and Allen, 2011), drying of Paratethys across central Asia
 498 (Ramstein et al., 1997) and transgression across the East Asian continental margin
 499 (Zhang et al., 2007).

500

501 The Early Miocene was globally a time of relatively warm, humid climate
 502 (e.g. Zachos et al., 2001), albeit punctuated by Antarctic glaciations (Miller et al.,
 503 1991). Amongst other possible mechanisms, this mild global climate would have been
 504 promoted by high rates of combined metamorphism and exhumation in the
 505 Himalayas, via rapid fluxes of metamorphic CO₂ to the atmosphere (applying the
 506 reasoning of Becker et al., 2008; Skelton, 2011). The decrease in Himalayan
 507 metamorphism and exhumation rates in the middle Miocene would presumably have
 508 reduced this CO₂ flux, at the time the global climate swung in to a colder mode.

509

510 This leaves the question of the significance of climatic changes reported or
 511 inferred for ~8 Ma. Molnar et al. (1993) proposed a correspondence between tectonic
 512 events in the Himalayan/Tibetan system at roughly this time and the climatic changes
 513 in Asia. In this model increased surface elevations of the Tibetan plateau, perhaps
 514 caused by partial loss of the lower lithosphere, would lead to an intensified monsoon
 515 system whether this was related to the physical barrier presented by the elevated
 516 topography or the increased heating of the atmosphere (Molnar et al., 2010). Increased
 517 seasonality at ~8 Ma fits this scenario, whether it is interpreted from the marine
 518 sedimentary record (Kroon et al., 1991) or terrestrial deposits (Quade et al., 1989; An
 519 et al., 2001). However, the evidence for lithosphere loss and surface uplift at 8 Ma is
 520 now debatable (Molnar, 2005), as described above, which casts doubt on the model.

521

Also, the story at 8 Ma appears more complex than a simple, regional intensification of the monsoon in all of its forms. Sedimentation rates in the basins receiving Himalayan detritus record a slow down in the late Miocene (Clift et al., 2008). Multi-isotope (Sr, Nd and Os) were used as proxies for the source of the sediments deposited in the Bengal Fan over the last 12 million years by Galy et al. (2010), and their data indicated a stable provenance over this time, without evidence for changing drainage or uplift histories. Biological productivity increased in the Atlantic, Pacific and Indian oceans far away from the influence of these Himalayan sediments (Dickens and Owen, 1999; Gupta et al., 2004), and wind-blown dust increased in the North Pacific (Rea et al., 1998). This geographical range of events begs the question whether changes in Asia's orography influenced global climate patterns, or global climate change affected the climate in the Himalaya, Tibet and adjacent regions.

There is another aspect to the 8 Ma palaeoclimate record that needs to be addressed. Much of the published evidence relates to increased seasonality and/or increased aridity at this time (Table 1). Not so many data point to increased overall precipitation. Several lines of evidence suggest the strengthening of the East Asian winter monsoon and weakening of the summer monsoon across northern and eastern China (e.g. Wan et al., 2007), consistent with a trend towards increased aridity over much of China. This is surprising, as so much of the modern monsoon impact on South and East Asia is through high precipitation and its effects on the landscape. The late Miocene decline in sedimentation rates across South and East Asia is likewise difficult to explain in terms of late Miocene monsoon enhancement, if that enhancement should mean increased precipitation (Clift and Gaedicke, 2002; Clift et

al., 2008). It is also notable that recent studies of palaeoclimate and palaeoaltitude within the interior of Tibet are concluding that altitudes were similar to the present and climate was similar to the present since the late Miocene or earlier (Fig. 1; DeCelles et al., 2007; Saylor et al., 2009).

The spatial separation of the ITCZ and the Himalaya/Tibet may provide an explanation for the climatic and depositional patterns through the Miocene (Figs 3 and 4). Fig. 3 shows the increase in the latitudinal separation of the Central Pacific ITCZ and the Himalaya increased by roughly 1500-1900 km by the late Miocene, compared with the early Miocene. It is possible that the increasing separation of the ITCZ and the Himalaya led to a more seasonal and arid climate, as the ITCZ was only moved northwards for a few months of the year, leading to heavy summer monsoon rains and a relatively arid winter monsoon climate south of the Himalaya (like the modern case; Fig. 2) and a more arid subtropical climate generally across north and east China. The southward movement of the descending limb of the Hadley cell is also important in controlling climatic shifts: as the mean ITCZ position moved south through the Miocene, so presumably did each Hadley cell to its north and south, bringing drier winter climates to regions such as the Himalayan foreland. This climatic evolution is summarised on Fig. 4.

The explanation for the late Cenozoic variation in the latitude of the ITCZ is likely to be the emergence of a bipolar world (Miller et al., 1991; Rea, 1994; Zachos et al., 2001), as increased northern hemisphere glaciation produced hemispheric wind systems that were closer to mirror images of each other. The mean position of the modern ITCZ some 6 degrees north of the equator is a consequence of the greater size

of the Antarctic ice sheet than its Arctic counterpart. Pre-Pleistocene climates had more a northerly ITCZ, caused by the greater asymmetry between the north and south polar regions and the lesser strength of northern hemisphere wind systems. A world lacking ice sheets at either pole would likewise have an ITCZ located nearer the equator.

There is no step-change in the separation of the ITCZ and the Himalaya in the late Miocene, at least not in available data (Fig. 3). One possibility is that a threshold was crossed in the climate system, such that an incremental increase in the ITCZ/Himalayan separation caused a climate re-organisation. Another possibility is that the increased latitudinal separation acted in concert with global trends towards a cooler late Cenozoic climate, but there is no clear shift in the global oceanic $\delta^{18}\text{O}$ proxy at this specific time (Zachos et al., 2001). Either way, we propose that the 8 Ma climatic shifts be regarded as the intensification of the both the South and East Asian monsoon's seasonal aspect, but a decline in terms of precipitation rates.

Our emphasis in this paper and Armstrong and Allen (2011) has been on the effect of the distance between the mean ITCZ and the Himalaya in terms of climate. It is worth remembering the tectonic changes in the Himalaya and southern Tibet through the Miocene, as over the middle Miocene the regime changed from the synchronous ductile shear along the MCT and STDS, with co-eval leucogranite generation and rapid exhumation, to the pattern of ~north-south normal faulting in southern Tibet and thrusting along the MBT and MFT that continues to the present day. As more age constraints are published the time gap between these contrasting styles reduces: the existing data suggest that at least a modest overlap of no more than

a few million years on the scale of the entire orogen (Blisniuk et al., 2001; Searle et al., 2003; Harris, 2007), but in each individual area ~east-west extension postdates the top-to-north extensional shear along the STDS (Leloup et al., 2010).

We speculate that the reduction in precipitation over the Himalaya, driven by southwards ITCZ migration, may have played a role in thrust migration in the Himalaya. The mid Miocene saw a reconfiguration of the Himalaya, with new thrusting (MBT) breaking to the south of the MCT (e.g. Meigs et al., 1995). Previous models have made a link between the changing tectonics in southern Tibet and the reorganisation of thrusting in the Himalaya, with Tibetan surface uplift (suggested to be driven by lower lithosphere detachment) driving the migration of Himalayan thrusting southwards so as to re-establish a critically-tapered orogenic wedge (Harrison et al., 1992; Huyghe et al., 2001). The lack of identified surface uplift within southern Tibet, of the right age, is a problem for this model, as discussed above. However, it is possible that thrust migration was at least a partial response to changing Himalayan climate, and a represented a shift to a drier, wider, orogenic belt (e.g. Jamieson and Beaumont, 1988; Whipple and Meade, 2006).

Acknowledgements

“Nos esse quasi nanos, gigantium humeris insidentes”. We are also grateful for the reviews and the editor’s comments, which improved the first version of the manuscript.

References

- Aikman, A.B., Harrison, T.M., Lin, D., 2008. Evidence for Early (> 44 Ma) Himalayan Crustal Thickening, Tethyan Himalaya, southeastern Tibet. *Earth and Planetary Science Letters*, 274, 14-23.
- An, Z., Kutzbach, J.E., Prell, W.L., Porter, S.C., 2001. Evolution of Asian monsoons and phased uplift of the Himalayan Tibetan plateau since Late Miocene times. *Nature*, 411, 62-66.
- Armstrong, H.A., Allen, M.B., 2011. Shifts in the Intertropical Convergence Zone, Himalayan exhumation and late Cenozoic climate. *Geology*, 39, 11-14.
- Barry, J.C., Johnson, N.M., Raza, S.M., Jacobs, L.L., 1985. Neogene mammalian faunal change in southern Asia - correlations with climatic, tectonic and eustatic events. *Geology*, 13, 637-640.
- Barry, J.C., Morgan, M.L.E., Flynn, L.J., Pilbeam, D., Behrensmeyer, A.K., Raza, S.M., Khan, I.A., Badgley, C., Hicks, J., Kelley, J., 2002. Faunal and environmental change in the late Miocene Siwaliks of northern Pakistan. *Paleobiology*, 28, 1-71.
- Beaumont, C., Jamieson, R.A., Nguyen, M.H., Lee, B., 2001. Himalayan tectonics explained by extrusion of a low-viscosity crustal channel coupled to focused surface denudation. *Nature*, 414, 738-742.
- Becker J.A., Bickle M.J., Galy A., Holland, T.J.B., 2008. Himalayan metamorphic CO₂ fluxes: Quantitative constraints from hydrothermal springs. *Earth and Planetary Science Letters*, 265, 616-629.
- Blisniuk, P.M., Hacker, B.R., Glodny, J., Ratschbacher, L., Bi, S.W., Wu, Z.H., McWilliams, M.O., Calvert, A., 2001. Normal faulting in central Tibet since at least 13.5 Myr ago. *Nature*, 412, 628-632.

- 647 Boos, W.R., Kuang, Z.M., 2010. Dominant control of the South Asian monsoon by
648 orographic insulation versus plateau heating. *Nature*, 463, 218-U102.
- 649 Booth, A.L., Chamberlain, C.P., Kidd, W.S.F., Zeitler, P.K., 2009. Constraints on the
650 metamorphic evolution of the eastern Himalayan syntaxis from geochronologic
651 and petrologic studies of Namche Barwa. *Geological Society of America*
652 *Bulletin*, 121, 385-407.
- 653 Burbank, D.W., Derry, L.A., Francelanord, C., 1993. Reduced Himalayan sediment
654 production 8 Myr ago despite an intensified monsoon. *Nature*, 364, 48-50.
- 655 Burchfiel, B.C., Chen, Z., Hodges, K.V., Liu, Y., Royden, L.H., Deng, C., Xu, J.,
656 1992. The South Tibetan Detachment System, Himalayan orogen: extension
657 contemporaneous with and parallel to shortening in a collisional mountain belt.
658 *Special Paper of the Geological Society of America*, 269, 41 pp.
- 659 Cande, S.C., Kent, D.V., 1995. Revised calibration of the geomagnetic polarity
660 timescale for the Late Cretaceous and Cenozoic. *Journal of Geophysical*
661 *Research-Solid Earth*, 100, 6093-6095.
- 662 Catlos, E.J., Dubey, C.S., Harrison, T.M., Edwards, M.A., 2004. Late Miocene
663 movement within the Himalayan Main Central Thrust shear zone, Sikkim, north-
664 east India. *Journal of Metamorphic Geology*, 22, 207-226.
- 665 Clark, M.K., House, M.A., Royden, L.H., Whipple, K.X., Burchfiel, B.C., Zhang, X.,
666 Tang, W., 2005. Late Cenozoic uplift of southeastern Tibet. *Geology*, 33, 525-
667 528.
- 668 Clift, P., Gaedicke, C., 2002. Accelerated mass flux to the Arabian Sea during the
669 middle to late Miocene. *Geology*, 30, 207-210.
- 670 Clift, P.D., 2006. Controls on the erosion of Cenozoic Asia and the flux of clastic
671 sediment to the ocean. *Earth and Planetary Science Letters*, 241, 571-580.

- 672 Clift, P.D., 2010. Enhanced global continental erosion and exhumation driven by
 673 Oligo-Miocene climate change. *Geophysical Research Letters*, 37, art. no.
 674 L09402, doi: 10.1029/2010gl043067.
- 675 Clift, P.D., Hodges, K.V., Heslop, D., Hannigan, R., Van Long, H., Calves, G., 2008.
 676 Correlation of Himalayan exhumation rates and Asian monsoon intensity.
 677 *Nature Geoscience*, 1, 875-880.
- 678 Clift, P.D., Layne, G.D., Blusztajn, J., 2004. Marine sedimentary evidence for
 679 Monsoon strengthening, Tibetan uplift and drainage evolution in East Asia. In:
 680 Clift, P., Kuhnt, W., Wang, P., Hayes, D. (Eds.), *Continent-Ocean Interactions*
 681 *within East Asian Marginal Seas*, American Geophysical Union, Washington,
 682 vol. 149, pp. 255-282.
- 683 Coleman, M., Hodges, K., 1995. Evidence for Tibetan plateau uplift before 14 Myr
 684 ago from a new minimum age for east-west extension. *Nature*, 374, 49-52.
- 685 Cook, K.L., Royden, L.H., 2008. The role of crustal strength variations in shaping
 686 orogenic plateaus, with application to Tibet. *Journal of Geophysical Research-*
 687 *Solid Earth*, 113.
- 688 DeCelles, P.G., Quade, J., Kapp, P., Fan, M.J., Dettman, D.L., Ding, L., 2007. High
 689 and dry in central Tibet during the Late Oligocene. *Earth and Planetary Science*
 690 *Letters*, 253, 389-401.
- 691 Dettman, D.L., Fang, X.M., Garzzone, C.N., Li, J.J., 2003. Uplift-driven climate
 692 change at 12 Ma: a long delta O-18 record from the NE margin of the Tibetan
 693 plateau. *Earth and Planetary Science Letters*, 214, 267-277.
- 694 Dettman, D.L., Kohn, M.J., Quade, J., Ryerson, F.J., Ojha, T.P., Hamidullah, S.,
 695 2001. Seasonal stable isotope evidence for a strong Asian monsoon throughout
 696 the past 10.7 m.y. *Geology*, 29, 31-34.

- 697 Dickens, G.R., Owen, R.M., 1999. The Latest Miocene-Early Pliocene biogenic
698 bloom: a revised Indian Ocean perspective. *Marine Geology*, 161, 75-91.
- 699 Dupont-Nivet, G., Lippert, P.C., van Hinsbergen, D.J.J., Meijers, M.J.M., Kapp, P.,
700 2010. Palaeolatitude and age of the Indo-Asia collision: palaeomagnetic
701 constraints. *Geophysical Journal International*, 182, 1189-1198.
- 702 England, P., Houseman, G., 1989. Extension during continental convergence, with
703 application to the Tibetan plateau. *Journal of Geophysical Research-Solid Earth*
704 and Planets, 94, 17561-17579.
- 705 Fan, M.J., Dettman, D.L., Song, C.H., Fang, X.M., Garzione, C.N., 2007. Climatic
706 variation in the Linxia basin, NE Tibetan Plateau, from 13.1 to 4.3 Ma: The
707 stable isotope record. *Palaeogeography Palaeoclimatology Palaeoecology*, 247,
708 313-328.
- 709 Farrell, J.W., Raffi, I., Janecek, T.C., Murray, D.W., Levitan, M., Dadey, K.A.,
710 Emeis, K.-C., Lyle, M., Flores, J.-A., Hovan, S., 1995. Late Neogene
711 sedimentation patterns in the eastern equatorial Pacific Ocean. In: Pisias, N. G.,
712 Mayer, L. A., Janecek, T. R., Palmer-Julson, A., van Andel, T. H. (Eds.),
713 *Proceedings of the Ocean Drilling Program, Scientific Results*, vol. 138, pp.
714 717–756.
- 715 Flynn, L.J., Jacobs, L.L., 1982. Effects of changing environments on Siwalik rodent
716 faunas of northern Pakistan. *Palaeogeography Palaeoclimatology Palaeoecology*,
717 38, 129-138.
- 718 Galy, V., France-Lanord, C., Peucker-Ehrenbrink, B., Huyghe, P., 2010. Sr-Nd-Os
719 evidence for a stable erosion regime in the Himalaya during the past 12 Myr.
720 *Earth and Planetary Science Letters*, 290, 474-480.

- 721 Garzione, C.N., Quade, J., DeCelles, P.G., English, N.B., 2000. Predicting
 722 paleoelevation of Tibet and the Himalaya from delta O-18 vs. altitude gradients
 723 in meteoric water across the Nepal Himalaya. *Earth and Planetary Science*
 724 *Letters*, 183, 215-229.
- 725 Guo, Z.T., Ruddiman, W.F., Hao, Q.Z., Wu, H.B., Qiao, Y.S., Zhu, R.X., Peng, S.Z.,
 726 Wei, J.J., Yuan, B.Y., Liu, T.S., 2002. Onset of Asian desertification by 22 Myr
 727 ago inferred from loess deposits in China. *Nature*, 416, 159-163.
- 728 Gupta, A.K., Singh, R.K., Joseph, S., Thomas, E., 2004. Indian Ocean high-
 729 productivity event (10-8 Ma): Linked to global cooling or to the initiation of the
 730 Indian monsoons? *Geology*, 32, 753-756.
- 731 Harris, N., 2007. Channel flow and the Himalayan-Tibetan orogen: a critical review.
 732 *Journal of the Geological Society*, 164, 511-523.
- 733 Harris, N., Massey, J., 1994. Decompression and anatexis of Himalayan metapelites.
 734 *Tectonics*, 13, 1537-1546.
- 735 Harrison, T.M., Copeland, P., Kidd, W.S.F., Yin, A., 1992. Raising Tibet. *Science*,
 736 255, 1663-1670.
- 737 Harrison, T.M., Copeland, P., Hall, S.A., Quade, J., Burner, S., Ojha, T.P., Kidd,
 738 W.S.F., 1993. Isotopic preservation of Himalayan-Tibetan uplift, denudation and
 739 climatic histories of two molasse deposits. *Journal of Geology*, 101, 157-175.
- 740 Herman, F., Copeland, P., Avouac, J.P., Bollinger, L., Maheo, G., Le Fort, P., Rai, S.,
 741 Foster, D., Pecher, A., Stuwe, K., Henry, P., 2010. Exhumation, crustal
 742 deformation, and thermal structure of the Nepal Himalaya derived from the
 743 inversion of thermochronological and thermobarometric data and modeling of
 744 the topography. *Journal of Geophysical Research-Solid Earth*, 115, art. no.
 745 B06407, doi: 10.1029/2008jb006126.

- 746 Hodges, K., Bowring, S., Davidek, K., Hawkins, D., Krol, M., 1998. Evidence for
747 rapid displacement on Himalayan normal faults and the importance of tectonic
748 denudation in the evolution of mountain ranges. *Geology*, 26, 483-486.
- 749 Hodges, K.V., 2000. Tectonics of the Himalaya and southern Tibet from two
750 perspectives. *Bulletin of the Geological Society of America*, 112, 324-350.
- 751 Houseman, G.A., McKenzie, D.P., Molnar, P., 1981. Convective instability of a
752 thickened boundary layer and its relevance for the thermal evolution of
753 continental convergent belts. *Journal of Geophysical Research*, 86, 6115-6132.
- 754 Huyghe, P., Galy, A., Mugnier, J.L., France-Lanord, C., 2001. Propagation of the
755 thrust system and erosion in the Lesser Himalaya: Geochemical and
756 sedimentological evidence. *Geology*, 29, 1007-1010.
- 757 Hyeong, K., Park, S.H., Yoo, C.M., Kim, K.H., 2005. Mineralogical and geochemical
758 compositions of the eolian dust from the northeast equatorial Pacific and their
759 implications on paleolocation of the Intertropical Convergence Zone.
760 *Paleoceanography*, 20, art. no. Pa1010, doi: 10.1029/2004pa001053.
- 761 Jamieson, R.A., Beaumont, C., 1988. Orogeny and metamorphism - a model for
762 deformation and pressure-temperature-time paths with applications to the central
763 and southern Appalachians. *Tectonics*, 7, 417-445.
- 764 Jiang, H.C., Ji, J.L., Gao, L., Tang, Z.H., Ding, Z.L., 2008. Cooling-driven climate
765 change at 12-11 Ma: Multiproxy records from a long fluviolacustrine sequence
766 at Guyuan, Ningxia, China. *Palaeogeography Palaeoclimatology Palaeoecology*,
767 265, 148-158.
- 768 Jiang, H., Ding, Z., 2009. Spatial and temporal characteristics of Neogene palynoflora
769 in China and its implication for the spread of steppe vegetation. *Journal of Arid*.
770 *Environments*, 73, 765-772.

- 771 Jiang, H.C., Ding, Z.L., 2010. Eolian grain-size signature of the Sikouzi lacustrine
 772 sediments (Chinese Loess Plateau): Implications for Neogene evolution of the
 773 East Asian winter monsoon. *Geological Society of America Bulletin*, 122, 843-
 774 854.
- 775 Kali, E., Leloup, P.H., Arnaud, N., Maheo, G., Liu, D.Y., Boutonnet, E., Van der
 776 Woerd, J., Liu, X.H., Jing, L.Z., Li, H.B., 2010. Exhumation history of the
 777 deepest central Himalayan rocks, Ama Drime range: Key pressure-temperature-
 778 deformation-time constraints on orogenic models. *Tectonics*, 29, art. no. Tc2014
 779 doi: 10.1029/2009tc002551.
- 780 Kalnay, E., Kanamitsu, M., Kistler, R., Collins, W., Deaven, D., Gandin, L., Iredell,
 781 M., Saha, S., White, G., Woollen, J., Zhu, Y., Chelliah, M., Ebisuzaki, W.,
 782 Higgins, W., Janowiak, J., Mo, K.C., Ropelewski, C., Wang, J., Leetmaa, A.,
 783 Reynolds, R., Jenne, R., Joseph, D., 1996. The NCEP/NCAR 40-year reanalysis
 784 project. *Bulletin of the American Meteorological Society*, 77, 437-471.
- 785 Kellett, D.A., Grujic, D., Erdmann, S., 2009. Miocene structural reorganization of the
 786 South Tibetan detachment, eastern Himalaya: Implications for continental
 787 collision. *Lithosphere*, 1, 259-281.
- 788 Kent-Corson, M.L., Ritts, B.D., Zhuang, G.S., Bovet, P.M., Graham, S.A.,
 789 Chamberlain, C.P., 2009. Stable isotopic constraints on the tectonic,
 790 topographic, and climatic evolution of the northern margin of the Tibetan
 791 Plateau. *Earth and Planetary Science Letters*, 282, 158-166.
- 792 Kim, J., Hyeong, K., Jung, H.S., Moon, J.W., Kim, K.H., Lee, I., 2006. Southward
 793 shift of the Intertropical Convergence Zone in the western Pacific during the late
 794 Tertiary: Evidence from ferromanganese crusts on seamounts west of the

- 795 Marshall Islands. *Paleoceanography*, 21, art. no. Pa4218, doi:
 796 10.1029/2006pa001291.
- 797 Kirby, E., Reiners, P.W., Krol, M.A., Whipple, K.X., Hodges, K.V., Farley, K.A.,
 798 Tang, W.Q., Chen, Z.L., 2002. Late Cenozoic evolution of the eastern margin of
 799 the Tibetan Plateau: Inferences from Ar-40/Ar-39 and (U-Th)/He
 800 thermochronology, *Tectonics*, 21, doi: 10.1029/2000TC001246.
- 801 Kroon, D., Steens, T., Troelstra, S.R., 1991. Onset of monsoonal related upwelling in
 802 the western Arabian Sea as revealed by planktonic foraminifera, *Proceedings*
 803 *of the Ocean Drilling Program, Scientific Results*, 117, 257-263.
- 804 Kutzbach, J.E., Guetter, P.J., Ruddiman, W.F., Prell, W.L., 1989. Sensitivity of
 805 climate to Late Cenozoic uplift in southern Asia and the American west -
 806 numerical experiments. *Journal of Geophysical Research-Atmospheres*, 94,
 807 18393-18407.
- 808 Lave, J., Avouac, J.P., 2000. Active folding of fluvial terraces across the Siwaliks
 809 Hills, Himalayas of central Nepal. *Journal of Geophysical Research-Solid Earth*,
 810 105, 5735-5770.
- 811 Leloup, P.H., Maheo, G., Arnaud, N., Kali, E., Boutonnet, E., Liu, D.Y., Liu, X.H.,
 812 Li, H.B., 2010. The South Tibet detachment shear zone in the Dinggye area
 813 Time constraints on extrusion models of the Himalayas. *Earth and Planetary*
 814 *Science Letters*, 292, 1-16.
- 815 Lyle, M.W., Wilson, P.A., Janecek, T.R., and shipboard party, 2002. Leg 199
 816 Summary. In: Lyle, M. W., et al. (Eds.), *Proceedings of the Ocean Drilling*
 817 *Program, Initial reports*, vol. 199, pp. 1-89.
- 818 Maheo, G., Leloup, P.H., Valli, F., Lacassin, R., Arnaud, N., Paquette, J.L.,
 819 Fernandez, A., Haibing, L., Farley, K.A., Tapponnier, P., 2007. Post 4 Ma

- 820 initiation of normal faulting in southern Tibet. Constraints from the Kung Co
 821 half graben. *Earth and Planetary Science Letters*, 256, 233-243.
- 822 Meigs, A.J., Burbank, D.W., Beck, R.A., 1995. Middle-Late Miocene (greater than 10
 823 Ma) formation of the Main Boundary Thrust in the western Himalaya. *Geology*,
 824 23, 423-426.
- 825 Miller, K.G., Fairbanks, R.G., Mountain, G.S., 1987. Tertiary oxygen isotope
 826 synthesis, sea level history, and continental margin erosion. *Paleoceanography*,
 827 2, 1-19.
- 828 Miller, K.G., Wright, J.D., Browning, J.V., 2005. Visions of ice sheets in a
 829 greenhouse world. *Marine Geology*, 217, 215-231.
- 830 Miller, K.G., Wright, J.D., Fairbanks, R.G., 1991. Unlocking the ice house -
 831 Oligocene-Miocene oxygen isotopes, eustasy, and margin erosion. *Journal of*
 832 *Geophysical Research-Solid Earth and Planets*, 96, 6829-6848.
- 833 Molnar, P., 2005. Mio-pliocene growth of the Tibetan Plateau and evolution of East
 834 Asian climate, *Palaeontol. electron.*, 8(1):2A.
- 835 Molnar, P., Boos, W.R., Battisti, D.S., 2010. Orographic controls on climate and
 836 paleoclimate of Asia: Thermal and mechanical roles for the Tibetan Plateau.
 837 *Annual Review of Earth and Planetary Sciences*, 38, 77-102.
- 838 Molnar, P., England, P., Martinod, J., 1993. Mantle dynamics, uplift of the Tibetan
 839 plateau and the Indian monsoon. *Reviews of Geophysics*, 31, 357-396.
- 840 Molnar, P., Lyon-caen, H., 1989. Fault plane solutions of earthquakes and active
 841 tectonics of the Tibetan Plateau and its margins. *Geophysical Journal*
 842 *International*, 99, 123-153.

- 843 Molnar, P., Stock, J.M., 2009. Slowing of India's convergence with Eurasia since 20
 844 Ma and its implications for Tibetan mantle dynamics. *Tectonics*, 28, doi:
 845 10.1029/2008tc002271.
- 846 Molnar, P., Tapponnier, P., 1978. Active tectonics of Tibet. *Journal of Geophysical*
 847 *Research*, 83, 5361-5375.
- 848 Murphy, M.A., Saylor, J.E., Ding, L., 2009. Late Miocene topographic inversion in
 849 southwest Tibet based on integrated paleoelevation reconstructions and
 850 structural history. *Earth and Planetary Science Letters*, 282, 1-9.
- 851 Najman, Y., Bickle, M., BouDagher-Fadel, M., Carter, A., Garzanti, E., Paul, M.,
 852 Wijbrans, J., Willett, E., Oliver, G., Parrish, R., Akhter, S.H., Allen, R., Ando,
 853 S., Chisty, E., Reisberg, L., Vezzoli, G., 2008. The Paleogene record of
 854 Himalayan erosion: Bengal Basin, Bangladesh. *Earth and Planetary Science*
 855 *Letters*, 273, 1-14.
- 856 Ouimet, W., Whipple, K., Royden, L., Reiners, P., Hodges, K., Pringle, M., 2010.
 857 Regional incision of the eastern margin of the Tibetan Plateau. *Lithosphere*, 2,
 858 50-63.
- 859 Prell, W.L., Murray, W.D., Clemens, S.C., Anderson, D.M., 1992. Evolution and
 860 variability of the Indian Ocean summer monsoon: evidence from the western
 861 Arabian Sea drilling program. In: Duncan, R. A., Rea, D. K., Kidd, R. B., von
 862 Rad, U., Weissel, J. K. (Eds.), *Synthesis of Results from Scientific Drilling in*
 863 *the Indian Ocean*, Geophysical Monograph 70. American Geophysical Union,
 864 Washington, DC, pp. 447-469.
- 865 Priestley, K., McKenzie, D., 2006. The thermal structure of the lithosphere from shear
 866 wave velocities. *Earth and Planetary Science Letters*, 244, 285-301.

- 867 Qiang, X.K., Li, Z.X., Powell, C.M., Zheng, H.B., 2001. Magnetostratigraphic record
868 of the Late Miocene onset of the East Asian monsoon, and Pliocene uplift of
869 northern Tibet. *Earth and Planetary Science Letters*, 187, 83-93.
- 870 Quade, J., Cater, J.M.L., Ojha, T.P., Adam, J., Harrison, T.M., 1995. Late Miocene
871 environmental change in Nepal and the northern Indian subcontinent - stable
872 isotopic evidence from paleosols. *Geological Society of America Bulletin*, 107,
873 1381-1397.
- 874 Quade, J., Cerling, T.E., Bowman, J.R., 1989. Development of Asian monsoon
875 revealed by marked ecological shift during the latest Miocene in northern
876 Pakistan. *Nature*, 342, 163-166.
- 877 Ramstein, G., Fluteau, F., Besse, J., Joussaume, S., 1997. Effect of orogeny, plate
878 motion and land sea distribution on Eurasian climate change over the past 30
879 million years. *Nature*, 386, 788-795.
- 880 Raymo, M.E., Ruddiman, W.F., 1992. Tectonic forcing of late Cenozoic climate.
881 *Nature*, 359, 117-122.
- 882 Raymo, M.E., Ruddiman, W.F., Froelich, P.N., 1988. Influence of Late Cenozoic
883 mountain building on ocean geochemical cycles. *Geology*, 16, 649-653.
- 884 Rea, D.K., 1994. The paleoclimatic record provided by eolian deposition in the deep
885 sea - the geologic history of wind. *Reviews of Geophysics*, 32, 159-195.
- 886 Rea, D.K., Snoeckx, H., Joseph, L.H., 1998. Late Cenozoic eolian deposition in the
887 North Pacific: Asian drying, Tibetan uplift, and cooling of the northern
888 hemisphere. *Paleoceanography*, 13, 215-224.
- 889 Richter, F.M., Rowley, D.B., Depaolo, D.J., 1992. Sr isotope evolution of seawater -
890 the role of tectonics. *Earth and Planetary Science Letters*, 109, 11-23.

- 891 Rieser, A.B., Bojar, A.V., Neubauer, F., Genser, J., Liu, Y.J., Ge, X.H., Friedl, G.,
 892 2009. Monitoring Cenozoic climate evolution of northeastern Tibet: stable
 893 isotope constraints from the western Qaidam Basin, China. *International Journal*
 894 *of Earth Sciences*, 98, 1063-1075.
- 895 Robinson, D.M., DeCelles, P.G., Copeland, P., 2006. Tectonic evolution of the
 896 Himalayan thrust belt in western Nepal: Implications for channel flow models.
 897 *Geological Society of America Bulletin*, 118, 865-885.
- 898 Roe, G., 2009. On the interpretation of Chinese loess as a paleoclimate indicator.
 899 *Quaternary Research*, 71, 150-161.
- 900 Rowley, D.B., Currie, B.S., 2006. Palaeo-altimetry of the late Eocene to Miocene
 901 Lunpola basin, central Tibet. *Nature*, 439, 677-681.
- 902 Rowley, D.B., Pierrehumbert, R.T., Currie, B.S., 2001. A new approach to stable
 903 isotope-based paleoaltimetry: implications for paleoaltimetry and
 904 paleohypsometry of the High Himalaya since the Late Miocene. *Earth and*
 905 *Planetary Science Letters*, 188, 253-268.
- 906 Ruddiman, W.F., Kutzbach, J.E., 1989. Forcing of Late Cenozoic Northern
 907 Hemisphere climate by plateau uplift in southern Asia and the American West.
 908 *Journal of Geophysical Research-Atmospheres*, 94, 18409-18427.
- 909 Sanyal, P., Bhattacharya, S.K., Kumar, R., Ghosh, S.K., Sangode, S.J., 2004. Mio-
 910 Pliocene monsoonal record from Himalayan foreland basin (Indian Siwalik) and
 911 its relation to vegetational change. *Palaeogeography Palaeoclimatology*
 912 *Palaeoecology*, 205, 23-41.
- 913 Saylor, J.E., Quade, J., Dellman, D.L., DeCelles, P.G., Kapp, P.A., Ding, L., 2009.
 914 The Late Miocene through present paleoelevation history of southwestern Tibet.
 915 *American Journal of Science*, 309, 1-42.

- 916 Schelling, D., 1992. The tectonostratigraphy and structure of the eastern Nepal
917 Himalaya. *Tectonics*, 11, 925-943.
- 918 Searle, M.P., 2010. Low-angle normal faults in the compressional Himalayan orogen;
919 Evidence from the Annapurna-Dhaulagiri Himalaya, Nepal. *Geosphere*, 6, 296-
920 315.
- 921 Searle, M.P., Godin, L., 2003. The South Tibetan Detachment and the Manaslu
922 Leucogranite: A structural reinterpretation and restoration of the Annapurna-
923 Manaslu Himalaya, Nepal. *Journal of Geology*, 111, 505-523.
- 924 Searle, M.P., Law, R.D., Godin, L., Larson, K.P., Streule, M.J., Cottle, J.M., Jessup,
925 M.J., 2008. Defining the Himalayan Main Central Thrust in Nepal. *Journal of*
926 *the Geological Society*, 165, 523-534.
- 927 Searle, M.P., Waters, D.J., Dransfield, M.W., Stephenson, B.J., Walker, C.B., Walker,
928 J.D., Rex, D.C., 1999. Thermal and mechanical models for the structural and
929 metamorphic evolution of the Zaskar High Himalaya. In: MacNiocaill, C.,
930 Ryan, P. (Eds.), *Continental Tectonics*. Geological Society, London, Special
931 Publication vol. 164, pp. 139-156.
- 932 Seward, D., Burg, J.P., 2008. Growth of the Namche Barwa Syntaxis and associated
933 evolution of the Tsangpo Gorge: Constraints from structural and
934 thermochronological data. *Tectonophysics*, 451, 282-289.
- 935 Skelton, A., 2011. Flux rates for water and carbon during greenschist facies
936 metamorphism. *Geology*, 39, 43–46, doi:10.1130/G31328.1.
- 937 Spicer, R.A., Harris, N.B.W., Widdowson, M., Herman, A.B., Guo, S.X., Valdes, P.J.,
938 Wolfe, J.A., Kelley, S.P., 2003. Constant elevation of southern Tibet over the
939 past 15 million years. *Nature*, 421, 622-624.

- 940 Steinke, S., Groeneveld, J., Johnstone, H., Rendle-Buhring, R., 2010. East Asian
 941 summer monsoon weakening after 7.5 Ma: Evidence from combined planktonic
 942 foraminifera Mg/Ca and delta O-18 (ODP Site 1146; northern South China Sea).
 943 *Palaeogeography Palaeoclimatology Palaeoecology*, 289, 33-43.
- 944 Sun, J.M., Ye, J., Wu, W.Y., Ni, X.J., Bi, S.D., Zhang, Z.Q., Liu, W.M., Meng, J.,
 945 2010. Late Oligocene-Miocene mid-latitude aridification and wind patterns in
 946 the Asian interior. *Geology*, 38, 515-518.
- 947 Sun, X.J., Wang, P.X., 2005. How old is the Asian monsoon system? *Palaeobotanical*
 948 records from China. *Palaeogeography Palaeoclimatology Palaeoecology*, 222,
 949 181-222.
- 950 Tapponnier, P., Xu, Z.Q., Roger, F., Meyer, B., Arnaud, N., Wittlinger, G., Yang,
 951 J.S., 2001. Oblique stepwise rise and growth of the Tibet plateau. *Science*, 294,
 952 1671-1677.
- 953 Treloar, P.J., Rex, D.C., Guise, P.G., Wheeler, J., Hurford, A.J., Carter, A., 2000.
 954 Geochronological constraints on the evolution of the Nanga Parbat syntaxis,
 955 Pakistan Himalaya. In: Khan, M. A., Treloar, P. J., Searle, M. P., Jan, M. Q.
 956 (Eds.), *Tectonics of the Nanga Parbat Syntaxis and the Western Himalaya*.
 957 Geological Society, London Special Publication vol. 170, pp. 137-162.
- 958 Wan, S.M., Li, A.C., Clift, P.D., Stuut, J.B.W., 2007. Development of the East Asian
 959 monsoon: Mineralogical and sedimentologic records in the northern South
 960 China Sea since 20 Ma. *Palaeogeography Palaeoclimatology Palaeoecology*,
 961 254, 561-582.
- 962 Wang, J., Wang, Y.J., Liu, Z.C., Li, J.Q., Xi, P., 1999. Cenozoic environmental
 963 evolution of the Qaidam Basin and its implications for the uplift of the Tibetan

- 964 Plateau and the drying of central Asia,. *Palaeogeography Palaeoclimatology*
 965 *Palaeoecology*, 152, 37-47.
- 966 Wei, G., Li, X.H., Liu, Y., Shao, L., Liang, X.R., 2006. Geochemical record of
 967 chemical weathering and monsoon climate change since the early Miocene in
 968 the South China Sea. *Paleoceanography*, 21, doi: 10.1029/2006pa001300.
- 969 Whipple, K.X., Meade, B.J., 2006. Orogen response to changes in climatic and
 970 tectonic forcing. *Earth and Planetary Science Letters*, 243, 218-228.
- 971 White, N.M., Pringle, M., Garzanti, E., Bickle, M., Najman, Y., Chapman, H., Friend,
 972 P., 2002. Constraints on the exhumation and erosion of the High Himalayan
 973 Slab, NW India, from foreland basin deposits. *Earth and Planetary Science*
 974 *Letters*, 195, 29-44.
- 975 Williams, H., Turner, S., Kelley, S., Harris, N., 2001. Age and composition of dikes
 976 in Southern Tibet: New constraints on the timing of east-west extension and its
 977 relationship to postcollisional volcanism. *Geology*, 29, 339-342.
- 978 Xu, Y., Yue, L.P., Li, J.X., Sun, L., Sun, B., Zhang, J.Y., Ma, J., Wang, J.Q., 2009.
 979 An 11-Ma-old red clay sequence on the Eastern Chinese Loess Plateau.
 980 *Palaeogeography Palaeoclimatology Palaeoecology*, 284, 383-391.
- 981 Yin, A., 2006. Cenozoic tectonic evolution of the Himalayan orogen as constrained by
 982 along-strike variation of structural geometry, exhumation history, and foreland
 983 sedimentation. *Earth-Science Reviews*, 76, 1-131.
- 984 Yin, A., 2010. Cenozoic tectonic evolution of Asia: A preliminary synthesis.
 985 *Tectonophysics*, 488, 293-325.
- 986 Yin, A., Harrison, T.M., 2000. Geologic evolution of the Himalayan-Tibetan orogen.
 987 *Annual Review of Earth and Planetary Sciences*, 28, 211-280.

- 988 Zachos, J., Pagani, M., Sloan, L., Thomas, E., Billups, K., 2001. Trends, rhythms, and
 989 aberrations in global climate 65 Ma to present. *Science*, 292, 686-693.
- 990 Zaleha, M.J., 1997. Siwalik paleosols (Miocene, northern Pakistan): Genesis and
 991 controls on their formation. *Journal of Sedimentary Research*, 67, 821-839.
- 992 Zeitler, P.K., Chamberlain, C.P., Smith, H.A., 1993. Synchronous anatexis,
 993 metamorphism and rapid denudation at Nanga Parbat (Pakistan Himalaya).
 994 *Geology*, 21, 347-350.
- 995 Zhang, Z.S., Wang, H.J., Guo, Z.T., Jiang, D.B., 2007. Impacts of tectonic changes on
 996 the reorganization of the Cenozoic paleoclimatic patterns in China. *Earth and*
 997 *Planetary Science Letters*, 257, 622-634.
- 998 Zheng, H.B., Powell, C.M., Rea, D.K., Wang, J.L., Wang, P.X., 2004. Late Miocene
 999 and mid-Pliocene enhancement of the East Asian monsoon as viewed from the
 1000 land and sea. *Global and Planetary Change*, 41, 147-155.
- 1001

1002 **Figure captions**

1003 Fig. 1. A. Compilation of localities relevant to Miocene climate change, and/or the
 1004 palaeoaltitude history of Tibet. See text for discussion. B. Geographic areas
 1005 mentioned in the text.

1006 Fig. 2. Modern seasonal behaviour of the ITCZ. Modified from Kalnay et al. (1996)
 1007 as represented on
 1008 http://geography.uoregon.edu/envchange/clim_animations/index.html (accessed
 1009 6.9.2010). Global topographic base map from
 1010 <http://www.ngdc.noaa.gov/mgg/topo/globega2.html> (accessed 16.9.2010). Boxes
 1011 show ITCZ study areas of Lyle et al. (2002) and Kim et al (2006).

1012 Fig. 3. Correlation of global isotopic data and Asian tectonics and climatic events, and
 1013 the reconstructed positions of the Himalaya and the ITCZ for the past 35 million
 1014 years. A. Isotopic synthesis for the past 35 million years. Global deep sea oxygen
 1015 isotope records are from Miller et al., (1987; 2005) and Zachos et al. (2001). Seawater
 1016 strontium isotope records from Richter et al. (1992). Vertical lines delimiting ice sheet
 1017 state are from Miller et al. (2005). Abbreviations: MMCO, Mid-Miocene Climate
 1018 Optimum; Mi-1, glaciation during oxygen isotope excursion Mi-1; Oi-1, glaciation
 1019 during oxygen isotope excursion Oi-1; P-P, Pliocene-Pleistocene. B. Asian tectonic
 1020 and climatic events summarised from sources discussed in the text. C. Reconstructed
 1021 trajectories of the ITCZ. Black dots are from ODP Leg 199 (Lyle et al., 2002); open
 1022 squares are from Kim et al. (2006). NW and NE Himalayan trajectories (open circles)
 1023 are re-calculated from Molnar and Stock (2009) by moving their Pakistan and
 1024 Bangladesh (stable Indian plate) reference points 500 km northwards to the Himalaya.
 1025 Palaeogeographic errors fall within the open circle or are expressed as bars. Adapted
 1026 from Armstrong and Allen (2011).

1027 Fig. 4. Conceptual diagrams for ITCZ evolution and the Himalayan/Tibetan system
1028 through the late Cenozoic. Location of the Himalaya/S. Tibet is generalised to a
1029 position of the central part of the range: Present day = 30° N; 8 Ma approximately 25°
1030 N; 20 Ma approximately 20° N.

1031


1032

1033


1034

1035

Figure




A.



B.


Allen & Armstrong Figure 1

Figure




Allen & Armstrong Figure 2

Figure



Allen & Armstrong Figure 3

Figure



Allen & Armstrong Figure 4

Table 1. Evidence for Miocene climate change in South, Central and East Asia.

Reference	Region	Timing (Ma)	Inferred climate change	Evidence
Quade et al (1995)	Southeast Nepal	7-6	Increased seasonality	Changes in soil carbonate $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$
Harrison et al (1993)	Southeast Nepal	7	Increased seasonality	Changes in soil carbonate $\delta^{13}\text{C}$
Flynn and Jacobs (1982)	Northern Pakistan	7	Increased aridity	Vertebrate palaeontology
Barry et al (1985)	Northern Pakistan	7	Increased aridity	Vertebrate palaeontology
Galy et al (2010)	Bengal fan	7.4	Increased aridity?	Change in $\delta^{13}\text{C}$ of organic matter
Dettman et al (2001)	Southeast Nepal	7.5	Increased aridity	$\delta^{18}\text{O}$ trends in shells and mammal teeth
Steinke et al (2010)	South China Sea	7.5	Decreased summer monsoon intensity	Change in seawater $\delta^{18}\text{O}$
Hoorn et al (2001)	Southeast Nepal	8-6.5	Increased aridity?	Vegetation changes
Wan et al (2007)	South China Sea	8	Increased winter monsoon intensity	Chemical weathering proxies; sediment composition
Rea et al (1998)	North Pacific Ocean	8	Increased wind	Dust pulse
Zheng et al (2004)	South China Sea	8	Intensified winter monsoon wind	Foraminifera trends/productivity shifts
Kroon et al (1991)	Arabian Sea	8	Summer monsoon wind	Increase in <i>Globigerina bulloides</i>
Prell et al (1992)	Arabian Sea	8	Summer monsoon wind	Increase in <i>Globigerina bulloides</i>
Quade et al (1989)	Northern Pakistan	8	Increased seasonality	Changes in soil carbonate $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$
Burbank et al (1993)	Bengal fan	8	Unclear – reduced glaciation?	Reduced sediment flux
Qiang et al (2001)	Northern China, loess plateau	8.35	Onset of monsoonal wind system	Onset of loess deposition
Wei et al (2006)	South China Sea	8.4	Increased winter monsoon intensity	Chemical weathering proxies; sediment composition
Fan et al (2007)	NE Tibet	9.6-8.5	Short-lived aridity	Sedimentary $\delta^{18}\text{O}$ excursion
Sanyal et al (2004)	Northern India	10-6	Increased seasonality	Changes in soil carbonate $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$
Clift et al (2008)	Indus fan, Bengal fan, South China Sea	10-8	Weaker summer monsoon	Reduced sediment flux; chemical weathering indices
Barry et al (2002)	Northern Pakistan	10.1-9	Increased seasonality and aridity	Changes in fluvial sedimentology; vertebrate palaeontology
Xu et al (2009)	Northern China, loess plateau	11	Onset of monsoonal wind system	Onset of loess deposition
Jiang and Ding (2010)	Northern China	12	Stronger winter monsoon, weaker summer monsoon	Increase in 10-70 μm grain-size fraction
Dettman et al (2003)	NE Tibet	12	Increased aridity (Tibetan rain shadow?)	Increase in lacustrine carbonate $\delta^{18}\text{O}$
Jiang and Ding (2009)	China	14.25-11.35	Weaker summer monsoon	Pollen assemblage shift
Zaleha (1997)	Northern Pakistan	15-9	Strong seasonality	Variations in paleosols
Wan et al (2007)	South China Sea/SE China	15	Stronger winter monsoon; increased aridity	Chemical weathering proxies; sediment composition
Clift et al (2004)	East China Sea/NE China, South China Sea/SE China, Burma	~15	Stronger monsoon (increased precipitation)	High sediment flux
Wei et al (2006)	South China Sea/SE China	15.7	Stronger winter monsoon; increased aridity	Chemical weathering proxies; sediment composition
Clift and Gaedicke (2002)	Arabian Sea, Indus fan	16-11	Stronger monsoon (increased precipitation)	High sediment flux
Clift et al (2008)	Indus fan, Bengal fan, South China Sea	16-11	Stronger monsoon (increased precipitation)	High sediment flux; chemical weathering proxies
Jiang and Ding (2009)	Northern China, loess plateau	20.13-14.25	Humid climate	Pollen assemblages
Guo et al (2002)	Northern China, loess plateau	22	Onset of monsoonal wind system	Onset of loess deposition
Kent-Corson et al (2009)	Northern Tibet, Qaidam Basin	<~23	Aridification	Increase in sedimentary $\delta^{18}\text{O}$
Sun and Wang (2005)	China	~23	Stronger East Asian monsoon	Shifts in floral types and ranges
Wang et al (1999)	Qaidam Basin, northern China	~23	Shift to more humid climate	Decrease in xerophyte pollen
Clift et al (2004)	Mekong Delta	~23	Increased precipitation	Increased sediment flux
Wei et al (2006)	South China Sea	24-10	Increased precipitation	Increased chemical weathering
Clift et al (2008)	South China Sea	24-10	Increased precipitation	Increased chemical weathering
Sun et al (2010)	Junggar Basin, northern China	24	Aridification; change to modern wind patterns	Onset of aeolian deposition

Palaeolatitudes of the Pacific ITCZ and the Himalayas diverged through the Miocene

Himalayan climate changed to the modern South Asian monsoon in the Late Miocene

Himalayan tectonics may have adjusted to lower precipitation in the Late Miocene