

Anoxia in the terrestrial environment during the Late Mesoproterozoic

Vivien M. Cumming^{1*}, Simon W. Poulton², David Selby¹, Alan D. Rooney³

¹ *Department of Earth Sciences, Durham University, South Road, Durham, DH1 3LE, UK*

² *School of Earth and Environment, University of Leeds, Leeds, LS2 9JT, UK*

³ *Department of Earth and Planetary Sciences, Harvard University, Cambridge, Massachusetts 02138, USA*

**Corresponding Author: v.m.cumming@durham.ac.uk*

ABSTRACT

A significant body of evidence suggests that the marine environment remained largely anoxic throughout most of the Precambrian. In contrast, the oxygenation history of terrestrial aquatic environments has received little attention, despite the significance of such settings for early eukaryote evolution. To address this, we provide here a geochemical and isotopic assessment of sediments from the Late Mesoproterozoic Nonesuch Formation of central North America. We utilize rhenium-osmium (Re-Os) geochronology to yield a depositional age of 1078 ± 24 Ma, while Os isotope compositions support existing evidence for a lacustrine setting. Fe-S-C systematics suggest that the Nonesuch Formation was deposited from an anoxic Fe-rich (ferruginous) water column. Thus, similar to the marine realm, anoxia persisted in terrestrial aquatic environments in the mid-late Proterozoic, but sulfidic water column conditions were not ubiquitous. Our data suggest that oxygenation of the terrestrial realm was not pervasive at this time and may not have preceded oxygenation of the

marine environment, signifying a major requirement for further investigation of links between the oxygenation state of terrestrial aquatic environments and eukaryote evolution.

Keywords: oxygenation; Precambrian; terrestrial environment; Re-Os; Fe speciation

INTRODUCTION

Recent reconstructions suggest that the global ocean remained anoxic between Earth's two major periods of rising atmospheric oxygen at either end of the Proterozoic Eon (2500 – 542 Ma). In detail, sulfidic water column conditions were prevalent along productive continental margins, overlying deeper waters that contained dissolved Fe (ferruginous) (Canfield et al., 2008; Poulton et al., 2010; Poulton and Canfield, 2011; Planavsky et al., 2011). However, in contrast to these advances in our understanding of the evolution of mid-Proterozoic ocean chemistry, much less is known about oxygenation of terrestrial aquatic environments during this period.

Recently, the determination of sulfur isotope fractionations between sulfate and sulfide ($\Delta^{34}\text{S}$) of >50‰ in lacustrine sediments from the Mesoproterozoic Torridon and Stoer Groups of NW Scotland, were interpreted to suggest that oxygenation of terrestrial aquatic environments preceded oxygenation of the oceans (Parnell et al., 2010). Such fractionations are generally considered to require an active oxidative sulfur cycle driven by disproportionation reactions involving sulfide-oxidising bacteria, and are interpreted to reflect a major rise in oxygen in the late Neoproterozoic (Canfield and Teske, 1996). If correct, this suggests that, unlike the marine realm, the late Mesoproterozoic terrestrial environment was sufficiently oxidized to support a complex biota adapted to an oxygen-rich atmosphere (Parnell et al., 2010). However, $\Delta^{34}\text{S}$ fractionations of 60-70‰ have recently been measured in an anoxic, low sulfate lake in the absence of oxidative sulfur cycling (Canfield et al.,

2010). This builds upon theoretical calculations of the magnitude of fractionation possible by bacterial sulfate reduction alone (Brunner and Bernasconi, 2005), and is supported by direct measurements of fractionations obtained during growth of a pure culture of bacterial sulfate reducers (Sim et al., 2011).

Considering the evolutionary significance of possible early oxygenation of terrestrial environments, coupled with these recent developments in our understanding of sulfur isotope fractionation, a more direct assessment of the redox state of the terrestrial realm during the late Mesoproterozoic is clearly warranted. Our focus is on the Nonesuch Formation, deposited within the ~1100 Ma intracratonic Mid-Continent Rift System of central North America (Ojakangas et al., 2001). Like the Torridon Group of NW Scotland, the Nonesuch Formation contains a rich record of eukaryotic life (Pratt et al., 1991; Strother and Wellman, 2010) and thus represents an ideal locality to assess possible links to early terrestrial oxygenation. We utilize Fe-S-C systematics to assess water column redox conditions, coupled with Re-Os geochronology to provide a depositional age for the Nonesuch Formation, and Os isotope systematics to yield insight into the nature of the depositional setting.

GEOLOGICAL SETTING

The Mid-Continent Rift System of central North America was one of the world's largest continental rifts (Ojakangas et al., 2001). In the Lake Superior region, up to 30 km of volcanic and sedimentary rift-fill sequences make up the Keweenawan Supergroup (Ojakangas et al., 2001; Fig. 1). The Oronto Group consists of fluvial and alluvial volcanoclastics, with the exception of the Nonesuch Formation, a 40 to 200 m thick succession of organic-rich siliciclastics. Geochronology of the Keweenawan Supergroup is based upon U-Pb zircon ages of rift-related volcanics, with the overlying post-rift sediments

poorly constrained temporally. An existing U-Pb zircon age of 1087.2 ± 1.6 Ma from the final andesite flow in the Copper Harbour Conglomerate (Davis and Paces, 1990; Fig. 1) provides a maximum age for the Nonesuch Formation.

To allow redox assessment using Fe-S-C systematics we have sampled well-preserved drillcore (PI-1) across a 60 m interval that covers the entire Nonesuch Formation (Pratt et al., 1991). These samples are augmented by Re-Os geochronology and Os isotope analyses of outcrop samples collected ~30 m above the Copper Harbour Conglomerate (Fig. 1; full details of the sampling and analytical protocols are provided in the GSA Data Repository). The Re-Os age for all the outcrop samples is 1040 ± 78 Ma (2σ , $n = 16$, Mean Squared of Weighted Deviation [MSWD] = 10.6; Fig. DR1; Table DR1). The large MSWD and uncertainty on this Re-Os date is due to variation in the initial $^{187}\text{Os}/^{188}\text{Os}$ (Os_i) values rather than disturbance to the isotope system, as suggested by the isochronous nature of the samples and agreement with the underlying U-Pb date (see GSA Data Repository). In fact, the Os_i values form two distinct groups, with eleven of the 16 samples yielding the most precise depositional age of 1078 ± 24 Ma (2σ , $n = 11$, MSWD = 1.05; Fig. DR1).

DEPOSITIONAL ENVIRONMENT

Sedimentological characteristics of the Nonesuch Formation and proximity to continental red-beds (Fig. 1), coupled with paleogeographic reconstructions suggesting that the nearest coastline was ~800 km away, indicating that the Nonesuch Formation was likely deposited in a lacustrine environment (Elmore et al., 1989; Imbus et al., 1992; Ojakangas et al., 2001). However, a marine embayment or estuarine environment has also been suggested based on the presence of specific biomarkers and S/C ratios (Pratt et al., 1991; Hieshima and Pratt, 1991). Biomarkers extracted from the Nonesuch Formation include low levels of 24-*n*-propylcholestane (Pratt et al., 1991), which is commonly, but not uniquely, found in rocks of

marine origin (Moldowan et al., 1990) and may in fact relate to the presence of eukaryotic organisms in the Nonesuch Formation. Thus, the presence of 24-*n*-propylcholestane at low concentration does not unambiguously denote a marine depositional setting for the Nonesuch Formation.

Enrichments in S/C ratios have previously been interpreted to reflect deposition of the Nonesuch Formation under sulfate-rich marine conditions (Fig. DR2; Hieshima and Pratt, 1991; Imbus et al., 1992). However, enhanced fixation of sulfide as a result of the absence of bioturbation in the Precambrian and (potentially) non-Fe limited conditions during ferruginous deposition, combined with differences in the metabolisability of Precambrian organic matter, could readily lead to the observed enrichments in sulfur and decoupling between sulfur and carbon under lacustrine conditions (e.g., Raiswell and Canfield, 2012).

Application of Re-Os geochronology provides an additional, novel constraint on the nature of the depositional environment by yielding the Os_i composition of the water column at the time of deposition, which can be used to distinguish between lacustrine and marine settings (Poirier and Hillaire-Marcel, 2011; Cumming et al., 2012). The Os_i derived from the Re-Os isochrons are 0.81 ± 0.28 and 0.49 ± 0.13 (Fig. DR1), which shows variations reflecting a balance between weathering of radiogenic Archean-Proterozoic cratonic lithosphere (~1.01; upper continental crust calculated at 1100 Ma; Esser and Turekian, 1993) and unradiogenic rift-related flood basalts (~0.13; Allegre et al., 1999). At 1100 Ma, seawater $^{187}\text{Os}/^{188}\text{Os}$ was relatively unradiogenic at ~0.3 (a value derived from two separate margins; Rooney et al., 2010; Azmy et al., 2008), and therefore the more radiogenic values measured for the Nonesuch Formation (0.81 and 0.49) suggest that minimal Os was sourced from the marine realm. Thus, the Os isotope data support sedimentological and paleogeographic evidence for lacustrine depositional conditions.

REDOX CONDITIONS

Water column redox conditions during deposition of the Nonesuch Formation were evaluated using Fe speciation. A biogeochemically 'highly reactive' Fe pool (Fe_{HR}) is analysed through a series of extractions (Poulton and Canfield, 2005). Modern and ancient sediments deposited under anoxic conditions commonly have $\text{Fe}_{\text{HR}}/\text{total Fe}$ (Fe_{T}) ratios >0.38 , in contrast to oxic depositional conditions, where ratios are consistently below 0.22 (Poulton and Canfield, 2011). About 40% of drill core samples (Fig. 2), and all outcrop samples (Table DR2), were clearly deposited from an anoxic water column with $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ >0.38 , while others fall within the equivocal range ($\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}} = 0.22\text{--}0.38$). These equivocal samples may reflect the masking of water column Fe_{HR} enrichments due to rapid sedimentation or transformation of non-sulfidized Fe_{HR} to clay minerals during burial diagenesis or metamorphism (Poulton and Raiswell, 2002; Poulton et al., 2010). The latter possibility can be evaluated by considering Fe/Al ratios (Fig. 2), whereby significant enrichments in Fe relative to both average shale (Lyons and Severmann, 2006) and typical oxic lacustrine sediments (e.g., Kemp and Thomas, 1976; Fagel et al., 2005) throughout the Nonesuch Formation provide strong evidence for anoxic depositional conditions. Post-depositional loss of Fe_{HR} can also be assessed through an extraction that targets Fe associated with clay minerals (termed poorly reactive silicate Fe; Fe_{PRS}) (Poulton et al., 2010). Extreme enrichment in Fe_{PRS} in the Nonesuch Formation ($\text{Fe}_{\text{PRS}}/\text{Fe}_{\text{T}}$ values are well above the modern and Phanerozoic averages; Fig. 2; Poulton and Raiswell, 2002) suggests that significant loss of Fe_{HR} through authigenic clay mineral formation was responsible for reducing original depositional $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ ratios (Poulton et al., 2010), supporting anoxic deposition for all Nonesuch Formation samples.

For samples showing evidence of anoxic deposition, the extent to which the Fe_{HR} pool has been pyritized ($\text{Fe}_{\text{py}}/\text{Fe}_{\text{HR}}$) can then distinguish euxinic ($\text{Fe}_{\text{py}}/\text{Fe}_{\text{HR}} >0.7\text{--}0.8$) from

ferruginous ($\text{Fe}_{\text{py}}/\text{Fe}_{\text{HR}} < 0.7\text{--}0.8$) depositional conditions (Poulton and Canfield, 2011). All of the Nonesuch Formation samples have $\text{Fe}_{\text{py}}/\text{Fe}_{\text{HR}}$ ratios well below the euxinic threshold (Fig. 2), indicating anoxic ferruginous depositional conditions throughout and thus refuting the suggestion of a possible euxinic depositional setting (c.f. Imbus et al., 1992).

IMPLICATIONS FOR TERRESTRIAL BIOSPHERIC OXYGENATION

The upper Nonesuch Formation comprises fining and coarsening upward packages interpreted to be the transition between lacustrine deposition and fluvial conditions of the conformably overlying Freda Sandstone (Elmore et al., 1989). This suggests that the Nonesuch Formation was likely deposited under progressively more oxic conditions as water depth decreased towards the top of the succession (Elmore et al., 1989). Oxygenated surface waters in ~ 1.1 Ga terrestrial aquatic environments would be expected, particularly since shallow marine waters were apparently oxygenated much earlier in Earth history (~ 2.7 Ga; Kendall et al., 2010; Zerkle et al., 2012). However, we find no evidence for oxic deposition in core PI-1, suggesting that anoxia was a persistent feature throughout the depositional period, and any oxygenation was likely restricted to surface waters.

The observation of persistent ferruginous water column conditions suggests that the flux of Fe_{HR} to the lake overwhelmed the flux of sulfate (Poulton and Canfield, 2011). We note here that our Os_i values suggest that any hydrothermal fluid contribution to the lake from rifting or extensive weathering of rift-related basalts would have been negligible, as Os_i values closer to ~ 0.13 (Allegre et al., 1999; Miesel et al., 2001) would otherwise be expected. Hence there is no evidence to support a particularly enhanced influx of Fe due to rifting. Instead, the prevalence of ferruginous water column conditions is consistent with low rates of oxidative pyrite weathering driven by only modest levels of atmospheric oxygen (Canfield and Raiswell, 1999). Although it is not possible to directly determine water column sulfate

concentrations, we note that pyrite sulfur isotope compositions are relatively heavy ($8.5 \pm 6.3\text{‰}$) throughout most of the Nonesuch Formation (Fig. 2), while a compilation of pyrite sulfur isotope data for the Nonesuch Formation shows a similar distribution (possibly skewed towards slightly heavier values) to 1.5 - 1.0 Ga marine sediments (Fig. 3). This is entirely consistent with a relatively low sulfate environment (e.g., Canfield and Raiswell, 1999), and further supports efficient trapping of sulfide (and hence high S/C ratios; Fig. DR2) driven by a lack of bioturbation, and deposition from a ferruginous system which was sulfur (rather than reactive Fe) limited (e.g., Raiswell and Canfield, 2012).

The distribution of the Nonesuch Formation sulfur isotope data is very different to that observed for 1.18-1.0 Ga lacustrine sediments from NW Scotland (Fig. 3), with no indication of fractionations of the magnitude that would indicate significant oxidative sulfur cycling through microbial disproportionation. This is likely a consequence of the restriction of oxic conditions to surface waters, and limitation of sulfate reduction to sediment porewaters beneath the ferruginous water column of the Nonesuch basin. Together, this would limit formation of the oxidized sulfur species required for disproportionation (Thamdrup et al., 1993). In addition, the low sulfate concentrations inferred for the Nonesuch basin would limit the maximum expression of sulfur isotope fractionation even if oxidative sulfur cycling and disproportionation reactions were prevalent (Canfield and Raiswell, 1999). This is consistent with multiple ($^{32/33/34}\text{S}$) sulfur isotope systematics in marine sediments, which suggest that microbial sulfur disproportionation was prevalent by at least 1.3 Ga (Johnston et al., 2005), but was only manifest in the major ($^{32/34}\text{S}$) sulphur isotope record after a later Neoproterozoic rise in atmospheric oxygen led to a significant increase in seawater sulfate concentrations (Canfield and Teske, 1996). Thus, rather than reflecting high sulfate concentrations (Hieshima and Pratt, 1991) potentially attributable to extensive oxygenation of the continents under high atmospheric oxygen, Fe-S-C systematics in the Nonesuch

Formation suggest low sulfate concentrations, ferruginous conditions and only modest atmospheric oxygenation.

In light of this, the sulfur isotope data for the Stoer and Torridon Groups may reflect either an unusually sulfate-rich environment that allowed maximum expression of sulfur isotope fractionations (Parnell et al., 2010), or the preservation of large sulfur isotope fractionations through bacterial sulfate reduction alone (Canfield et al., 2010; Sim et al., 2011). Thus, while it remains possible that sulfur isotope systematics in the Stoer and Torridon Groups may be providing an early record of terrestrial oxygenation; our redox and sulfur isotope data for the Nonesuch Formation suggest that early terrestrial oxygenation was not a pervasive feature of localities that preserve evidence for diverse eukaryotic life. Instead, the redox characteristics display strong similarities to the marine realm (Poulton and Canfield, 2011), implying that in terms of water column and atmospheric oxygenation, terrestrial environmental conditions were likely no more pre-disposed towards eukaryote evolution than shallow marine environments. We suggest that the identification and detailed redox evaluation of other late Mesoproterozoic terrestrial sediments should be a priority in order to more precisely evaluate potential links between the redox evolution of aquatic systems and the rich terrestrial biological record.

ACKNOWLEDGEMENTS

This research was funded by a Lundin Petroleum CeREES PhD scholarship awarded to VMC and the Natural Environment Research Council (Life and the Planet Scheme). Three anonymous reviewers are thanked for helpful comments that have much improved the manuscript.

REFERENCES

225 Allegre, C. J., Birck, J. L., Capmas, F., and Courtillot, V., 1999, Age of the Deccan traps
 226 using ^{187}Re - ^{187}Os systematics: *Earth and Planetary Science Letters*, v. 70, p. 197-204.
 227 Azmy, K., Kendall, B., Creaser, R. A., Heaman, L., and de Oliveira, T. F., 2008, Global
 228 correlation of the Vazante Group, Sao Francisco Basin, Brazil: Re-Os and U-Pb
 229 radiometric age constraints: *Precambrian Research*, v. 164, no. 3-4, p. 160-172.
 230 Brunner, B., and Bernasconi, S. M., 2005, A revised isotope fractionation model for
 231 dissimilatory sulfate reduction in sulfate reducing bacteria: *Geochimica et*
 232 *Cosmochimica Acta*, v. 69, no. 20, p. 4759-4771.
 233 Canfield, D. E., Farquhar, J., and Zerkle, A. L., 2010, High isotope fractionations during
 234 sulfate reduction in a low-sulfate euxinic ocean analog: *Geology*, v. 38, no. 5, p. 415-
 235 418.
 236 Canfield, D. E., Poulton, S. W., Knoll, A. H., Narbonne, G. M., Ross, G., Goldberg, T., and
 237 Strauss, H., 2008, Ferruginous Conditions Dominated Later Neoproterozoic Deep-
 238 Water Chemistry: *Science*, v. 321, no. 5891, p. 949-952.
 239 Canfield, D. E., and Raiswell, R., 1999, The evolution of the sulfur cycle: *American Journal*
 240 *of Science*, v. 299, no. 7-9, p. 697-723.
 241 Canfield, D. E., and Teske, A., 1996, Late Proterozoic rise in atmospheric oxygen
 242 concentration inferred from phylogenetic and sulphur-isotope studies: *Nature*, v. 382,
 243 no. 6587, p. 127-132.
 244 Cumming, V.M., Selby, D., Lillis, P.G., Re-Os geochronology of lacustrine organic-rich
 245 sedimentary rocks: Insights from the Eocene Green River Formation: *Earth and*
 246 *Planetary Science Letters*; in press.
 247 Davis, D. W., and Paces, J. B., 1990, Time resolution of geologic events on the Keweenaw
 248 Peninsula and implications for development of the Midcontinent Rift system: *Earth*
 249 *and Planetary Science Letters*, v. 97, no. 1-2, p. 54-64.

250 Elmore, D. R., Milavec, G. J., Imbus, S. W., and Engel, M. H., 1989, The Precambrian
 251 nonesuch formation of the North American mid-continent rift, sedimentology and
 252 organic geochemical aspects of lacustrine deposition: *Precambrian Research*, v. 43,
 253 no. 3, p. 191-213.

254 Esser, B. K., and Turekian, K. K., 1993, The osmium isotopic composition of the continental-
 255 crust: *Geochimica et Cosmochimica Acta*, v. 57, no. 13, p. 3093-3104.

256 Fagel, N., Alleman, L. Y., Granina, L., Hatert, F., Thamo-Bozso, E., Cloots, R., and André,
 257 L., 2005, Vivianite formation and distribution in Lake Baikal sediments: *Global and*
 258 *Planetary Change*, v. 46, no. 1–4, p. 315-336.

259 Hieshima, G. B., and Pratt, L. M., 1991, Sulfur/carbon ratios and extractable organic matter
 260 of the middle proterozoic nonesuch formation, north american midcontinent rift:
 261 *Precambrian Research*, v. 54, no. 1, p. 65-79.

262 Imbus, S. W., Macko, S. A., Elmore, R. D., and Engel, M. H., 1992, Stable isotope (C, S, N)
 263 and molecular studies on the Precambrian Nonesuch Shale (Wisconsin-Michigan,
 264 USA) – Evidence for differential preservation rates, depositional environment and
 265 hydrothermal influence: *Chemical Geology*, v. 101, no. 3-4, p. 255-281.

266 Johnston, D. T., Wing, B. A., Farquhar, J., Kaufman, A. J., Strauss, H., Lyons, T. W., Kah, L.
 267 C., and Canfield, D. E., 2005, Active Microbial Sulfur Disproportionation in the
 268 Mesoproterozoic: *Science*, v. 310, no. 5753, p. 1477-1479.

269 Kemp, A. L., and Thomas, R. L., 1976, Cultural impact on geochemistry of sediments of
 270 lakes Ontario, Erie and Huron: *Geoscience Canada*, v. 3, no. 3, p. 191-207.

271 Kendall, B., Reinhard, C. T., Lyons, T. W., Kaufman, A. J., Poulton, S. W., and Anbar, A.
 272 D., 2010, Pervasive oxygenation along late Archaean ocean margins: *Nature*
 273 *Geoscience*, v. 3, no. 9, p. 647-652.

274 Lyons, T. W., and Severmann, S., 2006, A critical look at iron paleoredox proxies: New
 275 insights from modern euxinic marine basins: *Geochimica et Cosmochimica Acta*, v.
 276 70, no. 23, p. 5698-5722.

277 Meisel, T., Walker, R. J., Irving, A. J., and Lorand, J.-P., 2001, Osmium isotopic
 278 compositions of mantle xenoliths: a global perspective: *Geochimica et Cosmochimica*
 279 *Acta*, v. 65, no. 8, p. 1311-1323.

280 Moldowan, J. M., Fago, F. J., Lee, C. Y., Jacobson, S. R., Watt, D. S., Slougui, N.-E.,
 281 Jeganathan, A., and Young, D. C., 1990, Sedimentary 12-n-Propylcholestanes,
 282 Molecular Fossils Diagnostic of Marine Algae: *Science*, v. 247, no. 4940, p. 309-312.

283 Ojakangas, R. W., Morey, G. B., and Green, J. C., 2001, The Mesoproterozoic Midcontinent
 284 Rift System, Lake Superior Region, USA: *Sedimentary Geology*, v. 141-142, p. 421-
 285 442.

286 Parnell, J., Boyce, A. J., Mark, D., Bowden, S., and Spinks, S., 2010, Early oxygenation of
 287 the terrestrial environment during the Mesoproterozoic: *Nature*, v. 468, no. 7321, p.
 288 290-293.

289 Planavsky, N. J., McGoldrick, P., Scott, C. T., Li, C., Reinhard, C. T., Kelly, A. E., Chu, X.,
 290 Bekker, A., Love, G. D., and Lyons, T. W., 2011, Widespread iron-rich conditions in
 291 the mid-Proterozoic ocean: *Nature*, v. 477, no. 7365, p. 448-451.

292 Poirier, A., and Hillaire-Marcel, C., 2011, Improved Os-isotope stratigraphy of the Arctic
 293 Ocean: *Geophysical Research Letters*, v. 38.

294 Poulton, S. W., and Canfield, D. E., 2005, Development of a sequential extraction procedure
 295 for iron: implications for iron partitioning in continentally derived particulates:
 296 *Chemical Geology*, v. 214, no. 3-4, p. 209-221.

297 Poulton, S. W., and Canfield, D. E., 2011, Ferruginous Conditions: A Dominant Feature of
 298 the Ocean through Earth's History: *Elements*, v. 7, no. 2, p. 107-112.

299 Poulton, S. W., Fralick, P. W., and Canfield, D. E., 2010, Spatial variability in oceanic redox
 300 structure 1.8 billion years ago: *Nature Geosci*, v. 3, no. 7, p. 486-490.

301 Poulton, S. W., and Raiswell, R., 2002, The low-temperature geochemical cycle of iron:
 302 From continental fluxes to marine sediment deposition: *American Journal of Science*,
 303 v. 302, no. 9, p. 774-805.

304 Pratt, L. M., Summons, R. E., and Hieshima, G. B., 1991, Sterane and triterpane biomarkers
 305 in the Precambrian Nonesuch Formation, North American Midcontinent Rift:
 306 *Geochimica et Cosmochimica Acta*, v. 55, no. 3, p. 911-916.

307 Raiswell, R., and Canfield, D. E., 2012, The Iron Biogeochemical Cycle Past and Present:
 308 *Geochemical Perspectives*, v. 1, no. 1.

309 Rooney, A. D., Selby, D., Houzay, J.-P., and Renne, P. R., 2010, Re-Os geochronology of a
 310 Mesoproterozoic sedimentary succession, Taoudeni basin, Mauritania: Implications
 311 for basin-wide correlations and Re-Os organic-rich sediments systematics: *Earth and*
 312 *Planetary Science Letters*, v. 289, no. 3-4, p. 486-496.

313 Sim, M. S., Bosak, T., and Ono, S., 2011, Large Sulfur Isotope Fractionation Does Not
 314 Require Disproportionation: *Science*, v. 333, no. 6038, p. 74-77.

315 Strother, P. K., and Wellman, C. H., 2010, Paleobiology of a Nonmarine Precambrian Shale:
 316 Well-Preserved Eukaryotes from the 1.1 Ga Nonesuch Formation, in *Proceedings The*
 317 *Palaeontological Association 54th Annual Meeting*, Ghent University.

318 Thamdrup, B., Finster, K., Hansen, J. W., and Bak, F., 1993, Bacterial Disproportionation of
 319 Elemental Sulfur Coupled to Chemical Reduction of Iron or Manganese: *Applied and*
 320 *Environmental Microbiology*, v. 59, no. 1, p. 101-108.

321 Zerkle, A. L., Claire, M. W., Domagal-Goldman, S. D., Farquhar, J., and Poulton, S. W.,
 322 2012, A bistable organic-rich atmosphere on the Neoarchaeon Earth: *Nature*
 323 *Geoscience*, v. 5, no. 5, p. 359-363.

FIGURE CAPTIONS

Figure 1: A. Geological map of the Lake Superior region showing the location of the main Keweenawan Supergroup units. The Nonesuch Formation is part of the Oronto Group situated between continental red-beds of the Copper Harbour Conglomerate and the Freda Sandstone and outcrops on the southern shores of Lake Superior. B. Schematic stratigraphy of the Keweenawan Supergroup, focusing on post-rift sediments. The right hand panel illustrates depth variations for each unit. Locations of the Re-Os (1078 ± 24 Ma; this study) and U-Pb (1087.2 ± 1.6 Ma; Davis and Paces, 1990) ages are labelled. Figures adapted from Elmore et al. (1989) and Ojakangas et al. (2001).

Figure 2: Geochemical profiles for the Nonesuch Formation core samples. For the $\text{Fe}_{\text{HR}}/\text{Fe}_{\text{T}}$ and $\text{Fe}_{\text{Py}}/\text{Fe}_{\text{HR}}$ graphs the dashed lines show the fields for oxic or anoxic deposition and ferruginous or euxinic deposition, respectively (Poulton and Canfield, 2011). On the $\text{Fe}_{\text{T}}/\text{Al}$ graph the dashed line represents average anoxic shale (Lyons and Severmann, 2006). On the $\text{Fe}_{\text{PRS}}/\text{Fe}_{\text{T}}$ graph the dashed lines represent modern (Raiswell and Canfield, 1998) and Phanerozoic (Poulton and Raiswell, 2002) averages. CHC stands for Copper Harbour Conglomerate.


Figure 3: Probability density plot of pyrite sulfur isotope data for the Nonesuch Formation (this study; Imbus et al., 1992), the Torridon and Stoer Groups (Parnell et al., 2010) and 1.5 – 1.0 Ga marine sediments (Canfield and Raiswell, 1999).

¹GSA Data Repository item 2009xxx, which includes a detailed sampling and methodology section, a detailed Re-Os geochronology results section and results tables, is available online

348 at www.geosociety.org/pubs/ft2009.htm, or on request from editing@geosociety.org or
349 Documents Secretary, GSA, P.O. Box 9140, Boulder, CO 80301, USA.

350

351 **FIGURE 1**




352

353

354 **FIGURE 2**

355



356

357

358 **FIGURE 3**



359

360

361

362