

Rank-driven Markov processes

Michael Grinfeld Philip A. Knight Andrew R. Wade
University of Strathclyde

18th September 2011

Abstract

We study a class of Markovian systems of N elements taking values in $[0, 1]$ that evolve in discrete time t via randomized replacement rules based on the ranks of the elements. These rank-driven processes are inspired by variants of the Bak–Sneppen model of evolution, in which the system represents an evolutionary ‘fitness landscape’ and which is famous as a simple model displaying self-organized criticality. Our main results are concerned with long-time large- N asymptotics for the general model in which, at each time step, K randomly chosen elements are discarded and replaced by independent $U[0, 1]$ variables, where the ranks of the elements to be replaced are chosen, independently at each time step, according to a distribution κ_N on $\{1, 2, \dots, N\}^K$. Our main results are that, under appropriate conditions on κ_N , the system exhibits threshold behaviour at $s^* \in [0, 1]$, where s^* is a function of κ_N , and the marginal distribution of a randomly selected element converges to $U[s^*, 1]$ as $t \rightarrow \infty$ and $N \rightarrow \infty$. Of this class of models, results in the literature have previously been given for special cases only, namely the ‘mean-field’ or ‘random neighbour’ Bak–Sneppen model. Our proofs avoid the heuristic arguments of some of the previous work and use Foster–Lyapunov ideas. Our results extend existing results and establish their natural, more general context. We derive some more specialized results for the particular case where $K = 2$. One of our technical tools is a result on convergence of stationary distributions for families of uniformly ergodic Markov chains on increasing state-spaces, which may be of independent interest.

Keywords: Bak–Sneppen evolution model; self-organized criticality; Markov process on order statistics; phase transition; interacting particle system.

AMS 2010 Subject Classifications: 60J05 (Primary) 60J10, 60K35, 82B26, 92D15 (Secondary)

1 Introduction

Bak and Sneppen [5] introduced a simple stochastic model of evolution which initiated a considerable body of research by physicists and mathematicians. The Bak–Sneppen model has proved so influential because it is simple to describe and not difficult to simulate, and, while being challenging to analyse rigorously, demonstrates highly non-trivial behaviour: it is said to exhibit ‘self-organized criticality’ (see e.g. [20]).

The Bak–Sneppen model is as follows. Consider the sites $1, 2, \dots, N$ arranged cyclically, so that site k has neighbours $k - 1$ and $k + 1$ (working modulo N). Each site,

corresponding to a species in the evolution model, is initially assigned an independent $U[0, 1]$ random variable representing a ‘fitness’ value for the species; here and subsequently $U[a, b]$ stands for the uniform distribution on the interval $[a, b]$. The Bak–Sneppen model is a discrete-time Markov process, where at each step the minimal fitness value and the values at the two neighbouring sites are replaced by three independent $U[0, 1]$ random variables. A variation on the model is the (maximal) *anisotropic* Bak–Sneppen model [19] in which, at each step, only the *right* neighbour of the site with minimal fitness is updated along with the minimal value.

A large physics literature is devoted to these models. Simulations suggest that the equilibrium distribution of the fitness at any particular site approaches $U[s^*, 1]$ in the $N \rightarrow \infty$ limit, for some threshold value s^* ; simulations give $s^* \approx 0.667$ for the original Bak–Sneppen model and $s^* \approx 0.724$ for the anisotropic model [13]. There is a much smaller number of mathematical papers on the model and its variants: see e.g. [15, 16, 23, 24]; see also the thesis [14]. It is a challenge to obtain further rigorous results for such models.

A simpler model can be formulated by removing the underlying topology, and such ‘mean field’ or ‘random neighbour’ versions of the model have also received attention in the literature: see e.g. [7, 8, 12, 20, 21, 29]. For example, the mean-field version of the anisotropic Bak–Sneppen model again has N sites each endowed with a fitness in $[0, 1]$. At each step, the minimal fitness is replaced, along with one other fitness chosen *uniformly at random* from the remaining $N - 1$ sites. Again the replacement fitnesses are independent $U[0, 1]$ variables.

Such mean-field models display some features qualitatively similar to the original Bak–Sneppen model, but give little indication of how the distinctive asymptotics of the Bak–Sneppen model, in which the topology plays a key role, might arise. In particular, changing the topology of the model changes the value of the threshold s^* in a way that the mean-field models cannot account for. In the present paper we study some generalizations of the mean-field model described informally above, which we call *rank-driven processes*. These models represent one possibility for showing how the influence of topology might be replicated by simpler features.

In these more general models, we again have N sites, and at each time step some fixed number of fitness values are selected for replacement, but for this selection process we allow general stochastic rules based on the *ranks* of the values. These rank-driven processes are Markov processes on *ranked* sequences, or *order statistics* (see Sections 2 and 3 for formal definitions).

To give a concrete example, we could, at each step, replace the minimal fitness along with the R th ranked value, where R is chosen independently at each step from some distribution on $\{2, \dots, N\}$, with $R = 2$ corresponding to the second smallest value, and so on. This model generalizes that of [7], which has R *uniform* on $\{2, \dots, N\}$, and exhibits much richer behaviour. Specifically, the threshold s^* depends explicitly on the distribution chosen for R : in this way, the distribution of R is playing a role analogous to the underlying topology in the Bak–Sneppen models.

Such rank-driven processes are of interest in their own right, but our motivation for studying them also arises from an attempt to understand the original Bak–Sneppen model, where the topology plays a key role. While the Bak–Sneppen model can be viewed as a Markov process on the space $[0, 1]^N$, it gives rise to a decidedly non-Markovian process on order statistics. At the same time, as we explore in detail in [17], there is an algorithmic way to associate to the Bak–Sneppen model a rank-driven process (a process that *is* Markovian on order statistics) and which, according to numerical evidence, shares

a number of important properties with the Bak–Sneppen model. The aim of the present paper is to present a rigorous analysis of rank-driven processes. Ultimately we hope to show that if a process, such as the BS one, gives rise to a rank-driven process, the two processes must share observables, such as the location of thresholds in the invariant distribution; see [17] for supporting numerical evidence.

The outline of the remainder of the paper is as follows. In Section 2 we discuss an introductory example in which a single value is updated at each step. In Section 3 we describe the general rank-driven processes that we consider and state our main theorems on asymptotic behaviour. In Section 4 we focus on a specific class of examples, generalizing the mean-field anisotropic Bak–Sneppen model [7, 12], and give some more detailed results. We emphasize the difference in nature of the results in Sections 3 and 4: in the former, the results cover a very general class of processes and the proofs are robust, using Foster–Lyapunov arguments and general theory of Markov processes, while in the latter, we specialize to a narrower class of processes and exploit their special structure. It is likely that analogues of our results from Section 4 could be obtained for other processes, but the details would depend on the particular processes studied. In Section 5 we make some further remarks and state some open problems. Section 6 is devoted to the proofs of the main results in Section 3, while Section 7 is devoted to the proofs of the results in Section 4. The Appendix, Section 8, gives one of our key technical tools on the asymptotics of families of Markov chains that are uniformly ergodic in a precise sense.

2 Warm-up example: Replace the k th-ranked value

We start by describing a particularly simple model, which can be solved completely, to demonstrate some ideas that will be useful in greater generality later on. It will be convenient to view all of our models as Markov chains on *ranked* sequences, or *order statistics*. Fix $N \in \mathbb{N} := \{1, 2, \dots\}$. Given a vector (x^1, x^2, \dots, x^N) we write the corresponding increasing order statistics as

$$(x^{(1)}, x^{(2)}, \dots, x^{(N)}) = \text{ord}(x^1, \dots, x^N),$$

where $x^{(1)} \leq x^{(2)} \leq \dots \leq x^{(N)}$. Let Δ_N denote the ‘simplex’

$$\Delta_N := \{(x^1, \dots, x^N) \in [0, 1]^N : x^1 \leq x^2 \leq \dots \leq x^N\}.$$

We study stochastic processes on Δ_N indexed by discrete time $\mathbb{Z}^+ := \{0, 1, 2, \dots\}$.

Let U_1, U_2, \dots denote a sequence of independent $U[0, 1]$ random variables. Define a Markov process X_t on Δ_N with transition rule such that, given X_t ,

$$X_{t+1} = \text{ord}\{U_{t+1}, X_t^{(2)}, X_t^{(3)}, \dots, X_t^{(N)}\};$$

in other words, at each step, discard the *smallest* value and replace it by a $U[0, 1]$ random variable. To make clear the dependence on the model parameter N , we write \mathbb{P}_N for the probability measure associated with this model and \mathbb{E}_N for the corresponding expectation.

It is natural to anticipate that X_t should approach (as $t \rightarrow \infty$) a limiting (stationary) distribution; we show in this section that this is indeed the case. Assuming such a stationary distribution exists, and is unique, we can guess what it must be: the distribution of the random vector $(U, 1, 1, 1, \dots, 1)$ (a $U[0, 1]$ variable followed by $N-1$ units) is invariant under the evolution of the Markov chain. The process X_t itself lives on a relatively

complicated state-space, and at first sight it might seem that some fairly sophisticated argument would be needed to show that it has a unique stationary distribution. In fact, we can reduce the problem to a simpler problem on a finite state-space as follows.

For each $s \in [0, 1]$, define the *counting function*

$$C_t^N(s) := \#\{i \in \{1, 2, \dots, N\} : X_t^{(i)} \leq s\} = \sum_{i=1}^N \mathbf{1}\{X_t^{(i)} \leq s\}, \quad (2.1)$$

i.e., $C_t^N(s)$ is the number of values of magnitude at most s in the system at time t . (Here and throughout we use $\#A$ to denote the number of elements of a finite set A .) Then, for a fixed t , X_t is characterized by the counting functions $(C_t^N(s))_{s \in [0, 1]}$. For a specific s , $C_t^N(s)$ encodes marginal information about the $X_t^{(k)}$, since the two events $\{C_t^N(s) \geq k\}$ and $\{X_t^{(k)} \leq s\}$ are equivalent. By an analysis of the auxiliary stochastic processes $C_t^N(s)$ we will prove the following result, which deals with the more general model in which, at each time step, the point with rank k is replaced.

Proposition 2.1. *Let $N \in \mathbb{N}$ and $k \in \{1, 2, \dots, N\}$. For the model in which at each step we replace the k th-ranked value by an independent $U[0, 1]$ value, we have that, as $t \rightarrow \infty$,*

$$(X_t^{(1)}, X_t^{(2)}, \dots, X_t^{(N)}) \xrightarrow{d} (0, \dots, 0, U, 1, \dots, 1),$$

where U , the k th coordinate of the limit vector, has a $U[0, 1]$ distribution.

If $k = k(N)$ is such that $k(N)/N \rightarrow \theta \in [0, 1]$ as $N \rightarrow \infty$, a consequence of Proposition 2.1 is that the distribution of a uniformly chosen point converges (as $t \rightarrow \infty$ and then $N \rightarrow \infty$) to the distribution with an atom of mass θ at 0 and an atom of mass $1 - \theta$ at 1. For example, if we always replace a *median* value, $\theta = 1/2$ and the limit distribution has two atoms of mass $1/2$ at 0 and 1.

Proof of Proposition 2.1. It is not hard to see that $C_t^N(s)$ is a Markov chain on $\{0, 1, 2, \dots, N\}$. The transition probabilities $p_N^s(n, m) := \mathbb{P}_N[C_{t+1}^N(s) = m \mid C_t^N(s) = n]$ are given for $n \in \{0, \dots, k-1\}$ by $p_N^s(n, n) = 1 - s$ and $p_N^s(n, n+1) = s$, and for $n \in \{k, \dots, N\}$ by $p_N^s(n, n) = s$ and $p_N^s(n, n-1) = 1 - s$. For $s \in (0, 1)$ the Markov chain is reducible and has a single recurrent class consisting of the states $k-1$ and k . It is easy to compute the stationary distribution and for $s \in (0, 1)$ we obtain,

$$\lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) = n] = \begin{cases} 1 - s & \text{if } n = k - 1 \\ s & \text{if } n = k \\ 0 & \text{if } n \notin \{k - 1, k\} \end{cases},$$

by standard Markov chain theory. In particular, for $s \in (0, 1)$,

$$\lim_{t \rightarrow \infty} \mathbb{P}_N[X_t^{(k)} \leq s] = \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \geq k] = s.$$

That is, $X_t^{(k)}$ converges in distribution to a $U[0, 1]$ variable. Moreover, if $k > 1$, for any $s \in (0, 1)$, $\mathbb{P}[X_t^{(k-1)} \leq s] = \mathbb{P}[C_t^N(s) \geq k-1] \rightarrow 1$, which implies that $X_t^{(k-1)}$ converges in probability to zero. Similarly, if $k < N$, for any $s \in (0, 1)$, $\mathbb{P}[X_t^{(k+1)} \leq s] = \mathbb{P}[C_t^N(s) \geq$

$k+1] \rightarrow 0$, which implies that $X_t^{(k+1)}$ converges in probability to 1. Thus we have proved the marginal result that, as $t \rightarrow \infty$, for U a $U[0, 1]$ random variable,

$$X_t^{(i)} \rightarrow 0, \quad (i < k), \quad X_t^{(k)} \rightarrow U, \quad X_t^{(i)} \rightarrow 1, \quad (i > k),$$

in distribution. Then the Cramér–Wold device (convergence in distribution of an N -dimensional random vector is implied by convergence in distribution of all linear combinations of its components: see e.g. [10, p. 147]) together with Slutsky’s theorem (if Y_n converges in distribution to a random limit Y and Z_n converges in distribution to a deterministic limit z , then $Y_n + Z_n$ converges in distribution to $Y + z$: see e.g. [10, p. 72]) enable us to deduce the joint convergence. \square

Remark 2.1. *This simple example shows special features that will not recur in the general case. (i) Here we obtained a result for any fixed $N \geq 1$; in the general case, we will typically state results as $N \rightarrow \infty$. (ii) Since all but one of the $X_t^{(i)}$ had a degenerate limit distribution, we were able to use a soft argument to deduce convergence of the joint distribution of $(X_t^{(1)}, \dots, X_t^{(N)})$ from the convergence of the marginal distributions.*

3 Rank-driven processes and threshold behaviour

In this section we give a general definition of a *rank-driven process* and present some fundamental results on its asymptotic properties. Fix N (the number of points) and $K \in \{1, \dots, N\}$ (the number of replacements at each step). Define the set

$$\mathcal{I}_N^K := \{1, 2, \dots, N\}^K.$$

The model will be specified by a *selection distribution*. Let R^N denote a random K -vector with distinct components in $\{1, \dots, N\}$. In components, write $R^N = (R_1^N, \dots, R_K^N)$. We suppose that R^N is *exchangeable*, i.e., its distribution is invariant under permutations of its components. The distribution of R^N can be described by a probability mass function $\kappa_N : \mathcal{I}_N^K \rightarrow [0, 1]$ that is symmetric under permutations of its arguments, so $\mathbb{P}_N[R_1^N = i_1, \dots, R_K^N = i_K] = \kappa_N(i_1, \dots, i_K)$.

We define a Markov chain $(X_t)_{t \in \mathbb{Z}^+}$ of ranked sequences $X_t = (X_t^{(1)}, \dots, X_t^{(N)})$. The initial distribution X_0 can be arbitrary. The randomness of the process will be introduced via independent $U[0, 1]$ random variables U_1, U_2, \dots and independent copies of R^N , which we denote by $R^N(1), R^N(2), \dots$. In components, write $R^N(t) = (R_1^N(t), \dots, R_K^N(t))$. The transition law of the Markov chain is as follows.

Given X_t , discard the elements of (distinct) ranks specified by $R_1^N(t+1), \dots, R_K^N(t+1)$ and replace them by K new independent $U[0, 1]$ random variables, namely $U_{Kt+1}, \dots, U_{Kt+K}$; then rank the new sequence. That is, we take X_{t+1} to be

$$\text{ord} \left(X_t^{(1)}, X_t^{(2)}, \dots, X_t^{(R_1^N(t+1)-1)}, U_{Kt+1}, X_t^{(R_2^N(t+1)+1)}, \dots, U_{Kt+K}, X_t^{(R_K^N(t+1)+1)}, \dots, X_t^{(N)} \right).$$

Note that ties are permitted.

For $i \in \mathbb{N}$ let $g_N(i) := \mathbb{P}_N[R_1^N = i]$, the marginal distribution of a specific component of R^N . Denote the corresponding distribution function by

$$G_N(n) := \mathbb{P}_N[R_1^N \leq n] = \sum_{i=1}^n g_N(i) = \sum_{i_1=1}^n \sum_{i_2=1}^N \cdots \sum_{i_K=1}^N \kappa_N(i_1, i_2, \dots, i_K). \quad (3.1)$$

We make some further assumptions on the selection distribution. Assumption (A1) will ensure that an irreducibility property holds, excluding some degenerate cases, while (A2) regulates the $N \rightarrow \infty$ behaviour of the selection rule.

(A1) If $K = 1$, suppose that $g_N(i) > 0$ for all $i \in \{1, \dots, N\}$. If $K \geq 2$, suppose that $g_N(1) > 0$.

(A2) Suppose that for any $k \leq K$, for all distinct $i_1, \dots, i_k \in \mathbb{N}$, the limit $\kappa(i_1, \dots, i_k) := \lim_{N \rightarrow \infty} \mathbb{P}_N[R_1^N = i_1, \dots, R_k^N = i_k]$ exists.

Note that the limits in (A2) need not constitute proper distributions on \mathbb{N}^k : there may be some loss of mass. Indeed, the possibility of a defective distribution as a limit in (A2) plays a central role in the asymptotics of the rank-driven process, as we shall describe below. In the example of replacing the minimum element and one uniformly random other element, $\mathbb{P}[R_1^N = i, R_2^N = j] = 0$ if neither i nor j is 1, and this probability is $\frac{1}{2(N-1)}$ otherwise (see Example (E2) below). Hence (A2) holds and for all $i, j \in \mathbb{N}$, $\kappa(i, j) = 0$, so that κ is (maximally) defective. On the other hand, if we always choose the smallest and the second smallest elements, $\kappa_N(1, 2) = \kappa_N(2, 1) = 1/2$, and the limits in (A2) are indeed probability distributions. In general, a proper distribution can be recovered on $(\mathbb{N} \cup \{\infty\})^K$ by correctly accounting for the lost mass, and then (A2) can be interpreted as saying that R^N converges in distribution to a random vector on $(\mathbb{N} \cup \{\infty\})^K$: see Section 6.3 for details. A consequence of (A2) is that

$$\lim_{N \rightarrow \infty} g_N(n) = g(n) \text{ and } \lim_{N \rightarrow \infty} G_N(n) = G(n) \quad (3.2)$$

exist for all $n \in \mathbb{N}$; then G is a (possibly defective) distribution function on \mathbb{N} . (Note that $g(i) = \kappa(i)$.) Given (A2), we make an assumption on g analogous to (A1):

(A3) If $K = 1$, suppose that $g(i) > 0$ for all $i \in \mathbb{N}$. If $K \geq 2$, suppose that $g(1) > 0$.

We will show that a crucial parameter for the asymptotics of the process is

$$s^* := \lim_{n \rightarrow \infty} G(n) = \lim_{n \rightarrow \infty} \lim_{N \rightarrow \infty} G_N(n). \quad (3.3)$$

If (A2) holds, then the N -limit exists, and $s^* \in [0, 1]$ is well-defined. The value of s^* captures the ‘asymptotic atomicity’ of G_N in a certain sense.

Before stating our first results, we describe some concrete examples. For specifying some of the examples, it is more convenient to work with a version of κ_N on ranked sequences, namely γ_N defined for $i_1 < \dots < i_K$ by $\gamma_N(i_1, \dots, i_K) = K! \kappa_N(i_1, \dots, i_K)$. With this notation, note that

$$g_N(i) = \frac{1}{K} \left(\sum_{i < i_2 < i_3 < \dots < i_K} \gamma_N(i, i_2, i_3, \dots, i_K) + \sum_{i_2 < i < i_3 < \dots < i_K} \gamma_N(i_2, i, i_3, \dots, i_K) \right. \\ \left. + \dots + \sum_{i_2 < i_3 < \dots < i_K < i} \gamma_N(i_2, i_3, \dots, i_K, i) \right), \quad (3.4)$$

where each sum is over $i_2, i_3, \dots, i_K \in \{1, \dots, N\}$ satisfying the given rank constraints.

We describe three examples, by giving the non-zero values of either γ_N or κ_N , as convenient; (E1) was discussed in Section 2, while we study examples (E2) and (E3) in detail in Section 4.

Example (E1). Take $K = 1$ and $\gamma_N(k) = 1$, i.e., replace the k th ranked element only each time. In this case $g_N(k) = 1$ as well.

Example (E2). Let $K = 2$ and $\gamma_N(1, j) = \frac{1}{N-1}$ for $j \in \{2, \dots, N\}$, i.e., each time we replace the minimal element and one other uniformly chosen point. This model has been studied by [7] and others. From (3.4) we have that in this case $g_N(1) = 1/2$ and, for $i \in \{2, \dots, N\}$, $g_N(i) = \frac{1}{2(N-1)}$. Moreover, $g(1) = 1/2$ and $g(i) = 0$ for $i \neq 1$.

Example (E3). (A generalization of (E2).) Let $K \geq 2$ and let ϕ_N be a symmetric probability mass function on \mathcal{I}_N^{K-1} . Set $\kappa_N(1, i_2, \dots, i_K) = K^{-1}\phi_N(i_2, \dots, i_K)$. So now we replace the minimal element and $K - 1$ other randomly chosen points, where the distribution on the ‘other’ points is given by ϕ_N . Then $g_N(1) = 1/K$ and, for $i \in \{2, \dots, N\}$, $g_N(i) = \frac{K-1}{K}f_N(i)$ where $f_N(i) = \sum_{i_3, \dots, i_K} \phi_N(i, i_3, \dots, i_K)$. Write $F_N(n) = \sum_{i=1}^n f_N(i)$. Assume that $F(n) = \lim_{N \rightarrow \infty} F_N(n)$ exists for all n , and set $\alpha = \lim_{n \rightarrow \infty} F(n) \in [0, 1]$.

The assumptions (A1) and (A3) are satisfied by (E2) and (E3), but not (E1), while (A2) is satisfied by (E1), (E2), and (E3).

We will work with the counting functions defined by (2.1). Our first main result, Theorem 3.1 below, demonstrates a phase transition in the asymptotic behaviour of the system at the threshold value $s = s^*$; note that part of the theorem is the non-trivial statement that $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)]$ exists in $[0, \infty]$.

Theorem 3.1. *Suppose that (A1), (A2), and (A3) hold, so that s^* given by (3.3) exists in $[0, 1]$. Then*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] \begin{cases} < \infty & \text{if } s < s^* \\ = \infty & \text{if } s > s^* \end{cases}. \quad (3.5)$$

Theorem 3.1 shows that s^* is a threshold value for the model in the sense that

$$s^* = \sup\{s \geq 0 : \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] < \infty\}$$

is well defined (with the convention $\sup \emptyset = 0$). Example (E1) has $s^* = 1$, although Theorem 3.1 does not apply directly (since (A1) and (A3) fail). Example (E2) has $s^* = 1/2$, while Example (E3) has $s^* = \frac{1}{K}(1 + (K-1)\alpha)$.

For the next result we assume that the distribution G_N given by (3.1) is ‘eventually uniform’ in a certain sense; roughly speaking we will suppose that $g_N(n) \approx \frac{1-s^*}{N}$ for n large enough. The precise condition that we will use is as follows.

(A4) Suppose that there exists $n_0 \in \{2, 3, \dots\}$ such that

$$\lim_{N \rightarrow \infty} \sup_{n_0 \leq n \leq N} \left| \frac{N(G_N(n) - s^*)}{n - n_0 + 1} - (1 - s^*) \right| = 0.$$

For instance, Example (E2) satisfies condition (A4) with $s^* = 1/2$ and $n_0 = 2$, since $G_N(n) = \frac{1}{2} + \frac{n-1}{2(N-1)}$. In Example (E3), $G_N(n) = \frac{1}{K} + \frac{K-1}{K}F_N(n)$, so that condition (A4) holds if $F_N(n)$ satisfies a similar condition, namely

$$\lim_{N \rightarrow \infty} \sup_{n_0 \leq n \leq N} \left| \frac{N(F_N(n) - \alpha)}{n - n_0 + 1} - (1 - \alpha) \right| = 0. \quad (3.6)$$

Our next result shows the threshold phenomenon at the ‘ $O(N)$ ’ scale. We can define a threshold parameter

$$s^\# := \sup\{s \geq 0 : \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} (N^{-1} \mathbb{E}_N[C_t^N(s)]) = 0\}. \quad (3.7)$$

If both s^* and $s^\#$, as given by (3.3) and (3.7) respectively, are well-defined, then clearly $s^* \leq s^\#$. Theorem 3.2 shows that, under assumption (A4), $s^* = s^\#$; in other words, the transition is sharp. We use the notation

$$V(s) := \begin{cases} 0 & \text{if } s < s^* \\ \frac{s-s^*}{1-s^*} & \text{if } s \geq s^* \end{cases}. \quad (3.8)$$

Theorem 3.2. *Suppose that (A1), (A2), (A3), and (A4) hold. With $V(s)$ as given by (3.8), we have that for any $s \in [0, 1]$,*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \left(\frac{\mathbb{E}_N[C_t^N(s)]}{N} \right) = V(s). \quad (3.9)$$

Another way to interpret Theorem 3.2 is as follows. Let X_t^* denote $X_t^{(M)}$ where M is a random variable with $\mathbb{P}_N[M = j] = 1/N$ for $j \in \{1, \dots, N\}$. Then

$$\mathbb{P}_N[X_t^* \leq s] = N^{-1} \sum_{i=1}^N \mathbb{P}[X_t^{(i)} \leq s] = N^{-1} \mathbb{E} \sum_{i=1}^N \mathbf{1}\{X_t^{(i)} \leq s\} = N^{-1} \mathbb{E}_N[C_t^N(s)],$$

by (2.1), so that the conclusion of Theorem 3.2 is equivalent to, for $s \in [0, 1]$,

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[X_t^* \leq s] = V(s);$$

in other words, the marginal distribution of a ‘typical’ point converges (as $t \rightarrow \infty$ then $N \rightarrow \infty$) to a $U[s^*, 1]$ distribution. Note that some condition along the lines of (A4) is needed for this result to hold: see the example in Remark 3.2 below.

Remark 3.1. *In this paper we restrict attention to the case where the distribution of replacements is $U[0, 1]$, but instead of U_1, U_2, \dots one could take independent copies of a nonnegative random variable W with distribution function ρ . The results with the $U[0, 1]$ distribution immediately generalize to distributions ρ that are continuous, supported on a single interval, and strictly increasing on that interval: to see this, note that $(x_1, \dots, x_N) \mapsto (\rho(x_1), \dots, \rho(x_N))$ preserves ranks and $\rho(W)$ has a $U[0, 1]$ distribution, so that the dynamics of the process are preserved, up to the change of scale $s \mapsto \rho(s)$. Thus our results immediately extend to this class of distributions W .*

Remark 3.2. *Our results can be translated into complementary results by reversing the ranking and looking at $N - C_t^N(s)$. Indeed, $N - C_t^N(s)$ counts the number of points in $(s, 1]$; translating Theorem 3.1 shows that under appropriate versions of (A1)–(A3) (in which conditions on $g_N(n)$ are replaced by conditions on $g_N(N - n + 1)$) the threshold*

$$s_* = \inf\{s \leq 1 : \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[N - C_t^N(s)] < \infty\}$$

is given by

$$s_* = \lim_{n \rightarrow \infty} \lim_{N \rightarrow \infty} G_N(N - n).$$

For example, suppose that $K = 2$ and we always replace the smallest and the largest points, i.e., $g_N(1) = g_N(N) = 1/2$. Then $G_N(n) = (1 + \mathbf{1}\{n = N\})/2$, so that $G(n) = 1/2$ and $s^* = 1/2$. But also, $s_* = 1/2$. So the expected number of points in any interval not containing $1/2$ remains finite as $N \rightarrow \infty$; in other words, the marginal distribution of a typical point converges to a unit point mass at $1/2$. This example also serves to demonstrate that Theorem 3.2 may fail if (A4) does not hold.

4 Detailed example: Replace the minimum and one other

In this section we present some more specific results to complement our general results from Section 3. To do so, we specialize to the $K = 2$ case of Example (E3) from Section 3, in which we replace the smallest value and choose the other value to replace from $\{2, \dots, N\}$ according to a probability distribution f_N . Write

$$F_N(k) := \sum_{j=2}^k f_N(j), \quad (4.1)$$

for the corresponding distribution function, adopting the usual convention that an empty sum is zero, so that $F_N(0) = F_N(1) = 0$.

In the general set-up of Section 3, we have $\kappa_N(1, i) = f_N(i)/2$, $g_N(1) = 1/2$ and $g_N(i) = f_N(i)/2$ for $i \in \{2, \dots, N\}$. Here, assumption (A1) and (A3) are automatically satisfied; Assumption (A2) becomes a condition on f_N (or F_N), namely that

$$\lim_{N \rightarrow \infty} F_N(n) = F(n) \quad (4.2)$$

exists for all $n \geq 2$. The present version of (A2) is then:

(A2') Suppose that (4.2) holds.

Under (A2'),

$$\alpha := \lim_{n \rightarrow \infty} F(n) \quad (4.3)$$

exists in $[0, 1]$. Indeed, since $G_N(n) = (1 + F_N(n))/2$, we have that (A2') implies that s^* given by (3.3) satisfies $s^* = \frac{1+\alpha}{2}$.

Before stating our results, we comment briefly on their relation to previous work in the literature. The model of this section includes that studied by de Boer *et al.* [7] amongst others (see e.g. [20, §5.2.5]); the model of [7] is the special case where $F_N(n) = \frac{n-1}{N-1}$, which satisfies (A2) with $\alpha = 0$. Thus the $\alpha = 0$ cases of our results are not surprising in view of the (not completely rigorous) arguments in [7], or the heuristic analysis in [20, §5.2.5] that neglects correlations between the $X_t^{(k)}$, but our results are more general even in the case $\alpha = 0$, and we show explicitly the dependence of the phase transition on F_N via the parameter α . Moreover, one aim of the present work is to give a more rigorous approach to the results of [7] in the present considerably more general setting.

In this setting, the following result is immediate from Theorem 3.1.

Theorem 4.1. *Suppose that (A2') holds. Then*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] \begin{cases} < \infty & \text{if } s < \frac{1+\alpha}{2} \\ = \infty & \text{if } s > \frac{1+\alpha}{2} \end{cases}. \quad (4.4)$$

Similarly, we have the following translation of Theorem 3.2 into this setting. The appropriate version of condition (A4) is:

(A4') Suppose that F_N satisfies (3.6).

Theorem 4.2. *Suppose that (A2') and (A4') hold. With $V(s)$ as given by (3.8),*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \left(\frac{\mathbb{E}_N[C_t^N(s)]}{N} \right) = V(s), \quad s \in [0, 1].$$

Remark 4.1. *If instead of a $U[0, 1]$ distribution we use a distribution ρ for replacement points, as described in Remark 3.1, then the threshold exhibited in Theorem 4.1 becomes $s^* = \rho^{-1}(\frac{1+\alpha}{2})$; the inverse ρ^{-1} is well-defined when ρ satisfies the conditions described in Remark 3.1.*

Now we move on to our detailed results concerning the case $\alpha = 0$; note that $\alpha = 0$ if and only if $f_N(n) \rightarrow 0$ as $N \rightarrow \infty$ for any n . The case $\alpha = 0$ includes the discrete uniform case (as considered in [7]) in which $f_N(n) = \frac{1}{N-1}$, but includes many other possibilities. Theorem 4.1 shows that when $\alpha = 0$ the phase transition occurs at $s^* = 1/2$. The next result gives more information, giving an explicit expression for the limiting equilibrium expectation in the case in which it is finite.

Theorem 4.3. *Suppose that (A2') holds and that $\alpha = 0$. Then*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] = \begin{cases} 2s + \frac{s^2}{1-2s} & \text{if } 0 \leq s < 1/2 \\ \infty & \text{if } s \geq 1/2 \end{cases}. \quad (4.5)$$

We also have the following explicit description of the limit distribution.

Theorem 4.4. *Suppose that (A2') holds and that $\alpha = 0$. If $s < 1/2$, then for any $n \in \mathbb{Z}^+$,*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) = n] = \pi^s(n),$$

where

$$\begin{aligned} \pi^s(0) &= 1 - 2s; \\ \pi^s(1) &= 2s - \left(\frac{s}{1-s} \right)^2; \\ \pi^s(n) &= \left(1 - \left(\frac{s}{1-s} \right)^2 \right) \left(\frac{s}{1-s} \right)^{2(n-1)}, \quad (n \geq 2). \end{aligned} \quad (4.6)$$

On the other hand, if $s \geq 1/2$, then for any $n \in \mathbb{Z}^+$, $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \leq n] = 0$.

Remark 4.2. (i) *The corresponding stationary probabilities put forward in [7] do not sum to 1 (see equations (10)–(12) in [7]). Our argument is similar (based on the use of the ‘ $N = \infty$ ’ Markov chain) but we try to give a fuller justification. (ii) Let*

$$\tau_N(s) := \min\{t \in \mathbb{N} : C_t^N(s) = 0\}. \quad (4.7)$$

By standard Markov chain theory, $\pi_N^s(n) = (\mathbb{E}_N[\tau_N(s) \mid C_0^N(s) = 0])^{-1}$. So an immediate consequence of Theorem 4.4 is that (cf equation (16) of [7])

$$\lim_{N \rightarrow \infty} \mathbb{E}_N[\tau_N(s) \mid C_0^N(s) = 0] = \begin{cases} \frac{1}{1-2s} & \text{if } s < 1/2 \\ \infty & \text{if } s \geq 1/2 \end{cases}.$$

We also prove explicit limiting (marginal) distributions for the lower order statistics themselves. We use the notation

$$h_n(s) := \begin{cases} 2s & \text{if } n = 1 \\ \left(\frac{s}{1-s}\right)^{2(n-1)} & \text{if } n \geq 2 \end{cases}. \quad (4.8)$$

Theorem 4.5. *Suppose that (A2') holds and that $\alpha = 0$. Then for $n \in \mathbb{N}$,*

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[X_t^{(n)} \leq s] = \begin{cases} 0 & \text{if } s \leq 0 \\ h_n(s) & \text{if } 0 \leq s \leq 1/2 \\ 1 & \text{if } s \geq 1/2 \end{cases}. \quad (4.9)$$

The $n = 1$ case of (4.9) says that the large N , long-time distribution of the smallest component approaches a $U[0, 1/2]$ distribution. The distributions arising for $n \geq 2$ are not so standard, but, as $n \rightarrow \infty$, they approach a unit point mass at $1/2$.

Also note that Theorem 4.5 yields convergence of moments of the $X_t^{(n)}$. For example, for any $k \in \mathbb{N}$, $\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[(X_t^{(1)})^k] = 2^{-k}/(k+1)$, and for $n \geq 2$ and any $k \in \mathbb{N}$,

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[(X_t^{(n)})^k] = 2^{-k} - \frac{k2^{-(2n+k-2)}}{2n+k-2} {}_2F_1(2n-2, 2n+k-2; 2n+k-1; 1/2). \quad (4.10)$$

To see this, note that since $X_t^{(n)}$ is uniformly bounded, its moments converge to those of the distribution h_n by bounded convergence, so we have

$$\begin{aligned} \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[(X_t^{(n)})^k] &= k \int_0^{1/2} s^{k-1} \left(\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[X_t^{(n)} > s] \right) ds \\ &= k \int_0^{1/2} s^{k-1} (1 - h_n(s)) ds. \end{aligned}$$

When $n = 1$ this is $k \int_0^{1/2} (1 - 2s)s^{k-1} ds$ which yields the claimed result. When $n \geq 2$, using the substitution $u = 2s$, the limit becomes

$$2^{-k} - k2^{-2(n-1)-k} \int_0^1 u^{2(n-1)+k-1} (1 - (u/2))^{-2(n-1)} du,$$

which gives (4.10) via the integral representation of the hypergeometric function.

5 Further remarks and open problems

A multidimensional model

Allowing more general distributions W , as described in Remark 3.1, enables some multi-dimensional models to fit within the scope of our results. We describe one example. Let Z be a uniform random vector on $[0, 1]^2$, and let $\|\cdot\|$ denote the Euclidean norm. Starting with N points in $[0, 1]^2$, iterate the following Markovian model: at each step in discrete time, replace the minimal-ranked point, where the ranking is in order of increasing Euclidean distance from the origin, and another point (chosen uniformly at random) with independent copies of Z . This model corresponds to the model described in Section 4 but with the U_i replaced by copies of $W = \|Z\|$, and with $\alpha = 0$. Elementary calculations show that $\rho(x) := \mathbb{P}[W \leq x] = \frac{\pi x^2}{4}$ for $x \in [0, 1]$ ($\rho(x)$ is more complicated for $x \geq 1$), so that the phase transition (see Remark 4.1) occurs at $s^* = \rho^{-1}(1/2) = \sqrt{2/\pi} \approx 0.80$. See Figure 1 for a simulation.




Figure 1: Simulation of the model in which, at each step, the closest point to the origin and one uniformly random other point are replaced by independent uniform random points on $[0, 1]^2$, with $N = 10^4$ points and $t = 10^6$ steps. The initial distribution was N independent uniform points on $[0, 1]^2$. Also shown in the figure is part of the circle centred at the origin with radius $\sqrt{2}/\pi$. [colour online]

A partial-order-driven process

Here is a variation on the multidimensional model of the previous example governed by a *partial order* rather than a total order. Again consider a system of N points in $[0, 1]^2$. Consider the co-ordinatewise partial order ' \preceq ' under which $(x_1, y_1) \preceq (x_2, y_2)$ if and only if $x_1 \leq x_2$ and $y_1 \leq y_2$; a point x of a finite set $\mathcal{X} \subset [0, 1]^2$ is *minimal* if and only if there is no $y \in \mathcal{X} \setminus \{x\}$ for which $y \preceq x$. Now define a discrete-time Markov process as follows: at each step, replace a minimal element of the N points (chosen uniformly at random from amongst all possibilities) and a non-minimal element (again, chosen uniformly at random); all new points are independent and uniform on $[0, 1]^2$. This model seems more difficult to study than the previous one, although simulations suggest qualitatively similar asymptotic behaviour: see Figure 2.

A repeated beauty contest

We describe a process of a different flavour to those previously considered, in which the update rule depends not only on the ranks of the points; this is a variation on a *Keynesian*




Figure 2: Simulation of the model in which, at each step, one \preccurlyeq -minimal element and one non-minimal element (each uniformly chosen) are replaced by independent uniform random points on $[0, 1]^2$, with $N = 10^4$ points and $t = 10^6$ steps. The initial distribution was N independent uniform points on $[0, 1]^2$. Can the threshold curve be characterized? [colour online]

beauty contest.

Fix a parameter $p > 0$. Start with a uniform array of N elements on $[0, 1]$. At each step, compute the mean μ of the N elements, and replace by a $U[0, 1]$ random variable the element that is farthest (amongst all the N points) from $p\mu$. Thus at each step, either the minimum or maximum is replaced, depending on the current configuration.

This is related to the “ p -beauty contest” [27, p. 72] in which N players choose a number between 0 and 100, the winner being the player whose choice is closest to p times the average of all the N choices. The stochastic process described above is a repeated, randomized version of this game (without any learning, and with random player behaviour) in which the worst performer is replaced by a new player.

According to simulations and heuristic considerations, the equilibrium distribution of a typical point approaches, for large N , a point mass at 0 (1) in the case $p < 1$ ($p > 1$). The case $p = 1$ is more subtle, and is reminiscent of a Pólya urn. Stochastic approximation ideas (see e.g. [28]) may be relevant in studying this model.

6 General thresholds: Proofs for Section 3

6.1 Overview

This section contains the proofs of our general results from Section 3, and is arranged as follows. In Section 6.2 we give a basic result on the Markov chains $C_t^N(s)$. To study the $N \rightarrow \infty$ asymptotics of these Markov chains, at least when $s < s^*$, we introduce an ‘ $N = \infty$ ’ Markov chain $C_t(s)$. In Section 6.3 we show that we can define $C_t(s)$ in a consistent way, and we prove some of its basic properties. In Section 6.4 we relate the asymptotic properties of the finite- N chains $C_t^N(s)$ with $s < s^*$ to the chain $C_t(s)$, making use of our technical results from Section 8. Then in Section 6.5 we complete the proofs of Theorems 3.1 and 3.2.

6.2 The Markov chain $C_t^N(s)$

We have the following basic result.

Lemma 6.1. *Fix $N \in \mathbb{N}$. Suppose that (A1) holds. Suppose that $s \in (0, 1)$. Then $C_t^N(s)$ is an irreducible, aperiodic Markov chain on $\{0, 1, \dots, N\}$ with uniformly bounded jumps: $\mathbb{P}_N[|C_{t+1}^N(s) - C_t^N(s)| > K] = 0$. There exists a unique stationary distribution π_N^s , with $\pi_N^s(n) > 0$ for all $n \in \{0, 1, \dots, N\}$ and $\sum_{n=0}^N \pi_N^s(n) = 1$, such that*

$$\lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) = n] = \pi_N^s(n), \quad (6.1)$$

for any initial distribution $C_0^N(s)$. Moreover,

$$\lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] = \sum_{n=1}^N n \pi_N^s(n). \quad (6.2)$$

Finally, with G_N as defined at (3.1), the one-step mean drift of $C_t^N(s)$ is given by

$$\mathbb{E}_N[C_{t+1}^N(s) - C_t^N(s) \mid C_t^N(s) = n] = K(s - G_N(n)). \quad (6.3)$$

Note that the degenerate cases $s \in \{0, 1\}$ are excluded from Lemma 6.1: $C_t^N(1) = N$ a.s. for all t , while $C_t^N(0) \rightarrow 0$ a.s. as $t \rightarrow \infty$ for any initial distribution $C_0^N(0)$.

Proof of Lemma 6.1. Note that

$$\{C_t^N(s) = n\} = \{X_t^{(1)} \leq s, \dots, X_t^{(n)} \leq s, X_t^{(n+1)} > s, \dots, X_t^{(N)} > s\};$$

the distribution of $C_{t+1}^N(s)$ depends only on $(X_t^{(1)}, \dots, X_t^{(N)})$ through events of the form on the right-hand side of the last display. Specifically, given $C_t^N(s) = n$, we have that the increment $C_{t+1}^N(s) - C_t^N(s)$ is the number of the K new $U[0, 1]$ -distributed points that fall in $[0, s]$ minus the number of the K points selected for removal whose rank was at most n . That is, with the notation

$$\begin{aligned} B_{t+1}(s) &:= \#\{i \in \{Kt + 1, \dots, Kt + K\} : U_i \in [0, s]\}, \\ A_{t+1}^N(n) &:= \#\{i \in \{1, \dots, K\} : R_i^N(t + 1) \leq n\}, \end{aligned}$$

we have that, given $C_t^N(s) = n$,

$$C_{t+1}^N(s) = n + B_{t+1}(s) - A_{t+1}^N(n). \quad (6.4)$$

Thus, given $C_t^N(s) = n$, the increment depends only on n and the variables $R^N(t+1)$, $U_{Kt+1}, \dots, U_{Kt+K}$, which are all independent of $C_t^N(s)$. This demonstrates the Markov property.

The bounded jumps property is clear by definition, and can also be seen from (6.4). To show irreducibility and aperiodicity, we show that

$$\begin{aligned}\mathbb{P}_N[C_{t+1}^N(s) = n \mid C_t^N(s) = n] &> 0, \quad (n \in \{0, 1, \dots, N\}), \\ \mathbb{P}_N[C_{t+1}^N(s) = n+1 \mid C_t^N(s) = n] &> 0, \quad (n \in \{0, 1, \dots, N-1\}), \\ \mathbb{P}_N[C_{t+1}^N(s) = n-1 \mid C_t^N(s) = n] &> 0, \quad (n \in \{1, 2, \dots, N\}).\end{aligned}$$

Since $B_{t+1}(s)$ and $A_{t+1}^N(n)$ are independent given $C_t^N(s) = n$, it suffices to show that $\mathbb{P}_N[B_{t+1}(s) = i] > 0$ for any $i \in \{0, 1, \dots, K\}$, and that $\mathbb{P}_N[A_{t+1}^N(n) = i] > 0$ for: (i) $i \in \{0, 1\}$ if $K = 1$; or (ii) $i = 1$ if $K \geq 2$. Then the intersection of two independent events of positive probability will yield any increment of $C_t^N(s)$ in $\{-1, 0, 1\}$, as required. First consider $B_{t+1}(s)$: this has a $\text{Bin}(K, s)$ distribution, and so takes any value in $\{0, 1, \dots, K\}$ with positive probability, provided $s \in (0, 1)$. Now consider $A_{t+1}^N(n)$. Then

$$A_{t+1}^N(n) = \sum_{i=1}^n \mathbf{1}\{i \in \{R_1^N(t+1), \dots, R_K^N(t+1)\}\}. \quad (6.5)$$

It follows from (6.5) that, for $n \geq 1$, $A_{t+1}^N(n) \geq \mathbf{1}\{1 \in \{R_1^N(t+1), \dots, R_K^N(t+1)\}\}$, so

$$\mathbb{P}_N[A_{t+1}^N(n) = 1] \geq \mathbb{P}_N[1 \in \{R_1^N(t+1), \dots, R_K^N(t+1)\}] \geq \mathbb{P}_N[R_1^N = 1].$$

This latter probability is $g_N(1)$, which is positive by (A1). This completes the proof of irreducibility and aperiodicity in the case $K \geq 2$; it remains to show that $\mathbb{P}_N[A_{t+1}^N(n) = 0] > 0$ for $n \geq 0$ when $K = 1$. Using the $K = 1$ case of (6.5), we obtain

$$\mathbb{P}_N[A_{t+1}^N(n) = 0] = \mathbb{P}_N[R_1^N(t+1) > n] = 1 - G_N(n),$$

by (3.1), and $1 - G_N(n) > 0$ since (A1) implies that in this case $g_N(i) > 0$ for some $i > n$.

Thus the Markov chain is irreducible and aperiodic; it has a finite state-space, and so standard Markov chain theory implies the existence of a unique stationary distribution, for which (6.1) holds. Moreover, since $C_t^N(s)$ is bounded by N , (6.2) follows from (6.1).

Finally we prove the statement (6.3). We take expectations in (6.4); $B_{t+1}(s)$ has mean Ks , and taking expectations in (6.5) we obtain

$$\mathbb{E}[A_{t+1}^N(n)] = \sum_{i=1}^n \mathbb{P}[i \in \{R_1^N(t+1), \dots, R_K^N(t+1)\}] = K \sum_{i=1}^n \mathbb{P}[R_1^N = i],$$

by exchangeability. Thus from (3.1) we obtain (6.3). \square

A key step in our analysis is to study the stationary distributions π_N^s of the Markov chains $C_t^N(s)$, $s \in (0, 1)$, whose existence is proved in Lemma 6.1. We consider π_N^s as $N \rightarrow \infty$. One tool that we will use is a Markov chain $C_t(s)$ on the whole of \mathbb{Z}^+ that can be viewed in some sense as the $N \rightarrow \infty$ limit of the Markov chains $C_t^N(s)$: this Markov chain we call the ‘ $N = \infty$ ’ chain, and we describe it in Section 6.3; in Section 6.4 we make precise the sense in which the ‘ $N = \infty$ ’ chain is a limit of the finite- N chains.

6.3 The ‘ $N = \infty$ ’ chain $C_t(s)$

Our asymptotic analysis makes use of an ‘ $N = \infty$ ’ analogue of the Markov chain $C_t^N(s)$. The case $N = \infty$ does not make sense directly in terms of the original model X_t , but (A2) can be used to define a Markov chain on the whole of \mathbb{Z}^+ , which we can relate to our finite- N Markov chains, at least when $s < s^*$.

We use $C_t(s)$ to denote our new Markov chain, now defined on the whole of \mathbb{Z}^+ , and we write \mathbb{P} for the associated probability measure and \mathbb{E} for the corresponding expectation. The idea is to define transition probabilities via

$$\mathbb{P}[C_{t+1}(s) = m \mid C_t(s) = n] = \lim_{N \rightarrow \infty} \mathbb{P}_N[C_{t+1}^N(s) = m \mid C_t^N(s) = n];$$

to show that this is legitimate under suitable assumptions, we need the following result.

Lemma 6.2. *Suppose that (A2) holds. Let $s \in [0, 1]$. Then for any $n, m \in \mathbb{Z}^+$,*

$$p^s(n, m) := \lim_{N \rightarrow \infty} \mathbb{P}_N[C_{t+1}^N(s) = m \mid C_t^N(s) = n]$$

is well-defined, and $\sum_{m \in \mathbb{Z}^+} p^s(n, m) = 1$.

Proof. We show that the increment distribution, conditional on $\{C_t^N(s) = n\}$, given by (6.4) in the finite N case converges (as $N \rightarrow \infty$), using assumption (A2), to an appropriate limiting distribution, which will serve as the increment distribution $p^s(n, \cdot)$. This convergence is clear for the term $B_{t+1}(s)$, which has no N -dependence. Moreover, given $C_t^N(s) = n$, the terms $B_{t+1}(s)$ and $A_{t+1}^N(n)$ are independent. Thus it suffices to show that $A_{t+1}^N(n)$ converges in distribution to a proper random variable. We show that this follows from (A2), although care is needed to correctly account for lost mass in (A2).

To proceed, it is useful to introduce more notation. Let $R = (R_1, \dots, R_K)$ denote the $N \rightarrow \infty$ distributional limit of R^N : given (A2), this limit exists but is not necessarily a proper distribution on \mathbb{N}^K , but we recover a proper distribution by expanding the state-space to $(\mathbb{N} \cup \{\infty\})^K$. Thus components of R may take the value ∞ : this cannot be directly interpreted in terms of rank distributions, but is convenient for correctly accounting for the lost mass in (A2). Concretely, the distribution of R is given, for any $k \leq K$ and any distinct $i_1, i_2, \dots, i_k \in \mathbb{N}$, by

$$\begin{aligned} & \mathbb{P}[R_1 = i_1, \dots, R_k = i_k, R_{k+1} = \infty, \dots, R_K = \infty] \\ &= \lim_{N \rightarrow \infty} \mathbb{P}_N[R_1^N = i_1, \dots, R_k^N = i_k] - \sum_{i_{k+1} \in \mathbb{N}} \cdots \sum_{i_K \in \mathbb{N}} \lim_{N \rightarrow \infty} \mathbb{P}_N[R_1^N = i_1, \dots, R_K^N = i_K] \\ &= \kappa(i_1, \dots, i_k) - \sum_{i_{k+1} \in \mathbb{N}} \cdots \sum_{i_K \in \mathbb{N}} \kappa(i_1, \dots, i_K), \end{aligned} \tag{6.6}$$

using (A2). Note that since R^N is exchangeable on $\{1, \dots, N\}^K$, it follows that R is exchangeable on $(\mathbb{N} \cup \{\infty\})^K$.

Now we can define the $N = \infty$ analogue of $A_{t+1}^N(s)$ to be an independent copy of $\#\{i \in \{1, \dots, K\} : R_i \leq n\}$, i.e., for $R(t+1) = (R_1(t+1), \dots, R_K(t+1))$ an independent copy of R , with distribution given by (6.6), we take

$$A_{t+1}(n) := \sum_{i=1}^K \mathbf{1}\{R_i(t+1) \leq n\}.$$

Then we can construct $C_t(s)$ via its increments

$$C_{t+1}(s) - C_t(s) = B_{t+1}(s) - A_{t+1}(C_t(s)). \quad (6.7)$$

Since R^N converges in distribution to R as $N \rightarrow \infty$, $A_{t+1}^N(n) = \sum_{i=1}^K \mathbf{1}\{R_i^N(t+1) \leq n\}$ converges in distribution to $A_{t+1}(n)$; specifically, using exchangeability,

$$\begin{aligned} \mathbb{P}_N[A_{t+1}^N(n) = k] &= \binom{K}{k} \mathbb{P}_N[R_1^N \leq n, \dots, R_k^N \leq n, R_{k+1}^N > n, \dots, R_K^N > n] \\ &\rightarrow \binom{K}{k} \mathbb{P}[R_1 \leq n, \dots, R_k \leq n, R_{k+1} > n, \dots, R_K > n], \end{aligned}$$

as $N \rightarrow \infty$. This completes the proof. \square

The following result gives some basic properties of the Markov chain defined above.

Lemma 6.3. *Suppose that (A2) and (A3) hold. Then for any $s \in (0, 1)$, $C_t(s)$ is an irreducible, aperiodic Markov chain on \mathbb{Z}^+ , with uniformly bounded jumps: $\mathbb{P}[|C_{t+1}(s) - C_t(s)| > K] = 0$. The one-step mean drift of $C_t(s)$ is given by*

$$\mathbb{E}[C_{t+1}(s) - C_t(s) \mid C_t(s) = n] = K(s - G(n)). \quad (6.8)$$

Proof. The boundedness of the increments follows from the construction in (6.7). The irreducibility and aperiodicity follow from a similar argument to that used in the proof of Lemma 6.1 in the finite- N case, now using (A3) in place of (A1). The drift (6.8) also follows similarly to the proof of (6.3) in Lemma 6.1; in the present case

$$\mathbb{E}[A_{t+1}(n)] = \sum_{i=1}^n \mathbb{P}[i \in \{R_1, \dots, R_K\}] = K \sum_{i=1}^n \mathbb{P}[R_1 = i],$$

by exchangeability of R (see the comment after (6.6)). But $\sum_{i=1}^n \mathbb{P}[R_1 = i] = G(n)$, by (A2) and the definition of $G(n)$ at (3.2). \square

6.4 Large- N asymptotics

We show that properties of the Markov chains $C_t^N(s)$, described in Section 6.2, in the large N limit can (at least when $s < s^*$) be described using the ‘ $N = \infty$ ’ Markov chain $C_t(s)$, described in Section 6.3. The main tool is Theorem 8.1 stated and proved in Section 8. Recall the definition of π_N^s from Lemma 6.1.

Lemma 6.4. *Suppose that (A1), (A2), and (A3) hold, and that $s \in (0, s^*)$. There exists a unique stationary distribution π^s for $C_t(s)$, with $\pi^s(n) > 0$ for all $n \in \mathbb{Z}^+$ and $\sum_{n \in \mathbb{Z}^+} \pi^s(n) = 1$, such that, for all $n \in \mathbb{Z}^+$,*

$$\lim_{t \rightarrow \infty} \mathbb{P}[C_t(s) = n] = \pi^s(n), \quad (6.9)$$

for any initial distribution $C_0(s)$. In addition, the following results hold.

- (a) *There exist $c > 0$ and $C < \infty$ such that, for all $n \in \mathbb{Z}^+$, $\pi_N^s(n) \leq Ce^{-cn}$ and $\pi^s(n) \leq Ce^{-cn}$.*
- (b) *For any $n \in \mathbb{Z}^+$, $\lim_{N \rightarrow \infty} \pi_N^s(n) = \pi^s(n)$.*

(c) As $t \rightarrow \infty$, $\mathbb{E}_N[C_t^N(s)] \rightarrow \sum_{n=0}^N n\pi_N^s(n)$ and $\mathbb{E}[C_t(s)] \rightarrow \sum_{n \in \mathbb{Z}^+} n\pi^s(n) < \infty$.

Proof. We will show that we can apply Theorem 8.1 with $Y_t^N = C_t^N(s)$, $Y_t = C_t(s)$, $S_N = \{0, 1, \dots, N\}$, and $S = \mathbb{Z}^+$. Since $s \in (0, 1)$ and (A1) holds, Lemma 6.1 shows that $C_t^N(s)$ is an irreducible Markov chain on $\{0, 1, \dots, N\}$, while, since (A2) and (A3) hold, Lemma 6.3 implies that $C_t(s)$ is an irreducible Markov chain on \mathbb{Z}^+ . Lemmas 6.1 and 6.3 also imply that the increments of $C_t^N(s)$ and $C_t(s)$ are uniformly bounded in absolute value (by K) almost surely. Thus (8.1) holds.

Next we verify the drift conditions in (8.2). Since $s < s^*$, there exists $\varepsilon > 0$ such that $s < s^* - 2\varepsilon$. First consider the finite- N case. By (A2) and the definition of s^* at (3.3), given ε , we can take N_0 and n_0 such that for any $N \geq N_0$ and any $n \geq n_0$,

$$G_N(n) > s^* - \varepsilon > s + \varepsilon.$$

So we have from (6.3) that, for all $N \geq N_0$ and $n \geq n_0$,

$$\mathbb{E}_N[C_{t+1}^N(s) - C_t^N(s) \mid C_t^N(s) = n] \leq -\varepsilon K.$$

A similar argument holds for $C_t(s)$, using (6.8). Thus (8.2) is satisfied. Finally, we verify (8.3) by Lemma 6.2. Thus Theorem 8.1 applies, yielding the claimed results. \square

The next result deals with the case $s > s^*$. Recall that $\tau_N(s)$ defined by (4.7) denotes the time of the first return of $C_t^N(s)$ to 0.

Lemma 6.5. *Suppose that (A1), (A2), and (A3) hold, and that $s > s^*$. Then $\lim_{N \rightarrow \infty} \mathbb{E}_N[\tau_N(s)] = \infty$.*

Proof. Suppose that $s > s^*$. Then, for some $\varepsilon > 0$, $s - s^* - \varepsilon > \varepsilon$. Fix $x \in \mathbb{N}$. Since, by (A2), $\lim_{N \rightarrow \infty} G_N(n) = G(n) \leq s^*$ for any n , we can find $N_0(x)$ such that $G_N(n) \leq s^* + \varepsilon$ for any $n \leq x$ and any $N \geq N_0(x)$. Hence, by (6.3),

$$\mathbb{E}_N[C_{t+1}^N(s) - C_t^N(s) \mid C_t^N(s) = n] \geq K\varepsilon, \quad (6.10)$$

for any $N \geq N_0(x)$ and any $n \leq x$, where $\varepsilon > 0$ does not depend on x . We show that (6.10) implies that $C_t^N(s)$ has a positive probability (uniform in x) of reaching x before returning to 0, which will imply the result. It suffices to suppose that $C_0^N(s) \geq 1$.

To ease notation, write $\tau := \tau_N(s)$ for the remainder of this proof. To estimate the required hitting probability, set $W_t := \exp\{-\delta C_t^N(s)\}$, for $\delta > 0$ to be chosen later. Now

$$\begin{aligned} W_{t+1} - W_t &= \exp\{-\delta C_t^N(s)\} (\exp\{-\delta(C_{t+1}^N(s) - C_t^N(s))\} - 1) \\ &\leq \exp\{-\delta C_t^N(s)\} (-\delta(C_{t+1}^N(s) - C_t^N(s)) + M\delta^2), \end{aligned}$$

for some absolute constant M , using the fact that the increments of $C_t^N(s)$ are uniformly bounded. Taking expectations and using (6.10), we have that, on $\{C_t^N(s) \leq x\}$,

$$\mathbb{E}_N[W_{t+1} - W_t \mid C_t^N(s)] \leq \exp\{-\delta C_t^N(s)\} (-K\varepsilon\delta + M\delta^2) \leq 0,$$

for $\delta \leq \delta_0$ small enough, where $\delta_0 > 0$ depends only on ε and not on x or N . Let $\nu_x := \min\{t \in \mathbb{Z}^+ : C_t^N(s) \geq x\}$. Then we have shown that $W_{t \wedge \tau \wedge \nu_x}$ is a nonnegative supermartingale, which converges a.s. to $W_{\tau \wedge \nu_x}$. It follows that

$$e^{-\delta} \geq W_0 \geq \mathbb{E}_N[W_{\tau \wedge \nu_x}] \geq \mathbb{P}_N[\tau < \nu_x],$$

so that $\mathbb{P}_N[\nu_x < \tau] \geq 1 - e^{-\delta} =: p$, where $p > 0$ does not depend on x or on N . The fact that $C_t^N(s)$ has increments of size at most K implies that on $\{\nu_x < \tau\}$ we have $\{\tau \geq x/K\}$. Hence $\mathbb{P}_N[\tau \geq x/K] \geq \mathbb{P}_N[\nu_x < \tau] \geq p$, so that $\mathbb{E}_N[\tau] \geq px/K$ for all $N \geq N_0(x)$. Since x was arbitrary, the result follows. \square

6.5 Proofs of Theorems 3.1 and 3.2

Proof of Theorem 3.1. First suppose that $s < s^*$. Then Lemma 6.4 applies. By Lemma 6.4(a), $\pi_N^s(n) \leq Ce^{-cn}$ where $C < \infty$ and $c > 0$ do not depend on N or n . In particular, for any $p > 0$, $\sup_N \sum_{n \in \mathbb{Z}^+} n^p \pi_N^s(n) < \infty$. Moreover, by Lemma 6.4(b), $\pi_N^s(n) \rightarrow \pi^s(n)$ as $N \rightarrow \infty$. Hence for any $p > 0$, by uniform integrability, $\sum_n n^p \pi_N^s(n) \rightarrow \sum_n n^p \pi^s(n)$ as $N \rightarrow \infty$. Together with Lemma 6.4(c), this implies that, for $s < s^*$,

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] = \lim_{N \rightarrow \infty} \sum_{n \in \mathbb{Z}^+} n \pi_N^s(n) = \sum_{n \in \mathbb{Z}^+} n \pi^s(n) < \infty. \quad (6.11)$$

Next suppose that $s > s^*$. Then, for fixed $x > 0$ and some $\varepsilon > 0$ (not depending on x) we have that (6.10) holds for any $N \geq N_0(x)$ and any $n \leq x$. On the other hand, if $C_t^N(s) > x$, we have that $C_{t+1}^N(s) \geq x - K$ (by bounded jumps). It follows that

$$\mathbb{E}_N[C_{t+1}^N(s) - C_t^N(s) \mid C_t^N(s)] \geq K(1 + \varepsilon) \mathbf{1}\{C_t^N(s) \leq x\} - K.$$

Taking expectations implies that

$$\mathbb{E}_N[C_{t+1}^N(s)] - \mathbb{E}_N[C_t^N(s)] \geq K(1 + \varepsilon) \mathbb{P}_N[C_t^N(s) \leq x] - K.$$

By (6.2), the left-hand side of the last display tends to 0 as $t \rightarrow \infty$. It follows that, for some $\delta > 0$ that depends on ε but not on x , $\mathbb{P}_N[C_t^N(s) \geq x] \geq \delta$ for all t large enough. Hence $\mathbb{E}_N[C_t^N(s)] \geq x\delta$, for all N and t sufficiently large. Since x was arbitrary, and δ did not depend on x , the second part of the theorem follows. \square

Proof of Theorem 3.2. For $s < s^*$, the statement follows immediately from Theorem 4.1.

Suppose that $s > s^*$. For the duration of this proof, we write τ for $\tau_N(s)$ to ease notation. In this case, Lemma 6.5 applies, showing that $\lim_{N \rightarrow \infty} \mathbb{E}_N[\tau] = \infty$. We claim that $C_t^N(s)$ is asymptotically null in the sense that, for any $n \in \mathbb{Z}^+$,

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \leq n] = 0. \quad (6.12)$$

Indeed, (6.12) follows from Lemma 6.5 and the occupation-time representation for the stationary distribution of an irreducible, positive-recurrent Markov chain (see e.g. [1, Corollary I.3.6, p. 14]), which implies that

$$\lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \leq n] = \sum_{x=0}^n \pi_N^s(x) = \frac{\mathbb{E}_N \sum_{t=0}^{\tau-1} \mathbf{1}\{C_t^N(s) \leq n\}}{\mathbb{E}_N[\tau]},$$

in the final fraction, the denominator tends to infinity with N (by Lemma 6.5) while the numerator is uniformly bounded in N since the expected number of visits to any bounded interval stays bounded, by irreducibility (uniform in N). Thus (6.12) holds for $s > s^*$.

Taking expectations in (6.3) yields

$$\mathbb{E}_N[C_{t+1}^N(s)] - \mathbb{E}_N[C_t^N(s)] = Ks - K\mathbb{E}_N[G_N(C_t^N(s))]. \quad (6.13)$$

The left-hand side of (6.13) tends to 0 as $t \rightarrow \infty$ by (6.2). Also, for n_0 as in (A4),

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[G_N(C_t^N(s)) \mathbf{1}\{C_t^N(s) < n_0\}] \leq \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \leq n_0],$$

which is 0 by (6.12). Hence, taking limits in (6.13), we obtain

$$s = \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[G_N(C_t^N(s)) \mathbf{1}\{C_t^N(s) \geq n_0\}].$$

By condition (A4), a.s.,

$$G_N(C_t^N(s)) \mathbf{1}\{C_t^N(s) \geq n_0\} = s^* + (1 - s^*) \frac{C_t^N(s)}{N} + \varepsilon_N,$$

where $\varepsilon_N = o(1)$ as $N \rightarrow \infty$, uniformly in $C_t^N(s)$ (and hence uniformly in t). Thus

$$s = \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} (s^* + (1 - s^*) N^{-1} \mathbb{E}_N[C_t^N(s)]),$$

which yields the result (3.9) for $s > s^*$.

Finally, suppose that $s = s^*$. Then $C_t^N(s) \leq C_t^N(r)$ for all t, N , and $r > s^*$. Hence

$$\begin{aligned} \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} (N^{-1} \mathbb{E}[C_t^N(s)]) &\leq \lim_{r \downarrow s} \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} (N^{-1} \mathbb{E}[C_t^N(r)]) \\ &= \lim_{r \downarrow s} V(r), \end{aligned}$$

by the previous part of the proof, since $r > s^*$. The latter limit is $V(s^*) = 0$, so the result (3.9) is proved for $s = s^*$ as well. \square

7 Proofs for Section 4

7.1 Overview

In this section we study the $K = 2$ case of Example (E3), working towards proofs of the results in Section 4. The organization of the section broadly mirrors that of Section 6. In Sections 7.2 and 7.3 we return to the finite- N Markov chain $C_t^N(s)$ and the limit chain $C_t(s)$, respectively, describing their properties more explicitly in this special case, for which exact computations are available. Then in Section 7.4 we give the proofs of our remaining results, Theorems 4.3, 4.4, and 4.5.

7.2 The Markov chain $C_t^N(s)$

For this model, the following result complements the general Lemma 6.1. Write $p_N^s(n, m) := \mathbb{P}_N[C_{t+1}^N(s) = m \mid C_t^N(s) = n]$. Recall the definition of F_N from (4.1), and that $F_N(0) = F_N(1) = 0$. In the case where $F_N(n) = \frac{n-1}{N-1}$, $p_N^s(n, m)$ was written down in equations (1)–(3) in [7].

Lemma 7.1. *For any $s \in [0, 1]$, $(C_t^N(s))_{t \in \mathbb{Z}^+}$ is a Markov chain on $\{0, 1, 2, \dots, N\}$ under \mathbb{P}_N . The transition probabilities are given by*

$$p_N^s(0, 0) = (1 - s)^2, \quad p_N^s(0, 1) = 2s(1 - s), \quad p_N^s(0, 2) = s^2,$$

and for $n \geq 1$,

$$\begin{aligned} p_N^s(n, n - 2) &= (1 - s)^2 F_N(n) \\ p_N^s(n, n - 1) &= 2s(1 - s) F_N(n) + (1 - s)^2 (1 - F_N(n)) \end{aligned}$$

$$\begin{aligned} p_N^s(n, n) &= s^2 F_N(n) + 2s(1-s)(1-F_N(n)) \\ p_N^s(n, n+1) &= s^2(1-F_N(n)). \end{aligned} \quad (7.1)$$

Moreover, for $n \geq 0$,

$$\mathbb{E}_N[C_{t+1}^N(s) - C_t^N(s) \mid C_t^N(s) = n] = 2s - (1 + F_N(n))\mathbf{1}\{n \neq 0\}. \quad (7.2)$$

Proof. Suppose that $C_t^N(s) = n$. If $n = 0$, then the two points that we select come from $(s, 1)$, and $C_{t+1}^N(s)$ is 1 or 2 according to whether 1 or 2 of the new points land in $[0, s]$: each does so, independently, with probability s . This gives $p_N^s(0, m)$.

Suppose that $n \geq 1$. In this case, $X_t^{(1)} \leq s$ is always removed. Then $C_{t+1}^N(s)$ is either (i) $n-2$; (ii) $n-1$; or (iii) n according to whether (i) the second point selected for removal is one of $\{X_t^{(2)}, \dots, X_t^{(n)}\}$, and both new points fall in $(s, 1)$; (ii) the second point is one of $\{X_t^{(2)}, \dots, X_t^{(n)}\}$, and exactly one of the two new points falls in $(s, 1)$, or the second point is one of $\{X_t^{(n+1)}, \dots, X_t^{(N)}\}$, and both new points fall in $(s, 1)$; or (iii) the second point is one of $\{X_t^{(n+1)}, \dots, X_t^{(N)}\}$, and both new points fall in $[0, s]$. Thus we obtain the expressions in (7.1), noting that the probability that one of $\{X_t^{(2)}, \dots, X_t^{(n)}\}$ is selected as the second point for removal is $F_N(n)$. We then obtain (7.2) from (7.1). \square

7.3 The ‘ $N = \infty$ ’ chain $C_t(s)$

Again we consider the ‘ $N = \infty$ ’ chain $C_t(s)$ as described in Section 6.3. In the special case where (A2') holds, so that F is the limiting distribution given by (4.2), and $\alpha = 0$, the transition probabilities $p^s(n, m) = \mathbb{P}[C_{t+1}(s) = m \mid C_t(s) = n]$ are given for $n = 0$ by

$$p^s(0, 0) = (1-s)^2, \quad p^s(0, 1) = 2s(1-s), \quad p^s(0, 2) = s^2,$$

and for $n \in \mathbb{N}$ by

$$p^s(n, n-1) = (1-s)^2, \quad p^s(n, n) = 2s(1-s), \quad p^s(n, n+1) = s^2.$$

In their analysis, de Boer *et al.* [7] discuss this Markov chain, although they do not give full justification that it can be used to describe the asymptotics of the finite- N chains $C_t^N(s)$; this is justified in a specific sense by our results from Section 6.4, which rely on the technical tools from Section 8. In the remainder of this section we present some basic properties of $C_t(s)$.

The Markov chain $C_t(s)$ is *almost* a nearest-neighbour random walk (or birth-and-death chain), apart from the fact that from 0 we can make a jump of size 2. However, the form of the transition probabilities allows us to use a trick to transform this into a nearest-neighbour process (see the proof of Lemma 7.2 below). We will prove the following result, which corrects an error in the stationary distribution proposed in [7].

Lemma 7.2. *Suppose that $\alpha = 0$ and $s < 1/2$. Then for any $n \in \mathbb{Z}^+$,*

$$\lim_{t \rightarrow \infty} \mathbb{P}[C_t(s) = n] = \pi^s(n), \quad (7.3)$$

and the stationary distribution π^s satisfies (4.6). Moreover,

$$\lim_{t \rightarrow \infty} \mathbb{E}[C_t(s)] = 2s + \frac{s^2}{1-2s}. \quad (7.4)$$

Proof. Observe that the probability of a jump from 0 to 2 is the same as that from 1 to 2 (namely, s^2), while the probability of a jump from 0 into the set $\{0, 1\}$ is the same as that from 1 into $\{0, 1\}$ ($1 - s^2$). So we can merge $\{0, 1\}$ into a single state and preserve the Markov property. (A formal verification of the preservation of the Markov property under this transformation is provided by, for example, [9, Corollary 1].) This gives a genuine birth-and-death chain on a state-space isomorphic to \mathbb{Z}^+ . Call this new state-space $\{\bar{0}, \bar{1}, \bar{2}, \dots\}$, so that $\bar{0}$ corresponds to $\{0, 1\}$ and \bar{n} for $n \geq 1$ corresponds to $n + 1$ in the original state-space. This new Markov chain has transition probabilities

$$q^s(\bar{0}, \bar{0}) = 1 - s^2, \quad q^s(\bar{0}, \bar{1}) = s^2, \quad \text{and for } \bar{n} \geq 1, \quad q^s(\bar{n}, \bar{m}) = p^s(n + 1, m + 1).$$

This Markov chain is reversible and solving the detailed balance equations (cf e.g. [6, §I.12, pp. 71–76]) we obtain the stationary distribution $\bar{\pi}^s$ for $s < 1/2$ as

$$\bar{\pi}^s(\bar{n}) = \left(1 - \left(\frac{s}{1-s}\right)^2\right) \left(\frac{s}{1-s}\right)^{2n},$$

for all $n \geq 0$. To obtain $\pi^s(n)$, the stationary distribution for the original Markov chain, we need to disentangle the composite state $\bar{0}$. We have that $\pi^s(0) + \pi^s(1) = \bar{\pi}^s(\bar{0})$ and, by stationarity,

$$\pi^s(0) = (1 - s)^2 \pi^s(0) + (1 - s)^2 \pi^s(1).$$

Solving these equations we obtain (4.6). Some algebra then yields the mean of the distribution π^s (when $s < 1/2$), giving

$$\sum_{n \in \mathbb{Z}^+} n \pi^s(n) = 2s + \frac{s^2}{1 - 2s} < \infty, \tag{7.5}$$

since $s < 1/2$. Hence (7.4) follows from (7.5) and Lemma 6.4(c). \square

7.4 Proofs of Theorems 4.3, 4.4, and 4.5

Now we can complete the proofs of our remaining theorems.

Proof of Theorem 4.3. The $s < 1/2$ statement follows from (6.11) and (7.5). On the other hand, for any $s \geq 1/2$ and any $r < 1/2$, we have $C_t^N(s) \geq C_t^N(r)$, so

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{E}_N[C_t^N(s)] \geq \lim_{r \uparrow 1/2} \left(2r + \frac{r^2}{1 - 2r}\right) = \infty,$$

as required. \square

Proof of Theorem 4.4. The $s < 1/2$ part of the theorem follows from Lemma 7.2 and Lemma 6.4(b). On the other hand, for any $s \geq 1/2$ and any $r < 1/2$, $\mathbb{P}_N[C_t^N(s) \leq n] \leq \mathbb{P}_N[C_t^N(r) \leq n]$ so that, again by Lemmas 7.2 and 6.4(b),

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[C_t^N(s) \leq n] \leq \lim_{r \uparrow 1/2} \sum_{m=0}^n \pi^r(m) = 0,$$

by (4.6). \square

Proof of Theorem 4.5. Since $\mathbb{P}_N[X_t^{(n)} \leq s] = \mathbb{P}_N[C_t^N(s) \geq n]$, we have

$$\lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \mathbb{P}_N[X_t^{(n)} \leq s] = 1 - \lim_{N \rightarrow \infty} \lim_{t \rightarrow \infty} \sum_{m=0}^{n-1} \mathbb{P}_N[C_t^N(s) = m] = 1 - \sum_{m=0}^{n-1} \pi^s(m),$$

by Theorem 4.4. The result (4.9) now follows from (4.6) and some algebra. \square

8 Appendix: Markov chain limits

In relating the asymptotics of the Markov chains $C_t^N(s)$ to the ‘ $N = \infty$ ’ Markov chain $C_t(s)$, we need the following general result (Theorem 8.1) on a form of analyticity for families of Markov chains that are uniformly ergodic in a certain sense (but not the sense used in Chapter 16 of [26], where the uniformity is over all possible starting states of a single Markov chain; our ‘uniformity’ is over a family of Markov chains all starting at the same point). Theorem 8.1 is related to material in Chapters 6 and 7 of [11], although our setting is somewhat different and the proof we give uses different ideas. Our context differs from the set-up in [11], most notably in that our state-space changes with N , unlike in [11]. It is likely that the methods of [11] could be adapted to our setting. However, it is simpler to proceed directly; we use, in part, a coupling approach. Recall that a subset S of \mathbb{R} is locally finite if $S \cap R$ is finite for any bounded set R .

Theorem 8.1. *Fix $N_0 \in \mathbb{N}$. For each integer $N \geq N_0$ let Y_t^N be an irreducible, aperiodic Markov chain under P_N on S_N a countable subset of $[0, \infty)$, where $0 \in S_N$, $S_N \subseteq S_{N+1}$, and $\limsup_{N \rightarrow \infty} S_N = \infty$. Also suppose that Y_t is an irreducible, aperiodic Markov chain under P on $S := \cup_N S_N$. Suppose that S is locally finite. Write E_N and E for expectation under P_N and P respectively. Suppose that there exists $B < \infty$ such that for all $N \geq N_0$,*

$$P_N[|Y_{t+1}^N - Y_t^N| > B] = 0, \text{ and } P[|Y_{t+1} - Y_t| > B] = 0. \quad (8.1)$$

Suppose also that there exist $A_0 \in (0, \infty)$ and $\varepsilon_0 > 0$ for which

$$\begin{aligned} \sup_{N \geq N_0} \sup_{x \in S_N \cap [A_0, \infty)} E_N[Y_{t+1}^N - Y_t^N \mid Y_t^N = x] &\leq -\varepsilon_0; \\ \sup_{x \in S \cap [A_0, \infty)} E[Y_{t+1} - Y_t \mid Y_t = x] &\leq -\varepsilon_0. \end{aligned} \quad (8.2)$$

Let $q_N(x, y) := P_N[Y_{t+1}^N = y \mid Y_t^N = x]$ and $q(x, y) := P[Y_{t+1} = y \mid Y_t = x]$. Suppose that

$$\lim_{N \rightarrow \infty} [q_N(x, y) \mathbf{1}\{x \in S_N\}] = q(x, y), \quad (8.3)$$

for all $x, y \in S$. Then the following hold.

- (a) *The Markov chain Y_t is ergodic on S and, for any $N \geq N_0$, Y_t^N is ergodic on S_N . Let $\tau_N := \min\{t \in \mathbb{N} : Y_t^N = 0\}$ denote the time of the first return to 0 by the process Y_t^N ; similarly let $\tau := \min\{t \in \mathbb{N} : Y_t = 0\}$. There exists $\delta > 0$ such that*

$$E[e^{\delta\tau}] < \infty, \text{ and } \sup_{N \geq N_0} E_N[e^{\delta\tau_N}] < \infty. \quad (8.4)$$

(b) There exist stationary distributions ν_N on S_N and ν on S such that

$$\lim_{t \rightarrow \infty} P_N[Y_t^N = x] = \nu_N(x), \text{ and } \lim_{t \rightarrow \infty} P[Y_t = x] = \nu(x).$$

Moreover, there exist $c > 0$ and $C < \infty$ such that for all $N \geq N_0$ and all $x \in S_N$, $\nu_N(x) \leq Ce^{-cx}$ and, for all $x \in S$, $\nu(x) \leq Ce^{-cx}$.

(c) For any $x \in S$, $\lim_{N \rightarrow \infty} \nu_N(x) = \nu(x)$.

(d) Finally,

$$\lim_{t \rightarrow \infty} E_N[Y_t^N] = \sum_{x \in S_N} x \nu_N(x) < \infty, \text{ and } \lim_{t \rightarrow \infty} E[Y_t] = \sum_{x \in S} x \nu(x) < \infty.$$

Before getting into the details, we sketch the outline of the proof. The Foster-type condition (8.2) will enable us to conclude that the Markov chains have a uniform (in N) ergodicity property implying parts (a) and (b). We then couple Y_t and Y_t^N on an interval $[0, A]$ where A is chosen large enough so that the processes reach 0 before leaving $[0, A]$ with high probability. Given such an A , we choose N large enough so that on this finite interval (8.3) ensures that the two Markov chains can, with high probability, be coupled until the time that they reach 0. This strategy, which succeeds with high probability, ensures that the two processes follow identical paths over an entire excursion; using the excursion-representation of the stationary distributions will yield part (c).

An elementary but important consequence of the conditions of Theorem 8.1 is a ‘uniform irreducibility’ property that we will use repeatedly in the proof; we state this property in the following result. Note that the condition (8.3) is stronger than is necessary for parts (a) and (b) of Theorem 8.1: in the proof of Theorem 8.1 (a) and (b) below, we use only the uniform irreducibility property given in Lemma 8.1.

Lemma 8.1. *Under the conditions of Theorem 8.1, for any $A \in (0, \infty)$ there exist $\varepsilon_1 := \varepsilon_1(A) > 0$, $N_1(A) \in \mathbb{N}$, and $n_0(A) \in \mathbb{N}$ such that, for all $N \geq N_1(A)$ and all $x, y \in S_N \cap [0, A]$ there exists $n := n(x, y) \leq n_0(A)$ for which*

$$P_N[Y_n^N = y \mid Y_0^N = x] \geq \varepsilon_1. \quad (8.5)$$

Proof. Fix $A \in (0, \infty)$. Local finiteness implies that $S \cap [0, A]$ is finite. Take N large enough so that $S_N \cap [0, A] = S \cap [0, A]$. Irreducibility of Y_t implies that for any $x, y \in S$, there exists $n(x, y) < \infty$ such that $P[Y_{n(x, y)} = y \mid Y_0 = x] > 0$. There are only finitely many pairs $x, y \in S \cap [0, A]$, so for such x, y , in fact $P[Y_{n(x, y)} = y \mid Y_0 = x] \geq \varepsilon_2(A) > 0$ where $n(x, y) \leq n_0(A)$ and ε_2 and n_0 depend only on A , not on x, y . Moreover, for any $x, y \in S \cap [0, A]$, there are only finitely many paths of length at most $n_0(A)$ from x to y . It follows that for any $x, y \in [0, A]$ we can find a sequence of states of S , $x_0 = x, x_1, \dots, x_{n-1}, x_n = y$ with $n = n(x, y) \leq n_0(A)$ for which

$$P[Y_1 = x_1, \dots, Y_n = x_n \mid Y_0 = x_0] \geq \varepsilon_3(A) > 0.$$

However, by (8.3) we have that, as $N \rightarrow \infty$,

$$\begin{aligned} P_N[Y_1^N = x_1, \dots, Y_n^N = x_n \mid Y_0^N = x_0] &= q_N(x_0, x_1)q_N(x_1, x_2) \cdots q_N(x_{n-1}, x_n) \\ &\rightarrow q(x_0, x_1)q(x_1, x_2) \cdots q(x_{n-1}, x_n) = P[Y_1 = x_1, \dots, Y_n = x_n \mid Y_0 = x_0]. \end{aligned}$$

So for all N large enough, $P_N[Y_n^N = y \mid Y_0^N = x] \geq \varepsilon_3(A)/2$, say. \square

Now we move on to the proof of Theorem 8.1. The theorem and its proof are in parts closely related to existing results in the literature, in particular certain results from [2–4, 11, 18, 22, 25, 26, 30] amongst others. However, none of the existing results that we have seen fits exactly into the present context, and rather than try to adapt various parts of these existing results we give a largely self-contained proof. In the course of the proof we give some more details of how the arguments relate to existing results.

Proof of Theorem 8.1 (a) and (b). First we prove part (a). Since Y_t^N is an irreducible, aperiodic Markov chain on the finite or countably infinite state-space S_N , the drift condition (8.2) enables us to apply Foster’s criterion (see e.g. [11]) to conclude that Y_t^N is positive-recurrent (ergodic) and in particular, since (8.2) is uniform in N , $E_N[\tau_N]$ is uniformly bounded (independently of N).

In fact, we have the much stronger result (8.4). The exponential moments result (8.4) for a specific N is essentially a classical result, closely related to results in [11, 22, 26], for instance, and follows from the drift condition (8.2) together with the bounded jumps condition (8.1) and irreducibility: concretely, one may use, for example, Theorem 2.3 of [18] or the $a = 0$ case of Corollary 2 of [2]. The uniformity in (8.4) follows from the fact that (8.2) and (8.1) hold uniformly in N , and that the irreducibility is also uniform in the sense of Lemma 8.1. Indeed, the results of [2, 18] apply not to τ_N itself but to $\sigma_N := \min\{t \in \mathbb{Z}^+ : Y_t^N \leq A_0\}$ where A_0 is the constant in (8.2): standard arguments using the uniform irreducibility condition extend the uniform bound on $E_N[e^{\delta\sigma_N}]$ to the desired uniform bound on $E_N[e^{\delta\tau_N}]$. In particular, (8.4) implies that for any $k \in \mathbb{N}$ there exists $C_k < \infty$ such that

$$E[\tau^k] \leq C_k, \text{ and } \sup_{N \geq N_0} E_N[\tau_N^k] \leq C_k, \quad (8.6)$$

a fact that we will need later. This completes the proof of part (a).

By positive-recurrence, there exist (unique) stationary distributions ν_N on S_N and ν on S such that $\lim_{t \rightarrow \infty} P_N[Y_t^N = x] = \nu_N(x)$ and $\lim_{t \rightarrow \infty} P[Y_t = x] = \nu(x)$. Next we prove the uniform exponential decay of ν_N and ν . Again these results are closely related to existing results in the literature, such as those in [18, 25, 30], Chapters 6 and 7 of [11], Section 2.2 of [3], or Section 16.3 of [26].

For $\delta \in (0, 1)$, let $W_t := e^{\delta Y_t}$. We show that W_t has negative drift outside a finite interval, provided $\delta > 0$ is small enough. We have that

$$E[W_{t+1} - W_t \mid Y_t = x] = e^{\delta x} E[e^{\delta(Y_{t+1} - Y_t)} - 1 \mid Y_t = x].$$

Taylor’s theorem with Lagrange remainder implies that for all $y \in [-B, B]$ and all $\delta \in (0, 1)$, $e^{\delta y} - 1 \leq \delta y + K\delta^2$, where $K := K(B) < \infty$. Using this inequality and the bounded jumps assumption (8.1), we obtain

$$\begin{aligned} E[W_{t+1} - W_t \mid Y_t = x] &\leq \delta e^{\delta x} (E[Y_{t+1} - Y_t \mid Y_t = x] + K\delta) \\ &\leq \delta e^{\delta x} (-\varepsilon_0 + K\delta), \end{aligned}$$

when $x > A_0$, by (8.2). Hence, for $\delta := \delta(B, \varepsilon_0) \in (0, 1)$ sufficiently small, we have that

$$E[W_{t+1} - W_t \mid Y_t = x] < 0, \quad (8.7)$$

for $x > A_0$.

Let $\sigma := \min\{t \in \mathbb{Z}^+ : Y_t \leq A_0\}$ and $\nu_x := \min\{t \in \mathbb{Z}^+ : Y_t \geq x\}$. By irreducibility, $\sigma \wedge \nu_x < \infty$ a.s.. Moreover, by (8.7), $W_{t \wedge \sigma \wedge \nu_x}$ is a nonnegative supermartingale. Hence $W_{t \wedge \sigma \wedge \nu_x} \rightarrow W_{\sigma \wedge \nu_x}$ a.s. as $t \rightarrow \infty$, and

$$e^{\delta Y_0} = W_0 \geq E[W_{\sigma \wedge \nu_x}] \geq E[W_{\nu_x} \mathbf{1}\{\sigma > \nu_x\}] \geq e^{\delta x} P[\sigma > \nu_x].$$

The same argument holds for $W_t^N := e^{\delta Y_t^N}$, uniformly in $N \geq N_0$. Thus we have

$$P[\nu_x < \sigma \mid Y_0 = y] \leq e^{-\delta(x-y)}, \text{ and } P_N[\nu_{N,x} < \sigma_N \mid Y_0^N = y] \leq e^{-\delta(x-y)}, \quad (8.8)$$

where $\nu_{N,x} := \min\{t \in \mathbb{Z}^+ : Y_t^N \geq x\}$ and $\sigma_N := \min\{t \in \mathbb{Z}^+ : Y_t^N \leq A_0\}$.

We deduce from (8.8), with the uniform irreducibility property described in Lemma 8.1, that the probability of reaching $[x, \infty)$ before returning to 0 decays exponentially in x , uniformly in N . We will show that

$$P[\nu_x < \tau] \leq C e^{-\delta x}, \text{ and } P_N[\nu_{N,x} < \tau_N] \leq C e^{-\delta x}. \quad (8.9)$$

By uniform irreducibility (Lemma 8.1) and the bounded jumps assumption (8.1), we have that there exist $A_1 \in (A_0, \infty)$ and $\theta > 0$ for which

$$\inf_{y \in S \cap [0, A_0]} P[\tau < \nu_{A_1} \mid Y_0 = y] > \theta, \text{ and } \min_{N \geq N_0} \inf_{y \in S_N \cap [0, A_0]} P_N[\tau_N < \nu_{N,A_1} \mid Y_0^N = y] > \theta.$$

Together with (8.8), this will yield the result (8.9): the idea is that each time the process enters $[0, A_0]$, it has uniformly positive probability of reaching 0 before it exits $[0, A_1]$, otherwise, by (8.8), starting from $[A_1, A_1 + B]$ the process reaches $[x, \infty)$ before its next return to $[0, A]$ with an exponentially small probability, and (8.9) follows. We write out a more formal version of this idea for Y_t only; a similar argument holds for Y_t^N .

Let $\kappa_0 := 0$ and for $n \in \mathbb{Z}^+$ define iteratively the stopping times $\eta_n := \min\{t \geq \kappa_n : Y_t > A_1\}$ and $\kappa_{n+1} := \min\{t \geq \eta_n : Y_t \leq A_0\}$. By successively conditioning at these times (all of which are a.s. finite), we have

$$\begin{aligned} P[\nu_x < \tau] &\leq P[\nu_x < \kappa_1 \mid Y_{\eta_0}] + E[P[\nu_x < \tau \mid Y_{\kappa_1}] \mid Y_{\eta_0}] \\ &\leq P[\nu_x < \kappa_{n+1} \mid Y_{\eta_0}] + E[P[\eta_1 < \tau \mid Y_{\kappa_1}] E[P[\nu_x < \tau \mid Y_{\eta_1}] \mid Y_{\kappa_1}] \mid Y_{\eta_0}] \\ &\leq C e^{-\delta x} (1 + (1 - \theta) + (1 - \theta)^2 + \dots), \end{aligned}$$

since $P[\eta_n < \tau \mid Y_{\kappa_n}] \leq 1 - \theta$ a.s., and $P[\nu_x < \kappa_{n+1} \mid Y_{\eta_n}] \leq C e^{-\delta x}$ by (8.8) and the fact that $Y_{\eta_n} \leq A_1 + B$ a.s., by (8.1). Thus we verify (8.9).

Let $L_N(x)$ denote the total occupation time of state $x \in S_N$ by Y_t^N before time τ_N , i.e., during the first excursion of Y_t^N ; similarly for $L(x)$ with respect to Y_t . That is,

$$L_N(x) := \sum_{t=0}^{\tau_N-1} \mathbf{1}\{Y_t^N = x\}, \text{ and } L(x) := \sum_{t=0}^{\tau-1} \mathbf{1}\{Y_t = x\}.$$

Standard theory for irreducible, positive-recurrent Markov chains (see e.g. [1, Corollary I.3.6, p. 14]) gives

$$\nu_N(x) = \frac{E_N[L_N(x)]}{E_N[\tau_N]}, \text{ and } \nu(x) = \frac{E[L(x)]}{E[\tau]}. \quad (8.10)$$

By (8.9), the probability of visiting x during a single excursion decays exponentially. In order to bound the expected occupation time, we need an estimate for the probability of returning to x starting from x . We claim that there exists $\varepsilon_2 > 0$ for which

$$\max_{N \geq N_0} \max_{x \in S_N \cap [A_0, \infty)} P_N[\text{return to } x \text{ before hitting } 0 \mid Y_0^N = x] \leq 1 - \varepsilon_2, \quad (8.11)$$

and also $\max_{x \in S \cap [A_0, \infty)} P[\text{return to } x \text{ before hitting } 0 \mid Y_0 = x] \leq 1 - \varepsilon_2$. To verify (8.11), note that from (8.2) and (8.1) there exists $\varepsilon' > 0$ such that $P[Y_{t+1} - Y_t \leq -\varepsilon' \mid Y_t = x] \geq \varepsilon'$ for all $x \geq A_0$, and the same for Y_t^N (uniformly in N). Then (8.8) yields (8.11). It follows from (8.11) that, starting from x , the number of returns (before hitting 0) of Y_t or Y_t^N to x is stochastically dominated (uniformly in N and x) by a geometric random variable. In particular,

$$E_N[L_N(x)] \leq CP_N[\nu_{N,x} < \tau_N], \text{ and } E[L(x)] \leq CP[\nu_x < \tau],$$

which, with (8.10), yields the claimed tail bounds on ν_N and ν . \square

Proof of Theorem 8.1 (c) and (d). First we prove part (c). We will again use the representation (8.10). We use a coupling argument to show that, as $N \rightarrow \infty$, for any $x \in S$,

$$E_N[L_N(x)] \rightarrow E[L(x)], \text{ and } E_N[\tau_N] \rightarrow E[\tau]. \quad (8.12)$$

Let $\varepsilon > 0$. Take $A \in (0, \infty)$ large enough so that $BC_1/A < \varepsilon$, where C_1 is the constant in the $k = 1$ version of (8.6) and B is the bound in (8.1). Also, for convenience, choose A so that A/B is an integer. We claim that

$$\lim_{N \rightarrow \infty} \sup_{x \in S_N \cap [0, 2A]} \sum_{y \in S} |q_N(x, y) \mathbf{1}\{y \in S_N\} - q(x, y)| = 0. \quad (8.13)$$

To see this, note that since S (and hence also $S_N \subseteq S$) is locally finite, in the supremum in (8.13) x takes only finitely many values (uniformly in N), and, by (8.1), only finitely many terms in the sum are non-zero (again, uniformly in N). Hence by condition (8.3) we verify the claim (8.13). By (8.13), we can choose N large enough such that

$$\sup_{x \in S_N \cap [0, 2A]} \sum_{y \in S} |q_N(x, y) \mathbf{1}\{y \in S_N\} - q_\infty(x, y)| \leq \varepsilon B/A. \quad (8.14)$$

We couple Y_t^N and Y_t . We take N large enough so that $S_N \cap [0, 2A + B] = S \cap [0, 2A + B]$. We use the notation P_N^* for the probability measure on the space on which we are going to construct coupled instances of Y_t^N and Y_t , and write E_N^* for the corresponding expectation. We start at $Y_0^N = Y_0 \leq A$. We claim that one can construct (Y_t^N, Y_t) as a Markov chain under P_N^* so that $P_N^*[Y_t^N = y \mid Y_t^N = x] = q_N(x, y)$, $P_N^*[Y_t = y \mid Y_t = x] = q(x, y)$, and

$$P_N^*[(Y_t^N, Y_t) = (y, y) \mid (Y_t^N, Y_t) = (x, x)] \geq \min\{q_N(x, y), q(x, y)\}.$$

To see this, observe that when $Y_t^N = Y_t = x$ we may choose transition probabilities $r_x(y, z) = P_N^*[(Y_t^N, Y_t) = (y, z) \mid (Y_t^N, Y_t) = (x, x)]$ satisfying $r_x(y, y) = \min\{q_N(x, y), q(x, y)\}$, $\sum_{z \neq y} r_x(y, z) = q_N(x, y)$, and $\sum_{y \neq z} r_x(y, z) = q(x, z)$: these constraints can always be satisfied by some choice of r_x . If $Y_t^N \neq Y_t$, we define P_N^* by allowing the two processes to evolve independently.

Let $\sigma := \min\{t \in \mathbb{N} : Y_t^N \neq Y_t^\infty\}$ denote the time at which the processes first separate. For any $t \leq A/B$, we have from (8.1) that $\max\{Y_t^N, Y_t\} \leq 2A$ a.s., and together with the fact that the state-spaces of the two processes coincide on $[0, 2A + B]$, (8.14) implies that, for $t \leq A/B$, $P_N^*[\sigma > t + 1 \mid \sigma > t] \geq 1 - (B\varepsilon/A)$. Hence, for any $t \leq A/B$,

$$P_N^*[\sigma > t] \geq 1 - \left(\frac{tB\varepsilon}{A}\right). \quad (8.15)$$

Let E_t denote the event $E_t := \{\sigma > t\} \cap \{\tau_N \leq t\}$, i.e., that the paths of Y_t and Y_t^N coincide up until time t and visit zero by time t . Then, by (8.15),

$$P_N^*[E_{A/B}^c] \leq P_N^*[\sigma \leq A/B] + P_N^*[\tau_N > A/B] \leq \varepsilon + (BC_1/A) \leq 2\varepsilon, \quad (8.16)$$

using Markov's inequality and (8.6) to bound $P_N^*[\tau_N > A/B]$, and the choice of A to obtain the final inequality. On E_t , $\{\tau_N = \tau\}$, so that

$$\begin{aligned} E_N^*[|\tau_N - \tau|] &\leq E_N^*[|\tau_N - \tau| \mathbf{1}(E_{A/B}^c)] \\ &\leq (E_N^*[\tau_N^2] + E_N^*[\tau^2])^{1/2} (P_N^*[E_{A/B}^c])^{1/2}, \end{aligned}$$

by the Cauchy–Schwarz inequality. By (8.16) and the $k = 2$ case of (8.6), this last expression is bounded above by $\varepsilon^{1/2}$ times a constant not depending on N . Since $\varepsilon > 0$ was arbitrary, the second statement in (8.12) follows.

Similarly, on E_t , $\{L_N(x) = L(x)\}$ for any $x \in S$, so that

$$\begin{aligned} E_N^*[|L_N(x) - L(x)|] &\leq E_N^*[|L_N(x) - L(x)| \mathbf{1}(E_{A/B}^c)] \\ &\leq (E_N^*[\tau_N^2] + E_N^*[\tau^2])^{1/2} (P_N^*[E_{A/B}^c])^{1/2}, \end{aligned}$$

since $L_N(x) \leq \tau_N$ and $L(x) \leq \tau$ a.s.. Thus we obtain the first statement in (8.12). Combining the two statements in (8.12) with the representation in (8.10) we obtain $\nu_N(x) \rightarrow \nu(x)$ for any $x \in S$, completing the proof of part (c).

Finally we prove part (d). The convergence results follow from, for example, Theorem 2 of [30] once the integrability of the stationary distributions is established. But the fact that $\sum x \nu_N(x)$ and $\sum x \nu(x)$ are finite follows from the bounds in part (b). \square

Acknowledgements

AW is grateful to Edward Crane for suggesting the partial-order-driven model described in Section 5.

References

- [1] S. Asmussen, Applied Probability and Queues, 2nd edition, Springer-Verlag, New York, 2003.
- [2] S. Aspandiiarov and R. Iasnogorodski, General criteria of integrability of functions of passage-times for non-negative stochastic processes and their applications, *Theory Probab. Appl.* **43** (1999) 343–369; translated from *Teor. Veroyatnost. i Primenen.* **43** (1998) 509–539 (in Russian).
- [3] S. Aspandiiarov and R. Iasnogorodski, Asymptotic behaviour of stationary distributions for countable Markov chains, with some applications, *Bernoulli* **5** (1999) 535–569.
- [4] S. Aspandiiarov, R. Iasnogorodski, and M. Menshikov, Passage-time moments for nonnegative stochastic processes and an application to reflected random walks in a quadrant, *Ann. Probab.* **24** (1996) 932–960.

- [5] P. Bak and K. Sneppen, Punctuated equilibrium and criticality in a simple model of evolution, *Phys. Rev. Lett.* **71** (1993) 4083–4086.
- [6] K.L. Chung, Markov Chains with Stationary Transition Probabilities, 2nd edition, Springer-Verlag, Berlin, 1967.
- [7] J. de Boer, B. Derrida, H. Flyvbjerg, A.D. Jackson, and T. Wettig, Simple model of self-organized biological evolution, *Phys. Rev. Lett.* **73** (1994) 906–909.
- [8] J. de Boer, A.D. Jackson, and T. Wettig, Criticality in simple models of evolution, *Phys. Rev. E* **51** (1995) 1059–1074.
- [9] C.J. Burke and M. Rosenblatt, A Markovian function of a Markov chain, *Ann. Math. Statist.* **29** (1958) 1112–1122.
- [10] R. Durrett, Probability: Theory and Examples, Wadsworth & Brooks/Cole, Pacific Grove, CA, 1991.
- [11] G. Fayolle, V.A. Malyshev, and M.V. Menshikov, Topics in the Constructive Theory of Countable Markov Chains, Cambridge University Press, Cambridge, 1995.
- [12] H. Flyvbjerg, K. Sneppen, and P. Bak, Mean field theory for a simple model of evolution, *Phys. Rev. Lett.* **71** (1993) 4087–4090.
- [13] G.J.M. Garcia and R. Dickman, On the thresholds, probability densities, and critical exponents of Bak-Sneppen-like models, *Physica A* **342** (2004) 164–170.
- [14] A.J. Gillett, Phase Transitions in Bak–Sneppen Avalanches and in a Continuum Percolation Model, PhD thesis, Vrije Universiteit, Amsterdam, 2007.
- [15] A. Gillett, R. Meester, and M. Nuyens, Bounds for avalanche critical values of the Bak–Sneppen model, *Markov Process. Relat. Fields* **12** (2006) 679–694.
- [16] A. Gillett, R. Meester, and P. Van Der Wal, Maximal avalanches in the Bak–Sneppen model, *J. Appl. Probab.* **43** (2006) 840–851.
- [17] M. Grinfeld, P.A. Knight, and A.R. Wade, Bak–Sneppen type models and rank-driven processes. Preprint [arXiv:1011.1777](https://arxiv.org/abs/1011.1777) (2010).
- [18] B. Hajek, Hitting-time and occupation-time bounds implied by drift analysis with applications, *Adv. Appl. Probab.* **14** (1982) 502–525.
- [19] D.A. Head and G.J. Rodgers, The anisotropic Bak-Sneppen model, *J. Phys. A: Math. Gen.* **31** (1998) 3977–3988.
- [20] H.J. Jensen, Self-Organized Criticality, Cambridge University Press, Cambridge, 1998.
- [21] G.L. Labzowsky and Yu.M. Pis'mak, Exact analytical results for the Bak–Sneppen model with arbitrary number of randomly interacting species, *Phys. Lett. A* **246** (1998) 377–383.
- [22] J. Lamperti, Criteria for stochastic processes II: passage-time moments, *J. Math. Anal. Appl.* **7** (1963) 127–145.

- [23] R. Meester and D. Znamenski, Limit behavior of the Bak–Sneppen evolution model, *Ann. Probab.* **31** (2003) 1986–2002.
- [24] R. Meester and D. Znamenski, Critical thresholds and the limit distribution in the Bak–Sneppen model, *Commun. Math. Phys.* **246** (2004) 63–86.
- [25] M.V. Menshikov and S.Yu. Popov, Exact power estimates for countable Markov chains, *Markov Process. Relat. Fields* **1** (1995) 57–78.
- [26] S. Meyn and R.L. Tweedie, Markov Chains and Stochastic Stability, 2nd ed., Cambridge University Press, 2009.
- [27] H. Moulin, Game Theory for the Social Sciences, 2nd ed., New York University Press, New York, 1986.
- [28] R. Pemantle, A survey of random processes with reinforcement, *Probab. Surv.* **4** (2007) 1–79.
- [29] Yu.M. Pis'mak, Exact solution of master equations for a simple model of self-organized biological evolution, *J. Phys. A: Math. Gen.* **28** (1995) 3109–3115.
- [30] R.L. Tweedie, The existence of moments for stationary Markov chains, *J. Appl. Probab.* **20** (1983) 191–196.