

The onset of the North Atlantic Igneous Province in a rifting perspective

J. HANSEN*†, D. A. JERRAM*, K. McCAFFREY* & S. R. PASSEY‡

*Department of Earth Sciences, Durham University, South Road, DH1 3LE, Durham, United Kingdom

‡Jarðfeingi (Faroe Earth and Energy Directorate), Brekkutún 1, P.O. Box 3059, FO-110, Tórshavn, Faroe Islands

(Received 25 March 2008; accepted 17 February 2009; First published online 25 March 2009)

Abstract – The processes that led to the onset and evolution of the North Atlantic Igneous Province (NAIP) have been a theme of debate in the past decades. A popular theory has been that the impingement on the lower lithosphere of a hot mantle plume (the ‘Ancestral Iceland’ plume) initiated the first voluminous outbursts of lava and initiated rifting in the North Atlantic area in Early Palaeogene times. Here we review previous studies in order to set the NAIP magmatism in a time–space context. We suggest that global plate reorganizations and lithospheric extension across old orogenic fronts and/or suture zones, aided by other processes in the mantle (e.g. local or regional scale upwellings prior to and during the final Early Eocene rifting), played a role in the generation of the igneous products recorded in the NAIP for this period. These events gave rise to the extensive Paleocene and Eocene igneous rocks in W Greenland, NW Britain and at the conjugate E Greenland–NW European margins. Many of the relatively large magmatic centres of the NAIP were associated with transient and geographically confined doming in Early Paleocene times prior to the final break-up of the North Atlantic area.

Keywords: flood basalts, rifting, orogenic sutures, North Atlantic.

1. Introduction

The North Atlantic Igneous Province (NAIP) is a classic Large Igneous Province associated with a volcanic rifted margin. It has traditionally been considered to comprise the voluminous Palaeogene igneous rocks occurring at the conjugate E Greenland–NW European margins and in the W Greenland–Baffin Bay area (Upton, 1988; Saunders *et al.* 1997; Meyer, Van Wijk & Gernigon, 2007 and references therein). Other contemporaneous magmatism occurred in the northernmost parts of Greenland (Kap Washington Group at $c. 64 \pm 3$ Ma, (Estrada, Höhndorf & Henjes-Kunst, 2001) and in the W Barents Sea (Vestbakken Volcanic Province, $c. 54$ Ma, (Tsikalas, Eldholm & Faleide, 2002) (Fig. 1).

The major Early Palaeogene NAIP rocks can be regionally divided into: W Greenland–Baffin Island, SE Greenland, (central–east) CE Greenland, NE Greenland, Vøring margin, Møre margin, Faroe Islands, Rockall–Hatton area, Faroe–Shetland Basin, Rockall Trough and the NW British Isles (Saunders *et al.* 1997) (Fig. 1). Other contemporaneous, smaller and more isolated parts of the NAIP are also shown in Figure 1. The CE Greenland–Faroe Islands Ridge and Iceland formed subsequent to the onset of sea-floor spreading in the area (Meyer, Van Wijk & Gernigon, 2007). Exposed and submerged basaltic rocks of the NAIP extend roughly NE–SW for more than 2000 km along the conjugate East Greenland–NW European margins (Saunders *et al.* 1997) (Fig. 1).

The extrusive rocks of the province cover a surface area of at least $\sim 1.3 \times 10^6$ km², while the extrusive and intrusive rocks of the NAIP are together estimated to comprise a volume of $\sim 6.6 \times 10^6$ km³ (Eldholm & Grue, 1994). The majority of the extrusive rocks at the conjugate E Greenland–NW European margins (the Faroe Islands; Rockall–Hatton and Vøring–Møre) were extruded in subaerial or shallow-marine environments onto continental crust (e.g. Natland & Winterer, 2005). Similarly, the vast majority of the Early Palaeogene W Greenland igneous products were emplaced in continental crust (e.g. Larsen *et al.* 1999a) in a subaerial and/or in a shallow marine environment (Storey *et al.* 1998).

Here we use published studies to show that the formation of the NAIP could have been aided by the combined actions of a number of magmatic centres, whose initial actions in part were governed by regional and/or provincial plate tectonic reorganizations.

2. Geological setting prior to and during magmatism

In the context of a large igneous province such as the NAIP, it is pertinent to consider relevant tectonic events prior to the onset of magmatism and to consider possible temporal and spatial links between the igneous products.

2.a. Tectonic settings

Following the closure of the Iapetus Ocean and the collapse of the Caledonian Orogen in Silurian–Devonian times (Roberts, 2003), the proto-North

†Author for correspondence: jogvan.hansen@durham.ac.uk; jogvanha@post.olivant.fo




Figure 1. Simplified geological map of the North Atlantic Igneous Province and surrounding areas modified from Saunders *et al.* (1997); Nielsen, Larsen & Hopper (2002) and Nielsen, Stephenson & Thomson (2007). The ages are from the references in Table 1. The Early Palaeogene transient regional uplifts are from references in Table 2. The central igneous complexes and/or seamounts are modified from Bull & Masson (1996); Ritchie, Hitchen & Edwards (1997); Naylor *et al.* (1999); Edwards (2002); Hitchen (2004); Archer *et al.* (2005). Abbreviations: BJ – Bjørnøya; BJH – Bjørnøya High; CGFZ – Charlie Gibbs Fracture Zone; EEC – East Erlend Complex; FTF – Faroe Transform Fault; GFZ – Greenland Fracture Zone; JM – Jan Mayen; JMFZ – Jan Mayen Fracture Zone; JMR – Jan Mayen Ridge; MTFC – Møre-Trøndelag Fault Complex; RB – Rosemary Bank; SFZ – Senja Fracture Zone; UFZ – Ungava Fracture Zone; VV – Vestbrona Volcanic rocks; VVP – Vestbakken Volcanic Province; WEC – West Erlend Complex.

Atlantic area to the south of the Caledonian front (Figs 2, 3) was underlain by a patchwork of Archaean and Proterozoic terranes (Dickin, 1992).

Permian and Triassic broadly E–W-directed extension between Eurasia and Greenland resulted in the formation of numerous large half-graben basins widely distributed at the margins (Ziegler, 1989; Brekke *et al.* 1999; Doré *et al.* 1999; Surlyk, 1990). Jurassic E–W extension between Eurasia and North America–Greenland gave way to a dominantly NW–SE-directed extension in Early to Middle Cretaceous times (Doré *et al.* 1999). The Middle Cretaceous extension resulted in northwards-narrowing sea-floor spreading from the Rockall Trough to the Vøring Basin off W Norway (Price & Rattey, 1984). Renewed NW–SE-directed extension occurred in the proto-NAIP area from the Late Cretaceous to the Early Palaeogene (Doré *et al.* 1999). This event led to the initiation of sea-floor spreading in the Labrador Sea in the Early Paleocene (Chalmers & Laursen, 1995) and the initiation and northwards propagation of sea-floor spreading at the conjugate NW European–E Greenland margins in Early Eocene times (Ziegler, 1989, 1992; Doré *et al.* 1999). At around the same time, the Eurasian Basin began to open (Ziegler, 1989). Early Palaeogene exploitation/reactivation of Precambrian and Caledonian fault zones are inferred for the early phases of continental rifting and NAIP formation (Doré *et al.* 1997, 1999).

Regional vertical movements and the formation of transient domal uplifts preceded the main phases of Early Palaeogene igneous activities in many parts of the NAIP (e.g. MacLennan & Jones, 2006; Meyer, Van Wijk & Gernigon, 2007; Saunders *et al.* 2007). Published examples of some regional domal uplifts are listed in Table 2 (Fig. 2). Without constraining the depth of origin, Saunders *et al.* (2007) suggested that the ascent of narrow hot mantle jets and broadly contemporaneous rifting in areas of uplifts generated doming.

2.b. Igneous settings

While the igneous rocks of the NAIP cover a compositional spectrum from picrites to silicic rocks (Table 1), most of the rocks encountered in the province today are of basaltic composition (e.g. Saunders *et al.* 1997). Crustally contaminated rocks occur at or close to the base of volcanic successions in many parts of the basaltic sequences of the province (Gibson, 2002). The igneous products include both fissure or point-source fed lava-flows (Peate, Larsen & Lesher, 2003; Single & Jerram, 2004; Passey & Bell, 2007), water-influenced and water-lain volcanic successions (e.g. Peate, Larsen & Lesher, 2003; Jerram *et al.* 2009, this issue) and ignimbrites as well as plutonic or sheet intrusions (sills and/or dykes) (Table 1; Fig. 1), each reflecting the processes and crustal environment that prevailed in that particular area during melt emplacement. Most of the igneous activity of the

NAIP occurred in the time span from *c.* 62 to *c.* 53 Ma. Two main periods of melt emplacement have been inferred for the NAIP, with ages of *c.* 62 to 58 Ma and *c.* 57 to 53 Ma, and detectable peaks at *c.* 60 Ma and at *c.* 55 Ma, respectively (Saunders *et al.* 1997; Torsvik, Mosar & Eide, 2001; Jerram & Widdowson, 2005; Meyer, Van Wijk & Gernigon, 2007) (Table 1; Fig. 1). Smaller-scale igneous activity preceded these main periods in, for example, the N Rockall Trough (Morton *et al.* 1995; O'Connor *et al.* 2000), continued subsequently in parts of the NAIP area for tens of millions of years (e.g. Tegner & Duncan, 1999; O'Connor *et al.* 2000; Tegner *et al.* 2008), and is continuing on Iceland and on the island of Jan Mayen (e.g. Trønnes *et al.* 1999) (Table 1; Fig. 1).

3. The spatial distribution of known and inferred magmatic centres of the NAIP

It is noticeable that many of the inferred earliest igneous activities in the NAIP coincide well with some of the transient uplifts recorded for this period (Figs 1, 2; Tables 1, 2). Furthermore, it appears that a number of the uplifted regions and the parts of the NAIP with the most voluminous igneous production for this period, namely in the NW British Isles, the Faroe Islands, (central-east) CE Greenland and the Disko region in W Greenland (Upton, 1988; Saunders *et al.* 1997; Meyer, Van Wijk & Gernigon, 2007), were emplaced in the vicinity of old orogenic sutures and/or fronts from the Palaeozoic Caledonian Orogen, at suture zones between Archaean–Proterozoic terranes at the conjugate East Greenland–NW European margins and at the suture zone between the Archaean Nagssugtoqidian–Rinkian Orogen in the Disko–Baffin Island area (Figs 1, 2, 3).

In a reconstructed map of the NAIP region intended to show the spatial distribution of the igneous activities for the Middle Paleocene (Fig. 2; Table 1), the magmatic regions and/or centres at the conjugate E Greenland–NW European margins seem to form conspicuous double zigzag and roughly NNE–SSW-directed trends, from just to the north of Hold With Hope and southwards to the Ammassalik area along the E Greenland margin and from the Vøring area and southwards to the NW British Isles area at the NW European margin, converging at the CE Greenland–Faroe Islands area. According to current published data the igneous activities in N and W Greenland were spatially isolated from these events.

The suggested trends of igneous activities at the conjugate E Greenland–NW European margin from the Middle Paleocene seem to be more or less repeated in the Early Eocene (Fig. 3; Table 1), apart from the westward relocation of magmatism at the Vøring margin, the eastward relocation of magmatism at the Blossville Kyst and the establishment of volcanism in the W Barents Sea. Final sea-floor spreading at the




Figure 2. Simplified map of the NAIP at around 62 to 58 Ma modified from Torsvik *et al.* (2001) and Torsvik, Mosar & Eide (2001). The inferred locations of the Caledonian fronts and the Iapetus suture zone are from: Bott (1987); Soper *et al.* (1992); Ziegler (1992); Masson, Hauser & Jacob (1999); Skogseid *et al.* (2000); Hansen & Brooks (2002); Roberts (2003); Foulger, Natland & Anderson (2005a,b); Cocks (2005). The inferred Archaean-Proterozoic suture zone in the Rockall-Hatton-NW Britain area is modified from Dickin (1992). The inferred Nagssuqtocidian-Rinkian suture zone in the Disko region is modified from Krawiec (A. Krawiec, unpub. M.S. thesis, Univ. Texas Austin, 2003) and Connelly *et al.* (2006). The three major sinistral transforms are modified from Nielsen, Stephenson & Thomsen (2007). Broadly NW-trending lineaments at the NW European margin are modified from Kimbell *et al.* (2005). General spreading directions are from Harrison *et al.* (1999) and Nielsen, Stephenson & Thomsen (2007). Abbreviations: ADL – Anton Dohrn Lineament; CGFZ – Charlie Gibbs Fracture Zone; CL – Claire Lineament; JL – Judd Lineament; JML – Jan Mayen Lineament; FI – Faroe Islands; MF – Moray Firth; MFL – Marflo Lineament; MTFC – Møre-Trøndelag Fault Complex; PB – Porcupine Basin; SHL – South Hatton Lineament; SI – Shetland Islands; UFZ – Ungave Fault Zone; WTR – Wyville-Thomson Ridge; YR – Ymir Ridge. See text for further explanation.

Table 1. Summary of Early Palaeogene ages for the key regions of the North Atlantic Igneous Province (most studies are overlapping and fall within the time frames from *c.* 62 Ma to *c.* 53 Ma)

Regions: average emplacement ages	Rock compositions: modes of emplacements	Published examples
British Isles: <i>c.</i> 61 to <i>c.</i> 55 Ma	Ultramafic, mafic, silicic: volcanic, plutons, sills, dykes	Gamble, Wysoczanski & Meigham (1999); Chambers, Pringle & Parrish (2005); Storey, Duncan & Tegner (2007)
Rockall-Hatton margin: <i>c.</i> 58 to <i>c.</i> 53 Ma	Mafic, silicic: volcanic, plutons, sills, dykes	Sinton & Duncan (1998); Hitchen (2004)
Rockall Trough: <i>c.</i> 70 to <i>c.</i> 54 Ma	Mafic, silicic: volcanic, sills	Hitchen & Ritchie (1993); Morton <i>et al.</i> (1995); Sinton, Hitchen & Duncan (1998); O'Connor <i>et al.</i> (2000); Archer <i>et al.</i> (2005)
Faroe-Shetland Basin: <i>c.</i> 61 to <i>c.</i> 55 Ma	Mafic, silicic: volcanic, sills, dykes	Hitchen & Ritchie (1993); Trude <i>et al.</i> (2003)
Vøring margin: <i>c.</i> 61 to <i>c.</i> 55 Ma	Mafic, silicic: volcanic, sills, dykes	Skogseid <i>et al.</i> (1992); Sinton, Hitchen & Duncan (1998); Planke <i>et al.</i> (2005)
Møre Margin: <i>c.</i> 56 to <i>c.</i> 55 Ma	Mafic: sills	Planke <i>et al.</i> (2005)
Faroe Islands: <i>c.</i> 61 to <i>c.</i> 55 Ma	Ultramafic, mafic: volcanic, sills, dykes	Waagstein, Guise & Rex (2002); Storey, Duncan & Tegner (2007)
NE Greenland: <i>c.</i> 59 Ma to <i>c.</i> 53 Ma	Ultramafic, mafic: volcanic, sills, dykes	Upton <i>et al.</i> (1995); Price <i>et al.</i> (1997)
CE Greenland: <i>c.</i> 61 to <i>c.</i> 53 Ma	Ultramafic, mafic, silicic: volcanic, plutons, sills, dykes	Karson <i>et al.</i> (1998); Tegner <i>et al.</i> (1998); Hald & Tegner (2000); Lenoir, Féraud & Geoffroy (2003); Peate <i>et al.</i> (2003); Storey, Duncan & Tegner (2007)
SE Greenland: <i>c.</i> 62 Ma to <i>c.</i> 55 Ma	Ultramafic, mafic, silicic: volcanic, sills, dykes	Sinton & Duncan (1998); Sinton, Hitchen & Duncan (1998); Tegner & Duncan (1999); Storey, Duncan & Tegner (2007)
W Greenland: <i>c.</i> 61 Ma to <i>c.</i> 54 Ma	Ultramafic, mafic, silicic: volcanic, dykes	Storey <i>et al.</i> (1998); Larsen <i>et al.</i> (1999a); Geoffroy <i>et al.</i> (2001)
N Greenland: <i>c.</i> 64 Ma	Mafic, silicic: volcanic, dykes	Estrada, Höhndorf & Henjes-Kunst (2001)
Bjørnøya Marginal High: <i>c.</i> 54 Ma	Mafic: volcanic	Tsikalas, Eldholm & Faleide (2002)
Vestbrona, off SW Norway: <i>c.</i> 55 Ma	Mafic: volcanic	Bugge, Prestvik & Rokoengen (1980)

The ages presented in this table reflect only the initial main phases of NAIP magmatism. Subsequent magmatism occurred in many of the same regions as those presented in this table. For further information see e.g. Tegner *et al.* (2008), Morton *et al.* 1995 and O'Connor *et al.* (2000) and references in these papers.

Table 2. Early Palaeogene transient uplifts reported for regions within the North Atlantic Igneous Province

Regional locations	Cited example
Disko area, W Greenland	Japsen, Green & Chalmers (2005)
Ammassalik area, SE Greenland	Clift, Turner & ODP Leg 152 Scientific Party (1995); Larsen & Saunders (1998)
Kangerlussuaq area, CE Greenland	Peate, Larsen & Leshar (2003)
Scoresby Sund area, CE Greenland	Mathiesen, Bidstrup & Christiansen (2000)
Hold With Hope, NE Greenland	Thomson <i>et al.</i> (1999)
Vøring margin, off Norway	Ren <i>et al.</i> (2003)
Møre margin, off Norway	Brekke <i>et al.</i> (1999)
North N Sea Basin	Nadin, Kuszniir & Cheadle (1997)
Faroe-Shetland Basin	Nadin, Kuszniir & Cheadle (1997); Rudge <i>et al.</i> (2008)
North Rockall Trough	Archer <i>et al.</i> (2005)
Moray Firth to Shetland	Mackay <i>et al.</i> (2005); Rudge <i>et al.</i> (2008)
NW British Isles (Scotland)	Green <i>et al.</i> (1993); Mudge & Jones (2004)
Irish Sea	Cope (1994)
Porcupine Basin	Jones, White & Lovell (2001)

Bergerat & Angelier, 1996). In accepting the presence of a junction in this region, a broadly SE-trending failed rift arm or leaky transform with NE–SW to ENE–WSW-directed extension fits roughly with the observed orientation of dykes (NW–SE to NNW–SSE-directed) emplaced in NW Britain during this period (Speight *et al.* 1982; England, 1988; Geoffroy, Bergerat & Angelier, 1996). The later Eocene extension in NW Britain has been interpreted to result from broadly NW–SE-directed crustal extension associated with the opening of the North Atlantic (Geoffroy, Bergerat & Angelier, 1996). The Early Palaeogene magmatism in NW Britain has been associated with melting of the ‘Iceland Plume’ (Kent & Fitton, 2000; Upton *et al.* 2002), although Nadin, Kuszniir & Cheadle (1997)

tentatively suggested that a separate distinct mantle plume may have been active in the NW Britain area during this period. Tectonic activity has also been invoked by some authors to have facilitated melt generation in the area (Upton *et al.* 2002; Chambers, Pringle & Parrish, 2005).

3.b. The Faroe Islands–N Rockall Trough (Fig. 1; ‘II’ in Figs 2, 3)

Geoffroy, Bergerat & Angelier (1994) suggested that an Early Palaeogene triple junction was more or less centred on the Faroe Islands, while Burke & Dewey (1973) proposed a contemporaneous triple junction in the Faroes–N Rockall Trough area with magmatic

centres in the N Rockall Trough and to the SSW and/or SW off the Faroe Islands. This accords with the inferences by Waagstein (1988) regarding the depocentre of the Faroe Islands Beinivørð Formation (formerly lower basalt series: e.g. Passey & Bell, 2007) being located in the southern or central part of the Faroe Islands area. A NW-trending failed rift arm or leaky transform from this inferred junction(s) or from a junction that migrated within this region during Paleocene and Eocene times with relative extension directed towards the NE–SW and another rift arm or leaky transform with extension towards the NNW–SSE may explain the NW–SE and ENE–WSW sub-parallel igneous emplacement trends of contemporaneous central igneous complexes in the SW parts of the area (e.g. Archer *et al.* 2005), as well as NW-trending lineaments reported for this region (e.g. Johnson *et al.* 2005; Kimbell *et al.* 2005). A hypothetical connection between a Faroese and a NW British triple junction (Figs 2, 3) would presumably have been sub-parallel to the N–S-trending contemporaneous dykes in mainland Scotland to the E and S of Skye (e.g. Speight *et al.* 1982). Morton *et al.* (1995) tentatively suggested that volcanism at the Rosemary Bank (Fig. 1) in the N Rockall Trough was due to a separate underlying source, and Hitchen *et al.* (1997) likewise suggested a local source for the Early Palaeogene rocks in the area. Other authors have associated the Early Palaeogene magmatism in this area with the ‘Iceland plume’ (Holm, Hald & Waagstein, 2001; Archer *et al.* 2005).

3.c. The NE Faroe–Shetland Basin; N North Sea; offshore W Norway (Fig. 1; ‘III’ in Figs 2, 3)

An Early Palaeogene triple junction to the NNE off Shetland has been suggested by Burke & Dewey (1973), and an additional contemporaneous magmatic centre was active further to the NNE off the SW Norwegian coast. Based on reported Early Palaeogene uplifts and igneous activity in the area, Mudge & Jones (2004) and Rudge *et al.* (2008) suggested that the ‘Iceland Plume’ could be responsible for contemporaneous uplifts recorded in the northern North Sea and the NE Faroe–Shetland Basin area. Kanaris-Sotiriou, Morton & Taylor (1993) interpreted the Early Palaeogene basaltic and associated intermediate volcanic rocks of the Erlend Complex in the northern North Sea to be a result of extensional volcanism in the area. The Møre–Trøndelag Fault Complex, which extends offshore from the SW Norwegian coast, trends towards the area of this inferred junction and is thought to have been active in Early Palaeogene times (Doré *et al.* 1997; Redfield *et al.* 2004) and could have been linked to the contemporaneous igneous activity in the area. Torske & Prestvik (1991) tentatively suggested that the igneous products recorded off W Norway were related to an Early Palaeogene precursor to the subsequent Jan Mayen Fracture Zone (Fig. 1).

3.d. The Vøring margin; NE Greenland (Fig. 1; ‘IV’ in Figs 2, 3)

The inferred magmatic activity at the Vøring margin in Early Palaeogene times occurred at some distance from the future break-up zone in the region, but moved westwards with time (Eldholm, Thiede & Taylor, 1989). Early Eocene magmatism in NE Greenland and at the Vøring margin had a close spatial relationship (Viereck *et al.* 1988; Upton *et al.* 1995) and recent studies reveal a continuous Early Eocene igneous complex that directly linked these two regions together in the early stages of sea-floor spreading (Olesen *et al.* 2007). The igneous activities (volume and rock types) associated with these centres resemble those found in some places in the Rockall Trough (Upton, 1988) and on the NW British Isles (Viereck *et al.* 1988 and references therein). Volumes of the Paleocene to Early Eocene volcanism decreased from the central Vøring margin towards the south and north, respectively (Berndt *et al.* 2001), indicating melt supplies from a relatively confined magmatic source. An ‘Iceland Plume’ origin has been inferred for the NE Greenland magmatism (e.g. Upton *et al.* 1995) and for the igneous products at the Vøring margin by some authors (Skogseid *et al.* 1992). Conversely, Eldholm, Thiede & Taylor (1989) and Van Wijk *et al.* (2001) suggested that decompression melting triggered by rifting caused the magma generation at the Vøring margin. Recent re-interpretations of available magnetic, bathymetric, gravity and seismic data from the Vøring margin strongly suggest local Eocene magmatism related to an Azores-type triple junction linked to the embryonic stages of sea-floor spreading in the Norwegian–Greenland Sea (Gernigon *et al.* 2008).

3.e. The (central-east) CE Greenland (Fig. 1; ‘V’ in Figs 2, 3)

The voluminous and widespread igneous products in this region were probably the result of several contemporaneous magmatic centres (e.g. Callot, Geoffroy & Brun, 2002). The locations of hypothetical triple junctions at the CE Greenland margin have been estimated from Early Palaeogene magmatism and uplifts in the area (Larsen & Watt, 1985; Nielsen, 1987; Mathiesen, Bidstrup & Christiansen, 2000; Callot, Geoffroy & Brun, 2002; Peate, Larsen & Leshner, 2003) and from triple junction localities as suggested by Burke & Dewey (1973); Karson & Brooks (1999) and Tegner *et al.* (2008). This vast area was characterized by Early Palaeogene episodic igneous activity and frequent migration of magmatic centres (Larsen & Watt, 1985; Peate, Larsen & Leshner, 2003), and at least three separate rifting events have been recorded for this region, some of which occurred far inland (Nielsen, 1987; Olesen *et al.* 2007). The rifting associated with the bulk of the magmatism in CE Greenland and the Faroe Islands approximately at anomaly 24 (*c.* 55 Ma)

occurred close to the Blosseville Kyst (Larsen & Watt, 1985; Nielsen, 1987; Larsen *et al.* 1999b). This is in accordance with inferences that the magmas of the younger basalt formations of the then neighbouring Faroe Islands were supplied from the north during this period (Waagstein, 1988; Larsen *et al.* 1999b). The onset of final sea-floor spreading at the Blosseville Kyst latitude occurred further to the east along the now extinct Ægir Ridge at *c.* 54 Ma (Bott, 1985). On the one hand, Larsen & Marcussen (1992) and Hanghøj, Storey & Stecher (2003) considered the magmatism in CE Greenland to be related to extension in the area; on the other hand, authors such as Tegner *et al.* (2008 and references therein) suggested that the CE Greenland igneous products resulted from actions of the ‘Iceland Plume’.

3.f. The Hatton–Edoras margin; SE Greenland (Fig. 1; ‘VI’ in Figs 2, 3)

Only parts of this extensive area have been investigated in detail, but the close proximity in the Early Palaeogene suggest that these two margins perhaps shared some magmatic centres prior to the sea-floor spreading in the region (Figs 2, 3). Locations for some possible triple junctions in this region in Early Palaeogene times have been implied previously by Burke & Dewey (1973), Bull & Masson (1996), Karson & Brooks (1999), Nielsen, Larsen & Hopper (2002) and Nielsen, Stephenson & Thomsen (2007), and locations of some separate large magmatic centres and domal uplifts have been recorded by Morgan & Barton (1990), Barton & White (1997), Larsen & Saunders (1998) and Elliot & Parson (2008). A hypothetical SE-trending failed rift arm or transform from a triple junction in the southern parts of the Hatton Bank (Figs 2, 3) would be sub-parallel to the South Hatton Lineament (Johnson *et al.* 2005; Kimbell *et al.* 2005). Another major lineament intersecting the Hatton margin is the Anton Dohrn Lineament, which in part has been interpreted by Dickin (1992) to include an ancient orogenic suture zone (Figs 2, 3). Morgan & Barton (1990) detected a large separate Early Palaeogene magmatic centre on the NW Hatton Bank, and recent work by Elliot & Parson (2008) revealed that the Hatton rifted margin could be divided into three separate segments, each with a distinctive magmatic evolution. They tentatively suggested that the northern parts of the Hatton margin only experienced diffuse spreading in the Early Palaeogene prior to Chron 21 (*c.* 50 Ma) when regular coherent spreading was established. The phenomenon of diffuse sea-floor spreading has been inferred to reflect low obliquity rifting in a magmatically starved environment (e.g. Corti *et al.* 2001). In the southernmost parts of this margin, Elliot & Parson (2008) recorded relatively concentrated syn- to post-break-up volcanism. Most authors infer the ‘Iceland Plume’ to be the main source for the magmatism in these two margins (Barton & White, 1997; Fitton *et al.* 2000), but Edwards (2002)

considered any ‘Iceland Plume’-dominated processes further eastwards toward the Rockall–Hatton Basin to be problematic, and Barton & White (1997) suggested that there was no major long-distance lateral migration of the melts supplying the magmatism at the Edoras Bank.

3.g. The West Greenland–Baffin Island area (Fig. 1; ‘VII’ in Figs 2, 3)

Based on reported locations for large concentrations of Early Palaeogene igneous products (e.g. Chalmers, Larsen & Pedersen, 1995; Skaarup, Jackson & Oakey, 2006), large contemporaneous igneous centres (Callot, Geoffroy & Brun, 2002), doming (Japsen, Green & Chalmers (2005) and the trends of major faults thought to have been active in the same period (Chalmers, Larsen & Pedersen, 1995; Geoffroy *et al.* 2001; Callot, Geoffroy, Brun, 2002; Skaarup, Jackson & Oakey, 2006), the location of a hypothetical triple junction at the southern tip of the Ungava Fault System and another at Ubekendt Ejland around 100 km north of Disko seems to be reasonable. Another triple junction or kink between major faults further to the north between Baffin Island, Ellesmere Island and W Greenland reconstructed back at *c.* 60 Ma has been interpreted to have been active during the same period (Burke & Dewey, 1973; Torsvik *et al.* 2001; Nielsen, Stephenson & Thomsen, 2007). Gill, Holm & Nielsen (1995) associated a presumed high-temperature melting required for the generation of Early Palaeogene picrites in this region with a separate ‘Baffin Bay Plume’ rather than with a distant asymmetrical/irregular ‘Iceland Plume’ as suggested by Chalmers (1997) and Storey *et al.* (1998), among others. The generation of Eocene dykes in SW Greenland and the volcanism along the Ungava Fault System are supposed to have been facilitated by plate reorganizations in the area during that period (Storey *et al.* 1998; Larsen *et al.* 1999a; Skaarup & Pulvertaft, 2007 and references therein).

3.h. N Greenland (Fig. 1; ‘VIII’ in Fig. 2)

The Early Paleocene Kap Washington Group is thought to have been generated in response to continental rifting related to the break-up of the Laurasian plate (Estrada, Höhdorf & Henjes-Kunst, 2001). A contemporaneous triple junction off Kap Washington is in accordance with the study of Torsvik *et al.* (2001); Torsvik, Mosar & Eide (2001) and Nielsen, Stephenson & Thomsen (2007).

3.i. The W Barents region (Fig. 1; ‘IX’ in Fig. 3)

Volcanic rocks in the W Barents Sea (Vestbakken Volcanic Province) located at the inferred trace of the Caledonian suture zone are interpreted to have formed in response to Early Eocene transtension associated

with plate reorganizations in the area (Tsikalas, Eldholm & Faleide, 2002).

4. Discussion

4.a. Competing theories on the NAIP petrogenesis

A number of theories and geodynamic models (including mantle processes such as delamination, orogenic collapse, small-scale edge-driven convection, melting of fertile mantle, and melting of an individual large mantle plume) have previously been proposed to have caused the Early Palaeogene magmatism of the NAIP (e.g. Meyer, Van Wijk & Gernigon, 2007 and references therein). Other mantle processes sometimes thought to result in voluminous magmatism in general include decompression melting in response to global-scale extension (Ziegler, 1992), and melting resulting from spontaneous upwellings of near-solidus buoyant mantle material (Raddick, Parmentier & Scheirer, 2002). Other models applied to lithospheric processes, sometimes suggested to have influenced the NAIP genesis, include: (1) the 'soft-point model', where the lithosphere is pre-weakened locally by igneous activity and their sources at depth, thereby localizing extensional stresses, which in turn may control the rift propagation and geometry (Corti *et al.* 2001; Callot, Geoffroy & Brun, 2002; Geoffroy, 2005); (2) lithospheric extension due to relaxation of intra-plate tensional stress regimes, which in turn generates numerous centres of extension dispersed over plate-wide areas, perhaps ultimately facilitating rifting (Nielsen, Stephenson & Thomsen, 2007).

4.b. Exploring potential source regions for the NAIP magmas

Key issues for theories regarding the NAIP petrogenesis include the high temperatures necessary to explain the common occurrence of picrites and the nature of source rocks necessary to explain the heterogeneous compositions of many of the encountered basalts (Meyer, Van Wijk & Gernigon, 2007). Melting of peridotites, contaminated with various amounts of recycled oceanic crustal material, is frequently invoked to explain the geochemical variations in flood basalt provinces (Kogiso, Hirose & Takahashi, 1998; Yaxley, 2000; Green & Falloon, 2005). However, as subducted oceanic crustal material is thought to reside in the lower mantle (Zhao, 2004), in the middle mantle (Courtillet *et al.* 2003, Zhao, 2004) and in the upper mantle (Green *et al.* 2001; Donnelly *et al.* 2004), geochemical signatures do not necessarily constrain a certain level of the mantle as the source of origin. Picrites have been interpreted to form at $\sim 1440^\circ\text{C}$ and ~ 2 GPa (Green *et al.* 2001; Green & Falloon, 2005), and mid-ocean ridge basalts (MORB) are thought to form at temperatures above $\sim 1240^\circ\text{C}$ but below $\sim 1400^\circ\text{C}$ and at depths ranging from ~ 30 km to ~ 45 km (Hirose & Kawamoto, 1995; Presnall, Gudfinnsson &

Walter, 2002). Presnall, Gudfinnsson & Walter (2002) showed that a temperature increase of only $\sim 20^\circ\text{C}$ was required to increase melt productivity from 0 to 24% in a homogeneous peridotitic mantle at near-solidus temperatures. The presence of small amounts of recycled oceanic crust and/or water in peridotitic source rocks is inferred to increase the degree of melting at fixed temperatures (Kogiso, Hirose & Takahashi, 1998; Yaxley, 2000) and to lower the solidus temperature (Hirose & Kawamoto, 1995; Yaxley, 2000; Presnall, Gudfinnsson & Walter, 2002; Green & Falloon, 2005).

At average geothermal gradients for the upper mantle in an ocean ridge environment (e.g. Blatt & Tracy, 1995), adiabatic ascent of uncontaminated potential peridotitic source rocks from depths of > 400 km is required in order to produce picrites at $\sim 1440^\circ\text{C}$ and ~ 2 GPa (Fig. 4a). Substantially shallower depths may be required for picrite genesis from hypothetical assemblages of contaminated and/or hydrated source rocks (Fig. 4b, c). For flood basalts comparable in composition to oceanic island basalts (OIB), melting of enriched/hydrated source rocks would be expected to commence a few tens of kilometres deeper than similar melting of a pure peridotitic counterpart to produce MORB (Yaxley, 2000) (Fig. 4d, e). In summary, the adiabatic ascent of source material from depths of a few hundred kilometres is probably required in order to provide temperatures realistically needed to produce picrites.

4.c. The NAIP in the context of rift geometry and triple junctions

The geometry of the NAIP (Figs 2, 3) and the longevity of the igneous activity, together with the involvement of the Rockall–Faroe Islands microcontinent (Roberts & Searle, 1979; Edwards, 2002) and the Jan Mayen microcontinent (Kodaira *et al.* 1998; Mjelde *et al.* 2008) in the rift processes, suggest a complex and discontinuous break-up history. A comparable complex rifting evolution has been reported for the Afar Volcanic Province with migrating triple junctions and magmatism (Tesfaye, Harding & Kusky, 2003; Wolfenden *et al.* 2004) and where microcontinents (Danakil and Aisha) were involved in the rifting/igneous processes and commonly defined their own secondary triple junctions and associated magmatism (Garfunkel & Beyth, 2006). Individual large mantle plumes have commonly been linked to magmatism, rifting and triple junction formation in flood basalt provinces like the Afar Volcanic Province (Garfunkel & Beyth, 2006) and the NAIP (Section 3), but other authors have argued that the East African rift system in general developed in response to global plate reorganizations (e.g. Wolfenden *et al.* 2004). The common occurrence of dissimilar geochemical and isotopic signatures in rift-related basalts, within confined areas from, for example, the East African rift system (Barrat *et al.* 1998; Orihashi, Al-Jailani & Nagao, 1998; Rogers *et al.* 2000; George & Rogers, 2002; Keranen & Klemperer,




Figure 4. Simplified diagram showing depths versus Mid-Ocean Ridge mantle geotherms (right y axis) from (Blatt & Tracy, 1995). (a) Required depths of adiabatic ascent and melting of dry peridotitic source rocks to generate picrites at ~ 2.0 GPa (around 60 km depth) and at $\sim 1440^\circ\text{C}$ have been calculated/estimated. Adiabatic gradients of (i) $0.60^\circ\text{C km}^{-1}$ (McKenzie & Bickle, 1988), (ii) $0.54^\circ\text{C km}^{-1}$ (McKenzie, Jackson & Priestley, 2005) and (iii) $0.42^\circ\text{C km}^{-1}$ (Ichiki *et al.* 2006) have been applied in the calculations/estimations. In assuming lowering of solidus temperatures of $\sim 50^\circ\text{C}$ for source material containing H_2O (Green & Falloon, 2005) and a further decrease of solidus of $\sim 50^\circ\text{C}$ for fertile source material (Yaxley, 2000) calculations/estimations are carried out for (b) wet and (c) wet + fertile source rocks. However, the scenarios in (b) and (c) may produce normal basalts unless melting starts at deeper levels (e.g. Yaxley, 2000). (d) MORB generation from partial melting at ~ 30 to ~ 45 km depth. (e) Increased melting column for the generation of OIB (oceanic island basalts). See text for further explanation.

2008), from Iceland (Kitagawa *et al.* 2008) and from the Azores (Beier *et al.* 2008), suggests melting from distinct mantle reservoirs.

The association between enhanced magmatism and rift geometry, that is, triple junctions (Sears, George & Winne, 2005) or kinks in rifting trends (Abdel-Rahman & Nassar, 2004; Wolfenden *et al.* 2004), is well known. In this context the evolution of the proto-Iceland region may be of relevance for the Early Palaeogene NAIP magmatism, as the great increase in the volume of magma production in that area in Middle Palaeogene times (Foulger & Anderson, 2005) coincided with the establishment of the ridge–ridge–transform triple junction (Reykjanes ridge–Kolbeinsey ridge–Faroe transform fault) recorded by Bott (1985).

4.d. NAIP in the context of plate tectonic processes in adjacent areas

In the context of Early Palaeogene global plate-tectonic processes, it is noteworthy that the relative convergence and associated compression of Africa and Iberia with respect to W Europe came to a standstill from the earliest Paleocene to the Early Eocene (Rosenbaum, Lister & Duboz, 2002), that is, in the same time interval as the occurrence of the majority of Early Palaeogene NAIP magmatism and the initiation of the continental break-up of the proto-North Atlantic area (Table 1; Figs 1, 2, 3). The causal mechanism for this standstill of relative plate convergence has been tentatively interpreted to result from a contemporaneous continental collision in the Alps between the African

and European plates at around 65 Ma (Jolivet & Faccenna, 2000; Rosenbaum, Lister & Duboz, 2002). A recent complementary tectonic model inferred to have terminated compression and perhaps facilitated extension in the NW Atlantic area in the Early Palaeogene involves major left-lateral displacements between Greenland and Europe and within the NW parts of Europe (Fig. 2) that ultimately resulted in narrowing (contraction) and retreat of the European plate relative to the African plate (Nielsen, Stephenson & Thomsen, 2007).

In a rifting perspective, Lundin & Doré (2005) argued that the Early Palaeogene igneous–tectonic activities in the proto-North Atlantic area that generated the NAIP were merely a result/expression of the final phases of the ongoing break-up of Pangaea, spatially and temporally linking the Early Paleocene central Atlantic rifting (e.g. Ziegler, 1989, 1992) in the south with the Early Eocene rifting in the Eurasian Basin to the north (e.g. Srivastava, 1985; Brown, Parsons & Becker, 1987).

4.e. Lithospheric strength

An important issue to be addressed in complex large igneous provinces like the NAIP is: what caused the magmatism to be so widespread until a relatively narrow sea-floor-spreading zone was finally established? Clearly, the strength of certain parts of the lithosphere and its relative capability to resist stretching, rupture and/or intrusion of magmas must have played a major role. Studies on lithospheric strength in a laterally

homogeneous undeformed lithosphere have been dealt with in a number of studies (e.g. Kohlstedt, Evans & Mackwell, 1995; Hirth & Kohlstedt, 1996; Kuszniir & Park, 2002; Van Wijk & Cloetingh, 2002; Jackson *et al.* 2008). In brief, these authors concluded that increased heat flow and high rates of spreading generally resulted in a net weakening of the lithosphere and conversely, very slow spreading rates and low heat flows could result in a net strengthening. Increased fluid pressure further weakens all affected rock assemblages in the lithosphere (Hirth & Kohlstedt, 1996; Jackson *et al.* 2008). Upper mantle heterogeneity and the presence of old shear zones in the lithosphere may play a prominent role in incipient rifting both by enhancing partial melting and by reactivation of old shear zones (Holdsworth, Butler & Roberts, 1997; Ryan & Dewey, 1997; Kohlstedt, Evans & Mackwell, 1995).

5. Conclusions and closing remarks

In this contribution we have reviewed some key magmatic centres of the NAIP in a geodynamic framework, focusing on their interrelationships and the tectonic developments during the onset and development of the NAIP. The specific conditions directly prior to the onset of the NAIP and the continued development of the region during the Palaeogene, based on the findings of the present study, are highlighted as follows:

(1) The onset of the rifting and igneous activity of the NAIP area was a temporal and spatial continuation of the rifting in the adjacent central Atlantic Ocean to the south and a precursor for the rifting in the Eurasian Basin to the north. The main igneous and tectonic activities in the NAIP in Early Palaeogene times coincide with contemporaneous changes in the relative motion between the European and African plates, which possibly halted the previous compressional regimes in the NW Atlantic during this time span.

(2) Taken all together, the apparent geometry of the main igneous regions of the NAIP at the conjugate E Greenland–NW European margins in particular shows similarities with trends of the embryonic stages of classic continental rifting regimes consisting of numerous more or less interconnected triple junctions. However, the locations of numerous smaller central igneous complexes and/or seamounts at the NW European margin (Fig. 1) do not seem to fit with such a simple rifting model if they all formed contemporaneously with the larger igneous NAIP regions.

(3) An aiding factor to NAIP rifting could have been Early Palaeogene oblique extension and active sinistral transforms (reactivation?) at the conjugate E Greenland–NW European margins (Nielsen, Stephenson & Thomsen, 2007) (Fig. 2).

(4) While partial melting in the upper mantle to produce the bulk of the (normal) basaltic rocks of the NAIP is not necessarily dependent on deep mantle convection, some ascent of hot mantle jets seems to be required in order to generate the NAIP picrites (Fig. 4).

(5) Whether global plate reorganizations or a single large mantle plume was the driving force for the NAIP evolution, the close proximity of many parts of the NAIP magmatism and transient uplifts to ancient orogenic sutures and/or fronts (Figs 2, 3) suggests that lithospheric control was an important factor in the embryonic stages of magmatism and rifting.

Acknowledgements. This review was compiled by the first author as part of an ongoing Ph.D. project ‘Sub-seismic geometry and influence of intrusions in igneous-affected basins’, sponsored by Chevron, BP and Geysir through the Faroese Competence Development Fund. A. Saunders and J. MacLennan critically reviewed an earlier version of the manuscript. Two anonymous reviewers are thanked for their comments on the recent version of the paper.

References

- ABDEL-RAHMAN, A.-F. M. & NASSAR, P. 2004. Cenozoic volcanism in the Middle East: petrogenesis of alkali basalts from northern Lebanon. *Geological Magazine* **141**, 545–63.
- ARCHER, S. G., BERGMAN, S. C., ILIFFE, J., MURPHY, C. M. & THORNTON, M. 2005. Palaeogene igneous rocks reveal new insights into the geodynamic evolution and petroleum potential of the Rockall Trough, NE Atlantic Margin. *Basin Research* **17**, 171–201.
- BARRAT, J. A., FOURCADE, S., JAHN, B. M., CHEMINÉE, J. L. & CAPDEVILA, R. 1998. Isotope (Sr, Nd, O) and trace-element geochemistry of volcanics from the Erta’Ale range (Ethiopia). *Journal of Volcanology and Geothermal Research* **80**, 85–100.
- BARTON, A. J. & WHITE, R. S. 1997. Crustal structure of Eororas Bank continental margin and mantle thermal anomalies beneath the north Atlantic. *Journal of Geophysical Research* **102**(B2), 3109–29.
- BEIER, C., HAASE, K. M., ABOUCHAMI, W., KRIENITZ, M.-S. & HAUFF, F. 2008. Magma genesis by rifting of oceanic lithosphere above anomalous mantle: Terceira Rift, Azores. *Geochemistry Geophysics Geosystems* **9**(12), Q12013, pp. 1–26.
- BERNDT, C., PLANKE, S., ALVESTAD, E., TSICALAS, F. & RASMUSSEN, T. 2001. Seismic volcanostratigraphy off the Norwegian margin: constraints on tectonomagmatic break-up processes. *Journal of the Geological Society, London* **158**, 413–26.
- BLATT, H. & TRACY, R. J. 1995. *Petrology: Igneous, Sedimentary, and Metamorphic*. Second Edition. New York: W. H. Freeman and Company, 529 pp.
- BOTT, M. H. 1985. Plate tectonic evolution of the Icelandic Transverse Ridge and adjacent regions. *Journal of Geophysical Research* **90**(B12), 9953–60.
- BOTT, M. H. P. 1987. The continental margin of central East Greenland in relation to North Atlantic plate tectonic evolution. *Journal of the Geological Society, London* **144**, 561–8.
- BREKKE, H., DAHLGREN, S., NYLAND, B. & MAGNUS, C. 1999. The prospectivity of the Vøring and Møre basins on the Norwegian Sea continental margin. *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*, 261–74. Geological Society, London.
- BROWN, P. E., PARSONS, I. & BECKER, S. M. 1987. Peralkaline volcanicity in the Arctic Basin – the Kap Washington Volcanics, petrology and palaeotectonics. *Journal of the Geological Society, London* **144**, 707–15.

- BUGGE, T., PRESTVIK, T. & ROKOENGEN, K. 1980. Lower Tertiary volcanic rocks off Kristiansund, mid Norway. *Marine Geology* **35**, 277–86.
- BULL, J. M. & MASSON, D. G. 1996. The southern margin of the Rockall Plateau: stratigraphy, Tertiary volcanism and plate tectonic evolution. *Journal of the Geological Society, London* **153**, 601–12.
- BURKE, K. & DEWEY, J. F. 1973. Plume-generated triple junctions: key indicators in applying plate tectonics to old rocks. *Journal of Geology* **81**, 406–33.
- CALLOT, J.-P., GEOFFROY, L. & BRUN, J.-P. 2002. Development of volcanic passive margins: three-dimensional laboratory models. *Tectonics* **21**(6), 2.1–2.13.
- CHALMERS, J. A. 1997. The continental margin off southern Greenland: along-strike transition from an amagmatic to a volcanic margin. *Journal of the Geological Society, London* **154**, 571–6.
- CHALMERS, J. A., LARSEN, L. M. & PEDERSEN, A. K. 1995. Widespread Palaeocene volcanism around the North Atlantic and Labrador Sea: evidence for a large, hot, early plume head. *Journal of the Geological Society, London* **152**, 965–9.
- CHALMERS, J. A. & LAURSEN, K. H. 1995. Labrador Sea: the extent of continental and oceanic crust and the timing of the onset of seafloor spreading. *Marine and Petroleum Geology* **12**, 205–17.
- CHAMBERS, L. M., PRINGLE, M. S. & PARRISH, R. R. 2005. Rapid formation of the Small Isles Tertiary centre constrained by precise $^{40}\text{Ar}/^{39}\text{Ar}$ and U–Pb ages. *Lithos* **79**, 367–84.
- CLIFT, P. D., TURNER, J. & ODP Leg 152 Scientific Party. 1995. Dynamic support by the Iceland Plume and its effect on the subsidence of the northern Atlantic margins. *Journal of the Geological Society, London* **152**, 935–41.
- COCKS, L. R. M. 2005. Presidential Address 2005: where was Britain in the Palaeozoic? *Proceedings of the Geologists' Association* **116**, 117–27.
- CONNELLY, J. N., THRANE, K., KRAWIEC, A. W. & GARDE, A. A. 2006. Linking the Palaeoproterozoic Nagssugtoqidian and Rinkian orogens through the Disko Bugt region of West Greenland. *Journal of the Geological Society, London* **163**, 319–35.
- COPE, J. C. W. 1994. A Latest Cretaceous hotspot and the southeasterly tilt of Britain. *Journal of the Geological Society, London* **151**, 905–8.
- CORTI, G., BONINI, M., INNOCENTI, F., MANETTI, P. & MULUGETA, G. 2001. Centrifuge models simulating magma emplacement during oblique rifting. *Journal of Geodynamics* **31**, 557–76.
- COURTILLOT, V., DAVAILLE, A., BESSE, J. & STOCK, J. 2003. Three distinct types of hotspots in the Earth's mantle. *Earth and Planetary Science Letters* **205**, 295–308.
- DICKIN, A. P. 1992. Evidence for an Early Proterozoic crustal province in the North Atlantic Region. *Journal of the Geological Society, London* **149**, 483–6.
- DONNELLY, K. E., GOLDSTEIN, S. L., LANGMUIR, C. H. & SPIEGELMAN, M. 2004. Origin of enriched ocean ridge basalts and implications for mantle dynamics. *Earth and Planetary Science Letters* **226**, 347–66.
- DORÉ, A. G., LUNDIN, E. R., FICHLER, C. & OLESEN, O. 1997. Patterns of basement structure and reactivation along the NE Atlantic margin. *Journal of the Geological Society, London* **154**, 85–92.
- DORÉ, A. G., LUNDIN, E. R., JENSEN, L. N., BIRKELAND, Ø., ELIASSEN, P. E. & FICHLER, C. 1999. Principal tectonic events in the evolution of the Northwest European Atlantic margin. In *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference* (eds A. J. Fleet & S. A. R. Boldy), pp. 41–61. Geological Society, London.
- EDWARDS, J. W. F. 2002. Development of the Hatton–Rockall Basin, North-East Atlantic Ocean. *Marine and Petroleum Geology* **19**, 193–205.
- ELDHOLM, O. & GRUE, K. 1994. North Atlantic volcanic margins: dimensions and production rates. *Journal of Geophysical Research* **99**(B2), 2955–68.
- ELDHOLM, O., THIEDE, J. & TAYLOR, E. 1989. Evolution of the Vøring volcanic margin. *Proceedings of the Ocean Drilling Program, Scientific Results* **104**, 1033–65.
- ELLIOT, G. M. & PARSON, L. M. 2008. Influence of margin segmentation upon the break-up of the Hatton Bank rifted margin, NE Atlantic. *Tectonophysics* **457**(3–4), 161–76.
- ENGLAND, R. W. 1988. The Early Tertiary stress regime in NW Britain: evidence from the patterns of volcanic activity. In *Early Tertiary volcanism and the opening of the NE Atlantic* (eds A. C. Morton & L. M. Parson), pp. 381–9. Geological Society, London, Special Publication no. 39.
- ESTRADA, S., HÖHNDORF, A. & HENJES-KUNST, F. 2001. Cretaceous/Tertiary volcanism in North Greenland: the Kap Washington Group. *Polarforschung* **69**, 17–23.
- FITTON, J. G., LARSEN, L. M., SAUNDERS, A. D., HARDARSON, B. S. & KEMPTON, P. D. 2000. Paleogene continental to oceanic magmatism on the SE Greenland continental margin at 63°N: a review of the results of Ocean Drilling Program Legs 152 and 163. *Journal of Petrology* **41**(7), 951–66.
- FOULGER, G. R. & ANDERSON, D. L. 2005. A cool model for the Iceland hotspot. *Journal of Volcanology and Geothermal Research* **141**, 1–22.
- FOULGER, G. R., NATLAND, J. H. & ANDERSON, D. L. 2005a. Genesis of the Iceland melt anomaly by plate tectonic processes. *Geological Society of America, Special Paper* **388**, 595–625.
- FOULGER, G. R., NATLAND, J. H. & ANDERSON, D. L. 2005b. A source for Icelandic magmas in remelted Iapetus crust. *Journal of Volcanology and Geothermal Research* **141**, 23–44.
- GAMBLE, J. A., WYSOCZANSKY, R. J. & MEIGHAN, I. G. 1999. Constraints on the age of the British Tertiary Volcanic Province from ion microprobe U–Pb (SHRIMP) ages for acid igneous rocks from NE Ireland. *Journal of the Geological Society, London* **156**, 291–9.
- GARFUNKEL, Z. & BEYTH, M. 2006. Constraints on the structural development of Afar imposed by the kinematics of the major surrounding plates. In *The Afar Volcanic Province within the East African Rift System* (eds G. Yirgu, C. J. Ebinger & P. K. H. Maguire), pp. 23–42. Geological Society of London, Special Publication no. 259.
- GEOFFROY, L. 2005. Volcanic passive margins. *C. R. Geoscience* **337**, 1395–1408.
- GEOFFROY, L., BERGERAT, F. & ANGELIER, J. 1994. Tectonic evolution of the Greenland–Scotland ridge during the Paleogene: new constraints. *Geology* **22**, 653–6.
- GEOFFROY, L., BERGERAT, F. & ANGELIER, J. 1996. Brittle tectonism in relation to the Palaeogene evolution of the Thulean/NE Atlantic domain: a study in Ulster. *Geological Journal* **31**, 259–69.
- GEOFFROY, L., CALLOT, J.-P., SCAILLET, S., SKUCE, A., GÉLARD, J. P., RAVILLY, M., ANGELIER, J., BONIN, B., CAYET, C., PERROT, K. & LEPVRIER, C. 2001. Southeast Baffin volcanic margin and the North American–Greenland plate separation. *Tectonics* **20**(4), 566–84.

- GEORGE, R. M. & ROGERS, N. W. 2002. Plume dynamics beneath the African plate inferred from the geochemistry of the Tertiary basalts of southern Ethiopia. *Contributions to Mineralogy and Petrology* **144**, 286–305.
- GERNIGON, L., OLESEN, O., EBBING, J., WIENECKE, S., GAINA, C., MOGAARD, J. O., SAND, M. & MYKLEBUST, R. 2008. Geophysical insights and early spreading history in the vicinity of the Jan Mayen Fracture Zone, Norwegian–Greenland Sea. *Tectonophysics* **TECTO-124188** (21 pp.), doi:10.1016/j.tecto.2008.04.025
- GIBSON, S. A. 2002. Major element heterogeneity in Archean to recent mantle plume starting heads. *Earth and Planetary Science Letters* **195**, 59–74.
- GILL, R. C. O., HOLM, P. M. & NIELSEN, T. F. D. 1995. Was a short-lived Baffin Bay plume active prior to initiation of the present Icelandic plume? Clues from high-Mg picrites of West Greenland. *Lithos* **34**, 27–39.
- GREEN, P. F., DUDDY, I. R., BRAY, R. J. & LEWIS, C. L. E. 1993. Elevated palaeotemperatures prior to Early tertiary cooling throughout the UK region: implications for hydrocarbon generation. *Petroleum Geology of Northwest Europe: Proceedings of the 4th Conference*, 1067–74. Geological Society, London.
- GREEN, D. H. & FALLOON, T. J. 2005. Primary magmas at mid-ocean ridges, ‘hotspots’, and other intraplate settings: Constraints on mantle potential temperature. *Geological Society of America, Special Paper* **388**, 217–47.
- GREEN, D. H., FALLOON, T. J., EGGINS, S. M. & YAXLEY, G. M. 2001. Primary magmas and mantle temperatures. *European Journal of Mineralogy* **13**, 437–51.
- HALD, N. & TEGNER, C. 2000. Composition and age of tertiary sills and dykes, Jameson Land Basin, East Greenland: relation to regional flood volcanism. *Lithos* **54**, 207–33.
- HANGHØJ, K., STOREY, M. & STECHER, O. 2003. An isotope and trace element study of the East Greenland Tertiary dyke swarm: constraints on temporal and spatial evolution during continental rifting. *Journal of Petrology* **44**(11), 2081–2112.
- HANSEN, K. & BROOKS, C. K. 2002. The evolution of the East Greenland margin as revealed from fission-track studies. *Tectonophysics* **349**, 93–111.
- HARRISON, J. C., MAYR, U., MCNEIL, D. H., SWEET, A. R., MCINTYRE, D. J., EBERLE, J. J., HARRINGTON, C. R., CHALMERS, J. A., DAM, G. & NØHR-HANSEN, H. 1999. Correlation of Cenozoic sequences of the Canadian Arctic region and Greenland; implications for the tectonic history of northern North America. *Bulletin, Canadian Petroleum Geology* **47**(3), 223–54.
- HIROSE, K. & KAWAMOTO, T. 1995. Hydrous partial melting of lherzolite at 1 GPa: the effect of H₂O on the genesis of basaltic magmas. *Earth and Planetary Science Letters* **133**, 463–73.
- HIRTH, G. & KOHLSTEDT, D. L. 1996. Water in the oceanic upper mantle: implications for rheology, melt extraction and the evolution of the lithosphere. *Earth and Planetary Science Letters* **144**, 93–108.
- HITCHEN, K. 2004. The geology of the UK Hatton–Rockall margin. *Marine and Petroleum Geology* **21**, 993–1012.
- HITCHEN, K., MORTON, A. C., MEARN, E. W., WHITEHOUSE, M. & STOKER, M. S. 1997. Geological implications from geochemical and isotopic studies of Upper Cretaceous and lower Tertiary igneous rocks around the northern Rockall Trough. *Journal of the Geological Society, London* **154**, 517–21.
- HITCHEN, K. & RITCHIE, J. D. 1993. New K–Ar ages, and a provisional chronology, for the offshore part of the British Tertiary Igneous Province. *Scottish Journal of Geology* **29**(1), 73–85.
- HOLDSWORTH, R. E., BUTLER, C. A. & ROBERTS, A. M. 1997. The recognition of reactivation during continental deformation. *Journal of the Geological Society, London* **154**, 73–8.
- HOLM, P. M., HALD, N. & WAAGSTEIN, R. 2001. Geochemical and Pb–Sr–Nd isotopic evidence for separate hot depleted and Iceland plume mantle sources for the Paleogene basalts of the Faroe Islands. *Chemical Geology* **178**, 95–125.
- ICHIKI, M., BABA, K., OBAYASHI, M. & UTADA, H. 2006. Water content and geotherm in the upper mantle above the stagnant slab: Interpretation of electrical conductivity and seismic P-wave velocity models. *Physics of the Earth and Planetary Interiors* **155**, 1–15.
- JACKSON, J., MCKENZIE, D., PRIESTLEY, K. & EMMERSON, B. 2008. New views on the structure and rheology of the lithosphere. *Journal of the Geological Society, London* **165**, 453–65.
- JAPSEN, P., GREEN, P. F. & CHALMERS, J. A. 2005. Separation of Palaeogene and Neogene uplift on Nuussuaq, West Greenland. *Journal of the Geological Society, London* **162**, 299–314.
- JERRAM, D. A., SINGLE, R. T., HOBBS, R. & NELSON, C. 2009. Understanding the offshore flood basalt sequence using onshore volcanic facies analogues: an example from the Faroe–Shetland basin. *Geological Magazine* **146**, 353–67.
- JERRAM, D. A. & WIDDOWSON, M. 2005. The anatomy of Continental Flood Basalt Provinces: geological constraints on the processes and products of flood volcanism. *Lithos* **79**, 385–405.
- JOHNSON, H., RITCHIE, J. D., MCINROY, D. B. & KIMBELL, G. S. 2005. Aspects of the Cenozoic deformational history of the Northeast Faroe–Shetland Basin, Wyville-Thomson Ridge and Hatton Bank areas. *Petroleum Geology of Northwest Europe: Proceedings of the 6th Conference* 993–1007. Geological Society, London.
- JOLIVET, L. & FACCENNA, C. 2000. Mediterranean extension and the Africa–Eurasia collision. *Tectonics* **19**(6), 1095–1106.
- JONES, S. M., WHITE, N. & LOVELL, B. 2001. Cenozoic and Cretaceous uplift in the Porcupine Basin and its relationship to a mantle plume. In *The Petroleum Exploration of Ireland's Offshore Basins* (eds D. Corcoran, D. W. Haughton & P. M. Shannon), pp. 345–60. Geological Society of London, Special Publication no. 188.
- KANARIS-SOTIRIOU, R., MORTON, A. C. & TAYLOR, P. N. 1993. Palaeogene peraluminous magmatism, crustal melting and continental breakup: the Erlend complex, Faeroe–Shetland Basin, NE Atlantic. *Journal of the Geological Society, London* **150**, 903–14.
- KARSON, J. A. & BROOKS, C. K. 1999. Structural and magmatic segmentation of the Tertiary East Greenland Volcanic Rifted Margin. In *Continental Tectonics* (eds C. M. Niocaill & P. D. Ryan), pp. 313–38. Geological Society of London, Special Publication no. 164.
- KARSON, J. A., BROOKS, C. K., STOREY, M. & PRINGLE, M. S. 1998. Tertiary faulting and pseudotachylites in the east Greenland volcanic rifted margin: seismogenic faulting during magmatic construction. *Geology* **26**(1), 39–42.

- KENT, R. W. & FITTON, J. G. 2000. Mantle sources and melting dynamics in the British Paleogene Igneous Province. *Journal of Petrology* **41**(7), 1023–40.
- KERANEN, K. & KLEMPERER, S. L. 2008. Discontinuous and diachronous evolution of the main Ethiopian Rift: Implications for the development of continental rifts. *Earth and Planetary Science Letters* **265**, 96–111.
- KIMBELL, G. S., RITCHIE, J. D., JOHNSON, H. & GATLIFF, R. W. 2005. Controls on the structure and evolution of the NE Atlantic margin revealed by regional 3D gravity modelling. In *Petroleum Geology: North-West Europe and Global Perspectives – Proceedings of the 6th Petroleum Geology Conference* (eds A. G. Doré & B. A. Vining), pp. 933–47. Geological Society of London.
- KITAGAWA, H., KOBAYASHI, K., MAKISHIMA, A. & NAKAMURA, E. 2008. Multiple pulses of the Mantle Plume: Evidence from Tertiary Icelandic Lavas. *Journal of Petrology* **49**(7), 1365–96.
- KODAIRA, S., MJELDE, R., GUNNARSON, K., SHIOBARA, H. & SHIMAMURA, H. 1998. Structure of the Jan Mayen microcontinent and implications for its evolution. *Geophysical Journal International* **132**, 383–400.
- KOGISO, T., HIROSE, K. & TAKAHASHI, E. 1998. Melting experiments on homogeneous mixtures of peridotite and basalt: application to the genesis of ocean island basalts. *Earth and Planetary Science Letters* **162**, 45–61.
- KOHLSTEDT, D. L., EVANS, B. & MACKWELL, S. J. 1995. Strength of the lithosphere: constraints imposed by laboratory experiments. *Journal of Geophysical Research* **100**(B9), 17587–602.
- KUSZNIR, N. J. & PARK, R. G. 2002. The extensional strength of the continental lithosphere: its dependence on geothermal gradient, and crustal composition and thickness. In *Extensional Tectonics: Regional-scale Processes* (compilers R. E. Holdsworth & J. P. Turner), pp. 97–114. *The Geological Society, Key Issues in Earth Sciences*, **2**(1) (first published in *Continental Extensional Tectonics* (eds M. P. Coward, J. F. Dewey & P. L. Hancock), pp. 35–52. Geological Society of London, Special Publication no. 28. 1987).
- LARSEN, H. C. & MARCUSSEN, C. 1992. Sill intrusion, flood basalt emplacement and deep crustal structure of the Scoresby Sund region, East Greenland. In *Magmatism and the Causes of Continental Break-up* (eds B. C. Storey, T. Alabaster & R. J. Pankhurst), pp. 365–86. Geological Society of London, Special Publication no. 68.
- LARSEN, L. M., REX, D. C., WATT, W. S. & GUISE, P. G. 1999a. $^{40}\text{Ar}/^{39}\text{Ar}$ dating of alkali basaltic dykes along the south-west coast of Greenland: Cretaceous and Tertiary igneous activity along the eastern margin of the Labrador sea. *Geology of Greenland, Survey Bulletin* **184**, 19–29.
- LARSEN, H. C. & SAUNDERS, A. D. 1998. Tectonism and volcanism at the southeast Greenland rifted margin: a record of plume impact and later continental rupture. *Proceedings of the ODP, Scientific Results* **152**, 503–33.
- LARSEN, L. M., WAAGSTEIN, R., PEDERSEN, A. K. & STOREY, M. 1999b. Trans-Atlantic correlation of the Palaeocene volcanic successions in the Faeroe Islands and East Greenland. *Journal of the Geological Society, London* **156**, 1081–95.
- LARSEN, L. M. & WATT, W. S. 1985. Episodic volcanism during break-up of the North Atlantic: evidence from the East Greenland plateau basalts. *Earth and Planetary Science Letters* **73**, 105–16.
- LENOIR, X., FERAUD, G. & GEOFFROY, L. 2003. High-rate flexure of the East Greenland volcanic margin: constraints from $^{40}\text{Ar}/^{39}\text{Ar}$ dating of basaltic dykes. *Earth and Planetary Science Letters* **214**, 515–28.
- LUNDIN, E. R. & DORÉ, A. G. 2005. NE Atlantic break-up: a re-examination of the Iceland mantle plume model and the Atlantic–Arctic linkage. In *Petroleum Geology: North-West Europe and Global Perspectives – Proceedings of the 6th Petroleum Geology Conference* (eds A. G. Doré & B. A. Vining), pp. 739–54. Geological Society of London.
- MACKAY, L. M., TURNER, J., JONES, S. M. & WHITE, N. J. 2005. Cenozoic vertical motions in the Moray Firth Basin associated with initiation of the Iceland Plume. *Tectonics* **24**, TC5004 (pp. 1–23).
- MACLENNAN, J. & JONES, S. M. 2006. Regional uplift, gas hydrate dissociation and the origins of the Paleocene–Eocene thermal maximum. *Earth and Planetary Science Letters* **245**, 65–80.
- MASSON, F., HAUSER, F. & JACOB, A. W. B. 1999. The lithospheric trace of the Iapetus Suture in SW Ireland from teleseismic data. *Tectonophysics* **302**, 83–98.
- MATHIESEN, A., BIDSTRUP, T. & CHRISTIANSEN, G. 2000. Denudation and uplift history of the Jameson Land basin, East Greenland—constrained from maturity and apatite fission track data. *Global and Planetary Change* **24**, 275–301.
- MCKENZIE, D. & BICKLE, M. J. 1988. The volume and composition of melt generated by extension of the lithosphere. *Journal of Petrology* **29**(3), 625–79.
- MCKENZIE, D., JACKSON, J. & PRIESTLEY, K. 2005. Thermal structure of oceanic and continental lithosphere. *Earth and Planetary Science Letters* **233**, 337–49.
- MEYER, R., VAN WIJK, J. & GERNIGON, L. 2007. The North Atlantic Igneous Province: A review of models for its formation. *Geological Society of America, Special Paper* **430**, 525–52.
- MJELDE, R., BREIVIK, A. J., RAUM, T., MITTELSTAEDT, E., ITO, G. & FALEIDE, J. I. 2008. Magmatic and tectonic evolution of the North Atlantic. *Journal of the Geological Society, London* **165**, 31–42.
- MORGAN, J. V. & BARTON, P. J. 1990. A geophysical study of the Hatton Bank volcanic margin: a summary of the results from a combined seismic, gravity and magnetic experiment. *Tectonophysics* **173**, 517–26.
- MORTON, A. C., HITCHEN, K., RITCHIE, J. D., HINE, N. M., WHITEHOUSE, M. & CARTER, S. G. 1995. Late Cretaceous basalts from Rosemary Bank, Northern Rockall Trough. *Journal of the Geological Society, London* **152**, 947–52.
- MUDGE, D. C. & JONES, S. M. 2004. Paleocene uplift and subsidence events in the Scotland–Shetland and North Sea region and their relationship to the Iceland Plume. *Journal of the Geological Society, London* **161**, 381–6.
- NADIN, P. A., KUSZNIR, N. J. & CHEADLE, M. J. 1997. Early Tertiary plume uplift of the North Sea and Faeroe–Shetland Basins. *Earth and Planetary Science Letters* **148**, 109–27.
- NATLAND, J. H. & WINTERER, E. L. 2005. Fissure control on volcanic action in the Pacific. *Geological Society of America, Special Paper* **388**, 687–710.
- NAYLOR, P. H., BELL, B. R., JOLLEY, D. W., DURNALL, P. & FREDSTED, R. 1999. Palaeogene magmatism in the Faeroe–Shetland Basin: influences on uplift history and sedimentation. *Petroleum Geology of Northwest Europe: Proceedings of the 5th Conference*, pp. 545–58. Geological Society of London.
- NIELSEN, T. F. D. 1987. Tertiary alkaline magmatism in East Greenland: a review. In *Alkaline Igneous Rocks* (eds J.

- G. Fitton & B. G. J. Upton), pp. 489–515. Geological Society of London, Special Publication no. 30.
- NIELSEN, T. K., LARSEN, H. C. & HOPPER, J. R. 2002. Contrasting rifted margin styles south of Greenland: implications for mantle plume dynamics. *Earth and Planetary Science Letters* **200**, 271–86.
- NIELSEN, S. B., STEPHENSON, R. & THOMSEN, E. 2007. Dynamics of mid-Paleocene north Atlantic rifting linked with European intra-plate deformations. *Nature* **450**, 1071–3.
- O'CONNOR, J. M., STOFFERS, P., WIJBRANS, J. R., SHANNON, P. M. & MORRISSEY, T. 2000. Evidence from episodic seamount volcanism for pulsing of the Iceland plume in the past 70 Myr. *Nature* **408**, 954–8.
- OLESEN, O., EBBING, J., LUNDIN, E., MAURING, E., SKILBREI, J. R., TORSVIK, T. H., HANSEN, E. K., HENNINGSEN, T., MIDBØE, P. & SAND, M. 2007. An improved tectonic model for the Eocene opening of the Norwegian–Greenland Sea: Use of modern magnetic data. *Marine and Petroleum Geology* **24**, 53–66.
- ORIHASHI, Y., AL-JAILANI, A. & NAGAO, K. 1998. Dispersion of the Afar plume: Implications from the Spatiotemporal Distribution of the Late Miocene to Recent Volcanics, Southwestern Arabian Peninsula. *Gondwana Research* **1**(2), 221–34.
- PARK, R. G. 1995. *Geological structures and moving plates*. Blackie Academic & Professional, an Imprint of Chapman & Hall, 337 pp.
- PASSEY, S. R. & BELL, B. R. 2007. Morphologies and emplacement mechanisms of the lava flows of the Faroe Islands Basalt Group, Faroe Islands, NE Atlantic Ocean. *Bulletin of Volcanology* **70**, 139–56.
- PEATE, D. W., BAKER, J. A., BLICHERT-TOFT, J., HILTON, D. R., STOREY, M., KENT, A. J. R., BROOKS, C. K., HANSEN, H., PEDERSEN, A. K. & DUNCAN, R. A. 2003. The Prinsen of Wales Bjerge Formation lavas, East Greenland: the transition from tholeiitic to alkalic magmatism during Paleogene continental break-up. *Journal of Petrology* **44**(2), 279–304.
- PEATE, I. U., LARSEN, M. & LESHER, C. E. 2003. The transition from sedimentation to flood volcanism in the Kangerlussuaq Basin, East Greenland: basaltic pyroclastic volcanism during initial Palaeogene continental break-up. *Journal of the Geological Society, London* **160**, 759–72.
- PLANKE, S., RASMUSSEN, T., REY, S. S. & MYKLEBUST, R. 2005. Seismic characteristics and distribution of volcanic intrusions and hydrothermal vent complexes in the Vøring and Møre basins. In *Petroleum Geology: North-West Europe and Global Perspectives – Proceedings of the 6th Petroleum Geology Conference* (eds A. G. Doré & B. A. Vining), pp. 833–44. Geological Society of London.
- PRESNALL, D. C., GUDFINNSSON, G. H. & WALTER, M. J. 2002. Generation of mid-ocean ridge basalts at pressures from 1 to 7 GPa. *Geochimica et Cosmochimica Acta* **66**(12), 2073–90.
- PRICE, S., BRODIE, J., WHITHAM, A. & KENT, R. 1997. Mid-Tertiary rifting and magmatism in the Traill Ø region, East Greenland. *Journal of the Geological Society, London* **154**, 419–34.
- PRICE, I. & RATTEY, R. P. 1984. Cretaceous tectonics off mid-Norway: implications for the Rockall and Faeroe–Shetland troughs. *Journal of the Geological Society, London* **141**, 985–92.
- RADDICK, M. J., PARMENTIER, E. M. & SCHEIRER, D. S. 2002. Buoyant decompression melting: a possible mechanism for intraplate volcanism. *Journal of Geophysical Research* **107**(B10), ECV 7.1–7.14.
- REDFIELD, T. F., TORSVIK, T. H., ANDRIESEN, P. A. M. & GABRIELSEN, R. H. 2004. Mesozoic and Cenozoic tectonics of the Mære Trøndelag Fault Complex, central Norway: constraints from new apatite fission track data. *Physics and Chemistry of the Earth* **29**, 673–82.
- REN, S., FALEIDE, J. I., ELDHOLM, O., SKOGSEID, J. & GRADSTEIN, F. 2003. Late Cretaceous–Paleocene tectonic development of the NW Vøring Basin. *Marine and Petroleum Geology* **20**, 177–206.
- RITCHIE, J. D., HITCHEN, K. & EDWARDS, J. W. F. 1997. The Sigmundur Complex, a ? Tertiary igneous centre in the northern Rockall Trough. *Scottish Journal of Geology* **33**(2), 97–103.
- ROBERTS, D. 2003. The Scandinavian Caledonides: event chronology, palaeogeographic settings and likely modern analogues. *Tectonophysics* **365**, 283–99.
- ROBERTS, D. G. & SEARLE, R. C. 1979. The western Rockall Plateau: stratigraphy and structural evolution. *Initial Reports of the Deep Sea Drilling Project XLVIII*, 1061–88.
- ROGERS, N., MACDONALD, R., FITTON, J. G., GEORGE, R., SMITH, M. & BARREIRO, B. A. 2000. Two mantle plumes beneath the East African rift system: Sr, Nd and Pb isotope evidence from Kenya Rift basalts. *Earth and Planetary Science Letters* **176**, 387–400.
- ROSENBAUM, G., LISTER, G. S. & DUBOZ, C. 2002. Relative motions of Africa, Iberia and Europe during alpine orogeny. *Tectonophysics* **359**, 117–29.
- RUDGE, J. F., CHAMPION, M. E. S., WHITE, N., MCKENZIE, D. & LOVELL, B. 2008. A plume model of transient diachronous uplift at the Earth's surface. *Earth and Planetary Science Letters* **267**, 146–60.
- RYAN, P. D. & DEWEY, J. F. 1997. Continental eclogites and the Wilson Cycle. *Journal of the Geological Society, London* **154**, 437–42.
- SAUNDERS, A. D., FITTON, J. G., KERR, A. C., NORRY, M. J. & KENT, R. W. 1997. The North Atlantic Igneous Province. *Geophysical Monograph* **100**, 45–93.
- SAUNDERS, A. D., JONES, S. M., MORGAN, L. A., PIERCE, K. L., WIDDOWSON, M. & XU, Y. G. 2007. Regional uplift associated with continental large igneous provinces: the roles of mantle plumes and the lithosphere. *Chemical Geology* **241**, 282–318.
- SEARS, J. W., GEORGE, G. M. ST. & WINNE, J. C. 2005. Continental rift systems and anorogenic magmatism. *Lithos* **80**, 147–54.
- SINGLE, R. T. & JERRAM, D. A. 2004. The 3-D facies architecture of flood basalt provinces and their internal heterogeneity: examples from the Palaeogene Skye Lava Field. *Journal of the Geological Society, London* **161**, 911–26.
- SINTON, C. W. & DUNCAN, R. A. 1998. ⁴⁰Ar–³⁹Ar ages of lavas from the Southeast Greenland margin, ODP leg 152, and the Rockall Plateau, DSDP leg 81. *Proceedings of the ODP, Scientific Results* **152**, 387–401.
- SINTON, C. W., HITCHEN, K. & DUNCAN, R. A. 1998. ⁴⁰Ar/³⁹Ar geochronology of silicic and basic volcanic rocks on the margins of the North Atlantic. *Geological Magazine* **135**, 161–70.
- SKAARUP, N., JACKSON, H. R. & OAKEY, G. 2006. Margin segmentation of Baffin Bay/Davis Strait, eastern Canada based on seismic reflection and potential field data. *Marine and Petroleum Geology* **23**, 127–44.
- SKAARUP, N. & PULVERTAFT, C. R. 2007. Aspects of the structure on the coast of the West Greenland volcanic

- province revealed in seismic data. *Bulletin of the Geological Society of Denmark* **55**, 65–80.
- SKOGSEID, J., PEDERSEN, T., ELDHOLM, O. & LARSEN, B. T. 1992. Tectonism and magmatism during NE Atlantic continental break-up: the Vøring Margin. In *Magmatism and the Causes of Continental Break-up* (eds B. C. Storey, T. Alabaster & R. J. Pankhurst), pp. 305–20. Geological Society of London, Special Publication no. 68.
- SKOGSEID, J., PLANKE, S., FALEIDE, J. I., PEDERSEN, T., ELDHOLM, O. & NEVERDAL, F. 2000. NE Atlantic continental rifting and volcanic margin formation. In *Dynamics of the Norwegian Margin* (ed. A. Nøttvedt), pp. 295–326. Geological Society of London, Special Publication no. 167.
- SOPER, N. J., STRACHAN, R. E., HOLDSWORTH, R. E., GAYER, R. A. & GREILING, R. O. 1992. Sinistral transpression and the Silurian closure of Iapetus. *Journal of the Geological Society, London* **149**, 871–80.
- SPEIGHT, J. M., SKELHORN, R. R., SLOAN, T. & KNAAP, R. J. 1982. The dyke swarms of Scotland. *Igneous Rocks of the British Isles* (ed. D. S. Sutherland), pp. 449–59. John Wiley & Sons Ltd.
- SRIVASTAVA, S. P. 1985. Evolution of the Eurasian Basin and its implications to the motion of Greenland along Nares Strait. *Tectonophysics* **114**, 29–53.
- STOREY, M., DUNCAN, R. A., PEDERSEN, A. K., LARSEN, L. M. & LARSEN, H. C. 1998. $^{40}\text{Ar}/^{39}\text{Ar}$ geochronology of the West Greenland Tertiary volcanic province. *Earth and Planetary Science Letters* **160**, 569–86.
- STOREY, M., DUNCAN, R. A. & TEGNER, C. 2007. Timing and duration of volcanism in the North Atlantic Igneous Province: implications for geodynamics and links to the Iceland hotspot. *Chemical Geology* **241**, 264–81.
- SURLYK, F. 1990. Timing, style and sedimentary evolution of Late Palaeozoic–Mesozoic extensional basins of East Greenland. In *Tectonic Events Responsible for Britain's Oil and Gas Reserves* (eds R. F. P. Hardman & J. Brooks), pp. 107–25. Geological Society of London, Special Publication no. 55.
- TEGNER, C., BROOKS, C. K., DUNCAN, R. A., HEISTER, L. E. & BERNSTEIN, S. 2008. ^{40}Ar – ^{39}Ar ages of intrusions in East Greenland: Rift-to-drift transition over the Iceland hotspot. *Lithos* **101**, 480–500.
- TEGNER, C. & DUNCAN, R. A. 1999. ^{40}Ar – ^{39}Ar chronology for the volcanic history of the Southeast Greenland rifted margin. *Proceedings of the ODP, Scientific Results* **163**, 53–62.
- TEGNER, C., DUNCAN, R. A., BERNSTEIN, S., BROOKS, C. K., BIRD, D. K. & STOREY, M. 1998. ^{40}Ar – ^{39}Ar geochronology of Tertiary mafic intrusions along the East Greenland rifted margin: relation to flood basalts and the Iceland hotspot track. *Earth and Planetary Science Letters* **156**, 75–88.
- TESFAYE, S., HARDING, D. J. & KUSKY, T. M. 2003. Early continental breakup boundary and migration of the Afar triple junction, Ethiopia. *Geological Society of America Bulletin* **115**(9), 1053–67.
- THOMSON, K., GREEN, P. F., WHITHAM, A. G., PRICE, S. P. & UNDERHILL, J. R. 1999. New constraints on the thermal history of North-East Greenland from apatite fission track analysis. *Geological Society of America Bulletin* **111**(7), 1054–68.
- TORSKE, T. & PRESTVIK, T. 1991. Mesozoic detachment faulting between Greenland and Norway: inferences from Jan Mayen Fracture Zone system and associated alkalic volcanic rocks. *Geology* **19**, 481–4.
- TORSVIK, T. H., MOSAR, J. & EIDE, E. A. 2001. Cretaceous–Tertiary geodynamics: a North Atlantic Exercise. *Geophysical Journal International* **146**, 850–66.
- TORSVIK, T. H., VAN DER VOO, R., MEERT, J. G., MOSAR, J. & WALDERHAUG, H. J. 2001. Reconstructions of the continents around the North Atlantic at about the 60th parallel. *Earth and Planetary Science Letters* **187**, 55–69.
- TRUDE, J., CARTWRIGHT, J., DAVIES, R. J. & SMALLWOOD, J. 2003. New technique for dating igneous sills. *Geology* **31**(9), 813–6.
- TRØNNES, R. G., PLANKE, S., SUNDVOLL, B. & IMSLAND, P. 1999. Recent volcanic rocks from Jan Mayen: low-degree melt fractions of enriched northeast Atlantic mantle. *Journal of Geophysical Research* **104**(B4), 7153–68.
- TSIKALAS, F., ELDHOLM, O. & FALEIDE, J. I. 2002. Early Eocene sea floor spreading and continent–ocean boundary between Jan Mayen and Senja fracture zones in the Norwegian–Greenland Sea. *Marine Geophysical Researches* **23**, 247–70.
- UPTON, B. G. J. 1988. History of Tertiary igneous activity in the N Atlantic borderlands. In *Early Tertiary volcanism and the opening of the NE Atlantic* (eds A. C. Morton & L. M. Parson), pp. 429–53. Geological Society, London, Special Publication no. 39.
- UPTON, B. G. J., EMELEUS, C. H., REX, D. C. & THIRLWALL, M. F. 1995. Early Tertiary magmatism in NE Greenland. *Journal of the Geological Society, London* **152**, 959–64.
- UPTON, B. G. J., SKOVGAARD, A. C., MCCLURG, J., KIRSTEIN, L., CHEADLES, M., EMELEUS, C. H., WADSWORTH, W. J. & FALLICK, A. E. 2002. Picritic magmas and the Rum ultramafic complex, Scotland. *Geological Magazine* **139**(4), 437–52.
- VAN WIJK, J. W. & CLOETINGH, S. A. P. L. 2002. Basin migration caused by slow lithospheric extension. *Earth and Planetary Science Letters* **198**, 275–88.
- VAN WIJK, J. W., HUISMANS, R. S., TER VOORDE, M. & CLOETINGH, S. A. P. L. 2001. Melt generation at volcanic continental margins: no need for a mantle plume. *Geophysical Research Letters* **28**(20), 3995–8.
- VIERECK, L. G., TAYLOR, P. N., PARSON, L. M., HERTOGEN, J., GIBSON, I. L. & the ODP Leg 104 Scientific Party. 1988. Origin of the Paleogene Vøring Plateau volcanic sequence. In *Early Tertiary volcanism and the opening of the NE Atlantic* (eds A. C. Morton & L. M. Parson), pp. 69–83. Geological Society of London, Special Publication no. 39.
- WAAGSTEIN, R. 1988. Structure, composition and age of the Faeroe basalt plateau. In *Early Tertiary volcanism and the opening of the NE Atlantic* (eds A. C. Morton & L. M. Parson), pp. 225–38. Geological Society, London, Special Publication no. 39.
- WAAGSTEIN, R., GUISE, P. & REX, D. 2002. K/Ar and $^{39}\text{Ar}/^{40}\text{Ar}$ whole-rock dating of zeolite facies metamorphosed flood basalts: the upper Paleocene basalts of the Faroe Islands, NE Atlantic. In *The North Atlantic Igneous Province: Stratigraphy, Tectonic, Volcanic and Magmatic Processes* (eds D. W. Jolley & B. R. Bell), pp. 219–52. Geological Society of London, Special Publication no. 197.
- WOLFENDEN, E., EBINGER, C., YIRGU, G., DEINO, A. & AYALEW, D. 2004. Evolution of the northern Main Ethiopian rift: birth of a triple junction. *Earth and Planetary Science Letters* **224**, 213–28.

- YAXLEY, G. M. 2000. Experimental study of the phase and melting relations of homogeneous basalt + peridotite mixtures and implications for the petrogenesis of flood basalts. *Contributions to Mineralogy and Petrology* **139**, 326–38.
- ZHAO, D. 2004. Global tomographic images of mantle plumes and subducting slabs: insight into deep Earth dynamics.

- Physics of the Earth and Planetary Interiors* **146**, 3–34.
- ZIEGLER, P. A. 1989. Evolution of the North Atlantic – an overview. *The American Association of Petroleum Geologists (AAPG) Memoir* **46**, 111–29.
- ZIEGLER, P. A. 1992. Plate tectonics, plate moving mechanisms and rifting. *Tectonophysics* **215**, 9–34.