# Amalgamations of factorizations of complete graphs

Matthew Johnson
Department of Mathematics
London School of Economics
Houghton Street
London
WC2A 2AE
U.K.

e-mail: matthew@maths.lse.ac.uk

#### Abstract

Let t be a positive integer and let  $L = (l_1, \ldots, l_t)$  and  $K = (k_1, \ldots, k_t)$  be collections of nonnegative integers. A (t, K, L)-factorization of a graph is a decomposition of the graph into factors  $F_1, \ldots, F_t$  such that  $F_i$  is  $k_i$ -regular and at least  $l_i$ -edge-connected. In this paper we apply the technique of amalgamations of graphs to study (t, K, L)-factorizations of complete graphs. In particular, we describe precisely when it is possible to embed a factorization of  $K_m$  in a (t, K, L)-factorization of  $K_n$ .

keywords: factorizations, embeddings, amalgamations

### 1 Introduction

A factor of a graph is a subgraph with the same vertex set as the graph. A factorization of a graph is a set of factors with the property that the edge sets of the factors partition the edge set of the graph. In this paper we consider factorizations of complete graphs. Let t be a positive integer and let  $K = (k_1, k_2, \ldots, k_t)$  and  $L = (l_1, l_2, \ldots, l_t)$  be lists of nonnegative integers. We shall consider factorizations  $F_1, \ldots, F_t$  of the complete graph  $K_n$  in which, for  $1 \le i \le t$ ,  $F_i$  is a  $k_i$ -regular  $l_i$ -edge-connected graph. These are called (t, K, L)-factorizations. Johnstone [8] proved the following result that describes precisely when they exist.

**Theorem 1** A (t, K, L)-factorization of  $K_n$  exists if and only if

(A1) 
$$\sum_{i=1}^{t} k_i = n-1$$
,

- (A2) if n is odd, then each  $k_i$  is even,
- (A3) for  $1 \le i \le t$ ,  $l_i \le k_i$ , and

(A4) if 
$$n \ge 3$$
,  $l_i = 0$  if  $k_i = 1$ .

Johnstone proved Theorem 1 by constructing the factorizations. At the end of the next section we shall give a proof using amalgamations. Many combinatorial problems have been solved using amalgamations; see, for example, [1, 2, 3, 4, 6, 7, 11]. Let us sketch how the technique is used on graph factorizations. Consider a partition of a graph G's vertex set into subsets  $V_1, \ldots, V_r$ . Then an amalgamation of G has vertex set  $V_1, \ldots, V_r$ , and for each edge in G joining a pair of vertices in  $V_i$ ,  $1 \le i \le r$ , there is a loop on  $V_i$  in the amalgamation, and for each edge in G joining a vertex in  $V_i$  to a vertex in  $V_j$ ,  $1 \le i < j \le r$ , there is an edge  $V_i V_j$  in the amalgamation. (We can think of the amalgamation as being obtained from G by merging vertices that belong to the same subset whilst retaining all edges.)

If G has a factorization, then we can represent it as an edge-colouring with the factors as the colour classes (in this paper we frequently use the equivalence of factorizations and edge-colourings). This colouring can be transferred to an amalgamation of G—each edge of the amalgamation has the same colour as the corresponding edge of G. Henceforth when we refer to an amalgamation we mean a graph with an edge-colouring. Suppose that  $G = K_n$  and that it has a (t, K, L)-factorization. Then we can find some properties that an amalgamation of G must possess. For example we can find the number of loops on each vertex, the number of edges between each pair of vertices and the number of edges of each colour incident with each vertex. We call any edge-coloured graph that satisfies these properties an outline (t, K, L)-factorization of  $K_n$ . In Theorem 3 we prove that every outline (t, K, L)-factorization is an amalgamated graph. That is, given an outline graph G we find a (t, K, L)-factorization of  $K_n$  of which G is an amalgamation. This will allow us to give a simple proof of Theorem 1

This kind of outline/amalgamation result is a staple of papers on combinatorial amalgamations, but we were not able to apply the standard techniques (such as those used on problems on amalgamations of factorization of graphs in [4, 6, 7, 11]). An innovation of this paper is to show how a new technique for finding factorizations of graphs introduced by Hilton and Johnson [5] can be applied to amalgamations.

In the final section we use the outline/amalgamation result to solve the problem of embedding a factorization of  $K_m$  in a (t, K, L)-factorization of  $K_n$ . We describe briefly how this will be done. Suppose that we have a factorization (or an edge-colouring) of  $K_m$ . Add to it a vertex v. Join v to each vertex of  $K_m$  by (n-m) edges and put  $\binom{n-m}{2}$  loops on v to form a graph G. Complete the edge-colouring of G by colouring the edges incident with v. (Note that G can be seen to be  $K_n$  with (n-m) vertices merged.) If G is an outline (t, K, L)-factorization of  $K_n$ , then there is a (t, K, L)-factorization of  $K_n$  in which the factorization of  $K_m$  is embedded; we can think of this factorization of  $K_n$  as being obtained from G by splitting v into

(n-m) vertices. From the properties that define an outline factorization we can work back to find the properties that  $K_m$  must possess if it is to be embedded.

# 2 Amalgamated factorizations

Before we formally define amalgamations we require another definition. Let D and G be graphs. D is a detachment of G if there is a bijection  $\rho$ :  $E(D) \longrightarrow E(G)$  and a surjection  $\sigma$ :  $V(D) \longrightarrow V(G)$  such that

- if e is a loop on v in D, then  $\rho(e)$  is a loop on  $\sigma(v)$  in G,
- if e is an edge joining v and w in D and  $\sigma(v) = \sigma(w)$ , then  $\rho(e)$  is a loop on  $\sigma(v)$  in G, and
- if e is an edge joining v and w in D and  $\sigma(v) \neq \sigma(w)$ , then  $\rho(e)$  is an edge joining  $\sigma(v)$  and  $\sigma(w)$  in G.

We can think of D as being obtained from G by splitting vertices. A detachment is the opposite of an amalgamation, except that we define amalgamations on graphs which have an edge-colouring.

Let t be a positive integer. Let F and H be t-edge-coloured graphs. H is an amalgamation of F if there is a bijection  $\phi \colon E(F) \longrightarrow E(H)$  and a surjection  $\psi \colon V(F) \longrightarrow V(H)$  such that

- if e is a loop coloured i on v in F, then  $\phi(e)$  is a loop coloured i on  $\psi(v)$  in H,
- if e is an edge coloured i joining v and w in F and  $\psi(v) = \psi(w)$ , then  $\phi(e)$  is a loop coloured i on  $\psi(v)$  in H, and
- if e is an edge coloured i joining v and w in F and  $\psi(v) \neq \psi(w)$ , then  $\phi(e)$  is an edge coloured i joining  $\psi(v)$  and  $\psi(w)$  in H.

We can think of the set of vertices  $\{u : u \in V(K_n), \psi(u) = v\}$  as being merged to form v.

Let  $F_i$  and  $H_i$  be the subgraphs of F and H induced by edges coloured  $i, 1 \leq i \leq t$ . Then  $F_i$  is a detachment of  $H_i$ .

Let t, n, K and L be as defined in the Introduction. Suppose that  $F = K_n$  is t-edge-coloured and that  $F_i$  is  $k_i$ -regular and  $l_i$ -edge-connected,  $1 \le i \le t$ , (that is, the edge-colouring gives a (t, K, L)-factorization of  $K_n$ ). If H is an amalgamation of  $K_n$ , then define  $f: V(H) \longrightarrow \mathbb{N}$  by

$$f(v) = |\{u : u \in V(K_n), \psi(u) = v\}|.$$

So f counts the vertices that are merged to form v. Together H and f form an amalgamated (t, K, L)-factorization of  $K_n$ .

**Theorem 2** Let H and f be an amalgamated (t, K, L)-factorization of  $K_n$ . Then

- (B1) for all pairs of distinct vertices  $v, w \in V(H)$ , there are f(v)f(w) edges joining v to w,
- (B2) for all  $v \in V(H)$ , there are  $\binom{f(v)}{2}$  loops on v,
- (B3) for all  $v \in V(H)$ , for  $1 \le i \le t$ , v is incident with  $k_i f(v)$  edges of colour i (counting loops twice),
- (B4)  $\sum_{v \in V(H)} f(v) = n, \text{ and }$
- (B5) for  $1 \le i \le t$ ,  $H_i$  has an  $l_i$ -edge-connected  $k_i$ -regular detachment.

**Proof**: We know that f(v) vertices in  $K_n$  are merged to form v and f(w) vertices are merged to form w. In  $K_n$  there are f(v)f(w) edges between these two sets of vertices, and when the vertices are merged these edges join v to w. Hence we obtain (B1).

The subgraph of  $K_n$  induced by the f(v) vertices merged to form v is  $K_{f(v)}$  and contains  $\binom{f(v)}{2}$  edges. When the vertices are merged these edges become loops on v. Hence we obtain (B2).

In  $F_i$  the f(v) vertices merged to form v each have degree  $k_i$ . The sum of these degrees is the degree of v in  $H_i$ . Hence we obtain (B3).

As f counts the number of vertices merged to form each vertex of the amalgamation of  $K_n$  and as each vertex of  $K_n$  corresponds to exactly one of the vertices of the amalgamation, we obtain (B4).

As we noted before,  $F_i$  is an  $l_i$ -edge-connected  $k_i$ -regular detachment of  $H_i$  so (B5) is satisfied.

A t-edge-coloured-coloured graph H and a function  $f: V(H) \longrightarrow \mathbb{N}$  form an outline (t, K, L)-factorization of  $K_n$  if they satisfy (B1) to (B5). By Theorem 2, an amalgamated (t, K, L)-factorization of  $K_n$  is an outline (t, K, L)-factorization of  $K_n$ . We prove that the converse is true.

**Theorem 3** Let H and f be an outline (t, K, L)-factorization of  $K_n$ . Then H and f are an amalgamented (t, K, L)-factorization of  $K_n$ .

Before we prove Theorem 3, we must introduce an important tool first used in [5]. Let a and b be vertices each of degree d in a multigraph G. Let u be a neighbour of a and v be a neighbour of b in G. To (a, b)-swap the vertices u and v means to form a new graph from G by deleting the edges au and bv, and adding the edges av and bu. Clearly this manoeuvre leaves the degrees of all the vertices unaltered.

We can find d neighbours of a in G by counting a vertex u as a neighbour of a as many times as there are edges au. An (a,b)-swap-set is a collection of d pairs of vertices such that each neighbour of a is the first element of exactly one pair and each neighbour of b is the second element of exactly one pair. We call the pairs (a,b)-pairs. The proof of the following lemma uses an argument from [5]

**Lemma 4** If a and b are vertices each of degree d in a l-edge-connected multigraph G, then there exists an (a,b)-swap-set S such that a graph obtained from G by (a,b)-swapping any number of (a,b)-pairs in the swap-set is at least l-edge-connected.

**Proof**: First form S. In G we can find l edge-disjoint a-b paths  $au_j \cdots v_j b$ ,  $1 \leq j \leq l$ . Let  $(u_j, v_j)$  be a pair in S. For any edges ab in G not already considered as one of the paths, let (b, a) be a pair in S. Complete S by pairing off the remaining neighbours of a and b arbitrarily.

Consider a graph obtained from G by (a, b)-swapping pairs in S. It contains l edge-disjoint a-b paths since, for  $1 \leq j \leq l$ , it contains either  $au_j \cdots v_j b$  or  $bu_j \cdots v_j a$ . Now we use induction to prove the lemma. We know that G is l-edge-connected. Suppose that after some number of (a, b)swaps we have obtained a graph H that is l-edge-connected, and then we (a,b)-swap a further (a,b)-pair (u,v) to obtain a graph J. That is, au and bv are deleted in H and replaced by av and bu to obtain J. If J is not l-edge connected, then we can find a minimal edge-cutset E such that |E| < l. We show that H has an edge-cutset of the same size as E, a contradiction. Let  $C_1$  and  $C_2$  be the two connected components of J-E. In J there are l edge-disjoint a-b paths so a and b must be in the same component of J - E, say  $C_1$ . If u and v are also both in  $C_1$ , then in J-E we could reverse the (a,b)-swap of u and v to obtain H-E which would also have two components. If u and v are both in  $C_2$ , then av and bu must both be in E. Thus  $(E \setminus \{av, bu\}) \cup \{au, bv\}$  is an edge-cutset of H. Finally, suppose that u is in  $C_1$  and v is in  $C_2$ . Then  $av \in E$  and  $bu \in C_1$ . Let  $E' = (E \setminus \{av\}) \cup \{bv\}$ and  $C'_1 = (C_1 - \{bu\}) \cup \{au\}$ . Thus H - E' has two connected components,  $C_1'$  and  $C_2$ . 

**Proof of Theorem 3**: Given an outline graph H and f, we find a (t, K, L)-factorization of  $K_n$  of which H and f are an amalgamation.

By (B5), for  $1 \le i \le t$ ,  $H_i$  has an  $l_i$ -edge-connected  $k_i$ -regular detachment which we denote  $F_i$ . In this proof we refer to the subgraphs  $H_1, \ldots, H_t$  as

colour classes and to their detachments  $F_1, \ldots, F_t$  as factors. Each of the factors is a graph with n vertices so let the vertex set of each factor be  $V(K_n)$ . Label the vertices of each factor so that for each  $v \in H$  the set of vertices formed by the splitting of v when  $F_i$  is obtained from  $H_i$  is the same for each  $i, 1 \leq i \leq t$ . Let U be a graph on  $V(K_n)$  that contains each edge of each factor. Thus U has the same number of edges as  $K_n$ . To prove the theorem we show that we can alter the edge sets of some of the factors  $F_i$  in such a way that each of the graphs obtained is also an  $l_i$ -edge-connected  $k_i$ -regular detachment of  $H_i$  and the union of the new graphs is  $K_n$ .

(As we remarked in the Introduction, this method of proof differs from that used previously in outline/amalgamation theorems on graphs. In the standard proof (see, for example, [4, 6, 11]) the outline graph is "disentangled" by considering in turn each vertex v with f(v) > 1. A new graph is obtained by splitting v into two vertices  $v_1$  and  $v_2$  with  $f(v_1) = f(v) - 1$  and  $f(v_2) = 1$  in such a way that the new graph is also an outline graph. By repetition, an outline graph in which f(v) = 1 for every vertex v is obtained. Such a graph is the required factorization.)

Let  $V(H) = \{v_1, v_2, \ldots, v_r\}$ . Let  $V(K_n) = V_1 \cup V_2 \cup \cdots \cup V_r$ , where  $V_j$ ,  $1 \leq j \leq r$ , is the set of vertices  $\{u_{j1}, u_{j2}, \ldots, u_{jf(v_j)}\}$  that was formed by the splitting of the vertex  $v_j$  in each  $H_i$ . We call these smaller vertex sets sets of split vertices. Notice that two subgraphs of  $K_n$  are both detachments of the same colour class if and only if for each pair of sets of split vertices  $V_j$  and  $V_z$ ,  $1 \leq j \leq z \leq r$ , the number of edges that join a vertex in  $V_j$  to a vertex in  $V_z$  is the same in each subgraph. From the definition of an outline factorization we find that

- (B1') for all pairs of distinct sets of split vertices  $V_j$  and  $V_z$ , in U there are  $f(v_j)f(v_z)$  edges joining vertices in  $V_j$  to vertices in  $V_z$ , and
- (B2') for all sets of split vertices  $V_j$ , there are  $\binom{f(v_j)}{2}$  edges in the subgraph of U induced by the vertices of  $V_j$ .

Now we alter the factors to obtain a factorization of  $K_n$ . Note that we shall refer to  $F_i$  before and after each alteration by the same name, and we shall also refer to the altered graphs as factors and define U in terms of the altered graphs. Our aim is to alter the factors so that  $U = K_n$ . The factors may have loops, and removing them is the first alteration we make. Suppose that there is a loop on a vertex a in  $F_i$ . Let  $V_z$  be the set of split vertices that contains a. By (B2'),  $|V_z| \ge 2$  so there is a vertex  $b \in V_z$ ,  $a \ne b$ . If there is also a loop on b, then we can delete the loops and replace them with two edges joining a to b. Clearly  $F_i$  is still  $k_i$ -regular and its edge-connectivity has not decreased. If there is no loop on b, then find  $l_i$  disjoint a-b paths (choosing edges ab if possible). We must have  $l_i < k_i$  (else there could not be any loops), so we can find an edge bu that is not in one of these a-b paths and  $u \neq a$  (as there is a loop on a, and a and b have the same degree, b must be adjacent to a vertex other than a). Delete the loop on a and bu and add edges ab and au. The new graph is  $k_i$ -regular and has one fewer loop than the original graph. We must check that the new graph is  $l_i$ -edge-connected. If it is not, then there is a minimal edge-cutset E,  $|E| < l_i$ . Note that Ecannot separate a and b since they are joined by  $l_i$  disjoint paths. Thus  $ab \notin E$ . If  $au \notin E$ , then E is a cutset of the original graph, and if  $au \in E$ , then  $(E \setminus \{au\}) \cup bu$  is a cutset of the original graph; a contradiction since the original graph was  $l_i$ -edge-connected.

By repetition we obtain a set of loopless factors. Note that in each case the new factor is still a detachment of the corresponding colour class.

By Lemma 4 for  $1 \le i \le t$ , if a and b are vertices in  $F_i$ , then we can find a set  $S_i(a,b)$  that is a collection of  $k_i$  (a,b)-pairs such that

- each neighbour of a in  $F_i$  is the first element of exactly one pair and each neighbour of b is the second element of exactly one pair,
- there are  $l_i$  pairs  $(u_j, v_j)$  such that there exist in  $F_i$  edge-disjoint paths  $au_j \cdots v_j b$ ,  $1 \leq j \leq l_i$ , and
- for each edge ab in  $F_i$ , there is an (a, b)-pair (b, a).

Note that if a and b are in the same set of split vertices  $V_j$ , then any graph obtained from a factor  $F_i$  by (a, b)-swapping a pair (u, v) in  $S_i(a, b)$  is also a detachment of the corresponding colour class  $H_i$  since we delete an edge, au, that joins u to a vertex in  $V_j$  and replace it with another edge, bu, that also joins u to a vertex in  $V_j$ . Similarly for v. Also by Lemma 4, any graph obtained from  $F_i$  by (a, b)-swapping pairs in  $S_i(a, b)$  is  $l_i$ -edge-connected. So if we alter the factors using only (a, b)-swaps for pairs of vertices a and b in the same set of split vertices and we obtain a (t, K, L)-factorization of  $K_n$ , then H and f will be an amalgamation of this factorization. We show how this is done.

There are two further stages to the proof. We will often say informally that two disjoint sets of vertices V and V' are joined by the *correct* number of edges if they are joined by |V||V'| edges, that is, the number of edges between them in  $K_n$ . In the next stage of the proof we alter the factors so that each vertex is joined the correct number of times to each set of split vertices. That is, we alter the factors so that they satisfy

(C1) in U, for  $1 \le j \le r$ ,  $1 \le h \le f(v_j)$ ,  $u_{jh}$  is joined by  $f(v_j) - 1$  edges to vertices of  $V_j$  and, for  $1 \le z \le r$ ,  $z \ne j$ , by  $f(v_z)$  edges to vertices in  $V_z$ .

We then complete the proof by further altering the edge sets of the factors so that

(C2) in U each pair of distinct vertices is joined by exactly one edge.

In other words,  $U = K_n$ .

First we alter the factors so that (C1) is satisfied. For any vertex  $a \in V$ , for  $1 \le j \le r$ ,

- let  $p(a, V_j)$  be the number of edges in U that join a to a vertex in the set of split vertices  $V_j$ ,
- let  $q(a, V_i) = f(v_i)$  if  $a \notin V_i$ , let  $q(a, V_i) = f(v_i) 1$  if  $a \in V_i$ .

That is,  $q(a, V_j)$  is the number of edges that will join a to vertices in  $V_j$  in U when  $U = K_n$ . Thus to satisfy (C1) we must alter the factors so that for each vertex  $a \in V(K_n)$ , for  $1 \le j \le r$ ,  $p(a, V_j) = q(a, V_j)$ . Let the set-discrepancy  $\delta_s$  be defined by

$$\delta_s = \sum_{a \in V(K_n)} \sum_{j=1}^r |p(a, V_j) - q(a, V_j)|.$$

(C1) is satisfied when the set-discrepancy is zero. We describe a method that will reduce the set-discrepancy if it is greater than zero. By applying it repeatedly we obtain a set of factors that satisfies (C1).

Let j and z be fixed. By (B1') and (B2') each pair of sets of split vertices is joined by the correct number of edges. Thus

$$\sum_{a \in V_z} p(a, V_j) = \sum_{a \in V_z} q(a, V_j). \tag{1}$$

If the set-discrepancy is greater than zero, then for some vertex a and some  $z_1$ ,  $p(a, V_{z_1}) \neq q(a, V_{z_1})$ . We can assume that

$$p(a, V_{z_1}) > q(a, V_{z_1}),$$
 (2)

since by (1) this implies, and is implied by, the existence of a vertex b in the same set of split vertices as a such that

$$p(b, V_{z_1}) < q(b, V_{z_1}). (3)$$

Using the sets  $S_i(a,b)$ ,  $1 \le i \le t$ , we create a further set, S(a,b). For  $1 \le i \le t$ , if  $(c,d) \in S_i(a,b)$ , then  $(i,c,d) \in S(a,b)$ . So S(a,b) contains ordered triples (i,c,d) where c is a neighbour of a and d is a neighbour of b in  $F_i$ . Note that there is an obvious one-to-one relationship between the triples of S(a,b) and the neighbours, over all the factors, of a, and also between the triples of S(a,b) and the neighbours, over all the factors, of b.

Claim 5 There is a sequence of sets of split vertices

$$\Gamma = V_{z_1}, V_{z_2}, \dots, V_{z_m}$$

such that

- (D1)  $V_{z_{\alpha}} \neq V_{z_{\beta}}$  if  $\alpha \neq \beta$ ,
- (D2) either  $p(a, V_{z_m}) < q(a, V_{z_m})$  or  $p(b, V_{z_m}) > q(b, V_{z_m})$ , and
- (D3) for  $2 \leq j \leq m$ , there is a triple  $(i_j, c_j, d_j) \in S(a, b)$  where  $c_j \in V_{z_{j-1}}$  and  $d_j \in V_{z_j}$ .

The claim is proved below. First we use it to reduce  $\delta_s$ . For  $2 \leq j \leq m$ , we (a,b)-swap  $c_j$  and  $d_j$  in  $F_{i_j}$ : the edges  $ac_j$  and  $bd_j$  are deleted and replaced with the edges  $ad_j$  and  $bc_j$ . Each new factor  $F_i$  obtained is clearly  $k_i$ -regular and, by Lemma 4, it is  $l_i$ -edge-connected. It is also a detachment of the corresponding colour class  $H_i$  since the number of edges in the factor between each pair of sets of split vertices does not change.

For  $2 \leq j \leq m-1$ , an edge from a to a vertex,  $c_{j+1}$ , that is in  $V_{z_j}$ , has been deleted and an edge from a to a vertex,  $d_j$ , that is in  $V_{z_j}$  has been added. Thus  $p(a, V_{z_j})$  is unchanged. Similarly  $p(b, V_{z_j})$ ,  $2 \leq j \leq m-1$ , is unchanged.

The only neighbour of a in  $V_{z_1}$  involved in an (a, b)-swap is  $c_2$ . The edge  $ac_2$  is deleted so  $p(a, V_{z_1})$  is reduced by 1. Hence, by (2),  $\delta_s$  is also reduced by 1. The addition of  $bc_2$  causes  $p(b, V_{z_1})$  to increase by 1, so by (3),  $\delta_s$  decreases further by 1.

The only neighbour of b in  $V_{z_m}$  involved in an (a, b)-swap is  $d_m$ . Consider (D2). If  $p(a, V_{z_m}) < q(a, V_{z_m})$ , then the addition of  $ad_m$  causes  $p(a, V_{z_m})$  to increase by 1, and  $\delta_s$  is reduced further by 1. The deletion of  $bd_m$  may cause  $\delta_s$  to increase by 1, but at worst  $\delta_s$  is reduced by 2 overall. The only other possibility is that  $p(b, V_{z_m}) > q(b, V_{z_m})$ , and by a similar argument  $\delta_s$  is reduced overall by at least 2 in this case also.

We show that the factors remain loopless. A loop is put on a only if one of the triples is  $(i_j, c_j, d_j)$  with  $d_j = a$ . That is,  $(c_j, a)$  is a pair in  $S_{i_j}(a, b)$ . Recall that a is the second element of a pair in  $S_i(a, b)$  only if b is the first element. But if  $c_j = b$ , then  $c_j$  and  $d_j$  are in the same set of split vertices, a contradiction by (D1) and (D3). By a similar argument b also remains loopless.

**Proof of Claim 5**: In fact we shall prove that there is a sequence of sets of split vertices

$$\Delta = V_{g_1}, V_{g_2}, \dots, V_{g_{m'}}$$

such that

- (E1)  $V_{g_1} = V_{z_1}$ ,
- (E2)  $V_{g_{\alpha}} \neq V_{g_{\beta}}$  if  $\alpha \neq \beta$ ,
- (E3) either  $p(a, V_{g_{m'}}) < q(a, V_{g_{m'}})$  or  $p(b, V_{g_{m'}}) > q(b, V_{g_{m'}})$ , and
- (E4) for  $2 \leq j \leq m'$ , there is a triple  $(i_j, c_j, d_j) \in S(a, b)$  where  $c_j \in V_{g_h}$  for some  $h \in \{1, 2, \dots, j-1\}$  and  $d_j \in V_{g_j}$ .

It is easy to see that  $\Delta$  has a subsequence that has  $V_{g_1} = V_{z_1}$  as the first term and satisfies (D1), (D2) and (D3). (Let  $V_{g_{m'}}$  be the final term and work backwards. If  $V_{g_{\alpha}}$  is the last term reached, then if  $\alpha = 1$  the subsequence is found. Otherwise there is a triple  $(i_{\alpha}, c_{\alpha}, d_{\alpha})$ . Let the previous term of the sequence be the set of split vertices  $V_{g_{\beta}}$  that contains  $c_{\alpha}$ . As  $\beta < \alpha$  we must eventually get back to  $V_{g_1}$ .)

We find  $\Delta$ . The first term  $V_{g_1} = V_{z_1}$  was found before the claim was stated. Suppose that we have found the first  $\omega$  terms, and that this sequence of  $\omega$  terms satisfies (E1), (E2) and (E4) with  $m' = \omega$ . If for any  $\alpha \in \{1, 2, ..., \omega\}$ 

$$p(a, V_{g_{\alpha}}) < q(a, V_{g_{\alpha}}), \text{ or }$$
  
 $p(b, V_{g_{\alpha}}) > q(b, V_{g_{\alpha}}),$ 

then we pick the smallest such  $\alpha$  and let  $\Delta = V_{g_1}, V_{g_2}, \dots, V_{g_{\alpha}}$  as this also satisfies (E3). Otherwise, for  $1 \leq j \leq \omega$ ,

$$p(a, V_{g_j}) \geq q(a, V_{g_j}), \tag{4}$$

$$p(b, V_{g_j}) \leq q(b, V_{g_j}). \tag{5}$$

Let  $W = V_{g_1} \cup V_{g_2} \cup \cdots \cup V_{g_{\omega}}$ . As a and b are in the same set of split vertices,  $q(a, V_j) = q(b, V_j)$ ,  $1 \leq j \leq r$ . Therefore, by (2) and (3), a has more neighbours than b in  $V_{g_1}$  and, by (4) and (5), a has at least as many neighbours as b in  $V_{g_j}$ ,  $2 \leq j \leq \omega$ . Therefore over all the factors a has more neighbours than b in W. Recall that in S(a, b) there is a triple corresponding to each neighbour of a in each factor; similarly there is a triple corresponding to each neighbour of b. So there is a triple  $(i_{\omega+1}, c_{\omega+1}, d_{\omega+1}) \in S(a, b)$ , such that  $c_{\omega+1} \in W$  and  $d_{\omega+1} \notin W$ . Let the set of split vertices containing  $d_{\omega+1}$  be  $V_{g_{\omega+1}}$ . Then  $V_{g_{\omega+1}} \neq V_{g_j}$ ,  $1 \leq j \leq \omega$ , since  $V_{g_{\omega+1}} \not\subset W$ .

We must eventually find a set of split vertices that satisfies (E3): note that

$$\sum_{j=1}^{r} p(a, V_j) = \sum_{j=1}^{r} q(a, V_j), \tag{6}$$

since both sums are equal to n-1, the sum of the degrees of a taken over all the factors. As  $p(a, V_{z_1}) > q(a, V_{z_1})$ , there is at least one set of split vertices  $V_z$  such that  $p(a, V_z) < q(a, V_z)$  and therefore  $V_z$ , at least, satisfies (E3). This completes the proof of Claim 5.

We must now show that when (C1) is satisfied we can further alter the factors so that (C2) is also satisfied. For a pair of distinct vertices a and c, let p(a,c) be the number of edges in U from a to c. Note that p(a,c) = p(c,a).

Let the vertex-discrepancy  $\delta_v$  be defined by

$$\delta_v = \sum_{ac \in E(K_n)} |p(a, c) - 1|.$$

If (C2) is satisfied, then for all pairs of distinct vertices a and c, p(a, c) = 1, and the vertex-discrepancy is zero. We describe a method that will reduce the vertex-discrepancy if it is greater than zero. By applying it repeatedly we shall obtain a set of factors that satisfies (C2).

We can see that if c is the only vertex in a set of split vertices  $V_z$ , then p(a,c) = 1: let a be some other vertex; as (C1) is satisfied,  $p(a,V_z) = q(a,V_z) = f(v_z) = 1$ , and as  $p(a,c) = p(a,V_z)$ , we already have p(a,c) = 1.

**Claim 6** Suppose that a and b are vertices in the same set of split vertices, that  $c_1 \notin \{a, b\}$  and that

$$p(a, c_1) > 1, (7)$$

$$p(b, c_1) < 1. (8)$$

Let S(a,b) be defined as before. There is a sequence of vertices  $c_1, c_2, \ldots, c_m$  such that

- (F1)  $c_i \notin \{a, b\}, 2 \le j \le m$ ,
- (F2)  $c_{\alpha} \neq c_{\beta}$  if  $\alpha \neq \beta$ ,
- (F3) either  $p(a, c_m) < 1$  or  $p(b, c_m) > 1$ , and
- (F4) for  $1 \leq j \leq m-1$  there is a triple  $(i_j, c_i, c_{i+1}) \in S(a, b)$ .

**Proof**: The first term of the sequence is known by the hypothesis. Suppose that we have found the first  $\omega$  terms and that this sequence satisfies (F1), (F2) and (F4) with  $m = \omega$ . If for some  $h \in \{1, 2, ..., \omega\}$ 

$$p(a, c_h) < 1$$
, or  $p(b, c_h) > 1$ ,

then choose the smallest such h and let the complete sequence be  $c_1, c_2, \ldots, c_h$  since this also satisfies (F3) with m = h. Otherwise, for  $1 \le j \le \omega$ ,

$$p(a, c_j) \ge 1,$$
  
$$p(b, c_i) < 1.$$

As  $p(a, c_{\omega}) \geq 1$  we can find a triple  $(i_{\omega}, c_{\omega}, c_{\omega+1}) \in S(a, b)$ . As there are no loops and  $c_{\omega+1}$  is a neighbour of b,  $c_{\omega+1} \neq b$ . By (F1),  $c_{\omega} \neq b$  and a is the second element of a pair in  $S_{i_{\omega}}(a, b)$  only if b is the first element, so  $c_{\omega+1} \neq a$ . By (8),  $p(b, c_1) = 0$ , so  $c_{\omega+1} \neq c_1$ . As  $p(b, c_j) \leq 1$ ,  $2 \leq j \leq \omega$ , there is at most one triple in S(a, b) with  $c_j$  as the third element and we have already found one such triple (namely  $(i_{j-1}, c_{j-1}, c_j)$ ). Therefore  $c_{\omega+1} \neq c_j$ ,  $2 \leq j \leq \omega$ .

The sequence must terminate: there is a finite number of vertices and it is easily seen that  $p(a, c_1) > 1$  implies that for some vertex c, p(a, c) < 1 (that is, if a vertex a is joined too many times to one vertex, then it must be joined too few times to some other vertex). This completes the proof of Claim 6.

We describe how to use the claim to reduce the vertex discrepancy. First choose a set of split vertices  $V_z$  such that

(C1a) for every vertex 
$$c \notin V_z$$
,  $p(c, V_j) = q(c, V_j)$ ,  $1 \le j \le r$ .

Note that (C1) implies (C1a) so initially we can choose any set of split vertices as  $V_z$ . If possible choose a pair of vertices  $a \in V_z$ ,  $c_1 \notin V_z$  that satisfy (7). By (C1a) there is a vertex  $b \in V_z$ ,  $a \neq b$ , that satisfies (8). Therefore, by Claim 6, there is a sequence of vertices  $c_1, c_2, \ldots, c_m$  that satisfies (F1) to (F4). For  $1 \leq j \leq m-1$ , (a,b)-swap  $(c_j, c_{j+1})$  in  $F_{i_j}$ . For  $2 \leq j \leq m-1$ , we add  $ac_j$  to  $F_{i_{j-1}}$  and delete  $ac_j$  from  $F_{i_j}$ , so  $p(a,c_j)$  is unchanged. Similarly  $p(b,c_j)$  is unchanged,  $2 \leq j \leq m-1$ . By (7), the deletion of  $ac_1$  reduces  $\delta_v$  by 1, and, by (8), the addition of  $bc_1$  reduces  $\delta_v$  further by 1. By (F3), the addition of  $ac_m$  and the deletion of  $bc_m$  at worst has no net effect on  $\delta_v$ . So overall  $\delta_v$  is reduced by at least 2. As  $c_j \notin \{a,b\}$ ,  $1 \leq j \leq m$ , no loops are created.

Consider the effect of these (a, b)-swaps on the set discrepancy. Let  $V_{z_j}$  be the set of split vertices that contains  $c_j$ ,  $1 \le j \le m$ . For  $2 \le j \le m - 1$ ,  $p(a, c_j)$  and  $p(b, c_j)$  were unchanged so  $p(a, V_{z_j})$  and  $p(b, V_{z_j})$  are unchanged. Note that

$$p(a, V_{z_1})$$
 and  $p(b, V_{z_m})$  are reduced by 1, and (9)

$$p(a, V_{z_m})$$
 and  $p(b, V_{z_1})$  are increased by 1. (10)

As  $a, b \in V_z$ , (C1a) remains satisfied. So we can look for further pairs  $a \in V_z$ ,  $c_1 \notin V_z$  that satisfy (7) and repeat the procedure. When no such pairs

remain we have p(a,c)=1 for every  $a \in V_z$ ,  $c \notin V_z$ . For  $1 \leq j \leq r$ ,  $j \neq z$ ,  $p(a,V_j)=\sum_{c \in V_j} p(a,c)=|V_j|$ . Thus  $p(a,V_j)=q(a,V_j)$ ,  $1 \leq j \leq r$ ,  $j \neq z$ . By (6), this implies that  $p(a,V_z)=q(a,V_z)$  also. Thus

(C1b) for every vertex  $a \in V_z$ ,  $p(a, V_j) = q(a, V_j)$ ,  $1 \le j \le r$ .

Note that (C1a) and (C1b) together imply (C1).

Now if possible choose a pair  $a \in V_z$ ,  $c \in V_z$  that satisfies (7). By (C1b), there is a vertex  $b \in V_z$  that satisfies (8), so we can use the claim and the method of (a, b)-swapping just described to reduce  $\delta_v$ . Note that  $V_{z_1} = V_z$  (since  $V_{z_1}$  is the set that contains  $c_1$ ) and that  $V_{z_m} = V_z$  (since  $V_{z_m}$  is the set that contains  $c_m$ ,  $c_m$  satisfies (F3) and we know that p(a, c) = 1 for all  $a \in V_z$ ,  $c \notin V_z$ ). Thus (9) and (10) cancel each other out and (C1a) and (C1b) remain satisfied. Look for further pairs  $a, c_1 \in V_z$  that satisfy (7) and reduce  $\delta_v$  further. When no such pairs remain since (C1a) and (C1b) are satisfied, (C1) is satisfied and we can begin the process again with another choice of  $V_z$ . Eventually  $\delta_v$  is reduced to zero and (C2) is satisfied. This completes the proof of Theorem 3

**Proof of Theorem 1**: The following four sentences prove the necessity of (A1) to (A4). The sum, taken over all the factors, of the degrees of a vertex is equal to its degree in  $K_n$ . By the handshaking lemma, a regular graph with an odd number of vertices has even degree. As the set of all edges incident with a vertex forms an edge-cutset of a graph, the edge-connectivity of a graph is at most its minimum degree. A 1-regular graph is a set of independent edges and is not connected if it has more than 2 vertices.

By Theorem 2, we can show that the conditions are sufficient by finding an outline (t, K, L)-factorization, H and f. Let  $V(H) = \{v\}$ . Let there be  $\binom{n}{2}$  loops on v, and let  $nk_i/2$  of the loops be coloured i,  $1 \le i \le t$ . Let f(v) = n. It is easy to check that H and f satisfy (B1) to (B5).

## 3 Embedding factorizations

Here we answer this question: when can a factorization  $G_1, \ldots, G_t$  of  $K_m$  be embedded in a (t, K, L)-factorization  $F_1, \ldots F_t$  of  $K_n$ . By embed we mean that the vertices of  $K_m$  are identified with m of the vertices of  $K_n$  in such a way that  $G_i$  is a subgraph of  $F_i$ ,  $1 \leq i \leq t$ . Note that we can think of  $G_1, \ldots, G_t$  as the colour classes of a t-edge-colouring of  $K_m$ . In some cases a solution to the embedding problem is known. When each  $l_i = 0$ , that is, when there is no constraint on the connectivity of the factors, the solution was found by Andersen and Hilton [2] (and independently by Rodger and Wantland [11]). A solution in the case where each  $l_i = 1$  was found by Hilton, Johnson, Rodger and Wantland [6]. Solutions when each  $l_i = 2$  are also known: Hilton [4] solved the subcase where each  $k_i = 2$ , and this was generalized by Rodger and Wantland [11] (it also follows from a result of Nash-Williams [9]). Below in Theorem 8 we solve the general case where t, K and L are required to satisfy (G1) to (G4).

First we need a result about detachments. Recall that, by (B5), if H is a (t, K, L)-outline factorization, then each  $H_i$  (the subgraph induced by edges coloured i) has an  $l_i$ -edge-connected  $k_i$ -regular detachment. Theorem 7 is a result of Nash-Williams that characterizes those graphs that have such detachments (in fact, this is a specialization of a much more general result). We need some definitions first. Let G be a graph of which we seek to find a detachment. We define three functions  $f, c, e: \mathcal{P}(V(G)) \longrightarrow \mathbb{Z}$ ,  $(\mathcal{P}(V(G)))$  is the power set of V(G). For each set of vertices  $V \subseteq V(G)$ , let f(V) be the total number of vertices we wish to split the vertices of V into, let C(V) be the number of components in C - V, and let C(V) be the number of edges (including loops) that are incident with at least one vertex in V (loops and edges incident twice with vertices in V are only counted once).

**Theorem 7** [10] Let k and l be nonnegative integers. Let G be a graph in which the degree of each vertex is a multiple of k. Then G has an l-edge-connected k-regular detachment if and only if

- (X1) G is l-edge-connected,
- (X2) if l = 1, then for all  $V \subseteq V(G)$ ,  $f(V) + c(V) \le e(V) + 1$ ,
- (X3) if l is odd and l = k, then G has no cutvertex with degree 2l, and
- (X4) if l is odd and l = k, then G is not a loopless graph that contains exactly two vertices each with degree 2l.

We need some more definitions before we state the embedding result. Let  $\omega_i$  be the number of connected components of  $G_i$  and let these components be  $C_{i,1}, C_{i,2}, \ldots, C_{i,\omega_i}$ . Let  $\varepsilon_{i,j} = \sum_{v \in V(C_{i,j})} k_i - d_{G_i}(v)$ , and let  $\varepsilon_i = \sum_{j=1}^{\omega_i} \varepsilon_{i,j}$ . Let  $r_{i,j}$  be the number of minimal edge-cutsets of  $C_{i,j}$  that contain fewer than  $l_i$  edges, let these sets be  $E_1^{i,j}, E_2^{i,j}, \ldots, E_{r_{i,j}}^{i,j}$ , and let  $C_{m_1}^{i,j}$  and  $C_{m_2}^{i,j}$  be the connected components of  $C_{i,j} - E_m^{i,j}$ . Let  $\varepsilon_{i,j,m_p} = \sum_{v \in V(C_{m_p}^{i,j})} k_i - d_{G_i}(v)$ .

**Theorem 8** Let n, t, K and L satisfy (G1) to (G4) and let  $\alpha = n - m$ . A t-edge-coloured  $K_m$  can be embedded in a (t, K, L)-factorization of  $K_n$  if and only if

- (I)  $d_{G_i}(v) \leq k_i$  for each  $v \in V(K_m)$ , for  $1 \leq i \leq t$ ,
- (II)  $\varepsilon_{i,j} \geq l_i$  for  $1 \leq i \leq t$ ,  $1 \leq j \leq \omega_i$ ,
- (III)  $\alpha \ge \max\{\varepsilon_i/k_i : 1 \le i \le t\},\$

(IV) for 
$$1 \le i \le t$$
, if  $l_i = 1$ , then  $\alpha \ge \frac{2\omega_i - \varepsilon_i - 2}{k_i - 2}$ ,

- (V) for  $1 \le i \le t$ , if  $l_i = k_i$ ,  $l_i$  is odd and  $\omega_i \ge 2$ , then  $\alpha \ne 2$ , and
- (VI)  $\varepsilon_{i,j,m_p} \ge l_i |E_m^{i,j}|$ , for  $1 \le i \le t$ ,  $1 \le j \le \omega_i$ ,  $1 \le m \le r_{i,j}$ ,  $1 \le p \le 2$ .

**Proof**: Necessity: suppose that a t-edge-coloured  $K_m$  is embedded in an (t, K, L)-factorization of  $K_n$  (so each  $G_i$  is a subgraph of a  $k_i$  regular  $l_i$ -edge-connected graph  $F_i$ ). We show that the conditions of the theorem hold.

For  $1 \leq i \leq t$ , as  $G_i$  is a subgraph of a  $k_i$ -regular graph,  $d_{G_i}(v) \leq k_i$  for each  $v \in V(K_m)$ . So (I) holds.

By definition,  $\varepsilon_{i,j}$  is the number of edges incident with vertices of  $C_{i,j}$  in  $E(F_i) \setminus E(G_i)$ . These edges all join  $C_{i,j}$  to  $V(K_n) \setminus V(K_m)$  and form an edge-cutset so there must be at least  $l_i$  of them. So (II) holds.

Similarly,  $\varepsilon_i$  is the number of edges incident with vertices of  $G_i$  in  $E(F_i) \setminus E(G_i)$ , and all these edges join  $G_i$  to one of the  $\alpha$  vertices of  $V(K_n) \setminus V(K_m)$  which each have degree  $k_i$ . Thus  $\varepsilon_i \leq k_i \alpha$ . So (III) holds.

If  $l_i = 1$ , then from  $F_i$  form a graph J by merging vertices that belong to the same component in  $G_i$  and deleting any loops on these merged vertices. Thus J contains the  $\alpha$  vertices of  $V(K_n) \setminus V(K_m)$  plus  $\omega_i$  vertices corresponding to the  $\omega_i$  components of  $G_i$ . Its edge set contains  $\varepsilon_i$  edges corresponding to the  $\varepsilon_i$  edges in  $F_i$  that join vertices of  $V(K_m)$  to vertices of  $V(K_n) \setminus V(K_m)$ . It also contains the edges of  $F_i$  joining pairs of vertices in  $V(K_n) \setminus V(K_m)$ ; there are  $(\alpha k_i - \varepsilon_i)/2$  such edges since there are  $\alpha$  vertices with degree  $k_i$  and all but  $\varepsilon_i$  of the sum of their degrees is due to edges joining pairs of these vertices. As J is connected we must have that  $|V(J)| \leq |E(J)| + 1$ . Thus  $\alpha + \omega_i \leq \varepsilon_i + (\alpha k_i - \varepsilon_i)/2 + 1$ . Rearranging we see that (IV) holds.

Suppose that  $\alpha = 2$  and  $l_i = k_i$  is odd. If  $\omega_i \geq 2$ , then  $C_{i,1}$  and  $C_{i,2}$  are two components of  $G_i$ . There must be  $k_i$  distinct paths from  $C_{i,1}$  to  $C_{i,2}$  which each go through  $V(K_n) \setminus V(K_m) = \{w_1, w_2\}$ . We can assume that at least  $\left\lceil \frac{k_i}{2} \right\rceil$  of these paths contain  $w_1$ . But then  $d_{F_i}(w_1) \geq 2 \left\lceil \frac{k_i}{2} \right\rceil = k_i + 1$ , a contradiction. So (V) holds.

For  $1 \leq i \leq t$ ,  $1 \leq j \leq \omega_i$ ,  $1 \leq m \leq r_{i,j}$ , there must be  $l_i$  distinct paths from  $C_{m_1}^{i,j}$  to  $C_{m_2}^{i,j}$ . We know that  $|E_m^{i,j}|$  of these paths are in  $C_{i,j}$ . The remainder must go through  $V(K_n) \setminus V(K_m)$ . Therefore there must be at least  $l_i - |E_m^{i,j}|$  edges from each of  $C_{m_1}^{i,j}$  and  $C_{m_2}^{i,j}$  to  $V(K_n) \setminus V(K_m)$ . So (VI) holds as  $\varepsilon_{i,j,m_p}$  is the number of edges incident with vertices of  $C_{m_p}^{i,j}$  in  $E(F_i) \setminus E(G_i)$ .

Sufficiency: to complete the proof we must show we can find an embedding if the six conditions hold. From  $K_m$  we form H and f, an outline

(t, K, L)-factorization of  $K_n$ . Let  $V(H) = V(K_m) \cup \{v_0\}$ . Let  $f(v_0) = \alpha$ , let f(v) = 1 for all  $v \in V(K_m)$ . The edge set of H contains the edges of  $K_m$  (which are already coloured) and

- for  $1 \leq i \leq t$ , for each  $v \in V(K_m)$ , there are  $k_i d_{G_i}(v)$  edges coloured i from  $v_0$  to v, and
- for  $1 \le i \le t$ , there are  $(\alpha k_i \varepsilon_i)/2$  loops coloured i on  $v_0$ .

If we can prove that H and f are an outline (t, K, L)-factorization of  $K_n$ , then the proof is completed by applying Theorem 2 since any (t, K, L)-factorization  $F_1, \ldots, F_t$  of  $K_n$  of which H and f is an amalgamation is such that  $G_i$  is a subgraph of  $F_i$ .

First we check that the number of loops added of each colour is an integer. As  $\alpha = n - m$ ,

$$\frac{\alpha k_i - \varepsilon_i}{2} = \frac{(n-m)k_i - \varepsilon_i}{2}$$
$$= \frac{k_i n}{2} - \varepsilon_i - \frac{k_i m - \varepsilon_i}{2}$$

which is an integer since  $k_i n$  is even (by (G2)) and  $(k_i m - \varepsilon_i)/2 = |E(G_i)|$ .

We must show that H and f satisfy (B1) to (B5).

For  $v, w \in V(K_m)$ , there is 1 = f(v)f(w) edge joining v to w. For  $v \in V(K_m)$ , the number of edges from v to  $v_0$  is

$$\sum_{i=1}^{t} (k_i - d_{G_i}(v)) = \sum_{i=1}^{t} k_i - \sum_{i=1}^{t} d_{G_i}(v)$$

$$= (n-1) - (m-1)$$

$$= \alpha$$

$$= f(v) f(v_0).$$

So (B1) is satisfied.

For  $v \in V(K_m)$  there are  $0 = \binom{f(v)}{2}$  loops on v. The number of loops on  $v_0$  is

$$\sum_{i=1}^{t} \frac{\alpha k_i - \varepsilon_i}{2} = \sum_{i=1}^{t} \frac{\alpha k_i}{2} - \sum_{i=1}^{t} \sum_{v \in V(K_m)} \frac{k_i - d_{G_i}(v)}{2}$$

by the definition of  $\varepsilon_i$ . The order in which we evaluate the double sum is not important, and

$$\sum_{i=1}^{t} \frac{\alpha k_i}{2} - \sum_{v \in V(K_m)} \sum_{i=1}^{t} \frac{k_i - d_{G_i}(v)}{2} = \frac{\alpha(n-1)}{2} - \sum_{v \in V(K_m)} \frac{(n-1) - (m-1)}{2}$$

$$= \frac{\alpha(n-1)}{2} - \sum_{v \in V(K_m)} \frac{\alpha}{2}$$

$$= \frac{\alpha(n-1)}{2} - \frac{\alpha m}{2}$$

$$= \frac{\alpha(n-1) - m}{2}$$

$$= \frac{\alpha(\alpha-1)}{2}$$

$$= \frac{\alpha(\alpha-1)}{2}$$

$$= \binom{\alpha}{2}$$

$$= \binom{f(v_0)}{2}.$$

So (B2) is satisfied.

For  $v \in V(K_m)$  there are  $d_{G_i}(v) + (k_i - d_{G_i}(v)) = k_i = k_i f(v)$  edges of each colour incident with v. The number of edges of each colour incident with  $v_0$  is

$$\left(\sum_{v \in V(K_m)} (k_i - d_{G_i}(v))\right) + \alpha k_i - \varepsilon_i = \varepsilon_i + \alpha k_i - \varepsilon_i$$

$$= \alpha k_i$$

$$= k_i f(v_0).$$

So (B3) is satisfied.

As 
$$\sum_{v \in V(H)} f(v) = m + \alpha = m + n - m = n$$
, (B4) is satisfied.

Finally to show that (B5) is satisfied we must show that each  $H_i$  has an  $l_i$ -edge-connected  $k_i$ -regular detachment. Thus we must show that each  $H_i$  satisfies (X1) to (X4).

First we show that each  $H_i$  is  $l_i$ -edge-connected. Suppose that  $H_i$  is not  $l_i$ -edge-connected. Then there is a minimal edge-cutset E such that  $|E| < l_i$ . As E is minimal it will contain only edges from one component of  $G_i$ , say  $C_{i,1}$ , and perhaps also edges from  $v_0$  to  $C_{i,1}$ . It cannot contain only edges from  $v_0$  to  $C_{i,1}$  since it would need to contain all of them and there are  $\sum_{v \in V(C_{i,j})} (k_i - d_{G_i}(v)) = \varepsilon_{i,j} \text{ such edges and, by (II), } \varepsilon_{i,j} \geq l_i. \text{ The edges of } v \in V(C_{i,j})$ 

E contained in  $C_{i,1}$  form one of its minimal separating sets, say  $E_1^{i,1}$ , and we can assume that the two components of  $H_i - E$  are  $C_{1_1}^{i,1}$  and  $H_i - C_{1_1}^{i,1}$ . Therefore E must also contain all the edges from  $C_{1_1}^{i,1}$  to  $v_0$ . There are  $\sum_{v \in V(C^{1_1})} (k_i - d_{G_1}(v)) = \varepsilon_{i,1,1_1} \text{ such edges. So}$ 

$$|E| = |E_1^{i,1}| + \varepsilon_{i,1,1_1}$$
  
>  $l_i$ ,

by (VI), a contradiction. So each  $H_i$  satisfies (X1).

We show that (X2) is satisfied. First consider  $V \subseteq V(H_i)$  such that  $v_0 \notin V$ . Thus f(V) = |V|. From  $H_i$  form a graph J by merging vertices that belong to the same component of  $H_i - V$  and deleting any loops on these merged vertices. Thus J has f(V) + c(V) vertices and as it is connected,

$$f(V) + c(V) = |V(J)| \le |E(J)| + 1$$
  
  $\le e(V) + 1.$ 

So (X2) is satisfied in this case. Now let  $V = \{v_0\}$ . So  $f(v) = \alpha$ ,  $c(V) = \omega_i$  and  $e(V) = \varepsilon_i + (k_i\alpha - \varepsilon_i)/2$ . Then (X2) can be shown to hold by rearranging the inequality given in (IV). If  $\{v_0\} \subset V$ , then label the other vertices of V so that  $V = \{v_0, v_1, \ldots, v_s\}$ . We have just seen that  $\{v_0\}$  satisfies (X2) so we can show that V satisfies (X2) by proving that if  $V' = \{v_0, v_1, \ldots, v_\sigma\}$ ,  $\sigma < s$ , satisfies (X2), then so does  $V'' = \{v_0, v_1, \ldots, v_{\sigma+1}\}$ . This is done by examining how f, c and e change when the argument V' is replaced by V''. The change in f is clearly +1. Let C be the component of J - V' containing

 $v_{\sigma+1}$ , and let x be the number of components of  $C - v_{\sigma+1}$ . So the change in c is +(x-1). As  $v_{\sigma+1}$  is joined by at least one edge to each of the x components of  $C - v_{\sigma+1}$ , there at least x edges incident with V'' but not with V'. So the change in e is at least +x. So e increases by at least as much as f + c. Thus (X2) remains satisfied.

The only vertex that can have degree  $2k_i$  in  $H_i$  is  $v_0$ . It is a cutvertex if  $G_i$  has more than one component. By (V), if  $l_i = k_i$  and  $l_i$  is odd, then  $\alpha \neq 2$  and so  $d_{H_i}(v_0) \neq 2l_i$ . So (X3) is satisfied.

Finally, (X4) is satisfied since each  $H_i$  contains only one vertex with degree greater than  $k_i$ .

#### References

- L. D. Andersen and A. J. W. Hilton, Generalized latin rectangles, in: R. J. Wilson (Ed.), Research Notes in Mathematics, Vol 34, Pitman, New York, pp.1-17.
- [2] L. D. Andersen and A. J. W. Hilton, Generalized latin rectangles II: Embedding, Discrete Math. 31 (1980) 235-260.
- [3] H. L. Buchanan, Graph factors and Hamiltonian decompositions, Ph.D dissertation, West Virginia University.
- [4] A. J. W. Hilton, Hamiltonian decompositions of complete graphs, J. Combin. Theory Ser. B 36 (1984) 125-134.
- [5] M. Johnson and A.J.W. Hilton, An algorithm for finding factorizations of complete graphs, Journal of Graph Theory 43 (2003) 132-136.
- [6] A. J. W. Hilton, M. Johnson, C. A. Rodger and E. B. Wantland, Amalgamations of connected k-factorizations, submitted.
- [7] A. J. W. Hilton and C. A. Rodger, Hamiltonian decompositions of complete regular s-partite graphs, Discrete Math. **48** (1986) 63-78.

- [8] W. R. Johnstone, Decompositions of complete graphs, Bull. London Math. Soc. **32** (2000) 141-145.
- [9] C. St. J. A. Nash-Williams, Amalgamations of almost regular edge-colourings of simple graphs J. Combin. Theory Ser. B **43** (1987) 322-342.
- [10] C. St. J. A. Nash-Williams, Connected detachments of graphs and generalized Euler trails, J. London Math. Soc. 31 (1985) 17-29.
- [11] C. A. Rodger and E. B. Wantland, Embedding edge-colorings into 2-edge-connected k-factorizations of  $K_{kn+1}$ , J. Graph Theory **10** (1995) 169-185.