

# Past glaciation of temperate-continental mountains: A model for a debris-charged plateau icefield/cirque glacier landsystem in the Southern Carpathians, Romania

Cristina I. Balaban  | David H. Roberts  | David J. A. Evans  | Stewart S. R. Jamieson 

Department of Geography, Durham University, Durham, UK

## Correspondence

Cristina I. Balaban, Department of Geography, Durham University, Durham, DH1 3LE, UK.  
Email: [cristina-ioana.balaban@durham.ac.uk](mailto:cristina-ioana.balaban@durham.ac.uk)

## Funding information

Durham University; Ustinov College, Durham, Grant/Award Number: Norman Richardson Postgraduate Research Fund; Quaternary Research Association, Grant/Award Number: New Research Worker's Award; Royal Geographical Society, Grant/Award Number: Dudley Stamp Memorial Award (PRA 37.20)

## Abstract

Reconstructing the extent, style, timing and drivers of past mountain glaciation is crucial in both understanding past atmospheric circulation and predicting future climate change. Unlike in high-elevation mountains situated in maritime and continental climates, less is known of past glaciation in mid-altitude mountains, located in transitional climates, such as the Southern Carpathians of Romania. Despite these mountains harbouring a rich glacial geomorphology, this has never been systematically mapped according to well-established morphological criteria, nor confidently related to former styles of glaciation. Therefore, filling this gap is important for not only accurately identifying glacial extents but also for establishing past glaciation styles and relating them to past ice dynamics and climate. Here, we devise the first glacial landsystem model for the Southern Carpathians, to enhance our understanding of the glaciation style and dynamics of past mountain glaciers in temperate-continental climates. We present a geomorphological map, from which we infer the spatial and temporal evolution of glacial, periglacial and paraglacial landforms. We then assess the links between the mountain geomorphology and glaciation style and dynamics to construct a glacial landsystem model for debris-charged plateau icefields and cirque glaciers in temperate-continental mountain settings.

## KEYWORDS

cirque glaciers, glacial geomorphology, glacial landsystems, glaciation, plateau icefields, Romania, Southern Carpathians

## 1 | INTRODUCTION


Reconstructing the extent, style, timing and drivers of past mountain glaciation is crucial for both understanding past atmospheric circulation and predicting future climate change (Li et al., 2016). Unlike high-elevation mountains situated in maritime (e.g., Federici et al., 2017) and dry-continental climates (e.g., Xu et al., 2010), less is known of the style, extent and dynamics of past glaciation in mid-altitude mountains located in transitional climates (e.g., temperate-continental/subarctic type), such as the Carpathian Mountains, Romania (Figure 1a,b). This climate zone is characterised by long, cold winters and short, warm summers because of its continental location

and distance from the influence of oceanic air masses (Beck et al., 2018).

Although the Romanian Carpathians (particularly the Southern Carpathians) contain a rich glacial geomorphology (Urdea et al., 2022a, 2022b, 2022c, 2022d, 2022e; Urdea et al., 2022), these mountains have not been systematically mapped according to well-established diagnostic criteria (e.g., Chandler et al., 2018; Giles et al., 2017). Early (19th century) studies on the glacial geomorphology focused on landform descriptions in individual mountain ranges (e.g., de Martonne, 1900; Lehmann, 1881; Tietze, 1878). Later studies (20th century) provided more detailed inventories (e.g., Micalevich-Velcea, 1961; Nedelcu, 1967; Niculescu, 1965; Sîrcu, 1978). The most

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Earth Surface Processes and Landforms* published by John Wiley & Sons Ltd.



**FIGURE 1** Location maps of the study area. (a) The location of Romania within Europe; (b) the location of the Godeanu Mountains within the Southern Carpathians; (c) large scale map of the Godeanu Mountains, showing the highest elevation point (Gugu Peak, 2291 m a.s.l.), plateau surfaces (highlighted in yellow), the names of the main cirques and valleys in the range, as well as the neighbouring mountain ranges: Țarcu Mountains in the West and Retezat Mountains in the East. Cirque and valley toponymy derived or adapted from Niculescu (1965) and Mîndrescu (2016). Digital elevation model for Europe (TOPO-WMS and TOPO-OSM-WMS), sourced from Terrestrial, and that of Romania, Southern Carpathians, and the Godeanu Mountains sourced from EORC, JAXA. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1002/esp.5723)]

recent studies consider an alpine (glacial, periglacial and paraglacial) geomorphological model as a framework for palaeoglaciation and palaeoclimate reconstruction. This has either focused on applying absolute/relative dating methods and/or numerical ice modelling (Balaban, 2018; Gheorghiu, 2012; Gheorghiu et al., 2015; Ignéczi & Nagy, 2016; Kłapyta et al., 2021, 2022; Kłapyta, Mîndrescu, & Zasadni, 2023; Kłapyta, Zasadni, & Mîndrescu, 2023; Kuhlemann, Dobre, et al., 2013; László et al., 2013; Reuther et al., 2007; Ruszkiczay-Rüdiger et al., 2016, 2021; Tirlă et al., 2020; Urdea & Reuther, 2009), or conducting quantitative inventories of landform characteristics (Gunnell et al., 2022; Mîndrescu & Evans, 2014, 2017; Mîndrescu et al., 2010; Necșoiu et al., 2016; Onaca, Ardelean, et al., 2017; Onaca et al., 2013; Popescu, 2018; Popescu et al., 2021; Popescu, Onaca, Urdea & Vespremeanu-Stroe, 2017; Șerban et al., 2019; Vasile et al., 2022; Vespremeanu-Stroe et al., 2012). Despite this work, the relationships between palaeoclimate, geomorphology, and glaciation style and dynamics, particularly in a mountain landsystem context, remain poorly understood. Moreover, a glacial landsystems approach, informed by modern analogues, is necessary for assessing an alpine/cirque versus plateau icefield style of glaciatisation in mountain terrains, to facilitate more accurate palaeoglaciological models (cf. Benn & Lukas, 2006; Benn & Owen, 2002; Bickerdike, Evans, et al., 2018; Bickerdike, Ó Cofaigh, et al., 2018; Carr & Coleman, 2007; Golledge, 2007; Hulton & Sugden, 1997; Rea et al., 1998; Rea et al., 1999; Rea & Evans, 2003, 2007).

In this paper, we compile the first glacial landsystem model for the Romanian Carpathians, in order to understand glaciation style and past glacier dynamics. Based on the Godeanu Mountains in the

southwestern part of the Southern Carpathians, this is achieved through the following objectives: (1) identifying and interpreting the origin and spatial distribution of mountain landforms, based on a remote and ground-truthed geomorphological map; (2) determining the glacial debris transport pathways and genesis of selected glacial depositional landforms, using sedimentological analyses; (3) hypothesising on the relationship between the mountain topography and glaciation style and dynamics, based on the spatial relationships between the landforms (glacial landsystem) in the study area; and (4) assessing the value of the landsystem model as an analogue for similar glacierised uplands.

## 2 | STUDY AREA

The Romanian Carpathians occupy the centre of Romania and comprise three chains, including the Eastern Carpathians, the Southern Carpathians and the Apuseni Mountains (Figure 1a,b). They are composed of sedimentary, metamorphic and volcanic rocks (Urdea et al., 2011) and resulted from Mesozoic–Cenozoic continental collision and Pliocene–Quaternary erosion (15–2.6 Ma), including glaciation during cold climate cycles (Popescu, Urdea, & Vespremeanu-Stroe, 2017; Sanders et al., 2002; Sanders, Andriessen, & Cloetingh, 1999). Their position in Central–Eastern Europe marks a transition between the temperate-maritime climate of Western Europe, the Mediterranean climate of Southern Europe and their continental-arid counterpart of Central Asia (Mîndrescu et al., 2010; Reuther et al., 2007). The mountains are not currently glacierised but have numerous peaks above 2000 m a.s.l. (max: 2544 m a.s.l.—Moldoveanu Peak). In the present climate, the

0°C isotherm is situated at 1650 m a.s.l. in the Eastern Carpathians and 2050 m a.s.l. in the Southern Carpathians, and a mean precipitation above 2000 m a.s.l. of 1400 mm/year in the Eastern Carpathians and 1000 mm/year in the Southern Carpathians (Popescu, Urdea, & Vespremeanu-Stroe, 2017).

The Godeanu Mountains are a ~100 km<sup>2</sup>, northeast–southwest-orientated mountain ridge located in the southwestern part of the Southern Carpathians (45°15'N 22°42'E; Figure 1c). Part of the Retezat–Godeanu group, the mountains form flat/rounded plateau surfaces overlooking ~70 cirques and short (3–4 km) U-shaped valleys with a range of orientations (Mîndrescu, 2016; Niculescu, 1965). The mountains are composed of gneiss, paragneiss, crystalline schist, amphibolite and pegmatite lithologies, with sparse limestone. With a maximum elevation of 2291 m a.s.l. (Gugu Peak; Figure 1c), these mountains are influenced by a sub-Mediterranean climate, with modern prevailing winds from the north, west and southwest (Dragomir et al., 2016; Niculescu, 1965). The climate is warmer (0–2°C average/year) and wetter (1000–1400 mm/year) than other areas of the Southern Carpathians (Dragomir et al., 2016; Niculescu, 1965). Therefore, although the geology and topography of the Godeanu Mountains are similar to other mountains in the eastern Southern Carpathians (e.g., Făgăraş and Iezer), the present climate characteristics are unique to it.

The Godeanu Mountains have previously been the subject of geomorphological and palaeoglaciological research. Early studies identified glacial landforms and produced the first topographic maps (Czirbusz, 1905; de Martonne, 1900; Schafarik, 1899). More recent literature mapped landforms such as intact and relict rock glaciers (Onaca, Ardelean, et al., 2017; Urdea, 1992), cirques (Mîndrescu & Evans, 2014, 2017; Mîndrescu et al., 2010), block streams (Şerban et al., 2019) rock walls (Vasile et al., 2022) and rock slope failures (RSFs; Gunnell et al., 2022). The most comprehensive research has been that of Niculescu (1965), who produced the first field-based geomorphological map of the range. An electro-resistivity tomography profile of a moraine sequence in the Soarbele valley by Ardelean (2010; Figure 1c) confirmed a glacial depositional origin. Niculescu (1965) employed the glacial geomorphology as a morphostratigraphic tool, by relating the Soarbele valley moraine sequence to two ice advances, which he assigned to the Riss (MIS 8–6) and Würm (MIS 4–2) glaciations. The former yielded a palaeo-equilibrium line altitude (ELA) of 1750–1800 m a.s.l., and the latter 2050 m a.s.l. (Niculescu, 1965). However, this approach assumed a cirque/valley glaciation style, with little consideration for plateau icefield sources, despite widespread upland plateau surfaces.

This assumption was challenged by Ignéczi and Nagy (2016), who tested the hypothesis of a plateau icefield style of glaciation by performing numerical modelling of palaeoglacier extents. Using a steady-state plastic numerical model, they reconstructed 40 km<sup>2</sup> of ice over the range, constrained by the geomorphological map of Niculescu (1965). The reconstruction included plateau icefields, with an average palaeo-ELA of 1855 m a.s.l. on the northern slopes and valley glaciers predominant in the south-facing cirques in the west. These yielded an average palaeo-ELA of 1700 m a.s.l. Ignéczi and Nagy (2016) suggested that the discrepancy between the ELAs is related to missing geomorphological evidence of palaeoglacier extents in the northern valleys, possibly caused by postglacial erosion. This demonstrates that re-mapping of the area is necessary before further modelling and dating of past ice extents is attempted to derive paleoglaciological and palaeoclimate reconstructions.

## 3 | METHODS

### 3.1 | Geomorphological mapping

Guided by protocols for remote and field-based mapping of former glaciers (Chandler et al., 2018), we produced a geomorphological map of the Godeanu Mountains, at a scale of 1:120,000 in ArcMap 10.7.1. This was undertaken remotely for the entire range and field-checked in the area of the Scărişoara plateau and four surrounding valleys. Because of low-resolution digital elevation models (max = 30 m, ALOS DEM), mapping was undertaken on high-resolution (1 m) orthorectified aerial photographs from ANCP, ArcGIS World Imagery, as well as on imagery viewed on Google Earth Pro.

The mapping focused on landforms and sediments of glacial depositional origin (moraines and till veneers/drift limits), glacial/glacifluvial erosional origin (ice-moulded bedrock and ice-marginal meltwater channels), periglacial origin (pronival and protalus ramparts, relict rock glaciers, rock-glacierised moraines) and para-glacial landforms (RSFs). Plateau areas were also demarcated to assess the possibility of former plateau icefields. Landforms were initially classified using descriptive terms (e.g., ridges and channels), avoiding premature genetic classifications (Darvill et al., 2017; Evans, Dinnage, & Roberts, 2018). Genetic classifications were subsequently made based on morphological diagnostic criteria, as outlined in Table 1. Field investigations in the area of the Scărişoara Plateau and four surrounding valleys (Bulz, Vlăsia, Scărişoara and Scurtele; Figure 1c) were employed as control sites for extrapolation to the remote mapping for the entire range.

### 3.2 | Sedimentological analyses

Landform origins and potential glacial debris transport pathways were assessed in the field via sedimentological analyses on natural exposures in depositional features in the ground-truthed area. Specifically, this entailed sediment logging and descriptions using a lithofacies approach, augmented by clast shape analyses (cf. Benn & Ballantyne, 1994; Lukas et al., 2013; Evans & Benn, 2021). Clast shape indices include clast roundness, quantified using the RA (percentage of A and VA clasts) and the average roundness (AvgR; cf. Evans, 2010), and clast form, quantified using the C<sub>40</sub> index as an indicator of 'blockiness', and hence subglacial modification (Benn & Lukas, 2021). The degree of clast modification is then related to glacial debris transport pathways by employing the Type II RA-C<sub>40</sub> covariance plot (Lukas et al., 2013) for anisotropic (foliated) lithologies, similar to those of the crystalline schists of the Scărişoara plateau.

### 3.3 | Glacial landsystems reconstruction

Glacial landsystems are assemblages of sediments and landforms that are representative of specific glacial processes, styles and dynamics in the context of the topographic and climatic characteristics of the investigated terrain (Evans, 2003). These characteristics, and thus, the landsystems, may change spatio-temporally in response to climate drivers (e.g., Benn et al., 2003; Bickerdike, Evans, et al., 2018; Bickerdike, Ó Cofaigh, et al., 2018). The importance of glacial landsystems lies in their integration with geochronology and ice

**TABLE 1** Inventory of glacial depositional, glacial erosional, periglacial depositional and paraglacial landforms in the Godeanu Mountains, with details on their associated morphology, sedimentology, significance and key references.

Landforms	Morphology	Sedimentology	Significance	References
<b>Glacial depositional landforms</b>				
Latero-frontal terminal and recessional moraines	Linear, valley-side ridges with asymmetric cross profiles that descend downslope, sub-parallel to contours, and are linked to arcuate ridges crossing valley floors.	Poorly sorted, mostly rubbly diamicton with little to no significant matrix support. Clast forms change down-valley from VA/A and slabby to SA/SR and blocky, with increasing amounts of striation.	Demarcation of lateral and frontal, maximum and recessional glacier margins as well as vertical ice limits.	Benn et al. (2003); Lukas (2005); Glasser et al. (2008); Evans (2011, 2013a)
Drift limits and till veneers	Subdued mantles of debris above which slopes are characterised by bedrock and non-glacial regolith, or valley floor-based deposits.	As above	Demarcation of maximum elevations of lateral glacier margins in steep terrain.	Benn and Evans (2010); Rootes and Clark (2022)
Lateral moraines	Linear, valley-side ridges with asymmetric cross profiles that descend downslope, sub-parallel to contours.	As above	Demarcation of lateral, maximum and recessional glacier margins as well as vertical ice limits. Indicators of large extraglacial debris supply, often resulting in lateral moraine asymmetry.	Matthews and Petch (1982); Small (1983); Benn (1989); Bennett and Boulton (1993); Evans (1999); Quincey and Glasser (2009); Lukas et al. (2012); Boston and Lukas (2019)
Medial moraines	Subdued, valley-parallel, straight or slightly curvilinear ridges/ veneers on valley floors and/or at valley confluences.	Poorly sorted, mostly rubbly/bouldery diamicton with little to no significant matrix support.	Indications of ice flow unit confluences and/or rock avalanche input from backwalls; agents of supraglacial/englacial transport and deposition in valley and cirque glaciers.	Eyles and Rogerson (1978); Rogerson et al. (1986); Benn and Evans (2010); Evans (2013b); Giles et al. (2017); Ballantyne and Dawson (2019)
<b>Glacial erosional landforms</b>				
Ice-moulded bedrock	Striated and sometimes polished bedrock, with symmetric or asymmetric cross profiles (whalebacks and roches moutonnées) as well as abraded rock steps.	N/A	Indications of subglacial erosion and former ice flow directions; evidence of warm-based glacier thermal regime.	Gordon (1981); Evans (1996); Glasser (2002); Glasser and Bennett (2004); Benn and Evans (2010); Rea (2013)
Ice-marginal meltwater channels	Dry (relict) linear or sinuous channels on valley sides, often grouped <i>en echelon</i> and descending in altitude down-valley, crossing contours at oblique angles.	N/A	Large glacier meltwater volumes and high subglacial water pressures; indicate ice limits and recession patterns.	Maag (1969); England (1990); Evans (1990); Dyke (1993); Greenwood et al. (2007); Benn and Evans (2010); Livingstone et al. (2010); Evans, Hughes, et al. (2017); Giles et al. (2017)
<b>Periglacial depositional landforms</b>				
Pronival/ protalus ramparts	Discrete debris accumulations comprising single (protalus) and/or multiple inset (pronival), continuous ridges with asymmetric cross profiles. Maximum distance from talus foot ~30–70 m.	Openwork large clasts with pockets of poorly sorted finer debris but largely devoid of fine matrix.	Debris ridges deposited by perennial snowpatches, formed either outside or inside former glacial limits; not to be confused with moraines.	Ballantyne and Kirkbride (1986); Ballantyne and Benn (1994); Shakesby (1997); Shakesby and Matthews (1993); Shakesby et al. (1999); Matthews et al. (2011); Hedding (2011, 2016); Hedding and Sumner (2013)

TABLE 1 (Continued)

Periglacial depositional landforms				
Relict rock glaciers and rock-glacierised moraines	Discrete debris accumulations comprising spatulate or tongue-shaped assemblage of multiple lobate ridges. Rock glaciers emanate from cirques or plateau edge cliffs. Other rock-glacierised bodies occur within lateral or latero-frontal moraine ridges.	Poorly sorted, coarse, predominantly bouldery openwork debris with no fine-grained matrix. Clast forms predominantly VA/A in relict rock glaciers but becoming more A-SR in rock glacierised moraines.	Melt-out of debris-covered glaciers or creep of coarse debris bodies with interstitial ground ice (permafrost creep). Indicative of mountain setting with high debris turnover. Relict rock glaciers formed either inside or outside glacial limits. Rock-glacierised moraines formed within glacial limits.	Thompson (1957); Østrem (1964); England (1978); Wilson (1990); Ballantyne and Harris (1994); Hamilton and Whalley (1995); Harrison et al. (2008); Whalley (2009, 2012, 2015); Anderson et al. (2018); Jones et al. (2019); Leigh et al. (2021); Urdea et al. (2022)
Paraglacial landforms				
Rock slope failures	Discrete debris accumulations or partially intact bedrock masses comprising large debris mounds, often with no preferred orientation, but situated at the foot of failure-source rock walls. Arrested forms appear as slope deformations and/or <i>en echelon</i> anticarps at the margins of plateaux and cirque backwalls.	Debris mounds are predominantly poorly sorted, matrix-poor, coarse and boulder-rich, with A/VA clast forms.	Deposits related to, but not exclusively diagnostic of, slope re-adjustment following deglaciation; not to be confused with moraines or periglacial landforms, which they may overlie.	Shakesby and Matthews (1996); Clark and Wilson (2004); Ballantyne (2002, 2013, 2018a); Jarman (2006); Coleman and Carr (2008); Jarman et al. (2013); Ballantyne et al. (2014); Gunnell et al. (2022)

modelling to produce a holistic understanding of glaciation style, dynamics and palaeoclimate through time. However, their compilation in ancient settings requires the recognition of geomorphological process-form regimes, informed by suitable modern analogues. Hence, we consulted modern analogue landsystems from similar climatic and topographic settings, such as those for plateaux (Rea et al., 1998; Rea et al., 1999; Rea & Evans, 2003, 2007) and glaciated valley (Benn et al., 2003; Benn & Owen, 2002) settings where paraglacial and periglacial process-form regimes interact with those of glacial systems (Knight et al., 2019; Leigh et al., 2021).

## 4 | LANDFORMS


### 4.1 | Moraines, drift limits and till veneers

#### 4.1.1 | Morphology and genesis

The moraines mapped in the Godeanu Mountains are of several morphological and genetic types (Figures 1c and 2 and Table 1). The continuous linear valley-side depositional ridges and associated valley-floor arcuate ridges, forming inset sequences in an up-valley direction, are clearly latero-frontal moraines (Figures 3a and 4a). Occasionally, these are replaced by subdued, lobate-shaped glacial debris/till veneers (Figure 3b). Both landform types demarcate recessional, lobate glacier margins (Evans, 2013a). Terminal moraines and till veneers are more subdued than their recessional counterparts, which have sharper crests (Figure 3b and Table 1). Linear depositional ridges, perched on valley sides but not continuous with arcuate valley

floor ridges, are regarded as lateral moraines, which underpin lateral glacier extents (Lukas et al., 2012; Figure 3c). Subdued, linear or sinuous bouldery deposits or veneers, which are situated on valley floors, usually at ice flow confluences, are interpreted as medial moraines, signifying supraglacial/englacial transport of debris in the glacial system (Eyles & Rogerson, 1978; Figures 3d and 4b).

Sedimentological investigations were undertaken on exposures through moraines in the Bulz, Vlășia, Scărișoara and Scurtele valleys, surrounding the Scărișoara plateau (Figure 5a), including the lateral moraines BLZ1, BLZ2, BLZ4 and SCT1, and the frontal/recessional moraines BLZ3, BLZ5, VLS1, VLS2 and SCS1. The moraines are composed of coarse, bouldery, sandy-silty matrix or clast-supported diamictons (Figure 6a,b). The RA-C<sub>40</sub> covariance plot (Figure 6c) reveals the relatively high C<sub>40</sub> values, typical of anisotropic lithologies, indicating that roundness indices such as RA and AvgR are more appropriate in diagnosing clast modification within the former glaciers of the area. Down-valley trends can be assessed only in one location, the Bulz Valley, where the most intensive sampling was undertaken. Here, clast modification increases as expected in a down-valley direction (cf. Benn, 1989; Benn & Ballantyne, 1994; Evans, 1999; Matthews & Petch, 1982), with AvgR values increasing, and C<sub>40</sub> and RA values decreasing, based upon valley head samples BLZ3 and BLZ2, valley side samples BLZ5 and BLZ4 and middle valley floor sample BLZ1 (Figure 5b-f). Overall, the moraine deposits and their sedimentological characteristics (Figures 5 and 6) indicate mountain glacier systems with limited active subglacial transport and predominantly passive, supraglacial debris transfer (Benn et al., 2003; Boulton, 1978; Evans, Roberts, & Evans, 2016; Evans, Roberts, et al., 2018).




**FIGURE 2** The mountain geomorphology of the Godeanu Mountains, Southern Carpathians, Romania, focusing on the main glacial, periglacial, paraglacial and contextual landforms. The numbered dashed line rectangles reveal the location of landform examples featured in Figure 3. Produced in ArcGIS, using two hillshade models (transparency = 95%, illumination angles = 180° and 315°), derived from the ALOS DEM (EORC, JAXA, resolution = 30 m). High-resolution version (without the numbered rectangles) is available to consult in the [Supporting Information](#) section in the online version of the article. [Color figure can be viewed at [wileyonlinelibrary.com](#)]

#### 4.1.2 | Spatial distribution

Latero-frontal moraines occur in all glacial valleys and on the Borăscu plateau (Figures 1c and 2). Where they are scarce, particularly in north/northeast-facing valleys, such as Scurtele, Scărișoara, Borăscu Mic, Galbena North and Galbena South, former glacier extents are demarcated by drift limits/till veneers. Also diagnostic of a latero-frontal moraine genesis is the reduction in ridge lengths and widths between lower valley reaches (e.g.,  $L = 617$  m;  $W = 326$  m) and upper valley heads (e.g.,  $L = 100$  m;  $W = 98$  m) (Figure 3a), which reflects the reduction in former glacier size in an up-valley direction. In valleys with multiple cirques (e.g., Galbena North valley; Figures 1c and 2), recessional moraines can be traced back onto each cirque floor. With respect to valley aspects, latero-frontal and recessional moraines are smaller, more closely spaced (average = 189.5 m) and greater in number (average = 9.05 moraines/valley) in south-facing valleys than those in north-facing ones, which are larger, more widely spaced (average = 303.75 m) and lower in number (average = 7.16 moraines/valley). Additionally,

the latero-frontal moraines reach lower elevations in south-facing valleys (average = 1477 m) than in the north-facing counterparts (average = 1648 m). The position and change in size of latero-frontal moraines from down-valley settings to valley heads reflect changes in palaeoglacier extents through time (e.g., Benn et al., 2003; Evans et al., 2010; Owen et al., 2001; Table 1). The predominance of subdued drift limits in north and northeast-facing valleys reflects lower glacier debris turnover, as well as relatively greater postglacial erosion and sediment reworking, promoted by the steeper, stepped bedrock valley profiles (Ignécz & Nagy, 2016; Niculescu, 1965). The increased number of moraine ridges in proximity to cirques in valley complexes shows the separation of different ice sources and concomitant relative stabilization of the glaciers once they retreated to higher elevations (Glasser et al., 2006). Notwithstanding the differences in north- versus south-facing valley geometries, the difference in number, spacing and size of the latero-frontal and recessional moraines is possibly due to north-facing palaeoglaciers being larger (Ignécz & Nagy, 2016) and less responsive to climate fluctuations. Aspects of valley topography that are likely influential in this respect




**FIGURE 3** Orthophotograph–map pairs of mountain landforms in the Godeanu Mountains. Black arrows indicate examples of the relevant landform observed. The maps use the location names in Figure 1c, and the legend in Figure 2, and include specific examples for some of the statistics discussed in the text: (a) latero-frontal and recessional moraines in the Soarbele valley; (b) till veneers in the Scurtele valley; (c) lateral moraines in the Cărbunele valley; (d) medial moraines in the Bulz valley; (e) ice-moulded bedrock in the Bontica valley; (f) ice-marginal meltwater channels in the Vlășia valley; (g) pronival ramparts in the Borăscu Mic valley; (h) rock glacier in the Scărișoara valley; (i) rock-glacierised moraine in the Vlășia valley; (j) rock slope failure (rockslide) in the Borăscu Mare valley. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

are the bedrock steps, which can act as pinning points, solar insolation, and lee-side aspects relative to southerly winds (e.g., Barr & Lovell, 2014; Benn & Ballantyne, 2005; Coleman et al., 2009; Mitchell, 1996; Oerlemans et al., 1998).

Lateral moraines are uncommon, with more extensive examples occurring in the Olanu West, Vlășia, Cărbunele, Măneasa, Scurtele, Mățu and Branu North valleys. More fragmented examples occur in the Scurtele, Mățu and Scărița-Bulz valleys (Figures 1c, 2 and 3c). In the Cărbunele (Figure 3c) and Soarbele (Figure 3a) valleys, there is a preferential slope aspect for the occurrence of large lateral moraines. For example, in the Cărbunele valley, a prominent lateral moraine occurs on the northwest-facing slope, where RSFs are also common,

whereas there are no lateral moraines on the debris-poor southeast-facing equivalent. The variable occurrences and sizes of lateral moraines reflect the availability of paraglacial/supraglacial debris, resulting in lateral moraine asymmetry (Benn, 1989; Evans, 1999; Matthews & Petch, 1982; Table 1). This refers to the preferentially larger sizes of lateral moraines situated on steep, freeface-dominated and failure-prone valley sides (cf. Jarman, 2009), particularly in association with RSFs and talus cones, as observed in the Cărbunele and Soarbele valleys.


Despite their generally low preservation potential in glaciated terrains (Benn & Evans, 2010), medial moraines are present in the four south-facing valleys of Mocirlu, Bulz, Cărbunele and Soarbele and in



**FIGURE 4** Field photograph example of mountain landforms in the Godeanu Mountains. The white arrows indicate examples in each landscape context: (a) recessional moraine in the Bulz valley; (b) ‘ice-stream interaction’-type medial moraines in the Bulz valley; (c) ice-moulded bedrock in the Scărișoara valley; (d) ice-marginal meltwater channel in the Vlășia valley; (e) pronival ramparts in the Scurtele valley; (f) relict rock glacier in the Scărișoara valley; (g) rock-glacierised moraine in the Vlășia valley; (h) ice-moulded rock slope failure blocks in the Scurtele valley. [Color figure can be viewed at [wileyonlinelibrary.com](http://wileyonlinelibrary.com)]

the northeast-facing Galbena North valley (Figures 1c, 2, 3d and 4b and Table 1). These features occupy a specific elevation niche at 1630–1950 m a.s.l., where they grade down-valley into recessional latero-frontal moraines (Figure 3d). Medial moraine production appears to have taken place via two distinct modes, based upon the models of Eyles and Rogerson (1978) and Rogerson et al. (1986). Firstly, ‘ice stream interaction’ (ISI)-type medial moraines are formed through the coalescence of lateral moraines of two tributary glaciers

or ice flow units (Eyles & Rogerson, 1978). This model seems appropriate for the confluence between the two cirques comprising the Bulz valley, demarcated by three transverse ridges (Figures 2, 3d and 4b). Secondly, ‘avalanche-type’ medial moraines do not require the existence of coalescing ice flow units, but rather a supraglacial rockfall debris source, such as from a plateau or high cirque backwall nunatak (Rogerson et al., 1986). This model is most applicable to the eastern Bulz, Mocirlu, Cărbunele and Soarbele valleys (Figures 1c, 2 and 3a,d).



**FIGURE 5** Clast sedimentological data on moraines in the valleys surrounding the Scărișoara plateau: (a) Bulz, Scurtele, Scărișoara and Vlășia. Each sample is named after the valley of provenance and order of collection: (b) BLZ3, (c) BLZ2, (d) BLZ 1, (e) BLZ5, (f) BLZ4, (g) SCT1, (h) SCS1, (i) VLS1 and (j) VLS2. Various sedimentological statistics (clast form ternary diagrams, clast roundness,  $C_{40}$  index, relative angularity (RA), and average roundness (AvgR) are shown in individual boxes. The map legend can be consulted in the bottom right corner. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com)]

In both cases, with ice retreat and thinning, the debris position of each flow unit narrows and shifts towards the valley sides, forming inset ridges that represent parallel medial moraines in the main valley. The highly lobate bouldery ridge in the Bulz valley (Figure 3d) is more complicated in that it resembles the teardrop-shaped looped medial moraines of surging mountain glaciers (e.g., Ellis & Calkin, 1979; Gusmeroli et al., 2014; Meier & Post, 1969; Paul, 2015). Consequently, this reflects the differential speeds of individual flow units within the former valley glaciers of the study area rather than surging, as other features indicative of surging are lacking: for example, crevasse squeeze ridges, thrust moraines and zigzag eskers (Evans & Rea, 2003). We attribute the high preservation potential of medial moraines to large volumes of debris sourced supra-, sub-, and englacially (e.g., Eyles & Rogerson, 1978; Goodsell, Hambrey, & Glasser, 2005; Goodsell, Hambrey, Glasser, Nienow, et al., 2005; Small, Clark, & Cawse, 1979).

## 4.2 | Ice-moulded bedrock

### 4.2.1 | Morphology and genesis

Ice-moulded bedrock is represented by abraded and plucked outcrops that form roches moutonnées and whalebacks/rock drumlins. These features are indicators of subglacial erosion and former ice flow directions, and are generally reflective of a warm-based glacial thermal regime (Glasser, 2002; Glasser & Bennett, 2004; Rea, 2013; Table 1).


### 4.2.2 | Spatial distribution

In the Godeanu Mountains, ice-moulded bedrock is commonly found at altitudes of 1860–1920 m a.s.l. In areas of paragneiss and amphibolite in particular, roches moutonnées are widespread. Ice-moulded bedrock is common and prominent in the north-facing Bonțica, Gropița, Mătu, Scurtele and Scărișoara, as well as the southern-facing Balmoș East, Scărița, Micușă, Stâna, Cărbunele and Soarbele valleys (Figures 1c, 2, 3e and 4c). Niculescu (1965) suggested that ice-moulded bedrock outcrops in areas where more durable lithologies, such as paragneiss, amphibolite and pegmatite, resist glacial erosion better than the weaker lithologies, such as crystalline schist. The prevalence of ice-moulded bedrock forms in north-facing valleys has been explained by Niculescu (1965) as a result of valley orientation, and hence ice flow being oriented in the opposite direction to the dip of the underlying strata.

## 4.3 | Ice-marginal meltwater channels

### 4.3.1 | Morphology and genesis

Relict channels indicative of former glacial meltwater incision appear as elongate depressions or interlinked channel networks, interpreted as ice-marginal meltwater channels. They are interspersed with bedrock and sometimes sediment ridges and are present in locations where their genesis is incompatible with a fluvial origin (Dyke, 1993). The ice-marginal meltwater channels are useful in identifying former



**FIGURE 6** Key information for interpreting the glacial transport pathways in the study area. (a) Stream-cut exposure in moraine in the Bulz valley, where clast data for sample BLZ2 were collected. Note the poorly sorted, matrix-supported, predominantly A and VA clasts. (b) Close-up photograph into a stream-cut exposure in a till veneer in the Scărișoara valley, where the SCS1 sample was taken from. Notice the poorly sorted, clast-supported, sandy matrix, with predominantly SA and SR clasts. Trowel for scale. (c) Covariance plot, showing how the clast data samples in the study area plot in terms of transport pathways. Each sample can be correlated with the valley of provenance by a different symbology. The colour gradient indicates the samples' relative position in their associated valleys, from the lighter upper valley samples to the darker lower valley ones. [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/terms-and-conditions)]

ice limits, and they offer insight into glacial thermal regime, meltwater volumes and subglacial water pressures (Table 1). Previous research has used meltwater channels as a diagnostic criterion for cold-based ice margins, whereby glacial meltwater is diverted along frozen margins, carving channels corresponding to their retreat positions (Dyke, 1993; Evans, Hughes, et al., 2017; Greenwood et al., 2007; Livingstone et al., 2010; Maag, 1969). However, Dyke (1993) and Syverson and Mickelson (2009) also acknowledged meltwater channel initiation in warm-based thermal settings.

#### 4.3.2 | Spatial distribution

Ice-marginal meltwater channels are rare and specific only to the south-facing Mocirlu, Soarbele, Bulz and Vlășia valleys (Figures 1c, 2, 3f and 4d). Their planforms can be employed to demarcate ice limits and glacier recession patterns, especially where they are nested and/or associated with recessional moraine ridges (Table 1). Located in the elevation range 1530–1890 m a.s.l., the meltwater channels range from 60 to 330 m long, with one exception in the Soarbele valley, of 823 m. Ground-truthed examples in the Bulz and Vlășia valleys

are 2–5 m deep. We cannot exclude a cold-based thermal regime in the Godeanu Mountains under specific conditions (elevation range and slope location). However, the association of meltwater channels with other landforms indicative of active deposition (recessional moraines), as well as warm-based thermal regimes (ice-moulded bedrock), indicates that an active temperate or polythermal setting was operating for at least some of the time.

### 4.4 | Pronival and protalus ramparts

#### 4.4.1 | Morphology and genesis

Pronival and protalus ramparts are single, largely continuous ridges with asymmetric cross profiles, formed by the accumulation of rockfall debris at the foot of perennial snowpatches (Shakesby, 1997; Shakesby et al., 1999; Matthews et al., 2011; Table 1). Although some debate has surrounded a snowpatch versus glacial (morainic) genesis for many mountainside ridges (e.g., Shakesby, 1997; Shakesby & Matthews, 1993), there are specific diagnostic criteria for palaeo-protalus rampart interpretations (Table 1). Typically, the distance

between a protalus rampart and the talus foot that it formerly covered can only be 30–70 m, otherwise the snowpatch would have turned into glacier ice (Ballantyne & Benn, 1994). Pronival ramparts, which represent ridges accumulating on mid-slopes, would thus be related to even smaller snow accumulations. The sedimentology of a pronival rampart should also typically be predominantly composed of open-work clasts in places, including a fine matrix (Table 1).

#### 4.4.2 | Spatial distribution

We mapped 48 pronival and protalus ramparts in the Godeanu Mountains (Figures 2, 3g and 4e). Pronival and protalus ramparts are dominant in north-facing valleys ( $n = 31$ ) in contrast to south-facing valleys ( $n = 17$ ). They are situated at lower elevations in the south-facing valleys (1620–2100 m a.s.l.; average = 1790 m a.s.l.), compared with north-facing ones (1710–2100 m a.s.l.; average = 1945 m a.s.l.). They also lie upslope of moraine ridges in many locations (e.g., up-valley distance between moraine and pronival rampart of 14.5 m in the Olanu West, and 15.03 m in the Cârnea valleys, respectively) (Figures 1c and 2). The spatial distribution of pronival and protalus ramparts offers potential insight into palaeoclimatic trends (Ballantyne & Kirkbride, 1986). Firstly, favourable conditions, such as shading and windward-facing basins, are reflected in preferential locations on north-facing slopes (Ballantyne & Kirkbride, 1986; Colucci et al., 2016). The lower elevations at which the ramparts are located in south-facing valleys might suggest these formed at a different time and climatic conditions than their north-facing equivalents. Secondly, the close association of pronival and protalus ramparts with recessional moraines reflects a spatio-temporal transition from small glaciers to snowpatches (Ballantyne & Benn, 1994; Niculescu, 1965). Indeed, Hedding et al. (2007) noted pronival and protalus ramparts forming under decreasing precipitation at the inception/demise of regional glacier activity.

### 4.5 | Relict rock glaciers and rock-glacierised moraines

#### 4.5.1 | Morphology and genesis

Given the relict nature of the geomorphology and the absence of active permafrost in the Godeanu Mountains (Onaca, Urdea, et al., 2017), we refrain from subdividing rock glaciers by their hypothetical origin (rock glaciers or periglacial protalus lobes) (Harrison et al., 2008). Here, we employ the descriptive use of the term ‘rock glacier’, or a tongue-shape assemblage of multiple lobate ridges, composed of coarse, predominantly bouldery debris, regardless of their origin (e.g., permafrost/interstitial ice vs. glacier-derived genesis; cf. Ballantyne, 2018b; Berthling, 2011; Harrison et al., 2008; Humlum, 1996 Table 1). Hence, we endorse the concept of equifinality of rock glaciers (Knight et al., 2019; Whalley, 2009). Additionally, we follow the conventions of Thompson (1957), Østrem (1964), England (1978), Leigh et al. (2021) and Urdea et al. (2022) in recognising rock-glacierised moraines in situations where moraines appear to have been remobilized by creep, either due to interstitial pore ice accumulation or an initial glacier ice core (Table 1). In each case, rock glaciers and rock-glacierised moraines in the study area are relict features and, hence, are genetically classified based upon their more muted appearance after de-icing. This makes them

difficult to distinguish from some other landforms, especially RSF run-out deposits and closely spaced, bouldery latero-frontal moraines (e.g., Ballantyne, Schnabel, & Xu, 2009; Whalley, 2009, 2012).

#### 4.5.2 | Spatial distribution

We identified relict rock glaciers on valley sides, in association with talus cones (Figures 1c, 2, 3h and 4f), and rock-glacierised moraines (Figures 1c, 2, 3i and 4g) on the valley floors. We estimate that around 12 relict rock glaciers exist, with six located in the north-facing valleys of Scurtele, Scărișoara, Stâna Borăscu, Galbena North and Galbena South, and another six in the south-facing Balmoș East, Godeanu, Mocirlu, Scărița, Bulz and Vlășia valleys. In contrast, rock-glacierised moraines are mainly found in south-facing valleys (six examples), with only two examples located in the north-facing Mățu valley. Taken together, and similar to the distribution of pronival and protalus ramparts, the elevation ranges at which rock glaciers and rock-glacierised moraines are found are higher in the north-facing valleys (1690–2016 m a.s.l.; average = 1841 m a.s.l.) than in their south-facing equivalents (1650–1940 m a.s.l.; average = 1786 m a.s.l.). Previous research on relict rock glaciers in the Southern Carpathians (Onaca, Ardelean, et al., 2017; Popescu, Urdea, & Vespremeanu-Stroe, 2017; Vespremeanu-Stroe et al., 2012) hypothesised that they formed as a response to either Lateglacial climatic warming during the Bølling–Allerød Interstadial (14.6–12.9 ka), precipitation starvation during the Younger Dryas (12.9–11.7 ka) or high humidity during the Early Holocene (11.7–8.2 ka), when glacier retreat was underway or complete, and slope destabilisation, permafrost development and rockfall occurred (Rasmussen et al., 2014; Walker et al., 2012). In terms of rock glacier distribution, Popescu, Onaca, Urdea, and Vespremeanu-Stroe (2017) and Șerban et al. (2019) noted a discrepancy in rock glacier and block stream elevation between north- and south-facing slopes in the Southern Carpathians, with the northern slope features located at higher elevations than their southern counterparts. Popescu, Onaca, Urdea, and Vespremeanu-Stroe (2017) and Șerban et al. (2019) attributed this to south-facing valleys receiving higher solar insolation, and thereby earlier glacier retreat and slope debutting than in north-facing valleys. Because of their shading, the north-facing valleys might have preferentially supported higher elevation rock glaciers more recently. Although geochronology might constrain landform age and associated climate change, the incrementally polygenetic origin (glacial-to-periglacial) of many features makes their employment as palaeoclimatic indicators complicated (cf. Harrison et al., 2008; Jarman et al., 2013; Knight et al., 2019).

### 4.6 | Rock slope failures

#### 4.6.1 | Morphology and genesis

Depending on the failure style, travel distance and degree of disintegration, RSFs are represented by a range of discrete debris accumulations or partially intact bedrock masses, situated on, or at the foot of, steep rock walls. They can be classified as follows: (1) rock avalanches—discrete debris assemblages on valley floors, whose transport path can be traced directly uphill; (2) rockslides—arrested rock masses showing little deformation due to their short-travel distance; (3) rock slope deformations—subtle ruptures, bulges, tension fault ridges and troughs

(mini horsts and grabens) or anticarps on valley sides and plateau edges (Gunnell et al., 2022). The lithology of RSFs can be traced to a source depression immediately upslope (Table 1; Ballantyne, 2013, 2021; Jarman, 2006). Such landforms are thought to relate to slope instabilities and isostatic adjustment following the removal of ice weight upon deglaciation (Table 1; Ballantyne, 2002; Ballantyne et al., 2014). Because of their similarity to, and juxtaposition with glacial landforms, RSFs can be misidentified as former glacier ice limits; they can also obscure them in cases of postglacial failures (Table 1). Additionally, RSFs that pre-date ice advance can be overrun and streamlined by glacier flow (Ballantyne, 2018a; Figure 4h).

#### 4.6.2 | Spatial distribution

We identified 97 RSFs in the Godeanu Mountains. These landforms are predominantly distributed in north-facing valleys (65) rather than in south-facing ones (32; Figures 2, 3j and 4h). The RSFs cluster around valley heads and occur at a wide range of elevations (1650–2190 m). Compared with other features, RSFs are large (0.206–0.0005 km<sup>2</sup>; average = 0.038 km<sup>2</sup>) and classify as three rock avalanches, 39 rockslides and 55 rock slope deformations. Occasionally, RSFs appear to be associated with relict rock glaciers and rock-glacierised moraines, both in north- (e.g., Borăscu Mare) and south- (e.g., Vlăşia) facing valleys. However, a number of RSF-rich north-facing valleys are devoid of rock glaciers and rock-glacierised moraines, including the Izvoru Moraru, Branu North, Măţu, Scurtele, Zănoaga, Galbena North and Paltina valleys (Figures 1c and 2). Our detailed mapping approach differs from that of Gunnell et al. (2022), which recognised only 21 RSFs in the Godeanu Mountains amongst 215 in the Southern Carpathians.

The work of Gunnell et al. (2022) and that of Vasile et al. (2022) constitute the first inventory and application of <sup>10</sup>Be cosmogenic nuclide dating of RSFs in the Southern Carpathians, respectively. These pioneering studies suggest that Neogene tectonic uplift and climate warming in the Southern Carpathians drove RSF development and that periglacial, fluvial and parafluvial triggers were more important than glacial and paraglacial processes. The dominance of Early Holocene (pre-9 ka) dated boulders in RSF deposits in particular supported these proposals (Vasile et al., 2022). Indeed, the spatio-temporal relationships between glacial, periglacial and paraglacial landforms may indicate a deglacial/postglacial (Early Holocene) development period for RSFs. Furthermore, the higher number of RSFs on the north-facing slopes and lower association with periglacial landforms on this aspect than on the south-facing ones re-inforces the hypothesis of a later deglaciation on the north-facing valleys, which may still be undergoing failure, whereas their southern counterparts have already completed the paraglacial cycle (Ballantyne, 2002). Over the longer timescales of multiple Quaternary glaciations, however, RSF operation likely initiates and/or contributes to cirque development through headwall erosion into plateau surfaces (cf. Evans, 2021; Turnbull & Davies, 2006), sourcing debris for the development of rock glaciers and moraines.


## 5 | GLACIAL LANDSYSTEMS

The spatio-temporal relationships between the mapped landforms and the sedimentological evidence provide a clear signature of past glaciation style, debris turnover and moraine construction in the

Godeanu Mountains, thereby facilitating the compilation of a glacial landsystem model for the region of the Southern Carpathians. This landsystem model displays many of the characteristics of the conceptual plateau icefield (Rea & Evans, 2003), glaciated valley (Benn et al., 2003) and the cirque/niche (Bickerdike, Ó Cofaigh, et al., 2018) glacial landsystems, as well as elements of debris-charged geomorphology observed in a modern analogue in Norway (Leigh et al., 2021). Therefore, it is best classified as a debris-charged plateau icefield/cirque glacier landsystem. The most extensive phases of glaciation were characterised by simultaneous occupation of valleys by ice descending both from surrounding plateaux and from cirques, where narrow mountain ridges dominate. Conversely, phases of less extensive glaciation were characterised by localised cirque and niche glaciers, detached snouts/reconstituted glaciers and perennial snowpatches (e.g., Benn & Lehmkuhl, 2000; Evans, Ewertowski, & Orton, 2016; Evans et al., 2006; Gellatly et al., 1986; Rea & Evans, 2003), whereas plateau ice may have been restricted to upland surfaces only or completely disappeared. The latter phases may have occurred initially on the more exposed south-facing slopes, with the more shaded north-facing counterparts possibly deglaciating later. Deglaciation unfolded in a landscape prone to rock glacier development, driven by rapid paraglacial debris accumulation and debris-covered glacier snouts, possibly compatible with the oscillating Lateglacial climate in the region (Magyari et al., 2012; Onaca, Ardelean, et al., 2017; Popescu, Urdea, & Vespremeanu-Stroe, 2017; Vespremeanu-Stroe et al., 2012). The spatial and temporal evolution of this complex mountain-based landsystem is depicted in a four-phase schematic model for the Godeanu Mountains in Figure 7.

### 5.1 | Spatio-temporal change in the Godeanu Mountains debris-charged plateau icefield/cirque landsystem

Using the most extensive moraine limits mapped, the first phase (Figure 7a) depicts the plateau icefield at its maximum extent, with ice descending into two neighbouring valleys. The valleys are separated by an ice-free nunatak, their bedrock slopes feeding extraglacial debris via talus cones to lateral and medial moraines of the outlet glaciers. Plateau icefields were developed on the Moraru, Branu, Scărişoara, Borăscu, Paltina and Stănuleţi summits, where the lack of glacial landforms (except rare moraine ridges documenting ice recession from surrounding valley heads) indicates cold-based ice dispersal centres. Cirque glaciers accumulating in the more alpine terrain flow into the main valleys, coalescing with the plateau outlet glaciers. This results in the deposition of further medial moraines; the medial moraines on the cirque glaciers, created predominantly as avalanche types fed by couloirs at headwalls (Eyles & Rogerson, 1978; Rogerson et al., 1986), are also added to the valley glacier surface downflow of the coalescence zone. The transition from predominantly cold-based ice on the plateaux to warmer ice in the outlet glaciers was likely driven initially by the strain heating at the base of icefalls on the plateau edges (Rea et al., 1999; Rea & Evans, 2003; Whalley et al., 1996). Together with the subglacial modification of clasts down-valley, this explains the contrast between plateau summits devoid of glacial landforms and the widespread occurrence of glacial erosional landforms at lower elevations. Over time, the enlargement of cirque headwalls, as well as trough head development at plateau edges, promotes not only further



**FIGURE 7** Conceptual model for the debris-charged plateau icefield/cirque glacier landsystem based on the Godeanu Mountains. The model is based on amalgamated elements from different valleys. Four phases (a–d) are represented to show how the landsystem signature changes spatio-temporally during deglaciation. Key features and landforms are as follows: (1a) plateau icefield dispersal centre; (1b) plateau icefield; (2) cirque glacier; (3) medial moraines; (4) nunatak; (5) snowpatches with protalus lobes (a) and protalus/pronival ramparts (b); (6) rock slope failure; (7a) latero-frontal end moraine; (7b) recessional moraine loops; (8) proglacial streams, ice-marginal meltwater channels and outwash deposits; (9) debris covered glacier snout developing into a rock glacier; (10) protalus rock glacier; (11) lateral moraines; (12) rock-glacierised lateral moraines; (13) reconstituted or stagnant glacier; (14) blockfield; (15) glacially scoured bedrock outcrops (roches moutonnées and whalebacks). [Color figure can be viewed at [wileyonlinelibrary.com](https://onlinelibrary.wiley.com/doi/10.1002/esp.5723)]

glacial erosion but also slope destabilisation, and the initial triggering of RSFs, sourcing large volumes of extraglacial debris to the glacial system. In turn, this creates increasingly more debris-charged mountain glaciers (Benn et al., 2003; Bennett et al., 2010; Evans, Ewertowski, & Orton, 2016, 2017; Rea & Evans, 2003). Based upon the landsystem extent and the general elevation of the outermost mapped moraines ( $\leq 1500$  m a.s.l.), we suggest that this phase may be coeval with early post-Last Glacial Maximum deglaciation, especially as two moraine systems in the neighbouring Retezat Mountains at lower elevations (M1 at  $\sim 1000$ – $1100$  m a.s.l.; M2a at  $1200$ – $1400$  m a.s.l.) have been dated  $21.0^{+0.8/-1.5}$ – $20.6^{+0.8/-1.3}$  ka and  $18.6^{+0.9/-0.8}$ – $18.4^{+0.7/-1.1}$  ka, respectively, by Ruszkiczay-Rüdiger et al. (2016, 2021).

The second phase (Figure 7b) depicts initial ice retreat from the maximum extent. During this phase, the lowermost set of latero-frontal moraines were abandoned, and medial moraines were lowered onto the subglacial footprint. Also prominent are the looped latero-frontal and medial moraine signatures of constricted cirque glacier ice, flowing along the valley edge. Ice-marginal meltwater channels were excavated along lower valley slopes and the valley floor. Glacier snout thinning during retreat promoted further slope destabilisation and development of RSFs. The occurrence of continuous latero-frontal moraine ridges and ice-marginal meltwater channels indicates a (partially) coupled ice margin, with high ice and debris supplies, and the efficient meltwater transfer of glacial debris from the snout to proglacial outwash systems (Benn et al., 2003). Although ice-marginal meltwater channels typify cold-based plateau icefields, their lower valley position, the association with depressions and occasional intercalation with moraine ridges (Figure 3f) suggests a

warm-based or polythermal basal thermal regime (Dyke, 1993). This is supported by  $^{10}\text{Be}$  cosmogenic dating-inferred erosion rates in the Retezat Mountains, which indicate warm-based conditions at the ice margins and colder ice at the valley head during initial deglaciation (Ruszkiczay-Rüdiger et al., 2021). Additionally, the reconstructed coupled margin at this stage is a defining characteristic for glaciated valley landsystem settings with high debris turnover (e.g., New Zealand), whereby debris is efficiently evacuated from the margin by high subglacial meltwater pressures and volumes (Benn et al., 2003; Dyke, 1993). Phase 2 moraines are located at  $\sim 1500 \geq 1750$  m a.s.l. and may correspond to the M2b moraine system ( $1600$ – $1750$  m a.s.l.) in the Retezat Mountains, which yielded early Lateglacial ages of  $18.6^{+0.9/-0.8}$ – $16.9^{\pm 0.9}$  ka (Ruszkiczay-Rüdiger et al., 2016, 2021).

The third phase (Figure 7c) was characterised by continued recession of valley and cirque-based glaciers until the different glacier bodies became disconnected. Inset sequences of cirque moraines then started to record recession into localised alpine uplands. Overall recession was under active, warm-based conditions, as indicated by widespread ice-moulded bedrock and multiple inset latero-frontal moraines. Large longitudinal concentrations of supraglacial debris are indicated by the development of significant medial moraines, linked to tight arcuate or looped frontal moraines, each loop demarcating a separate flow unit within the valley outlet glacier (Eyles & Rogerson, 1978; Jennings et al., 2014; Roberson, 2008). As medial moraine preservation potential is usually very low after complete deicing (Benn & Evans, 2010), the prominence of such features in the Godeanu Mountains (especially in increasing numbers up-valley)

indicates significant debris provision via extraglacial pathways. This was presumably derived from RSF sources, which began dominating large, steep areas of uncovered trough heads surrounding the plateaux upon ice recession (Ballantyne & Dawson, 2019; Evans, Ewertowski, & Orton, 2016, 2017; Rea & Evans, 2003). Examples of lateral moraine asymmetry (e.g., Figure 3c) reflect the uneven distribution of freefaces delivering such material during deglaciation. Over time, the uncovering of increasingly larger areas of such steep bedrock outcrops during glacier recession and downwasting resulted in enhanced debris production that locally overwhelmed glacier surfaces. This resulted in the formation of rock glaciers and rock-glacierised moraines and a transition towards an uncoupled snout (Benn et al., 2003). Unlike coupled ice margins, the meltwater produced in uncoupled systems is inefficient at removing sediment from the foreland, resulting in gradual ice burial by debris (Benn et al., 2003). This is specific to glaciated mountain valleys under drier climate conditions, with lower ice supplies (Benn et al., 2003). In summary, this phase underscores the evolving importance of topography and debris provision in this landsystem. Located between ~1750 and 1900 m a.s.l., Phase 3 moraines may have been deposited between the construction of the M2b and M3 moraines (1900–2000 m a.s.l.) in the Retezat Mountains, and hence likely date to  $15.8^{+0.9/-0.6}$ – $15.2^{+0.7/-0.8}$  ka (Ruszkiczay-Rüdiger et al., 2016, 2021).

The gradual increasing importance of debris supply and topographic confinement for spatio-temporal landsystem change becomes more evident in Phase 4 (Figure 7d). As ice retreats to the plateau margin, possibly under acute climate amelioration, glacier disconnections occur, leaving dead ice masses, niche glaciers and reconstituted glaciers at the valley/trough heads and at the base of valley walls (Benn & Lehmkuhl, 2000; Boston & Lukas, 2019; Davies et al., 2022; Evans, Ewertowski, & Orton, 2016; Evans et al., 2006; Gellatly et al., 1986; Rea et al., 1999; Rea & Evans, 2003; Whalley et al., 1996). The final stages of deglaciation renewed the paraglacial cycle (Ballantyne, 2002). In other words, RSF activity intensified, leading to increased supraglacial inputs, which altered the glaciers' responses to climate drivers, by insulating ice surfaces (Benn et al., 2003; Benn & Owen, 2002). Where buried ice persisted, obscured and/or modified glacial landforms, it led to creep, resulting in the formation of looped rock-glacierised moraines and rock glaciers on valley and cirque floors. The occurrence of freshly uncovered valley-side hollows, often bounded downslope by lateral moraines, became prime sites for the development of snowpatches, allowing pronival and proglacial ramparts to accumulate. This phase is compatible with the cirque/niche glacial landsystem of Bickerdike, Ó Cofaigh, et al. (2018). Therein, the recession of topographically confined and small ice masses at valley heads and along valley sides is documented by sequences of recessional moraines, or single arcuate moraine ridges, and the gradual development of pronival/proglacial ramparts, rock glaciers and RSFs (Bickerdike, Ó Cofaigh, et al., 2018; Ellis & Calkin, 1979). In the Godeanu Mountains, the cirque/niche glacial landsystem is found in the north-facing Şincu cirque and the south-facing Olanu and Balmoş cirques (Figures 1c and 2).

Also evident during Phase 4 is the role of topography in creating a geomorphic contrast between north and south-facing valleys. Some valleys contain far fewer recessional moraines and are devoid of medial moraines (e.g., the north-facing Cârnea, Mâţu, Scurtele and Scărişoara valleys and the south-facing Scăriţa and Micuşa valleys;

Figures 1c and 2). This contrast is likely a product of the variable occurrence of bedrock steps, which may act as topographic pinning points. Bedrock steps provide marginal stability, limiting variations in glacier accumulation areas when ELAs are coincident with them (Barr & Lovell, 2014; Dugmore, 1989; Oerlemans, 1989). Such topographic pinning points favoured the formation of rare, large moraines in north-facing valleys, contrasting with the small, but more numerous moraines that accumulated in shallower and smoother south-facing valleys (Barr & Lovell, 2014; Figure 2). Situated at ~1900–1950 m a.s.l., Phase 4 moraines may be equivalent to the M3 moraines in the Retezat Mountains (Ruszkiczay-Rüdiger et al., 2016, 2021). A further phase of moraine deposition (M4 at 2100–2150 m a.s.l.) was dated at  $14.4^{±0.5}$ – $13.5^{+0.5/-0.4}$  ka in the Retezat Mountains, which have a higher elevation than the Godeanu Mountains (max = 2509 m a.s.l.; Ruszkiczay-Rüdiger et al., 2016, 2021). This suggests that topography may have an inhibiting role in sustaining glaciers, lower elevations promote higher temperatures and lower precipitation, which are not compatible with glacier growth/persistence (DeBeer & Sharp, 2009). This is especially relevant for the cirques in the Godeanu Mountains, generally peaking at lower elevations (~1750–2000 m a.s.l.) than the plateau-fed valleys in the Godeanu Mountains (~2000–2250 m a.s.l.) (Figures 1c and 2) and the cirque/valley glaciers described for the Retezat Mountains (Reuther et al., 2007; Ruszkiczay-Rüdiger et al., 2016, 2021). Secondly, plateau-fed glaciers alter ice dynamics by either (a) depressing ELAs, leading to a lower elevation moraines than in cirque/valley settings (Aa, 1996; Dahl & Nesje, 1992; Rea et al., 1999), or (b) rapidly retreating to their plateau surfaces during a warming climate, leaving disconnected ice masses in the ablation area, which may or may not imprint geomorphological evidence (Rea et al., 1999; Whalley et al., 1996). Therefore, undated M4 moraines may be found at lower elevations in the plateau-fed valleys in the Godeanu mountains than in the Retezat, subject to confirmation by an absolute dating programme therein.

## 5.2 | Wider implications for the debris-charged plateau icefield/cirque glacier landsystem

The debris-charged plateau icefield/cirque glacier landsystem developed here is the first glacial landsystem devised for mid-altitude mountain ranges situated in temperate-continental climates. Considering the similarity of topographic characteristics, this landsystem model will have applicability to other glaciated landscapes in the Carpathians and beyond. In the Romanian Carpathians, plateau icefield styles of glaciation will likely be relevant to mountain areas with existing plateau surfaces above 1800 m a.s.l., for example, in the Rodna, Maramureş and Călimani Mountains (Balaban, 2018; Kłapyta, Mîndrescu, & Zasadni, 2023; Mîndrescu, 2001; Sîrcu, 1978; Urdea et al., 2011) in the Eastern Carpathians and the Țarcu, Retezat, Şureanu, Cîndrel, Lotru, Făgăraş, Iezer and Bucegi Mountains in the Southern Carpathians (Urdea et al., 2022d). In the central-eastern region of southern Europe, plateau icefields have been reconstructed, for example, in the Pindus Mountains, the Dinaric Alps, and Rila Mountains in the Balkans (e.g., Hughes et al., 2006, 2011, 2022; Kuhlemann, Gachev, et al., 2013), and in the Geyikdağ Mountains in Anatolia (e.g., Akçar, 2022; Çiner et al., 2015). However, the extent and timing of operation of these plateau icefields may differ from that

of the Godeanu Mountains, and therefore, a careful assessment of topographic, lithological and climatic contexts is needed before making more direct comparisons. Likewise, the debris-rich cirque glacier component of the landsystem is not dissimilar to that of the majority of former cirque and valley glaciers in all the Romanian Carpathians (Urdea et al., 2022a, 2022b, 2022c, 2022d, 2022e; Urdea et al., 2022) and the Western Carpathians (e.g., Tatra Mountains) (Zasadni, Kłapyta, Kałuża, & Makos, 2022; Zasadni et al., 2022a, 2022b, 2022c; Zasadni, Kłapyta, Tołoczko-Pasek, & Makos, 2022; Zasadni, Makos, & Kłapyta, 2022). Indeed, previous research in the Romanian Carpathians highlighted a glacial geomorphology throughout the last deglaciation that was characterised by debris-charged palaeoglaciers (e.g., Balaban, 2018; Gheorghiu, 2012; Gheorghiu et al., 2015; Kłapyta et al., 2021, 2022; Kłapyta, Mîndrescu, & Zasadni, 2023; Kłapyta, Zasadni, & Mîndrescu, 2023; László et al., 2013; Reuther et al., 2007; Ruszkiczay-Rüdiger et al., 2016, 2021), but the exact timing and causes of glacial recession have not been assessed in detail (Popescu, Urdea, & Vespremeanu-Stroe, 2017). Consequently, there is a need to apply a landsystem approach to geomorphological mapping and couple it with radiometric dating programmes and numerical ice models to enable robust comparisons of spatio-temporal changes in glaciation style across a wider region.

## 6 | CONCLUSIONS

Drawing on geomorphological mapping and sedimentological investigations, we devise a glacial landsystem model for the Godeanu Mountains, Southern Carpathians, Romania, to understand past glaciation styles and dynamics of mid-altitude mountain glaciers located in temperate-continental climates. We derive the following conclusions:

Firstly, we identified glacial depositional (latero-frontal, recessional, lateral, medial moraines, drift limits/till veneers), glacial erosional (ice-moulded bedrock, ice-marginal meltwater channels), periglacial (pronival and protalus ramparts, relict rock glaciers and rock-glacierised moraines) and paraglacial (RSFs) landforms. The spatial distribution and association of landforms are compatible with a debris-rich glacier retreat, sourced from plateau surfaces/cirques.

Secondly, we performed sedimentological analyses on selected glacial landforms. These further support a debris-charged glacial environment, with transport pathways transitioning from passive to active with distance from valley heads and sides. Although active transport in the lower valleys relates to subglacial erosion from the bed mostly during the earlier stages of deglaciation, passive transport became more dominant in the latter stages, wherein abundant extraglacial sediment was sourced from cirque heads/valley slopes and transported via well-preserved medial moraines to the glacier foreland.

Thirdly, we produce a four-phase, time-transgressive, debris-charged plateau icefield/cirque glacial landsystem model. Phase 1 corresponds to the maximum glacial extent, when plateau ice coalesced with cirque ice, highlighting a transition from cold-based uplands to the warm/polythermal regimes in the valleys. Alongside material sourced from active bedrock erosion at lower elevations, extraglacial debris from RSFs at the valley heads was transported passively via medial moraines and deposited as latero-frontal ridges at the glacier snout. Phase 2 follows ice retreat from the maximum extent. The association of ice-marginal meltwater channels with recessional

moraines is evidence of a warm-based/polythermal coupled ice margin, with a high debris turnover. Phase 3 underpins continual active but increasingly debris-charged recession and disconnection between cirque glaciers and plateau-fed outlets. This is supported by recessional moraines and looped medial moraines pertaining to individual ice flow units and asymmetrical lateral moraines, all of which reflect additional sediment supply sourced from fresh RSFs and the gradually uncovered ice-moulded bedrock ridges. Phase 4 illustrates the final retreat of the cirque and plateau outlets to their upland sources, the increased confinement of cirque glaciers to their niches and the more important role of debris and topography in driving landscape evolution. The likely climatic amelioration and abundance of extraglacial debris from debuttressed slopes led to the burial and uncoupling of glacial snouts from their foreland, which then became inefficient at removing the sediment load. This is supported by the development of pronival and protalus ramparts, rock glaciers and rock-glacierised moraines, which became more dominant over recessional moraines. Ultimately, the role of bedrock steps as topographic pinning points, and the elevation and topographic contrasts between plateau and cirque-fed glaciers explain the contrasts in moraine size, spacing and elevation between different outlets of the Godeanu Mountains and higher alpine summits in other areas of the Southern Carpathians.

Finally, the new landsystem model can be used as an analogue for other plateau/cirque-flanked mountains in Central, Eastern and Southern Europe that exhibit a debris-rich geomorphic signature on their former glacier beds but ideally in conjunction with absolute dating and numerical modelling of glacier extents.

### AUTHOR CONTRIBUTIONS

Cristina I. Balaban co-designed the study, obtained funding for fieldwork, designed the methodology, conducted fieldwork and analysis and drafted the initial and final manuscript. David H. Roberts co-designed the study, supported the methodology and fieldwork, supervised CIB and helped draft the final manuscript. David J. A. Evans co-designed the study, supported the methodology, supervised CIB and helped draft the final manuscript. Stewart S. R. Jamieson co-designed the study, supported the methodology, supervised CIB and helped draft the final manuscript.

### ACKNOWLEDGEMENTS

CIB is in receipt of a Durham Doctoral Studentship, Durham University. Fieldwork was funded through a Norman Richardson Postgraduate Research Fund (Ustinov College, Durham), a New Research Worker's Award (Quaternary Research Association) and a Dudley Stamp Memorial Award (Royal Geographical Society-IBG). We thank Kelan Ford for field assistance and the Domogled-Valea Cernei and Retezat National Parks for access permits and logistical support. We acknowledge the thorough and helpful comments of the two anonymous reviewers and the editors, which improved the manuscript.

### DATA AVAILABILITY STATEMENT

A full resolution of the geomorphological map (without the dashed, numbered rectangles) is available in Figure S1 (Supporting Information). The shapefiles of the mapped landforms are available upon request.

## ORCID

Cristina I. Balaban  <https://orcid.org/0000-0003-4248-363X>  
 David H. Roberts  <https://orcid.org/0000-0002-5976-8423>  
 David J. A. Evans  <https://orcid.org/0000-0002-1840-2513>  
 Stewart S. R. Jamieson  <https://orcid.org/0000-0002-9036-2317>

## REFERENCES

- Aa, A.R. (1996) Topographic control of equilibrium-line altitude depression on reconstructed 'Little Ice Age' glaciers, Grovabreen, western Norway. *The Holocene*, 6(1), 82–89.
- Akçar, N. (2022) Chapter 63 - The Anatolian Mountains: glacial landforms from the Last Glacial Maximum. In: *European Glacial Landscapes: Maximum Extent of Glaciation*. Amsterdam: Elsevier, pp. 497–504. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00016-9>
- Anderson, R.S., Anderson, L.S., Armstrong, W.H., Rossi, M.W. & Crump, S.E. (2018) Glaciation of alpine valleys: the glacier–debris-covered glacier–rock glacier continuum. *Geomorphology*, 311, 127–142. Available from: <https://doi.org/10.1016/j.geomorph.2018.03.015>
- Ardelean, M. (2010) *Munții Piule-Iorgovanu: studiu geomorfologic*. Unpublished doctoral dissertation. “Babeş-Bolyai” University, Cluj-Napoca.
- Balaban, C.I. (2018) *The Glacial History of the Bistricioara-Izvorul Cailor Cirque Complex, Rodna Mountains, Eastern Carpathians, Romania: a glacial geomorphology and numerical modelling investigation*. (Unpublished undergraduate dissertation), Durham University, 65 pp.
- Ballantyne, C.K. (2002) Paraglacial geomorphology. *Quaternary Science Reviews*, 21(18–19), 1935–2017. Available from: [https://doi.org/10.1016/S0277-3791\(02\)00005-7](https://doi.org/10.1016/S0277-3791(02)00005-7)
- Ballantyne, C.K. (2013) Lateglacial rock slope failures in the Scottish Highlands. *Scottish Geographical Journal*, 129(2), 67–84. Available from: <https://doi.org/10.1080/14702541.2013.781210>
- Ballantyne, C.K. (2018a) Glacially moulded landslide runout debris in the Scottish Highlands. *Scottish Geographical Journal*, 134, 224–236. Available from: <https://doi.org/10.1080/14702541.2018.1501085>
- Ballantyne, C.K. (2018b) *Periglacial geomorphology*. Oxford: Wiley Blackwell.
- Ballantyne, C.K. (2021) Rock-slope failures in the North West Highlands. In: Ballantyne, C.K. & Gordon, J.E. (Eds.) *Landscapes and Landforms of Scotland*. Switzerland: Springer, pp. 271–287.
- Ballantyne, C.K. & Benn, D.I. (1994) Glaciological constraints on proglacial rampart development. *Permafrost and Periglacial Processes*, 5, 145–153. Available from: <https://doi.org/10.1002/ppp.3430050304>
- Ballantyne, C.K. & Dawson, A.G. (2019) Scottish landform examples 45: Sgriob na Caillich: a landslide-sourced medial moraine on the Isle of Jura. *Scottish Geographical Journal*, 135(1–2), 139–149. Available from: <https://doi.org/10.1080/14702541.2019.1669812>
- Ballantyne, C.K. & Harris, C. (1994) *The Periglaciation of Great Britain*. Cambridge: Cambridge University Press.
- Ballantyne, C.K. & Kirkbride, M.P. (1986) The characteristics and significance of some Lateglacial proglacial ramparts in upland Britain. *Earth Surface Processes and Landforms*, 11(6), 659–671. Available from: <https://doi.org/10.1002/esp.3290110609>
- Ballantyne, C.K., Sandeman, G.F., Stone, J.O. & Wilson, P. (2014) Rock-slope failure following Late Pleistocene deglaciation on tectonically stable mountainous terrain. *Quaternary Science Reviews*, 86, 144–157. Available from: <https://doi.org/10.1016/j.quascirev.2013.12.021>
- Ballantyne, C.K., Schnabel, C. & Xu, S. (2009) Exposure dating and reinterpretation of coarse debris accumulations ('rock glaciers') in the Cairngorm Mountains, Scotland. *Journal of Quaternary Science*, 24, 19–31. Available from: <https://doi.org/10.1002/jqs.1189>
- Barr, I.D. & Lovell, H. (2014) A review of topographic controls on moraine distribution. *Geomorphology*, 226, 44–64. Available from: <https://doi.org/10.1016/j.geomorph.2014.07.030>
- Beck, H.E., Zimmermann, N.E., McVicar, T.R., Vergopolan, N., Berg, A. & Wood, E.F. (2018) Present and future Köppen-Geiger climate classification maps at 1-km resolution. *Scientific Data*, 5, 1–12. Available from: <https://doi.org/10.1038/sdata.2018.214>
- Benn, D.I. (1989) Debris transport by Loch Lomond Readvance glaciers in Northern Scotland: basin form and the within-valley asymmetry of lateral moraines. *Journal of Quaternary Science*, 4, 243–254. Available from: <https://doi.org/10.1002/jqs.3390040305>
- Benn, D.I. & Ballantyne, C.K. (1994) Reconstructing the transport history of glacial sediments: a new approach based on the covariance of clast shape indices. *Sedimentary Geology*, 91, 215–227. Available from: [https://doi.org/10.1016/0037-0738\(94\)90130-9](https://doi.org/10.1016/0037-0738(94)90130-9)
- Benn, D.I. & Ballantyne, C.K. (2005) Palaeoclimatic reconstruction from Loch Lomond readvance glaciers in the west Drumochter Hills, Scotland. *Journal of Quaternary Science*, 20(6), 577–592. Available from: <https://doi.org/10.1002/jqs.925>
- Benn, D.I. & Evans, D.J.A. (2010) *Glaciers and Glaciation*, Second edition. London: Hodder.
- Benn, D.I., Kirkbride, M.P., Owen, L.A. & Brazier, V. (2003) Glaciated valley landsystems. In: Evans, D.J.A. (Ed.) *Glacial landsystems*. London: Arnold, pp. 372–406.
- Benn, D.I. & Lehmkuhl, F. (2000) Mass balance and equilibrium-line altitudes of glaciers in high-mountain environments. *Quaternary International*, 65, 15–29. Available from: [https://doi.org/10.1016/S1040-6182\(99\)00034-8](https://doi.org/10.1016/S1040-6182(99)00034-8)
- Benn, D.I. & Lukas, S. (2006) Younger Dryas glacial landsystems in North West Scotland: an assessment of modern analogues and palaeoclimatic implications. *Quaternary Science Reviews*, 25(17–18), 2390–2408. Available from: <https://doi.org/10.1016/j.quascirev.2006.02.015>
- Benn, D.I. & Lukas, S. (2021) Clast morphology. In: Evans, D.J.A. & Benn, D.I. (Eds.) *A Practical Guide to the Study of Glacial Sediments*. London: Quaternary Research Association, pp. 107–124.
- Benn, D.I. & Owen, L.A. (2002) Himalayan glacial sedimentary environments: a framework for reconstructing and dating the former extent of glaciers in high mountains. *Quaternary International*, 97, 3–25. Available from: [https://doi.org/10.1016/S1040-6182\(02\)00048-4](https://doi.org/10.1016/S1040-6182(02)00048-4)
- Bennett, G.L., Evans, D.J.A., Carbonneau, P. & Twigg, D.R. (2010) Evolution of a debris-charged glacier landsystem, Kviárjökull, Iceland. *Journal of Maps*, 2010, 40–76. Available from: <https://doi.org/10.4113/jom.2010.1114>
- Bennett, M.R. & Boulton, G.S. (1993) Deglaciation of the Younger Dryas or Loch Lomond Stadial ice-field in the northern Highlands, Scotland. *Journal of Quaternary Science*, 8(2), 133–145. Available from: <https://doi.org/10.1002/jqs.3390080206>
- Berthling, I. (2011) Beyond confusion: rock glaciers as cryo-conditioned landforms. *Geomorphology*, 131, 98–105. Available from: <https://doi.org/10.1016/j.geomorph.2011.05.002>
- Bickerdike, H.L., Evans, D.J.A., Stokes, C.R. & Ó Cofaigh, C. (2018) The glacial geomorphology of the Loch Lomond (Younger Dryas) Stadial in Britain. *Journal of Quaternary Science*, 33(1), 1–54. Available from: <https://doi.org/10.1002/jqs.3010>
- Bickerdike, H.L., Ó Cofaigh, C., Evans, D.J.A. & Stokes, C.R. (2018) Glacial landsystems, retreat dynamics and controls on Loch Lomond Stadial (Younger Dryas) glaciation in Britain. *Boreas*, 47(1), 202–224. Available from: <https://doi.org/10.1111/bor.12259>
- Boston, C.M. & Lukas, S. (2019) Topographic controls on plateau icefield recession: insights from the Younger Dryas Monadhliath Icefield, Scotland. *Journal of Quaternary Science*, 34, 433–451. Available from: <https://doi.org/10.1002/jqs.3111>
- Boulton, G.S. (1978) Boulder shapes and grain-size distributions of debris as indicators of transport paths through a glacier and till genesis. *Sedimentology*, 25(6), 773–799. Available from: <https://doi.org/10.1111/j.1365-3091.1978.tb00329.x>
- Carr, S.J. & Coleman, C.G. (2007) An improved technique for the reconstruction of former glacier mass-balance and dynamics. *Geomorphology*, 92, 76–90. Available from: <https://doi.org/10.1016/j.geomorph.2007.02.008>
- Chandler, B.M.P., Lovell, H., Boston, C.M., Lukas, S., Barr, I.D., Benediktson, Í.Ö. et al. (2018) Glacial geomorphological mapping: a review of approaches and frameworks for best practice. *Earth-Science Reviews*, 185, 806–846. Available from: <https://doi.org/10.1016/j.earscirev.2018.07.015>

- Çiner, A., Sarikaya, M.A. & Yildirim, C. (2015) Late Pleistocene piedmont glaciations in the Eastern Mediterranean; insights from cosmogenic <sup>36</sup>Cl dating of hummocky moraines in southern Turkey. *Quaternary Science Reviews*, 116, 44–56. Available from: <https://doi.org/10.1016/j.quascirev.2015.03.017>
- Clark, R. & Wilson, P. (2004) A rock avalanche deposit in Burtness Comb, Lake District, Northwest England. *Geological Journal*, 39, 419–430. Available from: <https://doi.org/10.1002/gj.965>
- Coleman, C.G. & Carr, S.J. (2008) Complex relationships between Younger Dryas glacial, periglacial and paraglacial landforms, Brecon Beacons, South Wales. *Proceedings of the Geologists' Association*, 119, 259–276. Available from: [https://doi.org/10.1016/S0016-7878\(08\)80305-7](https://doi.org/10.1016/S0016-7878(08)80305-7)
- Coleman, C.G., Carr, S.J. & Parker, A.G. (2009) Modelling topoclimatic controls on palaeoglaciers: implications for inferring palaeoclimate from geomorphic evidence. *Quaternary Science Reviews*, 28, 249–259. Available from: <https://doi.org/10.1016/j.quascirev.2008.10.016>
- Colucci, R.R., Boccali, C., Žebre, M. & Guglielmin, M. (2016) Rock glaciers, protalus ramparts and pronaival ramparts in the south-eastern Alps. *Geomorphology*, 269, 112–121. Available from: <https://doi.org/10.1016/j.geomorph.2016.06.039>
- Czirbusz, G. (1905) A Godjan Gugu hegység. *Földrajzi Közlemények*, t. XXXIII, fasc. 5, 190–201.
- Dahl, S. O. & Nesje, A. (1992) Paleoclimatic implications based on equilibrium-line altitude depressions of reconstructed Younger Dryas and Holocene cirque glaciers in inner Nordfjord, western Norway. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 94(1–4), 87–97. Available from: [https://doi.org/10.1016/0031-0182\(92\)90114-K](https://doi.org/10.1016/0031-0182(92)90114-K)
- Darvill, C.M., Stokes, C.R., Bentley, M.J., Evans, D.J.A. & Lovell, H. (2017) Dynamics of former ice lobes of the southernmost Patagonian Ice Sheet based on a glacial land systems approach. *Journal of Quaternary Science*, 32(6), 857–876. Available from: <https://doi.org/10.1002/jqs.2890>
- Davies, B., Bendle, J., Carrivick, J., McNabb, R., McNeil, C., Pelto, M. et al. (2022) Topographic controls on ice flow and recession for Juneau Icefield (Alaska/British Columbia). *Earth Surface Processes and Landforms*, 47, 2357–2390. Available from: <https://doi.org/10.1002/esp.5383>
- De Martonne, E. (1900) Contribution à l'étude de la période glaciaire dans les Karpatés méridionales, *Bulletin de la Société Géologique de France*, Paris, 3 série, t. XXVIII, p. 274–319.
- DeBeer, C.M. & Sharp, M.J. (2009) Topographic influences on recent changes of very small glaciers in the Monashee Mountains, British Columbia, Canada. *Journal of Glaciology*, 55, 691–700. Available from: <https://doi.org/10.3189/002214309789470851>
- Dragomir, G., Șerban, A., Năstase, G. & Brezeanu, A.I. (2016) Wind energy in Romania: a review from 2009 to 2016. *Renewable and Sustainable Energy Reviews*, 64, 129–143. Available from: <https://doi.org/10.1016/j.rser.2016.05.080>
- Dugmore, A.J. (1989) Tephrochronological studies of Holocene glacier fluctuations in south Iceland. In: Oerlemans, J. (Ed.) *Glacier Fluctuations and Climatic Change*. Dordrecht: Springer, pp. 37–55.
- Dyke, A.S. (1993) Landscapes of cold-centred Late Wisconsinan ice caps, Arctic Canada. *Progress in Physical Geography*, 17, 223–247. Available from: <https://doi.org/10.1177/030913339301700208>
- Ellis, J.M. & Calkin, P.E. (1979) Nature and distribution of glaciers, neoglacial moraines, and rock glaciers, east-central Brooks Range, Alaska. *Arctic and Alpine Research*, 11, 403–420.
- England, J. (1978) The glacial geology of northeastern Ellesmere Island, NWT, Canada. *Canadian Journal of Earth Sciences*, 15(4), 603–617. Available from: <https://doi.org/10.1139/e78-065>
- England, J. (1990) The late Quaternary history of Greely Fiord and its tributaries, west-central Ellesmere Island. *Canadian Journal of Earth Sciences*, 27, 255–270. Available from: <https://doi.org/10.1139/e90-025>
- Evans, D.J.A. (1990) The last glaciation and relative sea level history of northwest Ellesmere Island, Canadian High Arctic. *Journal of Quaternary Science*, 5(1), 67–82. Available from: <https://doi.org/10.1002/jqs.3390050107>
- Evans, D.J.A. (1999) Glacial debris transport and moraine deposition: a case study of the Jardalen cirque complex, Sogn-og-Fjordane, western Norway. *Zeitschrift für Geomorphologie*, 43, 203–234. Available from: <https://doi.org/10.1127/zfg/43/1999/203>
- Evans, D.J.A. (Ed). (2003) *Glacial Land systems*. London: Arnold.
- Evans, D.J.A. (2010) Controlled moraine development and debris transport pathways in polythermal plateau icefields: examples from Tungnafellsjökull, Iceland. *Earth Surface Processes and Landforms*, 35(12), 1430–1444. Available from: <https://doi.org/10.1002/esp.1984>
- Evans, D.J.A. (2011) Glacial land systems of Satujökull, Iceland: a modern analogue for glacial land system overprinting by mountain icecaps. *Geomorphology*, 129, 225–237. Available from: <https://doi.org/10.1016/j.geomorph.2011.01.025>
- Evans, D.J.A. (2013a) Geomorphology and retreating glaciers. In: Shroder, J., Giardino, R. & Harbor, J. (Eds.) *Treatise on Geomorphology, Volume 8, Glacial and Periglacial Geomorphology*. San Diego: Academic Press, pp. 460–478.
- Evans, D.J.A. (2013b) Moraine forms and genesis. In: Elias, S.A. & Mock, C.J. (Eds.) *The Encyclopedia of Quaternary Science*, Second edition. Amsterdam: Elsevier, pp. 769–779.
- Evans, D.J.A. & Benn, D.I. (Eds). (2021) *A Practical Guide to the Study of Glacial Sediments*, Second edition. London: QRA.
- Evans, D.J.A., Dinnage, M. & Roberts, D.H. (2018) Geomorphology of Teesdale, northern Pennines, England: implications for upland styles of ice stream operation and deglaciation in the British-Irish Ice Sheet. *Proceedings of the Geologists' Association*, 129, 697–735. Available from: <https://doi.org/10.1016/j.pgeola.2018.05.001>
- Evans, D.J.A., Ewertowski, M. & Orton, C. (2016) Eiríksjökull plateau icefield land system, Iceland. *Journal of Maps*, 12, 747–756. Available from: <https://doi.org/10.1080/17445647.2015.1072448>
- Evans, D.J.A., Ewertowski, M. & Orton, C. (2017) The glaciated valley land system of Morsárjökull, southeast Iceland. *Journal of Maps*, 13, 909–920. Available from: <https://doi.org/10.1080/17445647.2017.1401491>
- Evans, D.J.A., Hughes, A.L.C., Hansom, J.D. & Roberts, D.H. (2017) Scottish Landform Examples 43: Glacifluvial landforms of Strathallan, Perthshire. *Scottish Geographical Journal*, 133, 42–53. Available from: <https://doi.org/10.1080/14702541.2016.1254276>
- Evans, D.J.A. & Rea, B.R. (2003) Surging glacier land system. In: Evans, D.J.A. (Ed.) *Glacial Land systems*. London: Arnold, pp. 259–288.
- Evans, D.J.A., Roberts, D.H. & Evans, S.C. (2016) Multiple subglacial till deposition: a modern exemplar for Quaternary palaeoglaciology. *Quaternary Science Reviews*, 145, 1–21. Available from: <https://doi.org/10.1016/j.quascirev.2016.05.029>
- Evans, D.J.A., Roberts, D.H., Hiemstra, J.F., Nye, K.M., Wright, H. & Steer, A. (2018) Submarginal debris transport and till formation in active temperate glacier systems: the southeast Iceland type locality. *Quaternary Science Reviews*, 195, 72–108. Available from: <https://doi.org/10.1016/j.quascirev.2018.07.002>
- Evans, D.J.A., Shulmeister, J. & Hyatt, O. (2010) Sedimentology of latero-frontal moraines and fans on the west coast of South Island, New Zealand. *Quaternary Science Reviews*, 29, 3790–3811. Available from: <https://doi.org/10.1016/j.quascirev.2010.08.019>
- Evans, D.J.A., Twigg, D.R. & Shand, M. (2006) Surficial geology and geomorphology of the Þórisjökull plateau icefield, west-central Iceland. *Journal of Maps*, 2(1), 17–29. Available from: <https://doi.org/10.4113/jom.2006.52>
- Evans, I.S. (1996) Abraded rock landforms (whalebacks) developed under ice streams in mountain areas. *Annals of Glaciology*, 22, 9–16. Available from: <https://doi.org/10.3189/1996AoG22-1-9-16>
- Evans, I.S. (2021) Glaciers, rock avalanches and the 'buzzsaw' in cirque development: why mountain cirques are of mainly glacial origin. *Earth Surface Processes and Landforms*, 46(1), 24–46. Available from: <https://doi.org/10.1002/esp.4810>
- Eyles, N. & Rogerson, R.J. (1978) A framework for the investigation of medial moraine formation: Austerdalsbreen, Norway, and Berendon Glacier, British Columbia, Canada. *Journal of Glaciology*, 20, 99–113. Available from: <https://doi.org/10.3189/S0022143000021249>
- Federici, P.R., Ribolini, A. & Spagnolo, M. (2017) Glacial history of the Maritime Alps from the Last Glacial Maximum to the Little Ice Age. In: Hughes, P.D. & Woodward, J.C. (Eds.) *Quaternary Glaciation in the*

- Mediterranean Mountains*, Vol. 433. London, Special Publication: Geological Society, pp. 137–159. Available from: <https://doi.org/10.1144/SP433.9>
- Gellatly, A.F., Whalley, W.B. & Gordon, J.E. (1986) Topographic control over recent glacier changes in southern Lyngen Peninsula, North Norway. *Norsk Geografisk Tidsskrift - Norwegian Journal of Geography*, 40, 211–218. Available from: <https://doi.org/10.1080/00291958608552166>
- Gheorghiu, D.M. (2012) *Testing climate synchronicity between Scotland and Romania since the last glacial maximum*. Unpublished PhD thesis, University of Glasgow. Available from: <http://theses.gla.ac.uk/3362/>
- Gheorghiu, D.M., Hosu, M., Corpade, C. & Xu, S. (2015) Deglaciation constraints in the Parâng Mountains, Southern Romania, using surface exposure dating. *Quaternary International*, 388, 156–167. Available from: <https://doi.org/10.1016/j.quaint.2015.04.059>
- Giles, D.P., Griffiths, J.S., Evans, D.J.A. & Murton, J.B. (2017) Geomorphological framework: glacial and periglacial sediments, structures and landforms. In: Griffiths, J.S. & Martin, C.J. (Eds.) *Engineering Geology and Geomorphology of Glaciated and Periglaciated Terrains*, Vol. 28. Geological Society, London, Engineering Geology Special Publication, pp. 59–368. Available from: <https://doi.org/10.1144/EGSP28.3>
- Glasser, N.F. (2002) The large roches moutonnées of Upper Deeside. *Scottish Geographical Journal*, 118, 129–138. Available from: <https://doi.org/10.1080/00369220218737141>
- Glasser, N.F. & Bennett, M.R. (2004) Glacial erosional landforms: origins and significance for palaeoglaciology. *Progress in Physical Geography*, 28(1), 43–75. Available from: <https://doi.org/10.1191/0309133304pp401ra>
- Glasser, N.F., Harrison, S., Ivy-Ochs, S., Duller, G.A. & Kubik, P.W. (2006) Evidence from the Rio Bayo valley on the extent of the North Patagonian Icefield during the Late Pleistocene–Holocene transition. *Quaternary Research*, 65, 70–77. Available from: <https://doi.org/10.1016/j.yqres.2005.09.002>
- Glasser, N.F., Jansson, K.N., Harrison, S. & Kleman, J. (2008) The glacial geomorphology and Pleistocene history of South America between 38 S and 56 S. *Quaternary Science Reviews*, 27, 365–390. Available from: <https://doi.org/10.1016/j.quascirev.2007.11.011>
- Golledge, N.R. (2007) An ice cap landsystem for palaeoglaciological reconstructions: characterizing the Younger Dryas in western Scotland. *Quaternary Science Reviews*, 26, 213–229. Available from: <https://doi.org/10.1016/j.quascirev.2006.08.007>
- Goodsell, B., Hambrey, M.J. & Glasser, N.F. (2005) Debris transport in a temperate valley glacier: Haut Glacier d'Arolla, Valais, Switzerland. *Journal of Glaciology*, 51(172), 139–146. Available from: <https://doi.org/10.3189/172756505781829647>
- Goodsell, B., Hambrey, M.J., Glasser, N.F., Nienow, P. & Mair, D. (2005) The structural glaciology of a temperate valley glacier: Haut Glacier d'Arolla, Valais, Switzerland. *Arctic, Antarctic, and Alpine Research*, 37(2), 218–232. Available from: [https://doi.org/10.1657/1523-0430\(2005\)037\[0218:TSGOAT\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0218:TSGOAT]2.0.CO;2)
- Gordon, J.E. (1981) Glacier margin fluctuations during the 19th and 20th centuries in the Ikamiut Kangerdluarssuat area, west Greenland. *Arctic and Alpine Research*, 13(1), 47–62. Available from: <https://doi.org/10.2307/1550625>
- Greenwood, S.L., Clark, C.D. & Hughes, A.L. (2007) Formalising an inversion methodology for reconstructing ice-sheet retreat patterns from meltwater channels: application to the British Ice Sheet. *Journal of Quaternary Science*, 22(6), 637–645. Available from: <https://doi.org/10.1002/jqs.1083>
- Gunnell, Y., Blondeau, S. & Jarman, D. (2022) Rock slope failure in the Southern Carpathians (Romania): range-wide inventory and links with long-term mountain landscape evolution. *Geomorphology*, 418, 108433. Available from: <https://doi.org/10.1016/j.geomorph.2022.108433>
- Gusmeroli, A., Wolken, G.J. & Arendt, A.A. (2014) Helicopter-borne radar imaging of snow cover on and around glaciers in Alaska. *Annals of Glaciology*, 55, 78–88. Available from: <https://doi.org/10.3189/2014AoG67A029>
- Hamilton, S.J. & Whalley, W.B. (1995) Rock glacier nomenclature: a reassessment. *Geomorphology*, 14(1), 73–80. Available from: [https://doi.org/10.1016/0169-555X\(95\)00036-5](https://doi.org/10.1016/0169-555X(95)00036-5)
- Harrison, S., Whalley, B. & Anderson, E. (2008) Relict rock glaciers and protalus lobes in the British Isles: implications for Late Pleistocene mountain geomorphology and palaeoclimate. *Journal of Quaternary Science*, 23(3), 287–304. Available from: <https://doi.org/10.1002/jqs.1148>
- Hedding, D.W. (2011) Pronival rampart and protalus rampart: a review of terminology. *Journal of Glaciology*, 57, 1179–1180. Available from: <https://doi.org/10.3189/002214311798843241>
- Hedding, D.W. (2016) Pronival ramparts: a review. *Progress in Physical Geography*, 40, 835–855. Available from: <https://doi.org/10.1177/0309133316678148>
- Hedding, D.W. & Sumner, P.D. (2013) Diagnostic criteria for pronival ramparts: site, morphological and sedimentological characteristics. *Geografiska Annaler. Series A, Physical Geography*, 95, 315–322. Available from: <https://doi.org/10.1111/geoa.12021>
- Hedding, D.W., Sumner, P.D., Holness, S.D. & Meiklejohn, K.I. (2007) Formation of a pronival rampart on sub-Antarctic Marion Island. *Antarctic Science*, 19, 443–450. Available from: <https://doi.org/10.1017/S0954102007000582>
- Hughes, P.D., Allard, J.L. & Woodward, J.C. (2022) Chapter 43 - The Balkans: glacial landforms prior to the Last Glacial Maximum. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 323–332. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00034-0>
- Hughes, P.D., Woodward, J.C., Gibbard, P.L., Macklin, M.G., Gilmour, M.A. & Smith, G.R. (2006) The glacial history of the Pindus Mountains, Greece. *Journal of Geology*, 114, 413–434. Available from: <https://doi.org/10.1086/504177>
- Hughes, P.D., Woodward, J.C., Van Calsteren, P.C. & Thomas, L.E. (2011) The glacial history of the Dinaric Alps, Montenegro. *Quaternary Science Reviews*, 30(23–24), 3393–3412. Available from: <https://doi.org/10.1016/j.quascirev.2011.08.016>
- Hulton, N.R.J. & Sugden, D.E. (1997) Dynamics of mountain ice caps during glacial cycles: the case of Patagonia. *Annals of Glaciology*, 24, 81–89. Available from: <https://doi.org/10.3189/S0260305500011976>
- Humlum, O. (1996) Origin of rock glaciers: observations from Mellemfjord, Disko Island, central west Greenland. *Permafrost and Periglacial Processes*, 7, 361–380. Available from: [https://doi.org/10.1002/\(SICI\)1099-1530\(199610\)7:4%3C361::AID-PPP227%3E3.0.CO;2-4](https://doi.org/10.1002/(SICI)1099-1530(199610)7:4%3C361::AID-PPP227%3E3.0.CO;2-4)
- Ignénczi, Á. & Nagy, B. (2016) Former plateau ice fields in the Godeanu Mountains, Southern Carpathians: first evidence of glaciated peninsulas in the Carpathians. *Quaternary International*, 415, 74–85. Available from: <https://doi.org/10.1016/j.quaint.2015.09.058>
- Jarman, D. (2006) Large rock slope failures in the highlands of Scotland: characterisation, causes and spatial distribution. *Engineering Geology*, 83(1–3), 161–182. Available from: <https://doi.org/10.1016/j.enggeo.2005.06.030>
- Jarman, D. (2009) Paraglacial rock slope failure as an agent of glacial trough widening. *Geological Society, London, Special Publications*, 320(1), 103–131. Available from: <https://doi.org/10.1144/SP320.8>
- Jarman, D., Wilson, P. & Harrison, S. (2013) Are there any relict rock glaciers in the British mountains? *Journal of Quaternary Science*, 28(2), 131–143. Available from: <https://doi.org/10.1002/jqs.2574>
- Jennings, S.J.A., Hambrey, M.J. & Glasser, N.F. (2014) Ice flow-unit influence on glacier structure, debris entrainment and transport. *Earth Surface Processes and Landforms*, 39(10), 1279–1292. Available from: <https://doi.org/10.1002/esp.3521>
- Jones, D.B., Harrison, S. & Anderson, K. (2019) Mountain glacier-to-rock glacier transition. *Global and Planetary Change*, 181, 102999. Available from: <https://doi.org/10.1016/j.gloplacha.2019.102999>
- Kłapyta, P., Mîndrescu, M. & Zasadni, J. (2021) Geomorphological record and equilibrium line altitude of glaciers during the last glacial maximum in the Rodna Mountains (eastern Carpathians). *Quaternary Research*, 100, 1–20. Available from: <https://doi.org/10.1017/qua.2020.90>
- Kłapyta, P., Mîndrescu, M. & Zasadni, J. (2022) The impact of local topoclimatic factors on marginal Pleistocene glaciation in the Northern Romanian Carpathians. *Catena*, 210, 105873.1. Available from: <https://doi.org/10.1016/j.catena.2021.105873>

- Kłapyta, P., Mîndrescu, M. & Zasadni, J. (2023) Late Pleistocene glaciation in the headwaters of the Ceremuşul Alb valley (Maramureş Mountains, Romania). *Geographia Polonica*, 1(1), 13–28. Available from: <https://doi.org/10.7163/GPol.0243>
- Kłapyta, P., Zasadni, J. & Mîndrescu, M. (2023) Late Pleistocene glaciation in the Eastern Carpathians—a regional overview. *Catena*, 224, 106994. Available from: <https://doi.org/10.1016/j.catena.2023.106994>
- Knight, J., Harrison, S. & Jones, D.B. (2019) Rock glaciers and the geomorphological evolution of deglaciating mountains. *Geomorphology*, 324, 14–24. Available from: <https://doi.org/10.1016/j.geomorph.2018.09.020>
- Kuhlemann, J., Dobre, F., Urdea, P., Krumrei, I., Gachev, E., Kubik, P. et al. (2013) Last Glacial Maximum glaciation of the Central South Carpathian range (Romania). *Austrian Journal of Earth Sciences*, 106, 83–95.
- Kuhlemann, J., Gachev, E., Gikov, A., Nedkov, S., Krumrei, I. & Kubik, P. (2013) Glaciation in the Rila Mountains (Bulgaria) during the Last Glacial Maximum. *Quaternary International*, 293, 51–62. Available from: <https://doi.org/10.1016/j.quaint.2012.06.027>
- László, P., Kern, Z. & Nagy, B. (2013) Late Pleistocene glaciers in the western Rodna Mountains, Romania. *Quaternary International*, 293, 79–91. Available from: <https://doi.org/10.1016/j.quaint.2012.09.020>
- Lehmann, P.W. (1881) Beobachtungen über Tektonik und Gletscherspuren im Fogaraschen Gebirge. *Zeitschrift. Deutsche Geologische Gesellschaft*, XXXIII, 109–117.
- Leigh, J.R., Evans, D.J.A., Stokes, C.R., Andreassen, L.M. & Carr, R.J. (2021) Glacial and periglacial geomorphology of central Troms and Finnmark county, Arctic Norway. *Journal of Maps*, 17, 348–366. Available from: <https://doi.org/10.1080/17445647.2021.1950580>
- Li, Y., Harbor, J., Liu, G., Yi, C. & Caffee, M.W. (2016) Cosmogenic <sup>10</sup>Be constraints on Little Ice Age glacial advances in the eastern Tian Shan, China. *Quaternary Science Reviews*, 138, 105–118. Available from: <https://doi.org/10.1016/j.quascirev.2016.02.023>
- Livingstone, S.J., Evans, D.J.A., Ó Cofaigh, C. & Hopkins, J. (2010) The Bampton kame belt and Pennine escarpment meltwater channel system (Cumbria, UK): morphology, sedimentology and formation. *Proceedings of the Geologists' Association*, 121, 423–443. Available from: <https://doi.org/10.1016/j.pgeola.2009.10.005>
- Lukas, S. (2005) A test of the englacial thrusting hypothesis of 'hummocky' moraine formation: case studies from the northwest Highlands, Scotland. *Boreas*, 34(3), 287–307. Available from: <https://doi.org/10.1111/j.1502-3885.2005.tb01102.x>
- Lukas, S., Benn, D.I., Boston, C.M., Brook, M., Coray, S., Evans, D.J.A. et al. (2013) Clast shape analysis and clast transport paths in glacial environments: a critical review of methods and the role of lithology. *Earth-Science Reviews*, 121, 96–116. Available from: <https://doi.org/10.1016/j.earscirev.2013.02.005>
- Lukas, S., Graf, A., Coray, S. & Schlüchter, C. (2012) Genesis, stability and preservation potential of large lateral moraines of Alpine valley glaciers—towards a unifying theory based on Findelengletscher, Switzerland. *Quaternary Science Reviews*, 38, 27–48. Available from: <https://doi.org/10.1016/j.quascirev.2012.01.022>
- Maag, H. (1969) *Ice Dammed Lakes and Marginal Glacial Drainage on Axel Heiberg Island*. Axel Heiberg Island Research Report, McGill University, Montreal.
- Magyari, E.K., Jakab, G., Bálint, M., Kern, Z., Buczkó, K. & Braun, M. (2012) Rapid vegetation response to Lateglacial and early Holocene climatic fluctuation in the South Carpathian Mountains (Romania). *Quaternary Science Reviews*, 35, 116–130. Available from: <https://doi.org/10.1016/j.quascirev.2012.01.006>
- Matthews, J.A. & Petch, J.R. (1982) Within-valley asymmetry and related problems of Neoglacial lateral moraine development at certain Jotunheimen glaciers, southern Norway. *Boreas*, 11, 225–247. Available from: <https://doi.org/10.1111/j.1502-3885.1982.tb00716.x>
- Matthews, J.A., Shakesby, R.A., Owen, G. & Vater, A.E. (2011) Protralus rampart formation in relation to snow avalanche activity and Schmidt hammer exposure age dating: three case studies from southern Norway. *Geomorphology*, 130(3–4), 280–288. Available from: <https://doi.org/10.1016/j.geomorph.2011.04.010>
- Meier, M.F. & Post, A.S. (1969) What are glacier surges? *Canadian Journal of Earth Sciences*, 6, 807–819. Available from: <https://doi.org/10.1139/e69-081>
- Micalevich-Velcea, V. (1961) *Masivul Bucegi. Studiu Geomorfologic*. Bucureşti: Ed. Academiei, p. 151.
- Mîndrescu, M. (2001) Lacul Vinderel. Caracterizare hidro-geomorfologica (Muntii Maramuresului). *Analele Universitatii "Stefan cel Mare" Suceava, sectiunea Geografie – Geologie*, nr. X.
- Mîndrescu, M. (2016) *Circurile glaciare din Carpații românești* (geomorphological map).
- Mîndrescu, M. & Evans, I.S. (2014) Cirque form and development in Romania: allometry and the buzzsaw hypothesis. *Geomorphology*, 208, 117–136. Available from: <https://doi.org/10.1016/j.geomorph.2013.11.019>
- Mîndrescu, M. & Evans, I.S. (2017) Glacial cirques in the Romanian Carpathians and their climatic implications. In: Rădoane, M. & Vespremeanu-Stroe, A. (Eds.) *Landform dynamics and evolution in Romania*. Dordrecht: Springer, pp. 197–214. Available from: [https://doi.org/10.1007/978-3-319-32589-7\\_9](https://doi.org/10.1007/978-3-319-32589-7_9)
- Mîndrescu, M., Evans, I.S. & Cox, N.J. (2010) Climatic implications of cirque distribution in the Romanian Carpathians: palaeowind directions during glacial periods. *Journal of Quaternary Science*, 25, 875–888. Available from: <https://doi.org/10.1002/jqs.1363>
- Mitchell, W.A. (1996) Significance of snowblow in the generation of Loch Lomond Stadial (Younger Dryas) glaciers in the western Pennines, northern England. *Journal of Quaternary Science*, 11(3), 233–248. Available from: [https://doi.org/10.1002/\(SICI\)1099-1417\(199605/06\)11:3<233::AID-JQS240>3.0.CO;2-Q](https://doi.org/10.1002/(SICI)1099-1417(199605/06)11:3<233::AID-JQS240>3.0.CO;2-Q)
- Necşoiu, M., Onaca, A., Wigginton, S. & Urdea, P. (2016) Rock glacier dynamics in Southern Carpathian Mountains from high-resolution optical and multi-temporal SAR satellite imagery. *Remote Sensing of Environment*, 177, 21–36. Available from: <https://doi.org/10.1016/j.rse.2016.02.025>
- Nedelcu, E. (1967) Trăsăturile morfostructurale ale Munților Iezerului. *Studii și cercetări de geologie, geofizică, geografie*, 14, 183–191.
- Niculescu, G. (1965) *Munții Godeanu. Studiu geomorfologic*. Bucureşti: Edit. Academiei.
- Oerlemans, J. (1989) On the response of valley glaciers to climatic change. In: *Glacier fluctuations and climatic change*. Dordrecht: Springer, pp. 353–371. Available from: [https://doi.org/10.1007/978-94-015-7823-3\\_23](https://doi.org/10.1007/978-94-015-7823-3_23)
- Oerlemans, J., Anderson, B., Hubbard, A., Huybrechts, P., Johannesson, T., Knap, W.H. et al. (1998) Modelling the response of glaciers to climate warming. *Climate Dynamics*, 14(4), 267–274. Available from: <https://doi.org/10.1007/s003820050222>
- Onaca, A., Ardelean, F., Urdea, P. & Magori, B. (2017) Southern Carpathian rock glaciers: inventory, distribution and environmental controlling factors. *Geomorphology*, 293, 391–404. Available from: <https://doi.org/10.1016/j.geomorph.2016.03.032>
- Onaca, A., Urdea, P., Ardelean, A.C., Şerban, R. & Ardelean, F. (2017) Present-day periglacial processes in the alpine zone. In: Radoane, M. & Vespremeanu-Stroe, A. (Eds.) *Landform Dynamics and Evolution in Romania*. Springer, Dordrecht. Available from: <https://doi.org/10.1007/978-3-319-32589-7>
- Onaca, A.L., Urdea, P. & Ardelean, A.C. (2013) Internal structure and permafrost characteristics of the rock glaciers of Southern Carpathians (Romania) assessed by geoelectrical soundings and thermal monitoring. *Geografiska Annaler. Series A, Physical Geography*, 95, 249–266. Available from: <https://doi.org/10.1111/geoa.12014>
- Østrem, G. (1964) Ice-cored moraines in Scandinavia. *Geografiska Annaler*, 46, 282–337. Available from: <https://doi.org/10.1080/20014422.1964.11881043>
- Owen, L.A., Gualtieri, L., Finkel, C., Caffee, M.W., Benn, D.I. & Sharma, M.C. (2001) Cosmogenic radionuclide dating of glacial landforms in the Lahul Himalaya, northern India: defining the timing of late Quaternary glaciation. *Journal of Quaternary Science*, 16, 555–563. Available from: <https://doi.org/10.1002/jqs.621>
- Paul, F. (2015) Revealing glacier flow and surge dynamics from animated satellite image sequences: examples from the Karakoram. *The Cryosphere*, 9(6), 2201–2214. Available from: <https://doi.org/10.5194/tc-9-2201-2015>

- Popescu, R. (2018) Permafrost investigations in lezer Mountains, Southern Carpathians. *Revista de Geomorfologie*, 20(1), 102–122. Available from: <https://doi.org/10.21094/rg.2018.033>
- Popescu, R., Mîndrescu, M., Cruțeru, N., Calisevici, S. & Barbălată, D. (2021) Combining geomorphological approach and thermal monitoring for permafrost research in Rodna Mountains, Northern Romanian Carpathians. *Forum Geografic. Studii și cercetări de geografie și protecția mediului*, 20, 127–137. Available from: <https://doi.org/10.5775/fg.2021.071.d>
- Popescu, R., Onaca, A., Urdea, P. & Vespremeanu-Stroe, A. (2017) Spatial distribution and main characteristics of alpine permafrost from Southern Carpathians, Romania. In: Rădoane, M. & Vespremeanu-Stroe, A. (Eds.) *Landform Dynamics and Evolution in Romania*. Dordrecht: Springer, pp. 117–146. Available from: [https://doi.org/10.1007/978-3-319-32589-7\\_6](https://doi.org/10.1007/978-3-319-32589-7_6)
- Popescu, R., Urdea, P. & Vespremeanu-Stroe, A. (2017) Deglaciation history of high massifs from the Romanian Carpathians: towards an integrated view. In: Rădoane, M. & Vespremeanu-Stroe, A. (Eds.) *Landform Dynamics and Evolution in Romania*. Dordrecht: Springer, pp. 87–112.
- Quincey, D.J. & Glasser, N.F. (2009) Morphological and ice-dynamical changes on the Tasman Glacier, New Zealand, 1990–2007. *Global and Planetary Change*, 68, 185–197. Available from: <https://doi.org/10.1016/j.gloplacha.2009.05.003>
- Rasmussen, S.O., Bigler, M., Blockley, S.P., Blunier, T., Buchardt, S.L., Clausen, H.B. et al. (2014) A stratigraphic framework for abrupt climatic changes during the Last Glacial period based on three synchronized Greenland ice-core records: refining and extending the INTIMATE event stratigraphy. *Quaternary Science Reviews*, 106, 14–28. Available from: <https://doi.org/10.1016/j.quascirev.2014.09.007>
- Rea, B.R. (2013) Glacial landforms, erosional features: micro-to macroscale forms. In: Elias, S.A. (Ed.) *Encyclopedia of Quaternary Science*. Amsterdam: Elsevier, pp. 865–876. Available from: <https://doi.org/10.1016/B978-0-444-53643-3.00076-5>
- Rea, B.R. & Evans, D.J.A. (2003) Plateau icefield landsystems. In: Evans, D.-J.A. (Ed.) *Glacial landsystems*. London: Arnold, pp. 407–431.
- Rea, B.R. & Evans, D.J.A. (2007) Quantifying climate and glacier mass balance in North Norway during the Younger Dryas. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 246, 307–330. Available from: <https://doi.org/10.1016/j.palaeo.2006.10.010>
- Rea, B.R., Whalley, W.B., Dixon, T.S. & Gordon, J.E. (1999) Plateau icefields as contributing areas to valley glaciers and the potential impact on reconstructed ELAs: a case study from the Lyngen Alps, North Norway. *Annals of Glaciology*, 28, 97–102. Available from: <https://doi.org/10.3189/172756499781822020>
- Rea, B.R., Whalley, W.B., Evans, D.J.A., Gordon, J.E. & McDougall, D.A. (1998) Plateau icefields: geomorphology and dynamics. *Quaternary Proceedings*, 6, 35–54.
- Reuther, A., Urdea, P., Geiger, C., Ivy-Ochs, S., Niller, H.P., Kubik, P. et al. (2007) Late Pleistocene glacial chronology of the Pietrele Valley, Retezat Mountains, Southern Carpathians, constrained by 10Be exposure ages and pedological investigations. *Quaternary International*, 164–165, 151–159. Available from: <https://doi.org/10.1016/j.quaint.2006.10.011>
- Roberson, S. (2008) Structural composition and sediment transfer in a composite cirque glacier: glacier de St. Sorlin, France. *Earth Surface Processes and Landforms*, 33, 1931–1947. Available from: <https://doi.org/10.1002/esp.1635>
- Rogerson, R.J., Olson, M.E. & Branson, D. (1986) Medial moraines and surface melt on glaciers of the Torngat Mountains, northern Labrador, Canada. *Journal of Glaciology*, 32, 350–354. Available from: <https://doi.org/10.3189/S0022143000012028>
- Rootes, C.M. & Clark, C.D. (2022) On the expression and distribution of glacial trimlines: a case study of Little Ice Age trimlines on Svalbard. *E&G Quaternary Science Journal*, 71(1), 111–122. Available from: <https://doi.org/10.5194/egqsj-71-111-2022>
- Ruszkiczay-Rüdiger, Z., Kern, Z., Urdea, P., Braucher, R., Madarász, B., Schimmelpfennig, I. et al. (2016) Revised deglaciation history of the Pietrele–Stânișoara glacial complex, Retezat Mts, Southern Carpathians, Romania. *Quaternary International*, 415, 216–229. Available from: <https://doi.org/10.1016/j.quaint.2015.10.085>
- Ruszkiczay-Rüdiger, Z., Kern, Z., Urdea, P., Madarász, B., Braucher, R. & ASTER Team (2021) Limited glacial erosion during the last glaciation in mid-latitude cirques (Retezat Mts, Southern Carpathians, Romania). *Geomorphology*, 384, 107719. Available from: <https://doi.org/10.1016/j.geomorph.2021.107719>
- Sanders, C., Huisman, R., Van Wees, J.D. & Andriessen, P. (2002) The Neogene history of the Transylvanian basin in relation to its surrounding mountains. *EGU Stephan Mueller Special Publication Series, European Geoscience Union*, 3, 121–133. Available from: <https://doi.org/10.5194/smsps-3-121-2002>
- Sanders, C.A., Andriessen, P.A.M. & Cloetingh, S.A.P.L. (1999) Life cycle of the East Carpathian orogen: erosion history of a doubly vergent critical wedge assessed by fission track thermochronology. *Journal of Geophysical Research - Solid Earth*, 104(B12), 29095–29112. Available from: <https://doi.org/10.1029/1998JB900046>
- Schafarzik, F. (1899). Die geologische Verhältnisse der Umgebung von Borlova und Pojana Mörl, *Jahrbuch der Ungar. Geol. Landesanstalt*, f. 1897, 120–156.
- Șerban, R.D., Onaca, A., Șerban, M. & Urdea, P. (2019) Block stream characteristics in Southern Carpathians (Romania). *Catena*, 178, 20–31. Available from: <https://doi.org/10.1016/j.catena.2019.03.003>
- Shakesby, R.A. (1997) Pronival (proatalus) ramparts: a review of forms, processes, diagnostic criteria and palaeoenvironmental implications. *Progress in Physical Geography*, 21(3), 394–418. Available from: <https://doi.org/10.1177/030913339702100304>
- Shakesby, R.A. & Matthews, J.A. (1993) Loch Lomond Stadial glacier at Fan Hir, Mynydd Du (Brecon Beacons), south Wales: critical evidence and palaeoclimatic implications. *Geological Journal*, 28(1), 69–79. Available from: <https://doi.org/10.1002/gj.3350280106>
- Shakesby, R.A. & Matthews, J.A. (1996) Glacial activity and paraglacial landsliding in the Devensian Lateglacial: evidence from Craig Cerrig-gleisiad and Fan Dringarth, Fforest Fawr (Brecon Beacons), South Wales. *Geological Journal*, 31, 143–157. Available from: [https://doi.org/10.1002/\(SICI\)1099-1034\(199606\)31:2<143::AID-GJ704>3.0.CO;2-K](https://doi.org/10.1002/(SICI)1099-1034(199606)31:2<143::AID-GJ704>3.0.CO;2-K)
- Shakesby, R.A., Matthews, J.A., McEwan, L. & Berrisford, M.S. (1999) Snow-push processes in pronival (proatalus) rampart formation: geomorphological evidence from Smørbotn, Romsdalsalpane southern Norway. *Geografiska Annaler*, 81A, 31–45. Available from: <https://doi.org/10.1111/j.0435-3676.1999.00047.x>
- Sîrcu, I. (1978) *Munții Rodnei. Studiu morfogeografic*. București: Edit. Academiei, p. 112.
- Small, R.J. (1983) Lateral moraines of glacier de Tsidjiore Nouve: form, development, and implications. *Journal of Glaciology*, 29, 250–259. Available from: <https://doi.org/10.3189/S0022143000008303>
- Small, R.J., Clark, M.J. & Cawse, T.J.P. (1979) The formation of medial moraines on Alpine glaciers. *Journal of Glaciology*, 22(86), 43–52. Available from: <https://doi.org/10.3189/S0022143000014040>
- Syverson, K.M. & Mickelson, D.M. (2009) Origin and significance of lateral meltwater channels formed along a temperate glacier margin, Glacier Bay, Alaska. *Boreas*, 38(1), 132–145. Available from: <https://doi.org/10.1111/j.1502-3885.2008.00042.x>
- Thompson, H.R. (1957) The old moraines of Pangnirtung Pass, Baffin Island. *Journal of Glaciology*, 3(21), 42–49. Available from: <https://doi.org/10.1017/S0022143000024722>
- Tietze, K. (1878) Über das Vorkommen der Eiszeitspuren in den Ostkarpathen. *Verhandlungen der geologische Reichsanstalt, Wien*, 142–146.
- Tîrlă, L., Drăgușin, V., Bajo, P., Covaliov, S., Cruțeru, N., Ersek, V. et al. (2020) Quaternary environmental evolution in the South Carpathians reconstructed from glaciokarst geomorphology and sedimentary archives. *Geomorphology*, 354, 107038. Available from: <https://doi.org/10.1016/j.geomorph.2020.107038>
- Turnbull, J.M. & Davies, T.R. (2006) A mass movement origin for cirques. *Earth Surface Processes and Landforms*, 31(9), 1129–1148. Available from: <https://doi.org/10.1002/esp.1324>

- Urdea, P. (1992) Rock glaciers and periglacial phenomena in the Southern Carpathians. *Permafrost and Periglacial Processes*, 3(3), 267–273. Available from: <https://doi.org/10.1002/ppp.3430030317>
- Urdea, P., Ardelean, F., Ardelean, M. & Onaca, A. (2022a) Chapter 14-Glacial landscapes of the Romanian Carpathians. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 109–114. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00031-5>
- Urdea, P., Ardelean, F., Ardelean, M. & Onaca, A. (2022b) Chapter 38 - The Romanian Carpathians: glacial landforms prior to the Last Glacial Maximum. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 277–282. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00024-8>
- Urdea, P., Ardelean, F., Ardelean, M. & Onaca, A. (2022c) Chapter 57 - The Romanian Carpathians: glacial landforms from the Last Glacial Maximum (29–19 ka). In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 441–447. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00013-3>
- Urdea, P., Ardelean, F., Ardelean, M. & Onaca, A. (2022d) Chapter 19 - The Romanian Carpathians: glacial landforms during deglaciation (18.9–14.6 ka). In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: The Last Deglaciation*. Amsterdam: Elsevier, pp. 165–173. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00036-X>
- Urdea, P., Ardelean, F., Ardelean, M. & Onaca, A. (2022e) Chapter 36 - The Romanian Carpathians: glacial landforms during Bølling–Allerød Interstadial (14.6–12.9 ka). In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: The Last Deglaciation*. Amsterdam: Elsevier, pp. 347–353. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00010-3>
- Urdea, P., Ardelean, F., Ardelean, M., Onaca, A. & Berzescu, O. (2022) Chapter 54 - The Romanian Carpathians: glacial landforms from the Younger Dryas. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: The Last Deglaciation*. Amsterdam: Elsevier, pp. 517–524. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00059-0>
- Urdea, P., Onaca, A., Ardelean, F. & Ardelean, M. (2011) Chapter 24 - New evidence on the Quaternary glaciation in the Romanian Carpathians. In: Ehlers, J., Gibbard, P.L. & Hughes, P.D. (eds.) *Quaternary Glaciations—Extent and Chronology, A Closer Look*. Developments in Quaternary Science, Vol. 15. Amsterdam: Elsevier, pp. 305–322. Available from: <https://doi.org/10.1016/B978-0-444-53447-7.00024-6>
- Urdea, P. & Reuther, A. (2009) Some new data concerning the Quaternary glaciation in the Romanian Carpathians. *Geographia Pannonica*, 13(2), 41–52. Available from: <https://doi.org/10.5937/GeoPan0902041U>
- Vasile, M., Vespremeanu-Stroe, A., Pascal, D., Braucher, R., Pleşoiu, A., Popescu, R. et al. (2022) Rock walls distribution and Holocene evolution in a mid-latitude mountain range (the Romanian Carpathians). *Geomorphology*, 413, 108351. Available from: <https://doi.org/10.1016/j.geomorph.2022.108351>
- Vespremeanu-Stroe, A., Urdea, P., Popescu, R. & Vasile, M. (2012) Rock glacier activity in the Retezat mountains, Southern Carpathians, Romania. *Permafrost and Periglacial Processes*, 23(2), 127–137. Available from: <https://doi.org/10.1002/ppp.1736>
- Walker, M.J.C., Berkelhammer, M., Björck, S., Cwynar, L.C., Fisher, D.A., Long, A.J. et al. (2012) Formal subdivision of the Holocene series/epoch: a discussion paper by a working group of INTIMATE (Integration of ice-core, marine and terrestrial records) and the Subcommission on Quaternary Stratigraphy (International Commission on Stratigraphy). *Journal of Quaternary Science*, 27(7), 649–659. Available from: <https://doi.org/10.1002/jqs.2565>
- Whalley, W.B. (2009) On the interpretation of discrete debris accumulations associated with glaciers with special reference to the British Isles. *Geological Society, London, Special Publications*, 320(1), 85–102. Available from: <https://doi.org/10.1144/SP320.7>
- Whalley, W.B. (2012) Using discrete debris accumulations to help interpret upland glaciation of the Younger Dryas in the British Isles. In: Piacentini, T. & Miccadei, E. (Eds.) *Chapter 1, studies on environmental and applied geomorphology*. Rijeka: InTech Europe.
- Whalley, W.B. (2015) Discrete debris accumulations in the Lake District and their role in the interpretation of past processes and climate. In: McDougall, D.A. & Evans, D.J.A. (Eds.) *The Quaternary of the Lake District-Field Guide*. London: Quaternary Research Association, pp. 97–107.
- Whalley, W.B., Gordon, J.E., Gellatly, A.F. & Hansom, J.D. (1996) Plateau and valley glaciers in north Norway: responses to climate over the last 100 years. *Zeitschrift für Gletscherkunde und Glazialgeologie*, 31, 115–124.
- Wilson, P. (1990) Morphology, sedimentological characteristics and origin of a fossil rock glacier on Muckish Mountain, northwest Ireland. *Geografiska Annaler. Series A, Physical Geography*, 72(3–4), 237–247. Available from: <https://doi.org/10.1080/04353676.1990.11880319>
- Xu, X., Kleidon, A., Miller, L.E.E., Wang, S., Wang, L. & Dong, G. (2010) Late Quaternary glaciation in the Tianshan and implications for palaeoclimatic change: a review. *Boreas*, 39(2), 215–232. Available from: <https://doi.org/10.1111/j.1502-3885.2009.00118.x>
- Zasadni, J., Kłapyta, P., Kałuża, P. & Makos, M. (2022) Chapter 37 - The Tatra Mountains: glacial landforms prior to the Last Glacial Maximum. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M., & de Andrés, N. (Eds.) *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 271–275. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00059-5>
- Zasadni, J., Kłapyta, P. & Makos, M. (2022a) Chapter 56 - The Tatra Mountains: glacial landforms from the Last Glacial Maximum. In: *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 435–440. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00049-2>
- Zasadni, J., Kłapyta, P. & Makos, M. (2022b) Chapter 18 - The evolution of glacial landforms in the Tatra Mountains during the deglaciation. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: The Last Deglaciation*. Amsterdam: Elsevier, pp. 157–164. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00042-5>
- Zasadni, J., Kłapyta, P. & Makos, M. (2022c) Chapter 35 - The evolution of glacial landforms in the Tatra Mountains during the Bølling–Allerød Interstadial. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes*. Amsterdam: Elsevier, pp. 341–346. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00035-8>
- Zasadni, J., Kłapyta, P., Totoczko-Pasek, A. & Makos, M. (2022) Chapter 53 - The evolution of glacial landforms in the Tatra Mountains during the Younger Dryas. In: Palacios, D., Hughes, P.D., Garcia-Ruiz, J.M. & de Andrés, N. (Eds.) *European Glacial Landscapes: The Last Deglaciation*. Amsterdam: Elsevier, pp. 509–515. Available from: <https://doi.org/10.1016/B978-0-323-91899-2.00009-7>
- Zasadni, J., Makos, M. & Kłapyta, P. (2022) Chapter 13 - Glacial landscapes of the Tatra Mountains. In: *European Glacial Landscapes: Maximum Extent of Glaciations*. Amsterdam: Elsevier, pp. 103–107. Available from: <https://doi.org/10.1016/B978-0-12-823498-3.00046-7>

## SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

**How to cite this article:** Balaban, C.I., Roberts, D.H., Evans, D. J.A. & Jamieson, S.S.R. (2024) Past glaciation of temperate-continental mountains: A model for a debris-charged plateau icefield/cirque glacier landsystem in the Southern Carpathians, Romania. *Earth Surface Processes and Landforms*, 49(2), 601–621. Available from: <https://doi.org/10.1002/esp.5723>