

ECOGRAPHY

Research article

Invasion risk of the currently cultivated alien flora in southern Africa is predicted to decline under climate change

Ali Omer^{1,2,3,4}, Franz Essl¹, Stefan Dullinger⁴, Bernd Lenzner¹, Adrián García-Rodríguez¹, Dietmar Moser⁴, Trevor Fristoe^{2,5}, Wayne Dawson^{6,7}, Patrick Weigelt^{8,9,10}, Holger Kreft^{8,9,10}, Jan Pergl¹¹, Petr Pyšek^{11,12}, Mark van Kleunen^{2,13,*} and Johannes Wessely^{4,*}

¹Division of BioInvasions, Global Change & Macroecology, Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria

²Ecology, Department of Biology, University of Konstanz, Konstanz, Germany

³Department of Forest Management, Faculty of Forestry, University of Khartoum, North Khartoum, Sudan

⁴Division of Biodiversity Dynamics & Conservation, Department of Botany and Biodiversity Research, University of Vienna, Vienna, Austria

⁵Department of Biology, University of Puerto Rico - Río Piedras, San Juan, Puerto Rico

⁶Department of Biosciences, Durham University, Durham, UK

⁷Department of Evolution, Ecology and Behaviour, Institute of Infection, Veterinary and Ecological Sciences, University of Liverpool, Liverpool, UK

⁸Biodiversity, Macroecology & Biogeography, University of Goettingen, Göttingen, Germany

⁹Centre of Biodiversity and Sustainable Land Use (CBL), University of Goettingen, Göttingen, Germany

¹⁰Campus Institute Data Science (CIDAS), University of Goettingen, Göttingen, Germany

¹¹Czech Academy of Sciences, Institute of Botany, Department of Invasion Ecology, Průhonice, Czech Republic

¹²Department of Ecology, Faculty of Science, Charles University, Prague, Czech Republic

¹³Zhejiang Provincial Key Laboratory of Plant Evolutionary Ecology and Conservation, Taizhou University, Taizhou, China

Correspondence: Ali Omer (ali.haroon.ali.omer@univie.ac.at)

Ecography

2024: e07010

doi: [10.1111/ecog.07010](https://doi.org/10.1111/ecog.07010)

Subject Editor: Damien Fordham

Editor-in-Chief:

Christine N. Meynard

Accepted 07 February 2024



www.ecography.org

Alien species can have massive impacts on native biodiversity, ecosystem functioning, and human livelihoods. Assessing which species from currently cultivated alien floras may escape into the wild and naturalize is essential for efficient and proactive ecosystem management and biodiversity conservation. Climate change has already promoted the naturalization of many alien plants in temperate regions, but whether it is similar in (sub)tropical areas is insufficiently known. In this study, we used species distribution models for 1527 cultivated alien plants to evaluate current and future invasion risks across different biomes and 10 countries in southern Africa. Our results confirm that the area of suitable climate is a strong predictor of naturalization success among the cultivated alien flora. In contrast to previous findings from temperate regions, however, climatic suitability is generally predicted to decrease for potential aliens across our (sub)tropical study region. While increasingly hotter and drier conditions are likely to drive declines in suitability for potential aliens across most biomes of southern Africa, in some the number of potential invaders is predicted to increase under moderate climate change scenarios (e.g. in dry broadleaf forests and flooded grasslands). We found that climatic suitability is expected to decline less for aliens originating from continents with the tropical biome or from the Southern Hemisphere. In addition, we found that the climatically suitable area will decline less for aliens that have already naturalized in the region. While the number of potential invaders may decrease across

© 2024 The Authors. Ecography published by John Wiley & Sons Ltd on behalf of Nordic Society Oikos

*Shared last authorship.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

southern Africa under future climate change, our results suggest that already naturalized aliens will continue to threaten native species and ecosystems.

Keywords: biological invasion, climate change, habitat suitability, invasion risk, naturalization success, ornamental plants, species distribution models

Introduction

The number of species that are naturalizing outside of their native ranges continues to increase (Seebens et al. 2020). The associated ecological and economic costs make alien species management an urgent task (Pyšek et al. 2020). Estimations and projections of current and future distributions are important for alien species management, especially as their spread will certainly be affected by other drivers of global change, such as climate and land-use change (Bellard et al. 2015, Vilà and Hulme 2017, Northrup et al. 2019, Liu et al. 2023). These changes alter both biotic and abiotic ecosystem properties known to be critical for biological invasions (Rodríguez-Labajos et al. 2009, Bellard et al. 2016, Dullinger et al. 2017).

Predicting the species that could successfully escape into the wild and naturalize from a larger pool of deliberately introduced species (e.g. those cultivated in a region) is one of the biggest challenges in invasion ecology. Apart from specific functional traits and evolutionary history (e.g. seed mass, geographical origins, and phylogenetic composition; Maurel et al. 2016, Divíšek et al. 2018, Lenzner et al. 2021, Omer et al. 2021, 2022), climate matching between native and alien ranges has been demonstrated to be fundamental for the naturalization success of alien plants (Richardson and Pyšek 2012, Feng et al. 2016, Mayer et al. 2017, Fristoe et al. 2023). The link between environmental suitability and the ability of alien plants to naturalize has long been established (Darwin 1859, Elton 1958). However, with continuing climate change, invasion dynamics have become more complex to predict. For example, while current climate might be favorable for alien species that have already naturalized in a region, future climates may promote the establishment of alien species that have been introduced already but have not yet naturalized. A warming climate might hence constrain the spread of some naturalized alien species but simultaneously might foster expansions or new naturalizations of others.

Climate-suitability analyses have emerged as promising tools for predicting the naturalization risk of alien plants (Thuiller et al. 2005, Dullinger et al. 2017, Haeuser et al. 2018, van Kleunen et al. 2018, Pouteau et al. 2021, Oduor et al. 2023); with projections based on future climate scenarios becoming increasingly important for predicting the potential distributions of alien species. In temperate regions as found in Europe and northern America, climate warming is predicted to generally increase the likelihood of biological invasions (Bellard et al. 2013, Dullinger et al. 2017, Haeuser et al. 2018, Oduor et al. 2023). However, biological invasion studies, including those that use climate-suitability analyses, are geographically biased towards regions in the Northern Hemisphere (Pyšek et al. 2008). Most of

such studies have been conducted in intensively researched regions (e.g. Europe; Dullinger et al. 2017, Haeuser et al. 2018, Pouteau et al. 2021) or for a specific set of species (e.g. the 100 world's worst invasive species; Bellard et al. 2013). These biases in research might hinder a general understanding of how a changing climate will affect biological invasions across the globe. Indeed, some studies suggest that outside of temperate regions, and especially in areas with extreme climates (e.g. hot desert), the risk of alien plant naturalization will decrease rather than increase under climate change (Bellard et al. 2013, Fulgêncio-Lima et al. 2021).

Southern Africa has a tropical and subtropical climate, including large areas of (semi-)arid habitats (Engelbrecht and Engelbrecht 2016). It is not immediately clear how climate change will affect the potential future distribution of alien species across this diverse region. Here, we evaluate current and future invasion risks across biomes of southern Africa using species distribution modeling to investigate the climate suitability for 1527 alien species currently cultivated in the 10 countries comprising the region. Cultivation in domestic gardens for economic use represents the primary pathway for vascular plant introductions (Lambdon et al. 2008, Faulkner et al. 2020, van Kleunen et al. 2020), and the high prevalence of cultivated species among naturalized floras (van Kleunen et al. 2018) implies that future naturalizations will likely also emerge mainly from cultivated plant populations. Our specific objectives were 1) to predict the current potential distributions of cultivated alien plants of southern Africa; 2) to assess how these potential distributions could change under a changing climate in the future by using two climate change scenarios; 3) to assess whether naturalization status explains current and future potential range size of cultivated alien plants of southern Africa; 4) to compare the biomes within southern Africa and native origins of the species with respect to changes in climatic suitability for introduced cultivated plants; and 5) to identify hotspot areas with the highest suitability for cultivated alien plants under current and future climatic conditions in southern Africa.

Methods

Study area

Our study focused on the southern Africa region, comprising ten countries: Angola, Botswana, Eswatini, Lesotho, Malawi, Mozambique, Namibia, Republic of South Africa, Zambia, and Zimbabwe, with a land area of around 4 000 000 km² (Bezeng et al. 2015). The history of modern plant introductions in southern Africa dates back to the late 18th century

when European settlers arrived in the region (Wells et al. 1986). The latest global IPCC report (IPCC 2023) shows that southern Africa is likely to become substantially hotter, while precipitation is likely to decrease in most regions. With the predicted changes in temperature, precipitation regimes, and water availability, southern Africa is expected to become one of the global climate-change hotspots (Hoegh-Guldberg et al. 2019).

Species selection and occurrence records

For our study, we used a list of cultivated alien plants of southern Africa extracted from Glen (2002). The initial list included more than 5316 taxa that are described to be cultivated in at least one region in southern Africa. To harmonize the list of cultivated alien plants of southern Africa with other datasets used in this study (below), we standardized the names of the species following The Plant List (ver. 1.1; www.theplantlist.org) using the R package ‘Taxonstand’ (Cayuela et al. 2019). Intraspecific taxa (varieties and subspecies) were merged at the species level to reduce complexity. The resulting list, therefore, consisted of 5212 cultivated alien plants with accepted names.

We collected occurrence data on the global distribution of these species from the Global Biodiversity Information Facility (GBIF) (GBIF.org 2021; <https://doi.org/10.15468/dl.9jsscb>) using the ‘rgbif’ library in R (Chamberlain et al. 2021). To account for the full realized niche of the species, we considered native and introduced occurrences globally (Pearman et al. 2008, Early and Sax 2014, Fernández and Hamilton 2015). Erroneous records (e.g. those that occur on ocean surfaces due to possible georeferencing errors and those in capitals, where they might have been planted) were automatically removed using the ‘CoordinateCleaner’ library in R (Zizka et al. 2019). Additionally, we removed duplicate data points (that is, multiple occurrence records within each 10' × 10' grid cell, ~ 20 × 20 km) for bias correction. The resulting species list, therefore, consisted of 1527 species with at least 50 occurrences per species combined from their native and alien ranges.

Climatic data

We retrieved global climate data from WorldClim ver. 2.1 (10' resolution for the period 1970–2000) (Fick and Hijmans 2017). From all the available bioclimatic variables, we selected the following five: 1) temperature seasonality (SD × 100), 2) maximum temperature of warmest month (°C), (3) precipitation of wettest month (°C), 4) precipitation of driest month (mm), and 5) precipitation seasonality (coefficient of variation). We selected these variables because they are known to strongly affect plant distributions (Root et al. 2003). In addition, we used human population density (person/10' × 10' grid cell), available from the NASA Socioeconomic Data and Applications Center, as an interaction term with nativeness as an indicator of propagule pressure (Gao 2020). Moreover, all explanatory variables have pairwise Pearson's *r* values < 0.7

(Supporting information), limiting the risk of biased model estimates due to multicollinearity (Dormann et al. 2013).

To represent possible future climatic conditions, we used projected climate data for the period 2081–2100 (means of the above-listed climatic variables), again retrieved from WorldClim ver. 2.1 (Fick and Hijmans 2017). We also used human population density projection for the year 2100, retrieved from the NASA Socioeconomic Data and Applications Center. We used two shared socioeconomic pathways (SSPs) to characterize future climate conditions, specifically SSP_{1-2.6} and SSP_{5-8.5}, to represent a best-case scenario (the sustainability/taking the green road scenario) and a worst-case scenario (fossil-fueled development/taking the highway scenario), respectively (O'Neill et al. 2017, Riahi et al. 2017). Because different global circulation models (GCMs) significantly affect species range projections, we selected three GCMs for each SSP scenario, namely CanESM5, CNRM-ESM2-1, and MIROC6. According to The Inter-Sectoral Impact Model Intercomparison Project (Lange 2019), these GCMs represent relatively low, moderate, and high global projected mean precipitation and temperature.

Data on naturalization status, native origins, life forms, and biomes

To analyze whether the potential current and future climatic suitability differ according to plants' naturalization status, biogeographical origin, and biome within southern Africa, we first extracted the naturalization status of each species (that is, cultivated but not yet naturalized; or cultivated and naturalized) using the latest version of the Global Naturalized Alien Flora (GloNAF) database (van Kleunen et al. 2019). Second, we used the nine level-1 regions of the World Geographical Scheme for Recording Plant Distributions of the Taxonomic Databases Working Group (TDWG; Brummitt 2001) to identify the native geographical origin of each species. These data were extracted from the Germplasm Resources Information Network (GRIN; <https://ars-grin.gov>), the World Checklist of Selected Plant Families (WCSP; <http://apps.kew.org/wcsp>), and the Plants of the World Online database (POWO 2019); <http://www.plantsoftheworldonline.org/>). Moreover, we assigned each species to one or more major life forms using data on species life forms that we compiled from different data sources (Omer et al. 2021). Finally, we assigned each 10' × 10' grid cell in southern Africa to one of the biomes defined by Dinerstein et al. (2017) (Supporting information).

Species distribution modeling

To define the current and future potential climatic suitability for the cultivated alien plants in southern Africa, we combined the bioclimatic variables with presence records and randomly generated pseudo-absence data using the biomod2 platform as implemented in the ‘biomod2’ R package ver. 3.4–6 (Thuiller et al. 2020). We used four modeling algorithms: 1) two regression techniques (that is, i) generalized

linear model (GLM) and 2) general additive model (GAM)) and two classification techniques: 3) random forest (RF) and 4) boosted regression trees (BRT). We kept the default argument settings of these four modeling algorithms in `biomod2`. Since all models require presences and pseudo-absences (or background), we randomly generated 10 000 pseudo-absence records from all over the calibration area (i.e. in our case, the globe terrestrial surface). The random draw of pseudo-absence records was repeated three times, and equal weights were given in the models to presences and pseudo-absences. Finally, to evaluate our models, each model was separately run three times using a random split-sampling approach in which data were split into 80% calibration and 20% evaluation datasets for each of the three pseudo-absence datasets (resulting in nine models per modeling algorithm and a total of 36 models for each species). We used the true skill statistic (TSS) of [Allouche et al. \(2006\)](#) to assess the predictive performance of the SDMs. TSS values range from -1 to 1 , where 0 indicates a random prediction, negative values indicate that predictions perform worse than random, and 1 indicates perfect agreement.

We then used the calibrated models to project the current and future climatic suitability in southern Africa using a weighted mean ensemble forecast ([Thuiller et al. 2009](#)). To do that, we first aggregated all the models of the repeated pseudo-absences and split-sampling into an ensemble projection to reduce uncertainties associated with each technique. The contribution of each model was weighted according to its TSS score (we only included models with a TSS score > 0.5). Then, the mean weighted ensemble was transformed into binary maps using a threshold that maximized the TSS to predict presences and absences for the ‘current’ climate and each of the two climate change scenarios. Three binary projections were produced for each SSP scenario, one for each GCM. We then combined these three projections into one consensus map where each cell was identified as suitable when the majority of GCMs (that is, two of three) predicted it as suitable; otherwise, the cell was identified as unsuitable.

This subset of modeled species ($n = 1527$) effectively represents the cultivated flora of southern Africa. This conclusion is supported by the similar distribution of native origins between the modeled and unmodeled species ($n = 3685$) (Supporting information). With the exception of species native to Europe and southern America, the proportion of species from each native origin was quantitatively consistent in the modeled species subset when compared to the entire pool of the cultivated flora.

We explored the potential impact of each bioclimatic variable on the future distribution of the cultivated alien species. To do so, we made predictions for future climatic conditions fixing one of the five predictors at its value of the reference period 1970–2000 in turn. We then compared these predictions to those of the fully adapted model (i.e. all predictors set to future conditions) by computing the difference in the number of suitable cells. The rationale is that a predictor variable has more impact the more the two projected future distributions differ (Supporting information).

Invasion impact considers not only the count of naturalized species but also their ratio to native ones. A region with 10 naturalized species among 100 natives may seem more invaded than one with 10 naturalized species among 1000, but the latter could face stronger negative effects, especially in regions rich in native and endemic species. Therefore, we also explored the relative distribution patterns of cultivated alien plants compared to the richness of native species in southern Africa. To do this, we extracted the projected numbers of native species in each grid cell in southern Africa ([Cai et al. 2023](#)), and used these numbers to standardize the richness of cultivated alien plants in each specific grid cell (Supporting information).

Climatic suitability and naturalization status

Under both current and future climatic scenarios, 824 (which accounts for 53.9% of all modeled species) were predicted to lack suitable climatic conditions in southern Africa. Therefore, we will not include these species in our subsequent analysis. Species with modeled unsuitable climatic suitability may still be successfully cultivated in gardens and public spaces, as under cultivation, weeding, irrigation and tendering may allow them to persist and flourish.

We tested whether the naturalization status of cultivated alien plants in southern Africa is correlated with the size of the current potential range (= number of suitable $10' \times 10'$ cells) and whether cultivated alien and naturalized plants differ in their response to climate change (change in the future range size compared to the current range size). We first calculated the difference in potential range sizes between current and future climate scenarios. Then we divided that difference by the potential range size under current climatic conditions (proportion of change). Negative and positive values indicate a net reduction or expansion, respectively, in the climatically suitable area under climate change. Then, we fitted three generalized linear models (GLMs) with a binomial error distribution and a logit link function. For each GLM model, we set naturalization status as the binary response variable and the number of climatically suitable cells under the three climatic scenarios (that is, current and change in $SSP_{1-2.6}$ and $SSP_{5-8.5}$ to the current climate) as explanatory variables. To facilitate comparisons of the estimates within and between the models, we also scaled each explanatory variable to a mean of zero standard deviation of one ([Schielzeth 2010](#)).

Climatic suitability and native origins and life forms

We assessed whether current potential range sizes would, on average, increase, decrease, or remain constant under future climate change scenarios for species of different geographical origins and life forms. First, we calculated the mean proportion of change for each group and then calculated 95% confidence intervals around these means with 1000 bootstrap replications using the `boot.ci` function in the ‘boot’ R package ver. 1.3–28 ([Canty and Ripley 2021](#)). We considered the mean proportion of change to deviate from zero if the

confidence intervals did not overlap with zero. We also tested if there are differences in the projected range-size change among geographical origins and life forms. Again, to account for the fact that each species can be native to multiple continents and belong to multiple life form, we applied simple randomizations to determine whether the mean projected range size change of each geographical origin and life form deviated from those expected by chance ($p < 0.05$, two-tailed test; see Divíšek et al. 2018, Omer et al. 2021). Therefore, from the pool of all proportions of changes, we randomly drew 999 times as many species as are in each geographical origin and life form. We defined the observed mean proportions of change to differ from random expectations if it was in or beyond the lower 2.5% or upper 2.5% of the distributions of random draws.

Climatic suitability and biomes

We assessed the difference between biomes within southern Africa with respect to changes in climatic suitability under climate change. To do this, we calculated for each grid cell in the different biomes of southern Africa the number of alien species that encounter suitable climatic conditions under current and future scenarios. Then, we calculated the difference in the number of alien species between current and future climate scenarios and divided it by the number of alien species under current climatic scenario (proportion of change). To test whether the potential number of alien species in each biome will, on average, increase, decrease, or remain constant under future climate change scenarios, we calculated the mean proportion of change for each group and then calculated the 95% confidence intervals around these means with 1000 bootstrap replications using the *boot.ci* function in the 'boot' R package ver. 1.3–28 (Canty and Ripley 2021). We considered the mean proportion of change of each group to deviate from zero if the confidence intervals did not overlap with zero. We also tested if there are differences in the numbers of potential alien species among biomes. To account for the fact that each species can potentially occur in multiple biomes, we applied simple randomizations to determine whether the mean potential number of alien species in each biome deviated from those expected by chance ($p < 0.05$, two-tailed test; see Divíšek et al. 2018, Omer et al. 2021). Therefore, from the pool of all proportions of changes, we randomly drew 999 times as many species as are in each biome. We defined the observed mean proportions of change to differ from random expectations if it was in or beyond the lower 2.5% or upper 2.5% of the distributions of random draws.

Hotspot analysis

To identify potential invasion hotspots for cultivated alien plants in southern Africa for each climatic scenario, we stacked the binary consensus maps of all 1527 modeled species. We then calculated, for each grid cell ($10' \times 10'$), the number of cultivated species that find suitable climatic

conditions there. We determined current invasion hotspots to be grid cells that were projected as suitable for at least as many cultivated plants as identified by the 90% percentile of grid cells under the current climate; this corresponds to grid cells that were projected to be suitable to 128 cultivated alien plants or more. To depict potential future contractions or expansions of invasion hotspots under warming scenarios, we identified the high-risk region under future climatic scenarios by applying cut-off value determined under current conditions (i.e. 128 alien species) to the future climatic scenarios.

Unmodeled species climatic suitability imputation

Due to the limited availability of the species' geographical distribution data, we could only predict the current and future species distribution for 1527, which represents just 29.2% of our entire pool of 5212 cultivated species. Although we showed that this subset of modeled species is representative of the entire pool of cultivated flora (Supporting information), we conducted further analysis to evaluate the impact of the unmodeled species on our conclusions. To address this, we used the result of the 1527 modeled species to impute the climatic suitability for the unmodeled species in our pool of cultivated flora of southern Africa. To do so, we fitted three separate linear models using the number of climatically suitable cells under the three climatic scenarios (that is, current and change in SSP_{1-2.6} and SSP_{5-8.5} to the current climate) as response variables. Naturalization status and geographical origins were used as explanatory variables. We then used the fitted values to predict the number of climatically suitable cells under the three climatic scenarios for species that were not included in the SDMs. Finally, we redid the analysis of how the change in climatic suitability is related to naturalization status and native origins using the pool of all species (including imputed species). The results using the entire pool of cultivated species show a more or less similar trend for the effects of naturalization status (Supporting information) and native origins (Supporting information).

All analyses were done in R ver. 3.6.1 (www.r-project.org).

Results

All calibrated models performed well with an average TSS value above 0.8 (Supporting information). The results reported below were consistent across all GCMs explored (Supporting information). Across the 1527 cultivated alien species, the number of projected suitable grid cells under current conditions varied from 0 to 9244 (approximately 51% of southern Africa's area). As expected, the current area of suitable climate in southern Africa was positively related to the probability that a species has naturalized within the region (GLM: $z=9.64$, $p \leq 0.001$; Fig. 1a, Supporting information). Under future climate scenarios, the area of suitable climate was predicted to decrease for most species (SSP_{1-2.6}: 72.8%; SSP_{5-8.5}: 85.6%). The strongest driver of suitability




Figure 1. Current climatic suitability (number of grid cells; (a) and predicted change to current climatic suitability under moderate (SSP_{1-2,6}), (b) and severe (SSP_{5-8,5}), (c) climate change by 2081–2100 of cultivated non-naturalized and naturalized plants of southern Africa. The grey dots represent the number of grid cells predicted to be suitable for each species under current conditions (a) and are predicted to be lost or gained in the future (b–c). The thick horizontal line in each box indicates the median number of cells predicted to be suitable under the current climate (a) and changed to suitable or not suitable under future climate scenarios. The boxes indicate the interquartile range, and the whiskers extend outside the box to 1.5 times the interquartile range. Asterisks indicate significant differences between the compared means according to the GLMs models, with *** indicating $p < 0.001$ and * indicating $p < 0.05$, and na indicating non-significant.

contractions was increasing Maximum Temperature of the Warmest Month across the region (Supporting information). Under the moderate future climate scenario SSP_{1-2,6}, these contractions are projected to be less severe for already naturalized species (GLM: $z = 2.15$, $p = 0.031$; Fig. 1b, Supporting information). Under the worst-case scenario SSP_{5-8,5}, stronger declines in suitable area are expected in general, with similar changes for naturalized and non-naturalized species (GLM: $z = 1.87$, $p = 0.060$; Fig. 1c, Supporting information). While the average cultivated plant in southern Africa will experience a reduction of its potential range, we note that increases are projected for ~ 26.0% and 13.5% of species under the scenarios SSP_{1-2,6} and SSP_{5-8,5}, respectively.

Across southern Africa, the number of species projected to encounter climatically suitable conditions under current climate varied geographically, ranging from 0 to 313 species per grid cell (approximately 20% of the modeled cultivated alien flora; Fig. 2a). Under climate warming scenarios (SSP_{1-2,6} and SSP_{5-8,5}), numbers of species per cell were generally projected to decrease (Fig. 2b–e). Changes in the number of potentially naturalized species varied across biomes, with patterns differing between climate change scenarios (Fig. 3). Notably, under moderate climate change (SSP_{1-2,6}), tropical and subtropical dry broadleaf forests and flooded grassland savannas are expected to become climatically suitable for a higher number of potential invaders (Fig. 3a).

The area of suitable climate, current and projected future, also varied depending on species native geographic origins and life form. Species native to continents spanning primarily equatorial latitudes or located in the Southern Hemisphere (i.e. Pacific Islands, Australasia, tropical Asia and southern America) generally had a higher current suitability and were projected to experience smaller contractions in suitable climate compared to those originating from Northern Hemisphere continents (Fig. 4). Under moderate climate change (SSP_{1-2,6}), species native to the Pacific Islands and Australasia were even predicted to experience increases in suitable area (Fig. 4a). Epiphyte and woody species were less likely to lose suitable area under both climate scenarios. While climber, aquatic, and long-lived herb plants did not deviate from random expectations, short-lived herbs were expected to lose suitable area more than expected by chance (Supporting information).

We defined the top 10.0% of grid cells in southern Africa that were climatically suitable for the highest number of cultivated alien species as invasion hotspots (i.e. a threshold of 128 species from the pool of 1527 modeled cultivated aliens; Fig. 5a). However, when accounting for native richness, proportion of cultivated plants was reduced along the coastal area where native richness is higher (Supporting information). Until the end of the century, the number of cells meeting this invasion hotspot criterion were predicted to decrease slightly under the SSP_{1-2,6} climatic scenario (to 7.1%; Fig. 5b)




Figure 2. Current and future potential richness of cultivated alien plants of southern Africa. The maps show the predicted number of species that are expected to encounter suitable climatic conditions per $10'$ grid cell under current climatic conditions (a), moderate future climate change (SSP_{1-2.6}) (b), and severe climate change (SSP_{5-8.5}) by the end of the 21st century (2081–2100) (c); and the expected change to current species richness under future climate change scenarios: SSP_{1-2.6} (d), and SSP_{5-8.5} (e).

but substantially under the worst-case climatic scenario SSP_{5-8.5} (to 2.0%; Fig. 5c). Under increasingly severe climate scenarios, invasion hotspots were restricted further towards southern coastal regions.

Discussion

Numerous regional studies have indicated that alien plants are projected to experience increased range sizes due to climate change, particularly in the Northern Hemisphere (Bellard et al. 2013, Dullinger et al. 2017, Thapa et al. 2018, Adhikari et al. 2022). In contrast, our results for the subtropical semi-arid region of southern Africa indicate that increasingly hotter and drier future conditions will result in reduced climatic suitability for the majority of already naturalized

aliens and non-naturalized species that may escape cultivation. Suitability within the region is projected to decline more for species originating from continents of the Northern Hemisphere, but the effect will be less pronounced for cultivated aliens introduced from primarily tropical or subtropical continents, with climatic favorability even increasing for some of these species. Cultivated aliens that have already naturalized populations in southern Africa are also expected to maintain larger areas of suitable climate relative to non-naturalized species, indicating that these species will continue to threaten native species and ecosystems.

Our results are in line with projections for the world's 100 worst invaders that suggest suitability will decline across many tropical and subtropical regions for these species (Bellard et al. 2013). Similarly, (Bezeng et al. 2017) projected that climatically suitable areas for the majority of alien trees




Figure 3. Predicted change in current climatic suitability under moderate future climate change (SSP_{1-2.6}) (a) and severe climate change (SSP_{5-8.5}) (b) of cultivated alien plants of southern Africa by 2081–2100 separated by their biomes. The dots represent the mean of the predicted change of a certain group. The lines are the 95% bootstrapped confidence intervals of the means of 1000 resamples from the population of species of a certain group. Red, blue, and black lines indicate whether the mean of the group is significantly larger, smaller, or not different from zero. The violin plots show the distribution of the means of predicted changes sampled by bootstrapping from the population of all species. Red, blue, and black dots indicate whether the means of the predicted change are significantly higher, lower, or not different from the random expectations, respectively.

and shrubs in the country of South Africa will contract under climate change. Our assessment of the cultivated alien flora of southern Africa also parallels projections predicting that the native flora of this region will experience range losses under climate change. Particularly, endemic plant species in southern Africa are predicted to lose approximately 50% of

their suitable ranges by 2050, even under optimistic climate change scenarios (Broennimann et al. 2006). We found that richness of alien cultivated plants is highest in areas with medium native richness, particularly within the Montane grasslands and Shrublands biome (Supporting information). This might be because this area presents milder climatic




Figure 4. Predicted change in current climatic suitability under moderate (SSP_{1-2.6}) (a) and severe climate change (SSP_{5-8.5}) (b) by 2081–2100 of cultivated alien plants of southern Africa separated by their native origins. The dots represent the mean of the predicted change of a certain group. The lines are the 95% bootstrapped confidence intervals of the means of 1000 resamples from the population of species of a certain group. Red, blue, and black lines indicate whether the mean of the group is significantly larger, smaller, or not different from zero. The violin plots show the distribution of the means of predicted changes sampled by bootstrapping from the population of all species. Red, blue, and black dots indicate whether the means of the predicted change are significantly higher, lower, or not different from the random expectations, respectively.




Figure 5. Current and future potential invasion hotspots of cultivated alien plants in southern Africa. The maps represent current climatic conditions (a), moderate future climate change (SSP_{1-2.6}) (b), and severe climate change (SSP_{5-8.5}) (c) by the end of the 21st century (2081–2100). We stacked the binary distribution maps of the 1527 species and then identified high-risk regions defined as the top 10% of cells that were predicted to be suitable under current climatic conditions for the highest number of species (depicted in red); the same cut off value was then used for climate change scenarios.

conditions compared to the harsh conditions in (semi-)arid biomes, coupled with lower competition from native plants in contrast to the Tropical biomes. This emphasizes the importance of considering interactions between biotic and abiotic drivers when assessing future invasion risks. Thus, further comparisons between native and non-native floras are required to better understand how plant communities across southern Africa will change as warming continues.

While we found that much of southern Africa will become less suitable for cultivated alien plants under future climate warming, the effects of climate change were not uniform across the different biomes and growth forms. We observed large variation among biomes; for example, climate change is predicted to cause fewer losses, or even gains, of potentially establishing alien species in some tropical biomes. In contrast, other biomes, such as semi-deserts, are expected to undergo significant contractions of potentially suitable area for many cultivated alien plants. Overall, we identified the southeastern region of southern Africa as the major invasion hotspot for the cultivated alien flora, currently and in the future (Fig. 5a–c). These patterns might be explained by higher predicted increases in temperature and aridity in the

western parts of southern Africa (such as the Namib desert), which already experience extreme climatic conditions today and generally have low suitability for most aliens currently cultivated in southern Africa (Almazroui et al. 2020). This is consistent with our results identifying Maximum temperature of the warmest month as the most influential bioclimatic variable in determining climatically suitable areas of alien cultivated plants (Supporting information). Similarly, temperature was found to be a major macroecological factor reducing diversity in native savanna flora of Kruger National Park, Republic of South Africa (Hejda et al. 2022). We also found that woody and epiphyte species are predicted to experience reduced losses of climatic suitability compared to other growth forms (Supporting information). Woody species are generally less sensitive to climate change compared to herbaceous species (Lin et al. 2010, Wang et al. 2020). Contrasting responses of woody and herbaceous species to climate change were also reported along dry lands in Africa (Verbruggen et al. 2021) and the New World (Šimová et al. 2018).

Overall, our study highlights that the potential distribution of the cultivated alien flora in southern Africa is unlikely to be amplified by future climate changes. In contrast,

climatically suitable ranges are projected to shrink, particularly under severe climate change. The reduction in climatic suitability for cultivated alien plants in southern Africa can be attributed to increasingly hot, semiarid climates that will be unfavorable to their growth. However, it is essential to note that, by the end of this century, the region is projected to experience novel climatic conditions, which could affect species distributions in unexpected ways (Williams et al. 2007). It is possible that current species distribution models (SDMs) do not appropriately account for how these cultivated plants will respond to such novel conditions, potentially leading to an overestimation of the effect of future climate on species distribution (Fitzpatrick and Hargrove 2009, Early and Sax 2014).

Consistent with previously identified correlations between climatic suitability and naturalization success (Feng et al. 2016, Mayer et al. 2017, Haeuser et al. 2018), naturalized plants currently have significantly larger climatically suitable areas in southern Africa than non-naturalized cultivated aliens. The species in our analyses that have not yet naturalized are expected to experience the most severe declines in suitability, indicating generally declining opportunities for future invasions from within southern Africa's cultivated flora. However, we note that between 15% and 8% of non-naturalized species (111 and 61 species; SSP_{1-2.6} and SSP_{5-8.5}, respectively) are projected to experience increased suitability under warming. Our results suggest that future invaders are most likely to originate from tropical or subtropical native regions and have epiphyte and woody life forms, with southern Africa's tropical biomes most at risk. Plants originating from temperate continents have historically been prominent among introduced and naturalized aliens in the region (Omer et al. 2021). However, with declines in suitability projected to be steepest for these species (see also Pouteau et al. 2021), the composition of southern Africa's naturalized flora is likely to change under warming. These changes, and threats from future invasions more generally, are likely to become more pronounced as cultivators and farmers will use new species that are better adapted to future conditions than species introduced in the past, such as the ones examined in this study.

Acknowledgements – We thank C. Gommel, K. Mamonova, V. Pasqualetto, and B. Rüter for help with data extraction, and Lesley Henderson for providing lists of naturalized species for South Africa. **Funding** – AO thanks FWF-DFG for funding (grant no. I-5825 – B). MvK thanks the German Research Foundation DFG for funding (grant no. 264740629 and no. 432253815). PP and JP were supported by EXPRO grant no. 19-28807X (Czech Science Foundation) and long-term research development project RVO 67985939 (Czech Academy of Sciences). FE, AG-R, and BL acknowledge funding from the Austrian Science Foundation FWF (grant no. I-5825-B).

Author contributions

Ali Omer: Conceptualization (lead); Methodology (lead); Visualization (lead); Writing – original draft (lead); Writing

– review and editing (lead). **Franz Essl:** Conceptualization (equal); Methodology (equal); Writing – original draft (equal); Writing – review and editing (equal). **Stefan Dullinger:** Conceptualization (equal); Methodology (equal); Writing – original draft (equal); Writing – review and editing (equal). **Bernd Lenzner:** Writing – original draft (supporting); Writing – review and editing (supporting). **Adrián García-Rodríguez:** Writing – original draft (supporting); Writing – review and editing (supporting). **Dietmar Moser:** Methodology (supporting); Writing – original draft (supporting); Writing – review and editing (supporting). **Trevor Fristoe:** Writing – original draft (supporting); Writing – review and editing (equal). **Wayne Dawson:** Writing – original draft (supporting); Writing – review and editing (supporting). **Patrick Weigelt:** Writing – original draft (supporting); Writing – review and editing (supporting). **Holger Kreft:** Writing – original draft (supporting); Writing – review and editing (supporting). **Jan Pergl:** Writing – original draft (supporting); Writing – review and editing (supporting). **Petr Pysek:** Writing – original draft (supporting); Writing – review and editing (supporting). **Mark van Kleunen:** Conceptualization (equal); Supervision (equal); Writing – original draft (supporting); Writing – review and editing (supporting). **Johannes Wessely:** Conceptualization (equal); Methodology (equal); Supervision (equal); Writing – original draft (equal); Writing – review and editing (equal).

Transparent peer review

The peer review history for this article is available at <https://publons.com/publon/10.1111/ecog.07010>.

Data availability statement

Data are available from the Dryad Digital Repository: <https://doi.org/10.5061/dryad.c59zw3rg2> (Omer et al. 2024).

Supporting information

The Supporting information associated with this article is available with the online version.

References

- Adhikari, P., Lee, Y. H., Adhikari, P., Hong, S. H. and Park, Y. 2022. Climate change-induced invasion risk of ecosystem disturbing alien plant species: an evaluation using species distribution modeling. – *Front. Ecol. Evol.* 10: 880987.
- Allouche, O., Tsoar, A. and Kadmon, R. 2006. Assessing the accuracy of species distribution models: prevalence, kappa and the true skill statistic (TSS). – *J. Appl. Ecol.* 43: 1223–1232.
- Almazroui, M., Saeed, F., Saeed, S., Nazrul Islam, M., Ismail, M., Klutse, N. A. B. and Siddiqui, M. H. 2020. Projected change in temperature and precipitation over Africa from CMIP6. – *Earth Syst. Environ.* 4: 455–475.
- Bellard, C., Thuiller, W., Leroy, B., Genovesi, P., Bakkenes, M. and Courchamp, F. 2013. Will climate change promote future invasions? – *Global Change Biol.* 19: 3740–3748.

- Bellard, C., Leclerc, C. and Courchamp, F. 2015. Combined impacts of global changes on biodiversity across the USA. – *Sci. Rep.* 5: 11828.
- Bellard, C., Leroy, B., Thuiller, W., Rysman, J.-F. and Courchamp, F. 2016. Major drivers of invasion risks throughout the world. – *Ecosphere* 7: e01241.
- Bezeng, B. S., Morales-Castilla, I., van der Bank, M., Yessoufou, K., Daru, B. H. and Davies, T. J. 2017. Climate change may reduce the spread of non-native species. – *Ecosphere* 8: e01694.
- Bezeng, S. B., Davies, J. T., Yessoufou, K., Maurin, O. and Van der Bank, M. 2015. Revisiting Darwin's naturalization conundrum: explaining invasion success of non-native trees and shrubs in southern Africa. – *J. Ecol.* 103: 871–879.
- Broennimann, O., Thuiller, W., Hughes, G., Midgley, G. F., Alkemade, J. M. R. and Guisan, A. 2006. Do geographic distribution, niche property and life form explain plants' vulnerability to global change? – *Global Change Biol.* 12: 1079–1093.
- Brummitt, R. K. 2001. World geographical scheme for recording plant distributions. – Hunt Institute for Botanical Documentation.
- Cai, L. R. et al. 2023. Global models and predictions of plant diversity based on advanced machine learning techniques. – *New Phytol.* 237: 1432–1445.
- Canty, A. and Ripley, B. 2021. boot: Bootstrap R (S-Plus) functions. – R package ver. 1.3-28.1, <https://cran.r-project.org/web/packages/boot/index.html>.
- Cayuela, L. et al. 2019. Taxonstand: taxonomic standardization of plant species names. – R package 2.2, <https://CRAN.R-project.org/package=Taxonstand>.
- Chamberlain, S., Oldoni, D., Barve, V., Desmet, P., Geffert, L., Mcglinn, D. and Ram, K. 2021. rgbif: interface to the global biodiversity information facility API. – R package 3.5.2, <https://cran.r-project.org/web/packages/rgbif/index.html>.
- Darwin, C. 1859. On the origin of species by means of natural selection or the preservation of favoured races in the struggle for life. – John Murray.
- Dinerstein, E. et al. 2017. An ecoregion-based approach to protecting half the terrestrial realm. – *BioScience* 67: 534–545.
- Divíšek, J., Chytrý, M., Beckage, B., Gotelli, N. J., Lososová, Z., Pyšek, P., Richardson, D. M. and Molofsky, J. 2018. Similarity of introduced plant species to native ones facilitates naturalization, but differences enhance invasion success. – *Nat. Commun.* 9: 4631.
- Dormann, C. F., Elith, J., Bacher, S., Buchmann, C., Carl, G., Carré, G., Marquéz, J. R. G., Gruber, B., Lafourcade, B., Leitão, P. J., Münkemüller, T., McClean, C., Osborne, P. E., Reineking, B., Schröder, B., Skidmore, A. K., Zurell, D. and Lautenbach, S. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. – *Ecography* 36: 27–46.
- Dullinger, I., Wessely, J., Bossdorf, O., Dawson, W., Essl, F., Gatzinger, A., Klöner, G., Kreft, H., Kuttner, M., Moser, D., Pergl, J., Pyšek, P., Thuiller, W., van Kleunen, M., Weigelt, P., Winter, M., Dullinger, S. and Beaumont, L. 2017. Climate change will increase the naturalization risk from garden plants in Europe. – *Global Ecol. Biogeogr.* 26: 43–53.
- Early, R. and Sax, D. F. 2014. Climatic niche shifts between species' native and naturalized ranges raise concern for ecological forecasts during invasions and climate change. – *Global Ecol. Biogeogr.* 23: 1356–1365.
- Elton, C. S. 1958. The ecology of invasions by animals and plants. – The Univ. of Chicago Press.
- Engelbrecht, C. J. and Engelbrecht, F. A. 2016. Shifts in Köppen-Geiger climate zones over southern Africa in relation to key global temperature goals. – *Theor. Appl. Climatol.* 123: 247–261.
- Faulkner, K. T., Burness, A., Byrne, M., Kumschick, S., Peters, K., Robertson, M., Saccaggi, D. L., Weyl, O. and Williams, V. L. 2020. South Africa's pathways of introduction and dispersal and how they have changed over time. – In: van Wilgen, B. W., Measey, J., Richardson, D., Wilson, J. R. U. and Zengeya, T. (eds), *Biological invasions in South Africa*. Springer International Publishing, pp. 313–354.
- Feng, Y., Maurel, N., Wang, Z., Ning, L., Yu, F. and van Kleunen, M. 2016. Introduction history, climatic suitability, native range size, species traits and their interactions explain establishment of Chinese woody species in Europe. – *Global Ecol. Biogeogr.* 25: 1356–1366.
- Fernández, M. and Hamilton, H. 2015. Ecological niche transferability using invasive species as a case study. – *PLoS One* 10: e0119891.
- Fick, S. E. and Hijmans, R. J. 2017. WorldClim 2: new 1-km spatial resolution climate surfaces for global land areas. – *Int. J. Climatol.* 37: 4302–4315.
- Fitzpatrick, M. C. and Hargrove, W. W. 2009. The projection of species distribution models and the problem of non-analog climate. – *Biodivers. Conserv.* 18: 2255–2261.
- Fristoe, T. S., Bleilevens, J., Kinlock, N. L., Yang, Q., Zhang, Z., Dawson, W., Essl, F., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Dufour-Dror, J. M., Sennikov, A. N., Wasowicz, P., Westergaard, K. B. and van Kleunen, M. 2023. Evolutionary imbalance, climate and human history jointly shape the global biogeography of alien plants. – *Nat. Ecol. Evol.* 7: 1633–1644.
- Fulgêncio-Lima, L. G., Andrade, A. F. A., Vilela, B., Lima-Júnior, D. P., de Souza, R. A., Sgarbi, L. F., Simião-Ferreira, J., De Marco, P. and Silva, D. P. 2021. Invasive plants in Brazil: climate change effects and detection of suitable areas within conservation units. – *Biol. Invas.* 23: 1577–1594.
- Gao, J. 2020. Global 1-km Downscaled Population Base Year and Projection Grids Based on the Shared Socioeconomic Pathways, Revision 01. – NASA Socioeconomic Data and Applications Center (SEDAC).
- GBIF 2021. org 2021. – <https://www.gbif.org/occurrence/download/0293436-200613084148143>.
- Glen, H. F. 2002. Cultivated plants of southern Africa: botanical names, common names, origins, literature. – National Botanical Institute.
- Haeuser, E., et al. 2018. European ornamental garden flora as an invasion debt under climate change. – *J. Appl. Ecol.* 55: 2386–2395.
- Hejda, M., Čuda, J., Pyšková, K., Zambatis, G., Foxcroft, L. C., MacFadyen, S., Storch, D., Tropek, R. and Pyšek, P. 2022. Water availability, bedrock, disturbance by herbivores, and climate determine plant diversity in South-African savanna. – *Sci. Rep.* 12: 338.
- Hoegh-Guldberg, O. et al. 2019. The human imperative of stabilizing global climate change at 1.5°C. – *Science* 365: eaaw6974.
- IPCC 2023. 2023: synthesis report. A report of the Intergovernmental Panel on Climate Change. Contribution of working groups I, II and III to the sixth assessment report of the Intergovernmental Panel on Climate Change (Team, H. L. and Romero, J., eds). – IPCC.
- Lambdon, P.-W., et al. 2008. Alien flora of Europe: species diversity, temporal trends, geographical patterns and research needs. – *Preslia* 80: 101–149.

- Lange, S. 2019. Trend-preserving bias adjustment and statistical downscaling with ISIMIP3BASD (v1.0). – *Geosci. Model Dev.* 12: 3055–3070.
- Lenzner, B., Magallón, S., Dawson, W., Kreft, H., König, C., Pergl, J., Pyšek, P., Weigelt, P., van Kleunen, M., Winter, M., Dullinger, S. and Essl, F. 2021. Role of diversification rates and evolutionary history as a driver of plant naturalization success. – *New Phytol.* 229: 2998–3008.
- Lin, D., Xia, J. and Wan, S. 2010. Climate warming and biomass accumulation of terrestrial plants: a meta-analysis. – *New Phytol.* 188: 187–198.
- Liu, D. et al. 2023. The impact of land use on non-native species incidence and number in local assemblages worldwide. – *Nat. Commun.* 14: 2090.
- Maurel, N., Hanspach, J., Kühn, I., Pyšek, P. and van Kleunen, M. 2016. Introduction bias affects relationships between the characteristics of ornamental alien plants and their naturalization success. – *Global Ecol. Biogeogr.* 25: 1500–1509.
- Mayer, K., Haeuser, E., Dawson, W., Essl, F., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Winter, M., Lenzner, B. and van Kleunen, M. 2017. Naturalization of ornamental plant species in public green spaces and private gardens. – *Biol. Invas.* 19: 3613–3627.
- Northrup, J. M., Rivers, J. W., Yang, Z. and Betts, M. G. 2019. Synergistic effects of climate and land-use change influence broad-scale avian population declines. – *Global Change Biol.* 25: 1561–1575.
- O'Neill, B. C., Kriegler, E., Ebi, K. L., Kemp-Benedict, E., Riahi, K., Rothman, D. S., van Ruijven, B. J., van Vuuren, D. P., Birkmann, J., Kok, K., Levy, M. and Solecki, W. 2017. The roads ahead: narratives for shared socioeconomic pathways describing world futures in the 21st century. – *Global Environ. Change* 42: 169–180.
- Oduor, A. M. O., Yang, B. and Li, J. 2023. Alien ornamental plant species cultivated in Taizhou, southeastern China, may experience greater range expansions than native species under future climates. – *Global Ecol. Conserv.* 41: e02371.
- Omer, A., Fristoe, T., Yang, Q., Maurel, N., Weigelt, P., Kreft, H., Bleilevens, J., Dawson, W., Essl, F., Pergl, J., Pyšek, P. and van Kleunen, M. 2021. Characteristics of the naturalized flora of Southern Africa largely reflect the non-random introduction of alien species for cultivation. – *Ecography* 44: 1812–1825.
- Omer, A., Fristoe, T., Yang, Q., Razanajatovo, M., Weigelt, P., Kreft, H., Dawson, W., Dullinger, S., Essl, F., Pergl, J., Pyšek, P. and van Kleunen, M. 2022. The role of phylogenetic relatedness on alien plant success depends on the stage of invasion. – *Nat. Plants* 8: 906–914.
- Omer, A., Essl, F., Dullinger, S., Lenzner, B., García-Rodríguez, A., Moser, D., Fristoe, T., Dawson, W., Weigelt, P., Kreft, H., Pergl, J., Pyšek, P., van Kleunen, M. and Wessely, J. 2024. Data from: Invasion risk of the currently cultivated alien flora in southern Africa is predicted to decline under climate change. – *Dryad Digital Repository*, <https://doi.org/10.5061/dryad.c59zw3rg2>.
- Pearman, P. B., Guisan, A., Broennimann, O. and Randin, C. F. 2008. Niche dynamics in space and time. – *Trends Ecol. Evol.* 23: 149–158.
- Pouteau, R., et al. 2021. Potential alien ranges of European plants will shrink in the future, but less so for already naturalized than for not yet naturalized species. – *Divers. Distrib.* 27: 2063–2076.
- POWO 2019. Plants of the World Online. Facilitated by the Royal Botanic Gardens. – <http://www.plantsoftheworldonline.org/>.
- Pyšek, P., Richardson, D. M., Pergl, J., Jarosík, V., Sixtová, Z. and Weber, E. 2008. Geographical and taxonomic biases in invasion ecology. – *Trends Ecol. Evol.* 23: 237–244.
- Pyšek, P. et al. 2020. Scientists' warning on invasive alien species. – *Biol. Rev. Camb. Phil. Soc.* 95: 1511–1534.
- Riahi, K., et al. 2017. The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: an overview. – *Global Environ. Change* 42: 153–168.
- Richardson, D. M. and Pyšek, P. 2012. Naturalization of introduced plants: ecological drivers of biogeographical patterns. – *New Phytol.* 196: 383–396.
- Rodríguez-Labajos, B., Binimelis, R. and Monterroso, I. 2009. Multi-level driving forces of biological invasions. – *Ecol. Econ.* 69: 63–75.
- Root, T. L., Price, J. T., Hall, K. R., Schneider, S. H., Rosenzweig, C. and Pounds, J. A. 2003. Fingerprints of global warming on wild animals and plants. – *Nature* 421: 57–60.
- Schielzeth, H. 2010. Simple means to improve the interpretability of regression coefficients. – *Methods Ecol. Evol.* 1: 103–113.
- Seebens, H., Bacher, S., Blackburn, T. M., Capinha, C., Dawson, W., Dullinger, S., Genovesi, P., Hulme, P. E., van Kleunen, M., Kühn, I., Jeschke, J. M., Lenzner, B., Liebhold, A. M., Pattison, Z., Pergl, J., Pyšek, P., Winter, M. and Essl, F. 2020. Projecting the continental accumulation of alien species through to 2050. – *Global Change Biol.* 27: 970–982.
- Šimová, I., Violle, C., Svenning, J., Kattge, J., Engemann, K., Sandel, B., Peet, R. K., Wiser, S. K., Blonder, B., McGill, B. J., Boyle, B., Morueta-Holme, N., Kraft, N. J. B., van Bodegom, P. M., Gutiérrez, A. G., Bahn, M., Ozinga, W. A., Tószögyová, A. and Enquist, B. J. 2018. Spatial patterns and climate relationships of major plant traits in the New World differ between woody and herbaceous species. – *J. Biogeogr.* 45: 895–916.
- Thapa, S., Chitale, V., Rijal, S. J., Bisht, N. and Shrestha, B. B. 2018. Understanding the dynamics in distribution of invasive alien plant species under predicted climate change in western Himalaya. – *PLoS One* 13: e0195752.
- Thuiller, W., Georges, D., Gueguen, M., Engler, R., Breiner, F., Lafourcade, B. and Patin, R. 2020. biomod2: Ensemble Platform for Species Distribution Modeling. – R package, 3.4.6, <https://biomodhub.github.io/biomod2/>.
- Thuiller, W., Richardson, D. M., Pyšek, P., Midgley, G. F., Hughes, G. O. and Rouget, M. 2005. Niche-based modelling as a tool for predicting the risk of alien plant invasions at a global scale. – *Global Change Biol.* 11: 2234–2250.
- Thuiller, W., Lafourcade, B., Engler, R. and Araújo, M. B. 2009. BIOMOD – a platform for ensemble forecasting of species distributions. – *Ecography* 32: 369–373.
- van Kleunen, M., et al. 2018. The changing role of ornamental horticulture in alien plant invasions. – *Biol. Rev. Camb. Phil. Soc.* 93: 1421–1437.
- van Kleunen, M., Bossdorf, O. and Dawson, W. 2018. The ecology and evolution of alien plants. – *Annu. Rev. Ecol. Evol. Syst.* 49: 25–47.
- van Kleunen, M. et al. 2019. The Global Naturalized Alien Flora (GloNAF) database. – *Ecology* 100: e02542.
- van Kleunen, M., Xu, X., Yang, Q., Maurel, N., Zhang, Z., Dawson, W., Essl, F., Kreft, H., Pergl, J., Pyšek, P., Weigelt, P., Moser, D., Lenzner, B. and Fristoe, T. S. 2020. Economic use of plants is key to their naturalization success. – *Nat. Commun.* 11: 3201.
- Verbruggen, W., Schurgers, G., Horion, S., Ardö, J., Bernardino, P. N., Cappelaere, B., Demarty, J., Fensholt, R., Kergoat, L.,

- Sibret, T., Tagesson, T. and Verbeeck, H. 2021. Contrasting responses of woody and herbaceous vegetation to altered rainfall characteristics in the Sahel. – *Biogeosciences* 18: 77–93.
- Vilà, M. and Hulme, P. E. 2017. Non-native species, ecosystem services, and human well-being. – In: Vilà, M. and Hulme, P. E. (eds), *Impact of biological invasions on ecosystem services*. Springer, pp. 1–14.
- Wang, P., Huang, K. and Hu, S. 2020. Distinct fine-root responses to precipitation changes in herbaceous and woody plants: a meta-analysis. – *New Phytol.* 225: 1491–1499.
- Wells, M. J., Poynton, R. J., Balsinhas, A. A., Musil, K. J., Joffe, H., Hoepen, E. and Abbott, S. K. 1986. History of introduction of invasive alien plants to southern Africa. I. – In: Macdonald, A. W. and Kruger, A. A. (eds), *Ecology and management of biological invasions in southern Africa*, proceedings of the national synthesis symposium on the ecology of biological invasions. Oxford Univ. Press.
- Williams, J. W., Jackson, S. T. and Kutzbach, J. E. 2007. Projected distributions of novel and disappearing climates by 2100 AD. – *Proc. Natl Acad. Sci. USA* 104: 5738–5742.
- Zizka, A., Silvestro, D., Andermann, T., Azevedo, J., Duarte Ritter, C., Edler, D., Farooq, H., Herdean, A., Ariza, M., Scharn, R., Svantesson, S., Wengström, N., Zizka, V. and Antonelli, A. 2019. CoordinateCleaner: standardized cleaning of occurrence records from biological collection databases. – *Methods Ecol. Evol.* 10: 744–751.