

Assessing people with visual impairments' access to information, awareness and satisfaction with high-tech assistive technology

British Journal of Visual Impairment

2024, Vol. 42(1) 149–163

© The Author(s) 2022



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/02646196221131746

journals.sagepub.com/home/jvi**Isabelle Liang****Ben Spencer**

University of Bath, UK

Meike Scheller 

Durham University, UK

Michael J Proulx **Karin Petrini** 

University of Bath, UK

Abstract

Assistive technology (AT) devices are designed to help people with visual impairments (PVI) perform activities that would otherwise be difficult or impossible. Devices specifically designed to assist PVI by attempting to restore sight or substitute it for another sense have a very low uptake rate. This study, conducted in England, aimed to investigate why this is the case by assessing accessibility to knowledge, awareness, and satisfaction with AT in general and with sensory restoration and substitution devices in particular. From a sample of 25 PVI, ranging from 21 to 68 years old, results showed that participants knew where to find AT information; however, health care providers were not the main source of this information. Participants reported good awareness of different ATs, and of technologies they would not use, but reported poor awareness of specific sensory substitution and restoration devices. Only three participants reported using AT, each with different devices and varying levels of satisfaction. The results from this study suggest a possible breakdown in communication between health care providers and PVI, and dissociation between reported AT awareness and reported access to AT information. Moreover, awareness of sensory restoration and substitution devices is poor, which may explain the limited use of such technology.

Corresponding author:

Karin Petrini, Department of Psychology, University of Bath, Claverton Down, Bath BA2 7AY, UK.

Email: K.Petrini@bath.ac.uk

Keywords

Access, assistive technologies, awareness, satisfaction, sensory restoration devices, sensory substitution devices

Introduction

Visual impairments, or vision that cannot be corrected with lenses or surgery and blindness, affects a large proportion of the human population. Globally, it is estimated that 285 million people, or 3.7% of the population, are living with visual impairment (He et al., 2020; National Health Service, 2018; World Health Organization, 2021). Visual impairment affects interactions with the physical world (West et al., 2002), interactions with others, and can result in poor mental health (Demmin & Silverstein, 2020). Chia et al. (2004) found that visual impairment significantly decreased one's quality of life, on physical, social, emotional, and mental factors, compared to those without visual impairment and those with correctable vision. Thus, there is great potential for assistive devices in aiding people with visual impairments (PVIs) and improving quality of life.

The United Kingdom's Medicines & Health care products Regulatory Agency (2021) defines assistive technology (AT) as any product designed to assist individuals with impairments with actions that would otherwise be difficult or impossible. As this report is focusing on visual impairments, further references to ATs refer to those designed specifically to assist with visual impairment. AT can be classified into low-tech, medium-tech, and high-tech devices: low-tech ATs require no electricity to operate, medium-tech devices require electricity but are not computerised, and high-tech devices require both electricity and are computerised (Kaye et al., 2008). Examples of popular "low-tech" devices for PVIs include canes and guide dogs or Braille (Illinois Library, 2020). Medium-tech devices for PVIs are less common, an example being electricity powered cassette tape readers (Varghese, 1996). Technological advances have facilitated the use of computer processing in ATs to provide more advanced assistance for PVIs through the development of high-tech devices (Bhowmick & Hazarika, 2017). High-tech ATs include sensory restoration devices and devices that utilise sensory substitution technology, either within a dedicated device or integrated into existing devices such as a mobile phone.

Sensory restoration devices for PVIs aim to restore functionally useful visual perception through stimulation of visual neural pathways (Lewis et al., 2015). Most devices are invasive, requiring surgical implantation, though some non-invasive techniques are currently being investigated (Sabel et al., 2020). Four Conformité Européenne (CE) marked sensory restoration devices, all retinal prostheses, are: Argus II (Luo & Da Cruz, 2016), Intelligent Retinal Implant System (IRIS) II (Pixium Vision, 2016), Alpha IMS (Stingl et al., 2013), and Alpha AMS (Daschner et al., 2018). All devices produced better performance on visual tasks, such as object localisation, shape recognition, and letter reading, with the devices on rather than off (Bloch et al., 2019). However, their effectiveness in helping PVIs during spatial navigation and wayfinding remains unclear (e.g., Garcia et al., 2015).

Unlike sensory restoration devices, sensory substitution devices do not require surgery. Sensory substitution technologies for PVIs communicate visual information via another, intact sensory modality (Ward & Wright, 2014); for example, playing different sounds to communicate different visual features, such as elevation and horizontal location, within the visual field. The technology is grounded in the understanding that the brain can interpret sensory information from one modality (e.g., sound) as information from a different modality (e.g., visual) (Shimojo & Shams, 2001; Solomon-Harris et al., 2013). Numerous devices designed specifically for sensory substitution have been developed (Scheller et al., 2018), such as The vOICe (visual-to-auditory) (Meijer, 1992), and BrainPort (visual-to-tactile) (Fale et al., 2016). Many devices designed specifically for sensory substitution have been successful in improving functional outcomes where sensory restoration

devices have struggled (Erickson-Davis & Korzybska, 2021; Garcia et al., 2015). For example, sensory substitution devices have been found to improve spatial competence (Chebat et al., 2020; Jicol et al., 2020), object recognition (Arnoldussen & Rhode, 2010) and object localisation (Proulx et al., 2008).

Despite encouraging results from clinical tasks, sensory restoration and substitution devices are rarely used outside of the laboratory (Elli et al., 2014; Griffin-Shirley et al., 2017; Lloyd-Esenkaya et al., 2020). This may be because the devices do not improve processes such as object identification and navigation enough to support daily life. For example, statistically significant improvements on object identification may not represent large-enough benefits for use in everyday life. Luo et al. (2014) presented photos of three-dimensional objects and asked patients using the Argus II to identify the objects. They found a significant improvement in object identification when the device was switched on. However, correct identification with the system on was only 35.7% on average ($SD=14.6\%$), which may not be accurate enough to be useful in identifying objects in everyday life. With regards to navigation, retinal prostheses provide little useful spatial information (Horne et al., 2015), and some evidence suggests that users may not utilise the perceptual experiences provided by a sensory restoration device to aid in navigation (Garcia et al., 2015). The most popular assistive device used for navigation and mobility by PVI's to date is the long cane (Batterman et al., 2018; Dahlin-Ivanoff & Sonn, 2004). Sensory restoration and substitution devices may not replace the long cane because of the greater need for training and cognitive effort (Hamilton-Fletcher et al., 2016), in exchange for little additional benefits. Moreover, low-tech devices are difficult to surpass because of their reliability, lack of reliance on electricity, and relative inexpensiveness (Manduchi & Coughlan, 2012). Thus, the limited benefits of sensory restoration and substitution devices compared to low-tech ATs may explain their poor uptake.

On the contrary, research that has taken into account PVI's opinions suggests that the low uptake of sensory restoration and substitution devices may not be related to device functionality at all. Some PVI's have expressed disappointment about the neglect of user experience in such devices (Lenay et al., 2003). For example, in Hamilton-Fletcher et al.'s study (2016), while participants reported optimism for some visual-to-auditory sensory substitution devices' potential in aiding everyday life, some devices were also criticised for the unpleasant auditory aesthetics. Others were highly praised for their focus on providing PVI's with sensory information beyond function, such as colour; a feature absent from most sensory substitution devices. This highlights the crucial distinction between a functionally effective device, and a device that PVI's like and want to use; a distinction often neglected in the development of ATs (Manduchi & Coughlan, 2012). Accordingly, another essential feature of ATs is subtlety. Many PVI's refrain from using ATs such as the cane due to the feelings of shame and stigmatisation that arise with being identified as visually impaired (Hayeems et al., 2005). Some sensory substitution devices are characterised by the same limitation. For example, part of the BrainPort device is placed in the mouth (Wicab Inc., 2010) and has been criticised for being indiscrete as well as impractical (Upson, 2007). Moreover, sensory restoration and substitution devices are often very expensive, thus rendering these devices inaccessible. A survey by Manjari et al. (2020) assessed a wide range of sensory substitution devices and concluded that seven of the 37 devices assessed were economic, and all but two were user-friendly. However, this study did not involve PVI's in drawing these conclusions, thus these conclusions should be regarded cautiously. This highlights the need for the inclusion of PVI's voices in research regarding ATs to refocus the industry on PVI's needs and desires, rather than on bioengineering solutions which may have contributed to the low uptake of such devices (Manduchi & Coughlan, 2012).

Another barrier to AT access may be awareness. Low-tech ATs, such as the cane, have existed for many decades (Strong, 2009) and are established as common assistive tools, as well as symbols of visual impairment (Due & Lange, 2018). Hence, PVI's are more likely to be aware of such technology (Senjam et al., 2020). As many high-tech ATs are relatively new, it is possible that PVI's are

unaware of the existence of sensory restoration and substitution devices. Senjam et al. (2020) found that the majority of participants in a sample of visually impaired students in Delhi had awareness of Braille-related devices and long canes, but much fewer were aware of more high-tech devices such as electronic magnifier aids, talking calculators and mobile applications. However, visually impaired students showed high awareness for talking watches and screen readers. Almost all devices investigated by Senjam et al.'s (2020) study were designed to aid with very specific aspects of life; namely school life. Similarly, Okonji and Ogwezzy (2019) studied PVI's awareness of older ATs that specifically aid access to computers. Thus, a better understanding of AT awareness, especially for more recent devices in a wider range of contexts, is lacking. Furthermore, the studies by Senjam et al. (2020) and Okonji and Ogwezzy (2019) did not address possible barriers to accessing AT information, which could limit AT awareness. Hence, this study aimed to examine specifically: how accessible AT information is to PVI's, the extent of PVI's awareness of AT (particularly of sensory restoration and substitution devices), and how satisfied PVI's are with their AT (particularly with their sensory restoration or substitution device).

Method

Participants

Visually impaired participants were recruited via volunteer sampling by distributing the survey link to Facebook support-groups and through the Royal National Institute of Blind People. In total, 25 participants completed the study (11 males, 13 females, and 1 unspecified), ranging from 21 to 68 years old ($M=44.91$, $SD=17.27$). The sample included a range of visual impairments: 7 participants were completely blind, 12 were severely sight impaired, and 6 were sight impaired. Study information was provided and electronic consent was obtained prior to study initiation. Ethical approval was granted by the University of Bath's Psychology Research Ethics Committee (reference no: UG 19-038). The study was carried out in 2019/2020.

De-sign, material, and procedure

This study was an online questionnaire study made and published using Qualtrics (2021). The data were analysed in three domains: Accessing AT information, awareness, and satisfaction. No validated measures of awareness of ATs, or accessing AT information, exist yet. Thus, a set of questions and measures were created for this study. The accessing AT information questionnaire comprised questions about sources of AT information and the extent to which PVI's actively searched for information (Table 1). The awareness questionnaire comprised questions about general knowledge of sensory restoration and substitution devices, and questions probing the depth of such awareness (Table 3). These items were measured on a 5-point Likert-type scale with "1" = *strongly agree* and "5" = *strongly disagree* (Nadler et al., 2015).

Participants who answered "yes" to the additional question "Do you currently use an assistive sensory restoration or substitution device?" were further invited to complete the satisfaction and preference questionnaires. The satisfaction and preference questionnaires were analysed in conjunction with the qualitative responses collected on the same topics. The Quebec User Evaluation of Satisfaction with Assistive Technology (QUEST 2.0) (Demers et al., 2000) was used to assess device and service satisfaction. Previous studies have shown this scale to have excellent test-retest reliability (Demers et al., 2002) and robust internal consistency (Galeoto et al., 2018). The Assistive Technology Device Predisposition Assessment for device users (ATDPA) (Scherer, 2005) assessed satisfaction with positive impact made to different aspects of participants' lives. This scale also has an excellent test-retest reliability and a relatively high internal consistency (Koumpouros et al., 2017). Both questionnaires

Table 1. Questionnaire items included in the accessing information analysis.

Item	Frequency				
	1	2	3	4	5
I know where I can find out more about new ATs	11	8	2	4	0
My healthcare provider is my main source of information about new ATs	0	1	3	3	18
I often search for ATs on the internet	12	5	2	3	3
I am always up to date with the newest developments in ATs	4	9	4	6	2

Note. AT = assistive technology.

used a 5-point Likert-type scale where ‘1’ = *extremely satisfied* and ‘5’ = *extremely dissatisfied*. The five questions regarding AT preferences used a 3-point Likert-type scale with “1” = yes, “2” = maybe, and “3” = no, and can be found in the Supplemental Material.

Results

One-sample Wilcoxon’s signed rank tests were used to compare the hypothetical median (“3,” or “neither agree nor disagree”) against the observed median. This was done as the data were on an ordinal scale and mostly non-normally distributed ($W[21] \leq .868$, $p \leq .009$), as assessed by Shapiro–Wilk tests. All p values were Bonferroni corrected to account for familywise error, with this results section reporting the corrected p values.

Accessing AT information

Frequency distributions can be found in Table 1, results from the Wilcoxon tests are shown in Table 2, and boxplots for the data are presented in Figure 1. Responses in this domain indicated that participants significantly agreed that they knew where to find information about new AT but strongly disagreed that their health care provider was their main source of information. No significant difference from “neither agree or disagree” was found for the items regarding often searching the internet for ATs and for always being up to date with the newest developments in ATs.

Awareness

Frequency distributions of the responses in the domain of AT awareness can be found in Table 3, results from the Wilcoxon tests are shown in Table 4, and boxplots for the data are presented in Figure 2.

The observed medians for four out of ten items for awareness were significantly different from the hypothetical medians. The results found that participants were significantly more aware than ambivalent of different ATs and technologies they would not use. Participants also reported significantly poorer level of awareness than neutral for the specific sensory restoration and substitution devices.

Satisfaction

Only three participants reported using high-tech AT for visual impairment, these were an iPhone/laptop, a Miniguide (Hill & Black, 2003) and BrainPort Vision Pro; the latter two being sensory substitution devices. With only three participants reporting using an AT for visual assistance, group




Figure 1. Box plots of the median and interquartile range of Likert-type scores for items regarding accessing AT information.

Note. The circles with corresponding numbers represent outlying data points. AT = assistive technology.

Table 2. Test statistics for the domain accessing AT information.

Item	N	Z value	p value	Median
Know where to find out about new ATs	25	26.00	.006	2 (“Somewhat agree”)
Healthcare provider	25	250.50	<.001	5 (“Strongly disagree”)
Internet search	25	61.50	.224	–
Up to date with new AT	25	85.00	>.999	–

Note. AT = assistive technology.

analysis was not possible. However, Table 5 shows the individual satisfaction scores for device, service, and positive impact.

One participant reported using a bone-anchored hearing aid which is a sensory restoration device for auditory impairment, hence the data for this participant were not further analysed (but can be found in the supplemental material).

Device satisfaction measured the satisfaction with practical aspects of the device such as comfort, size, and effectiveness. Services satisfaction measured satisfaction with the customer service received for the device. Positive impact satisfaction measured the device’s ability to improve different areas of the users’ life.

The Miniguide user reported good device satisfaction, but was ambivalent regarding service and positive impact satisfaction. The BrainPort user reported ambivalence for device satisfaction, ambivalence/slight satisfaction for the services but expressed dissatisfaction for its positive impact. The Miniguide user reported that they would recommend their device, but the Brainport user reported that they would not.

In the preferences domain, the iPhone/laptop user and Miniguide user reported that they were disappointed by the promises made by companies developing new ATs. Both sensory substitution device users reported that they would prefer sensory restoration over substitution, and would invest time and effort into training for an AT they knew would improve their daily life. However, while the Miniguide user reported that they strongly agreed that they knew what sensory restoration was,




Figure 2. Box plots of the median and interquartile ranges of Likert-type scores for items regarding awareness.

Note. For these items, stronger disagreement to the statements indicates lower awareness stronger agreement indicate higher awareness. The circles with corresponding numbers represent outlying data points.

Table 3. Questionnaire items included in the awareness analysis.

Item	Frequencies				
	1	2	3	4	5
I have a good knowledge of the different ATs for blind and partially sighted persons	10	8	4	3	0
There a lot of ATs for blind and partially sighted persons that I would not use	5	10	6	3	0
I know what sensory restoration is (e.g. retinal implants)	4	10	5	2	6
I know what sensory substitution is.	2	8	2	4	8
I know what the IMS Alpha and Arus II are ^a	0	0	3	3	17
I know what The vOICE or BrainPort are	3	2	2	0	17
I am aware of the advantages of sensory restoration	2	8	5	3	7
I am aware of the disadvantages of sensory restoration	1	6	6	2	8
I am aware of the advantages of sensory substitution	3	3	8	2	8
I am aware of the disadvantages of sensory substitution	1	3	6	4	9

Note. Not all participants responded to each question, therefore some frequencies do not add to 25. AT= assistive technology.

^aDue to a spelling mistake, ARGUS II was not considered further.

the BrainPort user reported strongly disagreeing. This suggests that the BrainPort user’s reported preference for sensory restoration over substitution may be poorly informed and instead possibly due to interpreting sensory restoration as full visual recovery, as the name suggests.

There was not enough data to conduct a thematic analysis, with only the Miniguide user and iPhone/laptop user responding to open-ended questions. In response to how the device helped with their day-to-day life, the Miniguide user emphasised the enhanced spatial awareness regarding other people as useful (Extract 1), and later suggested that this resulted in a higher confidence and an increased sense of safety (Extract 2). Conversely, the iPhone/laptop user reported using their

Table 4. Test statistics for the domain 'awareness of ATs.

Item	N	Z value	p value	Median
Good knowledge	25	18.00	.007	2 ("Somewhat agree")
AT would not use	24	21.00	.042	2 ("Somewhat agree")
Sensory restoration	25	118.00	>.999	–
Sensory substitution	24	166.00	>.999	–
IMS Alpha	23	210.00	<.001	5 ("Strongly disagree")
vOICE or Brain Port	24	212.50	.028	5 ("Strongly disagree")
Advantages of sensory restoration	25	130.00	>.999	–
Disadvantages of sensory restoration	23	113.00	>.999	–
Advantages of sensory substitution	24	94.00	>.999	–
Disadvantages of sensory substitution	23	128.50	.154	–

Note. AT = assistive technology.

Table 5. Satisfaction scores.

Assistive device	Device satisfaction score (QUEST 2.0)	Services satisfaction (QUEST 2.0)	Positive impact satisfaction score (ATDPA)
Miniguide	4.00	2.75	3.42
BrainPort Vision Pro	3.37	3.75	1.63
iPhone/laptop	4.13	4.00	4.33

Note. QUEST = Quebec User Evaluation of Satisfaction with Assistive Technology; ATDPA = Assistive Technology Device Predisposition Assessment.

device to support with interactions with digital content more so than facilitating engagement with the physical world (Extract 3), though they did refer to Global Positioning System (GPS) use.

Extract 1. (from Miniguide user)

Question: How does this device help your day-to-day life?

"Helps me to be aware of people entering my space."

Extract 2. (from Miniguide user)

Question: What reasons do you have for using your assistive device?

"Increased awareness of surroundings resulting in higher confidence and security."

Extract 3. (from iPhone/laptop user)

Question: How does this device help your day-to-day life?

Emailing, using GPS, watching videos onLINE and much more

Discussion

This study examined how participants source information about AT information, their awareness of high-tech AT, and how satisfied they are with high-tech AT. Results showed that participants knew where to find information about ATs, but their health care providers were not their main source of this information. It is unknown whether this is due to the existence of more prominent sources of information, a limited awareness of different devices from health care providers, or barriers to PVI's accessing the information or health care services. There are numerous challenges that

hinder PVI's access to health care services and communication between PVI's and health care providers. These include, information being communicated via printed text (Cupples et al., 2012), limited information and assistance about health care entitlement (Sharts-Hopko et al., 2010), and a lack of sensitivity to PVI needs (Withers & Speight, 2017). Thus, these factors may not only prevent good communication between PVI's and health care providers, but limit access to the health care services altogether (Cupples et al., 2012). Understanding why health care providers are not the main hub of information about new ATs for PVI's and finding a targeted solution could be an easy barrier to overcome for PVI's, possibly increasing the uptake of some of available high-tech ATs.

Participants also remained neutral in agreeing or disagreeing with whether they often searched the internet for information about ATs and with whether they were always up to date with the latest developments in AT. Thus, despite knowing where they could obtain information from, they were not accessing it. One possible explanation is that there may be accessibility issues creating barriers to information. Luchtenberg et al. (2008) found that 82% of 139 websites containing medical information were not fully accessible to PVI's. Since 2008, issues with accessibility seem to have somewhat improved, with the implementation of text-to-speech functions on major web browsers (Ebden & Sproat, 2015). However, accessibility remains a barrier for accessing health information through difficulties in searching the internet (Sahib et al., 2012; Xie et al., 2018). Alternatively, Okonji and Ogwezzy (2019) found that some participants did not use high-tech AT because they had no need (29%) and lacked interest (22%), which may suggest a disinterest in AT. Lenay et al. (2003) found that PVI's were disappointed by ATs due to the expectations created around them. Similarly, the Miniguide user in this study expressed disappointment at the promises made by companies developing new ATs. Thus, this could contribute to the lack of interest. It is important to determine why participants are not seeking information about ATs as it indicates what PVI's want from ATs, if they want them at all, so that resources and research can be focused where they will be most useful (Hamilton-Fletcher et al., 2016). As this study found that participants were not particularly responsive to open-ended questionnaire questions, future research could involve interviewing PVI's to understand why they do or do not look for information on ATs.

The need to develop our understanding of why PVI's are not seeking AT information, nor receiving AT information from their health care providers is highlighted by the fact that, despite agreeing that they had good knowledge of different ATs, participants reported poor awareness of the IMS Alpha, The vOICE and BrainPort devices. While this suggests that participants may have good awareness of ATs other than those asked about, it also highlights the failure for information about specific relatively new high-tech ATs to reach PVI's. As participants are not receiving AT information primarily from health care providers and are not keeping up to date with AT development, this is perhaps unsurprising, further highlighting a need to understand why PVI's are not seeking or receiving as part of their health service AT information.

As only three participants reported using an AT device, limited generalisations about satisfaction with devices can be made; however, the findings are in line with previous research. The Miniguide user reported good device satisfaction. This is in line with Roentgen et al.'s (2012) findings that other users were satisfied with the device's qualities such as discrete size, lightweight nature, and the ability to switch between visual-to-auditory and visual-to-tactile substitution. The Miniguide user in this study reported that their device was used for spatial awareness, particularly awareness of other people to provide a higher sense of confidence and security. This sentiment is echoed by other PVI's as a reason for AT use (Hersh, 2015).

The BrainPort user's notable dissatisfaction with the positive impact of their device reflects criticism that despite being able to significantly improve ability on laboratory tasks (Grant et al., 2016), it does not aid functioning in daily life (Manduchi & Coughlan, 2012; Upson, 2007). The BrainPort user reported ambivalence towards device satisfaction, suggesting that the more physical

aspects of the device were neither satisfying nor dissatisfying. BrainPort has been criticised for being uncomfortable (Upson, 2007), though some users noted that the discomfort is not insurmountable (Kastrup et al., 2009), perhaps explaining the user's ambivalence.

While iPhone/laptop are not specific sensory substitution devices, this participant's report of their use as AT is valuable within the wider context of AT research, particularly regarding the integration of sensory substitution technology into existing devices. The iPhone/laptop user's satisfaction with their mobile devices reflects the findings of previous studies indicating that AT on mobile devices has been well received. Griffin-Shirley et al. (2017) found that 95% of PVI's used a mobile device, and that more than 90% of these participants reported using specialised applications. Of those individuals, 95.4% considered the applications to be useful, and 91.1% considered them to be accessible. This rate of uptake far exceeds that of rare dedicated sensory restoration or substitution device use (Elli et al., 2014; Griffin-Shirley et al., 2017). The participant in this study used their devices mostly for interaction with the digital world, but also for GPS navigation. Previous research has found that independence in mobility is highly valued by PVI's (Quinones et al., 2011). Thus, the fact that mobile devices can integrate this accessibility feature as well as others is highly beneficial; especially as PVI's prefer AT on their smartphones to integrate multiple assistive functions into one device (Hamilton-Fletcher et al., 2016). This benefit makes smartphone AT convenient (Senjam, 2021), economical, and relatively discrete in comparison to sensory substitution and restoration specific devices.

The benefits and popularity of integrating AT with common mobile devices, such as smartphones (Vailshery, 2021), indicate a valuable direction for AT development. Advances in computer science have created a rise in artificial intelligence (AI) AT applications for smartphones, often for free (Kugler, 2020). These aim to assist with functions such as object identification, facial identification, emotion identification, typed and handwritten text reading, and scene description (Parikh et al., 2018; Qureshi et al., 2021). Moreover, AI technology is being used to create ATs to assist with social interaction (Microsoft, 2021; Salido et al., 2016). This is particularly beneficial as sensory substitution devices are currently limited in their ability to assist with social interaction and aesthetic subtlety (Phillips & Proulx, 2018). Furthermore, these AI systems focus on providing practical aids to PVI's without struggling to follow through on promises of sensory experiences (Lenay et al., 2003), as sensory substitution and restoration devices do. While sensory substitution and restoration devices continue to improve to meet the needs of PVI's, a focus on the development of AI smartphone applications for PVI's may be highly valuable and preferable in the near future. The AI applications are simple in operation (Kugler, 2020), requiring less cognitive effort than sensory restoration and substitution devices where experiences have to be interpreted. Moreover, many applications with different purposes can be downloaded onto one device, which many PVI's already have: the smartphone (Griffin-Shirley et al., 2017). Much of these AI applications are new, thus little research has been conducted on user experiences of them. Therefore, future research should compare the functional capabilities and user satisfaction of these AI applications.

However, the results for AT satisfaction needs to be viewed cautiously as only three participants out of 25 reported using a high-tech AT device and were thus able to complete the satisfaction domain. Furthermore, only two participants reported using a sensory restoration and substitution device. While this highlights the limited uptake of sensory substitution and restoration devices (Elli et al., 2014; Griffin-Shirley et al., 2017), it does limit our understanding of PVI's attitudes towards the devices on a wider scale, thus limiting the generalisation of our conclusions. In particular, these participants reported using their ATs for various functions, such as online access, navigation and spatial awareness; however, inferences cannot be made regarding the functions PVI's find useful due to the small sample size. Therefore, future research aiming to further investigate device satisfaction should sample from a target population of device users, such that PVI preferences around

AT functions can be determined to guide future AT research and design. In addition, it is acknowledged that 25 participants is a small sample size, thus the proportion of AT users to non-AT users may not be representative of the PVI population. Moreover, a larger sample size would allow more confident conclusions to be drawn about the levels of awareness and of access to AT information within the PVI population.


Despite this limitation, this research still provides important insight into PVI perspectives on ATs and a basis for future. The strong disagreement of PVIs about health care being a main source of AT information may indicate a breakdown in communication between PVIs and health care providers that should be further investigated and improved. With PVIs reporting a lack of seeking new AT information yet reporting good knowledge of AT devices, future research could also investigate the dissociation between awareness and accessing of information. Focus group studies may be valuable in further investigating these findings, as PVI experience with accessing AT information and interactions with health care services are complex topics. Thus, the opportunity to engage in discussion with others may help PVIs to dissect their experiences and attitudes towards high-tech ATs.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iDs

Meike Scheller  <https://orcid.org/0000-0002-3021-5614>

Michael J Proulx  <https://orcid.org/0000-0003-4066-3645>

Karin Petrini  <https://orcid.org/0000-0001-5354-5600>

Supplemental material

Supplemental material for this article is available online.

References

- Arnoldussen, A., & Rhode, K. (2010). Enhanced perception with the BrainPort vision device. *Investigative Ophthalmology & Visual Science, 51*(13), 3634–3634.
- Batterman, J. M., Martin, V. F., Yeung, D., & Walker, B. N. (2018). Connected cane: Tactile button input for controlling gestures of iOS voiceover embedded in a white cane. *Assistive Technology, 30*(2), 91–99. <https://doi.org/10.1080/10400435.2016.1265024>
- Bhowmick, A., & Hazarika, S. M. (2017). An insight into assistive technology for the visually impaired and blind people: State-of-the-art and future trends. *Journal on Multimodal User Interfaces, 11*(2), 149–172. <https://doi.org/10.1007/s12193-016-0235-6>
- Bloch, E., Luo, Y., & da Cruz, L. (2019). Advances in retinal prosthesis systems. *Therapeutic Advances in Ophthalmology, 11*, 2515841418817501. <https://doi.org/10.1177/2515841418817501>
- Chebat, D. R., Schneider, F. C., & Ptito, M. (2020). Spatial competence and brain plasticity in congenital blindness via sensory substitution devices. *Frontiers in Neuroscience, 14*, Article 815. <https://doi.org/10.3389/fnins.2020.00815>
- Chia, E. M., Wang, J. J., Roctchina, E., Smith, W., Cumming, R. R., & Mitchell, P. (2004). Impact of bilateral visual impairment on health-related quality of life: The Blue Mountains Eye Study. *Investigative Ophthalmology & Visual Science, 45*(1), 71–76. <https://doi.org/10.1167/iovs.03-0661>
- Cupples, M. E., Hart, P. M., Johnston, A., & Jackson, A. J. (2012). Improving healthcare access for people with visual impairment and blindness. *British Medical Journal, 344*, Article e542. <https://doi.org/10.1136/bmj.e542>
- Dahlin-Ivanoff, S., & Sonn, U. (2004). Use of assistive devices in daily activities among 85-year-olds living at home focusing especially on the visually impaired. *Disability and Rehabilitation, 26*(24), 1423–1430. <https://doi.org/10.1080/09638280400000906>

- Daschner, R., Rothermel, A., Rudolf, R., Rudolf, S., & Stett, A. (2018). Functionality and performance of the subretinal implant chip Alpha AMS. *Sensors and Materials*, 30(2), 179–192. <https://doi.org/10.18494/SAM.2018.1726>
- Demers, L., Monette, M., Lapierre, Y., Arnold, D. L., & Wolfson, C. (2002). Reliability, validity, and applicability of the Quebec User Evaluation of Satisfaction with assistive Technology (QUEST 2.0) for adults with multiple sclerosis. *Disability and Rehabilitation*, 24(1–3), 21–30. <https://doi.org/10.1080/09638280110066352>
- Demers, L., Weiss-Lambrou, R., & Ska, B. (2000). *Quebec User Evaluation of Satisfaction with Assistive Technology* (Versione 2.0). The Institute for Matching Persons and Technology.
- Demmin, D. L., & Silverstein, S. M. (2020). Visual impairment and mental health: Unmet needs and treatment options. *Clinical Ophthalmology*, 14, 4229–4251. <https://doi.org/10.2147/OPHT.S258783>
- Due, B., & Lange, S. (2018). Semiotic resources for navigation: A video ethnographic study of blind people's uses of the white cane and a guide dog for navigating in urban areas. *Semiotica*, 2018(222), 287–312. <https://doi.org/10.1515/sem-2016-0196>
- Ebden, P., & Sproat, R. (2015). The Kestrel TTS text normalization system. *Natural Language Engineering*, 21(3), 333–353. <https://doi.org/10.1017/S1351324914000175>
- Elli, G. V., Benetti, S., & Collignon, O. (2014). Is there a future for sensory substitution outside academic laboratories? *Multisensory Research*, 27(5–6), 271–291. <https://doi.org/10.1163/22134808-00002460>
- Erickson-Davis, C., & Korzybska, H. (2021). What do blind people 'see' with retinal prostheses? Observations and qualitative reports of epiretinal implant users. *PLOS ONE*, 16(2), Article e0229189. <https://doi.org/10.1371/journal.pone.0229189>
- Fale, A., Naik, A., Asegaonkar, K., Landge, A., & Atkare, G. (2016). Brainport vision technology. *International Research Journal of Engineering and Technology*, 3, 475–483.
- Galeoto, G., Colucci, M., Guarino, D., Esposito, G., Cosma, E., De Santis, R., Grifoni, G., Valente, D., & Tofani, M. (2018). Exploring validity, reliability, and factor analysis of the Quebec user evaluation of satisfaction with assistive technology in an Italian population: A cross-sectional study. *Occupational Therapy in Health Care*, 32(4), 380–392. <https://doi.org/10.1080/07380577.2018.1522682>
- Garcia, S., Petrini, K., Rubin, G. S., Da Cruz, L., & Nardini, M. (2015). Visual and non-visual navigation in blind patients with a retinal prosthesis. *PLOS ONE*, 10(7), Article e0134369. <https://doi.org/10.1371/journal.pone.0134369>
- Grant, P., Spencer, L., Arnoldussen, A., Hogle, R., Nau, A., Szlyk, J., Nussdorf, J., Fletcher, D. C., Gordon, K., & Seiple, W. (2016). The functional performance of the BrainPort V100 device in persons who are profoundly blind. *Journal of Visual Impairment & Blindness*, 110(2), 77–88.
- Griffin-Shirley, N., Banda, D. R., Ajuwon, P. M., Cheon, J., Lee, J., Park, H. R., & Lyngdoh, S. N. (2017). A survey on the use of mobile applications for people who are visually impaired. *Journal of Visual Impairment & Blindness*, 111(4), 307–323. <https://doi.org/10.1177/0145482X1711100402>
- Hamilton-Fletcher, G., Obrist, M., Watten, P., Mengucci, M., & Ward, J. (2016, May). 'I always wanted to see the night sky' blind user preferences for sensory substitution devices. In *Proceedings of the 2016 CHI Conference on Human Factors in Computing Systems* (pp. 2162–2174). Association for Computing Machinery. <https://doi.org/10.1145/2858036.2858241>
- Hayeems, R. Z., Geller, G., Finkelstein, D., & Faden, R. R. (2005). How patients experience progressive loss of visual function: A model of adjustment using qualitative methods. *British Journal of Ophthalmology*, 89(5), 615–620.
- He, Y., Nie, A., Pei, J., Ji, Z., Jia, J., Liu, H., Wan, P., Ji, M., Zhang, C., Zhu, Y., & Wang, X. (2020). Prevalence and causes of visual impairment in population more than 50 years old: The Shaanxi Eye Study. *Medicine*, 99(20), Article e20109. <https://doi.org/10.1097/MD.00000000000020109>
- Hersh, M. (2015). Cane use and late onset visual impairment. *Technology and Disability*, 27(3), 103–116. <https://doi.org/10.3233/TAD-150432>
- Hill, J., & Black, J. (2003). The miniguide: A new electronic travel device. *Journal of Visual Impairment & Blindness*, 97(10), 1–6. <https://doi.org/10.1177/0145482X0309701010>
- Horne, L., Alvarez, J. M., McCarthy, C., & Barnes, N. (2015, August). Semantic labelling to aid navigation in prosthetic vision. In *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society* (pp. 3379–3382). IEEE. <https://doi.org/10.1109/EMBC.2015.7319117>

- Illinois Library. (2020, March 10). *Blind/visual impairment: Common assistive technologies*. University Library. <https://guides.library.illinois.edu/c.php?g=526852&p=3602299>
- Jicol, C., Lloyd-Esenkaya, T., Proulx, M. J., Lange-Smith, S., Scheller, M., O'Neill, E., & Petrini, K. (2020). Efficiency of sensory substitution devices alone and in combination with self-motion for spatial navigation in sighted and visually impaired. *Frontiers in Psychology, 11*, Article 1443. <https://doi.org/10.3389/fpsyg.2020.01443>
- Kastrup, V., Sampaio, E., Almeida, M. C. D., & Carijó, F. H. (2009). The learning of the use of the tactile-vision sensory substitution by people with visual disability: First experiences and methodological strategies. *Psicologia & Sociedade, 21*(2), 256–265. <https://doi.org/10.1590/S0102-71822009000200013>
- Kaye, H. S., Yeager, P., & Reed, M. (2008). Disparities in usage of assistive technology among people with disabilities. *Assistive Technology, 20*(4), 194–203. <https://doi.org/10.1080/10400435.2008.10131946>
- Koumpouros, Y., Papageorgiou, E., Karavasili, A., & Alexopoulou, D. (2017). Translation and validation of the assistive technology device predisposition assessment in Greek in order to assess satisfaction with use of the selected assistive device. *Disability and Rehabilitation: Assistive Technology, 12*(5), 535–542. <https://doi.org/10.3109/17483107.2016.1161088>
- Kugler, L. (2020). Technologies for the visually impaired. *Communications of the ACM, 63*(12), 15–17. <https://doi.org/10.1145/3427936>
- Lenay, C., Gapenne, O., Hanne-ton, S., Marque, C., & Genouëlle, C. (2003). Sensory substitution: Limits and perspectives. In Y. Hatwell, A. Streri, & E. Gentaz (Eds.), *Touching for knowing: Cognitive psychology of haptic manual perception* (pp. 276–292). John Benjamins Publishing Company. <https://doi.org/10.1075/aicr.53.22len>
- Lewis, P. M., Ackland, H. M., Lowery, A. J., & Rosenfeld, J. V. (2015). Restoration of vision in blind individuals using bionic devices: A review with a focus on cortical visual prostheses. *Brain Research, 1595*, 51–73. <https://doi.org/10.1016/j.brainres.2014.11.020>
- Lloyd-Esenkaya, T., Lloyd-Esenkaya, V., O'Neill, E., & Proulx, M. J. (2020). Multisensory inclusive design with sensory substitution. *Cognitive Research: Principles and Implications, 5*(1), 1–15. <https://doi.org/10.1186/s41235-020-00240-7>
- Lüchtenberg, M., Kuhli-Hattenbach, C., Sinangin, Y., Ohrloff, C., & Schalnus, R. (2008). Accessibility of health information on the internet to the visually impaired user. *Ophthalmologica, 222*(3), 187–193. <https://doi.org/10.1159/000126082>
- Luo, Y. H. L., & Da Cruz, L. (2016). The Argus® II retinal prosthesis system. *Progress in Retinal and Eye Research, 50*, 89–107. <https://doi.org/10.1016/j.preteyeres.2015.09.003>
- Luo, Y. H. L., Zhong, J., Merlini, F., Anafloos, F., Arsiero, M., Stanga, P. E., & Da Cruz, L. (2014). The use of Argus® II retinal prosthesis to identify common objects in blind subjects with outer retinal dystrophies. *Investigative Ophthalmology & Visual Science, 55*(13), 1834–1834.
- Manduchi, R., & Coughlan, J. (2012). (Computer) vision without sight. *Communications of the ACM, 55*(1), 96–104. <https://doi.org/10.1145/2063176.2063200>
- Manjari, K., Verma, M., & Singal, G. (2020). A survey on assistive technology for visually impaired. *Internet of Things, 11*, 100188. <https://doi.org/10.1016/j.iot.2020.100188>
- Medicines & Healthcare products Regulatory Agency. (2021). *Assistive technology: Definition and safe use*. <https://www.gov.uk/government/publications/assistive-technology-definition-and-safe-use/assistive-technology-definition-and-safe-use#definitions>
- Meijer, P. B. (1992). An experimental system for auditory image representations. *IEEE Transactions on Biomedical Engineering, 39*(2), 112–121. <https://doi.org/10.1109/10.121642>
- Microsoft. (2021). *Seeing AI app from Microsoft*. <https://www.microsoft.com/en-us/ai/seeing-ai>
- Nadler, J. T., Weston, R., & Voyles, E. C. (2015). Stuck in the middle: The use and interpretation of mid-points in items on questionnaires. *The Journal of General Psychology, 142*(2), 71–89. <https://doi.org/10.1080/00221309.2014.994590>
- National Health Service. (2018). *Blindness and vision loss*. <https://www.nhs.uk/conditions/vision-loss/>
- Okonji, P. E., & Ogwezy, D. C. (2019). Awareness and barriers to adoption of assistive technologies among visually impaired people in Nigeria. *Assistive Technology, 31*(4), 209–219. <https://doi.org/10.1080/10400435.2017.1421594>

- Parikh, N., Shah, I., & Vahora, S. (2018, April). Android smartphone based visual object recognition for visually impaired using deep learning. In *2018 International Conference on Communication and Signal Processing* (pp. 420–425). IEEE. <https://doi.org/10.1109/ICCSP.2018.8524493>
- Phillips, M., & Proulx, M. J. (2018). Social interaction without vision: An assessment of assistive technology for the visually impaired. *Technology & Innovation, 20*(1–2), 85–93. <https://doi.org/10.21300/20.1-2.2018.85>
- Pixium Vision. (2016). *Pixium Vision implants IRIS® II, first epi-retinal implant with 150 electrodes*. <https://www.pixium-vision.com/2016/02/pixium-vision-implants-iris-ii-first-epi-retinal-implant-with-150-electrodes/>
- Proulx, M. J., Stoerig, P., Ludowig, E., & Knoll, I. (2008). Seeing ‘where’ through the ears: Effects of learning by-doing and long-term sensory deprivation on localization based on image-to-sound substitution. *PLOS ONE, 3*(3), Article e1840. <https://doi.org/10.1371/journal.pone.0001840>
- Qualtrics. (2021). *Online survey software*. <https://www.qualtrics.com/uk/core-xm/survey-software/>
- Quinones, P-A., Greene, T., Yang, R., & Newman, M. (2011). Supporting visually impaired navigation: A needs-finding study. In *CHI '11 Extended Abstracts on Human Factors in Computing Systems (CHI EA '11)* (pp. 1645–1650). Association for Computing Machinery. <https://doi.org/10.1145/1979742.1979822>
- Qureshi, T. A., Rajbhar, M., Pisat, Y., & Bhosale, V. (2021). AI based app for blind people. *International Research Journal of Engineering and Technology, 8*(3), 2883–2887.
- Roentgen, U. R., Gelderblom, G. J., & de Witte, L. P. (2012). User evaluation of two electronic mobility aids for persons who are visually impaired: A quasi-experimental study using a standardized mobility course. *Assistive Technology, 24*(2), 110–120. <https://doi.org/10.1080/10400435.2012.659794>
- Sabel, B. A., Thut, G., Haueisen, J., Henrich-Noack, P., Herrmann, C. S., Hunold, A., Kammer, T., Matteo, B., Sergeeva, E. G., Waleszczyk, W., & Antal, A. (2020). Vision modulation, plasticity and restoration using non-invasive brain stimulation—An IFCN-sponsored review. *Clinical Neurophysiology, 131*(4), 887–911. <https://doi.org/10.1016/j.clinph.2020.01.008>
- Sahib, N. G., Tombros, A., & Stockman, T. (2012). A comparative analysis of the information-seeking behavior of visually impaired and sighted searchers. *Journal of the American Society for Information Science and Technology, 63*(2), 377–391. <https://doi.org/10.1002/asi.21696>
- Salido, J., Deniz, O., & Bueno, G. (2016, April). Sainet: An image processing app for assistance of visually impaired people in social interaction scenarios. In *International Conference on Bioinformatics and Biomedical Engineering* (pp. 467–477). Springer. https://doi.org/10.1007/978-3-319-31744-1_42
- Scheller, M., Petrini, K., & Proulx, M. J. (2018). Perception and interactive technology. *Stevens' Handbook of Experimental Psychology and Cognitive Neuroscience, 2*, 1–50. <https://doi.org/10.1002/9781119170174.epcn215>
- Scherer, M. J. (2005). Assessing the benefits of using assistive technologies and other supports for thinking, remembering and learning. *Disability and Rehabilitation, 27*(13), 731–739. <https://doi.org/10.1080/09638280400014816>
- Senjam, S. S. (2021). Smartphones as assistive technology for visual impairment. *Eye, 35*, 2078–2080. <https://doi.org/10.1038/s41433-021-01499-w>
- Senjam, S. S., Foster, A., Bascaran, C., Vashist, P., & Gupta, V. (2020). Assistive technology for students with visual disability in schools for the blind in Delhi. *Disability and Rehabilitation: Assistive Technology, 15*(6), 663–669. <https://doi.org/10.1080/17483107.2019.1604829>
- Sharts-Hopko, N. C., Smeltzer, S., Ott, B. B., Zimmerman, V., & Duffin, J. (2010). Healthcare experiences of women with visual impairment. *Clinical Nurse Specialist, 24*(3), 149–153. <https://doi.org/10.1097/nur.0b013e3181d82b89>
- Shimojo, S., & Shams, L. (2001). Sensory modalities are not separate modalities: Plasticity and interactions. *Current Opinion in Neurobiology, 11*(4), 505–509. [https://doi.org/10.1016/S0959-4388\(00\)00241-5](https://doi.org/10.1016/S0959-4388(00)00241-5)
- Solomon-Harris, L. M., Mullin, C. R., & Steeves, J. K. (2013). TMS to the ‘occipital face area’ affects recognition but not categorization of faces. *Brain and Cognition, 83*(3), 245–251.
- Stingl, K., Bartz-Schmidt, K. U., Besch, D., Braun, A., Bruckmann, A., Gekeler, F., . . . Zrenner, E. (2013). Artificial vision with wirelessly powered subretinal electronic implant alpha-IMS. *Proceedings of the Royal Society B: Biological Sciences, 280*(1757), 20130077. <https://doi.org/10.1098/rspb.2013.0077>

- Strong, P. (2009). *The history of the white cane*. Paths to Literacy. <https://www.pathstoliteracy.org/wp-content/uploads/2022/06/The-History-of-the-White-Cane.docx>
- Upson, S. (2007). *Loser: Tongue vision*. IEEE Spectrum. <https://spectrum.ieee.org/loser-tongue-vision>
- Vailshery, L. S. (2021). *Ownership of Smartphones in the United Kingdom (UK) 2008, 2019-2020*. Statista. <https://www.statista.com/statistics/956297/ownership-of-smartphones-uk/>
- Varghese, J. E. (1996). Independence to the blind and handicapped in Asia through modern assistive devices. *Computers in Human Services*, 12(1–2), 5–12. https://doi.org/10.1300/J407v12n01_02
- Ward, J., & Wright, T. (2014). Sensory substitution as an artificially acquired synaesthesia. *Neuroscience & Biobehavioral Reviews*, 41, 26–35. <https://doi.org/10.1016/j.neubiorev.2012.07.007>
- West, S. K., Rubin, G. S., Broman, A. T., Munoz, B., Bandeen-Roche, K., & Turano, K. (2002). How does visual impairment affect performance on tasks of everyday life? The SEE Project. *Archives of Ophthalmology*, 120(6), 774–780. <https://doi.org/10.1001/archophth.120.6.774>
- Wicab Inc. (2010). *BrainPort (registered trademark) technology tongue interface characterization*.
- Withers, J., & Speight, C. (2017). Health care for individuals with hearing loss or vision loss+2 minefield of barriers to accessibility. *North Carolina Medical Journal*, 78(2), 107–112. <https://doi.org/10.18043/ncm.78.2.107>
- World Health Organization. (2021, February 26). *Blindness and vision impairment*. <https://www.who.int/news-room/fact-sheets/detail/blindness-and-visual-impairment>
- Xie, I., Babu, R., Castillo, M. D., & Han, H. (2018). Identification of factors associated with blind users' help-seeking situations in interacting with digital libraries. *Journal of the Association for Information Science and Technology*, 69(4), 514–527. <https://doi.org/10.1002/asi.23982>