

# Planar graph coloring avoiding monochromatic subgraphs: trees and paths make it difficult \*

*Hajo Broersma*  
University of Durham, UK  
hajo.broersma@durham.ac.uk

*Fedor V. Fomin*  
University of Bergen, Norway  
fomin@ii.uib.no

*Jan Kratochvíl*  
Charles University, Czech Republic  
honza@kam.ms.mff.cuni.cz

*Gerhard J. Woeginger*  
Eindhoven University of Technology, The Netherlands  
gwoegi@win.tue.nl

## Abstract

We consider the problem of coloring a planar graph with the minimum number of colors so that each color class avoids one or more forbidden graphs as subgraphs. We perform a detailed study of the computational complexity of this problem.

We present a complete picture for the case with a single forbidden connected (induced or non-induced) subgraph. The 2-coloring problem is NP-hard if the forbidden subgraph is a tree with at least two edges, and it is polynomially solvable in all other cases. The 3-coloring problem is NP-hard if the forbidden subgraph is a path with at least one edge, and it is polynomially solvable in all other cases. We also derive results for several forbidden sets of cycles. In particular, we prove that it is NP-complete to decide if a planar graph can be 2-colored so that no cycle of length at most 5 is monochromatic.

**Keywords:** graph coloring; graph partitioning; forbidden subgraph; planar graph; computational complexity.

**AMS Subject Classifications:** 05C15,05C85,05C17

---

\*An extended abstract was presented at SWAT2002 [5]. The work of HB and FVF is sponsored by NWO-grant 047.008.006. Part of the work was done while FVF a visiting postdoc at DIMATIA-ITI (supported by GAČR 201/99/0242 and by the Ministry of Education of the Czech Republic as project LN00A056). JK acknowledges support by the Czech Ministry of Education as project LN00A056. GJW acknowledges support by the START program Y43-MAT of the Austrian Ministry of Science.

# 1 Introduction

We denote by  $G = (V, E)$  a finite undirected and simple graph with  $|V| = n$  vertices and  $|E| = m$  edges. For any non-empty subset  $W \subseteq V$ , the subgraph of  $G$  induced by  $W$  is denoted by  $G[W]$ . A *clique* of  $G$  is a non-empty subset  $C \subseteq V$  such that all the vertices of  $C$  are mutually adjacent. A non-empty subset  $I \subseteq V$  is *independent* if no two of its elements are adjacent. An  $r$ -*coloring* of the vertices of  $G$  is a partition  $V_1, V_2, \dots, V_r$  of  $V$ ; the  $r$  sets  $V_j$  are called the *color classes* of the  $r$ -coloring. An  $r$ -coloring is *proper* if every color class is an independent set. The *chromatic number*  $\chi(G)$  is the minimum integer  $r$  for which a proper  $r$ -coloring of  $G$  exists.

Evidently, an  $r$ -coloring is proper if and only if for every color class  $V_j$ , the induced subgraph  $G[V_j]$  does not contain a subgraph isomorphic to  $P_2$ . (We use  $P_k$  to denote the path on  $k$  vertices.) This observation leads to a number of interesting generalizations of the classical graph coloring concept. One such generalization was suggested by Harary [24]: Given a graph property  $\pi$ , a positive integer  $r$ , and a graph  $G$ , a  $\pi$   $r$ -coloring of  $G$  is a (not necessarily proper)  $r$ -coloring in which each subgraph induced by a color class has property  $\pi$ . This generalization has been studied for the cases where the graph property  $\pi$  is being acyclic, or planar, or perfect, or a path of length at most  $k$ , or a clique of size at most  $k$ . We refer the reader to the work of Brown & Corneil [8, 7, 9], Chartrand *et al.* [11, 12, 13], Farrigua [16] and Sachs [29] for more information on these variants.

In this paper, we will investigate graph colorings where the property  $\pi$  can be defined via some (maybe infinite) list of forbidden induced subgraphs. This naturally leads to the notion of  $\mathcal{F}$ -free colorings. Let  $\mathcal{F} = \{F_1, F_2, \dots\}$  be the set of so-called forbidden graphs. Throughout the paper we will assume that the set  $\mathcal{F}$  is non-empty, and that all graphs in  $\mathcal{F}$  are connected and contain at least one edge. Moreover, to avoid technical difficulties in the proofs we will assume that no graph of  $\mathcal{F}$  is a proper subgraph of another graph of  $\mathcal{F}$ . For a graph  $G$ , a (not necessarily proper)  $r$ -coloring with color classes  $V_1, V_2, \dots, V_r$  is called *weakly  $\mathcal{F}$ -free*, if for all  $1 \leq j \leq r$ , the graph  $G[V_j]$  does not contain any graph from  $\mathcal{F}$  as an *induced* subgraph. Similarly, we say that an  $r$ -coloring is *strongly  $\mathcal{F}$ -free* if  $G[V_j]$  does not contain any graph from  $\mathcal{F}$  as an (induced or non-induced) subgraph. The smallest possible number of colors in a weakly (respectively, strongly)  $\mathcal{F}$ -free coloring of a graph  $G$  is called the *weakly* (respectively, *strongly*)  *$\mathcal{F}$ -free chromatic number*; it is denoted by  $\chi^w(\mathcal{F}, G)$  (respectively, by  $\chi^s(\mathcal{F}, G)$ ).

In the cases where  $\mathcal{F} = \{F\}$  consists of a single graph  $F$ , we will sometimes simplify the notation and not write the curly brackets: We will write  $F$ -free short for  $\{F\}$ -free,  $\chi^w(F, G)$  short for  $\chi^w(\{F\}, G)$ , and  $\chi^s(F, G)$  short for  $\chi^s(\{F\}, G)$ . With this notation  $\chi(G) = \chi^s(P_2, G) = \chi^w(P_2, G)$  holds for every graph  $G$ , and hence also

$$\chi^w(\mathcal{F}, G) \leq \chi^s(\mathcal{F}, G) \leq \chi(G).$$

It is easy to construct examples where both inequalities are strict. For instance, for  $\mathcal{F} = \{P_3\}$  (the path on three vertices) and  $G = C_3$  (the cycle on three vertices) we have  $\chi(G) = 3$ ,  $\chi^s(P_3, G) = 2$ , and  $\chi^w(P_3, G) = 1$ .

Our main concern in the paper are planar graphs. Recall that a graph is *planar* if it can be drawn in the (Euclidean) plane without intersections of edges. Such a drawing is

referred to as a *plane* graph. Hence a graph  $G$  is planar if and only if there exists a plane graph isomorphic to  $G$ . A planar graph is called *outerplanar* if it has a drawing such that all vertices lie on the boundary of the unbounded face (this face is usually referred to as the outer face).

## 1.1 Previous results

The literature contains quite a number of papers on weakly and strongly  $\mathcal{F}$ -free colorings of graphs. One of the most general results is due to Achlioptas [1]: For any graph  $F$  with at least three vertices and for any  $r \geq 2$ , the problem of deciding whether a given input graph has a weakly  $F$ -free  $r$ -coloring is NP-hard. We often use weakly (strongly)  $\mathcal{F}$ -free  $r$ -coloring as shorthand for the corresponding decision problem.

The special case of weakly  $P_3$ -free coloring is known as the *subcoloring problem* in the literature. It has been studied by Broere & Mynhardt [4], by Albertson, Jamison, Hedetniemi & Locke [2], by Fiala, Jansen, Le & Seidel [18], Gimbel & Hartman [21], and by Broersma, Fomin, Nešetřil & Woeginger [6]. We will further utilize especially the following result:

**Proposition 1.1.** [Fiala, Jansen, Le & Seidel [18]]

*Weakly  $P_3$ -free 2-coloring is NP-hard for triangle-free planar graphs.*

A  $(1, 2)$ -subcoloring of  $G$  is a partition of  $V$  into two sets  $S_1$  and  $S_2$  such that  $S_1$  induces an independent set and  $S_2$  induces a subgraph consisting of a matching and some (possibly no) isolated vertices. Le & Le [27] proved that recognizing if a graph is  $(1, 2)$ -subcolorable is NP-hard even for cubic triangle-free planar graphs.

The case of weakly  $P_4$ -free coloring has been investigated by Gimbel & Nešetřil [22] who study the problem of partitioning the vertex set of a graph into induced cographs. Since cographs are exactly the graphs without an induced  $P_4$ , the graph parameter studied in [22] equals the weakly  $P_4$ -free chromatic number of a graph. In [22] it is proved that the problems of deciding  $\chi^w(P_4, G) \leq 2$ ,  $\chi^w(P_4, G) = 3$ ,  $\chi^w(P_4, G) \leq 3$  and  $\chi^w(P_4, G) = 4$  all are NP-hard and/or coNP-hard for planar graphs. The work of Hoàng & Le [25] on weakly  $P_4$ -free 2-colorings was motivated by the Strong Perfect Graph Conjecture. Among other results, they show that weakly  $P_4$ -free 2-coloring is NP-hard for comparability graphs.

A notion that is closely related to strongly  $F$ -free  $r$ -coloring is the so-called *defective* graph coloring. A defective  $(k, d)$ -coloring of a graph is a  $k$ -coloring in which each color class induces a subgraph with maximum degree at most  $d$ . Defective colorings have been studied for example by Archdeacon [3], by Cowen, Cowen & Woodall [14], and by Frick & Henning [19]. Cowen, Goddard & Jesurum [15] have shown that the defective  $(3, 1)$ -coloring problem and the defective  $(2, d)$ -coloring problem for any  $d \geq 1$  are NP-hard even for planar graphs. We observe that for any  $k$ , defective  $(k, 1)$ -coloring is equivalent to strongly  $P_3$ -free  $k$ -coloring, and hence we derive the following proposition.

**Proposition 1.2.** [Cowen, Goddard & Jesurum [15]]

(i) *Strongly  $P_3$ -free 2-coloring is NP-hard for planar graphs.*

(ii) *Strongly  $P_3$ -free 3-coloring is NP-hard for planar graphs.*

## 1.2 Our results

We perform a complexity study of weakly and strongly  $\mathcal{F}$ -free coloring problems for *planar* graphs. By the Four Color Theorem, every planar graph  $G$  satisfies  $\chi(G) \leq 4$ . Consequently, every planar graph also satisfies  $\chi^w(\mathcal{F}, G) \leq 4$  and  $\chi^s(\mathcal{F}, G) \leq 4$ , and so we may concentrate on 2-colorings and on 3-colorings. For the case of a single forbidden subgraph, we obtain the following results for 2-colorings:

- If the forbidden (connected) subgraph  $F$  is not a tree, then *every* planar graph is strongly and hence also weakly  $F$ -free 2-colorable. Therefore, the corresponding decision problems are trivially solvable.
- If the forbidden subgraph  $F = P_2$ , then  $F$ -free 2-coloring is equivalent to proper 2-coloring. It is well-known that this problem is polynomially solvable.
- If the forbidden subgraph is a tree  $T$  with at least two edges, then both weakly and strongly  $T$ -free 2-colorings are NP-hard for planar graphs. Hence, these problems are intractable.

For 3-colorings with a single forbidden subgraph, we obtain the following results:

- If the forbidden (connected) subgraph  $F$  is not a path, then *every* planar graph is strongly and hence also weakly  $F$ -free 3-colorable. Hence, the corresponding decision problems are trivially solvable.
- For every path  $P$  with at least one edge, both weakly and strongly  $P$ -free 3-colorings are NP-hard for planar graphs. Hence, these problems are intractable.

Moreover, we derive several results for 2-colorings with certain forbidden sets of cycles.

- For the forbidden set  $\mathcal{F}_{345} = \{C_3, C_4, C_5\}$ , both weakly and strongly  $\mathcal{F}_{345}$ -free 2-colorings are NP-hard for planar graphs. In fact for any finite set  $\mathcal{F}_{\geq 345} \supseteq \{C_3, C_4, C_5\}$  of cycles, both weakly and strongly  $\mathcal{F}_{\geq 345}$ -free 2-colorings are NP-hard for planar input graphs.
- Also for the forbidden set  $\mathcal{F}_{cycle}$  of all cycles, both weakly and strongly  $\mathcal{F}_{cycle}$ -free 2-colorings are NP-hard for planar graphs.
- For the forbidden set  $\mathcal{F}_{odd}$  of all cycles of odd lengths, *every* planar graph is strongly and hence also weakly  $\mathcal{F}_{odd}$ -free 2-colorable. This follows from (in fact, it is equivalent to) the Four Color Theorem.

## 2 The machinery for establishing NP-hardness

Throughout this section, let  $\mathcal{F}$  denote some fixed set of forbidden planar subgraphs. We assume that all graphs in  $\mathcal{F}$  are connected and contain at least two edges. We also assume that no graph of  $\mathcal{F}$  is a (not necessarily induced) proper subgraph of another graph from  $\mathcal{F}$ . We will develop a generic NP-hardness proof for certain types of weakly and strongly  $\mathcal{F}$ -free 2-coloring problems. The crucial concept is the so-called *equalizer* gadget. Before we

define this gadget, let us introduce the following technical concept of crossing graphs. We note that we distinguish between planar graphs and plane graphs (the latter being particular nonintersecting drawings of abstract planar graphs), but we use the same notation for a plane graph and its underlying abstract (planar) graph. When talking about more than one graph, we use subscripts to distinguish their vertex and edge sets (i.e.,  $V_G$  and  $E_G$  denote the vertex and edge sets of a graph  $G$ ).

**Definition 2.1.** Given a plane graph  $G$  with outer face  $C$  and a set  $S \subseteq V_G$  of vertices on the boundary of  $C$  (referred to as *contact points*), we say that another plane graph  $H$  is *crossing*  $G$  if the following assertions hold:

1.  $G \cup H$  is a plane graph (i.e., no edge of  $G$  crosses any edge of  $H$  in the simultaneous drawing of  $G$  and  $H$ ),
2. all edges of  $E_H \setminus E_G$  are drawn in  $C$ ,
3. no edge of  $E_H \setminus E_G$  is incident with a vertex of  $V_G \setminus S$ ,
4.  $V_H \cap (V_G \setminus S) \neq \emptyset$ .

If  $G$  is a plane graph with a set  $S$  of contact points, we say that a planar graph  $H$  *may cross*  $G$  if some nonintersecting planar drawing of a graph isomorphic to  $H$  is crossing  $G$ .

In the left half of Figure 1 the graph  $H$  induced by the vertices  $b_2, c$  and  $d$  is crossing the graph  $G$  induced by the vertices  $a, b_1, b_2, b_3$  and  $c$  with  $S = \{c\}$ .

**Definition 2.2.** (Equalizer)

An  $(a, b)$ -*equalizer* for  $\mathcal{F}$  is a plane graph  $\mathcal{E}$  with two nonadjacent contact points  $a$  and  $b$  on the boundary of the outer face, which satisfies the following properties:

- (i) In every weakly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{E}$ ,  $a$  and  $b$  receive the same color.
- (ii) There exists a strongly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{E}$  such that  $a$  and  $b$  receive the same color, whereas no monochromatic copy of a graph in  $\mathcal{F}$  may cross  $\mathcal{E}$ . Such a coloring is called a *good* 2-coloring of  $\mathcal{E}$ .

The graph  $\mathcal{E}$  in the right half of Figure 1 is an  $(a, b)$ -equalizer for  $P_3$ . In every weakly  $P_3$ -free 2-coloring of  $\mathcal{E}$  the vertices  $a$  and  $b$  should receive the same color; otherwise a monochromatic  $P_3$  is unavoidable if we extend the 2-coloring. A good coloring of  $\mathcal{E}$  can be obtained by assigning  $a$  and  $b$  the same color and all remaining vertices of  $\mathcal{E}$  the other color.

To better understand the definition of an equalizer  $\mathcal{E}$ , let us remark right away that if  $\mathcal{F}$  contains a graph with a leaf, then an  $\mathcal{F}$ -free 2-coloring of  $\mathcal{E}$  is good if and only if all neighbors of the contact points in  $\mathcal{E}$  have the color that is not assigned to the contact points.

The rest of this section is devoted to the proof of the following (technical) main theorem. This theorem is going to generate a number of NP-hardness statements in the subsequent sections of the paper.

**Theorem 2.3.** *Let  $\mathcal{F}$  be a set of connected planar graphs that all contain at least two edges, such that no graph of  $\mathcal{F}$  is a proper subgraph of another graph of  $\mathcal{F}$ . Suppose that*




Figure 1: Examples to illustrate Definitions 2.1 and 2.2.

- $\mathcal{F}$  contains a graph on at least four vertices with a cut vertex, or a 2-connected graph with a planar embedding with at least five vertices on the boundary of the outer face;
- there exists an  $(a, b)$ -equalizer for  $\mathcal{F}$ .

Then deciding weakly  $\mathcal{F}$ -free 2-colorability and deciding strongly  $\mathcal{F}$ -free 2-colorability are NP-hard problems for planar input graphs.

We postpone the proof of Theorem 2.3 to Section 2.2, but first introduce some additional tools.

## 2.1 Gadgets for the NP-hardness proof

We will design a series of gadgets that all use the equalizer gadget as an atomic component. In all constructions, the only connections between an equalizer and the rest of the constructed graph will always be via the contact points. The use of the equalizer gadget is justified (and motivated) by the following lemma.

**Lemma 2.4.** *Consider a nontrivial planar graph  $H$  and an edge  $xy \in E_H$ . Let the graph  $H^+$  result from  $H$  by adding a vertex-disjoint copy  $\mathcal{E}$  of an  $(a, b)$ -equalizer to  $H$  and then identifying vertex  $x$  with contact point  $a$ , and vertex  $y$  with contact point  $b$ .*

*Then  $H^+$  is a planar graph, and  $H^+$  has a strongly/weakly  $\mathcal{F}$ -free 2-coloring if and only if  $H$  has a strongly/weakly  $\mathcal{F}$ -free 2-coloring in which  $x$  and  $y$  both receive the same color.*

*Proof.* Since  $\mathcal{E}$  is a plane graph with  $a$  and  $b$  on the boundary of the outer face,  $H^+$  is also planar and it has a nonintersecting drawing such that all edges of  $H$  are drawn in the outer face of  $\mathcal{E}$ . For the proof of the ‘only if’ part, observe that every strongly/weakly  $\mathcal{F}$ -free 2-coloring of  $H^+$  induces a strongly/weakly  $\mathcal{F}$ -free 2-coloring of  $H$ . By property (i) in Definition 2.2, this induced coloring assigns the same color to  $x$  and  $y$ . For the proof of the ‘if’ part, we construct a strongly/weakly  $\mathcal{F}$ -free 2-coloring of  $H^+$ : We use the strongly/weakly  $\mathcal{F}$ -free 2-coloring for the subgraph  $H$ , and we color the  $(a, b)$ -equalizer  $\mathcal{E}$  using a good coloring in the sense of property (ii) in Definition 2.2.  $\square$

**The negator gadget.** An  $(a, b)$ -negator for  $\mathcal{F}$  is a plane graph  $\mathcal{N}$  with two nonadjacent contact points  $a$  and  $b$  on the boundary of the outer face, which satisfies the following properties:

- (i) In every weakly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{N}$ ,  $a$  and  $b$  receive different colors.
- (ii) There exists a strongly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{N}$  such that  $a$  and  $b$  receive different colors, whereas no monochromatic copy of a graph in  $\mathcal{F}$  may cross  $\mathcal{N}$ . Such a coloring is called a *good* 2-coloring of  $\mathcal{N}$ .

Next we show how to construct such an  $(a, b)$ -negator from  $(a, b)$ -equalizers. We choose an arbitrary graph  $F \in \mathcal{F}$ , and take some fixed planar embedding of  $F$  to form the so-called *skeleton* of the negator. Let  $a'$  and  $b'$  denote two vertices on the boundary of the outer face of  $F$ . We partition  $V_F$  into two disjoint sets  $V_1$  and  $V_2$  in such a way that both  $F[V_1]$  and  $F[V_2]$  (the subgraphs of  $F$  induced by  $V_1$  and  $V_2$ ) are connected, and so that  $a' \in V_1$  and  $b' \in V_2$ . For every edge  $xy \in E_{F[V_1]} \cup E_{F[V_2]}$ , we add an equalizer between  $x$  and  $y$  exactly in the way we described in Lemma 2.4. We introduce a new vertex  $a$  and connect it by an equalizer to  $a'$ ; we create a new vertex  $b$  and connect it by an equalizer to  $b'$ . This completes the construction of  $\mathcal{N}$ . To see that (i) is fulfilled, consider some weakly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{N}$ . Suppose that  $a$  and  $b$  receive the same color. Then the equalizers enforce that this color propagates to all vertices in the skeleton, and this yields a monochromatic induced copy of  $F$ , a contradiction. To see that (ii) is fulfilled, we may color  $\{a\} \cup V_1$  with one color, and  $\{b\} \cup V_2$  with the other color. The vertices inside the equalizer gadgets may be colored using a good coloring in the sense of Definition 2.2.(ii). Any monochromatic copy of a graph  $F' \in \mathcal{F}$  would either contain some edges of some equalizer gadget (which is impossible by the goodness of the equalizer coloring) or be a subgraph of  $F[V_1]$  or  $F[V_2]$  (which is impossible by the assumption we made on  $\mathcal{F}$ ).

In our constructions, the negator gadget will be used similarly as the equalizer gadget as described in Lemma 2.4. While the equalizer gadget can be used to enforce that a pair of vertices receives the same color, with the help of the negator gadget we can enforce that a pair of adjacent vertices in some planar graph must receive different colors in any weakly  $\mathcal{F}$ -free 2-coloring. We omit the details since the counterpart of Lemma 2.4 with respect to negators and its proof are straightforward variations on Lemma 2.4 and its proof.

For our NP-hardness proof (of Theorem 2.3) we need two additional gadgets.

**The clause gadget with four contact points.** The gadget  $\mathcal{C}_4(a, b, c, d)$  is a plane graph  $\mathcal{C}$  with pairwise nonadjacent contact points  $a, b, c$  and  $d$  that lie in this (cyclic) ordering on the boundary of the outer face of  $\mathcal{C}$ . It has the following properties:

- (i) In every weakly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{C}$ , not all four contact points receive the same color.
- (ii) Any 2-coloring of the four contact points that uses both colors, can be extended to a strongly  $\mathcal{F}$ -free 2-coloring of the gadget  $\mathcal{C}$ , in such a way that no monochromatic copy of a graph in  $\mathcal{F}$  may cross  $\mathcal{C}$ . Such a coloring is called a *good* 2-coloring of  $\mathcal{C}$ .

Now let us construct such a clause gadget  $\mathcal{C}_4(a, b, c, d)$ . Suppose we are in the case assumed in Theorem 2.3. Hence the set  $\mathcal{F}$  contains some graph  $F$  that can be planarly embedded such that there are four vertices  $a', b', c', d'$  on the boundary of the outer face. We choose this plane graph  $F$  to form the skeleton of the clause gadget. We create four new vertices  $a, b, c$ , and  $d$ . Each of these new vertices is connected by an equalizer to its corresponding primed vertex on the outer face of the skeleton. The vertices in the skeleton are partitioned into four components (with connecting edges between them) such that  $a', b', c', d'$  end up in different components. Within each component, we introduce equalizers along every edge in the way we described in Lemma 2.4. This completes the construction.

By now it is routine to verify that the construction indeed fulfills the two properties (i) and (ii). We leave the details to the reader.

**The clause gadget with five contact points.** The gadget  $\mathcal{C}_5(a, b, c, d_1, d_2)$  is a plane graph  $\mathcal{C}$  with pairwise nonadjacent contact points  $a, b, c, d_1$ , and  $d_2$  that lie in the (cyclic) ordering  $a - b - d_1 - c - d_2$  on the boundary of the outer face of  $\mathcal{C}$ . It has the following properties:

- (i) In every weakly  $\mathcal{F}$ -free 2-coloring of  $\mathcal{C}$ , the vertices  $d_1$  and  $d_2$  receive the same color, and at least one of the vertices  $a, b, c$  receives the opposite color.
- (ii) Any 2-coloring of the five contact points that assigns the same color to  $d_1$  and  $d_2$ , and the opposite color to at least one of  $a, b, c$ , can be extended to a strongly  $\mathcal{F}$ -free 2-coloring of the gadget  $\mathcal{C}$ , in such a way that no monochromatic copy of a graph in  $\mathcal{F}$  may cross  $\mathcal{C}$ . Such a coloring is called a *good* 2-coloring of  $\mathcal{C}$ .

Suppose we are in the case assumed in Theorem 2.3, hence the set  $\mathcal{F}$  contains a graph on at least four vertices with a cut vertex, or a 2-connected graph with a planar embedding with at least five vertices on the boundary of the outer face.

Let us first discuss the case of a graph  $F \in \mathcal{F}$  with a cut vertex  $d'$ . The skeleton of  $\mathcal{C}_5(a, b, c, d_1, d_2)$  is formed by a planar embedding of  $F$  where  $d'$  is on the boundary of the outer face. Choose three vertices  $a', b', c'$  that all lie on the boundary of the outer face, and that do not belong to the same component of  $F - d'$ , such that we can move around the boundary of the outer face starting at  $a'$ , then moving to  $b'$ , then to  $d'$ , then to  $c'$ , then to  $d'$  again, and then returning to  $a'$  (maybe meeting other vertices, including  $d'$  and  $b'$ , in between). For example, if  $F = K_{1,k}$  is a star with  $k \geq 3$  leaves, we choose  $d'$  as the center, and  $a', b'$  and  $c'$  as three successive end vertices in a cyclic ordering in a planar embedding of  $F$ . We can move around the boundary of the outer face from  $a'$  (via  $d'$ ) to  $b'$ , then to  $d'$  and to  $c'$ , and back to  $a'$  via  $d'$  (and alternating between  $d'$  and the possible other end vertices if  $k \geq 3$ ). The other cases are similar. We create five new vertices  $a, b, c, d_1$ , and  $d_2$ , and we connect them by equalizers to  $a', b', c', d'$ , and  $d'$ , respectively, at the place where we hit the primed vertices in the above ordering  $a' - b' - d' - c' - d'$  while moving around the boundary of the outer face in the way we described. The vertices in the skeleton are partitioned into four components such that  $a', b', c', d'$  end up in different components. Within each component, we introduce equalizers along every edge in the way we described in Lemma 2.4. This completes the construction for the first case.

Next we discuss the case of a 2-connected planar graph  $F \in \mathcal{F}$  that has a planar embedding with at least five vertices on the boundary of the outer face. We use such an embedding as the skeleton of  $\mathcal{C}_5(a, b, c, d_1, d_2)$ . Consider the cycle  $C$  that forms the boundary of the outer face. Choose five vertices  $v_0 - v_1 - v_2 - v_3 - v_4$  in this order along  $C$ . Because all these  $v$ -vertices are on the outer face, only two subcases are possible.

- (Subcase 1) *There is a face  $D$  inside  $C$  that touches all these  $v$ -vertices.* Then we choose two nonadjacent vertices  $d'_1$  and  $d'_2$  from these five and three additional appropriate vertices  $a', b', c'$  from  $C$  such that the cyclic ordering along the cycle  $C$  is  $a' - b' - d'_1 - c' - d'_2$ . Then we connect  $d'_1$  and  $d'_2$  by an equalizer that is put inside  $D$ . Notice that in the graph  $F - \{d'_1, d'_2\}$  the vertex  $c'$  is in a component different from the component containing  $a'$  or  $b'$ .
- (Subcase 2) *There is an  $i$  and a path  $P$  (possibly just one edge) internally-disjoint from  $C$  that connects two vertices  $v_i$  and  $v_{i+3}$  (where the indices are taken modulo 5).* We put  $d'_1 = v_i$ ,  $d'_2 = v_{i+3}$  and call the remaining three  $v$ -vertices  $a', b', c'$  in such a way that the cyclic ordering along  $C$  is  $a' - b' - d'_1 - c' - d'_2$ . For every edge of  $P$  we connect its incident vertices by an equalizer. Again notice that in the graph  $F - V_P$  the vertex  $c'$  is in a component different from the component containing  $a'$  or  $b'$ .

In either subcase, we create five new vertices  $a, b, c, d_1, d_2$ , and connect them by equalizers to their corresponding primed vertices on the outer face of the skeleton. Finally, we partition the vertices of the skeleton into five connected subgraphs, each containing one of the vertices  $a', b', c', d'_1, d'_2$ , and we introduce equalizers along the edges of these subgraphs as in Lemma 2.4. This completes the construction.

It can be verified that this construction in both cases and subcases indeed fulfills the two properties (i) and (ii).

## 2.2 The NP-hardness argument

Now we prove Theorem 2.3. The proof will be done by a reduction from an NP-hard variant of the 3-satisfiability problem: Let  $\Phi$  be a Boolean formula in conjunctive normal form over a set  $X$  of logical variables; every clause in  $\Phi$  contains exactly three variables. With  $\Phi$  we associate a graph  $Q_\Phi$ . The vertices of  $Q_\Phi$  are the clauses and the variables in  $\Phi$ . There are two types of edges in  $Q_\Phi$ . The first type belongs to a cycle that spans all the clauses in some ordering. The second type connects a variable  $x \in X$  to a clause  $\phi \in \Phi$  if and only if  $x$  or  $\bar{x}$  occurs as a literal in  $\phi$ . We call a formula  $\Phi$  *planar* if for some choice of the cycle spanning all the clauses of  $\Phi$  the associated graph  $Q_\Phi$  is planar. Fellows *et al.* [17] proved that the restriction of the 3-satisfiability problem to planar formulae is NP-hard. (To be precise, they only show the NP-hardness for formulas with *at most* three literals per clause. One may achieve exactly three literals per clause by dropping the requirement of distinctness of literals per clause. Since variable-clause incidences will later be replaced by gadgets with nonadjacent contact points, our final graph will have no multiple edges anyway.)

Consider an arbitrary planar formula  $\Phi$  as described above, and let  $Q_\Phi$  be an associated planar graph. We will construct in polynomial time a planar graph  $G_\Phi$  which has the following

two properties: If formula  $\Phi$  is satisfiable, then  $G_\Phi$  has a strongly  $\mathcal{F}$ -free 2-coloring. If  $G_\Phi$  has a weakly  $\mathcal{F}$ -free 2-coloring, then formula  $\Phi$  is satisfiable. This clearly will prove Theorem 2.3.

Fix a planar embedding of  $Q_\Phi$ . The cycle through the clause vertices divides the plane into a bounded and an unbounded region. Variables in  $X$  that are embedded in the unbounded region are called *outer* variables, and variables in the bounded region are called *inner* variables. As it is usual in reductions from planar SAT, we construct a graph from the planar drawing of  $Q_\Phi$  by a series of local replacements. Slightly informally described, we thicken the edges and the vertices in the planar embedding of  $Q_\Phi$  such that they become streets and squares; this yields a map into which we will put our gadgets. For every variable  $x \in X$ , we put a vertex  $v(x)$  into the square corresponding to  $x$ . For every clause  $\phi \in \Phi$ , we put a corresponding clause gadget into the square corresponding to  $\phi$  in the following way.

- If all three literals in clause  $\phi$  belong to inner variables, then the clause gadget for  $\phi$  is a clause gadget  $\mathcal{C}_4(a, b, c, d)$  with four contact points. The contact point  $d$  lies in the center of the square of  $\phi$ , and the contact points  $a, b, c$  lie at the beginning of the streets leading to these three inner variables.
- If two literals in clause  $\phi$  belong to inner variables and one literal belongs to an outer variable, then the clause gadget for  $\phi$  is a clause gadget  $\mathcal{C}_5(a, b, c, d_1, d_2)$  with five contact points. The contact points  $d_1$  and  $d_2$  lie at the beginning of the streets that lead to the left and the right neighbors of the clause  $\phi$  on the clause cycle. The contact points  $a$  and  $b$  lie at the beginning of the streets that lead to the two inner variables. The contact point  $c$  lies at the beginning of the street that leads to the outer variable.
- The case of three outer variables, and the case of one inner and two outer variables are handled symmetrically to the above two cases.

If the variable  $x$  occurs un-negated (respectively, negated) in the clause  $\phi$ , then we put an equalizer (respectively, a negator) from  $v(x)$  to the corresponding contact point in the clause gadget for  $\phi$ . Finally, we put an equalizer gadget between the  $d$ -vertices into every street that connects a clause square to another clause square, and thus connect all clause gadgets into a ring. These equalizer gadgets connect contact points  $d$  from clause gadgets  $\mathcal{C}_4(a, b, c, d)$ , and contact points  $d_1$  and  $d_2$  from clause gadgets  $\mathcal{C}_5(a, b, c, d_1, d_2)$  in an appropriate way. This completes the construction of the graph  $G_\Phi$  which is easily seen to be planar.

Assume that formula  $\Phi$  is satisfiable, and consider a satisfying truth assignment. Intuitively speaking, color 1 will correspond to TRUE and color 0 will correspond to FALSE. Color all contact points  $d, d_1$ , and  $d_2$  of clause gadgets by color 0. If variable  $x$  is TRUE, then color the vertex  $v(x)$  by color 1. If  $x$  is FALSE, then color  $v(x)$  by 0. The equalizers and negators propagate the colors (respectively opposite colors) of the variables to the corresponding contact points  $a, b, c$  in the clause gadgets. Since the truth assignment is a satisfying truth assignment, in every clause gadget at least one of the contact points  $a, b, c$  is colored 1. Moreover, in every clause gadget the contact points  $d$  (respectively,  $d_1$  and  $d_2$ ) are colored 0. Therefore, we can use property (ii) of the clause gadgets to get a strongly  $\mathcal{F}$ -free 2-coloring of all used clause gadgets. Altogether, this yields a strongly  $\mathcal{F}$ -free 2-coloring for the graph  $G_\Phi$ .

Now assume that  $G_\Phi$  has a weakly  $\mathcal{F}$ -free 2-coloring. Because of the ring of equalizer gadgets that connect the clause gadgets to each other and property (i) of the  $\mathcal{C}_5$ -gadgets, all contact points  $d, d_1, d_2$  of clause gadgets must receive the same color; without loss of generality we assume that this color is 0. We construct the following truth assignment for  $X$ : If  $v(x)$  is colored 1, then  $x$  is set to TRUE. If  $v(x)$  is colored 0, then  $x$  is set to FALSE. Suppose for the sake of contradiction that some clause  $\phi$  in  $\Phi$  is not satisfied by this truth setting. Then the three literals in  $\phi$  all are FALSE, and hence all three contact points  $a, b, c$  in the corresponding clause gadget are colored 0. But then *all* contact points of this clause gadget are colored 0, and by property (i) of the clause gadgets the coloring cannot be a weakly  $\mathcal{F}$ -free 2-coloring. This contradiction shows that  $\Phi$  is satisfiable.

This completes the proof of Theorem 2.3.

### 3 Tree-free 2-colorings of planar graphs

The main result of this section is an NP-hardness result for weakly and strongly  $T$ -free 2-colorings of planar graphs for the case where  $T$  is a tree with at least two edges (see Theorem 3.3). The proof of this result is based on an inductive argument over the number of edges in  $T$ . The following two propositions are used as the base case of the induction.

Let  $K_{1,k}$  denote the complete bipartite graph with one vertex in one color class and the other  $k \geq 1$  vertices in the other color class. The leftmost drawing in Figure 2 shows a  $K_{1,7}$ .

**Proposition 3.1.** *For every  $k \geq 2$ , it is NP-hard to decide whether a planar graph has a weakly (strongly)  $K_{1,k}$ -free 2-coloring.*

*Proof.* For  $k = 2$ , the statement for weakly  $K_{1,k}$ -free 2-colorings follows from Proposition 1.1, and the statement for strongly  $K_{1,k}$ -free 2-colorings follows from Proposition 1.2.(i).

For  $k \geq 3$ , we apply Theorem 2.3. The first condition in this theorem is fulfilled, since for  $k \geq 3$ , the star  $K_{1,k}$  is a graph on at least four vertices with a cut vertex.

For the second condition, we note that the complete bipartite graph  $K_{2,2k-1}$  with color classes  $I$  with  $|I| = 2k - 1$ , and  $\{a, b\}$ , is an  $(a, b)$ -equalizer for  $\mathcal{F} = \{K_{1,k}\}$ . This graph satisfies Definition 2.2.(i): In any 2-coloring, at least  $k$  of the vertices in  $I$  receive the same color, say color 0. If  $a$  and  $b$  are colored differently, then one of them is colored 0. This would yield an induced monochromatic  $K_{1,k}$ . A good coloring as required in Definition 2.2.(ii) results from coloring  $a$  and  $b$  by the same color, and all vertices in  $I$  by the opposite color. This coloring has no monochromatic copy of  $K_{1,k}$  itself, and since all neighbors of the contact points are colored with the other color than the contact points, no monochromatic copy of  $K_{1,k}$  may cross the equalizer.  $\square$

As we mentioned in Section 1.1, Cowen, Goddard & Jesurum [15] have shown that the defective  $(2, d)$ -coloring problem for any  $d \geq 1$  is NP-hard even for planar graphs. This implies that strongly  $K_{1,k}$ -free 2-coloring is NP-hard for planar graphs for any  $k \geq 2$ , so the above proof is needed for the weak case only.

For  $1 \leq k \leq m$ , a *double-star*  $X_{k,m}$  is a tree of the following form:  $X_{k,m}$  has  $k + m + 2$  vertices. There are two adjacent central vertices  $y_1$  and  $y_2$ . Vertex  $y_1$  is adjacent to  $k$  leaves, and  $y_2$  is adjacent to  $m$  leaves. In other words, the double-star  $X_{k,m}$  results from adding an




Figure 2: The graph  $K_{1,7}$  and the double-stars  $X_{4,5}$  and  $X_{1,1}$ .

edge between the centers (vertices of maximum degree) of  $K_{1,k}$  and  $K_{1,m}$ . See Figure 2 for an illustration. Note that  $X_{1,1}$  is isomorphic to the path  $P_4$ .

**Proposition 3.2.** *For every  $1 \leq k \leq m$ , it is NP-hard to decide whether a planar graph has a weakly (strongly)  $X_{k,m}$ -free 2-coloring.*

*Proof.* We apply Theorem 2.3. The first condition in this theorem is fulfilled, since  $X_{k,m}$  is a graph on at least four vertices with a cut vertex. For the second condition, we construct an  $(a, b)$ -equalizer.

The  $(a, b)$ -equalizer  $\mathcal{E} = (V', E')$  consists of  $2m + k - 1$  independent copies  $(V^i, E^i)$  of the double-star  $X_{k,m}$  where  $1 \leq i \leq 2m + k - 1$ . Moreover, there are five special vertices  $a, b, v_1, v_2$ , and  $v_3$ . We define

$$\begin{aligned}
 V' &= \{v_1, v_2, v_3, a, b\} \cup \bigcup_{1 \leq i \leq 2m+k-1} V^i \quad \text{and} \\
 E' &= \{v_1v_2, v_2v_3, v_1v_3, av_3, bv_3\} \cup \\
 &\quad \bigcup_{1 \leq i \leq 2m+k-1} E^i \cup \\
 &\quad \bigcup_{1 \leq i \leq m} \{v_1v : v \in V^i\} \cup \\
 &\quad \bigcup_{m+1 \leq i \leq 2m} \{v_2v : v \in V^i\} \cup \\
 &\quad \bigcup_{2m+1 \leq i \leq 2m+k-1} \{v_3v : v \in V^i\}.
 \end{aligned}$$

See Figure 3 for an illustration.

We claim that every 2-coloring of  $\mathcal{E}$  with  $a$  and  $b$  colored differently contains a monochromatic induced copy of  $X_{k,m}$ : Consider some weakly  $X_{k,m}$ -free 2-coloring of  $\mathcal{E}$ . Then each copy  $(V^i, E^i)$  of  $X_{k,m}$  must have at least one vertex that is colored 0 and at least one vertex that is colored 1. If  $v_1$  and  $v_2$  had the same color, then together with appropriate vertices in  $V^i$ ,  $1 \leq i \leq 2m$ , they would form a monochromatic copy of  $X_{k,m}$ . Hence, we may assume by symmetry that  $v_1$  is colored 1, and that  $v_2$  and  $v_3$  are colored 0. Suppose for the sake




Figure 3: An equalizer for the double-star  $X_{k,m}$ .

of contradiction that  $a$  and  $b$  are colored differently. Then one of them would be colored 0, and there would be a monochromatic copy of  $X_{k,m}$  with center vertices  $v_3$  and  $v_2$ . Thus  $\mathcal{E}$  satisfies property (i) in Definition 2.2.

To show that also property (ii) in Definition 2.2 is satisfied, we construct a good 2-coloring: The vertices  $a$ ,  $b$ ,  $v_1$  are colored 0, and  $v_2$  and  $v_3$  are colored 1. In every set  $V^i$  with  $1 \leq i \leq m$ , one vertex is colored 0 and all other vertices are colored 1. In every set  $V^i$  with  $m + 1 \leq i \leq 2m + k - 1$ , one vertex is colored 1 and all other vertices are colored 0. This coloring has no monochromatic copy of  $X_{k,m}$ , and since vertex  $v_3$  as the only neighbor of the contact points  $a, b$  is colored differently than  $a, b$ , no monochromatic copy of any tree may cross  $\mathcal{E}$ .  $\square$

Now we are ready to prove the main result of this section.

**Theorem 3.3.** *Let  $T$  be a tree with at least two edges. Then it is NP-hard to decide whether a planar input graph  $G$  has a weakly (strongly)  $T$ -free 2-coloring.*

*Proof.* By induction on the number  $\ell$  of edges in  $T$ . If  $T$  has  $\ell = 2$  edges, then  $T = K_{1,2}$ , and NP-hardness follows by Proposition 1.1. If  $T$  has  $\ell \geq 3$  edges, then we consider the so-called *shaved* tree  $T^*$  of  $T$  that results from  $T$  by removing all the leaves. If the shaved tree  $T^*$  is a single vertex, then  $T$  is a star, and NP-hardness follows by Proposition 3.1. If the shaved tree  $T^*$  is a single edge, then  $T$  is a double-star, and NP-hardness follows by Proposition 3.2.

Hence, it remains to settle the case where the shaved tree  $T^*$  contains at least two edges. In this case we know from the induction hypothesis that weakly (strongly)  $T^*$ -free 2-coloring is NP-hard. Consider an arbitrary planar input graph  $G^*$  for weakly (strongly)  $T^*$ -free 2-coloring. To complete the NP-hardness proof, we will construct in polynomial time a planar graph  $G$  that has a weakly (strongly)  $T$ -free 2-coloring if and only if  $G^*$  has a weakly (strongly)  $T^*$ -free 2-coloring: Let  $\Delta$  be the maximum number of leaves pending on a vertex of  $T$ . For

every vertex  $v$  in  $G^*$ , we create  $\Delta$  independent copies  $T_1(v), \dots, T_\Delta(v)$  of  $T$ , and we join  $v$  to all vertices of all these copies.

Assume first that  $G^*$  is weakly (strongly)  $T^*$ -free 2-colorable. We extend this coloring to a weakly (strongly)  $T$ -free 2-coloring of  $G$  by taking a proper 2-coloring of every subgraph  $T_i(v)$  in  $G$ , such that for every  $v \in V_{G^*}$ , exactly one vertex of each  $T_i(v)$  receives the same color as  $v$ . It can be verified that this extended 2-coloring for  $G$  does not contain any monochromatic copy of  $T$ .

Now assume that  $G$  is weakly (strongly)  $T$ -free 2-colorable, and let  $c$  be such a 2-coloring. Every subgraph  $T_i(v)$  in  $G$  must meet both colors. This implies that every vertex  $v$  in the subgraph  $G^*$  of  $G$  has at least  $\Delta$  neighbors of color 0 and at least  $\Delta$  neighbors of color 1 in the subgraphs  $T_i(v)$ . Any monochromatic (induced) copy of  $T^*$  in  $G^*$  would then extend to a monochromatic (induced) copy of  $T$  in  $G$ , and hence the restriction of the coloring  $c$  to the subgraph  $G^*$  is a weakly (strongly)  $T^*$ -free 2-coloring. This concludes the proof of the theorem.  $\square$

Using the same ideas as in the proofs of the previous theorem and propositions, one can prove the following more general statement about larger sets of forbidden graphs.

**Theorem 3.4.** *Let  $\mathcal{F}$  be a finite set of graphs containing at least one tree with at least two edges. Then both weakly and strongly  $\mathcal{F}$ -free 2-coloring are NP-hard.*

*Proof.* Let  $T \in \mathcal{F}$  be a tree in  $\mathcal{F}$  with the minimum number of edges. If  $T = P_3$ , then the remaining graphs in  $\mathcal{F}$  must be complete (every non-complete connected graph contains  $P_3$  as an induced subgraph), so they could be only  $K_3$  or  $K_4$ . The NP-hardness of  $\mathcal{F}$ -free 2-coloring then follows from Proposition 1.1, since for coloring triangle-free graphs, graphs containing triangles are irrelevant as forbidden subgraphs.

For  $T$  being a star (with at least three leaves) or a double-star, the result follows directly from the construction of equalizers in the proofs of Propositions 3.1 and 3.2, since the good colorings presented there are such that the neighbors of the contact points receive a different color from the contact points, and the only connected monochromatic subgraphs of the equalizers are singletons (in case of  $T = K_{1,k}$ ) or smaller double-stars ( $X_{k-1,m}$  in case of  $T = X_{k,m}$ ). Since  $T$  itself may be used as the skeleton of the negator and the clause gadgets, the good colorings of these gadgets also do not contain a monochromatic copy of any graph of  $\mathcal{F}$  (neither induced nor non-induced).

If  $T$  is such that the shaved tree  $T^*$  has at least two edges, we proceed by induction similarly as in the proof of Theorem 3.3. For any graph  $F \in \mathcal{F}$ , we denote by  $F^*$  the shaved copy of  $F$ , i.e., the graph obtained from  $F$  by removing all leaves (vertices of degree one). Let  $\Delta$  be the maximum number of leaves pending on a vertex of a graph from  $\mathcal{F}$ . Construct the graph  $G$  from a graph  $G^*$  as in the proof of Theorem 3.3 and use the fact that  $G$  has a weakly (strongly)  $\mathcal{F}$ -free 2-coloring if and only if  $G^*$  has a weakly (strongly)  $\mathcal{F}^*$ -free 2-coloring, where  $\mathcal{F}^* = \{F^* : F \in \mathcal{F}\}$ .  $\square$

## 4 Cycle-free 2-colorings of planar graphs

In this section we turn to the case when the forbidden graph  $F$  is not a tree and hence contains a cycle (we assume  $F$  is connected).

If  $F$  contains an odd cycle, then the Four Color Theorem (4CT) shows that any planar graph  $G$  has a strongly  $F$ -free 2-coloring: a proper 4-coloring of  $G$  partitions  $V_G$  into two sets  $S_1$  and  $S_2$  each inducing a bipartite graph. Coloring all the vertices of  $S_i$  by color  $i$  yields a strongly  $F$ -free 2-coloring of  $G$ . If we extend the set of forbidden graphs to all cycles of odd length, denoted by  $\mathcal{F}_{\text{odd}}$ , then the converse is also true: In any  $\mathcal{F}_{\text{odd}}$ -free 2-coloring of  $G$  both monochromatic subgraphs of  $G$  are bipartite, yielding a 4-coloring of  $G$ . To summarize we obtain the following.

**Proposition 4.1.** *The statement “ $\chi^S(\mathcal{F}_{\text{odd}}, G) \leq 2$  for every planar graph  $G$ ” is equivalent to the 4CT.*

In case  $F$  is just the triangle  $C_3$ , one can avoid using the heavy 4CT machinery to prove that  $\chi^S(C_3, G) \leq 2$  for every planar graph  $G$  by applying a result due to Burstein [10]. A brief sketch of the argument is as follows. Prove by induction that in any plane triangulation, any nonmonochromatic precoloring of the outer face (triangle) can be extended to a coloring which avoids monochromatic triangles.

If  $F$  contains no triangles, a result of Thomassen [31] can be applied. He proved that the vertex set of any planar graph can be partitioned into two sets each of which induces a subgraph with no cycles of length exceeding 3. Hence every planar graph is strongly  $\mathcal{F}_{\geq 4}$ -free 2-colorable, where  $\mathcal{F}_{\geq 4}$  denotes the set of all cycles of length exceeding 3. The following theorem summarizes the above observations.

**Theorem 4.2.** *If the forbidden connected subgraph  $F$  is not a tree, then every planar graph is strongly and hence also weakly  $F$ -free 2-colorable.*

The picture changes if one forbids all cycles, or a combination of cycles including the triangle. A result of Stein [30] states that the vertex set of a plane triangulation  $G$  can be partitioned into two sets each inducing a forest if and only if the plane dual of  $G$  is hamiltonian. Since deciding hamiltonicity of planar cubic graphs is NP-hard (see [20]), this implies that deciding whether a (maximal) planar graph has a weakly (strongly) cycle-free 2-coloring is NP-hard. We are able to slightly strengthen this statement.

**Theorem 4.3.** *Let  $\mathcal{F}_{345} = \{C_3, C_4, C_5\}$  be the set of cycles of lengths three, four, and five. Then the problem of deciding whether a given planar graph has a weakly (strongly)  $\mathcal{F}_{345}$ -free 2-coloring is NP-hard.*

We prove this theorem in a more general setting as a corollary of Theorem 4.4. But first let us note that  $\{C_3, C_4, C_5\}$  is a minimal set of cycles which determines an NP-hard instance of the  $\mathcal{F}$ -free 2-coloring problem. Indeed, if  $\mathcal{F} \subset \{C_3, C_4, C_5\}$  is a proper subset, then every planar graph is strongly  $\mathcal{F}$ -free 2-colorable. We have noted this already for  $\mathcal{F} \subseteq \{C_3, C_5\}$  and  $\mathcal{F} \subseteq \{C_4, C_5\}$ , and the last case  $\mathcal{F} = \{C_3, C_4\}$  is covered by the result of Kaiser & Škrekovski [26] who proved that every planar graph is strongly  $\{C_3, C_4\}$ -free 2-colorable.

**Theorem 4.4.** *Let  $\mathcal{F}$  be a finite set of planar 2-connected graphs. If there exists a planar graph which is not weakly (strongly)  $\mathcal{F}$ -free 2-colorable, then weakly (strongly)  $\mathcal{F}$ -free 2-coloring is NP-hard for planar input graphs.*

*Proof.* Consider the graphs of  $\mathcal{F}$  with some fixed plane embeddings. If every face of every graph  $F \in \mathcal{F}$  is  $C_3$  or  $C_4$ , then every planar graph is strongly  $\mathcal{F}$ -free 2-colorable by the main result in [26]. If not, there is an  $F \in \mathcal{F}$  with a face of size at least 5 and the first assumption of Theorem 2.3 is met.

Next we show how to construct an equalizer. Let  $H'$  be a smallest (by the number of edges) planar graph which is not weakly (strongly)  $\mathcal{F}$ -free 2-colorable. Take an edge  $xy \in E_{H'}$  and denote by  $H$  the graph obtained from  $H'$  by deleting this edge. Then  $H$  is weakly (strongly)  $\mathcal{F}$ -free 2-colorable, and in every weakly (strongly)  $\mathcal{F}$ -free 2-coloring of  $H$  the vertices  $x$  and  $y$  receive the same color. We construct an equalizer  $\mathcal{E}$  by concatenating sufficiently many copies of  $H$ . More formally, choose a number  $k$  to be larger than the order of any graph in  $\mathcal{F}$ . The copies of  $H$  will be  $H_i = (V_i, E_i)$  with  $V_i = \{v_i : v \in V_H\}$  and  $E_i = \{u_i v_i : uv \in E_H\}$ , for  $i = 1, 2, \dots, k$ . For  $i = 1, 2, \dots, k - 1$ , we identify  $y_i$  with  $x_{i+1}$ , and we set  $a = x_1$  and  $b = y_k$  to be the contact points.

Clearly  $\mathcal{E}$  is planar and in every weakly (strongly)  $\mathcal{F}$ -free 2-coloring of  $\mathcal{E}$  the vertices  $x_i, y_i$  for  $i = 1, 2, \dots, k$ , and hence also  $a$  and  $b$ , receive the same color.

Let  $c$  be a weakly (strongly)  $\mathcal{F}$ -free 2-coloring of  $H$ . Color  $\mathcal{E}$  using  $c$  on every  $H_i$ . Consider a graph  $F \in \mathcal{F}$ . No copy of  $F$  which lies entirely in  $\mathcal{E}$  is monochromatic, since the 2-connectedness of  $F$  implies that such a copy of  $F$  lies entirely within one of the  $H_i$ 's. Therefore this 2-coloring of  $\mathcal{E}$  is weakly (strongly)  $F$ -free. It also follows from the 2-connectedness of  $F$  that every copy of  $F$  which crosses  $\mathcal{E}$  contains a path from  $a$  to  $b$  through  $\mathcal{E}$ . But every such path has more vertices than  $F$ . Hence the 2-coloring of  $\mathcal{E}$  is good.  $\square$

To conclude the proof of Theorem 4.3, it would suffice to construct a planar graph which is not weakly  $\mathcal{F}_{345}$ -free 2-colorable. It is, however, equally simple to describe an equalizer for  $\mathcal{F}_{345}$  (and exploit the fact that  $C_5 \in \mathcal{F}_{345}$  is 2-connected and every plane drawing contains a face of size 5): Let  $\Theta(x, y)$  be the graph depicted in Figure 4. This graph has the following important property: In any weakly  $\mathcal{F}_{345}$ -free 2-coloring of  $\Theta(x, y)$ , the vertices  $x$  and  $y$  have different colors (we leave the simple proof of this fact to the reader). The  $(a, b)$ -equalizer is constructed from a graph  $\Theta(a, x)$  and a graph  $\Theta(b, y)$  by identifying the two vertices  $x$  and  $y$ . A good 2-coloring of the  $(a, b)$ -equalizer is induced by the 2-coloring indicated in Figure 4.

The following statement is now a direct corollary of Theorem 4.4 and Theorem 4.3.

**Corollary 4.5.** *For any finite set  $\mathcal{F}_{\geq 345} \supseteq \{C_3, C_4, C_5\}$  of cycles, both weakly and strongly  $\mathcal{F}_{\geq 345}$ -free 2-coloring are NP-hard for planar input graphs.*

## 5 3-colorings of planar graphs

A *linear forest* is a disjoint union of paths (some of which may be trivial). The following result was proved independently in [23] and [28].




Figure 4: A gadget for the forbidden set  $\mathcal{F}_{345}$  in Theorem 4.3.

**Proposition 5.1.** [Goddard [23] and Poh [28]]

Every planar graph has a partition of its vertex set into three subsets such that every subset induces a linear forest.

This result immediately implies that if a connected graph  $F$  is not a path, then  $\chi^w(F, G) \leq 3$  and  $\chi^s(F, G) \leq 3$  hold for *all* planar graphs  $G$ . Hence, these coloring problems are trivially solvable in polynomial time.




Figure 5: Example for the graph  $Y_3$ .

We now turn to the remaining cases of  $F$ -free 3-coloring for planar graphs where the forbidden graph  $F$  is a path. We start with a technical lemma that will yield a gadget for the NP-hardness argument.

**Lemma 5.2.** For every  $k \geq 2$ , there exists an outerplanar graph  $Y_k$  that satisfies the following properties.

- (i)  $Y_k$  is not weakly  $P_k$ -free 2-colorable.
- (ii) There exists a strongly  $P_k$ -free 3-coloring of  $Y_k$ , in which at least one color class being is independent set.

*Proof.* The skeleton of the graph  $Y_k$  is formed by a regular tree, in which every inner vertex has exactly  $k$  children, and all paths from the root to a leaf have exactly  $k$  vertices. Additionally to the edges in this regular tree, the children of every inner vertex are connected by a path. See Figure 5 for an illustration.

Suppose there was a  $P_k$ -free 2-coloring of  $Y_k$ . Since the children of any vertex induce a path  $P_k$ , both colors must show up at the children. Consequently, for every inner vertex  $v$  at least one of its children must get the same color as  $v$ . But this yields a monochromatic induced  $P_k$  running from the root to some leaf. This contradiction proves property (i). Property (ii) is straightforward to prove: The graph  $Y_k$  is outerplanar, and hence properly 3-colorable.  $\square$

**Theorem 5.3.** *For any path  $P_k$  with  $k \geq 2$ , it is NP-hard to decide whether a planar input graph has a weakly (strongly)  $P_k$ -free 3-coloring.*

*Proof.* We will use induction on  $k$ . The basic cases are  $k = 2$  and  $k = 3$ . For  $k = 2$ , both weakly and strongly  $P_2$ -free 3-colorings are equivalent to proper 3-coloring which is well-known to be NP-hard for planar graphs.

Next, consider the case  $k = 3$ . Proposition 1.2.(ii) yields NP-hardness of strongly  $P_3$ -free 3-coloring for planar graphs. For weakly  $P_3$ -free 3-coloring, we sketch a reduction from proper 3-coloring of planar graphs. As a gadget, we use the outerplanar graph  $Z$  depicted in Figure 6. The crucial property of  $Z$  is that it does not allow a weakly  $P_3$ -free 2-coloring, as is easily checked. Now consider an arbitrary planar graph  $G$ . From  $G$  we construct the planar graph  $G'$ : For every vertex  $v$  in  $G$ , create a copy  $Z(v)$  of  $Z$ , and add all possible edges between  $v$  and  $Z(v)$ . Since  $Z(v)$  is outerplanar, the new graph  $G'$  is planar. It is easy to verify that  $\chi(G) \leq 3$  if and only if  $\chi^w(P_3, G') \leq 3$ .




Figure 6: The graph  $Z$  in the proof of Theorem 5.3.

For  $k \geq 4$ , we will give a reduction from weakly (strongly)  $P_{k-2}$ -free 3-coloring to weakly (strongly)  $P_k$ -free 3-coloring. Consider an arbitrary planar graph  $G$ , and construct the following planar graph  $G'$ : For every vertex  $v$  in  $G$ , create a copy  $Y_k(v)$  of the graph  $Y_k$  from Lemma 5.2, and add all possible edges between  $v$  and  $Y_k(v)$ . Since  $Y_k$  is outerplanar, the new graph  $G'$  is planar. If  $G$  has a weakly (strongly)  $P_{k-2}$ -free 3-coloring, then this can be extended to a weakly (strongly)  $P_k$ -free 3-coloring of  $G'$  by coloring the subgraphs  $Y_k(v)$  according to Lemma 5.2.(ii). And if  $G'$  has a weakly (strongly)  $P_k$ -free 3-coloring, then by Lemma 5.2.(i) this induces a weakly (strongly)  $P_{k-2}$ -free 3-coloring for  $G$ .  $\square$

## 6 Concluding remarks and open problems

**6.1 Triangle-free graphs.** By modifying the gadgets for the equalizers in such a way that the planar graph  $G_{\Phi}$  constructed in the proof of Theorem 2.3 becomes triangle-free, one might be able to prove complexity results for weakly (strongly)  $F$ -free 2-coloring restricted to triangle-free planar graphs. In fact, it is not difficult to apply this method to prove that for  $F = K_{1,k}$  with  $k \geq 2$ , weakly (strongly)  $F$ -free 2-coloring remains NP-hard for triangle-free planar graphs.

**Problem.** Is it true that every triangle-free planar graph  $G$  is  $P_4$ -free 2-colorable? This would imply that for every connected graph  $F$  of diameter at least 3 there is a weakly  $F$ -free 2-coloring of  $G$ .

**6.2 Monotonicity.** All our NP-hardness techniques are such that hardness proofs for  $\mathcal{F}$ -free 2-colorability extend naturally to NP-hardness of  $\mathcal{F}'$ -free 2-colorability for any finite  $\mathcal{F}' \supseteq \mathcal{F}$ . This raises the following question.

**Problem.** For finite sets of graphs  $\mathcal{F}' \supseteq \mathcal{F}$ , is it true that  $\mathcal{F}' - F - 2 - CPG \propto \mathcal{F} - F - 2 - CPG$ ? ( $\mathcal{F} - F - 2 - CPG$  standing for  $\mathcal{F}$ -Free-2-Coloring-Planar-Graphs.)

Note that this is not necessarily true for infinite sets of forbidden graphs. The infinite set  $\mathcal{F}_{cycle}$  of all cycles has uncountably many subsets, and if each of these defines a different problem, infinitely many of them will have to be undecidable, whereas deciding the existence of an  $\mathcal{F}_{cycle}$ -free 2-coloring is surely in NP.

**6.3 Forbidden sets of cycles.** It would be interesting to characterize for which particular (finite) sets of forbidden cycles the  $\mathcal{F}$ -free 2-coloring problem on planar graphs is feasible and for which it is hard. In particular, for two cycles this question remains open if one of them is the triangle and the other one is an even cycle of length greater than 4.

**Problem.** For which  $k > 2$  does there exist a planar graph which is not  $\{C_3, C_{2k}\}$ -free 2-colorable?

**6.4 Equalizers.** Despite our inductive proof of NP-hardness for forbidden trees, it would be interesting to know whether one can use the equalizer gadget machinery directly.

**Problem.** Does there exist an equalizer for any tree  $T$ ?

## Acknowledgments

We are grateful to Oleg Borodin, Alesha Glebov, Sasha Kostochka, and Carsten Thomassen for fruitful discussions on the topic of this paper.

## References

- [1] D. ACHLIOPTAS, *The complexity of  $G$ -free colorability*, Discrete Math., 165/166 (1997), pp. 21–30.
- [2] M. O. ALBERTSON, R. E. JAMISON, S. T. HEDETNIEMI, AND S. C. LOCKE, *The subchromatic number of a graph*, Discrete Math., 74 (1989), pp. 33–49.
- [3] D. ARCHDEACON, *A note on defective colorings of graphs in surfaces*, J. Graph Theory, 11 (1987), pp. 517–519.
- [4] I. BROERE AND C. M. MYNHARDT, *Generalized colorings of outer-planar and planar graphs*, in Graph theory with applications to algorithms and computer science, Wiley, New York, 1985, pp. 151–161.
- [5] H. BROERSMA, F. V. FOMIN, J. KRATOCHVÍL, AND G. J. WOEGINGER, *Planar graph coloring with forbidden subgraphs: Why trees and paths are dangerous*, Proceedings of the 8th Scandinavian Workshop on Algorithm Theory (SWAT 2002), Springer-Verlag Lecture Notes in Computer Science, vol. 2368, 2002, pp. 160–169.
- [6] H. J. BROERSMA, F. V. FOMIN, J. NEŠETŘIL, AND G. J. WOEGINGER, *More about subcolorings*, Computing, 69 (2002), pp. 187–203.
- [7] J. I. BROWN AND D. G. CORNEIL, *On generalized graph colorings*, J. Graph Theory, 11 (1987), pp. 87–99.
- [8] J. I. BROWN AND D. G. CORNEIL, *Graph properties and hypergraph colorings*, Discrete Math., 98 (1991), pp. 81–93.
- [9] J. I. BROWN AND D. G. CORNEIL, *On uniquely  $-G$   $k$ -colorable graphs*, Quaestiones Math., 15 (1992), pp. 477–487.
- [10] M. I. BURSTEIN, *The bi-colorability of planar hypergraphs*, Sakhart. SSR Mecn. Akad. Moambe, 78 (1975), pp. 293–296.
- [11] G. CHARTRAND, D. P. GELLER, AND S. HEDETNIEMI, *A generalization of the chromatic number*, Proc. Cambridge Philos. Soc., 64 (1968), pp. 265–271.
- [12] G. CHARTRAND AND H. V. KRONK, *The point-arboricity of planar graphs*, J. London Math. Soc., 44 (1969), pp. 612–616.
- [13] G. CHARTRAND, H. V. KRONK, AND C. E. WALL, *The point-arboricity of a graph*, Israel J. Math., 6 (1968), pp. 169–175.
- [14] L. J. COWEN, R. H. COWEN, AND D. R. WOODALL, *Defective colorings of graphs in surfaces: partitions into subgraphs of bounded valency*, J. Graph Theory, 10 (1986), pp. 187–195.

- [15] L. J. COWEN, W. GODDARD, AND C. E. JESURUM, *Defective coloring revisited*, J. Graph Theory, 24 (1997), pp. 205–219.
- [16] A. FARRUGIA, *Uniqueness and complexity in generalized coloring*, PhD thesis, University of Waterloo, 2003.
- [17] M. R. FELLOWS, J. KRATOCHVÍL, M. MIDDENDORF, F. PFEIFFER, *The complexity of induced minors and related problems*, Algorithmica, 13 (1995), pp. 266–282.
- [18] J. FIALA, K. JANSEN, V. B. LE, AND E. SEIDEL, *Graph subcoloring: Complexity and algorithms*, SIAM J. Discrete Math., 16 (2003), pp. 635–650.
- [19] M. FRICK AND M. A. HENNING, *Extremal results on defective colorings of graphs*, Discrete Math., 126 (1994), pp. 151–158.
- [20] M. R. GAREY, D. S. JOHNSON, AND R. E. TARJAN, *The planar Hamiltonian problem is NP-complete*, SIAM J. Comput., 5 (1976), pp. 704–714.
- [21] J. GIMBEL AND C. HARTMAN, *Subcolorings and the subchromatic number of a graph*, Discrete Math., 272 (2003), pp. 139–154.
- [22] J. GIMBEL AND J. NEŠETŘIL, *Partitions of graphs into cographs*, Technical Report 2000-470, KAM-DIMATIA, Charles University, Czech Republic, 2000. To appear in Discrete Math.
- [23] W. GODDARD, *Acyclic colorings of planar graphs*, Discrete Math., 91 (1991), pp. 91–94.
- [24] F. HARARY, *Conditional colorability in graphs*, in Graphs and applications, Wiley, New York, 1985, pp. 127–136.
- [25] C. T. HOÀNG AND V. B. LE,  *$P_4$ -free colorings and  $P_4$ -bipartite graphs*, Discrete Math. Theor. Comput. Sci., 4 (2001), pp. 109–122.
- [26] T. KAISER AND R. ŠKREKOVSKI, *Planar graph colorings without short monochromatic cycles*, J. Graph Theory, 46 (2004), pp. 25–38.
- [27] H.-O. LE AND V. B. LE, *The NP-completeness of  $(1, r)$ -subcoloring of cubic graphs*, Information Proc. Letters, 81 (2002), pp. 157–162.
- [28] K. S. POH, *On the linear vertex-arboricity of a planar graph*, J. Graph Theory, 14 (1990), pp. 73–75.
- [29] H. SACHS, *Finite graphs (Investigations and generalizations concerning the construction of finite graphs having given chromatic number and no triangles)*, in Recent Progress in Combinatorics, Academic Press, New York, 1969, pp. 175–184.
- [30] S. K. STEIN, *B-sets and planar graphs*, Pac. J. Math., 37 (1971), pp. 217–224.
- [31] C. THOMASSEN, *Decomposing a planar graph into degenerate graphs*, J. Combin. Theory Ser., B 65 (1995), pp. 305–314.