

**Title:** Intraocular pressure responses to four different isometric exercises in men and women

**Running head:** Isometric exercise increases intraocular pressure

**Authors:** Jesús Vera<sup>a</sup>, PhD; Beatriz Redondo<sup>a</sup>, MS; George-Alex Koulieris<sup>b</sup>, PhD; Alejandro Torrejon<sup>c</sup>, MS; Raimundo Jiménez<sup>a</sup>, PhD; Amador Garcia-Ramos<sup>c,d</sup>, PhD.

**Affiliations:**

<sup>a</sup> Department of Optics, Faculty of Sciences, University of Granada, Granada, Spain.

<sup>b</sup> Department of Computer Science, Durham University, UK. <sup>c</sup> Department of Physical Education and Sport, Faculty of Sport Sciences, University of Granada, Spain.

<sup>c</sup> Department of Physical Education and Sport, Faculty of Sport Sciences, University of Granada, Spain.

<sup>d</sup> Department of Sports Sciences and Physical Conditioning, Faculty of Education, CIEDE, Catholic University of Most Holy Concepción, Concepción, Chile.

**Corresponding author:** Raimundo Jiménez, Department of Optics, University of Granada, Campus de la Fuentenueva 2, 18001 Granada, Spain. Tel: +34 958244067; fax: +34 958248533. E-mail: raimundo@ugr.es

## **Abstract**

**Significance:** The performance of resistance exercise has evidenced to induce abrupt intraocular pressure changes, which has been linked to the onset and progression of glaucoma. We found that four different isometric resistance exercises lead to an instantaneous and progressive intraocular pressure elevation, with these changes being independent on the type exercise.

**Purpose:** The impact of physical exercise on intraocular pressure has demonstrated to be dependent on exercise type and intensity, as well as individuals' characteristics. In this study, we aimed to explore the influence of the load, exercise type and participant's sex on the intraocular pressure behaviour during a 2-min isometric effort.

**Methods:** Twenty-eight physically active collegiate students performed 2-minutes of isometric exercise in the military press, biceps curl, leg extension and calf raise exercises against two different loads (high-load and low-load). Intraocular pressure was measured by rebound tonometry before, during (semi-continuous assessment [24 measurements]), and after 10 seconds of recovery in each of the eight (4 exercises x 2 loads) conditions.

**Results:** We found a statistically significant effect of load ( $p < .001$ ,  $\eta_p^2 = 0.906$ ), with greater intraocular pressure values when performing the isometric exercises against heavier loads. There was a positive intraocular pressure rise during the execution of isometric exercise in the high-load condition, returning to baseline levels after 10 seconds of passive recovery. The exercise type and participant's sex did not reveal statistically significant differences ( $p = .326$  and  $p = .558$ , respectively).

**Conclusions:** Our data evidenced an instantaneous and progressive intraocular pressure rise during the execution of isometric exercise leading to muscular failure, regardless of the exercise type and participant's sex. After exercise, intraocular pressure rapidly returned to baseline levels (within 10 seconds). The inclusion of glaucoma patients in future studies is guaranteed.

**Keywords:** resistance training; exercise physiology; ocular health; glaucoma management.

# 1        **Intraocular pressure responses to four different isometric exercises in men and women**

## 2        **Introduction**

3        The regular practice of physical activity is associated with an increase in life expectancy and a  
4        decrease in the development of mental conditions (e.g., depression, dementia or anxiety).<sup>1,2</sup> Recently,  
5        eye care specialists have begun to investigate the effects of physical exercise on ocular health, aiming  
6        to identify the most pertinent exercise programs for subjects with different ocular conditions.<sup>3-5</sup>  
7        Within the range of ocular conditions that have attracted researchers' attention, glaucoma is  
8        undoubtedly the most investigated visual condition due to both its high prevalence worldwide and the  
9        severity of the symptoms (irreversible vision loss).<sup>6,7</sup> The main variable related to the onset and  
10       progression of glaucoma is the intraocular pressure, which is defined as the pressure exerted by the  
11       aqueous humour against the outer layer of the eye.<sup>8</sup> Training programs should attempt to reduce  
12       baseline intraocular pressure values and avoid intraocular pressure peaks during exercise because  
13       these are the two most important factors for the management of glaucoma.<sup>9</sup>

14       Numerous studies have found acute intraocular pressure changes during or immediately after  
15       the performance of different physical efforts.<sup>10-15</sup> For example, resistance training against heavy loads  
16       acutely increases intraocular pressure values,<sup>16</sup> while running at different intensities seems to decrease  
17       intraocular pressure.<sup>11,12</sup> The impact of resistance training on intraocular pressure has been tested for  
18       dynamic and isometric exercises, being the intraocular pressure increments generally higher for the  
19       isometric exercises.<sup>14,17</sup> The results of previous studies suggest that performing resistance training  
20       against heavy loads, and mainly the execution of isometric exercises, should be discouraged for  
21       individuals at high risk for glaucoma onset or progression.

22       It is known that the changes in intraocular pressure during resistance training depend on  
23       several factors such as the magnitude of the load, participants' fitness level and sex, and the type of  
24       exercise.<sup>13,18,19</sup> Vera and colleagues reported a positive association between the intraocular pressure

25 rise and the load lifted during the dynamic bench press and back-squat exercises.<sup>18</sup> Regarding the  
26 exercise type, there is evidence suggesting that dynamic resistance training exercises involving larger  
27 muscle mass and the upper-body promote greater intraocular pressure rises compared to exercises  
28 involving smaller muscle groups and the lower-body, respectively.<sup>13</sup> Studies comparing the acute  
29 changes of intraocular pressure between men and women have reported contradictory results.  
30 Whereas women exhibited a greater intraocular pressure reduction during a high intensity interval  
31 training protocol,<sup>11</sup> the changes in intraocular pressure values during the execution of four different  
32 dynamic resistance training exercises (back squat, military press, biceps curl, and calf raise) were  
33 comparable for men and women.<sup>13</sup> In addition, it is unknown whether the the type of exercise and  
34 participants' sex may modulate the intraocular pressure behaviour during different isometric  
35 exercises. Therefore, it would be important to assess the intraocular pressure responses to isometric  
36 exercises that involve different amounts of muscle size and body parts in men and women.

37 To address the limitations previously mentioned, the intraocular pressure of healthy men and  
38 women was semi-continuously measured for 2 minutes during 4 isometric exercises (military press,  
39 biceps curl, leg extensions, and calf raises) performed against 2 loading conditions. Therefore, the  
40 main objective of the present study was to elucidate whether the intraocular pressure behaviour during  
41 a 2-minutes isometric effort is affected by (i) the magnitude of the load, (ii) the type of exercise, and  
42 (iii) the participant's sex. Based on previous findings, we hypothesized that (i) the use of greater loads  
43 would promote a higher intraocular pressure rise,<sup>18</sup> (ii) exercises performed with the upper-body  
44 involving large muscle groups would lead to greater intraocular pressure values,<sup>13</sup> and (iii) the lack  
45 of studies comparing the differences between men and women on the intraocular pressure responses  
46 to isometric exercises did not allow us to formulate any specific hypothesis.

## 47 **Methods**

### 48 *Participants*

49 Twenty-eight physically active collegiate students took part in this study (Table 1). All participants

	Total sample (n =28)	Men (n = 14)	Women (n = 14)
<b>Anthropometrical characteristics</b>			
Age (years)	22.4 ± 2.1	21.8 ± 2.1	23.0 ± 2.1
Height (cm)	172.4 ± 6.8	176.9 ± 5.6	167.8 ± 4.6
Body mass (kg)	67.6 ± 11.1	77.1 ± 5.7	58.2 ± 5.7
<b>Loads applied</b>			
Military press (kg)	14.9 ± 3.4	17.6 ± 1.8	12.2 ± 2.4
Biceps curl (kg)	10.0 ± 3.0	11.8 ± 2.4	8.2 ± 2.4
Leg extensions (kg)	22.6 ± 5.2	26.5 ± 3.3	18.6 ± 3.7
Calf raises (kg)	48.8 ± 12.1	55.4 ± 13.0	41.7 ± 5.5

50

51 had at least 2 years of resistance training experience and were familiar with the exercises used in this  
52 study. We excluded participants with any physical limitation that could compromise tested  
53 performance, as well as those with a history of any ocular or cardiovascular disease or surgery.

54 **Participants were not taking any medication that could affect eyes physiology and women were not**  
55 **assessed during the menstruation phase.** The study followed the guidelines of the Declaration of  
56 Helsinki and was approved by the Institutional Review Board.

57 **Table 1.** Mean ± standard deviation of anthropometrical characteristics and isometric  
58 exercise loads of all the participants included in this study.

59 \*\*Table 1 near here\*\*

### 60 *Experimental design*

61 We used a repeated measures design to explore the cumulative effect of four isometric exercises on  
62 intraocular pressure. Participants attended to the laboratory on two occasions separated by 48-72  
63 hours, **which were scheduled at the same time of the day (± 1 hour) in order to avoid the influence of**  
64 **circadian variations on physical performance.**<sup>20</sup> In the first session, they read and signed the consent  
65 form, and filled-in the demographic questionnaire. Participants were also instructed how to perform  
66 the different isometric exercises and the maximal load that participants could hold for approximately  
67 2 minutes was determined (see testing procedure below). The second session consisted of eight sets

68 (4 exercises [military press, biceps curl, leg extensions and calf raise] x 2 loads [heavy load: load that  
69 could be hold for 2 minutes; light load: no additional load was applied]). Each set was separated by  
70 10 minutes of passive rest. The order of the exercises and loads was randomised. Intraocular pressure  
71 was measured before exercise, during the 2-minutes of isometric effort (continuous measurement), and  
72 after 10 seconds of recovery (**Figure 1**). Both experimental sessions were conducted under controlled  
73 environmental conditions ( $\sim 22^{\circ}\text{C}$  and  $\sim 60\%$  humidity), and participants were not allowed to drink  
74 or eat during the experiment.

### 75 *Isometric exercises*

76 The maximum load that participants could hold for approximately 2 minutes in an isometric condition  
77 was determined in the first testing session for all exercises (military press, biceps curl, leg extension,  
78 and calf raise). After a warm-up consisting of 5 minutes of jogging and joint mobility exercises, an  
79 incremental loading test was performed to determine the maximum load. The initial load  
80 corresponded to the 75% of the load that participants believed that they could hold for 2 minutes.  
81 This load was progressively increased in agreement between the participant and an experienced  
82 researcher. To minimize fatigue, participants were instructed to stop the exercise when they or the  
83 researcher perceived that the applied load could be hold for more than 2 minutes. Three minutes were  
84 implemented between successive sets. The average number of sets needed to reach the maximum load  
85 was  $2.4 \pm 1.2$  for the military press,  $1.8 \pm 1.5$  for the biceps curl,  $2.8 \pm 1.6$  for the leg extension, and  
86  $2.5 \pm 1.5$  for the calf raise. The maximum load was determined at the same positions that were used  
87 for the measurement of intraocular pressure during session 2 (see **Figure 1**): (I) *military press*:  
88 participants seated with the back supported by a bench reclined  $5^{\circ}$  respect to the vertical direction and  
89 the elbows flexed with the hands positioned at the height of the chin; (II) *biceps curl*: upright position  
90 with elbows flexed at  $90^{\circ}$ ; (III) *leg extension*: participants seated with the back supported by a bench  
91 reclined  $15^{\circ}$  respect to the vertical direction and the knees fully extended; and (IV) *calf raise*: upright

92 position with hips, knees, and ankles fully extended. All exercises were performed bilaterally, and  
93 participants were asked to maintain a constant breathing pattern during exercise.

94 \*\*Figure 1 near here\*\*

#### 95 *Intraocular pressure assessment*

96 An Icare rebound tonometer (Icare, TiolatOy, INC. Helsinki, Finland) was used to assess intraocular  
97 pressure at the different time points. The main advantages of this apparatus include that it is portable  
98 and hand-held, allows a rapid acquisition of intraocular pressure measures, it is very well tolerated  
99 by patients, and does not require the use of topical anaesthesia.<sup>21</sup> Following the manufacturer's  
100 instructions, participants were instructed to look at a distant target while performing the isometric  
101 exercise and intraocular pressure was continuously measured against the central cornea by an  
102 experienced optometrist (JV or BR). Intraocular pressure values were vocalized to a research assistant  
103 for data logging, obtaining a minimum of 24 measurements during the isometric effort (range of data  
104 points from 24 to 28 measurements).

105 The Icare tonometer cannot acquire intraocular pressure measurements at exact time intervals  
106 and due to the manual logging of the values we lacked exact measurement timestamps. For this  
107 reason, we devised a process to overcome these technical restrictions and obtain a set of equally  
108 distributed values at regular intervals with exact timestamps that we describe in the data processing  
109 subsection. A baseline intraocular pressure was measured before each exercise, and we obtained a  
110 recovery measurement 10 seconds after the exercise. We always measured the right eye. The same  
111 procedure for intraocular pressure assessment was conducted in a recent study.<sup>22</sup>

112 We were able to semi-continuously measure intraocular pressure due to the inherent  
113 characteristics of the tonometer and the exercise (static exercise with neutral neck position). This is  
114 the main novelty of this study compared to previous investigations, where the effects of different  
115 types of strength or endurance exercises were evaluated using a simple pre/post design.<sup>14,16,18,19,23</sup>



116 During the 2-minutes isometric exercise, intraocular pressure values were acquired in a continuous  
117 fashion.

### 118 **Data processing**

119 To overcome the timestamping issue and the lack of automatic logging restrictions of the rebound  
120 tonometer, we obtain a set of equally distributed intraocular pressure values at regular intervals using  
121 the following procedure based on multi-rate digital signal processing. **Specifically, we use sample**  
122 **rate conversion to adjust the sampling rate of the discrete sampled signal (i.e., the intraocular pressure**  
123 **signal) in order to obtain a new discretised version of the original continuous signal at a different**  
124 **rate.**<sup>24,25</sup> When measuring intraocular pressure using the rebound tonometer, we sampled the  
125 continuous intraocular pressure function at slightly irregular intervals. The values measured were the  
126 values of the intraocular pressure function at those moments-in-time. But since the function is  
127 continuous, when intraocular pressure values rise and fall between two pressures, intraocular pressure  
128 will always take all intermediate values between these two pressures. **As such we can reconstruct the**  
129 **intraocular pressure function from the sample measurements by treating the obtained samples as**  
130 **geometric points and then creating the necessary new points by polynomially interpolating those**  
131 **values to obtain 24 discrete values for the 2-minute period, i.e., every 5 seconds.**

### 132 *Statistical analysis*

133 First, we applied a mixed analysis of variance (ANOVA) to intraocular pressure values considering  
134 the exercise type (military press, leg extensions, biceps curl, calf raises), load (low and high), and  
135 point of measure (baseline, 1 to 24, and recovery [a total of 26 measurements]) as the within-  
136 participants factors, and sex (men vs women) as the between-participants factor. Additionally, linear  
137 regression analyses, considering the 24 measurements taken during the 2-minutes isometric efforts,  
138 were carried out in order to assess the intraocular pressure behaviour during exercise. The magnitude

139 of the differences was reported by the partial eta squared ( $\eta_p^2$ ) and, statistical significance was set at  
140 an alpha level of .05.

## 141 **Results**

142 The first set of analysis evidenced a statistically significant effect of load ( $F_{1, 25} = 257.39$ ,  $p < .001$ ,  
143  $\eta_p^2 = 0.906$ ), point of measure ( $F_{25, 625} = 20.57$ ,  $p < .001$ ,  $\eta_p^2 = 0.443$ ), whereas no differences were  
144 observed for the exercise type ( $F_{3, 75} = 1.17$ ,  $p = .326$ ) and sex ( $F_{1, 25} = 0.30$ ,  $p = .588$ ). There were  
145 also statistically significant differences for the interactions *exercise type x sex* ( $F_{3, 75} = 3.76$ ,  $p = .014$ ,  
146  $\eta_p^2 = 0.126$ ), *exercise type x point of measure* ( $F_{3, 75} = 2.13$ ,  $p < .001$ ,  $\eta_p^2 = 0.08$ ), *load x point of*  
147 *measure* ( $F_{25, 625} = 23.68$ ,  $p < .001$ ,  $\eta_p^2 = 0.476$ ), and *exercise type x load x point of measure x sex* ( $F_{75,$   
148  $1875 = 1.34$ ,  $p = .030$ ,  $\eta_p^2 = 0.050$ ). The rest of interaction did not reach statistical significance (all p-  
149 values  $> .05$ ) (**Figures 2 and 3**).

150 In the high-load condition, the intraocular pressure behaviour showed a fairly linear and  
151 positive increase as a function of time-on-task during isometric effort, with Pearson correlation  
152 coefficients of 0.94, 0.80, 0.74, and 0.57 for the military press, biceps curl, leg extension, and calf  
153 raises exercises, respectively.

154 \*\*Figure 2 near here\*\*

155 \*\*Figure 3 near here\*\*

## 156 **Discussion**

157

158 The current study was designed to assess the influence of the load, exercise type and participants' sex  
159 on the intraocular pressure changes induced by isometric exercises. Our data revealed that performing  
160 2-minutes of isometric exercise leading to muscular failure (high-load condition) promoted a  
161 progressive intraocular pressure rise, regardless of exercise type and participants' sex. Namely, the  
162 military press, biceps curl, leg extensions, and calf raise exercises caused a comparable increment of

163 intraocular pressure levels during the high-load condition (average intraocular pressure rise of ~ 25%,  
164 range: 22% to 26%) for both men and women. When exercise ceased, intraocular pressure returned  
165 to baseline levels in the subsequent 10 seconds. This result highlights that valuable information could  
166 be missed in experimental studies using pre/post designs and, thus, intraocular pressure assessment  
167 during exercise should be recommended. The outcomes of this study may have important implications  
168 for the management of ocular conditions in which maintaining stable intraocular pressure levels are  
169 desirable, especially for subjects at high risk for glaucoma onset or progression.

170         During the execution of dynamic resistance training exercises, the intraocular pressure  
171 changes have been shown to be dependent on the type of exercise performed. For example, a recent  
172 investigation reported a higher intraocular pressure rise during the bench press compared to the back  
173 squat when both exercises are performed against the same relative load.<sup>19</sup> Similarly, Rüfer et al.<sup>14</sup>  
174 found a significant intraocular pressure rise after performing 20 repetitions with the butterfly  
175 machine, whereas the execution of 20 repetitions during the leg curl exercise did not induce a  
176 significant change in intraocular pressure. The results of the present study provide additional evidence  
177 highlighting that intraocular pressure changes associated with isometric exercise are not exercise-  
178 dependent, at least for the exercise types included in this study. It should be noted that we chose  
179 exercises involving a considerable amount of muscle mass (military press or leg extensions),  
180 however, other exercises such as the bench press or back-squat may be more physically demanding  
181 and could promote a more abrupt intraocular pressure response. Indeed, recently published data from  
182 our laboratory suggest that the execution of 1-minute isometric squat exercise leading to muscular  
183 failure promoted an intraocular pressure rise of approximately 8 mmHg at the end of the 1-min effort,  
184 whereas the greatest intraocular pressure increment found in the current study was of approximately  
185 5 mmHg.<sup>22</sup> Taken together, it seems reasonable to discourage the execution of very challenging and  
186 prolonged isometric efforts when intraocular pressure fluctuations are undesirable (e.g., glaucoma  
187 patients) because they have shown to provoke a meaningful, instantaneous and cumulative intraocular

188 pressure rise. It should be noted that the average IOP rise observed in this study may be comparable  
189 with the nocturnal elevation of IOP occurring in healthy subjects (2 to 6 mmHg).<sup>26,27</sup> Therefore, the  
190 possible detrimental effects of the IOP rises caused by resistance training in healthy individuals  
191 require further investigation.

192         There are controversial results regarding the sex-related differences in intraocular pressure  
193 responses to resistance training. For example, two studies have shown that males and females have  
194 similar intraocular pressure responses to different dynamic resistance training exercises,<sup>13,28</sup> whereas  
195 Vera and colleagues<sup>22</sup> observed that men had a more accentuated intraocular pressure rise in  
196 comparison to women, although these differences only reached statistical significance during the last  
197 seconds of an isometric squat exercise performed until muscular failure. In this study, our results  
198 demonstrated that the intraocular pressure response to isometric exercise does not systematically  
199 differ between men and women. However, in the military press exercise, the differences were  
200 considerable (ES [90% confidence intervals] = 1.62 [0.73 – 2.43], with men exhibiting a more  
201 accentuated intraocular pressure increase in comparison to women. Of note, previous studies have  
202 commonly considered 1-minute of isometric effort,<sup>22,29</sup> showing a linear increase of intraocular  
203 pressure over time. The results of the present study highlight that the intraocular pressure is  
204 progressively increased until at least 2 minutes. However, it should be noted that during the execution  
205 of the high-load condition the intraocular pressure rise in the four exercises after the first minute was  
206 22% (i.e., last point of measure of the first minute), whereas the percentage of intraocular pressure  
207 increment when considering the last point of measure of the 2-minutes effort was 29%. This analysis  
208 shows that the slope of the intraocular pressure rise during isometric exercise is slower when  
209 considering longer periods of time. Notably, when leading to muscular failure during an isometric  
210 effort, 1-minute of isometric squat exercise promoted an intraocular pressure increment of 59%.<sup>22</sup> In  
211 order to test the influence of exercise duration on intraocular pressure, the same isometric exercise

212 should be tested when the time under tension is different (i.e., 1 vs 2 minutes), but matching the level  
213 of effort (e.g. both exercises leading to muscular failure).

#### 214 *Current limitations and implications for future research*

215 The current investigation is not exempt of limitations, and they must be acknowledged. First, our  
216 experimental sample consisted of healthy and physically active collegiate students, and the main  
217 implications of our findings are targeted at glaucoma patients or those at risk of glaucoma. Therefore,  
218 the external validity of the present findings in glaucoma subjects needs to be addressed in future  
219 investigations. Second, we tested the intraocular pressure changes during four different isometric  
220 exercises, however, the effects of isometric efforts during other exercises, including those performed  
221 while adopting head down position (yoga or crossfit) or lying face down (abdominal planks), should  
222 be assessed. Third, fitness level has demonstrated to play a mediating role on the intraocular pressure  
223 responses to physical exercise,<sup>19,30</sup> and thus, it is our hope that future studies will assess whether  
224 intraocular pressure changes induced by isometric exercise are dependent on participants' fitness  
225 level. Lastly, the practice of physical exercise has numerous health benefits,<sup>31</sup> including a  
226 neuroprotective effect in different eye disorders.<sup>5</sup> **Based on this, it is necessary to determine the most  
227 beneficial physical activity in order to preserve the ocular health. In this regard, isometric resistance  
228 training may not be the most pertinent type of physical exercise, although it is clear that the health  
229 benefits associated with physical activity indicate that adopting a sedentary behaviour needs to be  
230 avoided.**

#### 231 **Conclusions**

232 The execution of 2-minutes isometric exercises leading to muscular failure promotes an instantaneous  
233 and progressive intraocular pressure rise in healthy adults, being these changes independent of the  
234 tested exercise (military press, leg extensions, biceps curl, calf raises). When physical effort ceased,  
235 intraocular pressure rapidly returned to baseline levels (within 10 seconds). Overall, there were no

236 meaningful differences between men and women in intraocular pressure responses to physical  
237 exercise with the only exception being the military press exercise in which men showed a greater  
238 intraocular pressure rise in comparison to women. The present outcomes may be of interest for eye  
239 care specialists, since the performance of isometric efforts should be discouraged when stable  
240 intraocular pressure levels are desirable, especially for individuals at high risk for glaucoma onset or  
241 progression. The inclusion of glaucoma patients in future investigations is needed to explore the  
242 generalizability of the current findings.

243

#### 244 **Acknowledgments**

245 The authors thank all the participants who selflessly collaborated in this research. The authors report  
246 no conflicts of interest and have no proprietary interest in any of the materials mentioned in this  
247 article. This research received no specific grant from any funding agency in the public, commercial,  
248 or not-for-profit sectors. This study complied with the Declaration of Helsinki and was approved the  
249 University of Granada Institutional Review Board (IRB approval: 438/CEIH/2017). All participants  
250 included in this study signed an informed consent form.

251

#### 252 **References**

- 253 1. Piercy KL, Troiano RP, Ballard RM, et al. The Physical Activity Guidelines for Americans.  
254 JAMA 2018;9762:1–9.
- 255 2. Hillman CH, Erickson KI, Kramer AF. Be smart, Exercise your Heart: Exercise Effects on  
256 Brain and Cognition. Nat Rev Neurosci 2008;9:58–65.
- 257 3. Mcguinness MB, Le J, Mitchell P, et al. Physical Activity and Age-related Macular  
258 Degeneration: A Systematic Literature Review and Meta-analysis. Am J Ophthalmol

- 259 2017;180:29–38.
- 260 4. Selin JZ, Orsini N, Lindblad BE, Wolk A. Long-Term Physical Activity and Risk of Age-  
261 Related Cataract: A Population-Based Prospective Study of Male and Female Cohorts.  
262 *Ophthalmology* 2015;122:274–80.
- 263 5. Zhu MM, Lai JS, Choy BN, et al. Physical Exercise and Glaucoma: A Review on the Roles  
264 of Physical Exercise on Intraocular Pressure Control, Ocular Blood Flow Regulation,  
265 Neuroprotection and Glaucoma-Related Mental Health. *Acta Ophthalmol* 2018;96:676–91.
- 266 6. Tham YC, Li X, Wong TY, et al. Global Prevalence of Glaucoma and Projections of  
267 Glaucoma Burden Through 2040: A Systematic Review and Meta-Analysis. *Ophthalmology*  
268 2014;121:2081–90.
- 269 7. Weinreb RN, Aung T, Medeiros FA. The Pathophysiology and Treatment of Glaucoma.  
270 *JAMA* 2014;311:1901–11.
- 271 8. Kiel J, Hollingsworth M, Rao M, et al. Ciliary Blood Flow and Aqueous Humor Production.  
272 *Prog Retin Eye Res* 2011;30:1–17.
- 273 9. De Moraes CG, Juthani VJ, Liebmann JM, et al. Risk Factors for Visual Field Progression in  
274 Treated Glaucoma. *Arch Ophthalmol* 2011;129:562–8.
- 275 10. Wylęgała A. The Effects of Physical Exercises on Ocular Physiology. *J Glaucoma*  
276 2016;25:e843–9.
- 277 11. Vera J, Jiménez R, Redondo B, et al. Acute Intraocular Pressure Responses to High-  
278 Intensity Interval-Training Protocols in Men and Women. *J Sports Sci* 2019;37:803–9.
- 279 12. Najmanova E, Pluhacek F, Botek M. Intraocular Pressure Response to Moderate Exercise  
280 During 30-Min Recovery. *Optom Vis Sci* 2016;93:281–5.
- 281 13. Vera J, Jiménez R, Redondo B, et al. Effect of the Level of Effort During Resistance

282 Training on Intraocular Pressure. *Eur J Sport Sci* 2019;19:394–401.

283 14. Rüfer F, Schiller J, Klettner A, et al. Comparison of the Influence of Aerobic and Resistance  
 284 Exercise of the Upper and Lower Limb on Intraocular Pressure. *Acta Ophthalmol*  
 285 2014;92:249–52.

286 15. Vera J, Jiménez R, García JA, et al. Baseline Intraocular Pressure Is Associated with  
 287 Subjective Sensitivity to Physical Exertion in Young Males. *Res Q Exerc Sport* 2018;89:25–  
 288 37.

289 16. Vieira G, Oliveira H, de Andrade D, et al. Intraocular Pressure Variation During Weight  
 290 Lifting. *Arch Ophthalmol* 2006;124:1251–4.

291 17. Vera J, Raimundo J, García-Durán B, et al. Acute Intraocular Pressure Changes During  
 292 Isometric Exercise and Recovery: The Influence of Exercise Type and Intensity, and  
 293 Participant´s Sex. *J Sports Sci* 2019.

294 18. Vera J, Garcia-Ramos A, Jiménez R, Cárdenas D. The Acute Effect of Strength Exercises at  
 295 Different Intensities on Intraocular Pressure. *Graefe´s Arch Clin Exp Ophthalmol*  
 296 2017;255:2211–7.

297 19. Vera J, Jiménez R, Redondo B, et al. Fitness Level Modulates Intraocular Pressure  
 298 Responses to Strength Exercises. *Curr Eye Res* 2018;43:740–6.

299 20. Drust B, Waterhouse J, Atkinson G, et al. Circadian Rhythms in Sports Performance - An  
 300 Update. *Chronobiol Int* 2005;22:21–44.

301 21. Pakrou N, Gray T, Mills R, et al. Clinical Comparison of the Icare Tonometer and  
 302 Goldmann Applanation Tonometry. *J Glaucoma* 2008;17:43–7.


303 22. Vera J, Jiménez R, Redondo B, et al. Investigating the Immediate and Cumulative Effects of  
 304 Isometric Squat Exercise for Different Weight Loads on Intraocular Pressure: A Pilot Study.



- 305 Sports Health 2019;11:247–53.
- 306 23. Vera J, Jiménez R, Redondo B, et al. Intraocular Pressure Responses to Maximal Cycling  
307 Sprints Against Different Resistances: The Influence of Fitness Level. *J Glaucoma*  
308 2017;26:881–7.
- 309 24. Crochiere RE, Rabiner LR. *Multirate Digital Signal Processing*. Vol 18. Prentice-Hall  
310 Englewood Cliffs, NJ; 1983.
- 311 25. Oppenheim AV, Buck JR, Schafer RW. *Discrete-Time Signal Processing, Volume 2*.  
312 Englewood Cliffs: Prentice Hall; 1989.
- 313 26. Liu JH, Zhang X, Kripke DF, Weinreb RN. Twenty-Four-Hour Intraocular Pressure Pattern  
314 Associated with Early Glaucomatous Changes. *Investig Ophthalmol Vis Sci* 2003;44:1586–  
315 90.
- 316 27. Agnifili L, Mastropasqua R, Frezzotti P, et al. Circadian Intraocular Pressure Patterns in  
317 Healthy Subjects, Primary Open Angle and Normal Tension Glaucoma Patients with a  
318 Contact Lens Sensor. *Acta Ophthalmol* 2015;93:14–21.
- 319 28. Chromiak J, Abadie B, Braswell R, et al. Resistance Training Exercises Acutely Reduce  
320 Intraocular Pressure in Physically Active Men and Women. *J Strength Cond Res*  
321 2003;17:715–20.
- 322 29. Bakke EF, Hisdal J, Semb SO. Intraocular Pressure Increases in Parallel with Systemic  
323 Blood Pressure During Isometric Exercise. *Investig Ophthalmol Vis Sci* 2009;50:760–4.
- 324 30. Vera J, Jiménez R, Redondo B, et al. Effect of a Maximal Treadmill Test on Intraocular  
325 Pressure and Ocular Perfusion Pressure: The Mediating Role of Fitness Level. *Eur J*  
326 *Ophthalmol* 2019:1120672119832840.
- 327 31. Warburton DE, Bredin SS. Reflections on Physical Activity and Health: What Should We

329 **Figure legends**


330 **Figure 1.** Schematic illustration of the experimental procedure (from left to right: military press,  
331 biceps curl, leg extension, and calf raise). The same protocol was repeated in eight different  
332 occasions (4 exercises x 2 loads), and the order of the exercises and loads was randomized. A 10  
333 minutes break was given between two sets. The recovery measurement of intraocular pressure was  
334 taken 10 seconds after the exercise cessation.



335


336 **Figure 2.** Effects of performing 2-minutes of isometric military press (panel A), biceps curl (panel  
337 B), leg extensions (panel C), and calf raises (panel D) against the maximum load that participants  
338 could hold for approximately 2 minutes in an isometric condition (high-load) and without applying  
339 any additional load (low-load) on intraocular pressure. The recovery (REC) value represents the

340 measurement taken 10 seconds after the exercise cessation. Error bars show the 95% confidence  
341 intervals.



342

343 **Figure 3.** Standardized differences (Cohen's d effect size) with the corresponding 90% confidence  
344 intervals in the intraocular pressure changes (average value of the 2-min isometric efforts in the low  
345 and high-load conditions) between men and women when performing the four different isometric  
346 exercises.



347