

Note: This article will be published in a forthcoming issue of the *International Journal of Sport Nutrition and Exercise Metabolism*. This article appears here in its accepted, peer-reviewed form; it has not been copyedited, proofed, or formatted by the publisher.

Section: Original Research

Article Title: Pre-Season Body Composition Adaptations in Elite Caucasian and Polynesian Rugby Union Athletes

Authors: Adam J. Zemski¹, Shelley E. Keating², Elizabeth M. Broad³, Damian J. March⁴, Karen Hind⁵, and Gary J. Slater¹

Affiliations: ¹School of Health and Sport Sciences, University of the Sunshine Coast, Maroochydore, Australia. ²School of Human Movement and Nutrition Sciences, The University of Queensland, St Lucia, Australia. ³US Olympic Committee, US Paralympics, Chula Vista, CA, USA. ⁴Fiji Rugby Union, Suva, Fiji. ⁵Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, West Yorkshire, UK.

Running Head: Pre-season body composition changes in rugby union

Journal: *International Journal of Sport Nutrition and Exercise*

Acceptance Date: April 16, 2018

©2018 Human Kinetics, Inc.

DOI: <https://doi.org/10.1123/ijsnem.2018-0059>

Title of the article

Pre-season body composition adaptations in elite Caucasian and Polynesian rugby union athletes

Submission type: Original research

Authors

Adam J Zemski
School of Health and Sport Sciences, University of the Sunshine Coast, Maroochydore, Australia

Shelley E Keating
School of Human Movement and Nutrition Sciences, The University of Queensland, St Lucia, Australia

Elizabeth M Broad
US Olympic Committee, US Paralympics, Chula Vista, CA, USA

Damian J Marsh
Fiji Rugby Union, Suva, Fiji

Karen Hind
Institute for Sport, Physical Activity and Leisure, Leeds Beckett University, Leeds, West Yorkshire, UK

Gary J Slater
School of Health and Sport Sciences, University of the Sunshine Coast, Maroochydore, Australia
Australian Rugby Union, Sydney, Australia

Contact details of corresponding author

Adam J Zemski
School of Health and Sport Sciences
Faculty of Science, Health, Education and Engineering
University of the Sunshine Coast
Maroochydore DC, QLD, 4558, Australia
Email: ajz006@student.usc.edu.au

Preferred running head

Pre-season body composition changes in rugby union

Abstract word count:

245 words

Text only word count (before referencing):

2971 words

Number of tables and figures

Tables – 3

Figures – 3

Abstract

During pre-season training, rugby union (RU) athletes endeavour to enhance physical performance characteristics that are aligned with on-field success. Specific physique traits are associated with performance, therefore body composition assessment is routinely undertaken in elite environments. This study aimed to quantify pre-season physique changes in elite RU athletes with unique morphology and divergent ethnicity. Twenty-two Caucasian and Polynesian professional RU athletes received dual-energy X-ray absorptiometry (DXA) assessments at the beginning and conclusion of an 11-week pre-season. Interactions between on-field playing position and ethnicity in body composition adaptations were explored, and the least significant change (LSC) model was used to evaluate variations at the individual level. There were no combined interaction effects with the variables position and ethnicity, and any body composition measure. After accounting for baseline body composition, Caucasians gained more lean mass during the pre-season than Polynesians (2425 ± 1303 g vs 1115 ± 1169 g; $F=5.4$, $p=0.03$). Significant main effects of time were found for whole body and all regional measures with fat mass decreasing ($F=31.1-52.0$, $p<0.01$), and lean mass increasing ($F=12.0-40.4$, $p<0.01$). Seventeen athletes (9 Caucasian, 8 Polynesian) had a reduction in fat mass, and 8 athletes (6 Caucasian, 2 Polynesian) increased lean mass. This study describes significant and meaningful physique changes in elite RU athletes during a pre-season period. Given the individualised approach applied to athletes in regards to nutrition and conditioning interventions, a similar approach to that used in this study is recommended to assess physique changes in this population.

Key words: dual-energy X-ray absorptiometry, fat mass, lean mass, training, ethnicity.

there are limits to relying on anthropometric measures for estimating body composition in athletes, given the regression equations haven't been validated for use in RU, or to track changes in body composition (Silva et al., 2009; Zemski et al., 2017). Over recent years, the use of dual-energy X-ray absorptiometry (DXA) for body composition assessment in elite RU has increased (Lees et al., 2017; Zemski et al., 2015). This technology provides an in-depth analysis of whole body and regional bone mineral content (BMC), FM and LM, and is recognised as a valid and precise body composition assessment tool (Harley et al., 2009; Van der Ploeg et al., 2003) when client presentation is standardised in accordance with best practice guidelines (Nana et al., 2015).

In recent years there has been a surge in the number of Polynesian athletes securing professional RU contracts. One study has investigated three-compartment body composition in Polynesian RU players and reported different distributions of regional FM and LM (Zemski et al., 2015). In non-athletes, large differences in physique have been reported between Caucasian and Polynesian individuals, with Polynesians having more LM and greater LM:FM ratios (Rush et al., 2004; Swinburn et al., 1996; Swinburn et al 1999). To date, no study has explored differences in physique adaptations to training by ethnicity in RU. Therefore, the aim of this study was to investigate pre-season team and individual athlete DXA body composition adaptations in elite RU athletes, with sub-group analysis to compare changes between Polynesian and Caucasian individuals.

Methods

Participants

Twenty-two professional male RU athletes were recruited via their involvement in a single Australian Super Rugby franchise, which is the premier professional RU competition in the southern hemisphere. All athletes provided informed consent to participate in the study,

Further, given there is no gold standard assessment of energy intake, any method employed would be subject to considerable error, particularly over a long period in an athletic population (Magkos & Yannakoulia, 2003). Such information may have provided further insight into the underlying reasons for the observed individual physique changes, and warrants consideration when appropriate and reliable technologies are available. Also, researchers were not made aware of individual athlete body composition goals over the pre-season, which may have added to the interpretation of results. Secondly, off-season changes and events likely to influence body composition were not taken into consideration when interpreting the results. An appreciation of such changes would allow for a more meaningful interpretation of the pre-season adaptations in the context of each individual athlete. Finally, associations between body composition and physical performance changes were not explored in this study. Future research investigating the association between physique adaptations and specific performance measures and fitness traits over a pre-season would be of great interest, in particular how these changes impact game performance in-season.

In conclusion, we identified significant whole body and regional body composition changes in elite RU athletes during a pre-season period, at both the team and individual level. Practitioners are encouraged to take an individualised approach to the interpretation of adaptations when tracking physique variables longitudinally, for which knowledge of LSC data is required. Future work exploring ethnicity differentiated body composition changes across the entire season, including the post-season period, would provide practitioners with valuable information allowing for a more personalised approach to athlete training and dietary interventions.

Hangartner, T. N., Warner, S., Braillon, P., Jankowski, L., & Shepherd, J. (2013). The official positions of the international society for clinical densitometry: acquisition of dual-energy x-ray absorptiometry body composition and considerations regarding analysis and repeatability of measures. *Journal of Clinical Densitometry*, 16, 520-536.

Harley, J. A., Hind, K., O'Hara, J., & Gross, A. (2009). Validation of the skin-fold thickness method and air displacement plethysmography with dual energy X-ray absorptiometry for the estimation of % body fat in professional male rugby football league players. *International Journal of Body Composition Research*, 7, 7-14.

Kouri, E. M., Pope, H. G., Katz, D. L., & Oliva P. (1995). Fat-free mass index in users and nonusers of anabolic-androgenic steroids. *Clinical Journal of Sports Medicine*, 5, 223-228.

Lees, M. J., Oldroyd, B., Jones, B., Brightmore, A., O'Hara, J. P., Barlow, M. J., Till, K., & Hind, K. (2017). Three-compartment body composition changes in professional rugby union players over one competitive season: A team and individual approach. *Journal of Clinical Densitometry*, 20, 50-57.

Magkos, F., & Yannakoulia, M. (2003). Methodology of dietary assessment in athletes: concepts and pitfalls. *Current Opinion in Clinical Nutrition and Metabolic Care*, 6, 539-549.

Nana, A., Salter, G. J., Stewart, A. D., & Burke, L. M. (2015). Methodology review: using dual-energy X-ray absorptiometry (DXA) for the assessment of body composition in athletes and active people. *International Journal of Sport Nutrition and Exercise Metabolism*, 25, 198-215.

Olds, T. (2001) The evolution of physique in male rugby union players in the twentieth century. *Journal of Sports Sciences*, 19, 253-262.

Quarrie, K. L., & Hopkins, W. G. (2007). Changes in player characteristics and match activities in Bledisloe Cup rugby union from 1972 to 2004. *Journal of Sports Sciences*, 25, 895-903.

Roberts, S. P., Trewartha, G., Higgitt, R. J., El-Abd, J., & Stokes, K. A. (2008). The physical demands of elite English rugby union. *Journal of Sports Sciences*, 26, 825-833.

Rush, E. C., Freitas, I., & Plank, L. D. (2009). Body size, body composition and fat distribution: comparative analysis of European, Maori, Pacific Island and Asian Indian adults. *British Journal of Nutrition*, 102, 632-641.

Rush, E., Plank, L., Chandu, V., Lалу, M., Simmons, D., Swinburn, B., & Yajnik, C. (2004). Body size, body composition, and fat distribution: a comparison of young New Zealand men of European, Pacific Island, and Asian Indian ethnicities. *New Zealand Medical Journal*, 117, U1203.

Selkirk, G. A., & McLellan, T. M. (2001). Influence of aerobic fitness and body fatness on tolerance to uncompensable heat stress. *Journal of Applied Physiology (1985)*, 91, 2055-2063.




Figure 1: Individual whole body and regional changes in bone mineral content by the least significant change (LSC) previously determined (Zemski et al., 2018) over a pre-season in elite rugby union athletes. Dashed lines indicate LSC-95% CI same day precision (technical error). Dotted lines indicate LSC-95% CI consecutive day precision (technical error and biological variation).




Figure 2: Individual whole body and regional changes in fat mass by the least significant change (LSC) previously determined (Zemski et al., 2018) over a pre-season in elite rugby union athletes. Dashed lines indicate LSC-95% CI same day precision (technical error). Dotted lines indicate LSC-95% CI consecutive day precision (technical error and biological variation).




Figure 3: Individual whole body and regional changes in lean mass by the least significant change (LSC) previously determined (Zemski et al., 2018) over a pre-season in elite rugby union athletes. Dashed lines indicate LSC-95% CI same day precision (technical error). Dotted lines indicate LSC-95% CI consecutive day precision (technical error and biological variation).

Table 1: Short-term prevision and corresponding SC in resistance trained athletes using the same Hologic Discovery A (Zemski et al., 2018)

	Same Day Technical Error				Consecutive Days Technical Error & Biological Variation			
	Precision		LSC-95% CI		Precision		LSC-95% CI	
	RMS-SD	%CV	RMS-SD	%CV	RMS-SD	%CV	RMS-SD	%CV
Whole body								
BMC (g)	21.1	0.6	59.0	1.7	25.2	0.7	80.5	1.9
Fat Mass (g)	238.4	1.8	660.4	5.1	455.2	2.9	1261.0	8.0
Lean Mass (g)	222.7	0.3	616.8	0.9	752.0	1.1	2083.0	3.2
Arms								
BMC (g)	5.6	1.1	15.5	3.0	6.8	1.3	18.9	3.7
Fat Mass (g)	43.5	2.5	120.5	6.8	89.1	5.3	246.8	14.5
Lean Mass (g)	101.1	1.2	279.9	3.3	154.1	1.9	426.7	5.2
Trunk								
BMC(g)	9.7	0.8	27.0	2.2	9.8	0.9	27.1	2.6
Fat Mass (g)	123.7	2.2	342.5	6.0	221.3	3.6	612.9	9.9
Lean Mass (g)	319.4	0.8	884.7	2.1	678.7	1.9	1880.0	4.1
Legs								
BMC(g)	20.2	1.5	56.1	4.2	18.6	1.5	51.6	4.1
Fat Mass (g)	146.0	2.7	404.4	7.5	230.7	3.4	639.1	9.5
Lean Mass (g)	335.6	1.1	929.6	3.0	406.5	1.5	1126.0	4.1

RMS-SD = root-mean-square standard deviation; CV = coefficient of variance; LSC = least significant change;
 BMC = bone mineral content

Table 2: Differences in surface anthropometry measures and indices, and dual-energy X-ray absorptiometry measured total and regional body composition characteristics of elite rugby union athletes over the course of a pre-season based on position and ethnicity.

	Position (n=22)				Ethnicity (n=22)			
	Forwards (n=11)		Backs (n=11)		Caucasians (n=11)		Polynesians (n=11)	
	Start Pre-Season	End Pre-Season	Start Pre-Season	End Pre-Season	Start Pre-Season	End Pre-Season	Start Pre-Season	End Pre-Season
Age (years)	22.9 ± 3.5	-	22.8 ± 3.0	-	22.1 ± 2.4	-	23.5 ± 3.8	-
Stature (cm) ^c	191.3 ± 7.5	-	182.2 ± 6.9	-	189.4 ± 8.7	-	184.1 ± 7.6	-
Mass (kg) ^{b,c}	112.5 ± 7.6	112.1 ± 7.6	90.5 ± 8.6	90.4 ± 8.1	101.2 ± 14.3	101.7 ± 14.0	101.8 ± 13.9	100.8 ± 13.6
FFMI (kg/m ²) ^{b,c}	26.1 ± 1.2	26.6 ± 1.1	23.8 ± 1.2	24.3 ± 1.1	24.4 ± 1.3	25.1 ± 1.2	25.5 ± 1.8	25.8 ± 1.9
WB BMC (g) ^c	4352 ± 439	4377 ± 437	3618 ± 379	3637 ± 364	4003 ± 553	4027 ± 548	3966 ± 571	3987 ± 569
WB FM (g) ^{b,c}	19629 ± 3879	17166 ± 3837	13438 ± 2723	11449 ± 1872	15495 ± 4839	13338 ± 4353	17572 ± 4214	15278 ± 3897
WB LM (g) ^{a,b,c}	91087 ± 5489	92912 ± 5711	75598 ± 6971	77312 ± 6436	84005 ± 10306	86430 ± 10447	82680 ± 10173	83795 ± 10431
Arms BMC (g) ^{b,c}	662 ± 76	661 ± 78	541 ± 68	535 ± 67	600 ± 106	602 ± 107	603 ± 85	594 ± 88
Arms FM (g) ^{b,c}	2287 ± 426	2038 ± 415	1470 ± 228	1304 ± 157	1759 ± 535	1601 ± 489	1999 ± 554	1741 ± 494
Arms LM (g) ^c	11698 ± 1098	12162 ± 928	9742 ± 1314	10198 ± 1454	10706 ± 1672	11191 ± 1636	10734 ± 1452	11169 ± 1556
Trunk BMC (g) ^{b,c}	1361 ± 179	1381 ± 193	1116 ± 117	1131 ± 127	1270 ± 186	1282 ± 196	1207 ± 205	1231 ± 219
Trunk FM (g) ^{b,c}	8594 ± 2392	7179 ± 2224	5419 ± 1191	5370 ± 824	6326 ± 2183	5079 ± 1910	7687 ± 2627	6470 ± 2293
Trunk LM (g) ^{b,c}	43339 ± 3136	44282 ± 3509	36259 ± 2564	36964 ± 2832	40498 ± 4305	41729 ± 4492	39100 ± 4939	39518 ± 5187
Legs BMC (g) ^c	1623 ± 183	1624 ± 175	1372 ± 162	1383 ± 148	1485 ± 212	1491 ± 198	1510 ± 222	1517 ± 212
Legs FM (g) ^{b,c}	7570 ± 1745	6765 ± 1755	5527 ± 1373	4757 ± 983	6348 ± 2191	5584 ± 2086	6749 ± 1529	5938 ± 1357
Legs LM (g) ^{b,c}	31977 ± 1872	32372 ± 1916	26056 ± 3616	26615 ± 3881	29126 ± 4553	29791 ± 4755	28908 ± 3888	29196 ± 3763

FFMI = fat-free mass index; WB = whole body; BMC = bone mineral content; FM = fat mass; LM = lean mass

Data presented as Mean ± Standard Deviation, significance set at 0.05

^a Significant interaction between time and ethnicity

^b Significant main effect for time

^c Significant difference between forwards and backs

