

**Linking Inter-annual River Flow River Variability Across
New Zealand to the Southern Annular Mode, 1979-2011**

Journal:	<i>Hydrological Processes</i>
Manuscript ID	HYP-16-0491.R2
Wiley - Manuscript type:	Research Article
Date Submitted by the Author:	n/a
Complete List of Authors:	Li, Ning; University of Auckland, School of Environment McGregor, Glenn; Durham University, Geography
Keywords:	Hydroclimatological variability, Southern Annular Mode, New Zealand, Climate and river flow

SCHOLARONE™
Manuscripts

Review

Linking Inter-annual River Flow River Variability Across New Zealand to the Southern Annular Mode, 1979-2011

N. Li¹, G. R. McGregor²

¹School of Environment, the University of Auckland, Auckland, New Zealand

²Department of Geography, Durham University, Durham, United Kingdom

Abstract

River flow constitutes an important element of the terrestrial branch of the hydrological cycle, yet knowledge regarding the extent to which its variability, at a range of time scales, is linked to a number of modes of atmospheric circulation is meagre. This is especially so in the Southern Hemisphere where strong candidates, such as El Niño Southern Oscillation (ENSO) and the Southern Annular Mode (SAM), for influencing climate and thus river flow variability can be found. This paper presents the results of an analysis of the impact of the SAM on winter and summer river flow variability across New Zealand, purposefully controlling for the influence of ENSO and the tendency for the SAM to adopt a positive phase over the last 10 – 20 years. Study results, based on identifying hydrological regions and applying circulation-to-environment and environment-to-circulation approaches commonly used in synoptic climatology, reveal a seasonal asymmetry of the response of river flow variability to the SAM; winter flows demonstrate a higher degree of statistical association with the SAM compared to summer flows. Further, because of the complex orography of New Zealand and its general disposition normal to zonal flows of moisture bearing winds, there are intra-seasonal spatial variations in river flow SAM associations with clear rain shadow effects playing out in resultant river flow volumes. The complexity of SAM river flow associations found in this study warns against using indices of large scale modes of atmospheric circulation as blunt tools for hydroclimatological prediction at scales beyond hydroclimatological regions or areas with internal hydrological consistency.

Keywords: river flow, climate, hydroclimatological variability, Southern Annual Mode, atmospheric circulation.

1. INTRODUCTION

River flow constitutes an important element of the terrestrial branch of the hydrological cycle. Understanding its inter-annual variability and associated driving mechanisms is crucial for at least two reasons. Firstly, variations in the strength of the terrestrial branch of the hydrologic cycle bears important consequences for global climate change via modification of hygrothermal properties at the catchment scale (Dettinger and Diaz 2000; Chiew and McMahon 2002; Prowse et al., 2011). Secondly, strong predictive skill derived from knowledge of climate-river flow co-variability helps inform water management practices. This is especially so in a country such as New Zealand where major sectors of the economy, such as agriculture and energy production, are water resource dependent and climatic variability presents a considerable risk management problem.

Notwithstanding the importance of understanding catchment scale processes for water resource management, large-scale modes of climatic variability are increasingly recognised as critical for determining river flow variability through their impact on regional climate variability. Amongst these modes, such as the Pacific Decadal Oscillation (PDO), the Pacific/North American Pattern (PNA), El Nino Southern Oscillation (ENSO) and the North Atlantic Oscillation (NAO), is the Southern Annular Mode (SAM) which, in the Southern Hemisphere, exerts an influence on hydroclimate at the intra-annual to inter-annual time scales (Lim and Hendon, 2015; Mariani and Fletcher, 2016; Kiem and Verdon-Kidd 2009; Raut et al., 2014; Swart et al., 2015).

The SAM (also known as the Antarctic Oscillation) is an important mode of atmospheric variability (Sallée et al., 2010; Thompson and Wallace, 2000) in the extratropical Southern Hemisphere. It involves periodic fluctuations in zonally averaged sea level pressure (SLP) between two latitudinal bands encompassing the subtropical high at around 40°S and the circumpolar trough at around 60°S. Positive polarity of the SAM is characterized by an amplified pressure gradient between the two aforementioned zonal bands, poleward migration of the mid-latitude jet stream and associated surface westerlies. It is also known to generate ocean circulation anomalies such that during a positive SAM, equatorward Ekman drift at all longitudes of the circumpolar ocean and poleward Ekman drift at around 30°S is strengthened, which in turn leads to anomalies in oceanic heat transport (Hall and Visbeck 2002). Opposite conditions prevail in the negative polarity phase of SAM. Several investigators have noted the trend towards a more positive phase of the SAM over recent decades (Marshall, 2003; Swart et al., 2015) but an unequivocal explanation for this observation remains elusive (Chang and Johnson, 2015; Ding et al., 2015; Solomon and Polvani., 2016).

An 'annular' or ring like feature depicts the annual mean state of the SAM. On seasonal timescales however, a zonally asymmetric structure emerges, which reaches its maximum in austral winter and spring with a minimum in summer (Fan, 2007). Fogt et al. (2012) argue this zonal asymmetry may alter regional SAM-temperature relationships. With regards to this, and despite its potential importance for determining inter-annual climate and thus hydrological variability, the nature of SAM-induced climate anomalies across New Zealand have been explored by only a few investigators. For example, Kidston et al. (2009) examined the climate anomaly response to the seasonality of SAM and found distinctly different outcomes for wind

1
2
3 direction, rainfall and temperature between summer and winter. Renwick (2011) identified a
4 connection between the negative SAM phase and a higher occurrence of trough-like weather
5 conditions while more zonal and blocking weather was associated with the SAM positive phase.
6 This work corroborates the earlier findings of Sinclair et al. (1997) who showed that the 1st EOF
7 Southern Hemisphere pressure pattern, which resembles SAM like pressure anomalies,
8 modulates cyclone activity in the Southern Ocean with increasing westerlies near 55°S
9 accompanied by more cyclones in high latitudes and fewer in middle latitudes, and vice versa.
10 At the annual time scale, Ummenhofer et al. (2007) found that during positive SAM years, New
11 Zealand as a whole is drier than normal, with the exception of the northernmost tips of both
12 islands. This they attribute to enhanced north-easterly flows over the country while during
13 negative SAM years the precipitation and wind anomaly patterns are reversed. Following on
14 from this work, Ummenhofer et al. (2009) have suggested that up to 80% of the reduction in
15 precipitation observed across New Zealand can be accounted for by the positive trend in the
16 SAM since the mid-1970s.

17
18
19
20
21 Given the extent to which SAM clearly exerts an influence on climatic variability it is surprising
22 that the linkage between the SAM and river flow across New Zealand remains unexplored.
23 Rather to date, modes of climatic variability of longer periodicity such as the IPO and ENSO
24 have received more attention in terms of New Zealand hydrology (McKerchar and Pearson 1994;
25 McKerchar et al., 1998; McKerchar and Henderson, 2003; McKerchar et al., 2010;; Mosley
26 2000), with the latter becoming the basis of seasonal river flow forecasts for New Zealand
27 (Bardsley, 2016; Pearson, 2008; Singh, 2016). The main aim of this paper is therefore to
28 investigate the extent to which seasonal austral summer (DJF) and winter (JJA) river flow at the
29 inter-annual time scale is linked to variations in the SAM. The chief justification for this is that
30 currently SAM possesses superior predictability to precipitation at intra-seasonal to seasonal
31 time scales (Hendon et al., 2014; Lim and Hendon, 2015; Marshall et al., 2012; Zheng et al.,
32 2015) as the accuracy of rainfall prediction decreases as the lead time increases beyond
33 hydrometeorological timescales of 5-10 days (Jang and Hong, 2014; Shukla et al., 2013). This
34 bears implications for the reliability of seasonal river flow forecasts and for assessments of the
35 likely impacts of climate change on precipitation (Lim et al., 2016) as determined by variability
36 and change of major modes of climatic variability. Further, as there is strong evidence that river
37 flow characteristics are modulated by modes of atmospheric-ocean circulation variability (Marti
38 et al., 2010; Rimbu et al., 2016; Ward et al., 2016), the efficacy of rainfall-runoff curves for river
39 flow prediction may be somewhat compromised as they are based on a deterministic modelling
40 of the long-term or average relationship between rainfall and runoff as opposed to a modelling
41 of hydrological response conditioned on hydrologically relevant phases of climate variability,
42 Indeed Beckers et al. (2016) have noted the general improvement of river flow prediction
43 achieved when ensemble stream prediction is conditioned on ENSO state. This resonates with
44 the general move to a seamless approach to both weather/climate and hydrological forecasting
45 based on predictors that transcend a range of timescales (Brown et al., 2012; Tang et al., 2016;
46 Yuan et al., 2014) with associated gains in predictability (Hoskins, 2012; Yuan et al., 2015). For
47 these reasons and with early warning systems for extremes in mind, particularly in relation to
48 managing hydroclimatological related risks, a number of studies have focused on detecting
49 SAM signals in extreme event chronologies (Min et al., 2013) such as wildfire (Holz and Veblen,
50
51
52
53
54
55
56
57
58
59
60

2011; Mariani and Fletcher, 2016), drought (Cai et al., 2011), tropical cyclones (Diamond and Renwick, 2014) and flood producing extreme rainfall (King et al., 2014). Moreover because of its predictability and its derivation from pressure fields, which in comparison to other climate fields can be recovered more easily from historic records (Jones et al., 2009), SAM can be used effectively for climate related reconstructions such as stream flow, weather types and glacier behaviour (Allen et al., 2015; Jiang et al., 2013; Purdie et al., 2011). Overall then, seeking to increase knowledge relating to the importance of modes of atmospheric circulation variability, for hydrological processes at timescales beyond the hydrometeorological, can provide insights into the causes of seasonal to decadal hydrological variability thus facilitating not only an explanation of a region's hydrological characteristics within a broader hydroclimatological context but concomitantly open up the potential for seasonal to decadal river flow prediction. Understanding SAM river flow associations also strikes a chord with the views expressed by a range of users of climate and hydrological products relating to the need to access information on climate and water resource associations at a range of temporal and spatial scales, in order to build as a complete picture as possible of the drivers of hydrological variability (Dravitzki and McGregor, 2011; Pappenberger et al., 2013; Purdie and Bardsley, 2010).

Associated with the paper's overall aim are a number of specific objectives. These include the assembly of the requisite SAM, climate and river flow data sets, the establishment of river flow responses under contrasting SAM phases and the atmospheric circulation anomaly patterns associated with anomalously high and low river flow conditions. Accordingly this paper is organised in the following way. Section 2 describes the data sources, followed by an explanation of the so-called environment-to-circulation and circulation-to-environment approaches for investigating SAM river flow linkages. Section 3 presents a description of river flow characteristics and outlines the development of a river flow regionalization which is used as a spatial framework for the exploration of hydro-climate linkages, the focus of which will be section 4. A discussion of results will be presented in section 5 and evaluated in the context of the current understanding of hydro-climate linkages in the broader New Zealand region and beyond. Conclusions are drawn in section 6. Only austral summer (DJF) and winter (JJA) river flow are investigated in this study because these seasons demonstrate the strongest climatic responses to contrasting SAM phases. Such strong climate contrasts between SAM phases should therefore be reflected in river flow responses.

2. DATA AND METHODS

In this section the data and methodology used to explore SAM river flow associations is described. Three broad types of data are used for the 1979 – 2011 study period including indices of major modes of atmospheric circulation and time series of winter and summer river flow and temperature and precipitation across New Zealand. Time series correlation, linear regression, principal components and composite analyses were the main techniques employed. A fuller description of data and methods is presented in the following sections.

2.1 Indices of atmospheric circulation modes

The two principal modes of atmospheric circulation focused on here are the Southern Oscillation (SO) and the Southern Annular Mode (SAM). From amongst a plethora of SO and SAM indices

1
2
3 (Hanley et al., 2003; Ho et al., 2012) we made use of the Troup (Troup, 1965) Southern
4 Oscillation Index (SOI) and Marshall's SAM index (Marshall, 2003). The Troup SOI (T_SOI
5 hereafter) is the standardized mean sea level pressure (SLP) difference between Tahiti and
6 Darwin. This was downloaded from the Australian Bureau of Meteorology (BoM) web site:
7 <ftp://ftp.bom.gov.au/anon/home/ncc/www/sco/soi/soiplaintext.html>. Marshall's SAM index
8 (M_SAMI hereafter) is a sea level pressure observation based index that uses normalized
9 monthly zonal mean sea level pressure (SLP) differences from a total of 12 stations located at
10 approximately 40°S and 65°S. The M_SAMI was downloaded from the British Antarctic Survey
11 web site <https://legacy.bas.ac.uk/met/gjima/sam.html>.
12
13
14

15 **2.2 River flow, temperature and precipitation data**

16
17 In selecting river flow gauging locations four criteria were applied namely: (1) flow records are
18 continuous over time (less than 5% of missing data), (2) river flow records must be at least 30
19 years in length, (3) catchment areas upstream of the river flow gauging sites have minimal
20 human modification due to flow diversion, damming and irrigation, and (4) the flow regime has
21 not undergone significant changes over time. The first two criteria ensures that there is a
22 balance between spatial and temporal coverage in that the selected gauging stations
23 adequately represent river flow patterns across the country, and possible inter-decadal
24 variability is captured. The third criterion was applied to ensure that the river flow data
25 represented conditions as close to "natural flow" as possible, while the last criterion
26 acknowledges the importance of the stationarity of the river flow depth–volume rating curve
27 such that erroneous discharge estimates do not arise due to major changes in the river channel
28 morphology and thus cross-sectional area. Based on these criteria 40 river flow gauging
29 locations representing 40 rivers were selected (Figure 1). For these stations daily river flow,
30 measured in cubic metres per second, were provided by New Zealand's National Institute of
31 Water and Atmosphere (NIWA). Daily flow data was subsequently converted into monthly and
32 seasonal means and then logarithmically transformed to ensure normality of distribution given
33 this is an underlying assumption in the case of the parametric statistical methods applied in this
34 research. The river flow datasets were truncated to a common period (1979-2011) so as to
35 maximise the number of gauging stations while meeting the minimum requirement of a 30 year
36 record. Details regarding catchment area, winter and summer river flow coefficient of variance
37 (CV) and location for the 40 gauging stations are listed in Table 1. Monthly mean temperature
38 (Tmean), maximum temperature (Tmax) and total rainfall (RainT) for five stations were
39 downloaded from the New Zealand National Climate Database available via NIWA at
40 <http://cliflo.niwa.co.nz/>.
41
42
43
44
45
46
47

48 **2.3 Approach and analytical methods**

49
50 We tackled the analysis of SAM river flow associations following two broad strategies referred to
51 as the environment-to-circulation (E-C) and circulation-to-environment (C-E) approaches
52 (Yarnal, 1993). In the case of the E-C approach years with winters and summers possessing
53 threshold based high and low river flows are first identified, followed by a consideration of the
54 atmospheric circulation characteristics for the set of high and low river flow years. For the C-E
55 approach, groups of years with strongly contrasting atmospheric circulation characteristics are
56
57
58
59
60

1
2
3 first identified. Subsequently the magnitude of winter and summer river flows for these two
4 contrasting groups of years is compared.
5

6
7 In order to reveal genuine hydro-climate linkages on the inter-annual timescale, linear trends in
8 river flow and indices of atmospheric circulation modes were removed prior to analysis. Also in
9 an attempt to understand the unfettered river flow response to the SAM, with respect to El Nino
10 Southern Oscillation (ENSO) as represented by the T-SOI, multiple linear regression (MLR) of
11 concurrent and lead SOI variables were applied to de-trended river flow time series and its
12 proxy in the form PC scores (see below). A subsequent stepwise variable elimination (SMLR)
13 was performed to retrieve a parsimonious model that accounts for the impact of ENSO on river
14 flow. The final modelled river flow residuals are considered as the variability not explained by
15 ENSO and therefore constitute the river flow time series that formed the basis of the analysis
16 presented here; Cox (2006) has discussed extensively the use of residuals in this way to de-
17 emphasise the importance of strong underlying drivers of environmental time series patterns.
18
19

20
21 As opposed to assessing SAM river flow associations on a SAM-individual gauging station basis,
22 winter and summer river flow regions were identified in order to provide a spatial framework for
23 the analysis. We used a S-mode decomposition Principal Component Analysis (PCA) based on
24 a correlation matrix with Varimax rotation (Yarnal 1993) in the first of a two-step process to
25 achieve a river flow regionalisation for New Zealand.
26
27

28
29 There are a number of subjective decisions to be made in applying PCA. Following Yarnal
30 (1993) a correlation dispersion matrix was selected as the basis for identifying river flow gauging
31 stations with covariant river flow. Of a range of rotation possibilities, including non-rotation, a
32 Varimax rotation was performed. A Varimax rotation was selected as it is more conducive to a
33 physical interpretation of the principal components (PC) compared to other rotations (Yarnal,
34 1993). Further, rotation is deemed necessary when PCA is applied in a spatial context
35 (Richman, 1986), as it is here. An outcome of PCA is the identification of a number of principal
36 components ($1 - n$, where n equals the number of original variables or gauging stations), which
37 in decreasing order of importance explain the variability of river flow attributable to groups of
38 gauging stations possessing similar flow variability. The relative importance of the PC in terms
39 of the proportion of the total river flow variability explained is represented by the magnitude of a
40 PCs' eigenvalue. In this analysis the number of PC to retain for further analysis and thus
41 explaining the largest proportion of variability, was selected based on Kaiser's eigenvector/PC
42 selection criterion (Jackson, 1993), that is, PCs with eigenvalues greater than unity were
43 retained. The retained PC were used to generate a time series of PC scores which represent
44 the covariant behaviour of river flow across the gauging stations that comprise each of the river
45 flow regions; PC scores for each region are in effect regional river flow series. PC Scores in
46 addition to values of the M-SAMI formed the basis of the E-C and C-E approaches applied in
47 the split composite analysis.
48
49
50
51
52

53 In the second step the resultant loadings associated with the retained PC were subject to
54 Cluster Analysis (CA) so as to identify groups of stations, or regions that possess river flow
55 affinity for winter and summer. As CA can be approached in a number of ways, both partitioning
56 and hierarchical methods were explored in the development of the final river flow regionalisation.
57
58
59
60

1
2
3 In doing so the following was considered: (1) K-means partitioning based on Euclidean distance,
4 (2) Hierarchical clustering based on Euclidean distance using Average Linkage, Complete
5 linkage or Ward's method. K-means partitioning is an iterative technique that is best used with
6 large datasets. It is sensitive to the starting strategy and when used on a small dataset it has
7 problems converging to a global optimum. Average linkage classifies according to the average
8 distances between members of paired groups, and complete linkage uses a distance measure
9 for the two farthest members in paired groups. The former does not cope well with overlap
10 between clusters while the latter is easily affected by outliers. Ward's method is the hierarchical
11 version of the k-means partitioning method. Although it has the tendency to form clusters of
12 equal size, it has been shown to perform well in simulations and in practice (e.g., Stahl and
13 Demuth, 1999; Kingston et al., 2009). By comparing the results arising from the range of
14 clustering methods described above it was decided that the Ward's method yielded the most
15 physically meaningful river flow regionalisation.
16
17
18
19
20

21 **3. RESULTS**

22 Three broad types of results are presented in this section. Firstly the outcome of the river flow
23 regionalisation based on PCA and CA is described. Following this, the results from an
24 assessment of the significance of the statistical association between times series of river flow,
25 climate variables and the M-SAMI are presented. Lastly results arising from the application of
26 split sample composite analyses in the broader context of the E-C and C-E approaches
27 described above are outlined.
28
29
30

31 **3.1 River flow regionalization**

32 The PCA of summer and winter river flows for the period 1979 – 2011 produced seven PC with
33 eigenvalues great than one, explaining 83.2% and 84% of the total river flow variance across
34 the 40 gauging stations respectively. For summer and winter PC1, or the dominant mode of
35 variability, accounted for 39.1 and 43.4 percent of the total variability respectively. A mapping of
36 PC1 loadings for DJF and JJA (Figure 2) shows for summer that river flow in the Bay of Plenty,
37 East Cape, Hawkes Bay region (Figure 2a) accounts for the greatest variability whereas in the
38 winter (Figure 2b), the most variable region shifts westward, encompassing the broader Waikato,
39 Taranaki and central North Island region. Although space restrictions prevent the presentation
40 of summer and winter loading maps for the other six PCs, Table 2 summarises the percentage
41 variance explained by each of the seven individual PC for both seasons.
42
43
44
45

46 The emergent river flow regionalisation based on CA of the PC loadings, comprising six winter
47 and summer river flow regions, is presented in Figure 3. The river flow regions resemble those
48 produced by Mosely (1981) and Beadle and McKerchar (1982), a convergence of results that
49 lends credence to the PCA/CA based regionalisation produced here. Table 3 presents a listing
50 of stations belonging to each river flow region. Moving from north to south the regions are
51 named in the following way: Northland and Coromandel (NICor), Bay of Plenty and Hawke's
52 Bay (BoPHB), Central Western Lower North Island (CWLNI), Upper South Island (UpSI),
53 Canterbury (CTB), and Lower South Island (LoSI). The summer/winter clusters are almost
54 identical except for a few rivers located exclusively in the broad central North Island regions
55
56
57
58
59
60

1
2
3 namely Tongariro, Waihothonu, Waikato and Waihou. These all become more “BoPHB like” in
4 winter.
5

6
7 The general hydrological characteristics of the emergent regions are encapsulated in plots of
8 the monthly distribution of river flow (Figure 4) for gauging stations near the statistical centroid
9 of each of the clusters. CWLNI and NICor demonstrate a strong seasonal cycle that closely
10 mirrors seasonal precipitation distribution in the North Island (Ummenhofer and England, 2007)
11 with a clear summer period of distinctly lower flows compared to the winter months. The drop in
12 flow in September suggests purely rain-fed river flow without snowmelt contribution in these two
13 regions. Relative to its counterparts the BoPHB region in the North Island possesses a
14 dampened seasonality of river flow, apart from the slight flow elevation July through to
15 September often due to storms which stall off the east coast of the North Island around the East
16 Cape region and advect moisture off the Pacific Ocean to the east. The South Island regions
17 demonstrate clear evidence of the importance that spring snow melt water plays in the seasonal
18 distribution of river flow. This is especially so for UpSI and LoSI where throughout the months
19 July to October there is a steady increase in flow indicating an increasing release of water from
20 melting high elevation snowpacks. Interestingly for UpSI and LoSI, June relative to July
21 demonstrates on average higher flows as in June rainfall dominates the precipitation mix and is
22 available to runoff immediately while in July much of the precipitation that falls is tied up in snow
23 with moisture released for runoff later in the winter or in early spring. Of note for all rivers is the
24 dominance of data points for high flow compared to low flow indicating that New Zealand rivers
25 demonstrate most of their variability due to high as opposed to low flows. This is especially so
26 for the winter months (Figure 4). A further explanation for the predominance of high flow data
27 points is that while river flow is bound at the lower limit it is not so at the upper limit with the
28 predominance of high flow data points indicating the skewed nature of monthly river flow, a
29 feature quite typical of rivers in maritime temperate climates (Moseley and Pearson, 1997;
30 Sturman and Tapper, 2006).
31
32
33
34
35
36

37 **Association between SAM, climate and regional river flow**

38
39 The response of regional river flow, as represented by PC scores, to the SAM and regional
40 climate variables is assessed in this section for both summer and winter.
41
42

43 Table 4 presents the correlation coefficients for monthly values of SAM and climate variables
44 correlated with river flow as represented by PC scores for DJF (Table 4a) and JJA (Table 4b)
45 (n= 96 months). Perhaps unsurprisingly river flow is positively correlated with RainT for all
46 regions in both summer and winter (Table 4). This contrasts with the situation regarding
47 temperature both in terms of the sign of correlation and the universality of the river flow
48 temperature association. For summer, although the sign of the correlation coefficients indicate a
49 general inverse association between temperature and river flow, this association is not
50 significant for all regions. Both BoPHB and CWLNI demonstrate no statistically significant
51 correlations between river flow and temperature. For UpSI, CTB and LoSI however the
52 significant inverse association between river flow and Tmean and Tmax points to the possible
53 role of reduced effective precipitation because of high evaporation rates and thus low (high)
54 river flows being linked with warm (cool) summers. This seems to be especially so for CTB, a
55
56
57
58
59
60

1
2
3 particularly drought prone region, which shows a very strong inverse association between river
4 flow and Tmax. For winter there is no consistency in the direction of association between river
5 flow and temperature. While CWLNI river flow exhibits no apparent sensitivity to temperature,
6 CTB and LoSI river flow demonstrate a significant inverse association with temperature, which
7 is distinctly different to the positive relation exhibited for UpSI and BoPHB. This seemingly
8 contradictory situation may be explained by possible differences in the way in which water is
9 stored within a catchment and eventually delivered to the river channel. For CTB and LoSI a
10 large proportion of winter precipitation is delivered as snow, the storage of which will be
11 favoured in cool winters with suppressed Tmax. In this case a deep snowpack and therefore high
12 water equivalents will be available for runoff during intermittent melt periods during the winter
13 with this reflected in elevated winter discharge levels. In the case of UpSI, which also receives
14 considerable winter snowfall but straddles a transitional zone between rain and snow along a
15 warming south to north temperature gradient, higher winter temperatures result in a great
16 proportion of the precipitation being delivered as rain which is directly available for runoff. For
17 BoPBH, in which the majority of catchments are rainfall dominated, warmer winters are
18 associated with higher river flows via increased rainfall from warm air masses emanating from
19 the Pacific Ocean to the east often allied with extra-tropical weather systems.

20
21
22 The link between the SAM and regional river flow is weak in summer. While only BoPHB shows
23 a statistically significant association, the inverse nature of which indicates summer high flows
24 are associated with a negative phase of SAM, the sign of the correlation suggests similar
25 associations for UpSI and LoSI (Table 4) Compared to summer, winter river flows appear to be
26 more responsive to variations in the SAM. The statistically significant inverse association
27 between CWLNI and LoSI river flows and the SAM indicates that vigorous westerly flows
28 associated with eastward moving cyclonic systems, as occur under a negative phase of the
29 SAM, are conducive to high river flows. In the case of CTB a positive phase of the SAM
30 increases the number of rain bearing weather systems arriving over CTB from the Pacific Ocean
31 to the east hence explaining the positive SAM river flow associations for this region, similar to
32 the SAM BoPHB association for summer.

33
34
35 Whereas correlations presented in Table 4 are for monthly values, Table 5 presents a similar
36 analysis but at the seasonal level ($n = 32$) for DJF and JJA for concurrent and one season in
37 advance associations between SAM and river flow as represented by PC scores. Spring (SON)
38 SAM is found to be strongly correlated with summer (DJF) CTB river flow ($r = 0.579$) (Table 5).
39 This may be explained with reference to regional climate as for the CTB region, a positive SAM
40 in SON leads to lower SON temperatures ($r = -0.297$, $p = 0.05$) which in turn suppresses
41 springtime snowmelt providing a relatively large reservoir of snow for summer melt and hence
42 elevated flow. The negative relationship between spring SAM and summer flow found for
43 CWLNI ($r = -0.437$), and to some extent NICor ($r = -0.210$) is most likely related to elevated soil
44 moisture in spring due to positive rainfall anomalies associated with a negative SAM and thus
45 elevated baseflow in summer. Although UpSI is the only region that displays statistically
46 significant concurrent associations between the SAM and summer river flow near significant
47 associations are evident for BoPHB and CTB, reflecting the situation found for monthly river flow
48 (Table 4). In the case of UpSI and BoPHB (CTB) the SAM river flow association is an inverse
49 (positive) one such that a positive (negative) SAM phase and thus below (above) average
50
51
52
53
54
55
56
57
58
59
60

1
2
3 summer rainfall and possible enhanced (suppressed) evaporation rates leads to reduced
4 (increased) river flow.
5

6
7 At the seasonal level winter river flow is found to be influenced by the state of the SAM in the
8 preceding and current season for both LoSI and CWLNI with both displaying a significant
9 inverse association (Table 5). The lagged and concomitant inverse SAM river flow associations
10 found for these two regions corroborate that found at the monthly time scale such that strong
11 westerlies and attendant positive precipitation anomalies (SAM negative phase) are conducive
12 to high winter flows. The positive concurrent SAM river flow association for winter BoPHB river
13 flow is opposite to the inverse association found for summer (Table 5) suggesting a strong
14 seasonal influence of the SAM on river flow in this region as hinted at by the monthly analyses
15 presented in Table 4.
16
17

18
19 From the preliminary results presented here, it is tentatively proposed that CWLNI, UpSI are
20 sensitive to changes in atmospheric circulation in both seasons while NICor, BoPHB, CTB and
21 LoSI show a seasonally varying response.
22

23 **3.2 Split sample and composite analyses**

24
25 River flow SAM associations were also investigated using the so called circulation-to-
26 environment(C-E) and environment-to-circulation (E-C) approaches (Yarnal, 1993). In the
27 former case, river flow response is assessed relative to the state of the atmospheric circulation,
28 while for the latter, atmospheric circulation fits criteria based on river flow. These two
29 approaches are realized by splitting and compositing the variables (circulation state or river flow)
30 of interest.
31
32

33 **3.3.1 Circulation-to-environment approach**

34
35 The aim of the C-E approach is to establish the nature of regional flow response associated with
36 firstly positive and negative phases of SAM for summer and winter and secondly pronounced
37 phases of SAM for the aforementioned seasons. The first aim was achieved by plotting the
38 distribution of PC scores (proxies of river flow) for broad-spectrum positive and negative phases
39 of the SAM as defined by SAM index values greater or less than zero respectively. The second
40 aim required sub-setting seasonal SAM indices into pronounced positive and negative phases
41 of SAM as defined by upper and lower quartile values of the seasonal SAM index, and then
42 evaluating the resultant distribution of PC scores. A standard t-test was employed to test for the
43 significance of any differences in river flow based on the broad-spectrum and pronounced
44 phases of SAM. The outcomes of the C-E and E-C approaches are both presented in the form
45 of boxplots showing the distribution of PC scores or SAM index values.
46
47
48
49

50
51 Figure 5 shows the distribution of PC scores for the broadly defined positive and negative
52 phases of SAM. For summer (Figure 5a) the only region that demonstrates a statistically
53 significant difference in flow between positive and negative phases of SAM is CTB such that
54 there is a tendency for river flow to be higher (lower) under a positive (negative) phase of SAM.
55 Although there appears to be visible SAM positive and negative phase related differences in the
56 distribution of river flow values for BoPHB, UpSI and CWLNI, large flow variances for the two
57
58
59
60

1
2
3 SAM states militate against any statistically significant differences. In contrast to summer, SAM
4 phase related differences in flow are stronger for winter (Figure 5b). T-tests reveal a strong
5 negative association, significant at the 5 percent level, for CWLNI and LoSI and a weak positive
6 association, significant at the 10 per cent level, for CTB and BoPHB. For pronounced phases of
7 the SAM, the association identified for the broad-spectrum positive and negative phases are
8 replicated with no additional regions demonstrating river flow contrasts for pronounced SAM
9 phase differences.
10
11

12 13 **3.3.2 Environment-to-Circulation Approach**

14
15 The aim of the E-C approach was to establish whether distinct phases of SAM are associated
16 with summers and winters possessing high or low river flows. A summer or winter high (low)
17 river flow state is defined as possessing PC score values greater (less) than the 75th (25th)
18 percentile value. An analysis of the difference in SAM index values between summer high and
19 low flow years revealed no statistically significant differences although the distribution of SAM
20 index values, as revealed by boxplots (Figure 6a), are suggestive of summer SAM river flow
21 associations for BoPHB, CWLNI and UpSI. In contrast to summer the situation is far less
22 equivocal for winter (Figure 6b). Statistically different distributions of SAM index values between
23 high and low river flow states are evident for CWLNI and LoSI such that low SAM index values
24 or a pronounced negative phase of SAM is associated with anomalously high river flows.
25 Although not statistically significant, CTB displays a similar tendency to that of CWLNI and LoSI
26 with the opposite evident for BoPHB.
27
28
29
30

31 **4. DISCUSSION**

32
33 A burgeoning number of studies on the association between large scale climate mechanisms
34 and hydrological response point to the importance of a range of modes of atmospheric
35 circulation in determining river flow variability across a number of timescales (Clarke et al, 2014;
36 Coleman et al., 2013; Fendekova et al., 2014; Hannaford et al., 2013). Given this and the fact
37 that precipitation and temperature variability across the southern hemisphere is responsive to
38 SAM (Cai et al., 2011; Diamond and Renwick, 2015; Feng et al., 2015; Hendon et al., 2014;
39 Holz et al., 2011; Lim and Hendon, 2015; Manatsa et al., 2015; Oliveira et al., 2014; Raut et al.,
40 2014; Zheng et al., 2014), the linkages between SAM and river flow across New Zealand
41 observed in this study appear to support the general claim that large scale climate forcings
42 serve as remote drivers of hydrological variability. Notwithstanding this generality, the situation
43 uncovered for New Zealand indicates clear geographical and seasonal variation in the nature of
44 SAM river flow associations.
45
46
47

48
49 Of the two seasons considered, SAM river flow associations are most pronounced in winter.
50 This seasonal asymmetry of the climatic impact of SAM has been noted by others in relation to
51 New Zealand winter precipitation and temperature patterns (Fogt et al., 2012) but as yet has not
52 been explored for river flow. The asymmetric seasonal response of winter flow to the SAM can
53 possibly be understood via seasonal SAM climate associations described by Kidston et al.
54 (2009) such that contrasting phases of the SAM have their greatest impacts on precipitation in
55 the winter via pronounced anomalies in the westerlies and advection of moisture from the west.
56 In addition to this, Fogt et al. (2012) hypothesise that the apparently stronger SAM climate
57
58
59
60

1
2
3 teleconnections in winter are possibly related to the more variable background state in winter
4 while recognising that the extent to which the seasonal zonal asymmetric structure in the SAM
5 impacts on teleconnections is equivocal. That the seasonal asymmetry in the hydrological
6 response to the SAM is not unique to New Zealand is supported by findings from elsewhere
7 (Cai et al., 2011; Liess et al., 2014; Wu et al., 2015) but, as implied by the work of Fogt et al.
8 (2012), a compelling explanation for this remains elusive.
9
10

11 Geographical contrasts of the impact of SAM on river flow can be explained through terrain
12 atmospheric circulation interactions and the marked impact that orography has on wind speed
13 and direction and thus climate anomalies across New Zealand (Kidson, 2000) and the resultant
14 precipitation anomalies that arise from distinct phases of the SAM (Kidston et al., 2009).
15 Focusing on winter as the season of greatest sensitivity of river flow to SAM, CWLNI and LoSI
16 display an inverse association with the SAM while BoPHB possesses a positive association.
17 Therefore in terms of SAM phases, when SAM is negative (positive) the CWLNI and the LoSI
18 receive anomalously strong flows of air from the west to northwest (north to northeast) exposing
19 these regions to moisture advection (rain shadow effect and subsidence) resulting in elevated
20 (reduced) precipitation levels and thus high (low) river flow. Typically moisture advection from
21 the west is associated with rapid eastwardly moving low pressure systems which track just to
22 the south of NZ (Li et al., 2016). For the BoPHB region, lying on the eastern side of the North
23 Island, the opposite is true such that a positive phase of the SAM results in anomalous easterly
24 flows such that the rain shadow effect associated with a negative phase is suppressed allowing
25 penetration of moisture off the Pacific Ocean to the east which is conducive to high river flows.
26 This resonates with the findings of Griffiths (2011) who found that the BoPHB region generally
27 experiences wet conditions during a positive phase of the SAM. Study findings have also
28 revealed that the summer and winter river flow response for some regions depends on the
29 phase of SAM in the preceding season, namely spring and autumn respectively. The one
30 season ahead associations may well relate to antecedent hydroclimatological factors related to
31 the SAM phase that prime hydrological regions for either anomalously high or low river flow.
32 This suggestion resonates with the findings of Liu et al. (2015; 2016) who have suggested that
33 boreal autumn ocean-atmosphere conditions influenced by SAM persist into the winter with a
34 knock-on effect regarding the nature of winter precipitation over land in the Northern
35 Hemisphere.
36
37
38
39
40
41
42

43 The regional contrasts in the response of winter river flow described above for CWLNI, LoSI and
44 BoPHB point to possible hydrological teleconnections between CWLNI/LoSI and BoPHB. This
45 possibility is borne out by examination of the covariant behaviour of river flow time series for
46 these regions as represented by PC scores (Figure 7). At times there is clear concomitant
47 inverse behaviour in regional river flow. For example, if BoPHB is compared with the LoSI
48 (Figure 7a) for summer, the period post 1995 is noteworthy for a number of years that show
49 reversals in river flow response with high flow in one region matched by low flow in the other.
50 This is also evident in winter for BoPHB and CWLNI which are geographically close and lie on
51 opposite sides of the central axis of the North Island (Figure 7b). Such possible hydrological
52 teleconnections or “seesaws” can be understood by examining, via an environment-to-
53 circulation perspective, the large scale atmospheric circulation situation associated with
54 anomalously high/low winter river flows for the CWLNI region as portrayed in Figures 8 and 9. In
55
56
57
58
59
60

1
2
3 Figure 8a (Figure 8b), the pressure anomaly pattern is shown for a composite of years for the
4 upper (lower) 25 per cent of CWLNI winter river flows. Immediately apparent is the typical winter
5 asymmetric structure of the pressure field associated with a negative SAM phase, as described
6 by Kidston et al (2009). Notwithstanding this, of interest is the relation of the CWLNI and BoPHB
7 regions relative to the extensive region of anomalously low pressure that stretches across NZ,
8 associated with which is an area of anomalously strong westerly winds (Figure 9a). Such an
9 anomalous circulation configuration will expose the west facing CWLNI region to advection of
10 moisture from the northwest to west while the BoPHB lies in a rain shadow area to the east
11 sheltered from rain bearing westerly flows. The pressure anomaly pattern in Figure 8b is that
12 associated with CWLNI winter low flows. It resembles to a very high degree the pressure
13 distribution accompanying a strong positive phase of the SAM with attendant weakened
14 (strengthened) westerly (easterly) flows across the North Island of NZ (Figure 9b). Under such a
15 pressure configuration and allied anomalous flow pattern, the situation as described for high
16 flows in the CWLNI region is reversed such that the BoPHB region is exposed to moisture
17 advection from the east, emanating from the region of anomalously high pressure over the
18 central South Pacific, while the CWLNI lies in a rain shadow to the west. Although statistically
19 significant concurrent correlations were not found between SAM index values and river flow for
20 the CTB region, the sign of the monthly and one season ahead correlations are strongly
21 suggestive of CTB high flows being consequential on a positive SAM phase (Tables 4 and 5).
22 Given its similar geographic setting to that of BoPHB, and the SAM river flow association
23 described above for that region as well as CWLNI and LoSI, it would appear that orographic
24 forcing/blocking or steering play a fundamental role in controlling the hydrological outcome of
25 strong positive and negative phases of SAM across New Zealand via its control on precipitation
26 delivery. This claim lends credence to the hydrological implications of climatological research
27 conducted on zonal precipitation gradients across NZ (Chater and Sturman, 1998; Kerr et al.,
28 2015; Webster et al., 2015). Further the posited significance of distinct synoptic scale
29 configurations of atmospheric circulation for river flow across New Zealand is supported by work
30 conducted for similar geographical settings that evaluate the importance of orography in
31 determining hydrological outcomes arising from contrasting phases of large scale modes of
32 climatic variability (Luce et al., 2013; Mass et al., 2015; Siler et al., 2013; Viale and Nunez,
33 2011).

34
35
36
37
38
39
40
41
42
43 Lastly as implied by the above discussion, this study's findings on the impact of the SAM on
44 river flow complement the SAM-rainfall relationships described by Kidston et al (2009) but
45 contrast with those outlined by Ummenhofer and England (2007). Two reasons account for the
46 discrepancy of our findings with those Ummenhofer and England (2007). Firstly we, as did
47 Kidston et al (2009), analysed potential relationships at the seasonal timescale in contrast to
48 Ummenhofer and England (2007) who evaluated the SAM-rainfall relationship at the annual
49 timescale. Secondly, we consider a symmetric response of river flow to the SAM while
50 Ummenhofer and England (2007) examined the two discrete phases independently.

54 5. CONCLUSIONS

55
56 This study has assessed the nature of the possible association between the SAM and river flow
57 across New Zealand at the inter-annual timescale by implementing an approach for controlling
58

1
2
3 for the likely influence of ENSO on river flow and the observed tendency for the SAM to adopt
4 an increasingly positive phase over the length of the study period. Other notable aspects of the
5 approach taken in exploring hydroclimatological linkages include the comparison of analysis
6 outcomes from the so called circulation-to-environment and environment-to-circulation
7 approaches to understanding climate-environment relationships.
8
9

10 Study findings confirm suspected linkages between the SAM and river flow based on known
11 SAM influences on temperature and precipitation patterns across New Zealand and in doing so
12 add to the growing international literature on the determinants of hydroclimatological variability. A
13 component of the analysis strategy that has assisted with revealing the nature of regional scale
14 impacts of the SAM on river flow is the development of a summer and winter river flow
15 regionalisation. This has provided an effective spatial framework for the analyses presented
16 here as it has facilitated the identification of groups of gauging stations that have a high degree
17 of affinity in relation to natural flow regimes and the possibility of SAM related hydrological
18 teleconnections across New Zealand.
19
20
21

22 Because of the complex orography of New Zealand and its general north-south orientation, in a
23 part of the world dominated by temporally and spatially variable westerlies, it is perhaps no
24 surprise that there is little season to season and geographical stationarity in the SAM river flow
25 relationship. Study results have revealed winter as the season with the clearest hydrological
26 response to SAM. These findings serve as a warning against applying knowledge based on
27 analyses of river flow response to large scale modes of climatic variability, conducted at coarse
28 geographical and temporal scales, to climate-informed long range hydrological forecasts.
29
30
31

32 Although it has been found that BoPHB and LoSI flows are correlated with SAM, as implied in
33 the limited SAM-climate literature for New Zealand, a hypothetical assumption of an out-of-
34 phase relationship between these two regions turns out to be too simplistic. Indeed the PC
35 score time series suggests periodical coupling and decoupling, indicating complex driving
36 mechanisms. Further the associations between SAM and river flow are seasonally different
37 (notably in the BoPHB and CWLNI regions), meaning that relationships derived from one
38 season, may not be extrapolated to other seasons. There is also clear evidence that large-scale
39 sea level pressure and zonal wind patterns associated with SAM phases modulate regional river
40 flow and that the SAM phase in the spring and autumn may well exert influence on summer and
41 winter river flow respectively. These findings provide a persuasive argument for the inclusion of
42 information related to the SAM in seamless river flow forecasting models for New Zealand as
43 autumn and spring SAM conditions have the potential to provide a significant forecasting signal
44 for NZ climate and thus river flow in the following winter and summer respectively.
45
46
47
48

49 This study has largely focused on the direct influence of SAM on river flow. However it is
50 acknowledged that precipitation, temperature and land surface conditions (e.g., vegetation and
51 soil moisture status) will moderate the influence of SAM on river flow as will topography. The
52 effects of these moderating or “hydrological memory” factors, require investigation in future
53 research. In a similar vein and acknowledging the likely conjoint influence of ENSO and SAM on
54 river flow, future research on the impact of synchronous and asynchronous ENSO/SAM phases
55 on river flow is warranted.
56
57
58
59
60

1
2
3 Although much remains to be uncovered about the hydroclimatology of New Zealand, this study
4 has facilitated the placement of the hydrology of a number of New Zealand regions in a wider
5 hydroclimatological context by shedding light on how the SAM, as one of a number of large
6 scale modes of climate variability, may influence winter and summer hydrological variability at
7 the inter-annual timescale. This type of knowledge opens up the potential for seasonal river flow
8 prediction based on the conditioning of catchment to regional scale hydrological modelling on
9 the large scale atmospheric circulation state.
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review

REFERENCES

- Allen KJ, Nichols SC, Evans R, Cook ER, Allie S, Carson G, Ling F, Baker PJ. 2015. Preliminary December-January inflow and streamflow reconstructions from tree rings for western Tasmania, southeastern Australia. *Water Resources Research*, 51: 5487-5503. doi: 10.1002/2015WR017062
- Bardsley E. 2016. Preliminary evaluation of New Zealand seasonal river flow forecasts. *Journal of Hydrology (NZ)*,55: 65-68.
- Beable MR, McKerchar AI. 1982. Regional Flood Estimation in New Zealand. Water & Soil Technical Publication 20. Wellington, National Water and Soil Conservation Organisation.
- Beckers JVL, Weerts AH, Tjiedeman E, Welles E. 2016. ENSO-conditioned weather resampling method for seasonal ensemble streamflow prediction. *Hydrology and Earth System Sciences*, 20: 3277-3287. DOI: 10.5194/hess-20-3277-2016
- Brown A, Milton S, Cullen M, Golding B, Mitchell J, Shelly A. 2012, Unified modelling and prediction of weather and climate: A 25-year journey, *Bulletin American Meteorological Society*, 93: 1865–1877.
- Cai W, van Rensch P, Borlace S, Cowan T. 2011. Does the Southern Annular Mode contribute to the persistence of the multi-decade-long drought over southwest Western Australia? *Geophysical Research Letters* 38: Article Number: L14712. DOI: 10.1029/2011GL047943
- Chang C-H, Johnson NC. 2015. The continuum of wintertime southern hemisphere atmospheric teleconnection patterns. *Journal of Climate* 28: 9507-9529. DOI: 10.1175/JCLI-D-14-00739.1
- Chater AM, Sturman AP. 1998. Atmospheric conditions influencing the spillover of rainfall to lee of the southern Alps, New Zealand *International Journal of Climatology*, 18: 77-92.
- Chiew FHS, McMahon TA. 2002. Global ENSO-streamflow teleconnection, streamflow forecasting and inter-annual variability. *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques* 47: 505-522. DOI: 10.1080/02626660209492950
- Clark, C III, C; Nnaji GA, Huang W. 2014. Effects of El-Nino and La-Nina sea surface temperature anomalies on annual precipitations and streamflow discharges in southeastern United States. *Journal of Coastal Research* 68: 113-120. DOI: 10.2112/SI68-015.1
- Coleman JSM, Budikova D. 2013. Eastern U.S. summer streamflow during extreme phases of the North Atlantic Oscillation. *Journal of Geophysical Research-Atmospheres* 118: 4181-4193. DOI: 10.1002/jgrd.50326
- Coleman JSM, Rogers JC. 2003. Ohio River Valley winter moisture conditions associated with the Pacific-North American teleconnection pattern. *Journal of Climate* 16: 969-98. DOI: 10.1175/1520-0442(2003)016<0969:ORVWMC>2.0.CO;2 1.

- 1
2
3 Cox NJ. 2006. Assessing agreement of measurements and predictions in geomorphology.
4 Geomorphology, 76:332-346.
5
6
7 Dettinger MD, Diaz H. 2000. Global characteristics of stream flow seasonality and variability.
8 Journal of Hydrometeorology 1: 289-310. doi: 10.1175/1525-
9 7541(2000)001<0289:GCOSFS>2.0.CO;2
10
11 Diamond HJ, Renwick JA. 2015. The climatological relationship between tropical cyclones in the
12 southwest Pacific and the southern annular mode. International Journal of Climatology 35: 613-
13 623. DOI: 10.1002/joc.4007
14
15
16 Ding H, Greatbatch RJ, Gollan G. 2015. Tropical impact on the interannual variability and long-
17 term trend of the Southern Annular Mode during austral summer from 1960/1961 to 2001/2002.
18 Climate Dynamics 44: 2215-2228. DOI: 10.1007/s00382-014-2299-x
19
20
21 Dravitzki S, McGregor J. 2011. Predictability of heavy precipitation in the Waikato River Basin of
22 New Zealand. Monthly Weather Review, 139: 2184-2197. DOI: 10.1175/2010MWR3137.1
23
24 Fan K. 2007. Zonal asymmetry of the Antarctic Oscillation. Geophysical Research Letters 34
25 Article Number: L02706. DOI: 10.1029/2006GL028045
26
27
28 Fendekova M, Pekarova P, Fendek M, Pekar J, Skoda P. 2014. Global drivers effect in multi-
29 annual variability of runoff. Journal of Hydrology and Hydromechanics 62: 169-176. DOI:
30 10.2478/johh-2014-0027
31
32 Feng J, Li JP, Yu L, Zhu JL, Xie F. 2015. Relationships among the monsoon-like southwest
33 Australian circulation, the Southern Annular Mode, and winter rainfall over southwest Western
34 Australia. Advances in Atmospheric Sciences 32: 1063-1076. DOI: 10.1007/s00376-014-4142-z
35
36
37 Fogt RL, Jones JM, Renwick J. 2012. Seasonal zonal asymmetries in the Southern Annular
38 Mode and their impact on regional temperature anomalies. Journal of Climate 25: 6253-6270.
39 DOI: 10.1175/JCLI-D-11-00474.1
40
41 Griffiths G. 2011. Drivers of extreme daily rainfalls in New Zealand. Weather and Climate 31:
42 24-49.
43
44
45 Hall A, Visbeck M. 2002. Synchronous variability in the southern hemisphere atmosphere, sea
46 ice, and ocean resulting from the annular mode. Journal of Climate 15: 3043-3057.
47
48 Hanley DE, Bourassa MA, O'Brien JJ. 2003. A quantitative evaluation of ENSO indices. Journal
49 of Climate 16: 1249-1258. doi: 10.1175/1520-0442(2003)16<1249:AQEOEI>2.0.CO;2
50
51 Hannaford J, Buys G, Stahl K, Tallaksen LM. 2013. The influence of decadal-scale variability on
52 trends in long European streamflow records. Hydrology and Earth System Sciences 17: 2717-
53 2733. doi: 10.5194/hess-17-2717-2013
54
55
56
57
58
59
60

- 1
2
3 Hendon HH, Lim EP, Nguyen H. 2014. Seasonal variations of subtropical precipitation
4 associated with the Southern Annular Mode. *Journal of Climate* 27: 3446-3460. DOI:
5 10.1175/JCLI-D-13-00550.1
6
7
8 Ho M, Kiem AS. 2012. The Southern Annular Mode: a comparison of indices. *Hydrology and*
9 *Earth System Sciences* 16: 967-982.
10
11 Holz A, Veblen TT. 2011. Variability in the Southern Annular Mode determines wildfire activity in
12 Patagonia. *Geophysical Research Letters* 38 Article Number: L14710 DOI:
13 10.1029/2011GL047674
14
15
16 Hoskins B. 2013. The potential for skill across the range of the seamless weather-climate
17 prediction problem: A stimulus for our science. *Quarterly Journal of the Royal Meteorological*
18 *Society*, 139: 573–584.
19
20
21 Jackson DA. 1993. Stopping rules in Principal Components Analysis: A Comparison of
22 heuristical and statistical approaches. *Ecology* 74: 2204-2214. DOI: 10.2307/1939574
23
24 Jang J, Hong SY. 2014. Quantitative forecast experiment of a heavy rainfall event over Korea in
25 a global model: horizontal resolution versus lead time issues. *Meteorology and Atmospheric*
26 *Physics*, 124: 113-127. DOI: 10.1007/s00703-014-0312-x
27
28
29 Jiang NB, Griffiths G, Lorrey A. 2013. Influence of large-scale climate modes on daily synoptic
30 weather types over New Zealand. *International Journal of Climatology* 33: 499-519. DOI:
31 10.1002/joc.3443
32
33
34 Jones JM, Fogt RL, Widmann M, Marshall GJ, Jones PD, Visbeck M. 2009. Historical SAM
35 variability. Part I: Century-length seasonal reconstructions. *Journal of Climate*, 22: 5319-5345.
36 doi:10.1175/2009JCLI2785.1
37
38 Kerr T, Srinivasan MS, Rutherford J. 2015. Stable water isotopes across a transect of the
39 Southern Alps, New Zealand. *Journal of Hydrometeorology*, 16: 702-715.
40
41
42 Kidson JW. 2000.. An analysis of New Zealand synoptic types and their use in defining weather
43 regimes. *International Journal of Climatology* 20: 299-316.
44
45
46 Kidston J, Renwick J, McGregor J. 2009.. Hemispheric-Scale Seasonality of the Southern
47 Annular Mode and Impacts on the Climate of New Zealand. *Journal of Climate* 22: 4759-4770.
48
49
50 Kiely G. 1999.. Climate change in Ireland from precipitation and streamflow observations.
51 *Advances in Water Resources* 23: 141-151.
52
53
54 Kiem AS, Verdon-Kidd DC. 2009.. Climatic drivers of Victorian streamflow: is ENSO the
55 dominant influence? *Australian Journal of Water Resources* 13: 17-29.
56
57
58 King AD, Klingaman NP, Alexander LV, Donat MG, Jourdain NC, Maher P. 2014. Extreme
59 rainfall variability in Australia: patterns, drivers, and predictability. *Journal of Climate* 27: 6035-
60 6050. DOI: 10.1175/JCLI-D-13-00715.1

1
2
3 Kingston DG, Hannah DM, Lawler DM, McGregor GR. 2009. Climate–river flow relationships
4 across montane and lowland environments in northern Europe. *Hydrological Processes* 23: 985-
5 996. DOI: 10.1002/hyp.7202
6

7
8 Li N, Bellon G, McGregor, GR. (2016). Tropospheric moisture in the Southwest Pacific as
9 revealed by homogenized radiosonde data: Climatology and decadal trend. *International*
10 *Journal of Climatology* (in press)
11

12 Liess S, Kumar A, Snyder PK, Kawale J, Steinhäuser K Semazzi FHM, Ganguly AR,
13 Samatova NF, Kumar V. 2014. Different modes of variability over the Tasman Sea: Implications
14 for regional climate. *Journal of Climate* 27: 8466-8486. DOI: 10.1175/JCLI-D-13-00713.1
15
16

17 Lim EP, Hendon HH. 2015. Understanding and predicting the strong Southern Annular Mode
18 and its impact on the record wet east Australian spring 2010. *Climate Dynamics* 44: 2807-2824.
19 DOI: 10.1007/s00382-014-2400-5
20

21 Lim EP, Hendon HH, Arblaster JM, Delage F, Nguyen H, Min SK, Wheeler MC. 2016. The
22 impact of the Southern Annular Mode on future changes in Southern Hemisphere rainfall.
23 *Geophysical Research Letters*, 43: 7160-7167. doi: 10.1002/2016GL069453
24
25

26 Liu T, Li, J, Zheng F. 2015 Influence of the Boreal Autumn Southern Annular Mode on Winter
27 Precipitation over Land in the Northern Hemisphere. *Journal of Climate*, 28: 8825-8839.
28

29 Liu T, Li JP, Feng J, Wang XF, Li Y, 2016. Cross-Seasonal Relationship between the Boreal
30 Autumn SAM and Winter Precipitation in the Northern Hemisphere in CMIP5. *Journal of Climate*,
31 29: 6617-6636. doi: 10.1175/JCLI-D-15-0708.1
32
33

34 Luce CH, Abatzoglou JT, Holden ZA. 2013. The missing mountain water: slower westerlies
35 decrease orographic enhancement in the Pacific Northwest USA. *Science* 342: 1360-1364. DOI:
36 10.1126/science.1242335
37

38 Manatsa D, Matarira C, Mushore TD, Mudavanhu C. 2015. Southern Africa winter temperature
39 shifts and their link to the Southern Annular Mode. *Climate Dynamics* 45: 2337-2350. DOI:
40 10.1007/s00382-015-2474-8
41
42

43 Mariani M, Fletcher M-S. 2016. The Southern Annular Mode determines interannual and
44 centennial-scale fire activity in temperate southwest Tasmania, Australia. *Geophysical*
45 *Research Letters* 43: 1702-1709. DOI: 10.1002/2016GL068082
46
47

48 Marshall GJ. 2003. Trends in the southern annular mode from observations and reanalyses.
49 *Journal of Climate* 16: 4134-4143. DOI: 10.1175/1520-0442(2003)016<4134:TITSAM>2.0.CO;2
50

51 Marshall AG, Hudson D, Wheeler MC, Hendon HH. 2012. Simulation and prediction of the
52 Southern Annular Mode and its influence on Australian intra-seasonal climate in POAMA.
53 *Climate Dynamics*, 38: 2483. doi:10.1007/s00382-011-1140-z
54
55

56 Marti AI, Yerdelen C, Kahya E. 2010. ENSO modulations on streamflow characteristics. *Earth*
57 *Sciences Research Journal*, 14: 31-43.
58
59
60

1
2
3 Mass C, Johnson N, Warner M, Vargas R. 2015 Synoptic control of cross-barrier precipitation
4 ratios for the Cascade Mountains. *Journal of Hydrometeorology* 16: 1014-1028. DOI:
5 10.1175/JHM-D-14-0149.1
6

7
8 McKerchar AI, Henderson RD. 2003. Shifts in flood and low-flow regimes in New Zealand due to
9 interdecadal climate variations. *Hydrological Sciences Journal-Journal Des Sciences*
10 *Hydrologiques* 48: 637-654.
11

12
13 McKerchar AI, Pearson CP, Fitzharris BB. 1998. Dependency of summer lake inflows and
14 precipitation on spring SOI. *Journal of Hydrology* 205: 66-80. DOI: 10.1016/S0022-
15 1694(97)00144-3
16

17
18 McKerchar AI, Pearson CP. 1994. Forecasts of seasonal river flows using Southern Oscillation
19 Index. *Journal of Hydrology (NZ)* 32: 16-29.
20

21
22 McKerchar AI, Renwick JA, Schmidt J. 2010. Diminishing streamflows on the east coast of the
23 South Island New Zealand and linkage to climate variability and change. *Journal of Hydrology*
24 *(NZ)* 49: 1-14.
25

26
27 Mosley MP 1981. Delimitation of New Zealand hydrologic regions. *Journal of Hydrology* 49:
28 173-192. DOI: 10.1016/0022-1694(81)90211-0
29

30
31 Mosley MP. 2000. Regional differences in the effects of El Nino and La Nina on low flows and
32 floods. *Hydrological Sciences Journal-Journal Des Sciences Hydrologiques* 45: 249-267. DOI:
33 10.1080/02626660009492323
34

35
36 Mosley MP, Pearson CP. 1997. *Floods and Droughts: The New Zealand Experience*. Wellington,
37 New Zealand Hydrological Society, 206pp
38

39
40 Min SK, Cai WJ, Whetton, P. 2013. Influence of climate variability on seasonal extremes over
41 Australia. *Journal of Geophysical Research-Atmospheres*, 118: 643-654. doi:
42 10.1002/jgrd.50164
43

44
45 Oliveira FNM, Carvalho LMV. 2014. A new climatology for Southern Hemisphere blockings in
46 the winter and the combined effect of ENSO and SAM phases. *International Journal of*
47 *Climatology* 34:1676–1692. DOI:10.1002/joc.3795
48

49
50 Pappenberger F, Wetterhall F, Dutra E, Di Giuseppe F, Bogner K, Alfieri L, Cloke HL. 2013.
51 Seamless forecasting of extreme events on a global scale. In: E. Boegh et al., (eds.) *Climate*
52 *and Land Surface Changes in Hydrology*, 3–10, IAHS Publication, Gothenburg, Sweden.
53

54
55 Pearson C.P., 2008. Short- and medium-term climate information for water management.
56 *World Meteorological Organisation Bulletin*, 57, 173-177.
57

58
59 Purdie JW, Bardsley WE. 2010. Seasonal prediction of lake inflows and rainfall in a hydro-
60 electricity catchment, Waitaki river, New Zealand. *International Journal of Climatology*, 30: 372–
389.

1
2
3 Purdie H, Mackintosh A, Lawson W, Anderson B, Morgenstern U, Chinn T, Mayewski P. 2011.
4 Inter-annual variability in net accumulation on Tasman Glacier and its relationship with climate.
5 *Global and Planetary Change*, 77: 142-152. doi: 10.1016/j.gloplacha.2011.04.004
6

7
8 Renwick JA. 2011. Kidson's synoptic weather types and surface climate variability over New
9 Zealand. *Weather and Climate* 31: 3-21.
10

11 Rimbu N, Czymzik M, Ionita M, Lohmann G, Brauer A. 2016. Atmospheric circulation patterns
12 associated with the variability of River Ammer floods: evidence from observed and proxy data.
13 *Climate of the Past*, 12: 377-385. DOI: 10.5194/cp-12-377-2016
14

15 Sallée JB, Speer KG, Rintoul SG. 2010. Zonally asymmetric response of the Southern Ocean
16 mixed-layer depth to the Southern Annular Mode. *Nature Geoscience* 3: 273 – 279.
17 DOI:10.1038/ngeo812.
18

19
20 Shukla S, Sheffield J, Wood EF, Lettenmaier DP. 2013. On the sources of global land surface
21 hydrologic predictability. *Hydrology and Earth System Science*, 17:2781–2796.
22

23
24 Siler N, Roe G, Durran D. 2013. On the dynamical causes of variability in the rain-shadow effect:
25 A case study of the Washington Cascades. *Journal Of Hydrometeorology* 14: 122-139. DOI:
26 10.1175/JBM-D-12-045.1.
27

28
29 Sinclair MR, Renwick JA, Kidson JW. 1997. Low-frequency variability of Southern Hemisphere
30 sea level pressure and weather system activity. *Monthly Weather Review* 125: 2531-2543. DOI:
31 10.1175/1520-0493(1997)125<2531:LFVOSH>2.0.CO;2
32

33
34 Singh SK. 2016. Long-term streamflow forecasting based on ensemble streamflow prediction
35 technique: a case study in New Zealand. *Water Resources Management*, 30:2295–2309. doi:
36 10.1007/s11269-016-1289-7
37

38
39 Solomon A. Polvani LM. 2016. Highly significant responses to anthropogenic forcings of the
40 midlatitude jet in the Southern Hemisphere. *Journal of Climate*, 29:3463-3470. Stahl K, Demuth
41 S. 1999. Linking streamflow drought to the occurrence of atmospheric circulation patterns.
42 *Hydrological Sciences Journal-Journal des Sciences Hydrologiques* 44: 467-482. DOI:
43 10.1080/02626669909492240
44

45
46 Sturman AP, Tapper NJ. 2006. *The Weather and Climate of Australia and New Zealand*. Oxford,
47 Oxford University Press, 541pp.
48

49
50 Swart NC, Fyfe JC, Gillett N, Marshall GJ. 2015. Comparing trends in the Southern Annular
51 Mode and surface westerly jet. *Journal of Climate* 28: 8840-8859. DOI: 10.1175/JCLI-D-15-
52 0334.1
53

54
55 Tang Q, Zhang X, Duan Q, Huang S, Yuan X, Cui H, Li Z, Liu X. 2016. Hydrological monitoring
56 and seasonal forecasting: Progress and perspectives. *Journal of Geographical Science*, 26: 904-
57 920. DOI: 10.1007/s11442-016-1306-z
58
59
60

- 1
2
3 Thompson DWJ, Wallace JM. 2000. Annular Modes in the Extratropical Circulation. Part I:
4 Month-to-Month Variability. *Journal of Climate*, 13: 1000-1016.
5
6
7 Troup AJ. 1965. The Southern Oscillation. *Quarterly Journal of the Royal Meteorological Society*
8 91: 490-506. DOI: 10.1002/qj.49709139009
9
10 Ummenhofer CC, England MH 2007. Interannual extremes in New Zealand precipitation linked
11 to modes of Southern Hemisphere climate variability. *Journal of Climate* 20: 5418-5440. DOI:
12 10.1175/2007JCLI1430.1
13
14 Ummenhofer CC, Sen Gupta A, England MH. 2009. Causes of Late Twentieth-Century Trends
15 in New Zealand Precipitation. *Journal of Climate* 22: 3-19. DOI: 10.1175/2008JCLI2323.1
16
17
18 Viale M, Nunez MN. 2011. Climatology of winter orographic precipitation over the subtropical
19 Central Andes and associated synoptic and regional characteristics. *Journal of*
20 *Hydrometeorology* 12: 481-507. DOI: 10.1175/2010JHM1284.1
21
22
23 Ward PJ, Kummerow M, Lall U. 2016. Flood frequencies and durations and their response to El
24 Nino Southern Oscillation: Global analysis. *Journal of Hydrology*, 539: 358-378.
25 DOI:10.1016/j.jhydrol.2016.05.045
26
27 Webster CS, Kingston DG, Kerr T. 2015. Inter-annual variation in the topographic controls on
28 catchment-scale snow distribution in a maritime alpine catchment, New Zealand. *Hydrological*
29 *Processes*, 29: 1096-1109.
30
31
32 Wu Z, Dou J, Lin H. 2015. Potential influence of the November-December Southern
33 Hemisphere annular mode on the East Asian winter precipitation: a new mechanism. *Climate*
34 *Dynamics* 44: 1215-1226. DOI: 10.1007/s00382-014-2241-2
35
36 Yarnal B. 1993. *Synoptic Climatology in Environmental Analysis*. London, Belhaven Press
37 195pp.
38
39 Yuan X, Wood EF, Liang M, 2014. Integrating weather and climate prediction: Toward
40 seamless hydrologic forecasting. *Geophysical Research Letters*, 41. 5891-5896,
41 doi:10.1002/2014GL061076.
42
43
44 Yuan X, Wood E, Ma Z. 2015. A review on climate-model-based seasonal hydrologic
45 forecasting: physical understanding and system development. *Wiley Interdisciplinary Reviews:*
46 *Water*, 2: 523-536. DOI: 10.1002/wat2.1088
47
48
49 Zheng F, Li JP, Liu T. 2014. Some advances in studies of the climatic impacts of the Southern
50 Hemisphere annular mode. *Journal of Meteorological Research* 28: 820-835. DOI:
51 10.1007/s13351-014-4079-2
52
53
54
55
56
57
58
59
60

Table 1: River flow gauging station details. DJF is December, January, February; JJA is June, July, August. CV is coefficient of variation.

Index	Site ID	River	Catchment area (km ²)	DJF CV	JJA CV	Latitude	Longitude	Missing in %
1	46618	Mangakahia	246	0.87	0.44	-35.6297	173.851	0.81
2	46644	Wairua	544	1.24	0.60	-35.6541	174.151	2.69
3	9301	Kauaeranga	121	0.94	0.59	-37.1588	175.591	0.28
4	9213	Ohinemuri	305	0.94	0.50	-37.4165	175.715	0.24
5	43420	Waikato	12429	0.24	0.29	-37.4335	175.131	0.76
6	9205	Waihou	1130	0.27	0.29	-37.5484	175.707	0.32
7	16501	Motu	1393	0.65	0.44	-37.8655	177.632	0.27
8	15302	Tarawera	914	0.16	0.19	-37.9405	176.767	0.34
9	15514	Whakatane	1560	0.78	0.60	-38.0070	176.995	0.08
10	15412	Rangitaiki	2893	0.28	0.33	-38.0414	176.800	0.27
11	15901	Waioeka	662	0.80	0.51	-38.2257	177.314	0.71
12	15410	Whirinaki	509	0.55	0.49	-38.4802	176.748	0.56
13	33316	Ongarue	1075	0.56	0.48	-38.8638	175.238	0.00
14	1043459	Tongariro	772	0.29	0.29	-38.9963	175.814	0.00
15	21801	Mohaka	2370	0.55	0.41	-39.0751	177.129	3.86
16	1043466	Waihohonu	95.8	0.22	0.29	-39.2180	175.736	0.85
17	33107	Whangaehu	471	0.21	0.28	-39.5032	175.463	0.25
18	33111	Mangawhero	511	0.72	0.45	-39.5697	175.264	0.36
19	33301	Whanganui	6643	0.67	0.46	-39.7778	175.145	0.00
20	32702	Rangitikei	2787	0.59	0.40	-39.8108	175.808	0.00
21	29224	Waiohine	184	0.62	0.34	-41.0181	175.400	0.16
22	29202	Ruamahanga	2340	0.76	0.45	-41.1986	175.440	0.22
23	57008	Motueka	163	0.59	0.50	-41.6348	172.913	1.18
24	93203	Buller	6350	0.49	0.43	-41.8360	171.699	0.80
25	60114	Wairau	505	0.45	0.47	-41.8948	172.921	1.42
26	93207	Inangahua	234	0.56	0.48	-42.1244	171.878	0.76
27	64606	Waiau	74	0.51	0.44	-42.2214	172.650	5.74
28	62105	Clarence	440	0.63	0.40	-42.4589	172.906	0.00
29	65104	Hurunui	1060	0.51	0.42	-42.7920	172.542	0.19
30	66204	Ashley	472	1.01	0.58	-43.2316	172.217	1.34
31	66401	Waimakariri	3210	0.46	0.39	-43.4165	172.652	0.41
32	68001	Selwyn	163	0.94	0.66	-43.4652	171.894	4.19
33	69618	Opihi	406	0.81	0.72	-44.1698	170.942	2.25
34	69621	Rocky Gully	22.4	0.83	0.79	-44.3260	170.774	4.86
35	71103	Hakataramea	899	1.01	0.96	-44.7262	170.490	0.46
36	75262	Kawarau	4302	0.34	0.33	-45.0081	168.868	0.70
37	78607	Oreti	1139	0.57	0.45	-45.7178	168.428	0.10
38	75232	Pomahaka	1924	0.81	0.45	-46.0607	169.401	0.00
39	77504	Mataura	766	0.62	0.42	-46.1000	168.949	0.05
40	75207	Clutha	20582	0.35	0.23	-46.2385	169.748	0.11

Table 2: Principal Components Analysis results showing percentage variance explained by seven principal components. DJF is December, January, February; JJA is June, July, August.

Principal Component	DJF	JJA
PC1	39.1	43.4
PC2	17.2	14.2
PC3	10.1	8.3
PC4	5.8	6.6
PC5	4.8	5.5
PC6	3.7	3.4
PC7	2.9	2.7
Total	83.5	84.1

For Peer Review

Table 3: River flow regions and associated gauging station membership. CWLNI is central west lower North Island, CTB is Canterbury, NICor is Northland Coromandel, LoSI is lower South Island, UpSI is upper South Island, BoPHB is Bay of Plenty Hawkes Bay.

CWLNI	CTB	NICor	LoSI	UpSI	BoPHB
Ongarue	Ashley	Mangakahia	Kawarau	Motueka	Motu
Whangaehu	Selwyn	Wairua	Oreti	Wairau	Waioeka
Mangawhero	Opihi	Kauaeranga	Mataura	Waiau	Whakatane
Whanganui	Rocky Gully	Ohinemuri	Pomahaka	Clarence	Rangitaiki
Rangitikei	Hakataramea	Waihou (6)	Clutha	Hurunui	Tarawera
Waiohine				Waimakariri	Whirinaki
Ruamahanga				Buller	Mohaka
Tongariro (6)				Inangahua	
Waihohonu (6)					
Waikato (6)					

Table 4: Climate and river flow correlations for (a) DJF and (b) JJA.. SAM is Southern Annular Mode, RainT is total rainfall, Tman is mean temperature, Tmax is maximum temperature. DJF is December, January, February; JJA is June, July, August. * and ** indicate statistical significance at the 0.05 and 0.01 levels respectively.

(a) DJF	SAM	RainT	Tmean	Tmax
NICor	-0.039	-	-	-
BoPHB	-0.190*	0.471**	-0.04	-0.165
CWLNI	0.127	0.197**	-0.099	-0.164
UpSI	-0.142	0.294**	-	-0.393**
CTB	0.11	0.559**	-	-0.618**
LoSI_{PC4}	-0.137	-0.114	-0.180*	-0.148
LoSI_{PC6}	-0.079	0.437**	-	-0.291**

(b) JJA	SAM	RainT	Tmean	Tmax
NICor	0.019	-	-	-
BoPHB	0.101	0.561**	0.207**	0.098
CWLNI_{PC1}	-	0.197**	-0.006	-0.109
CWLNI_{PC6}	-	0.424**	0.15	0.042
UpSI	0.005	0.572**	0.496**	0.441**
CTB	0.192*	0.549**	-0.12	-
LoSI	-0.177*	0.485**	-0.063	-

Table 5: Concurrent and one season ahead SAM river flow associations at the seasonal level for DJF and JJA (n = 32). * and ** indicate statistical significance at the 0.05 and 0.01 levels respectively. SON is September, October, November while MAM is March, April, May.

Region	DJF River flow		JJA River flow	
	SON SAM	DJF SAM	MAM SAM	JJA SAM
NICor	-0.210	-0.09	-0.053	0.117
BoPHB	0.180	-0.283	-0.055	0.354**
CWLNI	-0.437**	0.181	-0.207	-0.360**
UpSI	0.119	-303*	0.001	0.036
CTB	0.537**	0.267	-0.080	0.048
LoSI	0.176	-0.090	-0.455**	-0.312*

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



Figure 1: River flow gauging stations selected for the analysis




Figure 2: Principal component one loading maps for (a) DJF and (b) JJA.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 3: River flow regionalisation for (a) DJF and (b) JJA. See Table 3 for river flow region membership




Figure 4: Seasonal distribution of river flow as portrayed by representative gauging stations for the six river flow regions. Vertical axis is the logarithm of flow. (a) Kauarenga River, NICOR region; (b) Tarawera River, BOPHB region; (c) Ongarue River, CWLNI region; (d) Waimakariri River, UPSI region; (e) Hakataramea River, CTB region; (f) Oreti River, LOSI region. In the box plots the central bar is the median value while the upper and lower bars of the box plot are the 75th and 25th percentile respectively. The lines below and above the box represent the lowest datum still within the 1.5 inter-quartile range (IQR) of the lower quartile, and the highest datum still within the 1.5 IQR of the upper quartile. The open circles represent data points beyond the 1.5 quartile range.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 5: Principal component score (as a proxy of river flow) distribution by region for positive (+SAM) and negative (-SAM) phases of SAM for (a) DJF and (b) JJA as revealed by the circulation to environment approach. The form of the box plots is as described for Figure 4.




Figure 6: Distribution of SAM Index values for high and low PC scores (proxy for river flow) for (a) DJF and (b) JJA for six river flow regions as revealed by the environment to circulation approach. For DJF (a) (i) is BOPHB region; (ii) is NICOR region; (iii) is CTB region; (iv) is UPSI region; (v) is LOSI region; (vi) is CWLNI region. For JJA (b) (i) is CWLNI region; (ii) is UPSI region; (iii) is CTB region; (iv) is NICOR region; (v) is LOSI region; (vi) is BOPHB region. The form of the box plots is as described for Figure 4.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 7: Time series plots of PC scores (proxy of river flow) for (a) Summer: BoPHB and LoSI regions and (b) Winter: BoPHB and CWLNI regions.




Figure 8: Sea level pressure anomaly plots for (a) upper and (b) lower 25 percent of CWLNI river flows. Plots are based on 20th Century Reanalysis V2 data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <http://www.esrl.noaa.gov/psd/>




Figure 9: Wind anomaly plots for (a) upper and (b) lower 25 percent of CWLNI river flows. Plots are based on 20th Century Reanalysis V2 data provided by the NOAA/OAR/ESRL PSD, Boulder, Colorado, USA, from their Web site at <http://www.esrl.noaa.gov/psd/>