

1 **Deglaciation of the Laurentide Ice Sheet from the Last**

2 **Glacial Maximum**

3
4 Chris R. Stokes

5 *Department of Geography, Durham University, Durham, DH1 3LE, UK*

6 c.r.stokes@durham.ac.uk

7
8
9 **ABSTRACT:** The last deglaciation of the Laurentide Ice Sheet (LIS) was associated with
10 major reorganisations in the ocean-climate system and its retreat also represents a valuable
11 analogue for understanding the rates and mechanisms of ice sheet collapse. This paper reviews
12 the characteristics of the LIS at its Last Glacial Maximum (LGM) and its subsequent
13 deglaciation, with particular emphasis on the pattern and timing of ice margin recession and
14 the driving mechanisms of retreat. The LIS initiated over the eastern Canadian Arctic ~116-
15 110 ka (MIS 5d), but its growth towards the LGM was highly non-linear and punctuated by
16 several episodes of expansion (~65 ka: MIS 4) and retreat (~50-40 ka: MIS 3). It attained its
17 maximum position around 26-25 ka (MIS 2) and existed for several thousand years as an
18 extensive ice sheet with major domes over Keewatin, Foxe Basin and northern
19 Quebec/Labrador. It extended to the edge of the continental shelf at its marine margins and
20 likely stored a sea-level equivalent of around 50 m and with a maximum ice surface ~3,000 m
21 above present sea-level. Retreat from its maximum was triggered by an increase in boreal
22 summer insolation, but areal shrinkage was initially slow and the net surface mass balance was
23 positive, indicating that ice streams likely played an important role in reducing the ice sheet
24 volume, if not its extent, via calving at marine margins. Between ~16 and ~13 ka, the ice sheet

25 margin retreated more rapidly, particularly in the south and west, whereas the north and east
26 underwent only minimal recession. The overall rate of retreat decreased during the Younger
27 Dryas (YD), when several localised readvances occurred. Following the YD, the ice sheet
28 retreated two to five times faster than previously, and this was primarily driven by enhanced
29 surface melting while ice streams reduced in effectiveness. Final deglaciation of the Keewatin
30 and Foxe Domes, left a remnant Labrador Dome that disappeared ~6.7 ka.

31

32 **Keywords:** Laurentide Ice Sheet; Last Glacial Maximum; deglaciation; ice streams

33

34 **1. Introduction**

35 The North American Laurentide Ice Sheet (LIS) was the largest ice sheet to grow and decay
36 during the last glacial cycle, dominating Late Pleistocene fluctuations in global sea-level
37 (Lambeck *et al.*, 2014) and delivering the largest contribution to early Holocene sea level rise
38 (Tarasov *et al.*, 2012; Peltier, 2004). Accurate reconstructions of its extent, volume and
39 dynamics are, therefore, critical to our understanding of glacial-interglacial cycles and the
40 sensitivity of ice sheets to climate change (Clark *et al.*, 2009; Carlson and Clark, 2012).
41 Knowledge of its deglaciation is also required to understand the rates, magnitude and
42 mechanisms of ice sheet decay and associated impacts on sea level (Carlson *et al.*, 2008;
43 Carlson and Winsor, 2012; Kleman and Applegate, 2013; Stokes *et al.*, 2016), which is relevant
44 to assessments of the future stability of modern-day ice sheets in Greenland and Antarctica
45 (IPCC, 2013; Nick *et al.*, 2013; Ritz *et al.*, 2015). It is also clear that, in addition to responding
46 to climate forcing, the behaviour of the LIS was capable of driving abrupt climate change
47 through the delivery of both meltwater and icebergs that perturbed the ocean-climate system
48 (Barber *et al.*, 1999; Clark *et al.*, 2001). More broadly, the configuration and retreat history of

49 the LIS was an important constraint on the migration and dispersal of flora and fauna (Shapiro
50 *et al.*, 2004), including early humans (Goebel *et al.*, 2008; Eriksson *et al.*, 2012; Dixon, 2013;
51 Pedersen *et al.*, 2016).

52 Given its size and importance, the LIS is one of the most widely-studied palaeo-ice
53 sheets and there are hundreds of papers that have attempted to reconstruct its extent and
54 dynamics using a variety of both empirical and modelling approaches (see review in Stokes *et*
55 *al.*, 2015). However, the majority of papers, especially those taking an empirical approach,
56 have tended to focus on specific regions and time periods, and fewer papers have attempted to
57 summarise both the timing and driving mechanisms of deglaciation since the global Last
58 Glacial Maximum (gLGM). Building on several major syntheses over the last few decades
59 (Denton and Hughes, 1981; Dyke and Prest, 1987; Fulton, 1989; Dyke, 2004), this paper aims
60 to provide an up-to-date review of the LIS at the gLGM with an emphasis on the pattern and
61 timing of its deglaciation and the mechanisms that lead to its demise. Following an overview
62 of the characteristics of the LIS at its Local LGM (LLGM) in Section 2, Section 3 focusses on
63 the pattern and timing of deglaciation, followed by a discussion of the mechanisms that have
64 been invoked to explain deglaciation in Section 4. Some of the associated impacts of
65 deglaciation, such as the origin of Heinrich events (e.g. Andrews, 1998) and major meltwater
66 pulses and routing (e.g. Tarasov *et al.*, 2012; Gregoire *et al.*, 2012) are beyond the scope of the
67 present paper and will receive less attention (see comprehensive reviews by Hemming, 2004;
68 Carlson and Clark, 2012).

69 For the purposes of this paper, I use ‘Laurentide Ice Sheet’ in its broadest sense and,
70 except where indicated explicitly. I include the Innuitian Ice Sheet (IIS) (Dyke *et al.*, 2002)
71 and other small ice caps (e.g. in Newfoundland and the Appalachians) with which it was
72 contiguous for most of its history. This does not include the Cordilleran Ice Sheet (CIS), which
73 was a separate ice sheet except for brief periods during glacial maxima (Prest, 1969; Dyke and

74 Prest, 1987; Dyke *et al.*, 2002; Stokes *et al.*, 2012). For consistency, all dates are quoted in
75 thousands of calendar years (ka) before present. Where the original source used only
76 radiocarbon ages (e.g. Dyke and Prest, 1987), they have been converted to calendar years using
77 a mixed marine and Northern Hemisphere atmosphere calibration curve (Stuiver *et al.*, 2017)
78 and the original radiocarbon dates appear in parentheses (^{14}C ka).

79

80

81 **2. The Laurentide Ice Sheet at its Last Glacial Maximum**

82 *2.1. Inception and build-up to its Last Glacial Maximum*

83 Before describing the characteristics of the LIS at its LGM, it is useful to briefly outline its
84 inception and growth since the last interglacial during Marine Isotope Stage 5 (MIS 5).
85 Unfortunately, this aspect of the ice sheet's history is very poorly constrained compared to the
86 post-LGM period, largely because of the fragmentary nature of the terrestrial evidence relating
87 to ice sheet build-up, most of which was erased by the much larger Late Wisconsinan (MIS 2)
88 ice sheet. This has perhaps led to an over-reliance on numerical ice sheet models of pre-LGM
89 ice sheet configurations (e.g. Marshall *et al.*, 2000; Kleman *et al.*, 2002; Stokes *et al.*, 2012)
90 which are themselves limited by the availability of constraint data. However, the ocean-
91 sediment record has proved particularly useful for investigating pre-LGM iceberg fluxes and
92 meltwater events (e.g. Andrews and MacLean, 2003; Hemming, 2004), and there are pockets
93 of evidence in the glacial geomorphological and stratigraphic record (e.g. Kleman *et al.*, 2010)
94 that have survived modification and, in some places, been dated to periods prior to the LGM
95 (e.g. Allard *et al.*, 2012; Dalton *et al.*, 2016).

96 During the penultimate glacial maximum around 140 ka (MIS 6), the LIS is known to
97 have been smaller than its LGM (MIS 2) counterpart, and is thought to have been similar in

98 size to its extent around 13 ka (Colleoni *et al.*, 2016). This is consistent with global sea level
99 records and empirical evidence that indicates that the Eurasian Ice Sheet was larger during MIS
100 6 than during MIS 2 (Svendsen *et al.*, 2004). Indeed, the smaller size of the LIS is consistent
101 with changes in large-scale atmospheric circulation that facilitated the development of a larger
102 Eurasian Ice Sheet during MIS 6 (Colleoni *et al.*, 2016). Little is known about the deglaciation
103 of the LIS at the end of MIS 6, but a major glacial lake outburst flood has been reported from
104 a proximal marine core in the Labrador Sea around 124 ka (Nicholl *et al.*, 2012), which may
105 be analogous to the widely reported drainage of glacial Lake Agassiz during the final
106 deglaciation of the LIS around 8.2 ka (e.g. Barber *et al.*, 1999). Following this event, the
107 general consensus is that there was virtually no ice cover in North America during the peak of
108 the MIS 5 (Sangamonian) interglacial (~125-122 ka), which is primarily based on ages obtained
109 from organic-rich sediments in the Hudson Bay Lowlands (e.g. Allard *et al.*, 2011; Dalton *et*
110 *al.*, 2016) and a widespread acknowledgement that global sea levels were 6-9 m higher than
111 present during MIS 5e (e.g. Dutton *et al.*, 2015).

112 The consensus from both empirically-based arguments and numerical modelling is that
113 the LIS initiated over the Arctic/subarctic plateaux along the eastern seaboard of Canada (e.g.
114 Ives, 1957; Ives *et al.*, 1975; Marshall *et al.*, 2000; Marshall and Clark, 2002; Kleman *et al.*,
115 2002; Stokes *et al.*, 2012; Abe-Ouchi *et al.*, 2013). These are locations where only a small
116 decrease in temperature resulted in a large decrease in the equilibrium line altitude (ELA) – a
117 process termed ‘instantaneous glaciation’ (Koerner, 1980). It is thought that an embryonic
118 dome formed over Labrador during MIS 5d (cf. Andrews and Mahaffy, 1976; Boulton *et al.*,
119 1985; Vincent and Prest, 1987; Clark *et al.*, 1993; Marshall *et al.*, 2000; Kleman *et al.*, 2010),
120 possibly as early as 116-114 ka, and with some modelling (Stokes *et al.*, 2012) indicating a
121 large but thin ice sheet at 110 ka that covered 70-80% of the area occupied by the MIS 2 ice
122 sheet (see also Vincent and Prest, 1987; Boulton and Clark, 1990a, b; Clark *et al.*, 1993). This

123 “*explosive ice sheet growth*” (Marshall, 2002: p. 133) during MIS 5d is consistent with records
124 of a rapid fall in global sea level around that time (Marshall *et al.*, 2000; Cutler *et al.*, 2003),
125 but some workers suggest more minimal ice volumes in North America (~2-3 m of sea level
126 equivalent: Kleman *et al.*, 2002) and that the LIS did not grow substantially until MIS 4 (e.g.
127 Kleman *et al.*, 2002; Marshall and Clark, 2002; Kleman *et al.*, 2010). If the ice sheet was
128 relatively large during MIS 5d (e.g. the ~20 m of sea level equivalent modelled by Stokes *et*
129 *al.*, 2012), it had shrunk rapidly by 100 ka (MIS 5c) (cf. St-Onge, 1987), and likely existed
130 only as a small, thin ice sheet over the original inception grounds in north-eastern Canada by
131 ~80 ka (MIS 5a) (Marshall *et al.*, 2000; Stokes *et al.*, 2012). Thereafter, the LIS is thought to
132 have grown rapidly during MIS 4, reaching a maximum extent around 65 ka (Vincent and Prest,
133 1987; Marshall *et al.*, 2000; Kleman *et al.*, 2002; Stokes *et al.*, 2012), which coincides with the
134 oldest recognised Heinrich event (H6) and a marked increase in ice-rafted debris from that time
135 (Kirby and Andrews, 1999; Hemming, 2004, Bassis *et al.*, 2017).

136 Following an MIS 4 maximum that may have been almost as large as the MIS 2 (LGM)
137 volume according to some models (Marshall *et al.*, 2000; Stokes *et al.*, 2012), the ice sheet
138 retreated to a mid-Wisconsinan (early MIS 3) minimum at some point between 60 and 40 ka
139 (Dredge and Thorleifson, 1987; Clark *et al.*, 1993; Kleman *et al.*, 2010; Stokes *et al.*, 2012).
140 Indeed, the extent of the ice sheet during MIS 3 is very poorly constrained (e.g. see review in
141 Dredge and Thorleifson, 1987), with numerical modelling indicating a relatively large ice sheet
142 that stored up to 30 m of sea level equivalent at 55 ka (Marshall *et al.*, 2000; Stokes *et al.*,
143 2012), but with a suite of new dates raising the possibility that the Hudson Bay Lowlands, close
144 to the geographic centre of the ice sheet, were completely ice free between ~50 and ~40 ka (see
145 Dalton *et al.*, 2016). Following the MIS 3 minimum, the ice sheet underwent gradual expansion
146 that was punctuated by episodes of successively less recession (e.g. at 30 ka) before a final
147 rapid growth towards the maximum LGM position (Dyke *et al.*, 2002; Stokes *et al.*, 2012).

149 *2.2. The Timing of the Local Last Glacial Maximum Laurentide Ice Sheet (Late Wisconsinan)*

150 In its broadest sense, the global LGM (gLGM) is conventionally defined from sea-level records
151 “as the most recent interval in Earth history when global ice sheets reached their maximum
152 integrated volume” (Clark *et al.*, 2009: p. 710). It has been recognised for some time, however,
153 that because global sea levels are an integrated signal of ice volume, this does not imply that
154 all ice sheets, or even various sectors within the same ice sheet, reached their ‘Local’ Last
155 Glacial Maximum (from hereon LLGM) extent simultaneously (Clark *et al.*, 2009; Hughes *et*
156 *al.*, 2013). In a recent synthesis, Clark *et al.* (2009) constrained the timing of the gLGM period,
157 based on relative sea-level data, as occurring from 26.5 to 19.0 ka, and suggested that this
158 broadly coincided with the duration of maximum extent of most global ice sheets, including
159 the LIS. They noted, however, that the LLGM of the various sectors of the LIS were
160 asynchronous (albeit with large uncertainties), with some margins (e.g. in the south) potentially
161 reaching their maximum early, perhaps even prior to, the gLGM and others occurring much
162 later (e.g. the Maritime provinces in the south-east). Indeed, Dyke *et al.* (2002) suggested that
163 ice advanced to its Late Wisconsinan (MIS 2) limit in the northwest, northeast and south about
164 27-28 ka (23-24 ¹⁴C ka), and in the southwest and far north about ~24-25 ka (20-21 ¹⁴C ka).
165 More recently, a number of studies have shown that ice sheet margin in the far north-west, in
166 the vicinity of the Mackenzie River delta and along the Richardson Mountains, attained its
167 maximum position relatively late and certainly less than 20 ka (e.g. Murton *et al.*, 2007;
168 Kennedy *et al.*, 2010; Lacelle *et al.*, 2013), possibly as a short-lived advance between 17 and
169 15 ka (Murton *et al.*, 2015). Thus, the consensus is that - overall - the LIS reached its local
170 maximum extent early in the gLGM period (cf. Dyke *et al.*, 2002), but with some margins
171 advancing much later, e.g. in the far north-west. Most recent modelling experiments converge

172 on maximum volumes ~26-25 ka (e.g. Tarasov *et al.*, 2012; Stokes *et al.*, 2012; Abe-Ouchi *et*
173 *al.*, 2013), although some place it closer to 21-20 ka (e.g. Marshall *et al.*, 2000).

174 It is very likely that the LIS existed at its near-maximum extent for several thousand
175 years (cf. Dyke *et al.*, 2002; Tarasov *et al.*, 2012). Given that it grew to this position from a
176 relatively large ice sheet late in MIS 3 (Dyke *et al.*, 2002; Stokes *et al.*, 2012; Tarasov *et al.*,
177 2012), the prolonged duration of its maximal configuration suggests that, for the most part, it
178 had a surface geometry and mass balance in equilibrium with the gLGM climate for a few
179 thousand years (Dyke *et al.*, 2002).

180

181 *2.3. Extent and Thickness of the LIS at its Local Last Glacial Maximum (Late Wisconsinan)*

182 The maximum extent and thickness of the ice sheet during its Late Wisconsinan maximum has
183 been the subject of debate for over 150 years (e.g. Bell, 1884) and, despite numerous studies
184 on this subject, consensus has only recently emerged (Dyke *et al.*, 2002). A comprehensive
185 review of the literature on this subject is beyond the scope of this paper (see Ives (1978) and
186 Dyke *et al.* (2002) for authoritative reviews), but it is useful to summarise key areas of
187 contention and consider how different ideas have evolved and, more often than not, been
188 revisited.

189 Much of the early work on the extent and thickness of the LIS during its LLGM (e.g.
190 Bell, 1884; Daly, 1902; Coleman, 1920) focussed on the mountains of the east coast of Canada
191 and argued that many of the highest peaks (e.g. the Torngat of northern Labrador) either
192 remained as nunataks or were only affected by local ice caps or glaciers. This ‘minimum’ model
193 was based on the identification of erosional trimlines (e.g. at 650 m in the Torngat Mountains,
194 see Daly, 1902) and the presence of frost shattered bedrock and blockfields above these limits
195 (e.g. Coleman, 1920). These interpretations were first questioned by Odell (1933) who reported

196 high-level erratics and poorly preserved striations at 1,446 m in the Torngat Mountains, thus
197 arguing that the last ice sheet had overtopped the mountains, and that block-fields formed after
198 deglaciation. Similar observations informed similar interpretations by Flint *et al.* (1942) in the
199 Shickshock (Chic-Choc) Mountains and Flint's hypothesis for the inception and growth of the
200 LIS (Flint, 1943) called for a highland origin and windward-growth that subsequently
201 inundated the high coastal mountains. This 'maximum' model (Fig. 1) was adopted in a series
202 of major publications (e.g. Flint, 1947; 1971) and on a new 'Glacial Map of North America'
203 (Flint *et al.*, 1945), which was one of the first attempts (see also Chamberlin, 1913) to
204 synthesise the glacial features in detail and on a large scale. An important corollary of Flint's
205 'maximum' model was that the ice sheet was viewed as a monolithic single-domed ice sheet
206 centred over Hudson Bay, although earlier workers had suggested alternative multi-domed
207 configurations (Tyrell, 1898; Coleman, 1920).

208 As noted by Ives (1978), Flint's maximum model appeared to have been widely
209 accepted and clearly influenced the boundary conditions for the first CLIMAP (Climate: Long
210 range Investigation, MApping, and Prediction) reconstruction of the 'Ice-Age' earth
211 (CLIMAP, 1976; Denton and Hughes, 1981), in addition to becoming firmly entrenched in
212 high school and University curricula. Indeed, when Prest (1969) produced one of the most
213 detailed maps of the retreat of the ice sheet (discussed in Section 3; see Fig. 7), his Late
214 Wisconsinan limit extended, for the most part, on to the continental shelf along the east coast
215 from northern Baffin Island, all the way down to the Atlantic provinces, and covered most of
216 the Canadian Arctic Archipelago (apart from Banks Island). He also depicted an extensive
217 southern margin that transgressed well into northern USA and as far south at 40 degrees.

218 Despite the ascendancy of the maximum model from the 1940s, a number of papers in
219 the 1950s, 1960s and 1970s (Ives, 1957; Andrews and Miller, 1972; Miller and Dyke, 1974)
220 had questioned it on the basis that several different 'weathering zones' could be distinguished

221 by their relative maturity (see review in Ives, 1978), with the oldest weathering zones
222 interpreted to pre-date the last glacial maximum and indicating ice-free refugia. In some
223 locations, this interpretation was further strengthened by a small number radiocarbon dates that
224 gave ages much older than the Late Wisconsinan (e.g. Løken, 1966). Those arguing for a return
225 to a more minimal model interpreted the high-altitude erratics and evidence of glacial abrasion
226 (e.g. Odell, 1933) to be from a much older (pre-Late Wisconsinan) glaciation, although it was
227 become increasingly recognised that they might also have been preserved beneath more recent
228 cold-based ice (e.g. Sugden, 1977; Sugden and Watts, 1977).

229 Similar debates were being played out along other parts of the ice sheet margin, with
230 multiple weathering zones and, in some cases, correlative till sheets being used to infer reduced
231 ice sheet extent on the Queen Elizabeth Islands (England, 1976a, 1976b) and Banks Island
232 (Vincent, 1982); and even at the south-western margin of the ice sheet, where drift previously
233 thought to have been of Late Wisconsinan age was subdivided on the basis of morphological
234 degradation, with the fresher drift delimiting the last ice cover (Stalker, 1977). Thus, in a
235 comprehensive review of a rapidly-growing body of literature, and on the basis of a large
236 amount of fieldwork carried out since the 1950s, Ives (1978) called for a return to the minimum
237 model that had prevailed prior the 1940s and with Late Wisconsinan limits well behind those
238 proposed by Prest (1969) at the north, eastern and south-eastern margins of the ice sheet, i.e.
239 with localised refugia around much of the ice sheet's perimeter and on the continental shelf.
240 Ives (1978) also emphasised that the minimum model implied a much reduced ice thickness
241 and that it was unlikely to have been a simple, monolithic dome with maximum ice thicknesses
242 over Hudson Bay, as originally envisaged by Flint.

243 By the early 1980s, therefore, a large body of work had argued for a retraction of the
244 Late Wisconsinan limit and "*adherents of minimum ice sheet models*" (Boulton *et al.*, 1985: p.
245 452) adopted a more restricted margin, but this was not universally accepted and the debate

246 continued (e.g. Hughes *et al.*, 1977; Denton and Hughes, 1981). Indeed, Dyke and Prest, 1987)
247 noted that Prest (1984) was unable to portray a single Late Wisconsinan limit that met with any
248 consensus and he instead showed a minimum and maximum limit, with the maximum similar
249 to his 1969 reconstruction.

250 It was in the context of this highly contentious body of literature, that Dyke and Prest
251 (1987) produced one of the most influential reconstructions of the pattern and timing of the
252 LIS that would act as a benchmark for several decades. Their maximum extent at 21.4 ka (18
253 ¹⁴C ka BP) was clearly influenced by the growing body of evidence for a retracted ice margin
254 along the northern and eastern coasts of Canada, with the Torngat Mountains (and parts of the
255 Appalachians, including the Shickshock Mountains) protruding as nunataks, and with large
256 areas of Baffin Island and the Queen Elizabeth Islands ice-free, together with Banks Island (see
257 [Fig. 2](#)). Acknowledging much larger uncertainty, Dyke and Prest (1987) also depicted major
258 ice shelves in association with the Appalachian ice complex and others extending off the coast
259 of Labrador, together with ice shelves in the Gulf of Boothia/Lancaster Sounds and in M'Clure
260 Strait. Elsewhere, the southern margin (e.g. the Lake Michigan Lobe) extended south of 40
261 degrees (based on work by Clayton and Moran, 1982) and they depicted fully coalescent
262 Laurentide and Cordilleran ice sheets at this maximum extent that was, at that time, far more
263 controversial than it is now (cf. Stalker, 1977). A further significant component of the Dyke
264 and Prest (1987) reconstruction was that it clearly portrayed a multi-domed configuration at its
265 maximum, with centres of ice mass (domes) located over Labrador, Keewatin and Foxe Basin,
266 and with major ice divides emanating from them ([Fig. 2](#)). This geometry attempted to reconcile
267 new evidence from erratic dispersal trains that clearly indicated a complex multi-domed
268 configuration (Shilts *et al.*, 1979; Shilts, 1980). Dyke and Prest (1987) also discussed the
269 importance of ice streams and the availability of 'soft' deformable sediments (cf. Fisher *et al.*,
270 1985) in influencing the ice surface topography (see also Section 2.4), noting that many of the

271 ice lobes at the southern margin of the ice sheet had extremely low ice surface gradients
272 (Mathews, 1974).

273 The pendulum swung again in the in the mid-1990s (cf. Miller *et al.*, 2002) when new
274 lines of evidence were uncovered to interpret a more extensive Late Wisconsinan limit than
275 had been portrayed by Dyke and Prest (1987), particularly at its northern margin, but also along
276 the eastern margin and in the Atlantic provinces. As noted by Dyke *et al.* (2002: p. 11): “*after*
277 *a century of debate, intensively for the last 25 years, about the existence of an Innuitian Ice*
278 *Sheet during the LGM over the northern half of the Canadian Arctic Archipelago, a consensus*
279 *has emerged that such an ice sheet did in fact cover most of that region*”. This was based on
280 new glacial geological evidence of ice streams within several inter-island channels and large
281 fjord systems (e.g. Blake, 1992, 1993; Dyke 1999; Lamoureux and England 2000; Ó Cofaigh
282 *et al.* 2000) and numerous sets of lateral meltwater channels that descended to marine limits of
283 early Holocene age (Dyke, 1999; England, 1999; England *et al.*, 2000). This appeared to offer
284 conclusive evidence that the Innuitian Ice Sheet (IIS) extended offshore at its maximum extent
285 and that it was fully coalescent with the Greenland Ice Sheet along Nares Strait in the east, and
286 with Laurentide ice along Parry Chanel in the south, which was originally proposed by Blake
287 (1970) and illustrated by Prest (1969), see [Fig. 3](#). The limits of the ice sheet in the north-west
288 were much more uncertain, but Dyke *et al.* (2002) portrayed the whole of Prince Patrick Island
289 as ice-free, and large parts of Melville Island and most of Banks Island as unglaciated ([Fig. 3](#)).

290 The other region that benefitted from increased scrutiny from the mid-1980s onwards,
291 particularly on the continental shelf, was the south-eastern margin of the ice sheet (cf. Miller
292 *et al.*, 2002). Dyke and Prest (1987) had portrayed large ice-free areas at the ice sheet’s
293 maximum extent, but new cosmogenic dating of intensively weathered terrains suggested that
294 they could have been covered by cold-based ice (Gosse *et al.*, 1995). New ¹⁴C AMS
295 radiocarbon ages from marine sediments on the continental shelf off Nova Scotia and southern

296 Newfoundland also dated sediment above the youngest till to be of post-LGM ages (e.g. Amos
297 and Knoll, 1987; Bonifay and Piper, 1988; Gipp and Piper, 1989; Mosher *et al.*, 1989; Amos
298 and Miller, 1990; Forbes *et al.*, 1991; Gipp, 1994; King, 1996; Stea *et al.*, 1998; Josenhans and
299 Lehman, 1999). To the north, sedimentological studies and high-resolution AMS dating of
300 marine sediment cores from the SE Baffin and Labrador shelves, and adjacent slopes, led to a
301 reconsideration of LGM ice extent in that region (Dyke *et al.*, 2002). Jennings (1993), for
302 example, concluded that Cumberland Sound was filled by an ice stream until ~11.5 ka (~10
303 ¹⁴C ka) which Kaplan (1999) suggested may have extended onto the continental shelf (see also
304 Miller *et al.*, 2002). In northern Labrador, Clark and Josenhans (1990) combined marine and
305 terrestrial evidence to suggest that LGM ice was more extensive than previously mapped, with
306 the ice limit extending onto the continental shelf. Cosmogenic exposure dating (Marsella *et al.*,
307 2000) also confirmed extensive Late Wisconsinan outlet glaciers in the fiords of Cumberland
308 Peninsula.

309 Thus, numerous lines of evidence had been uncovered to suggest that previous assigned
310 'old' (pre-LGM) moraines on Cumberland Peninsula and northern Baffin Island were of Late
311 Wisconsinan age, leading to the most significant re-interpretations of the ice extent along the
312 north-eastern Laurentide margin for several decades, which was summarised in Dyke *et al.*
313 (2002). The Late Wisconsinan extent portrayed in Dyke *et al.* (2002) (Fig. 3) was subsequently
314 incorporated into the updated deglaciation sequence for North America (Dyke *et al.*, 2003).
315 This proposed LLGM extent has also been reproduced in more recent overviews and syntheses
316 (e.g. Dyke, 2004) and the margin positions have been used as constraint data for numerical
317 modelling of the LIS (e.g. Tarasov *et al.*, 2012; Peltier *et al.*, 2015).

318 The most dramatic changes to the LLGM extent of the LIS in the last decade has seen
319 the ice margin extended to cover the >70,000 km² Banks Island in the Western Canadian Arctic
320 (England *et al.*, 2009; Lakeman *et al.*, 2012, 2013) and the recognition that it likely extended

321 to (or close to) the continental shelf edge in Baffin Bay (Briner *et al.*, 2006) and in Atlantic
322 Canada (Shaw *et al.*, 2006) (see Fig 4). Banks Island had long been regarded as an ice-free
323 refugium (Prest, 1969; Vincent, 1982; Dyke and Prest, 1987) and was portrayed as such in the
324 most recent Dyke *et al.* (2003) synthesis (see also Dyke, 2004). A suite of new radiocarbon
325 dates and glacial geomorphological mapping, however, has clearly indicated that the LIS
326 inundated Banks Island during the LLGM (England *et al.*, 2009; Lakeman *et al.*, 2012, 2013).
327 Similar methods and new dates have also extended the ice margin over the entirety of Melville
328 Island and onto Eglinton Island (Nixon *et al.*, 2014) and it is likely that the ice also overran
329 Prince Patrick Island and extended onto the continental shelf along the entire IIS margin (see
330 Stokes *et al.*, 2016). At the same time, a large body of work has used cosmogenic dating of
331 high-elevation erratics close to fjord mouths (e.g. on Baffin Island) to demonstrate that a
332 relatively thick LIS must have terminated on the continental shelf during the LLGM (e.g.
333 Briner *et al.*, 2006), and with most high-elevation areas covered by non-erosive cold-based ice
334 that accounts for the preservation of highly weathered surfaces in those locations. In the
335 Atlantic provinces, a large body of work undertaken offshore has identified moraines, flutings,
336 till tongues, cross-shelf troughs and associated trough mouth fans, which attest to warm-based
337 ice streams extending to the edge of the continental shelf and separated by more stagnant ice
338 on the shallow banks (e.g. Mosher *et al.*, 1989; Piper and Skene, 1998; Schnikter *et al.*, 2001;
339 Shaw *et al.*, 2006). It is also noteworthy that data-calibrated modelling of the North American
340 Ice Sheet complex at the LLGM (Fig. 5) also generates a large multi-domed ice sheet that
341 extends to the edge of the continental shelf, with some of these major ice streams (see Fig. 4)
342 captured in the basal velocity pattern (Stokes and Tarasov, 2010).

343 In summary, after over 150 years, consensus appears to have been reached that at its
344 Late Wisconsinan maximum (~25-24 ka), the LIS was a large multi-domed ice sheet with a
345 southern margin that extended south of 40° in the Great Lakes region, with a western margin

346 that was fully coalescent with the Cordilleran Ice Sheet, and with northern and eastern margins
347 that extended to the edge of the continental shelf (Fig. 4).

348

349 2.4. Quantification of LIS volume at its Local Last Glacial Maximum (Late Wisconsinan)

350 Quantifying the volume of the LIS is important for reconciling records of global sea level
351 (Carlson and Clark, 2012; Lambeck *et al.*, 2017), but this has often proved difficult due to the
352 lack of nunataks and trimlines from interior regions (cf. Simon *et al.*, 2014). As noted by Dyke
353 *et al.* (2002: p. 20): “all that can be concluded from direct mapping is that the vast interior
354 region of the ice sheet, generally the part that was more than about 1000 km behind the margin,
355 lay more than 2000 m above present sea level”. Thus, despite numerous studies and debates
356 regarding the extent and geometry of the ice sheet (see Section 2.3), few have attempted to
357 quantify its volume.

358 Paterson (1972) was one of the first to consider the theory of ice flow and relate the
359 area of the ice sheet to its thickness and volume. Using previous maps of ice sheet extent (from
360 Prest, 1969), that were informed by Flint’s (1943) large monolithic reconstruction, he argued
361 that the LIS may have been up to 3.6 km thick at its maximum and comprised $26.5 \times 10^6 \text{ km}^3$
362 of ice (with volume errors estimated at 16%). During the 1980s, however, the importance of
363 ‘deforming beds’ was becoming increasingly recognised (cf. Alley *et al.*, 1986; Boulton and
364 Hindmarsh, 1987) and this clearly influenced attempts to reconstruct the LIS, with important
365 implications for its volume. In particular, numerical modelling experiments (e.g. Fisher *et al.*,
366 1985; Boulton *et al.*, 1985) clearly showed that the incorporation of deformable beds with low
367 basal shear stress beneath the ice sheet generated thinner ice and removed much of the radial
368 symmetry of some of the earlier reconstructions (e.g. Flint, 1943). Indeed, the difference
369 between Fisher *et al.*’s (1985) ‘soft bed’ and ‘hard bed’ models of the LIS was a volumetric

370 reduction of around 30% for the former. The incorporation of deforming beds also generated
371 multi-domed configurations that were deemed to be more compatible with the emerging
372 geological evidence at that time (e.g. Shilts *et al.*, 1979; Shilts, 1980; see Fig. 2). Later
373 modelling work by Clark *et al.* (1996) and Licciardi *et al.* (1998) also demonstrated that a
374 reduction in the effective viscosity of the till in regions underlain by ‘soft’ sediments generated
375 a multi-domed ice sheet with a large bowl-shaped depression over Hudson Bay and thin ice
376 (~1000 m above modern sea level) over the western and southern sectors of the ice sheet. A
377 thinner, multi-domed ice sheet was also consistent with inverse modelling of crustal rebound
378 and relative sea level data used in the early ICE-NG series (e.g. ICE-3G, Tushingham and
379 Peltier, 1991; ICE-4G, Peltier, 1994).

380 More recently, Peltier’s ICE-5G model of the ice load history (Peltier, 2004) indicated
381 much larger ice sheet thicknesses over the Keewatin region (> 4 km) and correspondingly
382 larger volumes for the LIS. Tarasov *et al.* (2012) also noted that their modelled ice thicknesses
383 over Hudson Bay were “possibly a kilometer too thick” (p. 37). Other work indicates thinner
384 ice in this region (e.g. Lambert *et al.*, 2006; Argus and Peltier, 2010; Mazzotti *et al.*, 2011),
385 including the most recent ICE-6G modelling (Peltier *et al.*, 2015), which shows ice thickness
386 over Keewatin that are 1.5 km thinner than ICE-5G (Vettoretti and Peltier, 2013). A suite of
387 new radiocarbon dates that constrain the relative sea-level history of Arviat on the west coast
388 of Hudson Bay are also consistent with a peak thickness of ~3.4 km at the LLGM (Simon *et*
389 *al.*, 2014).

390 Thus, although estimates of the LIS have varied quite dramatically over the last ~100
391 years, and this aspect of the ice sheet’s history remains poorly constrained (see Lambeck *et al.*,
392 2017), Table 1 indicates a convergence towards lower volumes in the more recent literature
393 (cf. Abe-Ouchi *et al.*, 2015), which are consistent with the influence of deforming beds and a
394 thinner, multi-domed configuration (Fisher *et al.*, 1985; Dyke and Prest, 1987; Clark *et al.*,

1996; Marshall *et al.*, 2000; Tarasov *et al.*, 2012; Lambeck *et al.*, 2017). At its Late Wisconsinan maximum, the LIS (including the IIS and the Appalachian Ice Complex) most likely contained around $20 \times 10^6 \text{ km}^3$, which is equivalent to $\sim 50 \text{ m}$ of global sea level (Clark *et al.*, 1996) (Table 1). This is around 500-1000 m lower in elevation than the original CLIMAP reconstructions (Denton and Hughes, 1981), which contained $34.2 \times 10^6 \text{ km}^3$ (85 m of global sea level), or around 75% more ice (Table 1).

401

Table 1: Chronological compilation of published estimates of the LIS extent, elevation and volume at its Local LGM (updated from Licciardi *et al.*, 1998). Where indicated (*), note that some higher values are due to the inclusion of the Cordilleran Ice Sheet because separate values for the LIS were not quoted.

Reference	Extent (10^6 km^2)	Maximum elevation (km above present sea level)	Volume ($\times 10^6 \text{ km}^3$)
Ramsay (1931)	15.75	2.9	45.45
Donn <i>et al.</i> (1962)	12.74	-	31.85 - 25.48
Andrews (1969)	11.82	-	26.0
Flint (1971)	13.39	-	29.46
Paterson (1972) ¹	11.6	2.7	26.5
Sugden (1977) ¹	-	3.5	37.0
Budd and Smith (1981) ¹	-	4-4.5*	<32*
Denton and Hughes (1981) (CLIMAP) ¹	-	3.8 (max. model) 3.5 (min. model)	34.2 (max. model) 30.5 (min. model)
Boulton <i>et al.</i> (1985) ¹	-	3-3.5 (hard bed model) >3 (soft bed model)	33-44 (hard bed model) (soft bed not reported)
Fisher <i>et al.</i> (1985) ¹	-	>3.2 (max. hard bed model) >3.2 (min. hard bed model) 2.8-3.2 (soft-bed model)	25.9 (max. hard bed model) 21.1 (min. hard bed model) 18.0 (soft-bed model)
Tushingham & Peltier (1991) (ICE-3G) ¹	-	>3	21.0
Peltier (1994) (ICE-4G) ¹	-	~ 3	19.0
Clark <i>et al.</i> (1996)	-	2-2.5	19.7

Marshall & Clark (1997a, b) ²	~14	4.2	36.4
Licciardi <i>et al.</i> (1998) ¹	-	3.1 (min. model) 3.6 (max. model)	15.9 (min. model) 19.7 (max. model)
Tarasov & Peltier (1999) ²	~13*	~3.8*	25*
Peltier (2004) (ICE-5G)	-	>4	-
Andrews (2006)	12	3-4	-
Tarasov <i>et al.</i> (2012)	-	-	28 (model nn9927)
Gregoire <i>et al.</i> (2012)	~16	>3	~35*
Lambeck <i>et al.</i> (2017)		≥3.5	

406 ¹Taken from Licciardi *et al.* (1998)

407 ²Cited in Marshall *et al.* (2000)

408

409

410 **3. Pattern and Timing of Deglaciation**

411 It was not until the 1960s that researchers attempted to systematically reconstruct the pattern
412 and timing of deglaciation at the scale of the entire ice sheet and produce maps of the ice
413 margins at specific time-steps (isochrones). One of the first to attempt to undertake this was
414 by Bryson *et al.* (1969) who utilised existing radiocarbon dates (289 in total) and geological
415 information to plot the ice sheet perimeter at 500 to 1,000 yr intervals through time from about
416 13 ka (Fig. 6). Key conclusions from that pioneering study were that the northern limit of the
417 ice sheet lay close to the Arctic mainland coast of Canada, now known to be incorrect (see
418 Section 2.3), and that the most dramatic retreat took place along the western margin, creating
419 an ice-free corridor from the Arctic Ocean to the Great Plains around 10.1 ka (9 ¹⁴C ka). They
420 also noted that the LIS “*retained its identity as a distinct unit*” (p. 1) until around 8.4 ka, which
421 they termed the Cockburn Phase. At the time, this was thought to be the “*only major glacial*
422 *pulsation*” (Bryson *et al.*, 1969: p. 7) that had been recognized stratigraphically and
423 geomorphologically (e.g. major moraine systems) over large areas of the eastern and central
424 Canadian Arctic (e.g. Falconer *et al.*, 1965). It was significant because it represented the final

425 phase of the contiguous LIS, with remnant domes over Keewatin, Labrador, and Foxe Basin-
426 Baffin Island. By around 8 ka, however, Bryson *et al.* (1969) argued that marine incursion into
427 Hudson Bay heralded the rapid disintegration of the ice sheet, with only a small ice cap
428 surviving to present day (Barnes Ice Cap).

429 Around the same time, Prest *et al.* (1968) produced their Glacial Map of Canada, which
430 summarised a vast amount of information and literature on the glacial geomorphology of the
431 ice sheet. This was soon followed by Prest's impressively detailed map of the 'Retreat of
432 Wisconsin and Recent Ice in North America' (Prest, 1969), which resembled Bryson *et al.*'s
433 (1969) synthesis in many respects, but was far more detailed and with isochrones portrayed at
434 a much higher temporal resolution (Fig. 7). So impressive was this compilation that an
435 anonymous author in the journal *Nature* wrote that it "*deserves a place on every class room*
436 *wall where earth sciences and American archaeology are taught...*" (Anonymous, 1970, p.
437 224). Denton and Hughes' (1981) impressive compendium on 'The Last Great Ice Sheets' also
438 presented a continental-scale synthesis of the LIS, but with a particular focus on its
439 configuration during the Late Wisconsinan and at 4-5 key time-steps during deglaciation (see
440 Chapter 2: Mayewski *et al.*, 1981). The next detailed synthesis of the ice retreat pattern was by
441 Boulton *et al.* (1985). A key conclusion from their reconstruction was the rapid retreat of the
442 southern margins and a very slow retreat at the northern margins of the LIS, perhaps reflecting
443 a strong N-S climate gradient during deglaciation. They also noted that overall ice margin
444 recession must have paused and that the margin maybe re-advanced at various locations during
445 overall deglaciation.

446 Building on a large number of reports and maps on the glacial geology of Canada, much
447 of it undertaken with impressive detail by the Geological Survey of Canada, Dyke and Prest
448 (1987) produced their influential reconstruction of the pattern and timing of the LIS retreat (see
449 Section 2.3). This comprised a series of palaeogeographic reconstructions at 11 time-steps (4

450 simplified examples are shown in Fig. 8) which included information on the ice sheet outline,
451 geometry and associated changes in proglacial lake drainage and relative sea-level oscillations;
452 and an accompanying map showing much higher resolution isochrones (Map 1702A: Dyke and
453 Prest, 1987). In part, the paper was motivated by several debates that had emerged since Prest's
454 (1969) map (Fig. 7), namely: (i) the location (extent) of the maximum Late Wisconsinan limit
455 (see Section 2.3), (ii) the surface geometry of the ice sheet (i.e. the location of major ice domes
456 and divides and their evolution through time), and (iii) the synchronicity of ice marginal
457 fluctuations in the north versus the south (Dyke and Prest, 1987).

458 In many respects, Dyke and Prest's (1987) reconstruction has remained the benchmark
459 for the last three decades, with the only major changes being the revisions to a more extensive
460 Late Wisconsinan maximum at the LGM (see Section 2.3) and a refined ice margin chronology
461 that has benefited from improvements in radiocarbon dating (mainly the advent of AMS dating
462 methods) and the 'retirement' of hundreds of earlier conventional radiocarbon dates (see Dyke,
463 2004). The updated ice margin chronology is described in Dyke (2004) and is available in
464 digital format in Dyke *et al.* (2003), which includes 36 time steps, starting 21.8 ka (18 ¹⁴C ka)
465 and ending at 0.9 ka (1 ¹⁴C ka). This new chronology is based on >4,000 dates that are spread
466 across the entire ice sheet bed and consist of mainly radiocarbon dates, supplemented with
467 varve and tephra dates, which constrain ice margin positions and shorelines of large glacial
468 lakes. Dates on problematic materials (e.g. bulk samples with probable blended ages) were
469 excluded in the Dyke *et al.* (2003) and Dyke (2004) chronologies and marine-shell dates, a
470 major component, were also adjusted for regionally variable marine-reservoir effects on the
471 basis of a new set of radiocarbon ages. The net effect is that deglaciation is delayed in most
472 places by 500-2000 years with respect to the Dyke and Prest (1987) reconstructions (cf. Dyke,
473 2004). However, the spatial pattern of ice recession resembles earlier reconstructions and the
474 pattern of deglaciation is described in Dyke and Prest (1987) and Dyke (2004), which form the

475 basis of the following discussion and to which the reader is referred for more detail (see also
476 the compendium in Fulton (1989), which covers several regions in impressive detail). The
477 following sections present a broad overview of the pattern and timing of deglaciation at the
478 continental scale, with potential driving mechanisms of these broad patterns discussed in
479 Section 4.

480

481 *3.1. Local LGM to early Late Glacial: ~25-17.6 ka (18-14.5 ¹⁴C ka)*

482 Dyke and Prest's (1987) reconstruction at the LLGM included ice flow patterns that were
483 informed by the distribution of glacial landforms (e.g. moraines, eskers and glacial lineations)
484 on Prest *et al.*'s (1968) Glacial Map of Canada. Ice flowlines intersected the margin at right
485 angles (unless they were in highly lobate areas with assumed divergent flow) and were
486 followed back toward the centre of the ice sheet until features orientated in a different direction
487 were encountered. A key feature of the LLGM ice flow pattern was the major ice stream in
488 Hudson Strait that issued from a catchment area centred over Hudson Bay and with major
489 domes over Quebec-Labrador to the south-east, Keewatin to the west and the Foxe-Baffin
490 dome to the north (see Fig. 2). A major 'Trans-Laurentide Ice Divide' extended from near
491 Victoria Island in the Canadian Arctic Archipelago south towards the Keewatin Dome and then
492 westwards to connect with the Labrador dome, with secondary ice divides emanating from
493 regional ice dispersal centres over the Queen Elizabeth Islands, Baffin Island, Newfoundland,
494 and the Appalachians (Fig. 2). Dyke and Prest (1987) also depicted several major ice streams
495 in regions where flow-lines exhibited strong convergence and they noted that some of these
496 coincided with distinctive erratic dispersal plumes (e.g. Dyke *et al.*, 1982; Dyke, 1984). Other
497 ice streams were invoked at the southern margin of the ice sheet, associated with the major ice
498 lobes that were known to have possessed very low ice surface gradients (cf. Mathews, 1974)

499 as a result of deformable bed conditions beneath the ice sheet (e.g. Boulton *et al.*, 1985; Fisher
500 *et al.*, 1985; see Section 2.4).

501 It is widely recognised that initial deglaciation from the LLGM configuration outlined
502 above was generally slow during the first part of the period known as the Late Glacial (Dyke
503 and Prest, 1987; Dyke *et al.*, 2002) and that some margins may even have been advancing to
504 their local maximum, e.g. at the far north-west (Kennedy *et al.*, 2010; Lacelle *et al.*, 2013;
505 Murton *et al.*, 2007; 2015). Dyke *et al.* (2002) pointed out that there is little evidence of regional
506 recession prior to 17 ka (14 ¹⁴C ka), with the exception of the Atlantic Provinces, the Lake
507 Michigan basin, the Mackenzie Lobe in the far north-west (cf. Harrington, 1989), and possibly
508 in Hudson Strait, following Heinrich Event 2 (H2) (Andrews *et al.*, 1998) (Fig. 8a). The
509 Atlantic provinces underwent the most dramatic retreat during this period, where deglaciation
510 was associated with a margin in deep water (Mosher *et al.*, 1989; Piper *et al.*, 1990; King, 1996;
511 Scnitker *et al.*, 2001; Shaw *et al.*, 2006), and was perhaps triggered by eustatic sea level rise
512 (Dyke, 2004).

513 Dyke (2004) also noted that the lobes of the southern margin were probably oscillating
514 during slow net recession, but had retreated more substantially by the culmination of the Erie
515 Interstadial, which is poorly dated, but which Dyke (2004) placed at around 18.8 ka (15.5 ¹⁴C
516 ka, see Barnett, 1992). It has then been suggested that several of the Great Lakes ice lobes
517 underwent a major readvance (several hundred kilometres) during the succeeding Port Bruce
518 Stadial (e.g. Erie/Huron lobe, Des Moines Lobe: Clayton and Moran, 1982; Clayton *et al.*,
519 1985) and produced the only net increase in ice extent during overall deglaciation (Dyke, 2004)
520 (see Fig. 9). It is difficult to date correlative advances elsewhere in the ice sheet (see discussion
521 in Dyke, 2004), but it is thought that they may also have taken place in Hudson Strait (Andrews
522 *et al.*, 2001) and perhaps in the Atlantic provinces (Miller *et al.*, 2001; Shaw, 2003). Elsewhere,
523 Dyke (2004) noted that an AMS date on wood from basal lake sediments in south-western

524 Alberta (Beierle and Smith, 1998) indicates that initial decoupling of Laurentide and
525 Cordilleran ice had begun by around 19 ka (15.7 ¹⁴C ka BP).

526 The fact that large-scale retreat of the LIS did not begin until around 16.8 ka (14 ¹⁴C
527 ka: Dyke *et al.*, 2002) is noteworthy because far-field sea-level records indicate that global sea
528 levels had begun to increase a few thousand years prior to that time (Clark *et al.*, 2009). Thus,
529 it has been argued that if the LIS was contributing to sea-level rise in the early Late Glacial,
530 then it must have been largely through thinning, rather than areal recession, and that this
531 thinning and drawdown may have been associated with a transition from a thick, cold-based
532 LGM ice sheet to thinner, warm-based ice sheet during early deglaciation (Marshall *et al.*,
533 2000; Marshall and Clark, 2002; Robel and Tziperman, 2016). This transition may also have
534 been manifest in a major internal flow re-organisation that may have been correlative with
535 Heinrich event 1 (H1) around 17.7 ka (14.5 ¹⁴C ka) (see Veillette *et al.*, 1999). Indeed, Dyke
536 *et al.* (2002) noted that if this reorganisation occurred, it is likely that the drawdown of central
537 ice surface would have promoted subsequent deglaciation by increasing the ELA. However,
538 Dyke and Prest (1987) noted that changes in the ice marginal configuration between the LLGM
539 and 17 ka (14 ¹⁴C ka) were insufficient to effect any permanent or substantial changes in the
540 position of the primary ice domes and divides.

541

542 *3.2. Late Glacial Interstadial: ~17.6-12.8 ka (14.5-11 ¹⁴C ka)*

543 This period includes the Bølling-Allerød warm interval, punctuated by the brief Older Dryas
544 cold event (Lowe *et al.*, 1994; Dyke, 2004). As noted by Dyke (2004), this period was
545 associated with a clear pattern of net retreat of the LIS, particularly along the southern and
546 western margins. It has also been pinpointed as a time of marked volume loss, particularly
547 between ~15 and 14.5 ka (Lambeck *et al.*, 2017). Indeed, the LIS had likely become fully

548 separated from the Cordilleran Ice Sheet (CIS) by the end of the Late Glacial Interstadial
549 (Bølling-Allerød), whilst the northern and eastern margins of the ice sheet underwent only
550 minimal recession (Fig. 8b) (Dyke and Prest, 1987; Dyke, 2004). Despite this asymmetric
551 retreat, the overall configuration of the ice sheet geometry is thought to have changed little,
552 with many of the major ice domes and divides remaining stable (Dyke and Prest, 1987).
553 However, the marked retreat of the western margin of the ice sheet is likely to have driven an
554 eastward migration of the main north-south M'Clintock Ice Divide and there were some
555 marked changes in the regional ice flow directions over the interior plains (Dyke and Prest,
556 1987; Ó Cofaigh *et al.*, 2009; Ross *et al.*, 2009). It is also noticeable that this broad time interval
557 was associated with the development of numerous glacial lakes along the western and southern
558 margin of the ice sheet (Dyke and Prest, 1987; Fig. 8b).

559 The timing of the opening of the ice-free corridor between the LIS and CIS has been
560 the subject of much debate, largely due to its importance as a potential route for the peopling
561 of North America (e.g. Dixon, 1999; Goebel *et al.*, 2008; Eriksson *et al.*, 2012; Pedersen *et al.*,
562 2016), but also in relation to the routing of meltwater from Glacial Lake Agassiz (Smith and
563 Fisher, 1993; Fisher and Smith, 1994; Fisher *et al.*, 2002; Tarasov and Peltier, 2005; Murton *et*
564 *al.*, 2010; Fisher and Lowell, 2012; Teller, 2013; see Section 3.3). The rapid collapse of the
565 saddle between the LIS and CIS has also been hypothesised as a potential source of meltwater
566 pulse 1A (Gregoire *et al.*, 2012, 2015a), although the precise contributions from North America
567 are subject to ongoing debate (Clark *et al.*, 2002; Carlson and Clark, 2012; Deschamps *et al.*,
568 2012).

569 Dyke and Prest's (1987) reconstruction showed the ice free corridor opening up ~15.6
570 ka (13 ¹⁴C ka) and with the western margin of the LIS some 200-600 km east of the Cordilleran
571 Mountains by ~12.9 ka (11 ¹⁴C ka) (Fig. 8b). This conflicts with some arguments that were
572 later put forward (e.g. Dixon, 1999) that suggested that the ice sheets must have been fully-

573 coalescent until quite late in the interstadial (~12.9 ka). Recently, Lambeck *et al.* (2017) have
574 also argued that any ice-free corridor is unlikely to have existed prior to 13 ka, using a new
575 model of glacial rebound based on relative sea level data and the tilting of glacial lake
576 shorelines. A similar conclusion was reached by Pedersen *et al.* (2016) who obtained
577 radiocarbon dates, pollen, macrofossils and metagenomic DNA from lake sediment cores along
578 the central portion of the corridor and found that it was not likely to be viable as a migration
579 route before 12.6 ka. Unfortunately, this ‘unzipping’ of the two ice sheet remains very poorly
580 dated. Dyke (2004) argued that the southern margin probably began opening around 18.2 ka
581 (~15 ¹⁴C ka) and that it is possible that it may have opened completely by ~16.3 ka (13.5 ¹⁴C
582 ka), based on radiocarbon dating of basal sediment in glacial Lake Peace (Catto *et al.*, 1996),
583 situated mid-way along the corridor from south to north. However, Dyke (2004) concluded that
584 this scenario is unlikely and that the northern end probably deglaciated later and around 14.7-
585 13.9 ka (12.5-12 ¹⁴C ka). A new analysis of available dates has attempted to constrain the
586 minimum timing of the opening of the ice-free corridor and suggests that it must have been
587 completed by 11 ka (Gowan, 2013).

588 Elsewhere, the Late Glacial Interstadial is characterised by oscillations of the ice lobes
589 at the southern margin of the ice sheet, superimposed on net recession, especially in the vicinity
590 of the Great Lakes (Dyke, 2004). Ice margin retreat has been tracked at a remarkably high
591 resolution (<100 years) in numerous glacial lake sequences that were deposited as various
592 basins became isolated (e.g. Karrow and Calkin, 1985). Many of these sequences record the
593 initiation of retreat at ~17 ka (14 ¹⁴C ka), but it was clearly punctuated by readvances of the ice
594 margin that blocked drainage routes of glacial lakes. For example, the large readvance of the
595 Lake Michigan Lobe around 13.6 to 12.3 ka (11.8 to 11.5 ¹⁴C ka) is thought to have diverted
596 water away from a westward route into the Labrador Sea and back towards the Mississippi
597 drainage basin and the Gulf of Mexico (Dyke, 2004).

598

599 *3.3. The Younger Dryas (YD): 12.9-11.7 ka (11-10 ¹⁴C ka)*

600 Ice recession during the YD was generally slow, particularly at the northern and eastern
601 margins of the ice sheet, where deglaciation mostly occurred after this period (Fig. 8)
602 (Andrews, 1973; Dyke, 2004). However, the clear asymmetry of the retreat pattern (cf.
603 Andrews, 1973) continued to drive the main M'Clintock Ice Divide (running north from the
604 Keewatin dome: Fig. 2) and the ancestral Keewatin Ice Divide eastward (Dyke and Prest,
605 1987). Indeed, Dyke and Prest (1987) also noted that the period beginning around 11.5 ka (10
606 ¹⁴C ka) marked the beginning of the demise of the main Trans Laurentide Ice Divide and
607 increased the autonomy of the regional ice dispersal centres.

608 Although it was barely mentioned in comprehensive treatments of North American
609 deglaciation prior to the late 1980s (Fulton, 1989), the Younger Dryas (YD) cold event is now
610 known (cf. Dyke, 2004) to have been characterised by a period of moraine construction and, in
611 several places, major readvances of the ice margin and ice marginal lobes (e.g. Dyke and
612 Savelle, 2000). Such readvances have been well-documented along several parts of the ice
613 sheet margin and include the large Gold Cove readvance from Labrador across Hudson Strait
614 (Miller and Kaufman, 1990) and major readvances at the north-western margin of the ice sheet
615 in the Canadian Arctic Archipelago (Dyke and Savelle, 2000). In many places, such readvances
616 were associated with major ice lobes/ice streams, such as the Cumberland Sound Ice Stream
617 on Baffin Island (Jennings *et al.*, 1996; Andrews *et al.*, 1998) and the M'Clintock Channel ice
618 stream on Victoria Island (Hodgson, 1994; Stokes *et al.*, 2009). As Dyke (2004) noted,
619 although some moraines are clearly distinguishable as of YD age, others are likely to be
620 correlative, but have not been precisely dated.

621 Any discussion of the LIS retreat during the YD warrants a mention of the drainage
622 routes of glacial Lake Agassiz, which has been implicated as causative mechanism of this
623 abrupt climatic reversal (Broecker *et al.*, 1989). Initially, glacial Lake Agassiz drained to the
624 south and into the Gulf of Mexico via the Mississippi River. The traditional model (cf. Dyke,
625 2004; Carlson and Clark, 2012) is that retreat of the Lake Superior Lobe after ~12.9 ka (Fig.
626 8b) allowed glacial Lake Agassiz to drain rapidly towards the east via the St Lawrence River
627 and into the North Atlantic Ocean (Broecker *et al.*, 1989; Dyke, 2004). This may have released
628 up to 9,500 km³ of water, which is thought to have been capable of disrupting the North
629 Atlantic's Meridional Overturning Circulation (AMOC) and instigating the YD cooling
630 (Broecker *et al.*, 1989; Dyke, 2004). Following the initial outburst, the eastward drainage is
631 thought to have continued until it was blocked by ice during the Marquette readvance, which
632 culminated around 11.5 ka (10 ¹⁴C ka) (Dyke, 2004). It is then thought that the drainage route
633 may have switched northwards via the Clearwater spillway and towards the Arctic Ocean, via
634 glacial Lake McConnell (Smith and Fisher, 1993; Fisher and Smith, 1994; Fisher *et al.*, 2002).
635 The drainage is then thought to have switched back to its original southward route until the
636 recession of ice north of Lake Superior once again opened the westward route (Teller and
637 Thorleifson, 1983). More recently, however, an alternative model has been suggested which
638 indicates that Lake Agassiz may have drained to the northwest and into the Arctic Ocean much
639 earlier than originally thought, and at the onset of the Younger Dryas (Murton *et al.*, 2010).
640 This is based on the dating of sands associated with the Mackenzie delta and upstream gravels
641 and erosional channels (Murton *et al.*, 2010), but numerical modelling has also indicated
642 increased runoff via this outlet around this time, even in the absence of any lake drainage
643 (Tarasov and Peltier, 2005). Furthermore, high-resolution ocean modelling indicates that
644 freshwater input to the Arctic Ocean is much more effective at perturbing the AMOC compared
645 to an input from the eastern drainage route (Condrón and Winsor, 2012). Field evidence for the

646 opening of the Clearwater spillway at the onset of the YD is, however, far from equivocal
647 (Fisher and Lowell, 2012) and the debate continues.

648

649 3.3. Final Deglaciation: 11.5-6 ka (10-5.2 ¹⁴C ka)

650 Final deglaciation of the LIS occurred during the early to middle Holocene (11.5 – 6.0 ka) in
651 response to increased summer insolation and increasing levels of carbon dioxide (CO₂)
652 (Carlson *et al.*, 2007; 2008; Marcott *et al.*, 2013). This warming led to the disappearance of
653 most Northern Hemisphere ice sheets, but Ullman *et al.* (2016) noted that, despite this strong
654 radiative and temperature forcing, global mean sea level (GMSL) was still around 60 m below
655 present at the start of the Holocene (Lambeck *et al.*, 2014), indicating a lag (of as much as 4
656 ka) between deglaciation of the LIS and peak insolation and CO₂ forcings (see also Section
657 4.1).

658 Retreat of the LIS was most dramatic along the northern and western margins of the ice
659 sheet. Recession of the northern margin of the ice sheet accelerated dramatically soon after
660 11.5 ka (10 ¹⁴C ka) and it is thought that the LIS and IIS had separated by 10.1 ka (9 ¹⁴C ka),
661 but that the IIS remained confluent with the Greenland Ice Sheet until 8.6 ka (7.8 ¹⁴C ka) (Fig.
662 8c, d) (England, 1999; Dyke, 2004). Dyke (2004) suggests that the IIS had fragmented by 9.5
663 ka (8.5 ¹⁴C ka) and had retreated close to modern ice margins by 8.6 ka (7.88 ¹⁴C ka). Dyke
664 (2004) also noted that the Keewatin Sector of the ice sheet had cleared the Canadian Arctic
665 Archipelago by 8.6 ka (7.8 ¹⁴C ka) (Dyke, 2004). Retreat across the mainland was also rapid,
666 but the ice sheet constructed a series of major moraine systems (e.g. the MacAlpine and
667 Chantrey moraines) that Dyke (2004) assigned to around 9.1 to 8.6 (8.2 to 7.8 ¹⁴C ka). These
668 moraines lie well inland of the marine limit and likely represent some form of readvance of the
669 ice margin, perhaps associated with ice streaming (e.g. Stokes and Clark, 2003), rather than

670 what Dyke (2004) referred to as ‘end-of-calving’ stabilisations in the Labrador Sector (see
671 below).

672 Recession of the southern margin of the ice sheet was also rapid and has been
673 reconstructed with impressive detail from tracing glacial lake shorelines to end moraines (e.g.
674 Barnett, 1992). As the ice margin retreated into an isostatically-depressed interior, numerous
675 lakes were decanted and their drainage re-routed, including a major eastward discharge of Lake
676 Agassiz at around 10.1 ka (9 ¹⁴C ka) (Dyke, 2004). These lakes may also have facilitated
677 localised readvances of the ice margin (often termed ‘surges’ or ice streaming), such as into
678 glacial Lake Ojibway (Thorleifson *et al.*, 1993) and help explain the contrasting dynamics of
679 neighbouring lobes during overall ice margin retreat (Cutler *et al.*, 2011).

680 The final evolution of lakes Agassiz and Ojibway after 8.9 ka (8 ¹⁴C ka) is more
681 speculative, but their northwards drainage into the Tyrrell Sea (ancestral Hudson Bay) is
682 evidenced by glacial bedforms, subglacial drainage channels and numerous iceberg scour-
683 marks (Josenhans and Zevenhuizen, 1990). Josenhans and Zevenhuizen (1990) argued that a
684 large calving bay opened up in western Hudson Bay and that glacial lakes Agassiz-Ojibway
685 initially drained into that region, rather than eastern Hudson Bay. The final catastrophic
686 drainage of the Agassiz-Ojibway and the full incursion of the Tyrrell Sea has been dated to
687 around 8.4 to 8.2 ka (Dyke, 2004), which is correlative with the ‘8.2 ka cold event’ seen in
688 Greenland ice cores (Alley *et al.*, 1997; Barber *et al.*, 1999; Rasmussen *et al.*, 2006). Indeed,
689 Barber *et al.* (1999) have argued that the sudden release of freshwater that accompanied this
690 event is likely to have disrupted the AMOC and lead to the abrupt but short-lived cooling seen
691 in numerous circum-North Atlantic records. Elsewhere, Dyke (2004) noted that the western
692 margin of the Québec-Labrador ice cap had stabilised at the 800-km long Sakami moraine (Fig.
693 10), most likely as a result of the sudden and large reduction of water depth that accompanied
694 by drainage of Lake Ojibway (Hardy, 1982).

695 In contrast to the rapid retreat seen elsewhere, Dyke (2004) suggests that the Baffin
696 Sector was still close to its maximum configuration at 11.5 ka (10 ¹⁴C ka) and that retreat of
697 many of its outlet glaciers proceeded slowly between 11.5 ka (10 ¹⁴C ka) and 9.5 ka (8.5 ¹⁴C
698 ka). A series of extensive moraine systems were also constructed around much of the Baffin
699 Sector between 9.5 ka (8.5 ¹⁴C ka) and 7.8 ka (7 ¹⁴C ka), which Dyke (2004) suggests may
700 reflect a mass balance that fluctuated from positive to slightly negative. That said, it is clear
701 that some major outlet glaciers retreated rapidly during this period, especially through deep
702 bathymetric troughs (Briner *et al.*, 2009), and Dyke and Prest (1987) portray the first major
703 recession of the terminus of the Hudson Strait Ice Stream at 10.1 ka (9 ¹⁴C ka).

704 Dyke (2004) argued that the final break-up of the Foxe-Baffin Sector likely involved
705 the northward progression of a calving bay from Hudson Bay between 7.8 and 6.9 ka (Fig. 8d)
706 (7 to 6 ¹⁴C ka), leaving residual ice caps on Baffin Island, Southampton Island, and Melville
707 Peninsula that remain the last major remnants of the LIS. Deglaciation of the remnant Keewatin
708 and Foxe Domes (Dyke, 2004; Ross *et al.*, 2012; Simon *et al.*, 2014) left a remnant Labrador
709 Dome that has been estimated to contain a sea level equivalent of 3.6 ± 0.4 m at ~8.2 ka (Ullman
710 *et al.*, 2016) (Fig. 10). Recently, Ullman *et al.* (2016) constrained the final retreat of this ice
711 mass using ¹⁰Be surface exposure dating and demonstrated that the ice margin may have been
712 highly sensitive to several abrupt climate events. Superimposed on overall retreat, they
713 demonstrated that the ice sheet deposited a series of moraine systems at ~10.3 ka (Paradise
714 Moraine), 9.3 ka (North Shore Moraine) and 8.2 ka (Sakami Moraine) (Fig. 10), which
715 coincided with North Atlantic cold events (Bond *et al.* 1997; Rasmussen *et al.*, 2006), and
716 which may have helped to stabilise the ice sheet. Following the widely-documented 8.2 ka
717 event (see also Alley *et al.*, 1997; Barber *et al.*, 1999) and the opening of Hudson Strait, they
718 suggest that Hudson Bay became seasonally ice-free and that the majority of the ice sheet
719 melted abruptly and within a few centuries, with deglaciation of the LIS completed by $6.7 \pm$

720 0.4 ka (Fig. 10) (cf. Carlson *et al.*, 2007; 2008). Indeed, using a Regional Climate Model,
721 Ullman *et al.* (2016) argued that the loss of ice over Hudson Bay would have been important
722 in driving negative mass balances in the surrounding ice masses, largely due to the increased
723 thermal capacity and reduced albedo of seasonally open water.

724

725 3.4. Summary

726 The pattern and timing of the LIS deglaciation is now reasonably well-known and is
727 characterised by a clear asymmetry whereby the western and southern margins retreated back
728 towards major dispersal centres over Foxe Basin and Labrador. In terms of the timing,
729 deglaciation is characterised by a period of very slow recession prior to ~17 ka, when it lost
730 <10% of its area, followed by a near-linear retreat until ~7.8 ka, when only 10% of the area
731 remained more glaciated than present (Fig. 8, 9) (Dyke, 2004). Although there were numerous
732 local-scale ice marginal fluctuations marked by rapid advance or retreat and internal flow
733 reorganisation, the two most important events that interrupted the overall linear recession were:
734 (i) the reduced rate of recession during the YD (including several well-documented
735 readvances), and (ii) the final increased rate of recession during marine incursion into Hudson
736 Bay (Dyke, 2004).

737

738

739 4. Climatic Forcing and Mechanisms of LIS Deglaciation

740 As noted in the Introduction, the deglaciation of palaeo-ice sheets represents a valuable
741 analogue for understanding the rates and mechanisms of ice sheet deglaciation. Mass loss from
742 ice sheets is complex, but can be broadly partitioned (cf. van den Broeke *et al.*, 2009) between
743 melting (mostly at the surface, but also under the ice sheet and where it meets the ocean), and

744 a ‘dynamic’ component whereby rapidly-flowing outlet glaciers transfer ice from the interior
745 to the oceans. In the Greenland Ice Sheet (GrIS), these processes are thought to be contributing
746 approximately equally to its recent negative mass balance (van den Broeke *et al.*, 2009). In
747 Antarctica, however, there is much less surface melt and dynamic changes, namely the
748 acceleration, thinning and retreat of outlet glaciers (Pritchard *et al.*, 2009), are a more important
749 influence on its mass balance, which is negative in West Antarctica but negligible or even
750 slightly positive in East Antarctica (Shepherd *et al.*, 2012). The extent to which surface mass
751 balance and ‘dynamic’ discharge (ice streaming) influenced the deglaciation of the Laurentide
752 Ice Sheet will now be discussed.

753

754 *4.1. Surface mass balance during deglaciation of the LIS*

755 In their discussion of the timing of the gLGM, Clark *et al.* (2009) argued that the primary
756 mechanism for triggering the onset of deglaciation in the Northern Hemisphere between 20 and
757 19 ka was increased insolation from orbital forcing. The efficacy of this external forcing is via
758 increased surface melt in marginal areas, particularly at the southern margin of the LIS. Clark
759 *et al.* (2009) also noted that once deglaciation had been initiated, it is likely that several
760 feedback mechanisms would have amplified the initial response (e.g. CO₂ and oceanic
761 feedbacks), including a delayed crustal rebound, which keeps the ice sheet elevation relatively
762 low and increases ablation (Abe-Ouchi *et al.*, 2013). Recent modelling by Gregoire *et al.*
763 (2015b) has attempted to partition the influence of increasing greenhouse gases (GHGs, e.g.
764 CO₂) and orbital forcing on North American deglaciation. They found that orbital forcing
765 explains around 50% of the reduction in ice volume during deglaciation, while GHGs explain
766 around 30%, but that the impact of GHGs lags orbital forcing. Orbital forcing begins around
767 23 ka and starts to impact the ice sheet from around 19 ka, but there is a delay of 3 ka before
768 CO₂ forcing has a noticeable influence from around 16 ka.

769 Recently, Ullman *et al.* (2015a) pinpointed the initial retreat of the southern margin of
770 the ice sheet in Wisconsin using a suite of ^{10}Be surface exposure ages from boulder surfaces in
771 terminal moraines. These ages dated the initial retreat of the ice margin from the LLGM
772 moraines to 23.0 ± 0.6 ka, which they noted was synchronous with several other locations along
773 the southern margin and coincided with the initial increase in summer insolation around 24-23
774 ka. They also pointed out that an acceleration in retreat after around 20.5 ka was likely driven
775 by an acceleration in boreal summer insolation and that this occurred before any increase in
776 atmospheric CO_2 , supporting an orbital forcing as the trigger for initial deglaciation (Clark *et*
777 *al.*, 2009; Gregoire *et al.*, 2015b). This response of the ice sheet to atmospheric forcing also
778 implies a higher sensitivity of land-terminating margins to small changes in climate forcing
779 than had hitherto been recognised, although it should be noted that overall recession of the ice
780 sheet was minimal (see Section 3.1). It is also interesting that whilst the southern margin was
781 beginning to retreat, there is strong evidence that the margin in the far north-west was still
782 advancing and likely attained its maximum position after 18.5 ka (e.g. Murton *et al.*, 2007;
783 Kennedy *et al.*, 2010; Lacelle *et al.*, 2013). It is not clear why the northwest margin advanced
784 to its maximum position a few millennia after the global LGM (*sensu* Clark *et al.*, 2009), but
785 Lacelle *et al.* (2013) suggested that sea-level rise and the opening of the Arctic Ocean along
786 the Beaufort Sea coastline may have provided a local source of moisture and increased
787 precipitation that enabled a advance of the LIS in this region. They also pointed out that the
788 abundance of deformable sediments in the region may also have facilitated a rapid advance of
789 the Mackenzie Lobe (ice stream) (cf. Beget, 1987).

790 Although increased atmospheric warming is thought to have triggered the initial retreat
791 of the LIS, surface energy balance modelling suggests that the ice sheet's overall net surface
792 mass balance remained positive for much of the early part of deglaciation (Ullman *et al.*,
793 2015b). Ullman *et al.* (2015b) used a surface energy balance model forced by climate data from

794 simulations with a fully coupled atmosphere-ocean General Climate Model (GCM) for key
795 time slices during the last deglaciation (24, 21, 19, 16.5, 15.5, 14, 13, 11.5 and 9 ka), see Fig.
796 11. They found that the net surface mass balance was positive until after 11.5 ka, which implied
797 that mass loss was primarily driven by dynamic discharge via calving at marine-terminating
798 ice streams (see Section 4.2). Only when summer temperatures increased by 6-7 °C (relative
799 to the gLGM) did the ice sheet's surface mass balance become increasingly negative in the
800 early Holocene. This occurred between 11.5 and 9 ka and was accompanied by an expansion
801 of the ablation area that was previously restricted to the low-gradient lobe of the southern
802 margin, but which expanded to most of the southern and western marginal areas by 9 ka (Fig.
803 11). Ullman *et al.* (2015b) noted that this time period also saw the LIS lose most of its marine
804 margin and would have coincided with a large reduction in dynamic discharge via calving
805 losses (cf. Stokes *et al.*, 2016).

806 It is worth noting, however, that the LIS had only lost around 40% of area in >10,000
807 years of deglaciation from the LLGM to around 9 ka, despite increasing boreal insolation and
808 a ~80 ppm increase in CO₂ (Ullman *et al.*, 2015b). Thus, Ullman *et al.* (2015b) noted that the
809 transition to a negative surface mass balance that occurred between ~11.5 and 9 ka, and the
810 very rapid retreat of the ice sheet after ~9 ka (two to five times faster than before ~11.5 ka),
811 suggests that some kind of instability threshold was crossed and that the final deglaciation of
812 60% of the ice sheet's area was driven by surface melt, rather than dynamic discharge. This is
813 supported by numerous studies that have shown that the rapid decay of the LIS in the early
814 Holocene was driven by enhanced boreal summer insolation and increased ablation (Carlson
815 *et al.*, 2007; 2008; 2009a). For example, Carlson *et al.* (2009a) used a surface energy balance
816 model, driven by atmosphere-ocean general circulation model at 9 ka, and calculated a net
817 surface mass balance of $-0.67 \pm 0.13 \text{ m a}^{-1}$. Given volume estimates of the LIS at this time, it

818 indicates that surface ablation accounted for $74 \pm 22\%$ of mass loss at that time, with the
819 remainder attributable to dynamic calving (Carlson *et al.*, 2009a).

820 The evolving pattern of the surface mass balance of the ice sheet (e.g. Fig. 11) partly
821 explains the clear asymmetry of retreat whereby the southern and western margins retreated
822 much more rapidly than those in the north and especially the east. Following the initial retreat
823 of the margins in the south, and the unzipping of the LIS and CIS in the east, surface-albedo
824 feedbacks are likely to have helped drive further retreat (e.g. the emergence of darker land-
825 surfaces). The formation of glacial lakes along the southern and western margins may also have
826 enhanced localised calving and ice sheet draw-down (e.g. Andrews, 1973; Cutler *et al.*, 2001;
827 Stokes and Clark, 2004). Indeed, Andrews (1973) was one of the first to point out that the
828 retreat of the LIS could not be explained solely by surface mass balance forcing and that an
829 important process during deglaciation was mass loss associated with calving in both marine
830 and lacustrine settings. However, Dyke and Prest (1987) noted that whilst calving was
831 undoubtedly an important means of ice sheet ablation, its role should not be overemphasized.
832 They pointed out that glacial lakes on the Prairies were small and that, until the formation of
833 Lake Agassiz, calving cannot account for deglaciation for most of that region prior to its
834 development. They also noted that the southern and eastern margins of the Labrador Sector had
835 some of the longest marine margins, but that these margins retreated more slowly than the
836 contemporaneous western margin of the Keewatin Sector, even though it occurred largely on
837 dry land. It is also noteworthy that the marine-based part of the LIS over Hudson Bay and Foxe
838 Basin were among the last to deglacialate, despite their susceptibility to calving and the fact that
839 their central areas were isostatically-depressed hundreds of metres below sea level (Dyke and
840 Prest, 1987). Indeed, even using an extreme ‘calving instability’, Marshall *et al.* (2000) were
841 unable to evacuate a significant volume of ice from Hudson Strait in their modelling
842 experiments, although more recent modelling (Bassis *et al.*, 2017) suggests that this ice stream

843 may have been particularly vulnerable to calving triggered by subsurface ocean warming (see
844 Section 4.2).

845 A further explanation for the asymmetric retreat may also relate to the release and
846 routing of meltwater during deglaciation and its subsequent impact on ocean circulation
847 (Carlson *et al.*, 2009b; Hoffman *et al.*, 2012; Jennings *et al.*, 2015; Gregoire *et al.*, 2015b).
848 Climate modelling indicates that meltwater discharge events routed into the Labrador Sea (e.g.
849 from meltwater runoff and glacial lakes) could cause a cooling of up to 1.5 °C over the Labrador
850 Dome (Morrill *et al.*, 2014), but with minimal cooling (<0.5°C) along the western margin of
851 the ice sheet, west of Hudson Bay. This negative feedback mechanism has thus been invoked
852 to explain the relatively stability of the Labrador Dome, whilst the western margin of the ice
853 sheet continued to retreat (Ullman *et al.*, 2016).

854

855 *4.2. The role of ice streaming during deglaciation of the LIS*

856 Ice streams are the key drainage routes of an ice sheet (Bamber *et al.*, 2000) and are known to
857 exert a considerable influence on ice sheet configuration and mass balance (Nick *et al.*, 2013;
858 Ritz *et al.*, 2015). It has been recognised for some time, therefore, that accurate reconstructions
859 of the LIS require a detailed knowledge of the location of ice streams. The first use of the term
860 ‘ice stream’ in relation to the LIS was by Bell (1895: p. 352-353), who inferred the presence
861 of a “*great ice stream*” passing through Hudson Strait (see Brookes, 2007). However, it was
862 not until 1981 that Denton and Hughes (1981) attempted to incorporate ice streams into a
863 reconstruction of the entire LIS. They were the first to portray an extensive network of ice
864 streams in the Northern Hemisphere ice sheets (Fig. 12a), which was clearly influenced by
865 Hughes’ knowledge of West Antarctic ice streams (e.g. Hughes, 1977). It is not clear what
866 evidence was used to locate the ice streams, but several major ice streams were depicted in

867 large topographic troughs, such as Hudson Strait, and others were depicted on low relief
868 terrains where they appeared as a regularly-spaced network, perhaps hinting at some notion of
869 spatial organisation. Dyke and Prest (1987) also incorporated ice stream flow-lines in their
870 reconstruction of the LIS, some of which had earlier been recognised from erratic dispersal
871 trains (Dyke *et al.*, 1982; Dyke, 1984). Around that time, Dyke and Morris (1988) published a
872 classic paper that would be one of the first to describe, in detail, the geomorphological
873 ‘footprint’ of an ice stream on Prince of Wales Island in the Canadian Arctic Archipelago. They
874 reported evidence for a convergent flow pattern of highly elongate drumlins associated with an
875 erratic dispersal train with abrupt lateral margins.

876 Many of the lobes of the southern margin were also attracting attention as possible
877 zones of ice streaming (Dredge and Cowan, 1989; Alley, 1991), largely because of their low
878 surface slopes (Mathews, 1974) and earlier suppositions about the possibility of rapid flow in
879 the form of surging (Wright, 1973; Clayton *et al.*, 1985). Hicock and co-workers reported
880 dispersal trains and tills associated with drumlins around the Great Lakes region that were
881 interpreted to reflect ice streaming (Hicock, 1988, 1992; Hicock and Dreimanis, 1992). Later
882 work by Patterson (1997; 1998) recognised entire landform assemblages that were interpreted
883 to result from ice streaming and which included suites of level-to-streamlined fine-grained till,
884 sometimes associated with highly elongate drumlins (Bluemle *et al.*, 1993) that typically
885 terminated towards the lobate margins, where thrusting of glacial sediment was evident in
886 association with hummocky topography and major moraine systems. Similar, albeit much
887 larger, patterns of streamlining were also reported by Clark (1993). Using Landsat satellite
888 imagery, he identified a “*hitherto undocumented and much larger form of ice moulded*
889 *landscape*” (p. 1) which comprised streamlined glacial lineations with typical lengths of
890 between 8 and 70 km, widths between 200 and 1300 m, and spacings between 300 and 5 km.
891 Clark (1993) termed these features ‘mega-scale glacial lineations’ (MSGs) and discussed a

892 variety of possible origins, concluding that were likely to form under conditions of extremely
893 rapid flow such as ice streams or surges.

894 Thus, by the late 1990s, significant progress had been made in terms of identifying the
895 glacial geological evidence of ice streaming on the bed of the LIS. These studies suggested that
896 ice streaming should leave behind sedimentological evidence of fast ice flow in the form of
897 heavily deformed tills and distinctive erratic dispersal trains that often depicted convergent
898 flow-patterns (e.g. Dyke and Morris, 1988; Hicock, 1988; Alley, 1991; Patterson, 1997; 1998).
899 Many of these flow-patterns, or fans (cf. Kleman and Borgström, 1996), also contained highly
900 elongate glacial lineations, which were postulated to reflect rapid ice velocities (e.g. Clark,
901 1993); and some were characterised by abrupt lateral margins (e.g. Hodgson, 1994) and lateral
902 shear margin moraines (Dyke and Morris, 1988). Taken together, these were argued to
903 represent the key ‘geomorphological criteria’ for identifying palaeo-ice streams, which Stokes
904 and Clark (1999) formalised in a series of landsystems models depending on whether the ice
905 stream terminated in water or on land, and whether the glacial lineations were formed rapidly
906 and synchronously or slowly and time-transgressively.

907 Despite much progress, however, the first systematic literature review of ice streams in
908 the LIS (Stokes and Clark, 2001) found only 10 hypothesised ice streams that had been
909 identified based on unambiguous glacial geological evidence (Fig. 12b). However, a large
910 number of ice streams were uncovered in the early 2000s (e.g. Clark and Stokes, 2001; Shaw,
911 2003), such that by 2004, a new map of ice streams in the LIS depicted a total of 49 ice streams,
912 34 of which had good evidence, with the remainder more uncertain (Winsborrow *et al.*, 2004).
913 Much of the evidence was based on the terrestrial glacial geological record, but the recognition
914 of discrete layers of ice rafted debris (IRD) in North Atlantic sediment cores (Heinrich, 1988)
915 had also begun to implicate episodic ice streaming, particularly in Hudson Strait, as being
916 primarily responsible for their deposition (Bond *et al.*, 1992; Andrews and Tedesco, 1992;

917 MacAyeal, 1993; Marshall and Clark, 1997a, b; Andrews, 1998). In addition, the burgeoning
918 growth of marine geophysical techniques saw a large number of ice stream footprints identified
919 in offshore settings, particularly in Atlantic Canada (Shaw, 2003; Shaw *et al.*, 2006; Todd *et*
920 *al.*, 2007; Shaw *et al.*, 2009), but also in the Canadian Arctic Archipelago (MacLean *et al.*,
921 2010) and in Hudson Bay (Ross *et al.*, 2011). These techniques, allied with the growth of
922 remote sensing studies across large regions of the ice sheet bed enabled a large number of ice
923 streams to be identified (e.g. De Angelis and Kleman, 2005, 2007; Evans *et al.*, 2008; Ross *et*
924 *al.*, 2009; Stokes *et al.*, 2009; Ó Cofaigh *et al.*, 2010).

925 Most recently, Margold *et al.* (2015a) compiled a new inventory of ice streams in the
926 LIS based on an up-to-date review of the literature and systematic mapping from across the
927 entire ice sheet bed using both terrestrial and offshore datasets. Their map (see also Fig. 4)
928 includes 117 ice streams and each ice stream is categorised according to the type of evidence
929 it left behind, with an acknowledgement that some locations are more speculative than others.
930 Indeed, identifying ice streams on more resistant ‘hard-bed’ terrain, such as the Canadian
931 Shield, is more difficult, but recent work (e.g. Eyles 2012; Eyles and Putkinen, 2014;
932 Krabbendam *et al.*, 2016) has described rock drumlins, megaflutes and mega-lineated terrain,
933 which likely represent a hard-bedded landform assemblage cut by ice streams. Thus, it is
934 unlikely that any major ice streams are missing (Margold *et al.*, 2015a, b). Indeed, most of the
935 major ice streams are also captured in numerical modelling of the ice sheet, although the
936 dynamics of land-terminating ice streams are much harder to reproduce (Stokes and Tarasov,
937 2010).

938 In a review of the spatial distribution of Laurentide ice streams, Margold *et al.* (2015b)
939 noted that the pattern of ice streams (Fig. 4) during the LLGM resembled the present day
940 velocity patterns in modern ice sheets (Fig. 13a). They estimated that around a third of the LIS
941 margin perimeter was drained (intersected) by ice streams at the LLGM, which is a very similar

942 value for the present-day Antarctic ice sheets. Large ice streams had extensive onset zones fed
943 by multiple tributaries and, where ice drained through regions of high relief, the spacing of ice
944 streams appears to show a degree of spatial self-organisation which was hinted at in the earlier
945 work by Denton and Hughes (1981), but which has perhaps not been fully appreciated and
946 explored. It is also clear that whilst topography exerted a primary control on fixing the location
947 of ice many streams in the LIS, there were large areas along the western and southern margin
948 of the ice sheet where networks of ice streams operated over soft sediments and switched
949 direction repeatedly and probably over short (centennial) time scales (cf. Ó Cofaigh *et al.*,
950 2009; Ross *et al.*, 2009). As the ice sheet retreated on to its low relief interior, however,
951 Margold *et al.* (2015a, b) noted that several ice streams showed no correspondence with
952 topography or underlying geology, and were perhaps facilitated by localised build-up of
953 pressurised subglacial meltwater (e.g. Stokes and Clark, 2003). Margold *et al.* (2015b) also
954 highlighted that there have been very few attempts to date the initiation and cessation of the
955 vast majority of ice streams, but that it is clear that they must have switched on and off during
956 deglaciation, rather than maintaining the same trajectory as the ice margin retreated.

957 The extent to which changes in the ice stream drainage network were a cause or effect
958 of ice sheet deglaciation is a key question. Put another way, does the drainage network of ice
959 streams (Fig. 4) arise as a result of climatically-driven changes in ice sheet mass balance or
960 could ice streams evolve to drive changes beyond that which might be expected from climate
961 forcing alone? This question has rarely been addressed with respect to the LIS, but it is clear
962 that ice streaming led to major reorganisations in the flow pattern of the ice sheet (Mooers *et al.*
963 *et al.*, 1997; Veillette *et al.*, 1999; Ó Cofaigh *et al.*, 2009; Ross *et al.*, 2009; Stokes *et al.*, 2009).
964 It is also clear that ice streaming is capable of rapidly lowering the ice sheet surface profile to
965 lower elevations, where ablation will be increased (Robel and Tziperman, 2016). However,
966 even where major reorganisations took place as a result of ice streaming, there is little evidence

967 that deglaciation proceeded more rapidly. For example, if the Hudson Strait Ice Stream was
968 responsible for major ice discharge events into the North Atlantic (e.g. Andrews and Tedesco,
969 1992; Andrews, 1998; Andrews and MacLean, 2003), there is tentative evidence for a
970 reorganisation of the internal geometry and flow patterns of the ice sheet (Dyke and Prest,
971 1987; Mooers *et al.*, 1997; Veillette *et al.*, 1999; Dyke, 2004), but little evidence that these
972 events resulted in a more rapid deglaciation of the area of the ice sheet, which was mostly linear
973 through time (Fig. 9). The only exception to this is where individual outlet glaciers retreated
974 through deep troughs that resulted in a localised acceleration in retreat until the margin was
975 able to re-stabilise on higher ground (Briner *et al.*, 2009). It is also possible that eustatic sea-
976 level rise early in deglaciation may have increased water depths close to marine-terminating
977 ice streams, enhancing their discharge and leading to rapid draw-down of ice. As noted above,
978 this might partly explain the relatively early deglaciation of the Atlantic provinces, where
979 deglaciation was associated with a margin in deep water (Andrews, 1973; Mosher *et al.*, 1989;
980 Piper *et al.*, 1990; King, 1996; Scnitker *et al.*, 2001; Shaw *et al.*, 2006), but it appears to have
981 had less effect at the marine margin in the Canadian Arctic Archipelago.

982 In a recent study, Stokes *et al.* (2016) examined whether the cumulative impact of ice
983 streams was able to increase and sustain rates of mass loss during deglaciation of the LIS
984 beyond those that might be expected from climate forcing alone. They used the Dyke *et al.*
985 (2003) ice margin chronology to bracket the duration of the 117 ice streams in the inventory
986 from Margold *et al.* (2015a: see Fig. 4). They found that as the ice sheet retreated, ice streams
987 activated and deactivated in different locations (see Fig. 13) and, unsurprisingly, their overall
988 number decreased. Perhaps more surprising was that ice streams occupied a progressively
989 smaller percentage of the ice sheet perimeter during deglaciation. At its maximum,
990 approximately 27% of the LIS margin was streaming, but this value decreased to between 25%
991 and 20% from 16 ka to 13 ka, and then rapidly dropped to ~5% at 11 ka (Stokes *et al.*, 2016).

992 This implies that the final 4 to 5 ka of deglaciation was largely driven by surface melt, which
993 is corroborated surface mass balance modelling (see Section 4.1) and inferences based on the
994 density of subglacial meltwater conduits (eskers) (Storrar *et al.*, 2014). Stokes *et al.* (2016) also
995 used a simple scaling relationship based on the width and discharge of modern ice streams to
996 estimate the potential cumulative discharge from Laurentide ice streams through time, and
997 found that this decreased and was strongly scaled to the ice sheet's volume. This scaling is also
998 found in numerically modelled estimates of ice stream discharge (Stokes *et al.*, 2012). They
999 concluded that whilst the underlying geology and topography clearly influenced ice stream
1000 activity (cf. Marshall *et al.*, 1996), the drainage network of ice streams – at the ice sheet scale
1001 – appears to have adjusted in response to ice sheet volume. Thus, contrary to the view that sees
1002 ice streams as unstable entities that can accelerate ice-sheet deglaciation, Stokes *et al.* (2016)
1003 found that ice streams exerted progressively less influence on deglaciation of the LIS.

1004 This is not to say, however, that ice streams did not play any role in deglaciation. They
1005 were likely to be very important in reducing the volume (if not the area) of the ice sheet in early
1006 deglaciation, when large parts of the LIS had a marine margin (Andrews, 1973), which is
1007 known to be a key control on ice streaming (Winsborrow *et al.*, 2010). Tanner (1965) was one
1008 of the first to recognise the importance of glacial isostatic depression in generating relatively
1009 higher sea levels at the marine margin when the ice sheet was at near-maximum configurations.
1010 This would have increased calving losses and encouraged ice stream draw-down. More
1011 recently, numerical modelling has shown that a simple feedback between ocean forcing and
1012 isostatic adjustment can explain the observed magnitude and timing of Heinrich events from
1013 the Hudson Strait Ice Stream (Bassis *et al.*, 2017). Bassis *et al.* (2017) showed that when the
1014 LIS is at its near-maximum extent, the terminus of the ice stream remains grounded on bed
1015 topography depressed about 300 m below sea level, rendering it particularly vulnerable to sub-
1016 surface ocean warming. They argued that a small warming in the subsurface ocean is enough

1017 to trigger rapid retreat of the ice sheet into the over-deepened bed (generating a Heinrich event)
1018 and that retreat continues until isostatic adjustment allows the bed to uplift, isolating the
1019 terminus from ocean forcing. At this point, they noted that retreat ceases and, with the ice sheet
1020 at its minimum extent, bed uplift facilitates regrowth on a slower timescale than collapse.

1021 Other numerical modelling experiments suggest that ice streaming may render large ice
1022 sheet more sensitive to Milankovitch forcing (Marshall and Clark, 2002; Robel and Tziperman,
1023 2016). Large ice sheets are likely to have a larger proportion of their bed above the pressure
1024 melting point and this is a first order control on the likelihood of generating ice streams
1025 (MacAyeal, 1993; Marshall and Clarke, 1997a, b; Stokes *et al.*, 2012; Marshall and Clark, 200;
1026 Tarasov *et al.*, 2012; Robel and Tziperman, 2016). In their numerical modelling experiments,
1027 Robel and Tziperman (2016) found that when ice streams are sufficiently developed, an upward
1028 shift in the ELA caused by external climatic (Milankovitch) forcing results in rapid
1029 deglaciation. However, when the same shift in the ELA was applied to an ice sheet without
1030 fully-formed ice streams, it led to continued ice sheet growth or slower deglaciation. These
1031 idealised experiments were also repeated using key aspects of the climatic and geographic
1032 complexity of the LIS and generated similar results: enhanced discharge caused by ice stream
1033 acceleration is the primary source of mass loss during the early part of deglaciation in response
1034 to orbital forcing. An interesting corollary is that these processes also explain why
1035 Milankovitch forcing late in a 100 ka glacial cycle leads to full deglaciation, when the ice sheet
1036 is large, isostatically-depressed, and has developed numerous large ice streams; while the same
1037 forcing does not produce deglaciation early in the glacial cycle when the ice sheet is small and
1038 without ice streams (Marshall and Clark, 2002; Abe-Ouchi *et al.*, 2013; Robel and Tziperman,
1039 2016).

1040
1041

1042 **5. Conclusions and Outlook**

1043 The LIS is thought to have initiated from an ice-free state over North America during MIS 5d,
1044 around 116-114 ka (e.g. Marshall *et al.*, 2000; Stokes *et al.*, 2012). It grew rapidly from its
1045 initial inception over the Arctic/subarctic plateaux along the eastern seaboard of Canada and
1046 likely attained an MIS 4 maximum around 65-60 ka (Marshall *et al.*, 2000; Stokes *et al.*, 2012).
1047 Its extent during MIS 3 is uncertain (Dredge and Thorleifson, 1987; Stokes *et al.*, 2012; Dalton
1048 *et al.*, 2016), but it grew rapidly to its Local Last Glacial Maximum, which was attained around
1049 26-25 ka (e.g. Dyke, 2002; Clark *et al.*, 2009), although with some regions advancing much
1050 later, such as in the far north-west (Lacelle *et al.*, 2013). After over a century of debate, a
1051 consensus has emerged that it existed as an extensive, multi-domed ice sheet that extended to
1052 the edge of the continental shelf at its marine margins, but that it was thinner (~3,000 m) than
1053 some earlier work had suggested and consumed a sea-level equivalent of around 50 m (Clark
1054 *et al.*, 1996). It is thought that it existed at or close to this maximal configuration for several
1055 thousand years (Dyke *et al.*, 2002).

1056 Our understanding of the pattern and timing of deglaciation is due in no small part to
1057 several major syntheses that have remained benchmark reconstructions for several decades
1058 (e.g. Bryson *et al.*, 1969; Denton and Hughes, 1981; Boulton *et al.*, 1995; Dyke and Prest,
1059 1987; Dyke, 2004), augmented by numerical modelling, which has seen rapid developments
1060 over the last two decades (Marshall *et al.*, 1996; Marshall *et al.*, 2000; Tarasov *et al.*, 2012;
1061 Peltier *et al.*, 2015, Gregoire *et al.*, 2015b). This body of work shows a clear asymmetry in
1062 retreat whereby the western and southern margins retreated back towards the major dispersal
1063 centres over Foxe Basin-Baffin Island and Quebec-Labrador. Ice margin retreat was relatively
1064 slow prior to ~17 ka, but it is clear that the ice sheet volume was decreasing. Between around
1065 16 and 13 ka, however, the margin retreated rapidly, particularly along the southern and
1066 western margins, which led to the separation of the Laurentide from the Cordilleran Ice Sheet.

1067 In contrast, the northern and eastern margins of the ice sheet underwent only minimal recession
1068 (e.g. Dyke and Prest, 1987; Dyke, 2004). During the Younger Dryas, the overall net recession
1069 was reduced and several notable readvances are known to have taken place (e.g. Dyke and
1070 Savelle, 2000; Jennings *et al.*, 1996; Andrews *et al.*, 1998). Following the Younger Dryas, the
1071 ice sheet retreated two to five times faster than previous rates (Ullman *et al.*, 2015b). Recession
1072 of the northern and eastern margins accompanied the continued rapid recession of the southern
1073 and western margins, although a series of moraine systems were built in some locations, which
1074 likely indicate temporary stabilisations (Dyke, 2004) or surges/ice streaming (Stokes and
1075 Clark, 2003). Final deglaciation of the remnant Keewatin and Foxe Domes (Dyke, 2004; Ross
1076 *et al.*, 2012; Simon *et al.*, 2014), left a remnant Labrador Dome that is thought to have
1077 deglaciated by around 6.7 ka (e.g. Ullman *et al.*, 2016).

1078 The pattern and timing of deglaciation of the LIS represents a valuable analogue for
1079 understanding the rates and mechanisms of ice sheet deglaciation (Denton and Hughes, 1981;
1080 Kleman and Applegate, 2014; Margold *et al.*, 2015b; Stokes *et al.*, 2016), which may be
1081 relevant to assessments of the future stability of modern-day ice sheets. In this context, it is
1082 generally accepted that the initial trigger for deglaciation was an increase in boreal summer
1083 insolation (Clark *et al.*, 2009; Ullman *et al.*, 2015a). However, modelling of the ice sheet's net
1084 surface mass balance indicates that it remained positive until around 11 ka (Ullman *et al.*,
1085 2015b). This suggests that the predominant source of mass loss was initially via rapidly-
1086 flowing ice streams, particularly at the ice sheet's marine margins (Andrews, 1973; Shaw *et*
1087 *al.*, 2006; De Angelis and Kleman, 2007; Stokes *et al.*, 2016). Indeed, our understanding of
1088 palaeo-ice streams has grown from almost completely ignorance in the early 1980s to the latest
1089 inventory of 117 ice streams that operated at various times during deglaciation (Margold *et al.*,
1090 2015a, b). Only when summer temperatures increased by 6-7 °C relative to the LLGM did the
1091 ice sheet's surface mass balance become increasingly negative in the early Holocene (around

1092 11.5 to 9 ka). Thereafter, deglaciation of the remaining 60% of the ice sheet's initial area was
1093 accomplished mostly by surface melt (Dyke, 2004; Carlson *et al.*, 2007, 2008, 2009a). This
1094 implies that 'dynamic discharge' via ice streams exerted progressively less influence on the
1095 deglaciation of the LIS (Stokes *et al.*, 2016).

1096 In his most recent synthesis, Dyke (2004) noted that the improved age control and more
1097 detailed mapping of deglacial patterns over the last few decades have enabled improved
1098 reconstructions of the LIS and a closer correlation between the deglaciation sequence and major
1099 climatic events recognised in the North Atlantic region and in the Greenland ice core record.
1100 That said, there remain areas of the ice sheet where deglaciation is relatively poorly
1101 constrained. The western margin of the LIS (and western interior) has emerged as a key area
1102 for a number of important debates, including the concept of an ice-free corridor and the
1103 peopling of North America (Dyke, 2004; Dixon, 2013; Pedersen *et al.*, 2016). The possibility
1104 of a rapid saddle-collapse and a large meltwater pulse (and sea-level jump) have also implicated
1105 this region (Gregoire *et al.*, 2012, 2015a), which has also been identified as a drainage route
1106 for a putative outburst flood from Glacial Lake Agassiz (Tarasov and Peltier, 2005; Murton *et*
1107 *al.*, 2010; Fisher and Lowell, 2012). It is interesting to note that, with admirable foresight, this
1108 region was specifically highlighted almost 50 years ago by Bryson *et al.* (1969: p. 5) who stated
1109 that "*it is unfortunate that there are so few available dates from the western interior; the*
1110 *corridor between the Cordillera and the retreating ice front is of great interest to the*
1111 *anthropologist, biologist, and climatologist*". It is perhaps more unfortunate that, almost five
1112 decades on, the scarcity of dates from this region persists, and this is a key area for future
1113 research to address. Improvements in the ice margin chronology and the quantification of its
1114 uncertainties will also provide tighter constraints for numerical modelling of the ice sheet,
1115 which will also require improvements in terms of model resolution and in the representation of
1116 the key physics, such as the simulation of ice streams, subglacial processes and changes in

1117 meltwater drainage routes (Marshall *et al.*, 2000; Hindmarsh, 2009; Stokes *et al.*, 2015;
1118 Kirchner *et al.*, 2016; Wickert, 2016). This should help resolve important debates about the
1119 role of the LIS during major reorganisations of the ocean-climate system (Broecker *et al.*, 1989;
1120 Bond *et al.*, 1992; Barber *et al.*, 1999; Carlson and Clark, 2012), and enable improved
1121 predictions of the response of modern-day ice sheets to future climate change.

1122

1123

1124 **Acknowledgements:**

1125 The writing of this paper was made considerably easier (and even more enjoyable) as a result
1126 of the remarkable quality and meticulous nature of previous reconstructions of the LIS and
1127 major review papers, most notably involving John Andrews, Art Dyke, Terry Hughes, Jack
1128 Ives, and Vic Prest. I would like to thank Martin Margold for discussions and comments on an
1129 earlier version of this manuscript, and my colleagues David Evans and Colm Ó Cofaigh for
1130 useful advice on the glacial history of this ice sheet. I am also grateful to John Andrews and an
1131 anonymous reviewer for their encouraging comments during the peer review process. My
1132 interest in the LIS was ignited in undergraduate lectures by Chris Clark, under whose tutelage
1133 I also conducted my Ph.D. I am grateful for his guidance and infectious enthusiasm about this
1134 giant amongst the ‘Last Great Ice Sheets’. I would also like to acknowledge Chris Orton,
1135 Department of Geography, Durham University, who re-drew many of the Figures for this
1136 paper.

1137

1138

1139 **References**

1140

- 1141 Abe-Ouchi, A., Saito, F., Kawamura, K., raymo, M.E., Okuno, J., Takahashi, K., Blatter, H.,
 1142 2013. Insolation-driven 100,000-year glacial cycles and hysteresis of ice-sheet
 1143 volume. *Nature*, 500, 190-194.
- 1144 Abe-Ouchi, A., Saito, F., Kageyama, M., Braconnot, P., Harrison, S.P., Lambek, K., Otto-
 1145 Bliesner, B.L., Peltier, W.R., Tarasov, L., Peterschmitt, J-Y., Takahashi, K., 2015.
 1146 Ice-configuration in the CMIP5/PMIP3 Last Glacial Maximum experiments. *Geosci.*
 1147 *Model Dev.*, 8, 3621-3637.
- 1148 Allard, G., Roy, M., Ghaleb, B., Richard, P.J.H., Larouche, A.C., Veillette, J.J., Parent, M.,
 1149 2012. Constraining the age of the last interglacial-glacial transition in the Hudson Bay
 1150 lowlands (Canada) using UeTh dating of buried wood. *Quat. Geochronol.* 7, 37-47.
- 1151 Alley, R.B., 1991. Deforming-bed origin for southern Laurentide till sheets? *J. Glaciol.* 37
 1152 (125), 67–76.
- 1153 Alley RB, Blankenship DD, Bentley CR, Rooney ST., 1986. Deformation of till beneath ice
 1154 stream B, West Antarctica. *Nature*, 322, 57-59.
- 1155 Alley, R.B., Mayewski, P.A., Sowers, T., Stuiver, M., Taylor, K.C., Clark, P.U., 1997.
 1156 Holocene climatic instability: a prominent, widespread event 8200 yr ago. *Geology*
 1157 25, 483-486.
- 1158 Amos, C.L., Knoll, R., 1987. The Quaternary sediments of Banquereau, Scotian Shelf.
 1159 *Geological Society of America Bulletin*, 99, 244–260.
- 1160 Amos, C.L., Miller, A.A.L., 1990. The Quaternary stratigraphy of southwest Sable Island
 1161 Bank, eastern Canada. *Geological Society of America Bulletin*, 102, 915–934.
- 1162 Andrews, J. T., 1969. The pattern and interpretation of restrained, post-glacial and residual
 1163 rebound in the area of Hudson Bay. In, Hood, P.J. (Ed) Earth Science Symposium on
 1164 Hudson Bay. *Geological Survey of Canada Paper* 68-53, p. 49.
- 1165 Andrews, J.T., 1973. The Wisconsin Laurentide Ice Sheet: dispersal centres, problems of
 1166 rates of retreat, and climatic implications. *Arctic and Alpine Research*, 5 (3), 185-199.
- 1167 Andrews, J.T., 1998. Abrupt changes (Heinrich events) in late Quaternary North Atlantic
 1168 marine environments: a history and review of data and concepts. *J. Quat. Sci.*, 13 (1),
 1169 3–16.
- 1170 Andrews, J.T., 2006. The Laurentide Ice Sheet: a review of history and processes. In, Knight,
 1171 P.G. (Ed) *Glacier Science and Environmental Change*. Blackwell Publishing, UK, p.
 1172 201-207.
- 1173 Andrews, J.T., Mahaffy, M.A., 1976. Growth rates of the Laurentide Ice Sheet and sea level
 1174 lowering (with emphasis on the 115,000 B.P. sea level low). *Quat. Res.*, 6, 167-183.
- 1175 Andrews, J.T., Miller, G.H., 1972. Quaternary history of northern Cumberland Peninsula,
 1176 Baffin Island, N.W.T., Canada. Part IV: maps of present glaciation limits and lowest
 1177 equilibrium line altitude for north and south Baffin Island. *Arctic and Alpine*
 1178 *Research*, 4, 45-59.
- 1179 Andrews, J.T., Tedesco, K., 1992. Detrital carbonate-rich sediments, northwestern Labrador
 1180 Sea: implications for ice-sheet dynamics and iceberg rafting (Heinrich) events in the
 1181 North Atlantic. *Geology*, 20 (12), 1087–1090.
- 1182 Andrews, J.T., MacLean, B., 2003. Hudson Strait ice streams: a review of stratigraphy,
 1183 chronology and links with North Atlantic Heinrich events. *Boreas* 32, 4-17
- 1184 Andrews, J.T., Kirby, M.E., Jennings, A.E., Barber, D.C., 1998. Late Quaternary stratigraphy,
 1185 chronology, and depositional processes on the SE Baffin Island slope, detrital
 1186 carbonate and Heinrich events: implications for onshore glacial history. *Géographie*
 1187 *physique et Quaternaire*, 52, 91-105.
- 1188 Andrews, J.T., Jennings, A.E., MacLean, B., 2001. Late Quaternary stratigraphy of Hatton
 1189 and Resolution basins, east of Hudson Strait. In, MacLean, B. (Ed.), *Marine Geology*
 1190 *of Hudson Strait and Ungava Bay, Eastern Arctic Canada: Late Quaternary*

- 1191 *Sediments, Depositional Environments, and Late Glacial-Deglacial History Derived*
 1192 *from Marine and Terrestrial Studies*. Geological Survey of Canada Bulletin, 566, 57-
 1193 64.
- 1194 Anonymous., 1970. Last throes of Laurentide Ice Sheet. *Nature*, 225 (5229), p. 224.
- 1195 Argus, D.F., Peltier, W.R., 2010. Constraining models of postglacial rebound using space
 1196 geodesy: a detailed assessment of model ICE-5G (VM2) and its relatives. *Geophysical*
 1197 *Journal International*, 181, 697–723.
- 1198 Bamber, J.L., Vaughan, D.G., Joughin, I., 2000. Widespread complex flow in the interior of
 1199 the Antarctic Ice Sheet. *Science*, 287 (5456), 1248–1250.
- 1200 Barber, D.C., Dyke, A., Hillaire-Marcel, C., Jennings, A.E., Andrews, J.T., Kerwin, M.W.,
 1201 Bilodeau, G., McNeely, R., Southon, J., Morehead, M.D., Gagnon, J.-M., 1999.
 1202 Forcing of the cold event of 8,200 years ago by catastrophic drainage of Laurentide
 1203 lakes. *Nature* 400, 344-348.
- 1204 Barnett, P.J., 1992. Quaternary geology of Ontario. In: Thurston, P.C., Williams, H.R.,
 1205 Sutcliffe, R.H., Scott, G.M. (Eds.), *Geology of Ontario*. Toronto, Ontario Geological
 1206 Survey, Special Volume 4, 1011-1088.
- 1207 Bassis, J.N., Petersen, S.V., Cathles, L.M., 2017. Heinrich events triggered by ocean forcing
 1208 and modulated by isostatic adjustment. *Nature*, 542, 332-334.
- 1209 Beierle, B., Smith, D.G., 1998. Severe drought in the early Holocene (10,000-6800 BP)
 1210 interpreted from lake sediment cores, southwestern Alberta. *Palaeogeography,*
 1211 *Palaeoclimatology, Palaeoecology*, 140, 75-83.
- 1212 Beget, J., 1987. Low profile of the northwest Laurentide map sheets. *Arctic and Alpine*
 1213 *Research* 19, 81–87.
- 1214 Bell, R., 1884. Observations on geology, minerology, zoology and botany of the Labrador
 1215 coast, Hudson's Strait and Bay. *Canada, Geological Survey, Report of Progress,*
 1216 1882-83-84, Part DD.
- 1217 Bell, R., 1895. The Labrador Peninsula. *Scot. Geogr. Mag.*, 11 (7), 335–361.
- 1218 Blake Jr., W., 1970. Studies of glacial history in arctic Canada. I. Pumice, radiocarbon dates,
 1219 and differential postglacial uplift in the eastern Queen Elizabeth Islands. *Canadian*
 1220 *Journal of Earth Sciences*, 7, 634–664.
- 1221 Blake Jr., W., 1992. Shell-bearing till along Smith Sound, Ellesmere Island–Greenland: age
 1222 and significance. *Sveriges Geologiska Undersökning, Series Ca*, 81, 51–58.
- 1223 Blake Jr., W., 1993. Holocene emergence along the Ellesmere Island coasts of northernmost
 1224 Baffin Bay. *Norsk Geologisk Tidsskrift*, 73, 147–160.
- 1225 Bluemle JP, Lord LM, Hunke NT., 1993. Exceptionally long, narrow drumlins formed in
 1226 subglacial cavities, North Dakota. *Boreas*, 22, 15-24.
- 1227 Bond, G., Heinrich, H., Broecker, W., Labeyrie, L., Mcmanus, J., Andrews, J., Huon, S.,
 1228 Jantschik, R., Clasen, S., Simet, C., Tedesco, K., Klas, M., Bonani, G., Ivy, S., 1992.
 1229 Evidence for massive discharges of icebergs into the North-Atlantic ocean during the
 1230 last glacial period. *Nature*, 360 (6401), 245–249.
- 1231 Bond, G., Showers, W., Cheseby, M., Lotti, R., Almasi, P., deMenocal, P., Priore, P., Cullen,
 1232 H., Hajda, I., Bonani, G., 1997. A pervasive millennial-scale cycle in North Atlantic
 1233 Holocene and glacial climates. *Science* 278, 1257-1266.
- 1234 Bonifay, D., Piper, D.J.W., 1988. Probable Late Wisconsinan ice margin on the upper
 1235 continental slope off St. Pierre Bank, eastern Canada. *Canadian Journal of Earth*
 1236 *Sciences*, 25, 853–865.
- 1237 Boulton, G.S. and Hindmarsh, R.C.A. (1987) Sediment deformation beneath glaciers:
 1238 Rheology and glaciological consequences. *Journal of Geophysical Research*, 92,
 1239 9059-9082.

- 1240 Boulton, G.S., Clark, C.D., 1990a. A highly mobile Laurentide Ice Sheet revealed by satellite
1241 images of glacial lineations. *Nature*, 346, 813e817.
- 1242 Boulton, G.S., Clark, C.D., 1990b. The Laurentide Ice Sheet through the last glacial cycle -
1243 the topology of drift lineations as a key to the dynamic behaviour of former ice sheets.
1244 *Transactions of the Royal Society of Edinburgh – Earth Sciences*, 81, 327-347.
- 1245 Boulton, G.S., Smith, G.D., Jones, A.S., Newsome, J., 1985. Glacial geology and glaciology
1246 of the last mid-latitude ice sheets. *J. Geol. Soc. Lond.*, 142, 4467-474.
- 1247 Briner, J. P., Miller, G. H., Davis, P. T., Finkel, R. C., 2006. Cosmogenic radionuclides from
1248 fjord landscapes support differential erosion by overriding ice sheets. *Geol. Soc. Am.*
1249 *Bull.* 118, 406–420.
- 1250 Briner, J.P., Bini, A.C., Anderson, R.S., 2009. Rapid early Holocene retreat of a Laurentide
1251 outlet glacier through an Arctic fjord. *Nat. Geosci.*, 2, 496-499.
- 1252 Broecker, W.S., Kennett, J., Flower, B., Teller, J., Trumbore, S., Bonani, G. & Wolfli, W.,
1253 1989. Routing of meltwater from the Laurentide Ice Sheet during the Younger Dryas
1254 cold episode. *Nature*, 341, 318-321.
- 1255 Brookes, I.A., 2007. First recognition of a Laurentide Ice Stream : Robert Bell on Hudson
1256 Strait. *Géog. Phys. Quatern.* 61 (2–3), 211–216.
- 1257 Bryson, R.A., Wendland, W.M., Ives, J.D., Andrews, J.T., 1969. Radiocarbon isochrones on
1258 the disintegration of the Laurentide Ice Sheet. *Arctic and Alpine Research*, 1 (1), 1-13.
- 1259 Budd, W.F., Smith, I.N., 1981. The growth and retreat of ice sheets in response to orbital
1260 radiation changes. In, Allison, I. (Ed.), *Sea level, Ice, and Climatic Change*,
1261 international Association of Hydrological Sciences Publication, No. 131, p. 369-409.
- 1262 Carlson, A.E., Winsor, K., 2012. Northern Hemisphere ice-sheet response to past climate
1263 warming. *Nature Geoscience*, 5, 607-613.
- 1264 Carlson, A.E., Clark, P.U., 2012. Ice sheet sources of sea level rise and freshwater discharge
1265 during the last deglaciation. *Rev. Geophys.* 50 (4), 4007-4072.
- 1266 Carlson, A.E., Clark, P.U., Raisbeck, G.M., Brook, E.J., 2007. Rapid Holocene deglaciation
1267 of the Labrador sector of the Laurentide ice sheet. *J. Clim.* 20, 5126-5133.
- 1268 Carlson, A.E., LeGrande, A.N., Oppo, D.W., Came, R.E., Schmidt, G.A., Anslow, F.S.,
1269 Licciardi, J.M., Obbink, E.A., 2008. Rapid early Holocene deglaciation of the
1270 Laurentide Ice Sheet. *Nature Geoscience*, 1, 620-624.
- 1271 Carlson, A.E., Anslow, F.S., Obbink, E.A., LeGrande, A.N., Ullman, D.J., Licciardi, J.M.,
1272 2009a. Surface-melt driven Laurentide Ice Sheet retreat during the early Holocene.
1273 *Geophys. Res. Lett.*, 36, L24502, doi: 10.1029/2009GL040948.
- 1274 Carlson, A.E., Clark, P.U., Haley, B.A., Klinkhammer, G.P., 2009b. Routing of western
1275 Canadian Plains runoff during the 8.2 ka cold event. *Geophys. Res. Lett.* 36, L14704.
- 1276 Catto, N.R., Liverman, D.G.E., Bobrowsky, P.T., Rutter, N.W., 1996. Laurentide,
1277 Cordilleran, and Montane glaciation in the western Peace River-Grande Prairie
1278 region, Alberta and British Columbia. *Quaternary International*, 32, 21-32.
- 1279 Chamberlin, T. C., 1913. *Map of North America during the great ice age*. Rand McNally &
1280 Co., Chicago. 57 x 40 inches. Scale: 1 inch = 104 miles. Colored.
- 1281 Clark, C.D., 1993. Mega-scale glacial lineations and cross-cutting ice-flow landforms. *Earth*
1282 *Surf. Process. Landf.*, 18 (1), 1–29.
- 1283 Clark, C.D., Stokes, C.R., 2001. Extent and basal characteristics of the M'Clintock Channel
1284 Ice Stream. *Quat. Int.* 86 (1), 81–101.
- 1285 Clark, P.U., Josenhans, H.W., 1990. Reconstructed ice-flow patterns and ice limits using drift
1286 pebble lithology, outer Nachvak Fiord, northern Labrador: discussion. *Canadian*
1287 *Journal of Earth Sciences*, 27, 1002–1006.
- 1288 Clark, P.U., Clague, J.J., Currey, B.B., Dreimanis, A., Hicock, S.R., Miller, G.H., Berger,
1289 G.W., Eyles, N., Lamothe, M., Miller, B.B., Mott, R.J., Oldale, R.N., Stea, R.R.,

- 1290 Szabo, J.P., Thorleifson, L.H., Vincent, J.-S., 1993. Initiation and development of the
1291 Laurentide and Cordilleran ice sheets following the last interglaciation. *Quaternary*
1292 *Science Reviews*, 12 (2), 79-114.
- 1293 Clark, P.U., Licciardi, J.M., MacAyeal, D.R., Jenson, J.W., 1996. Numerical reconstruction
1294 of a soft-bedded Laurentide Ice Sheet during the last glacial maximum. *Geology*, 24
1295 (8), 679-682.
- 1296 Clark, P.U., Marshall, S.J., Clarke, G.K.C., Hostetler, S.W., Licciardi, J.M., Teller, J.T.,
1297 2001. Freshwater forcing of abrupt climate change during the last glaciation. *Science*
1298 293, 283-287.
- 1299 Clark, P.U., Dyke, A.S., Shakun, J.D., Carlson, A.E., Clark, J., Wohlfarth, B., Mitrovica,
1300 J.X., Hostetler, S.W., McCabe, A., 2009. The Last Glacial Maximum. *Science*, 325,
1301 710-714.
- 1302 Clayton, L., Moran, S.R., 1982. Chronology of Late Wisconsin glaciation in middle North
1303 America. *Quaternary Science Reviews*, 1, 55-82.
- 1304 Clayton, L., Teller, J. T., Attig, J. W., 1985. Surging of the southwestern part of the
1305 Laurentide Ice Sheet. *Boreas*, 14, 235-241.
- 1306 CLIMAP Project Members., 1976. The surface of ice-age earth. *Science*, 191 (4232), 1131-
1307 1137.
- 1308 Coleman, A.P., 1920, Extent and thickness of the Labrador Ice Sheet. *Geological Society of*
1309 *America, Bulletin*, 31, 319-328.
- 1310 Colleoni, F., Wekerle, C., Näslund, J-O., Brandefelt, J., Masina, S., 2016. Constraint on the
1311 penultimate glacial maximum Northern Hemisphere ice topography (\approx 140 kyrs BP).
1312 *Quaternary Science Reviews*, 137, 97-112.
- 1313 Condron, A., Winsor, P., 2012. Meltwater routing and the Younger Dryas. *Proc. Nat. Acad.*
1314 *Sci.*, 109 (49), 19928-19933.
- 1315 Cutler, K.B., Edwards, R.L., Taylor, F.W., Cheng, H., Adkins, J., Gallup, C.D., Cutler, P.M.,
1316 Burr, G.S., Bloomo, A.L., 2003. Rapid sea-level fall and deep-ocean temperature
1317 change since the last interglacial period. *Earth and Planetary Science Letters*, 206,
1318 253-271.
- 1319 Cutler, P.I., Mickelson, D.M., Colgan, P.M., MacAyeal, D.R., Parizek, B.R., 2001. Influence
1320 of the Great Lakes on the dynamics of the southern Laurentide Ice sheet: numerical
1321 experiments. *Geology*, 29 (11), 1039-1042.
- 1322 Dalton, A.S., Finkelstein, S.A., Barnett, P.J., Forman, S.L., 2016. Constraining the Late
1323 Pleistocene history of the Laurentide Ice Sheet by dating the Missinaibi Formation,
1324 Hudson Bay Lowlands, Canada. *Quat. Sci. Rev.*, 146, 288-299.
- 1325 Daly, R.A., 1902. The geology of the northeast coast of Labrador. *Harvard University*
1326 *Museum of Comparative Zoology Bulletin*, 38, 205-270.
- 1327 De Angelis, H., Kleman, J., 2005. Palaeo-ice streams in the northern Keewatin sector of the
1328 Laurentide ice sheet. *Ann. Glaciol.*, 42, 135-144.
- 1329 De Angelis, H., Kleman, J., 2007. Palaeo-ice streams in the Foxe/Baffin sector of the
1330 Laurentide Ice Sheet. *Quat. Sci. Rev.*, 26 (9-10), 1313-1331.
- 1331 Denton, G.H., Hughes, T., 1981. *The Last Great Ice Sheets*. Wiley-Interscience, New York,
1332 484 pp.
- 1333 Deschamps, P., Durand, N., Bard, E., Hamelin, B., Camoin, G., Thomas, A.L., Henderson,
1334 G.M., Okuno, J., Yokoyama, Y., 2012. Ice-sheet collapse and sea-level rise at the
1335 Bølling warming 14,600 years ago. *Nature*, 483, 559-564)
- 1336 Dixon, E.J., 2013. Late Pleistocene colonization of North America from Northeast Asia: new
1337 insights from large-scale paleogeographic reconstructions. *Quaternary International*,
1338 285, 57-67.

- 1339 Donn, W. L., Farrand, W. R., Ewing, M., 1962. Pleistocene ice volumes and sea-level
1340 lowering. *J. Geol.*, 70, 206-214.
- 1341 Dredge, L.A., Thorleifson, L.H., 1987. The Middle Wisconsinan history of the Laurentide Ice
1342 Sheet. *Géographie physique et Quaternaire* , 41 (2), 215-235.
- 1343 Dredge, L.A., Cowan, W.R., 1989. Quaternary geology of the southwestern Canadian Shield.
1344 In, Fulton, R.J. (Ed.), *Quaternary Geology of Canada and Greenland*. Geological
1345 Survey of Canada, Ottawa, pp. 214–235.
- 1346 Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P.,
1347 Rahmstorf, S., Raymo, M.E., 2015. Sea-level rise due to polar ice-sheet mass loss
1348 during past warm periods. *Science*, 349, p. 153.
- 1349 Dyke, A.S., 1984. Quaternary geology of Boothia Peninsula and northern District of
1350 Keewatin, central Canadian Arctic. *Geol. Surv. Can. Mem.* 407, pp. 26.
- 1351 Dyke, A.S., 1999. The last glacial maximum of Devon Island, arctic Canada: support for an
1352 Inuitian Ice Sheet. *Quaternary Science Reviews*, 18, 393–420.
- 1353 Dyke, A.S., 2004. An outline of North American deglaciation with emphasis on central and
1354 northern Canada. In: Ehlers, J., Gibbard, P.L. (Eds.), *Quaternary Glaciations-extent
1355 and Chronology, Part II*. Elsevier Science and Technology Books, Amsterdam, pp.
1356 371-406.
- 1357 Dyke, A.S., Prest, V.K., 1987. Late Wisconsinan and Holocene history of the Laurentide Ice
1358 Sheet. *Géog. Phys. Quaternaire*, 41 (2), 237-263.
- 1359 Dyke, A.S., Morris, T.F., 1988. Canadian landform examples. 7. Drumlin fields, dispersal
1360 trains, and ice streams in Arctic Canada. *Can. Geogr.*, 32 (1), 86–90.
- 1361 Dyke, A.S., Savelle, J.M., 2000. Major end moraines of Younger Dryas age on Wollaston
1362 Peninsula, Victoria Island, Canadian Arctic: implications for paleoclimate and for
1363 formation of hummocky moraine. *Canadian Journal of Earth Sciences*, 37, 601-619.
- 1364 Dyke, A.S., Dredge, L.A., Vincent, J.-S., 1982. Configuration and dynamics of the
1365 Laurentide Ice Sheet during the Late Wisconsin maximum. *Géog. Phys. Quatern.* 36
1366 (1–2), 5–14.
- 1367 Dyke, A.S., Andrews, J.T., Clark, P.U., England, J.H., Miller, G.H., Shaw, J., Veillette, J.J.,
1368 2002. The Laurentide and Inuitian ice sheets during the Last Glacial Maximum.
1369 *Quat. Sci. Rev.*, 21, 9-31.
- 1370 Dyke, A.S., Moore, A., Robertson, L., 2003. Deglaciation of North America. Geological
1371 Survey of Canada Open File 1574.
- 1372 England, J.H. 1976a. Postglacial isobases and uplift curves from the Canadian and Greenland
1373 High Arctic. *Arctic and Alpine Research*, 8 (1), 61-78.
- 1374 England, J.H., 1976b. Late Quaternary glaciation of the eastern Queen Elizabeth Islands,
1375 N.W.T, Canada: Alternative models. *Quaternary Research* ,6, 185–202.
- 1376 England, J.H., 1999. Coalescent Greenland and Inuitian ice during the last glacial
1377 maximum: revising the quaternary of the Canadian high arctic. *Quaternary Science
1378 Reviews*, 18, 421–456.
- 1379 England, J.H., Smith, I.R., Evans, D.J.A., 2000. The last glaciation of east-central Ellesmere
1380 Island, Nunavut: ice dynamics, deglacial chronology, and sea level change. *Canadian
1381 Journal of Earth Sciences*, 37, 1355–1371.
- 1382 England, J.H., Furze, M.F.A., Doupé, J.P., 2009. Revision of the NW Laurentide Ice Sheet:
1383 implications for paleoclimate, the northeast extremity of Beringia, and Arctic Ocean
1384 sedimentation. *Quaternary Science Reviews*, 28, 1573–1596.
- 1385 Eriksson, A., Betti, L., Friend, A.D., Lycett, S.J., Singarayer, J.S., von Cramon-Taubadel, N.,
1386 Valdes, P.J., Balloux, F., Manica, A., 2012. Late Pleistocene climate change and the
1387 global expansion of anatomically modern humans. *Proceedings of the National
1388 Academy of Science USA*, 109, 16089-16094.

- 1389 Evans, D.J.A., Clark, C.D., Rea, B.R., 2008. Landform and sediment imprints of fast glacier
1390 flow in the southwest Laurentide Ice Sheet. *J. Quat. Sci.*, 23 (3), 249–272.
- 1391 Eyles, N., 2012. Rock drumlins and megaflutes of the Niagara Escarpment, Ontario, Canada:
1392 a hard bed landform assemblage cut by the Saginaw–Huron Ice Stream. *Quat. Sci.*
1393 *Rev.*, 55, 34–49.
- 1394 Eyles, N., Putkinen, N., 2014. Glacially-megalinedated limestone terrain of Anticosti Island,
1395 Gulf of St. Lawrence, Canada; onset zone of the Laurentian Channel Ice Stream.
1396 *Quat. Sci. Rev.*, 88, 125–134.
- 1397 Falconer, G., Ives, J. D., Løken, O. H., Andrews, J. T., 1965. Major end moraines in eastern
1398 and central Arctic Canada. *Geog. Bull.*, 7 (2), 137-153.
- 1399 Fisher, D.A., Reeh, N., Langley, K., 1985. Objective reconstructions of the Late Wisconsinan
1400 Laurentide Ice Sheet and the significance of deformable beds. *Géog. Phys. et Quat.*,
1401 39 (3), 229-238.
- 1402 Fisher, T.G., Smith, D.G., 1994. Glacial Lake Agassiz: its northwest maximum extent in
1403 Saskatchewan (Emerson phase). *Quaternary Science Reviews*, 13, 845-858.
- 1404 Fisher, T.G., Lowell, T.V., 2012. Testing northwest drainage from Lake Agassiz using extant
1405 ice margin and strandline data. *Quat. Int.* 260, 106-114.
- 1406 Fisher, T.G., Smith, D.G., Andrews, J.T., 2002. Preboreal oscillation caused by a glacial Lake
1407 Agassiz flood. *Quaternary Science Reviews*, 21, 873-878.
- 1408 Flint, R.F., 1943. Growth of the North American Ice Sheet during the Wisconsin Age.
1409 *Geological Society of America, Bulletin*, 54, 352-362.
- 1410 Flint, R.F., 1947. *Glacial Geology and the Pleistocene Epoch*. John Wiley, New York.
- 1411 Flint, R.F., 1971. *Glacial and Quaternary Geology*. John Wiley, New York.
- 1412 Flint, R.F., Demorest, M., Washburn, A.L., 1942. Glaciation of the Shickshock Mountains,
1413 Gaspé, Peninsula. *Geological Society of America, Bulletin*, 53, 1211-1230.
- 1414 Flint, R.F., et al., 1945. *Glacial Map of North America*. Geological Society of America,
1415 Special Paper 60.
- 1416 Forbes, D.L., Boyd, R., Shaw, J., 1991. Late Quaternary sedimentation and sea-level changes
1417 on the inner Scotian Shelf. *Continental Shelf Research*, 11, 1155–1179.
- 1418 Fulton, R.J., 1989. *Quaternary Geology of Canada and Greenland*. Geological Survey of
1419 Canada, Geology of Canada, No. 1, 839 pp.
- 1420 Gipp, M.R., 1994. Late Wisconsinan glaciation of Emerald Basin, Nova Scotia. *Canadian*
1421 *Journal of Earth Sciences*, 31, 554–566.
- 1422 Gipp, M.R., Piper, D.J.W., 1989. Chronology of Late Wisconsinan glaciation, Emerald
1423 Basin, Scotian Shelf. *Canadian Journal of Earth Sciences*, 26, 333–335.
- 1424 Goebel, T., Waters, M.R., O'Rourke, D.H., 2008. The Late Pleistocene dispersal of modern
1425 humans in the Americas. *Science*, 319, 1497-1502.
- 1426 Gosse, J., Grant, D.R., Klein, J., Lawn, B., 1995. Cosmogenic ¹⁰Be and ²⁶Al constraints on
1427 weathering zone genesis, ice cap basal conditions, and Long Range Mountain
1428 Newfoundland) glacial history. Programme, Abstracts, Field Guides, CANQUA
1429 CGRG Joint Meeting, St. John's, CA19.
- 1430 Gowan, E.J., 2013. An assessment of the minimum timing of ice free conditions of the
1431 western Laurentide Ice Sheet. *Quaternary Science Reviews*, 75, 100-113.
- 1432 Gregoire, L.J., Payne, A.J., Valdes, P.J., 2012. Deglacial rapid sea level rises caused by ice
1433 sheet saddle collapses. *Nature*, 487, 219-222.
- 1434 Gregoire, L.J., Otto-Bliesner, B., Valdes, P.J., Ivanovic, R., 2015a. Abrupt Bølling warming
1435 and ice saddle collapse contributions to the Meltwater Pulse 1a rapid sea level rise.
1436 *Geophysical Research Letters*, 43, 9130-9137, doi: 10.1002/2016GL070356.

- 1437 Gregoire, L.J., Valdes, P.J., Payne, A.J., 2015b. The relative contribution of orbital forcing
1438 and greenhouse gases to the North American deglaciation. *Geophysical Research*
1439 *Letters*, 42, doi: 10.1002/2015GL066005.
- 1440 Hardy, L., 1982. Le Wisconsinien supérieur à l'est de la baie James (Québec). *Naturaliste*
1441 *canadien*, 109, 333-351.
- 1442 Harington, C.R., 1989. Pleistocene vertebrate localities in the Yukon. In, Carter, L.D.,
1443 Hamilton, T.D., Galloway, J.P. (wds.), *Late Cenozoic History of the Interior Basins of*
1444 *Alaska and the Yukon*. United States Geological Survey Circular, 1026, 93-98.
- 1445 Heinrich, H., 1988. Origin and consequences of cyclic ice rafting in the Northeast Atlantic
1446 Ocean during the past 130,000 years. *Quat. Res.*, 29 (2), 142–152.
- 1447 Hemming, S.R., 2004. Heinrich events: massive late Pleistocene detritus layers of the North
1448 Atlantic and their global climate imprint. *Rev. Geophys.* 42, RG1005.
- 1449 Hicock, S.R., 1988. Calcareous till facies north of Lake Superior, Ontario: implications for
1450 Laurentide ice streaming. *Géog. Phys. Quatern.*, 42 (2), 120–135.
- 1451 Hicock, S.R., 1992. Lobal interactions and rheologic superposition in subglacial till near
1452 Bradtville, Ontario, Canada. *Boreas*, 21 (1), 73–88.
- 1453 Hicock, S.R., Dreimanis, A., 1992. Deformation till in the Great Lakes region: implications
1454 for rapid flow along the south-central margin of the Laurentide Ice Sheet. *Can. J.*
1455 *Earth Sci.*, 29 (7), 1565–1579.
- 1456 Hindmarsh, R.C.A., 2009. Consistent generation of ice streams via thermos-viscous
1457 instabilities modulated by membrane stresses. *Geophysical Research Letters*, 36 (6),
1458 doi: 10.1029/2008GL036877.
- 1459 Hodgson, D.A., 1994. Episodic ice streams and ice shelves during retreat of the
1460 northwesternmost sector of the late Wisconsinan Laurentide Ice Sheet over the central
1461 Canadian Arctic Archipelago. *Boreas*, 23 (1), 14–28.
- 1462 Hoffman, J.S., Carlson, A.E., Winsor, K., Klinkhammer, G.P., LeGrande, A.N., Andrews,
1463 J.T., Strasser, J.C., 2012. Linking the 8.2 ka event and its freshwater forcing in the
1464 Labrador Sea. *Geophys. Res. Lett.* 39, L18703.
- 1465 Hughes, T., 1977. West Antarctic ice streams. *Rev. Geophys.*, 15 (1), 1–46.
- 1466 Hughes, P. D., Gibbard, P. L., Ehlers, J., 2013. Timing of glaciation during the last glacial
1467 cycle: Evaluating the concept of a global 'Last Glacial Maximum' (LGM). *Earth-*
1468 *Science Reviews*, 125, 171-198.
- 1469 Hughes, T., Denton, G.H., Grosswald, M.G., 1977. Was there a late-Wiirm Arctic ice sheet?
1470 *Nature*, 266, 596-602.
- 1471 IPCC, 2013. Summary for Policymakers. In: *Climate Change 2013: The Physical Science*
1472 *Basis. Contribution of Working Group I to the Fifth Assessment Report of the*
1473 *Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner,
1474 M. Tignor, S. K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley
1475 (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York,
1476 NY, USA.
- 1477 Ives, J.D., 1957. Glaciation of the Torngat mountains, northern Labrador. *Arctic* 10, 67-87.
- 1478 Ives, J.D., 1978. The maximum extent of the Laurentide Ice Sheet along the East Coast of
1479 North America during the Last Glaciation. *Arctic*, 31 (1), 24-53.
- 1480 Ives, J.D., Andrews, J.T., Barry, R.G., 1975. Growth and decay of the Laurentide Ice Sheet
1481 and comparisons with Fenno-Scandinavia. *Die Naturwissenschaften*, 62, 118-125.
- 1482 Jennings, A.E., 1993. The Quaternary history of Cumberland Sound, southeast Baffin Island:
1483 the marine evidence. *Géographie Physique et Quaternaire*, 47, 21–42.
- 1484 Jennings, A.E., Tedesco, K.A., Andrews, J.T. & Kirby, M.E., 1996. Shelf erosion and glacial
1485 ice proximity in the Labrador Sea during and after Heinrich events (H-3 or 4 to H-0)
1486 as shown by foraminifera. In: Andrews, J.T., Austin, W.E.N., Bergstrom, H.,

- 1487 Jennings, A.E. (Eds.), *Palaeoceanography of the North Atlantic Margins*. Geological
 1488 Society Special Publication No. 111, 29-49.
- 1489 Jennings, A.E., Andrews, A., Pearce, C., Wilson, L., Ólfasdóttir, S., 2015. Detrital carbonate
 1490 peaks on the Labrador shelf, a 13-7 ka template for freshwater forcing from the
 1491 Hudson Strait outlet of the Laurentide Ice Sheet into the subpolar gyre. *Quat. Sci. Rev.*
 1492 107, 62-80.
- 1493 Josenhans, H.W., Zevenhuizen, J., 1990. Dynamics of the Laurentide Ice Sheet in Hudson
 1494 Bay, Canada. *Marine Geology*, 92, 1-26.
- 1495 Josenhans, H., Lehman, S., 1999. Quaternary stratigraphy and glacial history of the Gulf of
 1496 St. Lawrence, Canada. *Canadian Journal of Earth Sciences*, 36, 1327–1345.
- 1497 Kaplan, M.R., 1999. *The last glaciation of the Cumberland Sound region, Baffin Island,*
 1498 *Canada, based on glacial geology, cosmogenic dating, and numerical modeling.*
 1499 Unpublished Ph.D. Thesis, University of Colorado, Boulder, 206 pp.
- 1500 Karrow, P.F., Calkin, P.E., 1985. Quaternary evolution of the Great Lakes. *Geological*
 1501 *Association of Canada Special Paper*, 30, 258 pp.
- 1502 Kennedy, K.E., Froese, D.G., Zazula, G.D., Lauriol, B., 2010. Last Glacial Maximum age for
 1503 the northwest Laurentide maximum from the Eagle River spillway and delta complex,
 1504 northern Yukon. *Quaternary Science Reviews*, 29, 1288-1300.
- 1505 King, L.H., 1996. Late Wisconsinan ice retreat from the Scotian Shelf. *Geological Society of*
 1506 *America Bulletin*, 108, 1056–1067.
- 1507 Kirby, M.E., Andrews, J.T., 1999. Mid-Wisconsinan Laurentide Ice Sheet growth and decay:
 1508 implications for Heinrich events 3 and 4. *Paleoceanography* 14 (2), 211-223.
- 1509 Kirchner, N., Ahlkrona, J., Gowan, E.J., Lötstedt, P., Lea, J.M., Noormets, R., von Sydow,
 1510 L., Dowdeswell, J.A., Benham, T., 2016. Shallow ice approximation, second order
 1511 shallow ice approximation, and full Stokes models: A discussion of their roles in
 1512 palaeo-ice sheet modelling and development. *Quaternary Science Reviews*, 147, 136-
 1513 147.
- 1514 Kleman, J., Borgström, I., 1996. Reconstruction of palaeo-ice sheets: the use of
 1515 geomorphological data. *Earth Surf. Process. Landf.*, 21 (10), 893–909.
- 1516 Kleman, J., Applegate, P.J., 2014. Durations and propagation patterns of ice sheet instability
 1517 events. *Quaternary Science Reviews* 92, 32-39.
- 1518 Kleman, J., Fastook, J., Stroeven, A., 2002. Geologically and geomorphologically
 1519 constrained numerical model of Laurentide Ice Sheet inception and build-up.
 1520 *Quaternary International*, 95-6, 87-98.
- 1521 Kleman, J., Jansson, K., De Angelis, H., Stroeven, A.P., Hattestrand, C., Alm, G., Glasser,
 1522 N.F., 2010. North American Ice Sheet build-up during the last glacial cycle, 115-21
 1523 kyr. *Quaternary Science Reviews* 29, 17-18.
- 1524 Koerner, R.M., 1980. Instantaneous glacierization, the rate of albedo change, and feedback
 1525 effects at the beginning of an Ice Age. *Quaternary Research* 13, 153-159.
- 1526 Krabbendam, M., Eyles, N., Putkinen, N., Bradwell, T., Arbelaez-Moreno, L., 2016.
 1527 Streamlined hard beds formed by palaeo-ice streams: a review. *Sedimentary Geology*,
 1528 338, 24-50.
- 1529 Lacelle, D., Lauroil, B., Zazula, G., Ghaleb, B., Utting, N., Clark, I.D., 2013. Timing of
 1530 advance and basal condition of the Laurentide Ice Sheet during the last glacial
 1531 maximum in the Richardson Mountains, NWT. *Quaternary Research*, 80, 274-283.
- 1532 Lakeman, T. R., England, J. H., 2012. Palaeo-glaciological insights from the age and
 1533 morphology of the Jesse moraine belt, western Canadian Arctic. *Quat. Sci. Rev.* 47,
 1534 82–100.

- 1535 Lakeman, T. R., England, J. H., 2013. Late Wisconsinan glaciation and postglacial relative
1536 sea-level change on western Banks Island, Canadian Arctic Archipelago. *Quat. Res.*
1537 80, 99–112.
- 1538 Lambeck, K., Rouby, H., Purcell, A., Sun, Y., Sambridge, M., 2014. Sea level and global ice
1539 volumes from the last glacial maximum to the Holocene. *Proc. Natl. Acad. Sci.*
1540 *U.S.A.* 111, 15,296–15,303.
- 1541 Lambeck, K., Purcell, A., Zhao, S., 2017. The North American Late Wisconsin ice sheet and
1542 mantle viscosity from glacial rebound analyses. *Quaternary Science Reviews*, 158,
1543 172–210.
- 1544 Lambert, A., Courtier, N., James, T.S., 2006. Long-term monitoring by absolute gravimetry:
1545 tides to postglacial rebound. *Journal of Geodynamics*, 41, 307–317.
- 1546 Lamoureux, S.F., England, J.H., 2000. Late Wisconsinan glaciation of the central sector of
1547 the Canadian High Arctic. *Quaternary Research*, 54, 182–188.
- 1548 Licciardi, J.M., Clark, P.U., Jenson, J.W., MacAyeal, D.R., 1998. Deglaciation of a soft-
1549 bedded Laurentide Ice Sheet. *Quaternary Science Reviews*, 17, 427–448.
- 1550 Lowe, J.J., Ammann, B., Birks, H.H., Bjorck, S., Coope, G.R., Cwynar, L. Beaulieu, J-L.,
1551 Mott, R.J., Peteet, D.M., Walker, M.J.C., 1994. Climatic changes in areas adjacent to
1552 the North Atlantic during the last glacial-interglacial transition (14-9 ka BP): a
1553 contribution to IGCP-253. *Journal of Quaternary Science*, 9, 185–198.
- 1554 Løken, O.H., 1966. Baffin Island refugia older than 54,000 years. *Science*, 153, 1378–1380.
- 1555 MacAyeal, D., 1993. Binge/purge oscillations of the Laurentide ice sheet as a cause of the
1556 North Atlantic's Heinrich events. *Paleoceanography*, 8 (6), 775–784.
- 1557 MacLean, B., Blasco, S., Bennett, R., England, J., Rainey, W., Hughes-Clarke, J., Beaudoin,
1558 J., 2010. Ice keel seabed features in marine channels of the central Canadian Arctic
1559 Archipelago: evidence for former ice streams and iceberg scouring. *Quat. Sci. Rev.*,
1560 29 (17–18), 2280–2301.
- 1561 Margold, M., Stokes, C.R., Clark, C.D., Kleman, J., 2015a. Ice streams in the Laurentide Ice
1562 Sheet: a new mapping inventory. *Journal of Maps*, 11 (3), 380–395.
- 1563 Margold, M., Stokes, C.R., Clark, C.D., 2015b. Ice streams in the Laurentide Ice Sheet:
1564 identification, characteristics and comparison to modern ice sheets. *Earth-Science*
1565 *Reviews*, 143, 117–146.
- 1566 Marcott, S.A., Shakun, J.D., Clark, P.U., Mix, A.C., 2013. A reconstruction of regional and
1567 global temperature for the past 11,300 years. *Science* 339, 1198–1201.
- 1568 Marsella, K.A., Bierman, P.R., Davis, P.T., Caffee, M.W., 2000. Cosmogenic ¹⁰Be and ²⁶Al
1569 ages for the last glacial maximum, eastern Baffin Island, Arctic Canada. *Geological*
1570 *Society of America Bulletin*, 112, 1296–1312.
- 1571 Marshall, S.J., Clarke, G.K.C., 1997a. A continuum mixture model of ice stream
1572 thermomechanics in the Laurentide Ice Sheet. 1. Theory. *J. Geophys. Res. Solid*
1573 *Earth*, 102 (B9), 20599–20613.
- 1574 Marshall, S.J., Clarke, G.K.C., 1997b. A continuum mixture model of ice stream
1575 thermomechanics in the Laurentide Ice Sheet. 2. Application to the Hudson Strait Ice
1576 Stream. *J. Geophys. Res. Solid Earth*, 102 (B9), 20615–20637.
- 1577 Marshall, S.J., 2002. Modelled nucleation centres of the Pleistocene ice sheets from an ice
1578 sheet model with subgrid parameterizations. *Quaternary International*, 95–96, 125–
1579 137.
- 1580 Marshall, S.J., Clark, P.U., 2002. Basal temperature evolution of North American ice sheets
1581 and implications for the 100-kyr cycle. *Geophysical Research Letters*, 29, (4), 2214.
- 1582 Marshall, S.J., Clarke, G.K.C., Dyke, A.S., Fisher, D.A., 1996. Geologic and topographic
1583 controls on fast flow in the Laurentide and Cordilleran Ice Sheets. *J. Geophys. Res.*
1584 *Solid Earth*, 101 (B8), 17827–17839.

- 1585 Marshall, S.J., Tarasov, L., Clarke, G.K.C., Peltier, W.R., 2000. Glaciological reconstruction
1586 of the Laurentide Ice Sheet: physical processes and modelling challenges. *Canadian*
1587 *Journal of Earth Sciences* 37 (5), 769-793.
- 1588 Mathews, W. H., 1974. Surface profiles of the Laurentide Ice Sheet in its marginal areas,
1589 *Journal of Glaciology*, 13, 37-43.
- 1590 Mayewski, P.A., Denton, G.H., Hughes, T.J., 1981. Late Wisconsin ice sheets of North
1591 America. In, Denton, G.H., Hughes, T. (Eds) *The Last Great Ice Sheets*. Wiley-
1592 Interscience, New York, p. 67-178.
- 1593 Mazzotti, S., Lambert, A., Henton, J., James, T.S., Courtier, N., 2011. Absolute gravity
1594 calibration of GPS velocities and glacial isostatic adjustment in mid-continent North
1595 America. *Geophysical Research Letters* 38,.doi. 10.1029/2011GL049846.
- 1596 Miller, A.A.L., Fader, G.B.J., Moran, K., 2001. Late Wisconsinan ice advances, ice extent,
1597 and glacial regimes interpreted from seismic data, sediment physical properties, and
1598 foraminifera: Halibut Channel, Grand Banks of Newfoundland. In, Weddle, T.K.,
1599 Retelle, M.J. (Eds.), *Deglacial History and Relative Sea-Level Changes, Northern*
1600 *New England and Adjacent Canada*. Geological Society of America Special Paper,
1601 351, 51- 107.
- 1602 Miller, G.H., Dyke, A.S., 1974. Proposed extent of late Wisconsin ice on Baffin Island.
1603 *Geology* , 2, 125–130.
- 1604 Miller, G.H., Kaufman, D.S., 1990. Rapid fluctuations of the Laurentide Ice Sheet at the
1605 mouth of Hudson Strait: new evidence for ocean/ice-sheet interactions as a control on
1606 the Younger Dryas. *Paleoceanography*, 5, 907-919.
- 1607 Miller, G.H., Wolfe, A.P., Steig, E.J., Sauer, P.E., Kaplan, M.R., Briner, J.P., 2002. The
1608 Goldilocks dilemma: big ice, little ice, or “just right” ice in the Eastern Canadian
1609 Arctic. *Quaternary Science Reviews*, 21, 33-48.
- 1610 Mooers, H.D., Lehr, J.D., 1997. Terrestrial record of Laurentide Ice Sheet reorganization
1611 during Heinrich events. *Geology*, 25 (11), 987.
- 1612 Morrill, C., Ward, E.M., Wagner, A.J., Otto-Bliesner, B.L., Rosenbloom, N., 2014. Large
1613 sensitivity to freshwater forcing location in 8.2 ka simulations. *Paleoceanography* 29,
1614 930-945.
- 1615 Mosher, D.C., Piper, D.J.W., Vilks, G.V., Aksu, A.E., Fader, G.B.J., 1989. Evidence for
1616 Wisconsinan glaciations in the Verrill Canyon area, Scotian Slope. *Quaternary*
1617 *Research*, 31, 27–40.
- 1618 Murton, J.B., Frenchen, M., Maddy, D., 2007. Luminescence dating of mid- to Late
1619 Wisconsinan aeolian sand as a constraint on the last advance of the Laurentide Ice
1620 Sheet across the Tuktoyaktuk coastlands, western Arctic Canada. *Canadian Journal*
1621 *of Earth Sciences*, 44, 857-869.
- 1622 Murton, J.B., Bateman, M.D., Dallimore, S.R., Teller, J.T., Yang, Z., 2010. Identification of
1623 Younger Dryas outburst flood path from Lake Agassiz to the Arctic Ocean. *Nature*
1624 464, 740-743.
- 1625 Murton, J.B, Bateman, M.D, Waller, R.I., Whiteman, C.A., 2015. *Late Wisconsin glaciation*
1626 *of Hadwen and Summer islands, Tuktoyaktuk Coastlands, NWT, Canada*. In,
1627 GEOQuébec2015: 7th Canadian Permafrost Conference, 20–23 September, 2015,
1628 Quebec City, Canada.
- 1629 Nicholl, J.A.L., Hodell, D.A., Naafs, B.D.A., Hillaire-Marcel, C., Channell, J.E.T., Romero,
1630 O.E., 2012. A Laurentide outburst flooding event during the last interglacial period.
1631 *Nat. Geosci.* 5 (12), 901-904.
- 1632 Nick, F.M., Vieli, A., Anderson, M.L., Joughin, I., Payne, A., Edwards, T.J., Pattyn, F., van
1633 de Wal, R.S.W., 2013. Future sea-level rise from Greenland’s main outlet glaciers in a
1634 warming climate. *Nature*, 497, 235-238.

- 1635 Nixon, F. C., England, J. H., 2014. Expanded Late Wisconsinan ice cap and ice sheet margins
1636 in the western Queen Elizabeth Islands, Arctic Canada. *Quat. Sci. Rev.* 91, 146–164.
- 1637 Occhietti, S., Parent, M., Lajeunesse, P., Robert, F., Govare, E., 2011. Late Pleistocene-early
1638 Holocene decay of the Laurentide ice sheet in Quebec-Labrador. In, Ehlers, J.,
1639 Gibbard, P.L., Hughes, P.D. (Eds.) *Quaternary Glaciations - Extent and Chronology*,
1640 vol. 15. Elsevier Science and Technology Books, Amsterdam, pp. 601-630.
- 1641 Odell, N.E., 1933. The mountains of northern Labrador. *Geographical Journal*, 82(3-4), 193-
1642 211.
- 1643 O’Cofaigh, C., England, J., Zreda, M., 2000. Late Wisconsinan glaciation of southern Eureka
1644 Sound: evidence for extensive Innuitian ice in the Canadian high arctic during the last
1645 glacial maximum. *Quaternary Science Reviews*, 19, 1319–1341.
- 1646 Ó Cofaigh, C., Evans, D.J.A., Smith, I.R., 2010. Large-scale reorganization and
1647 sedimentation of terrestrial ice streams during late Wisconsinan Laurentide Ice Sheet
1648 deglaciation. *Geol. Soc. Am. Bull.*, 122 (5–6), 743–756.
- 1649 Patterson, C.J., 1997. Southern Laurentide ice lobes were created by ice streams: Des Moines
1650 lobe in Minnesota, USA. *Sediment. Geol.*, 111 (1–4), 249–261.
- 1651 Patterson, C.J., 1998. Laurentide glacial landscapes: the role of ice streams. *Geology*, 26 (7),
1652 643–646.
- 1653 Paterson, W.S.B., 1972. Laurentide Ice Sheet: estimated volumes during Late Wisconsin.
1654 *Reviews of Geophysics and Space Physics*, 10 (4), 885-917.
- 1655 Pedersen, M.W., Rueter, A., Schweger, C., Friebe, H., Staff, R.A., Kjeldsen, K.K., Mendoza,
1656 M.L.Z., Beaudoin, A.B., Zutter, C., Larsen, N.K., Potter, B.A., Nielsen, R., Rainville,
1657 R.A., Orlando, L., Meltzer, D.J., Kjaer, K.H., Willerslev, E., 2016. Postglacial
1658 viability and colonization in North America’s ice-free corridor. *Nature*, 537, 45-49.
- 1659 Peltier, W.R., 1994. Ice age paleotopography. *Science*, 265, 195–201.
- 1660 Peltier, W.R., 2004. Global glacial isostasy and the surface of the ice-age earth: The ICE-5G
1661 (VM2) model and GRACE. *Annu. Rev. Earth Planet. Sci.* 32, 111-149.
- 1662 Peltier, W.R., Argus, D.F., Drummond, R., 2015. Space geodesy constrains ice-age terminal
1663 deglaciation: the global ICE-6G_C (VM5a) model. *J. Geophys. Res. Solid Earth*, 120,
1664 450-487.
- 1665 Piper, D.J.W., Skene, K.I., 1998. Latest Pleistocene ice-rafting events on the Scotian Margin
1666 (eastern Canada) and their relationship to Heinrich events. *Palaeoceanography*, 13,
1667 205–214.
- 1668 Piper, D.J.W., Mudie, P.J., Fader, G.B., Josenhans, H.W., MacLean, B., Vilks, G., 1990.
1669 Quaternary geology. In, Keen, M.J. & Williams, G.L. (Eds.), *Geology of the*
1670 *Continental Margin of Eastern Canada*. Geological Survey of Canada, Geology of
1671 Canada, No. 2, 475-607.
- 1672 Prest, V.K., 1969. *Retreat of Wisconsin and Recent ice in North America*. Geological Survey
1673 of Canada Map, 1257A, scale 1:5 000 000.
- 1674 Prest, V.K., 1984. The Late Wisconsinan glacier complex. In, R. J. Fulton (Ed.) Quaternary
1675 Stratigraphy of Canada - a Canadian contribution to IGCP Project 24. *Geological*
1676 *Survey of Canada, Paper 84-10*, p. 22-36.
- 1677 Prest, V.K., Grant, D.R., Rampton, V.N., 1968. Glacial Map of Canada. *Geological Survey*
1678 *of Canada Map*, 1253A, scale 1:5 000 000.
- 1679 Pritchard, H.D., Arthern, R.J., Vaughan, D.G., Edwards, L., 2009. Extensive dynamic
1680 thinning on the margins of the Greenland and Antarctic ice sheets. *Nature*, 461, 971-
1681 975.
- 1682 Ramsay, W., 1931. Changes of sea level resulting from the increase and decrease of
1683 glaciations. *Fennia*, 52, 1-62.

- 1684 Rasmussen, S.O., Anderson, K.K., Svensson, A.M., Steffensen, J.P., Vinther, B.M., Clausen,
1685 H.B., Siggaard-Andersen, M.-L., Johnsen, S.J., Larsen, L.B., Dahl-Jensen, D., Bigler,
1686 M., Røstlisberger, R., Fischer, H., Goto-Azuma, K., Hansson, M.E., Ruth, U., 2006.
1687 A new Greenland ice core chronology for the last glacial termination. *J. Geophys.*
1688 *Res.* 111, D06102.
- 1689 Ritz, C., Edwards, T.L., Durand, G., Payne, A.J., Peyaud, V., Hindmarsh, R.C.A., 2015.
1690 Potential sea-level rise from Antarctic ice-sheet instability constrained by
1691 observations. *Nature*, 528, 115-118.
- 1692 Robel, A.A., Tziperman, E., 2016. The role of ice stream dynamics in deglaciation. *Journal*
1693 *of Geophysical Research – Earth Surface*, 121, doi:10.1002/2016JF003937.
- 1694 Ross, M., Campbell, J.E., Parent, M., Adams, R.S., 2009. Palaeo-ice streams and the
1695 subglacial landscape mosaic of the North American mid-continental prairies. *Boreas*
1696 38 (3), 421–439.
- 1697 Ross, M., Lajeunesse, P., Kosar, K.G.A., 2011. The subglacial record of northern Hudson
1698 Bay: insights into the Hudson Strait Ice Stream catchment. *Boreas*, 40 (1), 73–91.
- 1699 Ross, M., Utting, D.J., Lajeunesse, P., Kosar, K.G.A., 2012. Early Holocene deglaciation of
1700 northern Hudson Bay and Foxe Channel constrained by new radiocarbon ages and
1701 marine reservoir correction. *Quat. Res.* 78, 82-94.
- 1702 Schnikter, D., Belknap, D.F., Bacchus, T.S., Friez, J.K., Lusardi, B.A., Popek, D.M., 2001.
1703 Deglaciation of the Gulf of Maine. *Geological Society of America Special Paper*, 351,
1704 9–34.
- 1705 Shapiro, B., Drummond, A.J., Rambaut, A., Wilson, M.C., Matheus, P.E., Sher, A.V.,
1706 Pybus, M.C., Gilbert, M.T.P., Barnes, I., Binladen, J., Willerslev, E., Hansen, A.J.,
1707 Baryshnikov, G.F., Burns, J.A., Davydov, S., driver, J.C., Froese, D.G., Harington,
1708 C.R., Keddie, G., Kosintsev, P., Kunz, M.L., martin, L.D., Stephenson, R.O., Storer,
1709 J., Tedford, R., Zimov, S., Cooper, A., 2004. The rise and fall of the Beringian Steppe
1710 Bison. *Science*, 306, 1561-1565.
- 1711 Shaw, J., 2003. Submarine moraines in Newfoundland coastal waters: implications for
1712 deglaciation of Newfoundland and adjacent areas. *Quaternary International*, 99-100,
1713 115-134.
- 1714 Shaw, J., Piper, D.J.W., Fader, G.B.J., King, E.L., Todd, B.J., Bell, T., Batterson, M.J.,
1715 Liverman, D.G.E., 2006. A conceptual model of the deglaciation of Atlantic Canada.
1716 *Quaternary Science Reviews*, 25, 2059-2081.
- 1717 Shaw, J., Todd, B.J., Brushett, D., Parrott, D.R., Bell, T., 2009. Late Wisconsinan glacial
1718 landsystems on Atlantic Canadian shelves: new evidence from multibeam and single-
1719 beam sonar data. *Boreas*, 38 (1), 146–159.
- 1720 Shepherd, A., Ivins, E.R., Barletta, A.G., Bentley, V.R., Bettadpur, M.J., Briggs, S.,
1721 Bromwich, K.H., Forsberg, D.H., Galin, R., Horwath, N., Jacobs, M., Joughin, S.,
1722 King, I., Lenaerts, M.A., Li, J.T., Ligtenberg, J., Luckman, S.R., Luthcke, A.,
1723 McMillan, S.B., Meister, M., Milne, R., Mouginot, G., Muir, J., Nicolas, A., Paden,
1724 J.P., Payne, J., Pritchard, A.J., Rignot, H., Rott, E., Sorensen, H., Scambos, L.S.,
1725 Scheuchl, T.A., Schrama, B., Smith, E.J., Sundal, B., van Angelen, A.V., van de Berg,
1726 J.H., van den Broeke, W.J., Vaughan, M.R., Velicogna, D.G., Wahr, I., Whitehouse,
1727 J., Wingham, P.L., Yi, D.J., Young, D., Zwally, H.J., 2012. A reconciled estimate of
1728 ice-sheet mass balance. *Science*, 338 (6111), 1183–1189.
- 1729 Shilts, W.W., 1980. Flow patterns in the central North American ice sheet. *Nature*, 286, 213-
1730 218.
- 1731 Shilts, W.W., Cunningham, C.M., Kaszycki, C.A., 1979. Keewatin Ice Sheet - re-evaluation
1732 of the traditional concept of the Laurentide Ice Sheet. *Geology*, 537-541

- 1733 Simon, K., James, T.S., Forbes, D.L., Telka, A.M., Dyke, A.S., Henton, J.A., 2014. A relative
1734 sea-level history for Arviat, Nunavut, and implications for Laurentide ice sheet
1735 thickness west of Hudson Bay. *Quat. Res.* 82, 185-197.
- 1736 Smith, D.G., Fisher, T.G., 1993. Glacial Lake Agassiz: the northwestern outlet and
1737 palaeoflood. *Geology* 21, 9-12.
- 1738 Stalker, A., 1977. The probable extent of classical Wisconsin ice in southern and central
1739 Alberta, *Canadian Journal of Earth Sciences*, 14, 2614-2619.
- 1740 Stea, R.R., Mott, R.J., 1998. Deglaciation of Nova Scotia: stratigraphy and chronology of
1741 lake sediment cores and buried organic sections. *Géographie Physique et*
1742 *Quaternaire*, 52, 3–21.
- 1743 St-Onge, D.A., 1987. The Sangamonian stage and the Laurentide Ice Sheet. *Géographie*
1744 *physique et Quaternaire*, 41 (2), 189-198.
- 1745 Stokes, C.R., Clark, C.D., 1999. Geomorphological criteria for identifying Pleistocene ice
1746 streams. *Ann. Glaciol.*, 28 (1), 67–74.
- 1747 Stokes, C.R., Clark, C.D., 2001. Palaeo-ice streams. *Quat. Sci. Rev.* 20 (13), 1437–1457.
- 1748 Stokes, C.R., Clark, C.D., 2003. The Dubawnt Lake palaeo-ice stream: evidence for dynamic
1749 ice sheet behaviour on the Canadian Shield and insights regarding the controls on ice-
1750 stream location and vigour. *Boreas* 32 (1), 263–279.
- 1751 Stokes, C.R., Clark, C.D., 2004. Evolution of late glacial ice-marginal lakes on the
1752 northwestern Canadian Shield and their influence on the location of the Dubawnt
1753 Lake palaeo-ice stream. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 215 (1–2), 155–
1754 71.
- 1755 Stokes, C.R., Tarasov, L., 2010. Ice streaming in the Laurentide Ice Sheet: a first comparison
1756 between data-calibrated numerical model output and geological evidence.
1757 *Geophysical Research Letters*, 37, L01501.
- 1758 Stokes, C.R., Clark, C.D., Storrar, R., 2009. Major changes in ice stream dynamics during
1759 deglaciation of the north-western margin of the Laurentide Ice Sheet. *Quat. Sci. Rev.*
1760 28 (7–8), 721–738.
- 1761 Stokes, C.R., Tarasov, L., Dyke, A.S., 2012. Dynamics of the North American Ice Sheet
1762 Complex during its inception and build-up to the Last Glacial Maximum. *Quat. Sci.*
1763 *Rev.*, 50, 86-104.
- 1764 Stokes, C.R., Tarasov, L., Blomdin, R., Cronin, T., Fisher, T.G., Gyllencreutz, R.,
1765 Hättestrand, C., Heyman, J., Hindmarsh, R.C.A., Hughes, A.L.C., Jakobsson, M.,
1766 Kirchner, N., Livingstone, J.J., Margold, M., Murton, J.B., Noormets, R., Peltier,
1767 W.R., Peteet, D.M., Piper, D.J.W., Preusser, F., Renssen, H., Roberts, D.H., Roche,
1768 D.M., Saint-Ange, F., Stroeven, A.P., Teller, J.T., 2015. On the reconstruction of
1769 palaeo-ice sheets: recent advances and future challenges. *Quat. Sci. Rev.*, 125, 15-49.
- 1770 Stokes, C.R., Margold, M., Clark, C.D., Tarasov, L., 2016. Ice stream activity scaled to ice
1771 sheet volume during Laurentide Ice Sheet deglaciation. *Nature* 530, 322-326.
- 1772 Storrar, R.D., Stokes, C.R., Evans, D.J.A., 2014. Increased channelization of subglacial
1773 drainage during deglaciation of the Laurentide Ice Sheet. *Geology*, 42 (3), 239–242.
- 1774 Stuiver, M., Reimer, P.J., and Reimer, R.W., 2017, CALIB 7.1 [WWW program] at
1775 <http://calib.org>, accessed 2017-2-1
- 1776 Sugden, D.E., 1977. Reconstruction of the morphology, dynamics and thermal characteristics
1777 of the Laurentide Ice Sheet at its maximum. *Arctic and Alpine Research*, 9 (1), 21-47.
- 1778 Sugden, D.E., Watts, S., 1977. Tors, felsenmeer and glaciation in northern Cumberland
1779 Peninsula, Baffin Island. *Canadian Journal of Earth Sciences*, 14, 2817-2823.
- 1780 Svendsen, J., Alexanderson, H., Astakhov, V., Demidov, I., Julian, A., Funder, S., Gataulling,
1781 V., Henriksen, M., Hjort, J., Houmark-Nielsen, M., Hubberten, H., Ingulfsson, O.,
1782 Jakobsson, M., Kjer, K., Larsen, E., Lokrantz, H., Lunkka, J., Lys, A., Mangerud, J.,

- 1783 Matiouchkov, A., Murray, A., Muller, P., Niessen, F., Nikolskaya, O., Polyak, L.,
 1784 Saarnisto, M., Siegert, C., Siegert, M., Spielhagen, R., Ruedige, S., 2004. Late
 1785 Quaternary ice sheet history of Northern Eurasia. *Quat. Sci. Rev.* 23, 1229-1271.
- 1786 Tanner, W.F., 1965. Cause and development of an Ice Age. *The Journal of Geology*, 73, 413-
 1787 429.
- 1788 Tarasov, L., Peltier, W.R., 2005. Arctic freshwater forcing of the Younger Dryas cold
 1789 reversal. *Nature* 435, 662-665.
- 1790 Tarasov, L., Dyke, A.S., Neal, R.M., Peltier, W.R., 2012. A data-calibrated distribution of
 1791 deglacial chronologies for the North American ice complex from glaciological
 1792 modelling. *Earth and Planetary Science Letters*, 315-316, 30-40.
- 1793 Teller, J.T., 2013. Lake Agassiz during the Younger Dryas. *Quaternary Research*, 80, 361-
 1794 369.
- 1795 Teller, J.T., Thorleifson, L.H., 1983. The Lake Agassiz-Lake Superior connection. In: Teller,
 1796 J.T., Clayton, L. (Eds.), *Glacial Lake Agassiz*. Geological Association of Canada
 1797 Special Paper, 26, 261-290.
- 1798 Thorleifson, L.H., Wyatt, P.H., Warman, T.A., 1993. Quaternary stratigraphy of the Severn
 1799 and Winisk drainage basins, northern Ontario. *Geological Survey of Canada Bulletin*,
 1800 442, 59 pp.
- 1801 Todd, B.J., Valentine, P.C., Longva, O., Shaw, J., 2007. Glacial landforms on German Bank,
 1802 Scotian Shelf: evidence for Late Wisconsinan ice-sheet dynamics and implications for
 1803 the formation of De Geer moraines. *Boreas*, 36 (2), 148–169
- 1804 Tushingham, A. M., Peltier, W. R., 1991. ICE-3G: A new global model of late Pleistocene
 1805 deglaciation based upon geophysical predictions of post-glacial relative sea level
 1806 change: *Journal of Geophysical Research*, 96, 4497–4523.
- 1807 Tyrell, J.B., 1898. The glaciation of north-central Canada. *Journal of Geology*, 6, 147-160.
- 1808 Ullman, D.J., Carlson, A.E., LeGrande, A.N., Anslow, F.S., Moore, A.K., Caffee, M.,
 1809 Syverson, K.M., Licciardi, J.M., 2015a. Southern Laurentide ice-sheet retreat
 1810 synchronous with rising boreal summer insolation. *Geology*, 43 (1), 23-26.
- 1811 Ullman, D.J., Carlson, A.E., Anslow, F.S., LeGrande, A.N., Licciardi, J.M., 2015b.
 1812 Laurentide ice-sheet instability during the last deglaciation. *Nature Geoscience*, 8,
 1813 534-537.
- 1814 Ullman, D.J., Carlson, A.E., Hostetler, S.W., Clark, P.U., Cuzzone, J., Milne, G.A., Winsor,
 1815 K., Caffee, M. 2016. Final Laurentide ice-sheet deglaciation and Holocene climate-
 1816 sea level change. *Quaternary Science Reviews*, 152, 49-59.
- 1817 Van den Broeke, M., Bamber, J., Ettema, J., Rignot, E., Schrama, E., van de Berg, W.J., van
 1818 Meijgaard, E., Velicogna, I., Wouters, B., 2009. Partitioning recent Greenland mass
 1819 loss. *Science*, 326, 984-986.
- 1820 Veillette, J.J., Dyke, A.S., Roy, M., 1999. Ice-flow evolution of the Labrador Sector of the
 1821 Laurentide Ice Sheet: a review, with new evidence from northern Quebec. *Quat. Sci.*
 1822 *Rev.*, 18 (8–9), 993–1019.
- 1823 Vettoretti, G., Peltier, W.R., 2013. Last Glacial Maximum ice sheet impacts on North Atlantic
 1824 climate variability: the importance of a sea ice lid. *Geophysical Research Letters* 40,
 1825 6378–6383.
- 1826 Vincent, J.-S., 1982. The Quaternary history of Banks Island, NWT, Canada. *Géographie*
 1827 *physique et Quaternaire*, 36, 209-232.
- 1828 Vincent, J.-S., Prest, V.K., 1987. The early Wisconsinan history of the Laurentide Ice Sheet.
 1829 *Géographie physique et Quaternaire* 41 (2), 199-213.
- 1830 Wickert, A.D., 2016. Reconstruction of North American drainage basins and river discharge
 1831 since the Last Glacial Maximum. *Earth Surf. Dynam.*, 4, 831-869.

- 1832 Winsborrow, M.C.M., Clark, C.D., Stokes, C.R., 2004. Ice streams of the Laurentide Ice
1833 Sheet. *Géog. Phys. Quatern.* 58 (2–3), 269–280.
- 1834 Winsborrow, M.C.M., Clark, C.D., Stokes, C.R., 2010. What controls the location of ice
1835 streams? *Earth-Science Reviews*, 103, 45-59.
- 1836 Wright, H.E., 1973. Tunnel valleys, glacial surges, and subglacial hydrology of the Superior
1837 Lobe, Minnesota. In, Black, R.F., Goldthwait, R.B., Willman, H.B. (Eds.), *The*
1838 *Wisconsinan Stage*. Geological Society of America Memoirs. Geological Society of
1839 America, pp. 251–276.
- 1840
- 1841

1842 **Figures:**



1843
1844
1845
1846
1847
1848

Figure 1: An outline of Flint's (1971) portrayal of the maximum extent of the Laurentide Ice Sheet east of the Cordillera during the Quaternary (simplified and redrawn from Ives, 1978). Note that the Cordilleran Ice Sheet was not depicted in the original version.




1849
 1850
 1851
 1852
 1853
 1854
 1855

Figure 2: Reconstruction of the North American ice sheets at 21.8 ka (18 ¹⁴C ka), simplified and redrawn from Dyke and Prest (1987). The three major domes of the LIS over Quebec/Labrador (Q-L), Keewatin (K) and Foxe Basin (F) are labelled. Major ice divides shown in red with lower-lying ‘saddles’ in the ice sheet surface labelled ‘S’.




1856
 1857
 1858
 1859
 1860
 1861
 1862
 1863

Figure 3: Revised reconstruction of the North American ice sheets at 21.8 ka (18 ¹⁴C ka) simplified and redrawn from Dyke *et al.* (2002). Note the increase in the areal extent of the ice sheet compared to Fig. 2, especially over the Queen Elizabeth Islands and at the south-eastern margin (e.g. over Newfoundland), but with Banks Island and Prince Patrick Island remaining ice-free.



1864
 1865
 1866
 1867
 1868
 1869
 1870
 1871
 1872
 1873

Figure 4: A recent reconstruction of the Laurentide Ice Sheet redrawn from Stokes *et al.* (2016). Note the expansion of the ice sheet margin over Banks and Prince Patrick Island. This reconstruction also includes the location of 117 hypothesised ice streams (dark blue with flow-lines) in the Laurentide Ice Sheet based on published literature and new mapping in Margold *et al.* (2015a). Note that the ice streams did not all operate at the LLGM and that the inventory excluded the Cordilleran Ice Sheet.




1874
 1875
 1876
 1877
 1878
 1879
 1880
 1881
 1882

Figure 5: Weighted mean basal velocity and surface elevation of the North American Ice Sheet complex at 20 ka taken from Tarasov *et al.* (2012). Note that this ensemble is not representative of a single glaciologically-self-consistent model run and that the weighted averaging also blurs ice stream locations and magnitudes, and smooths ice surface topography. It is simply the expectation value. Note, however, that the mean captures some of the major ice-streams (compare with Fig. 4) and most of the key features of the geologically inferred reconstruction of Dyke and Prest (1987) (Fig. 3).



1883
 1884
 1885
 1886
 1887
 1888
 1889

Figure 6: Radiocarbon-constrained isochrones of the retreat of the Laurentide Ice Sheet redrawn from Bryson *et al.* (1969). Note that ages are in ¹⁴C years BP. Dots indicate location of radiocarbon dates and dashed lines indicate larger uncertainty.



1890
 1891
 1892
 1893
 1894

Figure 7: Reconstruction of the retreat pattern of the Laurentide and Cordilleran ice sheets redrawn from Prest (1969).




Figure 8: Examples of the reconstruction of the North American ice sheet during deglaciation redrawn from Dyke (2004): (a) 17 ka, (b) 12.8 ka, (c) 11.7 ka and (d) 7.8 ka.

1895
1896
1897
1898
1899





Figure 9: Percentage area deglaciated in North America compared to its Local Last Glacial Maximum, redrawn from Dyke (2004).

1900
 1901
 1902
 1903
 1904
 1905





1906
 1907
 1908
 1909
 1910
 1911
 1912
 1913

Figure 10: Reconstructed limits of the final deglaciation of the LIS in Quebec/Labrador, redrawn from Ullman *et al.* (2016). Blue lines show Dyke’s (2004) reconstruction at 8.4 – 8.2 ka and red lines show maximum (dashed) and minimum (dotted) 7.6 ka ice areas from Ullman *et al.* (2016) in relation to the Sakami, North Shore, and Paradise moraines (see Occhietti *et al.*, 2011).



1914
 1915
 1916
 1917
 1918
 1919
 1920
 1921

Figure 11: Modelled surface mass balance of the Laurentide Ice Sheet at various time-slices, with black lines demarcating various sectors of the ice sheet: St Lawrence (SL), Midwestern (MW), Southwestern (SW), Northwestern (NW), Hudson Bay (HB), Banks Island (BI) and Newfoundland (NL). Reprinted by permission from Macmillan Publishers Ltd: *Nature Geoscience* (Ullman *et al.*, 2015b), copyright (2015).



1922
 1923
 1924
 1925
 1926
 1927
 1928

Figure 12: (a) Denton and Hughes' (1981) reconstruction of the Northern Hemisphere ice sheets at the global LGM with their hypothesised ice streams as black arrows. (b) Hypothesised ice streams based on a review of the literature from Stokes and Clark (2001). Compare with the latest inventory of 117 Laurentide ice streams shown in Fig. 4.




Figure 13: Ice flow velocity of the Antarctic ice sheet compared (at the same spatial scale) with reconstructions of ice stream activity in the LIS at selected time steps, from Stokes *et al.* (2016). (a) Present-day Antarctic ice sheet velocity, with red lines indicating where ice streams intersect the grounding line, which is a similar in cumulative proportion (~30% of the perimeter) to the LIS (Margold *et al.*, 2015b). (b, c, d) Ice streams reconstructed for the LIS at its LLGM (approximately 21.8 kyr ago), 13.9 ka, and 10.1 ka. The locations of ice streams that were active at the given time are shown in blue and numbered in black. Those that switched off within the preceding 1 ka are shown in grey and those that switched on during the subsequent 1 ka are shown in dark blue with numbers in red. For cross-referencing, the numbers refer to the inventory numbers in Margold *et al.* (2015a, b).

1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941